

This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.

Patient Group Direction

Mounjaro (tirzepatide) Injection

for weight management

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	• Pharmacist registered and practicing with the GPhC • Pharmacy technician registered and practicing with the GPhC
Training and assessment	Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. • All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines • All users must be competent in weight management consultation, BMI calculation and the recognition of red flags requiring referral • All users must previously have used PGD's to supply medication • Must work in compliance with the SOPs of their own employer • All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	• All users must be up to date with CPD requirements and appraisal • All users must be up to date with MHRA and safety alerts with regards to medical products • All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005

Initial Assessment

Before commencing treatment with weight management therapies, the patient must be assessed by a registered healthcare professional. The initial assessment should be face to face. The assessment should include, but may not be limited to:

- Causes of weight gain. Refer the patient to their GP if prescribed medication is causing weight gain.
- The patient's lifestyle, diet and level of exercise.
- Has the patient tried to lose weight in the past? Discuss what has and has not worked.
- Other factors that may be causing weight gain, including mental health, environmental and psychological factors. Consider signposting to another health professional, for example a mental health team.
- Other disease states that may predispose to weight gain. Consider referring to the GP for further advice.
- The patient's expectations of weight loss, and whether they are realistic for their height and body frame.
- Calculate BMI and determine the ideal weight. Set a realistic target weight and agree review intervals.

Clinical condition or situation to which this PGD applies

PGD Indication	Mounjaro (tirzepatide) is a dual GIP/GLP-1 receptor agonist indicated as an adjunct to a reduced-calorie diet and increased physical activity for weight management, including weight loss and weight maintenance, in adults with an initial Body Mass Index (BMI) of 30 kg/m ² or above (obesity), or 27 kg/m ² to below 30 kg/m ² (overweight) in the presence of
----------------	--

	at least one weight-related comorbidity, for example dysglycaemia (pre-diabetes or type 2 diabetes mellitus), hypertension, dyslipidaemia, obstructive sleep apnoea or cardiovascular disease.
Inclusion criteria	• Adults aged 18 to 75 years (inclusive). • BMI 30 kg/m² or above, or BMI 27 kg/m² or above with at least one weight-related comorbidity (for example hypertension, type 2 diabetes mellitus, dyslipidaemia, obstructive sleep apnoea, established cardiovascular disease). • Patient is willing to follow a reduced-calorie diet and increase physical activity in line with the agreed lifestyle plan. • Initial assessment completed and documented (see the Initial Assessment section above). • Informed consent obtained, including off-label considerations where applicable, the side-effect profile and treatment expectations. • No exclusion criteria present.
Exclusion criteria	• Adults under 18 years of age. • Adults aged over 75 years (upper age limit under this PGD; refer to a specialist if treatment is being considered). • Pregnancy, breastfeeding, or planning pregnancy. Effective contraception is required; advise discontinuation at least 2 months before planned conception for semaglutide and 1 month for tirzepatide. • Known hypersensitivity to tirzepatide or to any of the excipients. • Personal or family history of medullary thyroid carcinoma (MTC) or Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). • History of pancreatitis, acute or chronic. • Severe gastrointestinal disease, including gastroparesis or severe persistent gastrointestinal disorder. • Current cholelithiasis (gallstones) or cholecystitis, or cholecystectomy within the last 3 months. • Obesity caused by an endocrinological disorder. If the patient was already overweight prior to that diagnosis, this exclusion may not apply. • Concurrent use of any other GLP-1 receptor agonist or insulin secretagogue used for weight management. • Type 1 diabetes mellitus. • Diabetic retinopathy. Treatment may worsen retinopathy; defer or refer to a specialist. • Patients on insulin or sulphonylureas whose GP is unwilling to monitor and adjust their hypoglycaemic regimen. • Severe renal impairment (eGFR below 30 mL/min/1.73 m²) or end-stage renal disease. • Severe hepatic impairment. • Known diagnosis of heart failure with reduced ejection fraction below 40%. • Active eating disorder (anorexia nervosa, bulimia, or binge-eating disorder under specialist care). • BMI below the PGD inclusion threshold. • Patients who, in the clinical judgement of the healthcare professional, are not suitable for the medication.
Cautions	• Counsel on gradual dose escalation and gastrointestinal side

	effects; consider delaying titration or reducing to the previous dose if significant symptoms occur. • History of suicidal ideation, or active severe mental illness. Ensure appropriate psychiatric oversight is in place, monitor mood at review, and refer if there is any concern. Do not supply where oversight is absent and concern exists. • Mild to moderate renal impairment: monitor for dehydration secondary to gastrointestinal side effects. • Concomitant oral medications: GLP-1 and GIP agents delay gastric emptying. Counsel patients on possible reduced absorption of oral medications, especially narrow therapeutic index drugs and oral contraceptives; non-oral contraception or a barrier method is advised during titration and for 4 weeks after each dose increase. • For individuals who take HRT, due to the lack of data regarding absorption, non-oral products (for example patch, gel, or levonorgestrel intrauterine device) may be considered. • Tirzepatide delays gastric emptying and thereby has the potential to impact the rate of absorption of concomitantly administered oral medicinal products, especially those with a narrow therapeutic index. Upon initiation of tirzepatide treatment in patients on warfarin, frequent monitoring of INR is recommended. • Cases of pulmonary aspiration have been reported in patients receiving GLP-1 receptor agonists undergoing general anaesthesia or deep sedation for surgery. The increased risk of residual gastric content due to delayed gastric emptying should therefore be considered before procedures under general anaesthesia or deep sedation. • Cases of tachycardia have been reported. In patients with a pre-existing increased heart rate, use with caution and seek specialist advice before use. If a patient experiences a clinically relevant sustained increase in resting heart rate, treatment should be discontinued and advice sought. • Counsel on the signs and symptoms of pancreatitis; discontinue and refer urgently if suspected. • Sulfonylurea or insulin co-use in patients with comorbid type 2 diabetes carries a hypoglycaemia risk. Refer to the prescribing GP or specialist before starting. • Risk of dehydration. Counsel on adequate fluid intake during gastrointestinal side effects. • Reassess the benefit-risk balance if there is no clinically meaningful weight loss, typically less than 5%, after 6 months on the maintenance dose. • Rotate injection sites to reduce local irritation.
Actions if patient is excluded or declines treatment	• Discuss the reason for exclusion with the patient and ensure they understand. • Advise on alternative treatment options and how these can be accessed, for example the GP, a specialist weight management service, or lifestyle programmes. • If the exclusion relates to an undiagnosed or unmanaged comorbidity, recommend GP review. • Document any advice given and the decision reached. Inform or refer to the GP as appropriate.

Details of the medicine

Name, form and strength of medicine	Mounjaro (tirzepatide) solution for injection in pre-filled pen (KwikPen or single-dose pen depending on the product supplied). Available strengths: 2.5 mg, 5 mg, 7.5 mg, 10 mg, 12.5 mg and 15 mg per 0.5 mL dose.
Legal category	POM
Storage	• Store in a refrigerator (2°C to 8°C). Do not freeze; discard if frozen. • Once removed from refrigeration, may be stored unrefrigerated below 30°C for up to 21 days. Discard if not used within this period. • Keep the pen in the outer carton in order to protect from light. • Travel: carry pens in hand luggage when travelling by air. Pens should not be placed in checked baggage where temperatures cannot be guaranteed. A travel letter from the prescriber or pharmacist may be required.
Route/method of administration	• Subcutaneous injection in the abdomen, thigh or upper arm. Rotate injection sites. • Administered once weekly, on the same day each week, with or without food, at any time of day. • The day of weekly administration can be changed if necessary as long as the time between two doses is at least 3 days (more than 72 hours). After selecting a new dosing day, once-weekly dosing should be continued. • Do not administer intravenously or intramuscularly.
Dose and frequency	• Starting dose: 2.5 mg once weekly for 4 weeks. This is a titration dose and is not intended for sustained weight management. • After 4 weeks at 2.5 mg, increase to 5 mg once weekly. • Further increases of 2.5 mg may be made after a minimum of 4 weeks on the current dose if additional weight management is required and the current dose is tolerated. • Recommended maintenance doses: 5 mg, 10 mg or 15 mg once weekly. Maximum dose 15 mg once weekly. • If a dose is missed, administer as soon as possible within 4 days (96 hours). If more than 4 days have passed, skip the missed dose and resume the usual schedule. Do not administer two doses within 3 days. • If significant gastrointestinal symptoms occur, consider delaying dose escalation or reducing to the previous dose until symptoms have improved. • If more than 2 doses are missed, reduce and re-escalate the dose to avoid unwanted side effects when treatment is re-initiated.
Quantity to be administered and/or supplied	• One pre-filled pen contains one weekly dose at the prescribed strength. • Supply 4 pens (one month of treatment) per patient appointment. • If the patient is going on holiday, their appointment may be brought forward by a few days to allow collection of medication and ensure continuity of treatment. • This PGD does not allow additional medication to be supplied to enable patients to stock up.
Monitoring and review	• Reassess clinical benefit, tolerability and target weight at each visit. • If the patient has not lost at least 5% of their initial body weight

	after 6 months on the maximum tolerated dose, a decision is required on whether to continue treatment, taking into account the benefit-risk profile in that patient. • When the patient reaches their target weight, discuss whether to continue treatment to maintain it. • If the patient has used Mounjaro in the past and wants to recommence treatment, the dose must be titrated again following the schedule above. The BMI inclusion criteria for initiation must be reapplied if more than 2 months have passed since discontinuing treatment.
Adverse effects	Very common: nausea, diarrhoea, vomiting, constipation and decreased appetite. Common: abdominal pain, dyspepsia, eructation, flatulence, abdominal distension, gastro-oesophageal reflux, injection site reactions, hypoglycaemia when used with a sulfonylurea or insulin, fatigue, dizziness, hair loss and tachycardia. Uncommon or rare: acute pancreatitis, cholelithiasis, cholecystitis, hypersensitivity reactions including anaphylaxis, and acute kidney injury usually secondary to dehydration. Counsel on warning symptoms, particularly severe abdominal pain, severe vomiting with dehydration and jaundice. Refer to the current SmPC for full safety information.
Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: • that valid informed consent was given • name of individual, address, date of birth and GP with whom the individual is registered • name of healthcare practitioner • name and brand of medication, and batch number • date of supply, dose, form, route and quantity supplied • height, weight and BMI at this visit, and the target weight agreed • advice given, including advice given if excluded or declines treatment • details of any adverse drug reactions and actions taken • that the medicine was supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. Adults aged 18 and over: keep records for 8 years.

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medicine, together with written lifestyle, diet and physical activity advice and the agreed target weight.
Follow-up advice to be given to patient or carer	Explain the expected pattern of weight loss and that the medicine works alongside diet and activity, not instead of them. Counsel on gastrointestinal side effects and how to manage them, on adequate fluid intake, and on the warning symptoms that need urgent attention: severe abdominal pain, persistent vomiting with dehydration, jaundice, sudden visual loss, or a sustained rise in resting heart rate. Advise when to return for review and that treatment will be reassessed if less than 5% of body weight has been lost after 6 months on the maintenance dose.

Key references

• Summary of Product Characteristics for the product supplied, current version
--

- NICE guidance on obesity management and on the relevant medicine
- MHRA Drug Safety Updates relevant to GLP-1 receptor agonists
- British National Formulary (BNF)
- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates	

Name of healthcare professional	Job title	Registration number	Signature

Valid from date	Expiry date

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or vaccine.

This section must be signed by the superintendent or clinical lead responsible for this service.

To be completed by the adopting pharmacy: the first block is signed by that pharmacy's superintendent or clinical lead, and the second by its professional group signatory. Get Real Health signs the authorship block below, not these.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date
Name of professional group signatory	Job title & name of organisation	Signature	Date

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		06/08/2026

Chris Pilkington	Pharmacist GPhC: 2046322		06/08/2026
------------------	-----------------------------	--	------------

Change history

Version number	Change details	Date
001	Development and issue of new PGD	1st May 2026
002	PPH clinical review (J. Wilkins with S. Passmore). Adjusted BMI thresholds for ethnic groups removed from the inclusion criteria as they are not in the SmPC. Gallbladder disease moved from cautions to exclusions as current cholelithiasis or cholecystitis, or cholecystectomy within 3 months. Obesity caused by an endocrinological disorder added as an exclusion. History of suicidal ideation or active severe mental illness moved from exclusions to cautions. Cautions added covering HRT absorption, delayed gastric emptying with INR monitoring on warfarin, pulmonary aspiration under general anaesthesia or deep sedation, and tachycardia. Dosing now states that if more than 2 doses are missed the dose is reduced and re-escalated.	6th August 2026