

This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.

Patient Group Direction

for the administration of Semaglutide (Wegovy) 0.25mg, 0.5mg, 1mg, 1.7mg, 2.4mg solution for injection in pre-filled pen for the treatment of Weight Management (GLP-1 Receptor Agonist Monitoring)

Summary of NICE / NICE CKS Guidelines for Weight Management (GLP-1 Receptor Agonist Monitoring)

Overview

GLP-1 receptor agonists require ongoing monitoring for efficacy and safety throughout treatment.

Assessment

- **Baseline and regular intervals:** Weight, BMI, blood pressure, blood glucose/HbA1c, renal function, liver function, lipid profile.

Management

- Dose titration monitoring with documented tolerability
- Ongoing adverse effect assessment
- Weight loss trajectory review
- Comorbidity improvement tracking (blood pressure, lipids, glucose control)

Red Flags

- Signs of pancreatitis (severe upper abdominal pain, vomiting)
- Thyroid nodules or enlargement
- Severe gastrointestinal symptoms (persistent vomiting, abdominal distension)
- Significant mood changes or suicidal ideation
- Renal impairment progression

Patient Group Direction

for the supply/administration of

Semaglutide (Wegovy) 0.25mg, 0.5mg, 1mg, 1.7mg, 2.4mg solution for injection in pre-filled pen

for the treatment of

Weight Management (GLP-1 Receptor Agonist Monitoring)

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	• Pharmacist registered and practicing with the GPhC

	Pharmacy technician registered and practicing with the GPhC
Training and assessment	Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. - All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines - All users must previously have used PGDs to supply medication - Must work in compliance with the SOPs of their own employer - All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	- All users must be up to date with CPD requirements and appraisal - All users must be up to date with MHRA and safety alerts with regards to medical products - All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	Adjunct to reduced-calorie diet and increased physical activity for weight management in adults with BMI ≥ 30 kg/m ² or BMI ≥ 27 kg/m ² with weight-related comorbidity (hypertension, type 2 diabetes, dyslipidaemia, cardiovascular disease). Ongoing monitoring and dose titration required throughout treatment.
Inclusion criteria	Adults aged 18 years and over Already commenced on or being newly commenced on semaglutide for weight management Meet BMI criteria: BMI ≥ 30 kg/m² or BMI ≥ 27 kg/m² with weight-related comorbidity Valid informed consent obtained Baseline assessment completed (weight, BMI, blood pressure, blood glucose/HbA1c, renal function, liver function, lipid profile)
Exclusion criteria	Known hypersensitivity to semaglutide or any excipients Personal or family history of medullary thyroid carcinoma (MTC) Personal or family history of multiple endocrine neoplasia type 2 (MEN2) Pregnancy or breastfeeding (effective contraception required) Severe hepatic impairment End-stage renal disease (eGFR < 15 mL/min/1.73m²) Concurrent use of other GLP-1 receptor agonists Type 1 diabetes mellitus

	Active malignancy or history of MTC
Cautions	History of pancreatitis (acute or chronic) Diabetic retinopathy (may worsen with rapid weight loss or if concurrent diabetes) Heart failure (New York Heart Association class III or IV) Depression, anxiety, or suicidal ideation (monitor closely) Renal impairment (eGFR 15-60 mL/min/1.73m²; caution if severe) Hepatic impairment (mild to moderate) Concurrent use of insulin or sulfonylureas (hypoglycaemia risk) Inflammatory bowel disease or history of bowel obstruction Severe dehydration or acute illness
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate.

Details of the medicine

Name, form and strength of medicine	Semaglutide (Wegovy) 0.25mg, 0.5mg, 1mg, 1.7mg, 2.4mg solution for injection in pre-filled pen
Legal category	POM
Storage	Before use: store at 2-8°C (in refrigerator). In use: store below 30°C for maximum 28 days. Do not freeze.
Route/method of administration	Subcutaneous injection once weekly into the abdomen, thigh, or upper arm
Dose and frequency	Dose titration schedule: Start 0.25mg weekly for 4 weeks, then 0.5mg for 4 weeks, then 1mg for 4 weeks, then 1.7mg for 4 weeks, then 2.4mg weekly maintenance dose. If dose escalation is not tolerated, delay by 4 weeks before attempting next dose increase. Injection on same day each week; time of day is flexible.
Quantity to be administered and/or supplied	Up to 4 pre-filled pens per supply (4-week supply at any titration stage)
Maximum or minimum treatment period	Ongoing with review every 4 weeks during titration phase, then 3-monthly during maintenance. Discontinue if less than 5% weight loss achieved after 6 months on maintenance dose (2.4mg weekly).
Adverse effects	Very common (≥1/10): Nausea, diarrhoea, vomiting, constipation Common (≥1/100, <1/10): Headache, dizziness, abdominal pain, fatigue, dyspepsia, flatulence, cholelithiasis (gallstones), injection site

	reactions, hypoglycaemia (in patients with type 2 diabetes on concomitant medications) Uncommon ($\geq 1/1000$, $< 1/100$): Pancreatitis, thyroid disorder, acute kidney injury Serious (report immediately): Acute pancreatitis, anaphylaxis, severe dehydration, diabetic retinopathy worsening, gallbladder-related complications, severe gastrointestinal obstruction, suicidal thoughts or behaviour
--	---

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: - that valid informed consent was given - name of individual, address, date of birth and GP - name of healthcare practitioner - name and brand of medication - date of supply dose, form and route quantity supplied - advice given, including advice given if excluded or declines treatment - details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: - Adults aged 18 years and over - keep records for 8 years - Children aged under 18 years - keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the Wegovy pen. Ensure patient understands the importance of regular monitoring appointments and reporting adverse effects.
Follow-up advice to be given to patient or carer	Rotate injection sites to prevent lipohypertrophy (fatty tissue buildup). Use abdomen, thigh, or upper arm in rotation. Report severe or persistent upper abdominal pain, vomiting, or signs of pancreatitis immediately. Attend all monitoring appointments every 4 weeks during dose titration, then 3-monthly during maintenance. Report any mood changes, depression, or suicidal thoughts to your healthcare provider.

	Report any thyroid symptoms (neck lump, voice change, difficulty swallowing) immediately. Maintain reduced-calorie diet and increase physical activity as advised. If dose escalation causes intolerable nausea or vomiting, inform your healthcare provider; dose increase can be delayed. Store pens in refrigerator (2-8°C) before first use; after first use, store below 30°C for up to 28 days. Do not freeze pens. Do not use if frozen. If you forget an injection, take it as soon as you remember, then resume normal schedule the following week. Report any new symptoms or worsening of existing health conditions at monitoring appointments. Seek immediate medical attention if you experience signs of allergic reaction (rash, difficulty breathing, swelling).
--	---

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance: Weight Management (GLP-1 Receptor Agonist Monitoring)
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature

Valid from date	Expiry date		
1/11/25	31/10/26		

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date
Chris Pilkington	Head Pharmacist, Get Real Health Limited (GPhC: 2046322)		19/06/2026
Name of professional group signatory	Job title & name of organisation	Signature	Date
Chris Pilkington	Head Pharmacist, Get Real Health Limited (GPhC: 2046322)		19/06/2026

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		2/11/25
Chris Pilkington	Pharmacist GPhC: 2046322		2/11/25

Change history

Version number	Change details	Date
001	Development & issue of new PGD	1st November 2025
