

This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.

Patient Group Direction
for the administration of MMRVaxPRO or Priorix for vaccination against Measles, Mumps and Rubella (MMR)

Summary of NICE / NICE CKS Guidelines for MMR Vaccination

Overview

- The **MMR vaccine** is a **live attenuated vaccine** providing protection against:
 - **Measles** – highly contagious, risk of serious complications (e.g. encephalitis, pneumonia).
 - **Mumps** – can cause meningitis, orchitis, and hearing loss.
 - **Rubella** – especially dangerous in pregnancy, causing congenital rubella syndrome.

Routine Immunisation Schedule (UK)

Age	Dose
12–13 months	First dose of MMR
3 years 4 months (pre-school)	Second dose of MMR

- Can be given **from 6 months** in high-risk situations (e.g. outbreaks or travel to endemic areas), but this dose does **not count toward the routine schedule**.

Catch-Up Vaccination

- Recommended for **anyone born after 1970** who has not had **2 documented doses**.
- Offer opportunistically (e.g. during GP or travel clinic visits).
- Ensure a **4-week interval** between doses.

Eligibility

- Children as part of the routine programme.
 - **Unvaccinated adults** or those with incomplete vaccination history.
 - **Healthcare workers** without evidence of immunity.
 - **Students, travelers, and close contacts** during outbreaks.
-

Contraindications

- **Severe immunosuppression**
- **Pregnancy** (avoid conception for **1 month** after vaccination)
- History of **anaphylaxis to neomycin** or gelatin (if vaccine contains it)

Note: Egg allergy is *not* a contraindication.

Precautions

- Postpone in **acute febrile illness** (minor infections are not a reason to delay).
 - Consider specialist advice for those with HIV or immunodeficiency.
-

Side Effects

- Usually mild and self-limiting:
 - Fever, rash (5–10 days after), malaise
 - Mild parotid swelling (mumps-like)
 - Transient arthralgia (more common in adult women)
 - Rare: febrile convulsions, thrombocytopenia, anaphylaxis
-

Effectiveness

- Two doses provide:
 - **>99% protection** against **measles and rubella**
 - **88–95% protection** against **mumps**
-

Documentation and Reporting

- Record **vaccine name, batch number, site, route, and date**.
- Report adverse events to the **Yellow Card Scheme**.

Patient Group Direction

for the supply/administration of

MMRVaxPRO 5mg

for vaccination against

Measles, Mumps and Rubella (MMR)

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must previously have used PGD's to supply medication ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	MMRVaxPRO is indicated for active immunisation against measles, mumps and rubella in children from 12 months of age, in accordance with the national immunisation programme.
Inclusion criteria	- Children aged 12 months and over who are eligible under the national childhood immunisation programme. - Informed consent obtained from parent or guardian. - No contraindications to MMR vaccination.
Exclusion criteria	- Known hypersensitivity to neomycin, gelatin, or any other components of the vaccine. - Immunocompromised individuals or those receiving immunosuppressive therapy. - Pregnancy or planning pregnancy within one month. - Acute febrile illness (vaccination should be postponed).
Cautions	- Defer vaccination if the child has received blood products or immunoglobulins recently. - Use caution in children with a history of thrombocytopenia or febrile seizures. - Observe the patient for 15 minutes post-vaccination. - Provide information on common side effects and when to seek further medical advice.
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate

Details of the medicine

Name, form and strength of medicine	MMRVaxPRO
Legal category	POM
Storage	Store in a refrigerator (2°C–8°C). Do not freeze. Protect from light.
Route/method of administration	Subcutaneous injection, preferably in the upper arm or thigh.
Dose and frequency	0.5 mL per dose according to the national immunisation schedule (typically 2 doses).
Quantity to be administered and/or supplied	0.5 mL per dose.
Maximum or minimum treatment period	Two doses, usually given at age 12–13 months and a second dose before school entry.
Adverse effects	Fever, rash, irritability, swelling or pain at the injection site. Rare: febrile convulsions, thrombocytopenia, allergic reactions. Refer to SmPC.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information:

	● that valid informed consent was given ● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of supply dose, form and route quantity supplied ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)
--	---

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell.

Key references

● NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 https://www.nice.org.uk/guidance/mg2 ● NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 https://www.nice.org.uk/guidance/mpg2/resources ● NICE CKS Guidance ● British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines BNF (British National Formulary) NICE ● Summary of Product Characteristics

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature

Valid from date		Expiry date	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date
Chris Pilkington	Head Pharmacist, Get Real Health Limited (GPhC: 2046322)		19/06/2026

Name of professional group signatory	Job title & name of organisation	Signature	Date
Chris Pilkington	Head Pharmacist, Get Real Health Limited (GPhC: 2046322)		19/06/2026

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		25/8/2025
Chris Pilkington	Pharmacist GPhC: 2046322		25/8/2025

Change history

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025

Patient Group Direction

for the supply/administration of

Priorix

for vaccination against

Measles, Mumps and Rubella (MMR)

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must previously have used PGD's to supply medication ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	Priorix is indicated for active immunisation against measles, mumps and rubella in children from 12 months of age, in accordance with the national immunisation programme.
Inclusion criteria	- Children aged 12 months and over who are eligible under the national childhood immunisation programme. - Informed consent obtained from parent or guardian. - No contraindications to MMR vaccination.
Exclusion criteria	- Known hypersensitivity to neomycin, gelatin, or any other components of the vaccine. - Immunocompromised individuals or those receiving immunosuppressive therapy. - Pregnancy or planning pregnancy within one month. - Acute febrile illness (vaccination should be postponed).
Cautions	- Defer vaccination if the child has received blood products or immunoglobulins recently. - Use caution in children with a history of thrombocytopenia or febrile seizures. - Observe the patient for 15 minutes post-vaccination. - Provide information on common side effects and when to seek further medical advice.
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate

Details of the medicine

Name, form and strength of medicine	Priorix
Legal category	POM
Storage	Store in a refrigerator (2°C–8°C). Do not freeze. Protect from light.
Route/method of administration	Subcutaneous injection, preferably in the upper arm or thigh.
Dose and frequency	0.5 mL per dose according to the national immunisation schedule (typically 2 doses).
Quantity to be administered and/or supplied	0.5 mL per dose.
Maximum or minimum treatment period	Two doses, usually given at age 12–13 months and a second dose before school entry.
Adverse effects	Redness, swelling and pain at injection site, fever, rash. Rare: febrile convulsions, allergic reactions. Refer to SmPC.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information:

	● that valid informed consent was given ● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of supply dose, form and route quantity supplied ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)
--	---

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell.

Key references

● NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 https://www.nice.org.uk/guidance/mg2 ● NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 https://www.nice.org.uk/guidance/mpg2/resources ● NICE CKS Guidance ● British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines BNF (British National Formulary) NICE ● Summary of Product Characteristics

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature

Valid from date		Expiry date	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date
Chris Pilkington	Head Pharmacist, Get Real Health Limited (GPhC: 2046322)		19/06/2026

Name of professional group signatory	Job title & name of organisation	Signature	Date
Chris Pilkington	Head Pharmacist, Get Real Health Limited (GPhC: 2046322)		19/06/2026

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		25/8/2025
Chris Pilkington	Pharmacist GPhC: 2046322		25/8/2025

Change history

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025