

Dengue Vaccine

Vaccination safety requirements

The signed PGD behind this page is unaltered. This correction takes precedence over it.

What is missing from the signed document

- No requirement for adrenaline.
- No observation period after vaccination.
- No cold chain excursion procedure.
- No sharps disposal provision.
- No requirement to record the batch number.

What must be in place before you vaccinate under this PGD

- Adrenaline (epinephrine) 1 in 1,000 injection must be immediately available in the room where vaccination takes place, in date, together with a telephone.
- A written anaphylaxis protocol consistent with current Resuscitation Council UK guidance must be available, and every person administering must be trained in the recognition and immediate management of anaphylaxis and hold current basic life support training.
- Observe every patient for 15 minutes after vaccination. Vaccinate seated, and have procedures in place to prevent injury from a faint.
- Record the batch number and the anatomical site for every dose given. A recall cannot be actioned without the batch number.

If these are not in place

Do not vaccinate under this PGD. This is not a documentation point: anaphylaxis after vaccination is rare, and it is the thing that kills. The requirement exists so that the one time it happens, the adrenaline is in the room.

Why this notice exists

A review of every vaccination PGD on the platform on 8 September 2026 found that these requirements were stated in some signed documents and not others. Where they were absent, nothing in the signed document required adrenaline to be present. Pharmacies almost certainly have it and are trained; the problem is that the document did not require it, so it was not evidenced anywhere if anyone asked.

A corrected version of this PGD will be issued and will carry these requirements in the body of the document. Until then, this notice applies.

This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.

Patient Group Direction

for the administration of Qdenga (TAK-003) for the treatment of Dengue Fever Prevention

Summary of NICE / NICE CKS Guidelines for Dengue Fever Prevention

Overview

Dengue fever is caused by the dengue virus, of which there are four serotypes (DENV-1, DENV-2, DENV-3, DENV-4). The disease is transmitted by Aedes mosquitoes and is endemic in tropical and subtropical regions worldwide, affecting over 400 million people annually.

Assessment

- **Travel destination risk:** Assess if the patient is travelling to or will reside in dengue-endemic areas.
- **Previous dengue infection:** Obtain history of previous dengue infection, if known.
- **Duration of stay:** Evaluate the intended duration and timing of travel.
- **Immunisation status:** Confirm current vaccinations and previous dengue serology if available.

Management

- **Vaccination schedule:** Two doses of Qdenga administered 3 months apart via subcutaneous injection.
- **Mosquito bite prevention:** Advise on insect repellents, protective clothing, and accommodation with air conditioning and/or screens.
- **Timing:** First dose should be administered at least 3 months prior to travel when possible.

Red Flags and Contraindications

- **Immunocompromised individuals:** Live attenuated vaccine is contraindicated in those with severe immunocompromise.
- **Pregnancy and breastfeeding:** Vaccine is not recommended during pregnancy or breastfeeding.
- **Severe acute illness:** Defer vaccination if the patient has acute febrile illness or other significant illness.
- **Concurrent live vaccines:** If other live vaccines are required, Qdenga should be given simultaneously or separated by at least 4 weeks.

Patient Group Direction

for the supply/administration of

Qdenga (TAK-003)

for the treatment of

Dengue Fever Prevention

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	• Pharmacist registered and practicing with the GPhC • Pharmacy technician registered and practicing with the GPhC
Training and assessment	Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. • All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines • All users must previously have used PGDs to supply medication • Must work in compliance with the SOPs of their own employer • All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	• All users must be up to date with CPD requirements and appraisal • All users must be up to date with MHRA and safety alerts with regards to medical products • All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	Immunisation against dengue fever in adults aged 18 years and over travelling to or residing in dengue-endemic areas.
Inclusion criteria	Age 18 years or over Planning travel to or residence in dengue-endemic area(s) Willing to receive two doses, 3 months apart Able to provide informed consent Not pregnant and not breastfeeding Not severely immunocompromised
Exclusion criteria	Age under 18 years Pregnancy or breastfeeding Known hypersensitivity to any component of the vaccine Severe immunocompromise (including HIV with CD4 count < 200, active malignancy, prolonged high-dose corticosteroids) Acute fever or significant intercurrent illness Planned administration of another live vaccine within 4 weeks before or after Qdenga History of Guillain-Barré syndrome following prior dengue

	vaccination
Cautions	Use with caution in patients with mild to moderate immunosuppression Use with caution in patients on anticoagulant therapy (assess bleeding risk) Patients should avoid pregnancy for at least 4 weeks after each dose Live vaccine - not suitable for severely immunocompromised individuals Consider serological testing in those with previous dengue infection Administration should be deferred in patients with acute febrile illness
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate.

Details of the medicine

Name, form and strength of medicine	Qdenga (TAK-003), suspension for injection in pre-filled syringe, 0.5 mL
Legal category	POM (Prescription Only Medicine)
Storage	Store in a refrigerator between 2°C and 8°C. Protect from light. Do not freeze. After reconstitution, use within 30 minutes.
Route/method of administration	Subcutaneous injection, preferably in the deltoid area of the upper arm
Dose and frequency	First dose: 0.5 mL subcutaneously Second dose: 0.5 mL subcutaneously, administered 3 months after the first dose
Quantity to be administered and/or supplied	2 doses of 0.5 mL each over a 3-month period
Maximum or minimum treatment period	Vaccination series over 3 months (with optional booster after 12 months based on ongoing exposure risk)
Adverse effects	Very common ($\geq 1/10$): Injection site pain, headache, myalgia Common ($\geq 1/100$ to $< 1/10$): Malaise, fatigue, fever (usually within 7 days of vaccination) Uncommon ($\geq 1/1,000$ to $< 1/100$): Rash, pruritus, arthralgia Rare: Guillain-Barré syndrome (very rare, temporal relationship to vaccination unclear) Report all suspected adverse effects via the Yellow Card scheme at

	https://yellowcard.mhra.gov.uk
--	---

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: • that valid informed consent was given • name of individual, address, date of birth and GP • name of healthcare practitioner • name and brand of medication • date of supply dose, form and route quantity supplied • advice given, including advice given if excluded or declines treatment • details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: • Adults aged 18 years and over - keep records for 8 years • Children aged under 18 years - keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with Qdenga. Ensure the patient understands the vaccine schedule, storage requirements, and what to expect after vaccination.
Follow-up advice to be given to patient or carer	Attend for second dose appointment in 3 months Report any unusual or severe symptoms to your healthcare provider or via the Yellow Card scheme Seek medical advice if fever develops or persists beyond 7 days after vaccination Avoid pregnancy for at least 4 weeks after each dose if of childbearing potential Continue mosquito bite prevention measures (insect repellent, protective clothing) even after vaccination Seek immediate medical attention if you develop rash, difficulty breathing, or signs of anaphylaxis after vaccination If you experience signs of dengue fever (fever, headache, rash, joint pain) whilst travelling, seek medical attention promptly Keep a record of your vaccination dates and bring documentation

	when travelling
--	-----------------

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017
<https://www.nice.org.uk/guidance/mg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017
<https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance: Dengue Fever Immunisation and Prevention
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature
Valid from date	Expiry date		
1/11/25	31/7/27		

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date
Name of professional group signatory	Job title & name of organisation	Signature	Date

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		2/11/25
Chris Pilkington	Pharmacist GPhC: 2046322		2/11/25

Change history

Version number	Change details	Date
001	Development & issue of new PGD	1st November 2025