

**This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.**

Patient Group Direction

for the administration of Qdenga (TAK-003) for the treatment of Dengue Fever Prevention

## Summary of NICE / NICE CKS Guidelines for Dengue Fever Prevention

### Overview

Dengue fever is caused by the dengue virus, of which there are four serotypes (DENV-1, DENV-2, DENV-3, DENV-4). The disease is transmitted by Aedes mosquitoes and is endemic in tropical and subtropical regions worldwide, affecting over 400 million people annually.

### Assessment

- **Travel destination risk:** Assess if the patient is travelling to or will reside in dengue-endemic areas.
- **Previous dengue infection:** Obtain history of previous dengue infection, if known.
- **Duration of stay:** Evaluate the intended duration and timing of travel.
- **Immunisation status:** Confirm current vaccinations and previous dengue serology if available.

### Management

- **Vaccination schedule:** Two doses of Qdenga administered 3 months apart via subcutaneous injection.
- **Mosquito bite prevention:** Advise on insect repellents, protective clothing, and accommodation with air conditioning and/or screens.
- **Timing:** First dose should be administered at least 3 months prior to travel when possible.

### Red Flags and Contraindications

- **Immunocompromised individuals:** Live attenuated vaccine is contraindicated in those with severe immunocompromise.
- **Pregnancy and breastfeeding:** Vaccine is not recommended during pregnancy or breastfeeding.
- **Severe acute illness:** Defer vaccination if the patient has acute febrile illness or other significant illness.
- **Concurrent live vaccines:** If other live vaccines are required, Qdenga should be given simultaneously or separated by at least 4 weeks.

### Patient Group Direction

for the supply/administration of

**Qdenga (TAK-003)**

for the treatment of

**Dengue Fever Prevention**

## Healthcare professionals covered and training

|                                              | Healthcare professionals allowed to work under this PGD and requirements                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|----------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Qualifications and Registration</b>       | <ul style="list-style-type: none"> <li>• Pharmacist registered and practicing with the GPhC</li> <li>• Pharmacy technician registered and practicing with the GPhC</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Training and assessment</b>               | <p>Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team.</p> <ul style="list-style-type: none"> <li>• All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines</li> <li>• All users must previously have used PGDs to supply medication</li> <li>• Must work in compliance with the SOPs of their own employer</li> <li>• All users must have appropriate indemnity in place to cover this service</li> </ul> |
| <b>Further training and responsibilities</b> | <ul style="list-style-type: none"> <li>• All users must be up to date with CPD requirements and appraisal</li> <li>• All users must be up to date with MHRA and safety alerts with regards to medical products</li> <li>• All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.</li> </ul>                                                                                                                                                                                                                                                             |

|                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|---------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>PGD Indication</b>     | Immunisation against dengue fever in adults aged 18 years and over travelling to or residing in dengue-endemic areas.                                                                                                                                                                                                                                                                                                                                                        |
| <b>Inclusion criteria</b> | <p>Age 18 years or over</p> <p>Planning travel to or residence in dengue-endemic area(s)</p> <p>Willing to receive two doses, 3 months apart</p> <p>Able to provide informed consent</p> <p>Not pregnant and not breastfeeding</p> <p>Not severely immunocompromised</p>                                                                                                                                                                                                     |
| <b>Exclusion criteria</b> | <p>Age under 18 years</p> <p>Pregnancy or breastfeeding</p> <p>Known hypersensitivity to any component of the vaccine</p> <p>Severe immunocompromise (including HIV with CD4 count &lt; 200, active malignancy, prolonged high-dose corticosteroids)</p> <p>Acute fever or significant intercurrent illness</p> <p>Planned administration of another live vaccine within 4 weeks before or after Qdenga</p> <p>History of Guillain-Barré syndrome following prior dengue</p> |

|                                                             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|-------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                             | vaccination                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Cautions</b>                                             | <p>Use with caution in patients with mild to moderate immunosuppression</p> <p>Use with caution in patients on anticoagulant therapy (assess bleeding risk)</p> <p>Patients should avoid pregnancy for at least 4 weeks after each dose</p> <p>Live vaccine - not suitable for severely immunocompromised individuals</p> <p>Consider serological testing in those with previous dengue infection</p> <p>Administration should be deferred in patients with acute febrile illness</p> |
| <b>Actions if patient is excluded or declines treatment</b> | <p>Advise on alternative treatment options and how these can be accessed.</p> <p>Document any advice given and the decision reached. Inform or refer to the GP as appropriate.</p>                                                                                                                                                                                                                                                                                                    |

#### *Details of the medicine*

|                                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|----------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Name, form and strength of medicine</b>         | Qdenga (TAK-003), suspension for injection in pre-filled syringe, 0.5 mL                                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Legal category</b>                              | POM (Prescription Only Medicine)                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Storage</b>                                     | Store in a refrigerator between 2°C and 8°C. Protect from light. Do not freeze. After reconstitution, use within 30 minutes.                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Route/method of administration</b>              | Subcutaneous injection, preferably in the deltoid area of the upper arm                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Dose and frequency</b>                          | <p>First dose: 0.5 mL subcutaneously</p> <p>Second dose: 0.5 mL subcutaneously, administered 3 months after the first dose</p>                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Quantity to be administered and/or supplied</b> | 2 doses of 0.5 mL each over a 3-month period                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Maximum or minimum treatment period</b>         | Vaccination series over 3 months (with optional booster after 12 months based on ongoing exposure risk)                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Adverse effects</b>                             | <p>Very common (<math>\geq 1/10</math>): Injection site pain, headache, myalgia</p> <p>Common (<math>\geq 1/100</math> to <math>&lt; 1/10</math>): Malaise, fatigue, fever (usually within 7 days of vaccination)</p> <p>Uncommon (<math>\geq 1/1,000</math> to <math>&lt; 1/100</math>): Rash, pruritus, arthralgia</p> <p>Rare: Guillain-Barré syndrome (very rare, temporal relationship to vaccination unclear)</p> <p>Report all suspected adverse effects via the Yellow Card scheme at</p> |

|  |                                                                             |
|--|-----------------------------------------------------------------------------|
|  | <a href="https://yellowcard.mhra.gov.uk">https://yellowcard.mhra.gov.uk</a> |
|--|-----------------------------------------------------------------------------|

|                              |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Yellow card reporting</b> | Report suspected adverse effects via <a href="https://yellowcard.mhra.gov.uk">https://yellowcard.mhra.gov.uk</a> and inform the GP as appropriate.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Records to be kept</b>    | <p>Record the following information:</p> <ul style="list-style-type: none"> <li>• that valid informed consent was given</li> <li>• name of individual, address, date of birth and GP</li> <li>• name of healthcare practitioner</li> <li>• name and brand of medication</li> <li>• date of supply dose, form and route quantity supplied</li> <li>• advice given, including advice given if excluded or declines treatment</li> <li>• details of any adverse drug reactions and actions taken supplied via PGD</li> </ul> <p>Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made.</p> <p>A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows:</p> <ul style="list-style-type: none"> <li>• Adults aged 18 years and over - keep records for 8 years</li> <li>• Children aged under 18 years - keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)</li> </ul> |

### ***Patient information***

|                                                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Written information to be given to patient or carer</b> | Supply the patient information leaflet (PIL) provided with Qdenga. Ensure the patient understands the vaccine schedule, storage requirements, and what to expect after vaccination.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Follow-up advice to be given to patient or carer</b>    | <p>Attend for second dose appointment in 3 months</p> <p>Report any unusual or severe symptoms to your healthcare provider or via the Yellow Card scheme</p> <p>Seek medical advice if fever develops or persists beyond 7 days after vaccination</p> <p>Avoid pregnancy for at least 4 weeks after each dose if of childbearing potential</p> <p>Continue mosquito bite prevention measures (insect repellent, protective clothing) even after vaccination</p> <p>Seek immediate medical attention if you develop rash, difficulty breathing, or signs of anaphylaxis after vaccination</p> <p>If you experience signs of dengue fever (fever, headache, rash, joint pain) whilst travelling, seek medical attention promptly</p> <p>Keep a record of your vaccination dates and bring documentation</p> |

|  |                 |
|--|-----------------|
|  | when travelling |
|--|-----------------|

## Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017  
<https://www.nice.org.uk/guidance/mg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017  
<https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance: Dengue Fever Immunisation and Prevention
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines

## Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

|                                                                                   |             |                     |           |
|-----------------------------------------------------------------------------------|-------------|---------------------|-----------|
| Name and address of pharmacy premises / clinic premises to which this PGD relates |             |                     |           |
|                                                                                   |             |                     |           |
| Name of healthcare professional                                                   | Job title   | Registration number | Signature |
|                                                                                   |             |                     |           |
|                                                                                   |             |                     |           |
|                                                                                   |             |                     |           |
|                                                                                   |             |                     |           |
|                                                                                   |             |                     |           |
|                                                                                   |             |                     |           |
|                                                                                   |             |                     |           |
|                                                                                   |             |                     |           |
| Valid from date                                                                   | Expiry date |                     |           |
| 1/11/25                                                                           | 31/10/26    |                     |           |

## PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

| Name of superintendent / clinical lead | Job title & name of organisation                         | Signature                                                                          | Date       |
|----------------------------------------|----------------------------------------------------------|------------------------------------------------------------------------------------|------------|
| Chris Pilkington                       | Head Pharmacist, Get Real Health Limited (GPhC: 2046322) |  | 19/06/2026 |
| Name of professional group signatory   | Job title & name of organisation                         | Signature                                                                          | Date       |
| Chris Pilkington                       | Head Pharmacist, Get Real Health Limited (GPhC: 2046322) |  | 19/06/2026 |

#### Signed on behalf of Get Real Health

| Name             | Qualification                  | Signature                                                                            | Date    |
|------------------|--------------------------------|--------------------------------------------------------------------------------------|---------|
| Nitin Shori      | Medical Doctor<br>GMC: 6047293 |    | 2/11/25 |
| Chris Pilkington | Pharmacist<br>GPhC: 2046322    |  | 2/11/25 |

#### Change history

| Version number | Change details                 | Date              |
|----------------|--------------------------------|-------------------|
| 001            | Development & issue of new PGD | 1st November 2025 |
|                |                                |                   |
|                |                                |                   |
|                |                                |                   |

## **VACCINE SAFETY REQUIREMENTS, ADDED AT VERSION V002, 9 September 2026**

The requirements on the following pages form part of this Patient Group Direction and must be met before any vaccine is administered under it. They were absent from the signed document and were added by this reissue after an estate-wide safety sweep of every vaccination PGD.

This reissue adds the safety requirements below and changes nothing else. It is NOT a clinical review of this PGD, and one remains outstanding.

**Anaphylaxis: what must be in place before you vaccinate**

ADRENALINE (EPINEPHRINE) 1 IN 1,000 INJECTION must be immediately available in the room where vaccination takes place, in date, together with a telephone.

A written anaphylaxis protocol consistent with current Resuscitation Council UK guidance must be available, and every person administering must be trained in the recognition and immediate management of anaphylaxis and hold current basic life support training.

Be able to distinguish a faint and a panic attack from anaphylaxis. Green Book chapter 8 sets out the clinical features of each. A faint after an injection is common; anaphylaxis is very rare.

Report any suspected anaphylaxis via the Yellow Card Scheme even where the diagnosis is uncertain.

**Observation after vaccination**

OBSERVE EVERY PATIENT FOR 15 MINUTES AFTER VACCINATION, SEATED.

Anxiety-related reactions including vasovagal syncope, hyperventilation and transient visual disturbance or paraesthesia can occur before or after any injection, and are commonest in adolescents. They are a response to the needle, not to the vaccine.

Have procedures in place to prevent injury from a faint.

Record that the observation period was completed.

**Cold chain and excursions**

Store at +2C to +8C in the original packaging to protect from light. Do not freeze, and discard any vaccine that has been frozen.

COLD CHAIN EXCURSION: any stock exposed to conditions outside +2C to +8C must be QUARANTINED and risk assessed in accordance with UKHSA Vaccine Incident Guidance before any further use.

Do not administer excursion stock until that assessment is complete and has been recorded.

Where the product SPC gives specific stability data for a temporary excursion, that data governs; otherwise quarantine and assess.

**Disposal**

Dispose of used syringes, needles, vials and any unused or reconstituted vaccine in a UN-approved puncture-resistant sharps container.

Follow local authority requirements and Health Technical Memorandum 07-01, Safe management of healthcare waste.

Never re-sheath a needle.

**Records: traceability**

RECORD THE VACCINE NAME AND BATCH NUMBER for every dose administered. This is a requirement for any biological product and it is what makes a recall actionable.

Record also the expiry date, the dose, the route, the anatomical site, and where more than one vaccine was given at the same visit, the site of each.

Record the name and registration number of the person administering, and that the vaccine was given under this PGD.

### **Consent in children and young people**

Where the individual is UNDER 16, valid consent must come from a person with PARENTAL RESPONSIBILITY, or from the young person where you assess them as GILICK COMPETENT.

Record which of the two applied. Where consent was given by a person with parental responsibility, record their name and relationship to the patient. Where the young person consented on the basis of Gillick competence, record the basis of that assessment.

A parent accompanying a child does not automatically hold parental responsibility. Ask.

Users must be competent in assessing consent in children and young people before working under this PGD.

## **Version and change record**

Version: v002

Issued: 9 September 2026

Supersedes: the previous signed version of this document.

Change: the vaccine safety requirements above were added following an estate-wide sweep of every vaccination PGD, which found that the signed documents were missing, between them, any requirement for adrenaline to be available, any anaphylaxis provision, a stated observation period, a cold chain excursion procedure, a sharps disposal provision, a batch number requirement, and any wording on consent in children and young people. The specific items added to THIS document are: Anaphylaxis: what must be in place before you vaccinate, Observation after vaccination, Cold chain and excursions, Disposal, Records: traceability, Consent in children and young people.

Authorised by Nitin Shori, Medical Director (GMC 6047293) and Chris Pilkington, Head Pharmacist (GPhC 2046322), on 9 September 2026. Signatures applied digitally on their joint instruction, which is the established process for Get Real Health PGDs.