

This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.

Patient Group Direction
for the administration of vaccines against Seasonal Influenza

Summary of NICE / NICE CKS Guidelines for Influenza

Overview

- **Influenza** is an **acute viral respiratory illness**, typically seasonal (autumn–winter in the UK).
- Caused by **influenza A or B viruses**, with variable severity.
- Most people recover in 1–2 weeks, but **complications** (e.g. pneumonia) can occur, especially in **high-risk groups**.

Symptoms

- Sudden onset of:
 - High fever
 - Headache
 - Myalgia (muscle aches)
 - Sore throat
 - Cough
 - Malaise and fatigue
- Often more severe than common cold.

Diagnosis

- Usually **clinical** during flu season.
- Laboratory confirmation (e.g. PCR) is used in hospitalised or high-risk cases.
- Consider differential diagnoses: COVID-19, bacterial pneumonia, RSV, etc.

At-Risk Groups (Higher Risk of Complications)

- Age ≥ 65
- **Pregnant women**
- **Children < 5 years**
- Chronic conditions (e.g. heart, lung, kidney, liver, diabetes)
- Immunocompromised individuals
- Residents in care homes
- People with learning disabilities or obesity (BMI ≥ 40)

Management

1. Symptomatic Treatment (Most Cases)

- Rest, fluids, paracetamol or ibuprofen for fever/pain
- Avoid unnecessary antibiotics
- Advise staying home until symptoms improve (usually 5–7 days)

2. Antiviral Treatment

Offer **antivirals (oseltamivir or zanamivir) within 48 hours of symptom onset** (or 36 hours for zanamivir in some cases) **if the patient is:**

- In a **clinical risk group** or severely ill AND
- Influenza is circulating in the community (confirmed by UKHSA)

Hospital patients may be treated beyond the 48-hour window if clinically indicated.

Antiviral	Route	Dose	Notes
Oseltamivir	Oral	75 mg BD for 5 days	Preferred in most cases
Zanamivir	Inhaled	10 mg BD for 5 days	Use if oseltamivir unsuitable or resistant virus suspected

Complications to Watch For

- Secondary bacterial infections (e.g. pneumonia, otitis media)
- Exacerbation of asthma or COPD
- Dehydration
- Hospitalisation (esp. elderly, frail)

Vaccination (Prevention)

Annual influenza vaccination is recommended for:

- All aged **65 and over**
- **Pregnant women**
- Children aged **6 months to 17 years** (nasal spray preferred)
- **Clinical risk groups**
- **Healthcare and social care workers**
- **Carers**, household contacts of immunocompromised

Vaccines used include:

- **Inactivated quadrivalent vaccines (QIV)**
 - **Live attenuated intranasal vaccine (LAIV)** for children
 - **Adjuvanted or high-dose vaccines** for older adults
-

When to Refer or Admit

- Suspected complications (e.g. pneumonia, sepsis)
 - Severe symptoms in high-risk patients
 - Rapid deterioration
 - Poor oral intake or dehydration
-

Infection Control

- Advise hand hygiene, respiratory etiquette
- Stay home until fever has settled and well enough to resume activities
- Avoid contact with high-risk individuals during acute illness

Patient Group Direction

for the supply/administration of



aTIV (Fluad)

for the treatment to

Seasonal Influenza

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must previously have used PGD's to supply medication ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	aTIV (Fluad) is indicated for the active immunisation of individuals aged 65 years and over against influenza caused by the three strains contained in the vaccine, in accordance with national influenza immunisation programmes.
Inclusion criteria	- Individuals aged 65 years and over eligible for influenza vaccination under the national programme. - Informed consent obtained. - No contraindications to aTIV.
Exclusion criteria	- Known hypersensitivity to the vaccine, its components (including egg protein), or previous severe reaction to influenza vaccines. - Acute febrile illness (postpone vaccination until recovery). - Bleeding disorders unless medically assessed as safe for IM injection. - Severe allergic reaction to any previous dose of influenza vaccine.
Cautions	- Observe patients for 15 minutes post-vaccination. - Record batch number, site of administration, and expiry date. - Advise on expected local/systemic reactions and when to seek help. - Use caution in patients with bleeding disorders or anticoagulation therapy.
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate

Details of the medicine

Name, form and strength of medicine	aTIV (Fluad)
Legal category	POM
Storage	Store in a refrigerator (2°C–8°C). Do not freeze. Protect from light.
Route/method of administration	Intramuscular injection in the deltoid muscle.
Dose and frequency	0.5 mL as a single dose during influenza season, annually.
Quantity to be administered and/or supplied	One 0.5 mL dose per eligible individual.
Maximum or minimum treatment period	One dose annually. Revaccination required each influenza season.
Adverse effects	Pain, redness, swelling at injection site; headache, fatigue, myalgia, fever. Rare: hypersensitivity reactions. Refer to SmPC.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: ● that valid informed consent was given

	● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of supply dose, form and route quantity supplied ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)
--	--

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell.

Key references

● NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 https://www.nice.org.uk/guidance/mg2 ● NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 https://www.nice.org.uk/guidance/mpg2/resources ● NICE CKS Guidance ● British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines BNF (British National Formulary) NICE ● Summary of Product Characteristics for drug
--

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature

Valid from date		Expiry date	
1/10/2025		1/6/2026	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		25/8/2025
Chris Pilkington	Pharmacist GPhC: 2046322		25/8/2025

Change history

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025

Patient Group Direction

for the supply/administration of

TIVc (Flucelvax)

for the treatment to

Seasonal Influenza

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must previously have used PGD's to supply medication ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	TIVc (Flucelvax) is indicated for the active immunisation of individuals aged 18 years and over against influenza caused by the three strains contained in the vaccine, in accordance with national influenza immunisation programmes.
Inclusion criteria	- Individuals aged 18 years and over eligible for influenza vaccination under the national programme. - Informed consent obtained. - No contraindications to TIVc.
Exclusion criteria	- Known hypersensitivity to the vaccine, its components, or previous severe reaction to influenza vaccines. - Acute febrile illness (postpone vaccination until recovery). - Bleeding disorders unless medically assessed as safe for IM injection. - Severe allergic reaction to any previous dose of influenza vaccine.
Cautions	- Observe patients for 15 minutes post-vaccination. - Record batch number, site of administration, and expiry date. - Advise on expected local/systemic reactions and when to seek help. - Use caution in patients with bleeding disorders or anticoagulation therapy.
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate

Details of the medicine

Name, form and strength of medicine	TIVc (Flucelvax)
Legal category	POM
Storage	Store in a refrigerator (2°C–8°C). Do not freeze. Protect from light.
Route/method of administration	Intramuscular injection in the deltoid muscle.
Dose and frequency	0.5 mL as a single dose during influenza season, annually.
Quantity to be administered and/or supplied	One 0.5 mL dose per eligible individual.
Maximum or minimum treatment period	One dose annually. Revaccination required each influenza season.
Adverse effects	Pain, redness, swelling at injection site; headache, fatigue, myalgia, fever. Rare: hypersensitivity reactions. Refer to SmPC.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: ● that valid informed consent was given

	● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of supply dose, form and route quantity supplied ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)
--	--

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell.

Key references

● NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 https://www.nice.org.uk/guidance/mg2 ● NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 https://www.nice.org.uk/guidance/mpg2/resources ● NICE CKS Guidance ● British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines BNF (British National Formulary) NICE ● Summary of Product Characteristics for drug
--

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature

Valid from date		Expiry date	
1/10/2025		1/6/2026	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		25/8/2025
Chris Pilkington	Pharmacist GPhC: 2046322		25/8/2025

Change history

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025

Patient Group Direction

for the supply/administration of

TIVe (Vaxigrip)

for the treatment to

Seasonal Influenza

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must previously have used PGD's to supply medication ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	TIVe (Vaxigrip) is indicated for the active immunisation of individuals aged 6 months and over against influenza caused by the three strains contained in the vaccine, in accordance with national influenza immunisation programmes.
Inclusion criteria	-Adults and children over 6 months old. (The vaccine to be administered must be licensed for the age of the patient. Follow the SPC for that product.) -Where valid consent has been obtained. -Where the patient requires immunisation but does not qualify for a free NHS vaccination. -Where the patient qualifies for a free NHS vaccination but prefers to have the vaccine privately. -No contraindications to influenza vaccination.
Exclusion criteria	- Known hypersensitivity to the vaccine, its components (including egg protein), or previous severe reaction to influenza vaccines. - Acute febrile illness (postpone vaccination until recovery). - Bleeding disorders unless medically assessed as safe for IM injection. - Severe allergic reaction to any previous dose of influenza vaccine.
Cautions	- Use caution in patients with bleeding disorders or anticoagulation therapy.
Actions if patient is excluded or declines treatment	Explain to the patient why the vaccine cannot be given Advise about the protective effects of the vaccine, the risks of infection and potential complications Advise on alternative options/treatments/prevention Document any advice given and the decision reached. Inform or refer to the GP as appropriate It may be appropriate to seek advice from the local Screening and Immunisation Team, local Health Protection Team or the individual's clinician as required

Details of the medicine

Name, form and strength of medicine	TIVe Sanofi Vaxigrip - split virion, egg cultured, inactivated
Legal category	POM
Storage	Store in a refrigerator (2°C–8°C). Do not freeze. Protect from light.
Route/method of administration	Intramuscular injection in the deltoid muscle.
Dose and frequency	0.5 mL as a single dose during influenza season, annually.
Quantity to be administered and/or supplied	One 0.5 mL dose per eligible individual.
Maximum or minimum treatment period	One dose annually. Revaccination required each influenza season.
Adverse effects	Pain, redness, swelling at injection site; headache, fatigue, myalgia, fever. Rare: hypersensitivity reactions. Refer to SmPC.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: ● Record batch number, site of administration, and expiry date. ● that valid informed consent was given ● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of supply dose, form and route quantity supplied ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell. Advise on expected local/systemic reactions and when to seek help. Individuals should be advised about possible adverse reactions to the vaccine and reassured that the inactivated vaccine cannot cause influenza Advise that the vaccine will not provide protection for about 14 days, and it will not protect against other respiratory viruses that often circulate during the flu season Immunosuppressed individuals should be advised that they may not make a full immune response to the vaccine. For immunosuppressed patients, consideration should also be given to vaccinating household contacts and other close contacts Patients should be warned that no vaccine is 100% effective. If systemic upset, acute illness or any other clinical issues develop which cause concern, the patient should be referred urgently for medical assessment.

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and

administration of medicines [BNF \(British National Formulary\)](#) | [NICE](#)

● Summary of Product Characteristics for drug

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature
Valid from date		Expiry date	
1/10/25		1/5/26	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		25/9/2025
Chris Pilkington	Pharmacist GPhC: 2046322		25/9/2025

Change history

Version number	Change details	Date
001	Development & issue of new PGD	25 th Sept 2025

Patient Group Direction

for the supply/administration of

IIVc (CSL Seqirus)

for vaccination against

Influenza

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must previously have used PGD's to supply medication ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	IIVc (CSL Seqirus) is indicated for active immunisation against influenza in adults and children over 6 months old to reduce the risk of influenza-related illness during seasonal outbreaks.
Inclusion criteria	-Adults and children over 6 months old. (The vaccine to be administered must be licensed for the age of the patient. Follow the SPC for that product.) -Where valid consent has been obtained. -Where the patient requires immunisation but does not qualify for a free NHS vaccination. -Where the patient qualifies for a free NHS vaccination but prefers to have the vaccine privately. -No contraindications to influenza vaccination.
Exclusion criteria	- Known hypersensitivity to any component of the vaccine - History of severe allergic reaction to previous influenza vaccination. - Acute febrile illness or systemic infection. - History of Guillain-Barré Syndrome following previous influenza vaccine.
Cautions	- Defer in cases of acute febrile illness until recovery. - Observe patient for immediate hypersensitivity reactions post-vaccination. - Advise patient that protection is generally achieved within 10–14 days. - Administer annually before or during influenza season.
Actions if patient is excluded or declines treatment	-Explain to the patient why the vaccine cannot be given -Advise about the protective effects of the vaccine, the risks of infection and potential complications -Advise on alternative options/treatments/prevention -Document any advice given and the decision reached. Inform or refer to the GP as appropriate -It may be appropriate to seek advice from the local Screening and Immunisation Team, local Health Protection Team or the individual's clinician as required

Details of the medicine

Name, form and strength of medicine	IIVc (CSL Seqirus)- cell cultured, inactivated influenza vaccine
Legal category	POM
Storage	Store in a refrigerator between 2°C–8°C. Do not freeze. Protect from light.
Route/method of administration	Intramuscular injection, preferably in the deltoid muscle.

Dose and frequency	Prevention: Single 0.5 mL dose administered during influenza season
Quantity to be administered and/or supplied	0.5 mL
Maximum or minimum treatment period	Administer annually before or during influenza season.
Adverse effects	Injection site pain, fatigue, headache, myalgia, fever. Rare: allergic reactions. Refer to SmPC for full details.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and advise the traveller to consult a healthcare professional if symptoms worsen.
Records to be kept	Record the following information: ● that valid informed consent was given ● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of supply dose, form and route quantity supplied ● Record batch number, site of administration and expiry date ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	-To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell. -Individuals should be advised about possible adverse reactions to the vaccine and reassured that the inactivated vaccine cannot cause influenza -Advise that the vaccine will not provide protection for about 14 days, and it will not protect against other respiratory viruses that often circulate during the flu season -Immunosuppressed individuals should be advised that they may not make a full immune response to the vaccine. For immunosuppressed patients, consideration

	should also be given to vaccinating household contacts and other close contacts -Patients should be warned that no vaccine is 100% effective. If systemic upset, acute illness or any other clinical issues develop which cause concern, the patient should be referred urgently for medical assessment.
--	--

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines [BNF \(British National Formulary\) | NICE](#)
- Summary of Product Characteristics for drug

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature
Valid from date		Expiry date	

1/10/2025	1/6/2026
-----------	----------

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or vaccine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		25/8/2025
Chris Pilkington	Pharmacist GPhC: 2046322		25/8/2025

Change history

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025

Patient Group Direction

for the supply/administration of

aIV (CSL Seqirus)

for vaccination against

Influenza (Ages 65+)

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must previously have used PGD's to supply medication ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	aIV (CSL Seqirus) is indicated for active immunisation against influenza in adults aged 65 years and over to reduce the risk of influenza-related illness during seasonal outbreaks.
Inclusion criteria	- Adults aged 65 years and over. - Eligible for NHS or private seasonal influenza vaccination. - Informed consent obtained. - No contraindications to influenza vaccination.
Exclusion criteria	- Known hypersensitivity to any component of the vaccine, including egg protein. - History of severe allergic reaction to previous influenza vaccination. - Acute febrile illness or systemic infection. - History of Guillain-Barré Syndrome following previous influenza vaccine.
Cautions	- Defer in cases of acute febrile illness until recovery. - Observe patient for immediate hypersensitivity reactions post-vaccination. - Advise patient that protection is generally achieved within 10–14 days. - Administer annually before or during influenza season.
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate

Details of the medicine

Name, form and strength of medicine	aIV (CSL Seqirus)
Legal category	POM
Storage	Store in a refrigerator between 2°C–8°C. Do not freeze. Protect from light.
Route/method of administration	Intramuscular injection, preferably in the deltoid muscle.
Dose and frequency	Prevention: Single 0.5 mL dose administered annually during influenza season
Quantity to be administered and/or supplied	0.5 mL
Maximum or minimum treatment period	Administer annually before or during influenza season.
Adverse effects	Injection site pain, fatigue, headache, myalgia, fever. Rare: allergic reactions. Refer to SmPC for full details.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and advise the traveller to consult a healthcare professional if symptoms worsen.
Records to be kept	Record the following information:

	● that valid informed consent was given ● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of supply dose, form and route quantity supplied ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)
--	---

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell.

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines [BNF \(British National Formulary\) | NICE](#)
- Summary of Product Characteristics for drug

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature

Valid from date		Expiry date	
1/10/2025		1/6/2026	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or vaccine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		25/8/2025
Chris Pilkington	Pharmacist GPhC: 2046322		25/8/2025

Change history

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025

Patient Group Direction

for the supply/administration of

IIVr (Supemtek, Sanofi)

for vaccination against Influenza

Influenza (18+)

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must previously have used PGD's to supply medication ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	IIVr (Supemtek, Sanofi) is indicated as a second-line option for active immunisation against influenza in adults aged 18 years and over when first-line vaccines are unavailable or unsuitable, to reduce the risk of influenza-related illness during seasonal outbreaks.
Inclusion criteria	- Adults aged 18 years and over. - Eligible for NHS or private seasonal influenza vaccination. - Informed consent obtained. - First-line influenza vaccine not available or contraindicated.
Exclusion criteria	- Known hypersensitivity to any component of the vaccine. - Severe allergic reaction to a previous influenza vaccine. - Acute febrile illness or systemic infection. - History of Guillain-Barré Syndrome following an influenza vaccine.
Cautions	- Defer in cases of acute febrile illness until recovery. - Observe patient for immediate hypersensitivity reactions post-vaccination. - Advise patient that protection is generally achieved within 10–14 days. - Administer annually before or during influenza season.
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate

Details of the medicine

Name, form and strength of medicine	IIVr (Supemtek, Sanofi)
Legal category	POM
Storage	Store in a refrigerator between 2°C–8°C. Do not freeze. Protect from light.
Route/method of administration	Intramuscular injection, preferably in the deltoid muscle.
Dose and frequency	Prevention: Single 0.5 mL dose administered annually during influenza season
Quantity to be administered and/or supplied	0.5 mL
Maximum or minimum treatment period	Administer annually before or during influenza season.
Adverse effects	Injection site pain, fatigue, headache, myalgia, fever. Rare: allergic reactions. Refer to SmPC for full details.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and advise the traveller to consult a healthcare professional if symptoms worsen.
Records to be kept	Record the following information:

	● that valid informed consent was given ● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of supply dose, form and route quantity supplied ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)
--	---

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell.

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mpg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines [BNF \(British National Formulary\) | NICE](#)
- Summary of Product Characteristics for drug

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature

Valid from date		Expiry date	
1/10/2025		1/6/2026	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or vaccine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		25/8/2025
Chris Pilkington	Pharmacist GPhC: 2046322		25/8/2025

Change history

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025