

This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.

Patient Group Direction
for the administration of Engerix B for vaccination against Hepatitis B

Patient Group Direction, version 006, issued 14 September 2026. Supersedes Version 005, 11 September 2026.

Owner and authorising organisation: Get Real Health Limited, company number 12744898, registered office Zenvite Health Unit 23, St. Asaph Business Park, St. Asaph, Denbighshire, LL17 0LJ. This Patient Group Direction is authorised on behalf of Get Real Health Limited by its Medical Director and its Head Pharmacist, whose signatures for this version are on the authorisation page at the end of the document. It applies to the registered healthcare professionals named under it by a pharmacy or clinic that has adopted it through a Get Real Health service agreement, at the premises stated in that adoption, and to no other organisation.

Summary of the governing guidance for hepatitis B vaccination: Green Book chapter 18 (Hepatitis B) and the Engerix B, HBvaxPRO and Fendrix SmPCs. NICE CKS 'Hepatitis B' covers the infection, not occupational vaccination.

Overview

- **Hepatitis B** is a **vaccine-preventable viral infection** of the liver, transmitted through **blood and body fluids**.
- The **Hepatitis B vaccine** is a **recombinant subunit vaccine** (non-live) that provides effective long-term protection.

Routine and Targeted Vaccination

1. Universal Childhood Programme

- Since August 2017, **Hepatitis B is included in the 6-in-1 (DTaP/IPV/Hib/HepB) vaccine.**
- Schedule: **8, 12, and 16 weeks of age**, as part of routine UK childhood immunisation.

2. Targeted Vaccination for At-Risk Groups

Recommended for individuals at higher risk, including:

- **Babies born to Hepatitis B positive mothers**
- **Healthcare workers and care workers** at risk of exposure
- **People who inject drugs** or have high-risk sexual behaviour
- **Household/sexual contacts of HBV carriers**
- Individuals with **chronic kidney disease** (especially on dialysis)
- **Prisoners**
- **Travellers to high-prevalence countries**
- People with **liver disease, HIV, or immunosuppression**

Vaccination Schedules

Standard Schedule (Adults and children ≥ 1 year):

- **0, 1, and 6 months** (3 doses)
- A **booster may be given** at 5 years for ongoing risk, though long-term protection is expected.

Accelerated Schedule (For rapid protection – e.g. travel or post-exposure):

- **0, 1, and 2 months** with a booster at 12 months

Ultra-Rapid Schedule (e.g. travellers):

- **Day 0, 7, and 21**, with a booster at 12 months
- **The ultra-rapid (day 0, 7 and 21) schedule is not authorised under this PGD; only the standard and accelerated schedules in the dose rows are.**

Use **monovalent Hepatitis B vaccine** for adults or for birth doses in infants born to infected mothers.

Post-Vaccination Serology

- **Test for anti-HBs** (antibody to Hepatitis B surface antigen) 1–4 months after final dose **if high-risk**, e.g.:

- Healthcare workers
 - Babies born to infected mothers
 - Immunocompromised individuals
 - **Anti-HBs ≥ 10 mIU/mL** = protective
 - If inadequate response: offer further doses (up to 6 total), then reassess.
-

Contraindications and Precautions

- Severe allergic reaction to a previous dose or any component.
 - Acute severe illness (defer until recovery).
 - Safe in pregnancy and breastfeeding.
-

Side Effects

- Usually mild: local pain, redness, low-grade fever.
 - Rare: hypersensitivity reactions.
-

Documentation

- Record **brand, dose, batch number, route, site, and date**.
- Report adverse events via the **Yellow Card Scheme**.

Patient Group Direction

for the supply/administration of

Engerix B

for vaccination against

Hepatitis B

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must have completed training in working under Patient Group Directions, for example the CPPE PGD e-learning and e-assessment or an equivalent, and have been assessed as competent to work under a PGD before first working under this one; previous experience of working under a PGD is not required ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	Engerix B is indicated for the active immunisation of individuals at risk of exposure to hepatitis B virus. Under this PGD: individuals aged 16 years and over at occupational, travel or lifestyle risk, including healthcare workers. Infants born to infected mothers, and anyone under 16, are outside this PGD.
Inclusion criteria	- Individuals aged 16 years and over at increased risk of hepatitis B infection (occupational, travel or lifestyle risk). Children under 16 are not vaccinated under this PGD; refer. - Eligible under national immunisation or occupational health guidance. - Informed consent obtained. - No contraindications to the vaccine.
Exclusion criteria	- Known hypersensitivity to the active substance or any excipients. - Known hypersensitivity to yeast (both vaccines are produced in yeast and may contain traces of yeast protein). - Previous allergic reaction to any hepatitis B vaccine. - Aged under 16 years. - Acute severe febrile illness (postpone until recovered).
Cautions	- Use with caution in individuals with bleeding disorders or on anticoagulants. - Observe every patient for 15 minutes after vaccination, seated, and record that the observation period was completed. - Ensure adherence to the complete vaccination schedule. - Provide follow-up advice and written patient information leaflet (PIL).
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate

Details of the medicine

Name, form and strength of medicine	Engerix B
Legal category	POM
Storage	Store in a refrigerator (2°C–8°C). Do not freeze. Protect from light.
Route/method of administration	Intramuscular injection, usually in the deltoid muscle.
Dose and frequency	Adults and adolescents (16+): 20 micrograms (1 mL) per dose. Standard schedule: 0, 1, and 6 months. Accelerated schedule: 0, 1, 2, and 12 months.
Quantity to be administered and/or supplied	1 mL per dose depending on the schedule.
Maximum or minimum treatment period	A course of 3 doses over 6 months (standard schedule) or 4 doses over

	12 months (accelerated schedule). Booster doses are not covered by this PGD.
Adverse effects	Injection site pain, redness, swelling, fatigue, headache, fever, gastrointestinal symptoms. Rare: allergic reactions. Refer to SmPC.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: ● that valid informed consent was given ● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of administration, dose, form and route, quantity administered ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and the actions taken, and that the vaccine was administered under this PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. Record also the vaccine brand, batch number, expiry date, dose number, anatomical site and the date the next dose is due. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	Attend for any further dose of the course on the date given. Seek medical advice for a severe or persistent injection site reaction, a fever that does not settle within 48 hours, or any sign of an allergic reaction. Call 999 for difficulty breathing, swelling of the face or throat, or collapse. Post-vaccination serology 1 to 4 months after the final dose where occupational health requires it.

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017
<https://www.nice.org.uk/guidance/mg2>

- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/ MPG2/resources>
- UKHSA, Immunisation Against Infectious Disease (the Green Book), chapter 18 Hepatitis B; Summaries of Product Characteristics for the vaccine administered, current version
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines [BNF \(British National Formulary\) | NICE](#)
- Summary of Product Characteristics

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature
Valid from date		Expiry date	
14/9/26		31/7/27	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service. To be completed by the adopting pharmacy. These are the adopting organisation's own signatories and must not be Get Real Health staff. To be completed by the adopting pharmacy. These are the adopting organisation's own signatories and must not be Get Real Health staff. To be completed by the adopting pharmacy. These are the adopting organisation's own signatories and must not be Get Real Health staff.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date

Name of professional group
signatory

Job title & name of
organisation

Signature

Date

Change history to version 001

Later versions are recorded in the Version and change record at the end of this document.

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025

Patient Group Direction

for the supply/administration of



HBvaxPRO

for vaccination against

Hepatitis B

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must have completed training in working under Patient Group Directions, for example the CPPE PGD e-learning and e-assessment or an equivalent, and have been assessed as competent to work under a PGD before first working under this one; previous experience of working under a PGD is not required ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	HBVax Pro is indicated for the active immunisation of individuals at risk of exposure to hepatitis B virus. Under this PGD: individuals aged 16 years and over at occupational, travel or lifestyle risk, including healthcare workers. Infants born to infected mothers, and anyone under 16, are outside this PGD.
Inclusion criteria	- Individuals aged 16 years and over at increased risk of hepatitis B infection (occupational, travel or lifestyle risk). Children under 16 are not vaccinated under this PGD; refer. - Eligible under national immunisation or occupational health guidance. - Informed consent obtained. - No contraindications to the vaccine.
Exclusion criteria	- Known hypersensitivity to the active substance or any excipients. - Known hypersensitivity to yeast (both vaccines are produced in yeast and may contain traces of yeast protein). - Previous allergic reaction to any hepatitis B vaccine. - Aged under 16 years. - Acute severe febrile illness (postpone until recovered).
Cautions	- Use with caution in individuals with bleeding disorders or on anticoagulants. - Observe every patient for 15 minutes after vaccination, seated, and record that the observation period was completed. - Ensure adherence to the complete vaccination schedule. - Provide follow-up advice and written patient information leaflet (PIL).
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate

Details of the medicine

Name, form and strength of medicine	HBVax Pro
Legal category	POM
Storage	Store in a refrigerator (2°C–8°C). Do not freeze. Protect from light.
Route/method of administration	Intramuscular injection, usually in the deltoid muscle.
Dose and frequency	HBVAXPRO 10 micrograms/1 mL: one dose per injection, individuals 16 years and over. Standard schedule: 0, 1, and 6 months. Accelerated schedule: 0, 1, 2, and 12 months. Pre-dialysis and dialysis patients (40 microgram presentation) are not covered by this PGD; refer.
Quantity to be	1 mL (10 micrograms) per dose.

administered and/or supplied	
Maximum or minimum treatment period	A course of 3 doses over 6 months (standard schedule) or 4 doses over 12 months (accelerated schedule). Booster doses are not covered by this PGD.
Adverse effects	Injection site reactions (pain, redness, swelling), fever, fatigue, headache, nausea. Rare: hypersensitivity reactions. Refer to SmPC.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: ● that valid informed consent was given ● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of administration, dose, form and route, quantity administered ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and the actions taken, and that the vaccine was administered under this PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. Record also the vaccine brand, batch number, expiry date, dose number, anatomical site and the date the next dose is due. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	Attend for any further dose of the course on the date given. Seek medical advice for a severe or persistent injection site reaction, a fever that does not settle within 48 hours, or any sign of an allergic reaction. Call 999 for difficulty breathing, swelling of the face or throat, or collapse. Post-vaccination serology 1 to 4 months after the final dose where occupational health requires it.

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- UKHSA, Immunisation Against Infectious Disease (the Green Book), chapter 18 Hepatitis B; Summaries of Product Characteristics for the vaccine administered, current version
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines [BNF \(British National Formulary\) | NICE](#)
- Summary of Product Characteristics

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature
Valid from date		Expiry date	
14/9/26		31/7/27	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service. To be completed by the adopting pharmacy. These are the adopting organisation's own signatories and must not be Get Real Health staff. To be completed by the adopting pharmacy.

These are the adopting organisation's own signatories and must not be Get Real Health staff. To be completed by the adopting pharmacy. These are the adopting organisation's own signatories and must not be Get Real Health staff.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date

Name of professional group signatory

Job title & name of organisation

Signature

Date

Change history to version 001

Later versions are recorded in the Version and change record at the end of this document.

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025



Vaccine safety requirements

The requirements on the following pages form part of this Patient Group Direction and must be met before any vaccine is administered under it.

Anaphylaxis: what must be in place before you vaccinate

ADRENALINE (EPINEPHRINE) 1 IN 1,000 INJECTION must be immediately available in the room where vaccination takes place, in date, together with a telephone.

A written anaphylaxis protocol consistent with current Resuscitation Council UK guidance must be available, and every person administering must be trained in the recognition and immediate management of anaphylaxis and hold current basic life support training.

Be able to distinguish a faint and a panic attack from anaphylaxis. Green Book chapter 8 sets out the clinical features of each. A faint after an injection is common; anaphylaxis is very rare.

Report any suspected anaphylaxis via the Yellow Card Scheme even where the diagnosis is uncertain.

Cold chain and excursions

Store at +2C to +8C in the original packaging to protect from light. Do not freeze, and discard any vaccine that has been frozen.

COLD CHAIN EXCURSION: any stock exposed to conditions outside +2C to +8C must be QUARANTINED and risk assessed in accordance with UKHSA Vaccine Incident Guidance before any further use.

Do not administer excursion stock until that assessment is complete and has been recorded.

Where the product SPC gives specific stability data for a temporary excursion, that data governs; otherwise quarantine and assess.



Disposal

Dispose of used syringes, needles, vials and any unused or reconstituted vaccine in a UN-approved puncture-resistant sharps container.

Follow local authority requirements and Health Technical Memorandum 07-01, Safe management of healthcare waste.

Never re-sheath a needle.

Consent in children and young people

This PGD excludes anyone under 16, so consent from a person with parental responsibility or on the basis of Gillick competence does not arise. A 16 or 17 year old may consent for themselves; assess capacity as for an adult.

Version and change record

Version: v006

Issued: 14 September 2026

Supersedes: Version 005, 11 September 2026

Valid from: 14 September 2026. Expiry: 31 July 2027. These dates govern this version and supersede any earlier date printed elsewhere in this document.

Changes in this version:

Identity and training, applied to every Get Real Health PGD on 14 September 2026 after two adopting pharmacies wrote: the Get Real Health logo is in the page header; the organisation that owns and authorises this direction, Get Real Health Limited, is named under the strapline with its company number and registered office; the requirement that practitioners must previously have used a PGD is replaced by a requirement for PGD training and a competence assessment, so that a newly registered pharmacist or technician who has completed the training may work under it; and the earlier change history table is headed with the versions it covers.

Previous version records

Version: v005

Issued: 11 September 2026

Supersedes: Version 004, 11 September 2026

Valid from: 11 September 2026. Expiry: 31 July 2027. These dates govern this version and supersede any earlier date printed elsewhere in this document.

Changes in this version:

The guidance summary now states that the ultra-rapid day 0, 7 and 21 schedule is not authorised under this PGD.

Both indications now say who this PGD covers (16 years and over at occupational, travel or lifestyle risk) and that infants born to infected mothers and anyone under 16 are outside it.

Known hypersensitivity to yeast, which both SmPCs name, is now an exclusion in both arms.

Both cautions rows now require the 15 minute seated observation to be recorded, as the other vaccine PGDs do.

The maximum treatment period rows now state the course (3 doses over 6 months or 4 doses over 12 months) instead of 'typically 6 to 12 months', and that boosters are not covered.

The records rows have their punctuation restored, and the follow-up advice rows, which carried the treatment template about 3 to 4 weeks, now give vaccination follow-up advice including post-vaccination serology where occupational health requires it.

The consent in children and young people block now states that it does not apply, because this PGD excludes anyone under 16.

Wording corrections from the tool alignment and adversarial review of the consultation tools, 11 September 2026. Each correction resolves a contradiction inside the document or a plain error, taking the safer reading; no indication is widened, no dose raised and no exclusion removed. Questions that need a clinical decision are recorded separately and are not changed here.

Previous version records

Version: v004

Issued: 11 September 2026

Supersedes: Version 003, 10 September 2026

Valid from: 11 September 2026. Expiry: 31 July 2027. These dates govern this version and supersede any earlier date printed elsewhere in this document.

Changes in this version:

The guidance summary and references are attributed to Green Book chapter 18 and the SmPCs rather than to NICE CKS, which does not cover occupational hepatitis B vaccination.

Structural clean-up applied to every live Get Real Health PGD on 11 September 2026, following the adversarial review of 10 September: the adopting pharmacy's authorisation block is blank, for the pharmacy's own signatories; one signed authorisation page for this version replaces the earlier signature blocks; every validity cell states this version's dates (valid from 11 September 2026, expiry 31 July 2027); narration about earlier versions, the consultation tool and individual customers is removed from the body of the document; the change records of earlier reissues are carried below as previous version records. No clinical criterion changes unless listed.

Previous version records

Version: v002

Issued: 9 September 2026

Supersedes: the previous signed version of this document.

Change: the vaccine safety requirements above were added following an estate-wide sweep of every vaccination PGD, which found that the signed documents were missing, between them, any requirement for adrenaline to be available, any anaphylaxis provision, a stated observation period, a cold chain excursion procedure, a sharps disposal provision, a batch number requirement, and any wording on consent in children and young people. The specific items added to THIS document are: Anaphylaxis: what must be in place before you vaccinate, Cold chain and excursions, Disposal, Consent in children and young people.

Authorised by Nitin Shori, Medical Director (GMC 6047293) and Chris Pilkington, Head Pharmacist (GPhC 2046322), on 9 September 2026. Signatures applied digitally on their joint instruction, which is the established process for Get Real Health PGDs.

Version: v003

Issued: 10 September 2026

Supersedes: Version 002, 9 September 2026

Valid from: 10 September 2026. Expiry: 31 July 2027. These dates govern this version and supersede any earlier date printed elsewhere in this document.

Changes in this version:

Validity: this version is valid from 10 September 2026 and expires on 31 July 2027. The signed master previously stated no period of validity, or an expiry of 31 October 2026; those are superseded. No period of validity was stated.


Inclusion, both arms: 'individuals of any age' with only the adult dose authorised meant a child received double the licensed dose or no valid dose. Restricted to 16 years and over; under 16 refers.

HBVAXPRO: the dose row named a 20 microgram strength that does not exist and a 0.5 mL volume that belongs to the paediatric product. Now HBVAXPRO 10 micrograms/1 mL for 16 years and over; dialysis patients (40 microgram product) refer.

Records: brand, batch number, expiry, dose number, site and next dose date added, as the document's own summary and both SmPCs require.

Signed on behalf of Get Real Health

Version v006, 14 September 2026.

Name	Qualification	Signature	Date
Nitin Shori	Medical Director GMC: 6047293		14 September 2026
Chris Pilkington	Head Pharmacist GPhC: 2046322		14 September 2026

Both authorising signatories reviewed this version together on 14 September 2026 and gave their authorisation for it to be issued. Signatures were applied digitally on their joint instruction, which is the established process for Get Real Health PGDs. This version is not valid without both signatures.