

This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.

Patient Group Direction

for the administration of Pneumovax 23 or Prevenar 13 to vaccinate against Pneumococcal Disease

Patient Group Direction, version 003, issued 10 September 2026. Supersedes Version 002, 9 September 2026.

Summary of NICE / NICE CKS Guidelines for Pneumococcal Vaccination

Overview

- Pneumococcal disease is caused by *Streptococcus pneumoniae*, leading to infections such as **pneumonia**, **septicaemia**, and **meningitis**.
- Vaccination is aimed at **preventing invasive pneumococcal disease (IPD)**, especially in high-risk groups.

Types of Pneumococcal Vaccines Used in the UK

1. **PCV15 or PCV13** (Prevenar 15 / 13):
 - **Conjugate vaccines** — used in infants and selected high-risk groups.
2. **PPSV23** (Pneumovax 23):
 - **Polysaccharide vaccine** — protects against 23 serotypes.
 - Used in adults and older children at risk of IPD.

Routine Childhood Schedule (UK)

- **PCV15 (or PCV13)** is given as:
 - **First dose at 12 weeks**
 - **Booster at 1 year** (as part of routine infant immunisation)

Note: The exact schedule may vary based on updated UKHSA recommendations.

At-Risk Groups (Eligible for Additional Vaccination)

Offer **PPSV23** (or PCV plus PPSV23) to individuals aged **2 years and over** with:

- **Chronic respiratory, heart, kidney, liver or neurological conditions**
- **Diabetes**
- **Immunosuppression** (e.g. HIV, chemotherapy, asplenia)
- **Cochlear implants**
- **CSF leaks**
- **Sickle cell disease or other haemoglobinopathies**

High-risk children under 2 years may receive **additional PCV doses** on a tailored schedule.

Older Adults

- **All adults aged 65 years** are offered a **single dose of PPSV23**, unless already vaccinated.

Vaccination Schedule by Risk Group (Simplified)

Group	Vaccine	Notes
Infants	PCV15/13 at 12 weeks + booster at 1 year	Routine
Adults ≥65	PPSV23 (one-off)	Even if healthy
At-risk ≥2 years	PPSV23	May also need PCV if never received
Severely immunocompromised	PCV followed by PPSV23 after ≥8 weeks	Requires specialist input

Contraindications

- Severe allergic reaction to previous dose or vaccine component.
- Acute febrile illness (defer until recovery).

Side Effects

- Mild: redness, swelling, pain at injection site; low-grade fever

- Rare: allergic reaction
-

Co-Administration

- Can be given with other vaccines (e.g. flu, COVID-19), in a different limb or site.
-

Documentation

- Record **vaccine name, batch number, site, date, and route.**
- Report adverse reactions via the **Yellow Card Scheme.**

Patient Group Direction

for the supply/administration of

Pneumovax 23

for vaccination against

Pneumococcal Disease

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must previously have used PGD's to supply medication ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	Pneumovax 23 is indicated for active immunisation against pneumococcal disease caused by the 23 serotypes included in the vaccine for individuals aged 2 years and over at increased risk, in accordance with national guidance.
Inclusion criteria	- Individuals aged 2 years and over eligible for pneumococcal vaccination under national guidance. - Informed consent obtained from the individual or guardian. - No contraindications to pneumococcal vaccination.
Exclusion criteria	- Known hypersensitivity to Pneumovax 23 or any of its components. - Previous severe allergic reaction to any pneumococcal vaccine. - PPV23 or PCV20 received within the last 5 years, unless the individual has asplenia, splenic dysfunction or chronic kidney disease and 5 years have elapsed. Revaccination is not recommended for any other group. - Acute illness with fever (postpone until recovered).
Cautions	- Use with caution in individuals with bleeding disorders. - Record batch number and site of administration. - Monitor for anaphylaxis for 15 minutes after administration. - Inform patient of possible side effects and when to seek help.
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate

Details of the medicine

Name, form and strength of medicine	Pneumovax 23
Legal category	POM
Storage	Store in a refrigerator (2°C–8°C). Do not freeze. Protect from light.
Route/method of administration	Intramuscular or subcutaneous injection.
Dose and frequency	Single 0.5 mL dose. Revaccination every 5 years ONLY for asplenia, splenic dysfunction or chronic kidney disease (Green Book chapter 25); not recommended for any other group, and never within 3 years of a previous dose.
Quantity to be administered and/or supplied	0.5 mL per dose.
Maximum or minimum treatment period	One lifetime dose in most cases. Repeat only under specific clinical recommendations.
Adverse effects	Injection site pain, redness, swelling, fever, fatigue, headache. Rare: allergic reactions. Refer to SmPC.

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: ● that valid informed consent was given ● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of supply dose, form and route quantity supplied ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell.

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mpg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines [BNF \(British National Formulary\) | NICE](#)
- Summary of Product Characteristics

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature
Valid from date		Expiry date	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date
Chris Pilkington	Head Pharmacist, Get Real Health Limited (GPhC: 2046322)		19/06/2026

Name of professional group signatory	Job title & name of organisation	Signature	Date
Chris Pilkington	Head Pharmacist, Get Real Health Limited (GPhC: 2046322)		19/06/2026

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		25/08/2025
Chris Pilkington	Pharmacist GPhC: 2046322		25/08/2025

Change history

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025

Patient Group Direction

for the supply/administration of

Prevenar 13

for vaccination against

Pneumococcal Disease

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	● Pharmacist registered and practicing with the GPhC ● Pharmacy technician registered and practicing with the GPhC
Training and assessment	● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. ● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines ● All users must previously have used PGD's to supply medication ● Must work in compliance with the SOPs of their own employer ● All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	● All users must be up to date with CPD requirements and appraisal ● All users must be up to date with MHRA and safety alerts with regards to medical products ● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	Prevenar 13 is indicated for active immunisation against pneumococcal disease caused by the 13 serotypes included in the vaccine, for infants, children, and adults in accordance with the national immunisation programme and clinical risk group recommendations.
Inclusion criteria	- Individuals aged 2 years and over who are eligible for pneumococcal vaccination under national guidance and in whom a conjugate vaccine is indicated. - Infant primary immunisation (under 2 years) is NOT given under this PGD. - Informed consent obtained from the individual or guardian. - No contraindications to Prevenar 13.
Exclusion criteria	- Known hypersensitivity to the active substance or any component of the vaccine, including hypersensitivity to diphtheria toxoid (CRM197 carrier protein). - Severe allergic reaction to a previous pneumococcal conjugate vaccine. - Aged under 2 years. - Acute febrile illness (postpone vaccination until recovery).
Cautions	- Observe the patient for 15 minutes post-vaccination. - Use with caution in patients with bleeding disorders. - Record batch number and expiry date in the patient record. - Advise on expected side effects and when to seek medical advice.
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate

Details of the medicine

Name, form and strength of medicine	Prevenar 13
Legal category	POM
Storage	Store in a refrigerator (2°C–8°C). Do not freeze. Protect from light.
Route/method of administration	Intramuscular injection.
Dose and frequency	Single 0.5 mL dose for individuals aged 2 years and over who have not previously received a pneumococcal conjugate vaccine. Where PPV23 is also indicated, give it at least 8 weeks after the conjugate vaccine.
Quantity to be administered and/or supplied	0.5 mL per dose.
Maximum or minimum treatment period	As per national immunisation schedule, including primary and booster doses.
Adverse effects	Irritability, drowsiness, loss of appetite, fever, injection site

	redness/swelling. Rare: hypersensitivity reactions. Refer to SmPC.
--	--

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: ● that valid informed consent was given ● name of individual, address, date of birth and GP ● name of healthcare practitioner ● name and brand of medication ● date of supply dose, form and route quantity supplied ● advice given, including advice given if excluded or declines treatment ● details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: ● Adults aged 18 years and over - keep records for 8 years ● Children aged under 18 years -keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell.

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mpg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines [BNF \(British National Formulary\) | NICE](#)
- Summary of Product Characteristics

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the

responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature
Valid from date		Expiry date	

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date
Chris Pilkington	Head Pharmacist, Get Real Health Limited (GPhC: 2046322)		19/06/2026

Name of professional group signatory	Job title & name of organisation	Signature	Date
Chris Pilkington	Head Pharmacist, Get Real Health Limited (GPhC: 2046322)		19/06/2026

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		25/08/2025
Chris Pilkington	Pharmacist GPhC: 2046322		25/08/2025

Change history

Version number	Change details	Date
001	Development & issue of new PGD	25 th June 2025



Vaccine safety requirements

The requirements on the following pages form part of this Patient Group Direction and must be met before any vaccine is administered under it.

Anaphylaxis: what must be in place before you vaccinate

ADRENALINE (EPINEPHRINE) 1 IN 1,000 INJECTION must be immediately available in the room where vaccination takes place, in date, together with a telephone.

A written anaphylaxis protocol consistent with current Resuscitation Council UK guidance must be available, and every person administering must be trained in the recognition and immediate management of anaphylaxis and hold current basic life support training.

Be able to distinguish a faint and a panic attack from anaphylaxis. Green Book chapter 8 sets out the clinical features of each. A faint after an injection is common; anaphylaxis is very rare.

Report any suspected anaphylaxis via the Yellow Card Scheme even where the diagnosis is uncertain.

Cold chain and excursions

Store at +2C to +8C in the original packaging to protect from light. Do not freeze, and discard any vaccine that has been frozen.

COLD CHAIN EXCURSION: any stock exposed to conditions outside +2C to +8C must be QUARANTINED and risk assessed in accordance with UKHSA Vaccine Incident Guidance before any further use.

Do not administer excursion stock until that assessment is complete and has been recorded.

Where the product SPC gives specific stability data for a temporary excursion, that data governs; otherwise quarantine and assess.



Disposal

Dispose of used syringes, needles, vials and any unused or reconstituted vaccine in a UN-approved puncture-resistant sharps container.

Follow local authority requirements and Health Technical Memorandum 07-01, Safe management of healthcare waste.

Never re-sheath a needle.

Consent in children and young people

Where the individual is UNDER 16, valid consent must come from a person with PARENTAL RESPONSIBILITY, or from the young person where you assess them as GILICK COMPETENT.

Record which of the two applied. Where consent was given by a person with parental responsibility, record their name and relationship to the patient. Where the young person consented on the basis of Gillick competence, record the basis of that assessment.

A parent accompanying a child does not automatically hold parental responsibility. Ask.

Users must be competent in assessing consent in children and young people before working under this PGD.

Version and change record

Version: v002

Issued: 9 September 2026

Supersedes: the previous signed version of this document.

Change: the vaccine safety requirements above were added following an estate-wide sweep of every vaccination PGD, which found that the signed documents were missing, between them, any requirement for adrenaline to be available, any anaphylaxis provision, a stated observation period, a cold chain excursion procedure, a sharps disposal provision, a batch number requirement, and any wording on consent in children and young people. The specific items added to THIS document are: Anaphylaxis: what must be in place before you vaccinate, Cold chain and excursions, Disposal, Consent in children and young people.

Authorised by Nitin Shori, Medical Director (GMC 6047293) and Chris Pilkington, Head Pharmacist (GPhC 2046322), on 9 September 2026. Signatures applied digitally on their joint instruction, which is the established process for Get Real Health PGDs.

Version and change record

Version: v003

Issued: 10 September 2026

Supersedes: Version 002, 9 September 2026

Valid from: 10 September 2026. Expiry: 31 July 2027. These dates govern this version and supersede any earlier date printed elsewhere in this document.

Changes in this version:

Validity: this version is valid from 10 September 2026 and expires on 31 July 2027. The signed master previously stated no period of validity, or an expiry of 31 October 2026; those are superseded. No period of validity was stated.


Pneumovax 23: exclusion added for PPV23 or PCV20 received within the last 5 years. The SmPC says revaccination within 3 years is not recommended and the Green Book restricts 5-yearly revaccination to asplenia, splenic dysfunction and chronic kidney disease. The vague 'may be considered' line is replaced with that rule.

Prevenar 13: hypersensitivity to diphtheria toxoid (the CRM197 carrier) added as an SmPC contraindication. Inclusion restricted to 2 years and over; infant primary immunisation is not given under this PGD and 'according to the national immunisation schedule' is replaced by a stated dose.

The guidance summary has not yet been rewritten against the July 2026 Green Book chapter (Prevenar 20, infant dose at 16 weeks). That is recorded in the review register for the next version.

Signed on behalf of Get Real Health

Version v003, 10 September 2026.

Name	Qualification	Signature	Date
Nitin Shori	Medical Director GMC: 6047293		10 September 2026
Chris Pilkington	Head Pharmacist GPhC: 2046322		10 September 2026

Both authorising signatories reviewed this version together on 10 September 2026 and gave their authorisation for it to be issued. Signatures were applied digitally on their joint instruction, which is the established process for Get Real Health PGDs. This version is not valid without both signatures.