

This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.

Patient Group Direction

for the administration of Phenoxymethylpenicillin 500mg tablets and Clarithromycin 250mg tablets for the treatment of Sore Throat Test & Treat

Summary of NICE / NICE CKS Guidelines for Sore Throat Test & Treat

Acute Sore Throat Assessment

- **Aetiology:** Most commonly viral (70-80%); Group A Streptococcus (GAS) responsible for 10-15% of bacterial sore throats.
- **Clinical presentation:** Sore throat, difficulty swallowing, pharyngitis, possible tonsillitis (exudate, erythema), fever.
- **Red flags:** Trismus, drooling, dysphagia, stridor, respiratory distress - refer for emergency assessment.

FeverPAIN and Centor Score

- **FeverPAIN score:** Fever, Purulence, Attend rapidly, No cough, Ill appearance. Score 0-5; score ≥ 4 suggests GAS infection.
- **Centor score:** Exudate, Fever, Nodes, No cough. Score 0-4; score ≥ 3 or rapid antigen test positive suggests GAS.
- **NICE guidance:** Offer antibiotics if FeverPAIN ≥ 4 or positive rapid GAS antigen test (RAST). Do not offer antibiotics for FeverPAIN 0-3 without test.

Group A Streptococcus Testing

- **RAST (rapid antigen test):** Can be used in primary care/pharmacy to guide antibiotic use (not part of this PGD).
- **Throat culture:** Gold standard but results delayed; not used for acute management.
- **Management:** Use FeverPAIN score or RAST to determine need for antibiotics.

Antibiotic Treatment

- **First-line (FeverPAIN ≥ 4 or RAST+):** Phenoxymethylpenicillin 500mg QDS for 5-10 days.
- **Penicillin allergy:** Use clarithromycin 250mg BD for 5 days.
- **No treatment:** FeverPAIN 0-3 without positive RAST - advise supportive care.

Patient Group Direction

for the supply/administration of

Phenoxymethylpenicillin (Penicillin V) 500mg tablets

for the treatment of

Sore Throat Test & Treat

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	- Pharmacist registered and practicing with the GPhC - Pharmacy technician registered and practicing with the GPhC
Training and assessment	Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. - All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines - All users must previously have used PGDs to supply medication - Must work in compliance with the SOPs of their own employer - All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	- All users must be up to date with CPD requirements and appraisal - All users must be up to date with MHRA and safety alerts with regards to medical products - All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	Treatment of acute sore throat caused by Group A Streptococcus (GAS) when FeverPAIN score ≥ 4 or positive RAST
Inclusion criteria	Adults aged 18 years and over Acute sore throat with FeverPAIN score ≥ 4 OR positive rapid antigen test for Group A Streptococcus Able to take oral medication No recent antibiotic use for this illness Capable of giving valid informed consent
Exclusion criteria	Known penicillin or beta-lactam allergy (use clarithromycin instead) Pregnancy requiring safer alternative (though penicillin V is generally safe) Severe hepatic or renal dysfunction
Cautions	Mild to moderate renal impairment: generally safe; monitor renal function Hepatic impairment: generally safe but monitor liver function Pregnancy and breastfeeding: penicillin V is generally safe; ensure informed consent

	Drug interactions: may reduce efficacy of oral contraceptives; advise additional contraception for 7 days Mononucleosis: avoid due to risk of rash (if uncertain of diagnosis, clarify before prescribing)
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate.

Details of the medicine

Name, form and strength of medicine	Phenoxymethylpenicillin 500mg tablets
Legal category	POM
Storage	Store below 25°C. Keep in original container. Protect from light and moisture.
Route/method of administration	Oral
Dose and frequency	500mg (one tablet) four times daily To be taken on an empty stomach (1 hour before or 2 hours after meals) for optimal absorption Continue for 5-10 days (typically 5 days minimum for GAS; longer courses sometimes used)
Quantity to be administered and/or supplied	20-40 tablets (5-10 day course)
Maximum or minimum treatment period	5-10 days
Adverse effects	Very common: nausea, diarrhoea Common: rash (non-allergic), vomiting, abdominal pain Uncommon: indigestion, oral candidiasis Rare: hypersensitivity reactions (urticaria, Stevens-Johnson syndrome), angioedema, anaphylaxis, C. difficile infection

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: - that valid informed consent was given - name of individual, address, date of birth and GP - name of healthcare practitioner

	• name and brand of medication • date of supply dose, form and route quantity supplied • advice given, including advice given if excluded or declines treatment • details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: • Adults aged 18 years and over - keep records for 8 years • Children aged under 18 years - keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)
--	---

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	Complete the full course of antibiotics as prescribed, even if symptoms improve within 2-3 days. Take phenoxymethylpenicillin (Penicillin V) on an empty stomach (1 hour before or 2 hours after food) for best absorption. Take clarithromycin with or without food; if GI upset occurs, take with food. If using oral contraception, use additional contraceptive methods during the antibiotic course and for 7 days afterwards. For pain relief, use paracetamol or ibuprofen as directed; a sore throat should improve within 3-5 days. Stay hydrated: drink plenty of water and other fluids to help soothe the throat. Use throat lozenges or warm salt water gargles for comfort (not part of medical treatment). Seek medical advice if symptoms worsen or do not improve after 3-5 days of treatment. Report any allergic reactions (rash, facial swelling, breathing difficulties) immediately. Inform your doctor if you develop persistent diarrhoea during or after clarithromycin (possible C. difficile infection). If you have a metallic taste with clarithromycin, this should resolve after stopping the medication. Do not share antibiotics with others; this course is prescribed for you alone.

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017
<https://www.nice.org.uk/guidance/mg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017
<https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance: Sore throat (acute): prescribing antibiotics
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature
Valid from date	Expiry date		
1/11/25	31/7/27		

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date
	(GPhC: 2046322)		
Name of professional group signatory	Job title & name of organisation	Signature	Date
	(GPhC: 2046322)		

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		2/11/25
Chris Pilkington	Pharmacist GPhC: 2046322		2/11/25

Change history

Version number	Change details	Date
001	Development & issue of new PGD	1st November 2025

Patient Group Direction

for the supply/administration of

Clarithromycin 250mg tablets

for the treatment of

Sore Throat Test & Treat

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	• Pharmacist registered and practicing with the GPhC • Pharmacy technician registered and practicing with the GPhC
Training and assessment	Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. • All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines • All users must previously have used PGDs to supply medication • Must work in compliance with the SOPs of their own employer • All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	• All users must be up to date with CPD requirements and appraisal • All users must be up to date with MHRA and safety alerts with regards to medical products • All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	Treatment of acute sore throat caused by Group A Streptococcus in patients with penicillin allergy (when FeverPAIN ≥ 4 or RAST positive)
Inclusion criteria	Adults aged 18 years and over Acute sore throat with FeverPAIN score ≥ 4 OR positive RAST Documented penicillin allergy or intolerance (non-anaphylaxis preferred for macrolide use) Able to take oral medication No contraindications to macrolide antibiotics Capable of giving valid informed consent
Exclusion criteria	Known hypersensitivity to macrolides (clarithromycin, erythromycin,

	azithromycin) Concurrent use of ergotamine, dihydroergotamine (risk of ergot toxicity) Concurrent use of certain statins (simvastatin, lovastatin) - increased statin levels QT prolongation or risk factors (hypokalaemia, hypomagnesaemia, cardiac arrhythmia history) Severe hepatic impairment Myasthenia gravis (macrolides can worsen muscle weakness)
Cautions	Hepatic impairment: mild-moderate use with caution; avoid if severe Renal impairment (eGFR <30 mL/min/1.73m²): dose adjustment or alternative needed QT interval risk: assess baseline risk; avoid in high-risk patients Drug interactions: significant interactions with statins, ergotamines, antiarrhythmics; check before prescribing Pregnancy and breastfeeding: relatively safe but ensure informed consent; small amount passes to breast milk Myasthenia gravis: may exacerbate; use with caution and close monitoring Anticoagulation: possible increased anticoagulant effect; monitor INR if on warfarin
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate.

Details of the medicine

Name, form and strength of medicine	Clarithromycin 250mg tablets
Legal category	POM
Storage	Store below 25°C. Keep in original container. Protect from moisture.
Route/method of administration	Oral
Dose and frequency	250mg (one tablet) twice daily May be taken with or without food Continue for 5 days
Quantity to be administered and/or supplied	10 tablets (5-day course)
Maximum or minimum	5 days

treatment period	
Adverse effects	Very common: GI upset (diarrhoea, nausea, abdominal pain), taste disturbance (metallic taste) Common: vomiting, indigestion, headache, abdominal pain, oral candidiasis Uncommon: hepatitis, jaundice, hypersensitivity reactions, dizziness Rare: QT prolongation, arrhythmias, Stevens-Johnson syndrome, C. difficile infection, severe hepatotoxicity

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: • that valid informed consent was given • name of individual, address, date of birth and GP • name of healthcare practitioner • name and brand of medication • date of supply dose, form and route quantity supplied • advice given, including advice given if excluded or declines treatment • details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: • Adults aged 18 years and over - keep records for 8 years • Children aged under 18 years - keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	Complete the full course of antibiotics as prescribed, even if symptoms improve within 2-3 days. Take phenoxymethylpenicillin (Penicillin V) on an empty stomach (1 hour before or 2 hours after food) for best absorption. Take clarithromycin with or without food; if GI upset occurs, take with food. If using oral contraception, use additional contraceptive methods during the antibiotic course and for 7 days afterwards.

	For pain relief, use paracetamol or ibuprofen as directed; a sore throat should improve within 3-5 days. Stay hydrated: drink plenty of water and other fluids to help soothe the throat. Use throat lozenges or warm salt water gargles for comfort (not part of medical treatment). Seek medical advice if symptoms worsen or do not improve after 3-5 days of treatment. Report any allergic reactions (rash, facial swelling, breathing difficulties) immediately. Inform your doctor if you develop persistent diarrhoea during or after clarithromycin (possible C. difficile infection). If you have a metallic taste with clarithromycin, this should resolve after stopping the medication. Do not share antibiotics with others; this course is prescribed for you alone.
--	---

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance: Sore throat (acute): prescribing antibiotics
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature

Valid from date	Expiry date		
1/11/25	31/10/26		

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date
Name of professional group signatory	Job title & name of organisation	Signature	Date

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		2/11/25
Chris Pilkington	Pharmacist GPhC: 2046322		2/11/25

Change history

Version number	Change details	Date
001	Development & issue of new PGD	1st November 2025
