

# SUPPLY PAUSED

Testosterone Replacement Therapy · Testogel, Tostran, Sustanon, Nebido

**Do not supply testosterone under this PGD until further notice.**

## Reason

Testosterone is a Schedule 4 Part II controlled drug. Whether a Patient Group Direction may lawfully be used to supply it is currently disputed and is being confirmed.

NHS Specialist Pharmacy Service guidance, updated 1 July 2026, states that unless a controlled drug appears on its permitted list it cannot be supplied under a PGD, and for Schedule 4 lists ·all drugs except anabolic steroids·. Schedule 4 Part II is the anabolic steroids part. On that reading this PGD has no lawful basis.

Earlier advice held the opposite. Because the two readings cannot both be right, and because the consequence of being wrong is unlawful supply of a controlled drug, supply is paused while a definitive answer is obtained.

**This replaces the correction notice issued earlier on 7 September 2026**

That notice restricted this PGD to pharmacists and stated that every other provision remained in force. That assurance was not verified and should not have been given.

## What to do now

Do not initiate or continue supply under this PGD. Patients already established on treatment should be referred to their prescriber or GP to arrange continuity. Do not stop **a patient's testosterone abruptly; arrange continuation through a prescriber.**

This is a pause, not a permanent withdrawal. The service will be restored if the legal position permits it.

Issued 7 September 2026 · Get Real Health

**This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.**

Patient Group Direction

for the administration of Testogel, Tostran, Sustanon and Nebido for the treatment of Low Testosterone

## Summary of NICE / NICE CKS Guidelines for Low Testosterone (Male Hypogonadism)

### Overview

- **Low testosterone** (hypogonadism) refers to **clinical symptoms** associated with **low serum testosterone levels**.
- Can be due to:
  - **Primary hypogonadism** (testicular failure)
  - **Secondary hypogonadism** (pituitary/hypothalamic dysfunction)
  - **Functional hypogonadism** (e.g. due to obesity, diabetes, chronic illness)

### Symptoms of Low Testosterone

- Reduced libido or erectile dysfunction
- Fatigue or low energy
- Low mood or depression
- Reduced muscle mass or strength
- Increased body fat
- Infertility
- Osteoporosis or low bone density

Symptoms are **non-specific**, and a **careful history and clinical judgement** are essential.

### Assessment

#### Indications for Testing

- Symptoms suggestive of hypogonadism
- Delayed or incomplete puberty
- Osteoporosis or low bone density in men
- Infertility

### *Investigations*

- **Early morning total testosterone** (7–11 am) — fasting if possible
    - If low ( $<12$  nmol/L), **repeat** to confirm
  - Consider **sex hormone-binding globulin (SHBG)** and **free testosterone** if total testosterone borderline (12–15 nmol/L)
  - **LH and FSH** to differentiate:
    - **High LH/FSH** = primary hypogonadism
    - **Low or normal LH/FSH** = secondary hypogonadism
  - Additional tests:
    - **Prolactin, pituitary function tests, MRI pituitary** if secondary cause suspected
    - **Karyotyping** if primary hypogonadism with small testes (e.g. Klinefelter's)
- 

## Management

### *Lifestyle and Treat Underlying Cause*

- Weight loss, reduce alcohol intake
- Optimise management of comorbidities (e.g. diabetes, sleep apnoea)
- Address medication causes (e.g. opioids, steroids)

### *Testosterone Replacement Therapy (TRT)*

- Consider if:
  - **Confirmed low testosterone AND**
  - **Consistent symptoms of hypogonadism**
- Not recommended for **age-related testosterone decline** alone without symptoms

### **Formulations:**

- **Transdermal gels/patches** (e.g. testosterone gel 50 mg daily)
- **Long-acting intramuscular injections** (e.g. testosterone undecanoate every 10–14 weeks)
- **Oral formulations** (less preferred)

Choice depends on patient preference, availability, and tolerability.

---

## Monitoring on TRT

- **Testosterone level** (aim for mid-normal range):
  - 2–3 months after starting
  - Then every 6–12 months
- Monitor:
  - **PSA** (especially in men  $>40$ )
  - **FBC** (check for polycythaemia)

- **LFTs, lipids, weight/BMI**
  - **Bone density** if osteoporosis present
- 

### Contraindications to TRT

- **Prostate or breast cancer**
  - **Elevated PSA without investigation**
  - **Severe untreated sleep apnoea**
  - **Polycythaemia (haematocrit >0.54)**
  - **Uncontrolled heart failure**
- 

### Referral to Endocrinology

- Diagnostic uncertainty
- Suspected pituitary disease
- Fertility concerns
- Contraindications to TRT or complex comorbidities

## Testing and monitoring regimen (all testosterone products)

Initial diagnosis: two early-morning fasting total testosterone measurements, taken at least four weeks apart, before treatment is offered.

Baseline panel (before first supply): total testosterone (early morning, fasting); free or calculated free testosterone; SHBG; LH and FSH; prolactin; full blood count and haematocrit; U&Es/eGFR; liver function tests; lipid profile; HbA1c or fasting glucose; PSA in men aged 40 and over.

Follow-up panel (6 to 12 weeks after initiation): testosterone level; full blood count and haematocrit; liver function tests; PSA where required (aged 40 and over).

Ongoing monitoring: every 3 to 6 months in the first year, then annually once stable — testosterone level; full blood count and haematocrit; liver function tests; lipid profile; PSA where required (aged 40 and over).

PSA is not routinely required for men under 40; it is offered as a separate add-on for men aged 40 and over.

Supply intervals: 28-day supplies for topical preparations aligned to review appointments (testosterone is a Schedule 4 controlled drug; 30 days is the good-practice maximum). Depot injection supply follows the product dosing interval.

# Patient Group Direction

for the supply/administration of

Testogel (testosterone gel)

for the treatment of

Low Testosterone

## ***Healthcare professionals covered and training***

|                                       | Healthcare professionals allowed to work under this PGD and requirements                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|---------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Qualifications and Registration       | <ul style="list-style-type: none"> <li>● Pharmacist registered and practicing with the GPhC</li> <li>● Pharmacy technician registered and practicing with the GPhC</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Training and assessment               | <ul style="list-style-type: none"> <li>● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team.</li> <li>● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines</li> <li>● All users must previously have used PGD's to supply medication</li> <li>● Must work in compliance with the SOPs of their own employer</li> <li>● All users must have appropriate indemnity in place to cover this service</li> </ul> |
| Further training and responsibilities | <ul style="list-style-type: none"> <li>● All users must be up to date with CPD requirements and appraisal</li> <li>● All users must be up to date with MHRA and safety alerts with regards to medical products</li> <li>● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.</li> </ul>                                                                                                                                                                                                                                                                  |

|                                                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PGD Indication                                       | Testogel is indicated for testosterone replacement therapy in adult men with confirmed primary or secondary hypogonadism associated with clinical symptoms and biochemical evidence of low serum testosterone.                                                                                                                                                                                                                                                                                                                                                 |
| Inclusion criteria                                   | <ul style="list-style-type: none"> <li>- Adult males aged 18 years and over.</li> <li>- Confirmed diagnosis of hypogonadism (low serum testosterone with symptoms).</li> <li>- Informed consent obtained.</li> <li>- No contraindications to testosterone therapy.</li> </ul>                                                                                                                                                                                                                                                                                  |
| Exclusion criteria                                   | <ul style="list-style-type: none"> <li>- Known or suspected carcinoma of the breast or prostate.</li> <li>- History of liver tumours or severe hepatic insufficiency.</li> <li>- Polycythaemia (haematocrit &gt;54%).</li> <li>- Untreated obstructive sleep apnoea.</li> <li>- Pregnancy or breastfeeding (contraindicated for female partners if risk of transference).</li> </ul> <p>Severe cardiac or renal disease.</p>                                                                                                                                   |
| Cautions                                             | <ul style="list-style-type: none"> <li>- Regularly monitor testosterone levels, haematocrit, PSA, and clinical response.</li> <li>- Avoid skin contact with others post-application; advise hand washing and covering site with clothing.</li> <li>- Discontinue if no clinical benefit or if significant adverse effects occur.</li> <li>- Counsel on application technique, safe handling, and disposal.</li> </ul> <p>Limited therapeutic experience in patients over 65 years of age — use with caution and review ongoing clinical benefit regularly.</p> |
| Actions if patient is excluded or declines treatment | <p>Advise on alternative treatment options and how these can be accessed.</p> <p>Document any advice given and the decision reached. Inform or refer to the GP as appropriate</p>                                                                                                                                                                                                                                                                                                                                                                              |

### *Details of the medicine*

|                                             |                                                                                                                              |
|---------------------------------------------|------------------------------------------------------------------------------------------------------------------------------|
| Name, form and strength of medicine         | Testogel                                                                                                                     |
| Legal category                              | POM                                                                                                                          |
| Storage                                     | Store below 25°C. Do not refrigerate or freeze.                                                                              |
| Route/method of administration              | Topical application to clean, dry, intact skin (shoulders, upper arms, or abdomen).                                          |
| Dose and frequency                          | Recommended starting dose is 50 mg of testosterone (one sachet or 1-2 pump actuation) once daily, preferably in the morning. |
| Quantity to be administered and/or supplied | 28-day supply based on 50–100 mg daily dose; adjusted according to serum testosterone levels.                                |



|                                     |                                                                                                                                            |
|-------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------|
| Maximum or minimum treatment period | Reassess at 3–6 months and then annually to confirm ongoing clinical benefit.                                                              |
| Adverse effects                     | Skin irritation, acne, increased red blood cell count, mood changes, prostate symptoms. Rare: polycythaemia, liver effects. Refer to SmPC. |

|                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|-----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Yellow card reporting | Report suspected adverse effects via <a href="https://yellowcard.mhra.gov.uk">https://yellowcard.mhra.gov.uk</a> and inform the GP as appropriate.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Records to be kept    | <p>Record the following information:</p> <ul style="list-style-type: none"> <li>● that valid informed consent was given</li> <li>● name of individual, address, date of birth and GP</li> <li>● name of healthcare practitioner</li> <li>● name and brand of medication</li> <li>● date of supply dose, form and route quantity supplied</li> <li>● advice given, including advice given if excluded or declines treatment</li> <li>● details of any adverse drug reactions and actions taken supplied via PGD</li> </ul> <p>Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made.</p> <p>A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows:</p> <ul style="list-style-type: none"> <li>● Adults aged 18 years and over - keep records for 8 years</li> <li>● Children aged under 18 years -keep records until the 25<sup>th</sup> birthday (if the patient was 17 years when treatment was finished, keep records until the 26<sup>th</sup> birthday)</li> </ul> |

## Patient information

|                                                     |                                                                                                                                           |
|-----------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| Written information to be given to patient or carer | Supply the patient information leaflet (PIL) provided with the medication.                                                                |
| Follow-up advice to be given to patient or carer    | To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell. |

## Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mpg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines [BNF \(British National Formulary\) | NICE](#)
- Summary of Product Characteristics for drug

### Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

| Name and address of pharmacy premises / clinic premises to which this PGD relates |           |                     |           |
|-----------------------------------------------------------------------------------|-----------|---------------------|-----------|
|                                                                                   |           |                     |           |
| Name of healthcare professional                                                   | Job title | Registration number | Signature |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
| Valid from date                                                                   |           | Expiry date         |           |
| 13/07/26                                                                          |           | 31/7/27             |           |

### PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

| Name of superintendent / clinical lead | Job title & name of organisation | Signature | Date |
|----------------------------------------|----------------------------------|-----------|------|
|                                        |                                  |           |      |

### Signed on behalf of Get Real Health

| Name             | Qualification                  | Signature                                                                          | Date      |
|------------------|--------------------------------|------------------------------------------------------------------------------------|-----------|
| Nitin Shori      | Medical Doctor<br>GMC: 6047293 |  | 13/07/26  |
| Chris Pilkington | Pharmacist<br>GPhC: 2046322    |  | 25/8/2025 |

### Change history

| Version number | Change details                                                                                                                                                                                                                                                              | Date                       |
|----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|
| 001            | Development & issue of new PGD                                                                                                                                                                                                                                              | 25 <sup>th</sup> June 2025 |
| 002            | Reviewed and authorised for the GRH platform; signed by Medical Director                                                                                                                                                                                                    | 13th July 2026             |
| 003            | Added shared testing and monitoring regimen (BSSM-aligned panels incl. U&Es baseline; PSA 40+ as separate add-on); severe cardiac or renal disease added to exclusions; over-65 caution added; supply interval guidance. Agreed with C. Pilkington and PPH clinical review. | 29th July 2026             |
|                |                                                                                                                                                                                                                                                                             |                            |

# Patient Group Direction

for the supply/administration of

Tostran (testosterone gel)

for the treatment of

Low Testosterone

## ***Healthcare professionals covered and training***

|                                       | Healthcare professionals allowed to work under this PGD and requirements                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|---------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Qualifications and Registration       | <ul style="list-style-type: none"> <li>● Pharmacist registered and practicing with the GPhC</li> <li>● Pharmacy technician registered and practicing with the GPhC</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Training and assessment               | <ul style="list-style-type: none"> <li>● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team.</li> <li>● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines</li> <li>● All users must previously have used PGD's to supply medication</li> <li>● Must work in compliance with the SOPs of their own employer</li> <li>● All users must have appropriate indemnity in place to cover this service</li> </ul> |
| Further training and responsibilities | <ul style="list-style-type: none"> <li>● All users must be up to date with CPD requirements and appraisal</li> <li>● All users must be up to date with MHRA and safety alerts with regards to medical products</li> <li>● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.</li> </ul>                                                                                                                                                                                                                                                                  |

|                                                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PGD Indication                                       | Tostran is indicated for testosterone replacement therapy in adult men with confirmed primary or secondary hypogonadism with clinical symptoms and laboratory confirmation of testosterone deficiency.                                                                                                                                                                                                                                                                                               |
| Inclusion criteria                                   | <ul style="list-style-type: none"> <li>- Adult males aged 18 years and over.</li> <li>- Confirmed diagnosis of hypogonadism (low serum testosterone with symptoms).</li> <li>- Informed consent obtained.</li> <li>- No contraindications to testosterone therapy.</li> </ul>                                                                                                                                                                                                                        |
| Exclusion criteria                                   | <ul style="list-style-type: none"> <li>- Known or suspected carcinoma of the breast or prostate.</li> <li>- History of liver tumours or severe hepatic impairment.</li> <li>- Polycythaemia (haematocrit &gt;54%).</li> <li>- Untreated obstructive sleep apnoea.</li> <li>- Risk of testosterone transference to others (household contact not manageable).</li> </ul> <p>Severe cardiac or renal disease.</p>                                                                                      |
| Cautions                                             | <ul style="list-style-type: none"> <li>- Monitor testosterone levels, haematocrit, PSA, and symptoms regularly.</li> <li>- Counsel on application technique, hygiene, and avoidance of transference.</li> <li>- Rotate application sites and avoid broken skin.</li> <li>- Reassess if inadequate clinical response or if adverse effects occur.</li> </ul> <p>Limited therapeutic experience in patients over 65 years of age — use with caution and review ongoing clinical benefit regularly.</p> |
| Actions if patient is excluded or declines treatment | <p>Advise on alternative treatment options and how these can be accessed.</p> <p>Document any advice given and the decision reached. Inform or refer to the GP as appropriate</p>                                                                                                                                                                                                                                                                                                                    |

### *Details of the medicine*

|                                             |                                                                                                                                             |
|---------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| Name, form and strength of medicine         | Tostran gel                                                                                                                                 |
| Legal category                              | POM                                                                                                                                         |
| Storage                                     | Store below 25°C. Do not refrigerate or freeze. Keep container tightly closed.                                                              |
| Route/method of administration              | Topical application to clean, dry, intact skin on the abdomen or inner thighs, preferably in the morning.                                   |
| Dose and frequency                          | Recommended starting dose is 3 g of gel (60 mg testosterone) once daily, adjusted based on clinical response and serum testosterone levels. |
| Quantity to be administered and/or supplied | One 60 g canister (approximately 30 doses), issued monthly or as appropriate to prescribed dose.                                            |

|                                     |                                                                                                                                                          |
|-------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Maximum or minimum treatment period | Reassess at 3–6 months and then annually to confirm continued need and benefit.                                                                          |
| Adverse effects                     | Application site reactions, acne, increased red blood cell count, mood changes, prostate enlargement. Rare: polycythaemia, liver effects. Refer to SmPC. |

|                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|-----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Yellow card reporting | Report suspected adverse effects via <a href="https://yellowcard.mhra.gov.uk">https://yellowcard.mhra.gov.uk</a> and inform the GP as appropriate.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Records to be kept    | <p>Record the following information:</p> <ul style="list-style-type: none"> <li>● that valid informed consent was given</li> <li>● name of individual, address, date of birth and GP</li> <li>● name of healthcare practitioner</li> <li>● name and brand of medication</li> <li>● date of supply dose, form and route quantity supplied</li> <li>● advice given, including advice given if excluded or declines treatment</li> <li>● details of any adverse drug reactions and actions taken supplied via PGD</li> </ul> <p>Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made.</p> <p>A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows:</p> <ul style="list-style-type: none"> <li>● Adults aged 18 years and over - keep records for 8 years</li> <li>● Children aged under 18 years -keep records until the 25<sup>th</sup> birthday (if the patient was 17 years when treatment was finished, keep records until the 26<sup>th</sup> birthday)</li> </ul> |

## ***Patient information***

|                                                     |                                                                                                                                           |
|-----------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| Written information to be given to patient or carer | Supply the patient information leaflet (PIL) provided with the medication.                                                                |
| Follow-up advice to be given to patient or carer    | To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell. |

## **Key references**

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> <li>● NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <a href="https://www.nice.org.uk/guidance/mpg2">https://www.nice.org.uk/guidance/mpg2</a></li> <li>● NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <a href="https://www.nice.org.uk/guidance/mpg2/resources">https://www.nice.org.uk/guidance/mpg2/resources</a></li> <li>● NICE CKS Guidance</li> <li>● British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines <a href="#">BNF (British National Formulary)   NICE</a></li> </ul> |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

● Summary of Product Characteristics for drug

**Agreement to practise by the Registered Healthcare Professional**

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

| Name and address of pharmacy premises / clinic premises to which this PGD relates |           |                     |           |
|-----------------------------------------------------------------------------------|-----------|---------------------|-----------|
|                                                                                   |           |                     |           |
| Name of healthcare professional                                                   | Job title | Registration number | Signature |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
| Valid from date                                                                   |           | Expiry date         |           |
| 13/07/26                                                                          |           | 31/7/27             |           |

**PGD adoption and authorisation by the employer/clinical lead**

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

| Name of superintendent / clinical lead | Job title & name of organisation | Signature | Date |
|----------------------------------------|----------------------------------|-----------|------|
|                                        |                                  |           |      |



### Signed on behalf of Get Real Health

| Name             | Qualification                  | Signature                                                                          | Date      |
|------------------|--------------------------------|------------------------------------------------------------------------------------|-----------|
| Nitin Shori      | Medical Doctor<br>GMC: 6047293 |  | 13/07/26  |
| Chris Pilkington | Pharmacist<br>GPhC: 2046322    |  | 25/8/2025 |

### Change history

| Version number | Change details                                                                                                                                                                                                                                                              | Date                       |
|----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|
| 001            | Development & issue of new PGD                                                                                                                                                                                                                                              | 25 <sup>th</sup> June 2025 |
| 002            | Reviewed and authorised for the GRH platform; signed by Medical Director                                                                                                                                                                                                    | 13th July 2026             |
| 003            | Added shared testing and monitoring regimen (BSSM-aligned panels incl. U&Es baseline; PSA 40+ as separate add-on); severe cardiac or renal disease added to exclusions; over-65 caution added; supply interval guidance. Agreed with C. Pilkington and PPH clinical review. | 29th July 2026             |
|                |                                                                                                                                                                                                                                                                             |                            |

# Patient Group Direction

for the supply/administration of

Sustanon (testosterone injection)

for the treatment of

Low Testosterone

## ***Healthcare professionals covered and training***

|                                       | Healthcare professionals allowed to work under this PGD and requirements                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|---------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Qualifications and Registration       | <ul style="list-style-type: none"> <li>● Pharmacist registered and practicing with the GPhC</li> <li>● Pharmacy technician registered and practicing with the GPhC</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Training and assessment               | <ul style="list-style-type: none"> <li>● Pharmacists and Pharmacy Technicians must have completed training with Get Real Health in order to use this PGD. This training will be documented and signed off by the Get Real Health team.</li> <li>● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines</li> <li>● All users must previously have used PGD's to supply medication</li> <li>● Must work in compliance with the SOPs of their own employer</li> <li>● All users must have appropriate indemnity in place to cover this service</li> </ul> |
| Further training and responsibilities | <ul style="list-style-type: none"> <li>● All users must be up to date with CPD requirements and appraisal</li> <li>● All users must be up to date with MHRA and safety alerts with regards to medical products</li> <li>● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.</li> </ul>                                                                                                                                                                                                                                                                       |

|                                                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PGD Indication                                       | Sustanon is indicated for testosterone replacement therapy in adult men with confirmed primary or secondary hypogonadism with clinical symptoms and biochemical evidence of testosterone deficiency.                                                                                                                                                                                                                                                                                                                      |
| Inclusion criteria                                   | <ul style="list-style-type: none"> <li>- Adult males aged 18 years and over.</li> <li>- Confirmed diagnosis of hypogonadism with low serum testosterone.</li> <li>- Informed consent obtained.</li> <li>- No contraindications to testosterone therapy.</li> </ul>                                                                                                                                                                                                                                                        |
| Exclusion criteria                                   | <ul style="list-style-type: none"> <li>- Known or suspected carcinoma of the prostate or breast.</li> <li>- Polycythaemia (haematocrit &gt;54%).</li> <li>- Severe cardiac, hepatic or renal impairment.</li> <li>- Untreated obstructive sleep apnoea.</li> <li>- Known hypersensitivity to any components of the injection.</li> </ul> <p>Severe cardiac or renal disease.</p>                                                                                                                                          |
| Cautions                                             | <ul style="list-style-type: none"> <li>- Monitor testosterone levels, haematocrit, PSA, lipids, and liver function regularly.</li> <li>- Educate on potential adverse effects and long-term monitoring.</li> <li>- Ensure injections are given intramuscularly and not intravenously.</li> <li>- Reassess benefit/risk balance periodically during long-term therapy.</li> </ul> <p>Limited therapeutic experience in patients over 65 years of age — use with caution and review ongoing clinical benefit regularly.</p> |
| Actions if patient is excluded or declines treatment | <p>Advise on alternative treatment options and how these can be accessed.</p> <p>Document any advice given and the decision reached. Inform or refer to the GP as appropriate</p>                                                                                                                                                                                                                                                                                                                                         |

### ***Details of the medicine***

|                                             |                                                                                                           |
|---------------------------------------------|-----------------------------------------------------------------------------------------------------------|
| Name, form and strength of medicine         | Sustanon injection                                                                                        |
| Legal category                              | POM                                                                                                       |
| Storage                                     | Store below 30°C. Do not refrigerate or freeze. Protect from light.                                       |
| Route/method of administration              | Intramuscular injection into the gluteal muscle by a trained healthcare professional.                     |
| Dose and frequency                          | Typically 1 mL (250 mg) every 2-3 weeks. Adjust based on serum testosterone levels and clinical response. |
| Quantity to be administered and/or supplied | One ampoule per injection; sufficient for 2-3 weeks of treatment.                                         |
| Maximum or minimum treatment period         | Reassess at 3–6 months and then annually to confirm continued need and effectiveness.                     |
| Adverse effects                             | Injection site pain, acne, mood changes, increased red blood cell count,                                  |

|  |                                                                            |
|--|----------------------------------------------------------------------------|
|  | fluid retention, prostate effects. Rare: liver dysfunction. Refer to SmPC. |
|--|----------------------------------------------------------------------------|

|                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|-----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Yellow card reporting | Report suspected adverse effects via <a href="https://yellowcard.mhra.gov.uk">https://yellowcard.mhra.gov.uk</a> and inform the GP as appropriate.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Records to be kept    | <p>Record the following information:</p> <ul style="list-style-type: none"> <li>● that valid informed consent was given</li> <li>● name of individual, address, date of birth and GP</li> <li>● name of healthcare practitioner</li> <li>● name and brand of medication</li> <li>● date of supply dose, form and route quantity supplied</li> <li>● advice given, including advice given if excluded or declines treatment</li> <li>● details of any adverse drug reactions and actions taken supplied via PGD</li> </ul> <p>Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made.</p> <p>A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows:</p> <ul style="list-style-type: none"> <li>● Adults aged 18 years and over - keep records for 8 years</li> <li>● Children aged under 18 years -keep records until the 25<sup>th</sup> birthday (if the patient was 17 years when treatment was finished, keep records until the 26<sup>th</sup> birthday)</li> </ul> |

## ***Patient information***

|                                                     |                                                                                                                                           |
|-----------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| Written information to be given to patient or carer | Supply the patient information leaflet (PIL) provided with the medication.                                                                |
| Follow-up advice to be given to patient or carer    | To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell. |

## **Key references**

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mpg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines [BNF \(British National Formulary\) | NICE](#)
- Summary of Product Characteristics for drug

## **Agreement to practise by the Registered Healthcare Professional**

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the

responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

| Name and address of pharmacy premises / clinic premises to which this PGD relates |           |                     |           |
|-----------------------------------------------------------------------------------|-----------|---------------------|-----------|
|                                                                                   |           |                     |           |
| Name of healthcare professional                                                   | Job title | Registration number | Signature |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
| Valid from date                                                                   |           | Expiry date         |           |
| 13/07/26                                                                          |           | 31/7/27             |           |

**PGD adoption and authorisation by the employer/clinical lead**

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

| Name of superintendent / clinical lead | Job title & name of organisation | Signature | Date |
|----------------------------------------|----------------------------------|-----------|------|
|                                        |                                  |           |      |

### Signed on behalf of Get Real Health

| Name             | Qualification                  | Signature                                                                          | Date      |
|------------------|--------------------------------|------------------------------------------------------------------------------------|-----------|
| Nitin Shori      | Medical Doctor<br>GMC: 6047293 |  | 13/07/26  |
| Chris Pilkington | Pharmacist<br>GPhC: 2046322    |  | 25/8/2025 |

### Change history

| Version number | Change details                                                                                                                                                                                                                                                              | Date                       |
|----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|
| 001            | Development & issue of new PGD                                                                                                                                                                                                                                              | 25 <sup>th</sup> June 2025 |
| 002            | Reviewed and authorised for the GRH platform; signed by Medical Director                                                                                                                                                                                                    | 13th July 2026             |
| 003            | Added shared testing and monitoring regimen (BSSM-aligned panels incl. U&Es baseline; PSA 40+ as separate add-on); severe cardiac or renal disease added to exclusions; over-65 caution added; supply interval guidance. Agreed with C. Pilkington and PPH clinical review. | 29th July 2026             |
|                |                                                                                                                                                                                                                                                                             |                            |

# Patient Group Direction

for the supply/administration of

**Nebido (testosterone undecanoate injection)**

for the treatment of

**Low Testosterone**



## ***Healthcare professionals covered and training***

|                                       | Healthcare professionals allowed to work under this PGD and requirements                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|---------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Qualifications and Registration       | <ul style="list-style-type: none"> <li>● Pharmacist registered and practicing with the GPhC</li> <li>● Pharmacy technician registered and practicing with the GPhC</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Training and assessment               | <ul style="list-style-type: none"> <li>● Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team.</li> <li>● All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines</li> <li>● All users must previously have used PGD's to supply medication</li> <li>● Must work in compliance with the SOPs of their own employer</li> <li>● All users must have appropriate indemnity in place to cover this service</li> </ul> |
| Further training and responsibilities | <ul style="list-style-type: none"> <li>● All users must be up to date with CPD requirements and appraisal</li> <li>● All users must be up to date with MHRA and safety alerts with regards to medical products</li> <li>● All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.</li> </ul>                                                                                                                                                                                                                                                                  |

|                                                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PGD Indication                                       | Nebido is indicated for testosterone replacement therapy in adult men with hypogonadism associated with clinical symptoms and confirmed by biochemical tests demonstrating testosterone deficiency.                                                                                                                                                                                                                                                                                                                                    |
| Inclusion criteria                                   | <ul style="list-style-type: none"> <li>- Adult males aged 18 years and over.</li> <li>- Confirmed diagnosis of testosterone deficiency (hypogonadism).</li> <li>- Informed consent obtained.</li> <li>- No contraindications to testosterone therapy.</li> </ul>                                                                                                                                                                                                                                                                       |
| Exclusion criteria                                   | <ul style="list-style-type: none"> <li>- Known or suspected prostate or breast carcinoma.</li> <li>- Severe cardiac, renal or hepatic disease.</li> <li>- Polycythaemia (haematocrit &gt;54%).</li> <li>- Untreated obstructive sleep apnoea.</li> <li>- Hypersensitivity to testosterone undecanoate or formulation ingredients.</li> </ul> <p>Severe cardiac or renal disease.</p>                                                                                                                                                   |
| Cautions                                             | <ul style="list-style-type: none"> <li>- Monitor serum testosterone, haematocrit, liver function, PSA, and clinical response routinely.</li> <li>- Counsel on long-term therapy, expected onset and duration of effects.</li> <li>- Ensure deep intramuscular injection; avoid intravenous use.</li> <li>- Monitor for symptoms of excessive androgen exposure or adverse effects.</li> </ul> <p>Limited therapeutic experience in patients over 65 years of age — use with caution and review ongoing clinical benefit regularly.</p> |
| Actions if patient is excluded or declines treatment | <p>Advise on alternative treatment options and how these can be accessed.</p> <p>Document any advice given and the decision reached. Inform or refer to the GP as appropriate</p>                                                                                                                                                                                                                                                                                                                                                      |

### *Details of the medicine*

|                                             |                                                                                                                           |
|---------------------------------------------|---------------------------------------------------------------------------------------------------------------------------|
| Name, form and strength of medicine         | Nebido injection                                                                                                          |
| Legal category                              | POM                                                                                                                       |
| Storage                                     | Store below 30°C. Do not refrigerate or freeze. Protect from light.                                                       |
| Route/method of administration              | Intramuscular injection into the gluteal muscle by a trained healthcare professional, administered slowly over 2 minutes. |
| Dose and frequency                          | 1000 mg (4 mL) every 10–14 weeks following an initial interval of 6 weeks after the first dose.                           |
| Quantity to be administered and/or supplied | One vial per injection, administered every 10–14 weeks.                                                                   |
| Maximum or minimum treatment period         | Reassess treatment at 3–6 months and annually thereafter based on                                                         |

|                 |                                                                                                                                                       |
|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|
|                 | testosterone levels and clinical response.                                                                                                            |
| Adverse effects | Injection site pain, acne, increased haematocrit, weight gain, mood changes, libido changes. Rare: polycythaemia, hepatic dysfunction. Refer to SmPC. |

|                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|-----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Yellow card reporting | Report suspected adverse effects via <a href="https://yellowcard.mhra.gov.uk">https://yellowcard.mhra.gov.uk</a> and inform the GP as appropriate.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Records to be kept    | <p>Record the following information:</p> <ul style="list-style-type: none"> <li>● that valid informed consent was given</li> <li>● name of individual, address, date of birth and GP</li> <li>● name of healthcare practitioner</li> <li>● name and brand of medication</li> <li>● date of supply dose, form and route quantity supplied</li> <li>● advice given, including advice given if excluded or declines treatment</li> <li>● details of any adverse drug reactions and actions taken supplied via PGD</li> </ul> <p>Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made.</p> <p>A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows:</p> <ul style="list-style-type: none"> <li>● Adults aged 18 years and over - keep records for 8 years</li> <li>● Children aged under 18 years -keep records until the 25<sup>th</sup> birthday (if the patient was 17 years when treatment was finished, keep records until the 26<sup>th</sup> birthday)</li> </ul> |

## Patient information

|                                                     |                                                                                                                                           |
|-----------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| Written information to be given to patient or carer | Supply the patient information leaflet (PIL) provided with the medication.                                                                |
| Follow-up advice to be given to patient or carer    | To seek medical advice if symptoms worsen rapidly or significantly, do not improve in 3–4 weeks, or they become systemically very unwell. |

## Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mpg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 <https://www.nice.org.uk/guidance/mpg2/resources>
- NICE CKS Guidance
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines [BNF \(British National Formulary\) | NICE](#)
- Summary of Product Characteristics for drug

### Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

| Name and address of pharmacy premises / clinic premises to which this PGD relates |           |                     |           |
|-----------------------------------------------------------------------------------|-----------|---------------------|-----------|
|                                                                                   |           |                     |           |
| Name of healthcare professional                                                   | Job title | Registration number | Signature |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
|                                                                                   |           |                     |           |
| Valid from date                                                                   |           | Expiry date         |           |
| 13/07/26                                                                          |           | 31/7/27             |           |

### PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

| Name of superintendent / clinical lead | Job title & name of organisation | Signature | Date |
|----------------------------------------|----------------------------------|-----------|------|
|                                        |                                  |           |      |

### Signed on behalf of Get Real Health

| Name             | Qualification                  | Signature                                                                          | Date      |
|------------------|--------------------------------|------------------------------------------------------------------------------------|-----------|
| Nitin Shori      | Medical Doctor<br>GMC: 6047293 |  | 13/07/26  |
| Chris Pilkington | Pharmacist<br>GPhC: 2046322    |  | 25/8/2025 |

### Change history

| Version number | Change details                                                                                                                                                                                                                                                              | Date                       |
|----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|
| 001            | Development & issue of new PGD                                                                                                                                                                                                                                              | 25 <sup>th</sup> June 2025 |
| 002            | Reviewed and authorised for the GRH platform; signed by Medical Director                                                                                                                                                                                                    | 13th July 2026             |
| 003            | Added shared testing and monitoring regimen (BSSM-aligned panels incl. U&Es baseline; PSA 40+ as separate add-on); severe cardiac or renal disease added to exclusions; over-65 caution added; supply interval guidance. Agreed with C. Pilkington and PPH clinical review. | 29th July 2026             |
|                |                                                                                                                                                                                                                                                                             |                            |

