

Tetanus, Diphtheria and Polio (Revaxis, Td/IPV)

Patient Group Direction, version 004, issued 9 September 2026. Revaxis, individuals aged 10 years and over.

Three changes in this version

- **PREGNANCY IS NOW AN UNQUALIFIED EXCLUSION.** Refer every pregnant individual to the GP or midwife, including after a tetanus-prone wound.
- **FIVE DOCUMENTED DOSES MEANS NO FURTHER VACCINE FOR A TETANUS-PRONE WOUND.** A fully immunised individual does not need another dose. Where the wound is high risk, the question is tetanus immunoglobulin, which is not a PGD supply. Refer.
- **A DOSE WITHIN THE LAST 12 MONTHS EXCLUDES.** Version 002 handled recent doses through an off-label note about the previous 5 years, which would have blocked a legitimate travel booster in someone who had a wound dose four years ago. 12 months is the operative figure.
- **Observe every patient for 15 minutes after vaccination.**

Revaxis (Td/IPV)

Patient Group Direction for the administration of Revaxis, adsorbed diphtheria (low dose), tetanus and inactivated poliomyelitis vaccine, to individuals aged 10 years and over.

Healthcare professionals covered, and training

- Pharmacist registered and practising with the GPhC.
- Pharmacy technician registered and practising with the GPhC.
- All users must be trained to the National Minimum Standards and Core Curriculum for Immunisation Training.
- All users must be competent in the recognition and immediate management of anaphylaxis, and in vaccine handling and cold chain management.
- All users must be able to apply Green Book chapter 30 table 30.1 to a tetanus-prone wound, and must know that a decision about tetanus immunoglobulin is a referral and not a PGD supply.
- All users must be able to count documented doses and to recognise when five have been given.
- All users must be familiar with Gillick competence and the assessment of consent in children and young people, and with the involvement of a person with parental responsibility.
- All users must previously have administered medicines under a PGD, must work in compliance with the SOPs of their own employer, and must hold appropriate indemnity covering this service.
- All users must be up to date with CPD and appraisal, and with MHRA safety alerts relating to this product.
- All users must understand capacity and consent and be aware of the Mental Capacity Act 2005.

The PGD

Indication	Active immunisation of individuals aged 10 years and over against tetanus, diphtheria and poliomyelitis, in accordance with chapters 15, 26 and 30 of Immunisation Against Infectious Disease (the Green Book). This PGD supports: the routine adolescent booster where it has been missed; completion of a primary course or booster where the immunisation history is incomplete, unknown or out of date; protection for travel in accordance with NaTHNaC recommendations; and a reinforcing dose for a tetanus-prone wound where immunoglobulin is not indicated.
Inclusion criteria	• Individuals aged 10 years and over. • Valid informed consent obtained from the individual, or from a person with parental responsibility where the individual is under 16 and not Gillick competent. • No tetanus, diphtheria or polio containing vaccine received in the last 12 months. • AND ONE OF THE FOLLOWING: ○ Requires a booster following a primary course against diphtheria, tetanus and poliomyelitis, usually offered at 13 to 18 years of age where not already completed; OR ○ Has no history, or an incomplete or uncertain history, of diphtheria, tetanus or poliomyelitis immunisation; OR ○ Is travelling to an area where medical attention may not be accessible should a tetanus-prone wound occur, or will be residing in an epidemic or endemic area, AND the final dose of the relevant antigen was received more than 10 years ago. This applies even where 5 doses have previously been received; OR ○ Has a tetanus-prone wound where a reinforcing dose is indicated under Green Book chapter 30 table 30.1, human tetanus immunoglobulin is NOT indicated, and the individual has FEWER than 5 documented doses. • No exclusion criterion present.
Five documented doses: what it means, and where it does not apply	Five documented doses of a tetanus-containing vaccine is a complete lifetime course. Understand where that exempts the individual and where it does not, because the two situations point in opposite directions. TETANUS-PRONE WOUND, 5 doses already given: NO further vaccine dose is needed. It adds nothing. Do not give one.

	• Where the wound is high risk, the question becomes whether tetanus immunoglobulin is indicated. That is a clinical assessment and an immunoglobulin supply, neither of which is covered by this PGD. Refer the same day. • Where the wound is not high risk and 5 doses have been given, no immunisation action is needed at all. Give wound care advice and record the assessment. TRAVEL, 5 doses already given: a booster IS indicated where the last dose was more than 10 years ago, and may be given under this PGD. • This is the one situation in which a fully immunised individual is boosted, and it exists because the traveller may not be able to reach medical care after a wound.
Exclusion criteria	Refer, do not vaccinate, where any of the following applies. • PREGNANCY, in all circumstances, including after a tetanus-prone wound. Refer to the GP or midwife the same day where the need is urgent. From week 16 of pregnancy a pertussis-containing vaccine is routinely indicated instead, which is a different product and a different decision. • A tetanus, diphtheria or polio containing vaccine received within the last 12 months. Refer to establish what was given and when. • Aged under 10 years. Refer to the GP or an appropriate immunisation service for an age-appropriate vaccine (DTaP/IPV/Hib/HepB or dTaP/IPV). • Confirmed anaphylactic reaction to a previous dose of a diphtheria, tetanus or poliomyelitis containing vaccine, including any conjugate vaccine in which diphtheria or tetanus toxoid is used as the carrier. • Confirmed anaphylactic reaction to any component of the vaccine, including neomycin, streptomycin or polymyxin B. • Acute severe febrile illness. The presence of a minor infection is not a contraindication. • A tetanus-prone wound where human tetanus immunoglobulin is indicated under Green Book chapter 30. Refer for same-day medical assessment. • A tetanus-prone wound in an individual with 5 or more documented doses. No further vaccine is needed; assess for immunoglobulin by referral. • Management of cases or contacts in an outbreak of diphtheria or poliomyelitis. These individuals must be managed by the local Health Protection Team and are outside this PGD. • Valid informed consent not given, or consent from a person with parental responsibility not obtained where required. • Current neurological deterioration. Defer and seek advice so that any change is not incorrectly attributed to the vaccine.
Cautions	Facilities and trained staff for the management of anaphylaxis must be available, with immediate access to adrenaline (epinephrine) 1 in 1,000 injection and a telephone (Green Book chapter 8 and Resuscitation Council UK guidance). OBSERVE EVERY PATIENT FOR 15 MINUTES AFTER VACCINATION. Vaccinate seated. Syncope can occur before or after vaccination, particularly in adolescents, and this PGD starts at 10 years of age. Ensure procedures are in place to avoid injury from a faint. A stable neurological condition is not a contraindication. Only current deterioration defers. Immunosuppression: the immune response may be reduced. Vaccination is still recommended and the individual should be advised that protection may be limited. Revaxis contains approximately 10 micrograms of phenylalanine per 0.5 mL dose. The NSPKU advises this amount is negligible and that individuals with

	phenylketonuria should take up the offer of immunisation. Bleeding disorders and anticoagulation: see route and method of administration. Where a person attends for a routine booster and has previously received a vaccine after a tetanus-prone wound, establish which vaccine was given and when. If it was the same vaccine and given after an appropriate interval, the routine booster is not required. If it was given within the last 12 months, this PGD excludes.
Actions if the patient is excluded or declines	Advise on alternative options and how to access them, including the GP practice and, for travel, an appropriate travel health service. Explain the protective effect of the vaccine, the risk of infection and the potential complications of disease. Where a tetanus-prone wound requires immunoglobulin or medical assessment, arrange same-day referral and do not delay. Where the individual is pregnant, refer to the GP or midwife and make the urgency explicit if a wound is involved. Document the advice given and the decision reached. Inform or refer to the GP as appropriate.
Yellow Card reporting	Report suspected adverse reactions via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	• That valid informed consent was given, and from whom. Where the individual is under 16, the name and relationship of the person with parental responsibility, or the basis of the Gillick assessment. • Patient name, address, date of birth, and the GP with whom they are registered. • THE NUMBER OF DOCUMENTED PRIOR DOSES, and the source of that information. • THE DATE OF THE MOST RECENT TETANUS-CONTAINING DOSE, and how it was established. • Which indication applied: adolescent booster, incomplete history, travel, or tetanus-prone wound. • Where a wound is involved, the assessment against Green Book chapter 30 table 30.1 and the conclusion on immunoglobulin. • Name and brand of vaccine, batch number and expiry date. • Date of administration, dose, route and anatomical site. • That the 15 minute observation period was completed. • Where off-label, that this was explained and consent given on that basis. • Name and registration number of the healthcare professional administering. • Advice given, including advice given if the individual is excluded or declines. • Details of any adverse drug reactions and the actions taken. • That the vaccine was administered under this PGD. Records signed, dated, legible and contemporaneous. Adults 18 and over: 8 years. Under 18s: until the 25th birthday, or the 26th where the individual was 17 when the course finished. Inform the GP.

The medicine

Name, form and strength	Revaxis, adsorbed diphtheria (low dose), tetanus and inactivated poliomyelitis vaccine, suspension for injection in a pre-filled syringe, 0.5 mL.
Legal category	POM.
Black triangle	No.
Off-label use	• Primary immunisation in individuals over 10 years of age is off-label but is administered in accordance with chapters 15, 26 and 30 of the Green Book.

	• Where a vaccine is administered off-label, inform the individual, parent or carer as part of the consent process that it is being given outside its product licence but in accordance with national guidance, and record that you did. • Version 002 also treated administration within 5 years of a previous diphtheria or tetanus toxoid containing vaccine as off-label. That is superseded: a dose within the last 12 months now excludes outright, and beyond 12 months no off-label statement is needed on that ground.
Dose and frequency of administration	• Single 0.5 mL dose per administration. • Adolescent booster: a single 0.5 mL dose at around 14 years of age, a minimum of 5 years after the pre-school booster. • Previously unimmunised, or unknown or incomplete history: a primary course of 3 doses with an interval of one month between doses. Where a course is interrupted it should be resumed, not restarted. • First booster: at least 5 years after the third primary dose. Second booster: a minimum of 5 years and ideally 10 years after the first booster, where fewer than 5 documented doses have been given. • Travel: a single booster where the last dose was more than 10 years ago, including where 5 doses have already been given. • Tetanus-prone wound: a single reinforcing dose where the last tetanus-containing dose was more than 10 years ago, or in accordance with Green Book chapter 30 table 30.1 where the history is incomplete or uncertain, AND fewer than 5 documented doses have been given. • In every case: no dose if a tetanus, diphtheria or polio containing vaccine has been given in the last 12 months.
Quantity to be administered	Single 0.5 mL dose per administration. This PGD does not permit supply of vaccine to the individual for administration elsewhere.
Route and method of administration	• Intramuscular injection, preferably into the deltoid muscle of the upper arm. • Shake the pre-filled syringe well before administration. The normal appearance is a cloudy white suspension that may sediment during storage. • Inspect visually for foreign particulate matter or any variation from the expected appearance. If present, do not administer and discard in accordance with local procedures. • Where given at the same time as other vaccines, use separate sites, preferably different limbs, or at least 2.5 cm apart in the same limb. Record the site of each. • Individuals on stable anticoagulation, including those on warfarin who are up to date with INR testing and whose latest INR was below the upper threshold of their therapeutic range, may be vaccinated intramuscularly using a 23 gauge or finer needle, followed by firm pressure without rubbing for at least 2 minutes. Advise on the risk of haematoma. Where the intramuscular route is not suitable, give by deep subcutaneous injection.
Storage and cold chain	Store at +2C to +8C. Do not freeze. Store in the original packaging to protect from light. Vaccine stored outside these conditions must be quarantined and risk assessed in accordance with UKHSA Vaccine Incident Guidance before any decision on continued use.
Disposal	Equipment used for immunisation, including used syringes and any discharged vaccine, must be disposed of in a UN-approved puncture-resistant sharps container in accordance with local arrangements and HTM 07-01.
Adverse effects	Very common and common: injection site pain, erythema and swelling, headache, malaise, myalgia, fatigue, fever. Uncommon: lymphadenopathy, nausea, dizziness. Rare: hypersensitivity reactions including urticaria and angioedema. Very rare: anaphylaxis, and extensive limb swelling which is self-

	limiting. Consult the current SPC.
--	------------------------------------

Information for the patient

Written information	Supply the manufacturer's patient information leaflet, and a written record of the vaccine given with the date, brand and batch number. Tell the individual to keep it: the number of doses they have had determines what happens if they are ever injured.
Counselling	• A sore arm, mild fever, headache or aching for a day or two is common and settles by itself. • Keep the record of this vaccination. If you are ever injured, whether you need anything depends on how many doses you have had and when. • For any dirty wound, a puncture wound, a burn, an animal bite, or a wound with soil or manure in it, clean it and seek medical advice, whatever vaccinations you have had. • Where a course is involved, come back for every dose. We have booked your next one. • Where you are travelling: this protects you, but it does not remove the need to get any significant wound cleaned and assessed while you are away.
Follow-up	Book the next dose of any course at this appointment. Seek medical advice for any tetanus-prone wound. For a routine query about the vaccine, contact the pharmacy.

Appendix 1: The three questions to settle before vaccinating

1. Is the individual pregnant?	• YES: EXCLUDE. Refer to the GP or midwife, same day where a wound is involved. • This is now unqualified. There is no "except where protection is required without delay" route under this PGD.
2. Has any tetanus, diphtheria or polio containing vaccine been given in the last 12 months?	• YES: EXCLUDE. Refer to establish what was given and when. • NO: proceed to question 3. • Record the date and how it was established.
3. How many documented doses have they had?	• FEWER THAN 5: proceed on the applicable indication. • 5 OR MORE, and the reason is a WOUND: no vaccine needed. Refer for an immunoglobulin assessment if the wound is high risk. • 5 OR MORE, and the reason is TRAVEL with the last dose over 10 years ago: a booster is indicated and may be given.

Appendix 2: Key references

- Immunisation Against Infectious Disease (the Green Book), chapter 15 (diphtheria), chapter 26 (poliomyelitis) and chapter 30 (tetanus), including table 30.1 on tetanus-prone wounds.
- Green Book chapter 8, immunisation procedures, for anaphylaxis provision.
- Summary of Product Characteristics for Revaxis, current version.
- NaTHNaC, travelhealthpro.org.uk, for travel recommendations.
- UKHSA Vaccine Incident Guidance.
- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions, <https://www.nice.org.uk/guidance/mpg2>

Authorisation

Version	v004
Supersedes	Version 003, 8 September 2026
Valid from date	9 September 2026
Expiry date	31 July 2027

Doctor	Name: Nitin Shori Job title: Medical Director, Get Real Health GMC: 6047293 Signed: N. Shori Date: 9 September 2026
Pharmacist	Name: Chris Pilkington Job title: Head Pharmacist, Get Real Health GPhC: 2046322 Signed: C. Pilkington Date: 9 September 2026

Both authorising signatories reviewed this version together on 9 September 2026 and gave their authorisation for it to be issued. Signatures were applied digitally on their joint instruction, which is the established process for Get Real Health PGDs.

Adoption by the employing organisation

To be completed by the adopting pharmacy. These signatories are the adopting organisation own signatories and must not be pre-filled by Get Real Health.

Superintendent / clinical lead	Name: Job title and organisation: Signature: Date:
Professional group signatory	Name: Job title and organisation: Signature: Date:

Change history

Version	v004
Date	8 September 2026
Changes	• The practitioner Agreement to practise page, the premises block and the practitioner signature table are restored. Every version 001 document carried them; no document produced by the current generator did, through one omission in one function, which left 15 live documents with no page for an individual practitioner to sign. Raised as blocking by an adopting pharmacy: NICE MPG2 expects a record of the individuals authorised to work under a PGD. The standard governance requirements are restored at the same time: SPC and BNF familiarity, MHRA safety alerts, CPD and appraisal, indemnity, and capacity and consent, all of which were in version 001 and were lost in the rewrite. This version changes no clinical content. • Pregnancy is an unqualified exclusion in all circumstances, including after a tetanus-prone wound. Version 002 excluded pregnancy "except where protection is required without delay" in its exclusion criteria while its cautions said to refer pregnant individuals to the GP or midwife, so the same page both permitted and refused the supply. • Five documented doses now handled explicitly, and differently for the two situations it arises in. For a tetanus-prone wound, 5 doses means no further vaccine is needed and the question becomes immunoglobulin, which is a referral. For travel with the last dose more than 10 years ago, a booster is still indicated. Version 002 stated only the travel case. • A tetanus, diphtheria or polio containing vaccine given within the last 12 months is now an outright exclusion. Version 002 handled recent doses through an off-label note about the previous 5 years, which would have blocked a legitimate travel booster in someone who had a wound dose four years earlier. 12 months was chosen so that it does not. • A 15 minute observation period after every vaccination is stated in the document, consistently with every other vaccination PGD in this estate. Version 002 noted the risk of syncope in adolescents without setting an observation period. • Records must now carry the number of documented prior doses and its source, the date of the most recent tetanus-containing dose, which indication applied, and, where a wound is involved, the assessment against table 30.1 and the conclusion on immunoglobulin. • Appendix 1 added: the three questions to settle before vaccinating, in order.

Previous versions

001	Development and issue of new PGD.
002, 6 August 2026	Revaxis Td/IPV, individuals aged 10 years and over, covering the adolescent booster, incomplete or unknown history, travel, and tetanus-prone wounds where immunoglobulin is not indicated.

Standard requirements for every practitioner working under this PGD

All users must be familiar with the current Summary of Product Characteristics for every product named in this PGD, and with current BNF and national guidance for the condition.

All users must be up to date with MHRA drug safety alerts and recalls relevant to these products.

All users must be up to date with their CPD requirements and appraisal.

All users must hold appropriate professional indemnity covering this service.

All users must understand capacity and consent, including the Mental Capacity Act 2005, and must be able to assess consent in children and young people where this PGD covers them.

By signing below you confirm that you agree to the contents of this PGD and that you will work within it. A PGD does not remove the inherent professional obligations or accountability of the individual practitioner. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Premises name
 Address
 Postcode:
 ODS code, if applicable

The employing organisation must keep this page, or an equivalent local register, and must be able to produce it on request. A practitioner who has not signed against the current version is not authorised to work under it.

[illegible]