


**+ - Test** - Works good for normal distribution.

- The t-test assesses whether the mean of two groups are statistically different from each other.



$$= M_2 - M_1$$

$$= 60 - 40$$

$$= 20$$

Does this 20 is significantly different or not?

$$t \text{ test} = \frac{\text{Difference between mean of two groups}}{\text{Standard error of difference between mean}} = \frac{M_2 - M_1}{(\text{SE of diff mean})}$$

- Why not directly find difference (i.e. 20) and say they are different. It means at the mean may show a difference, but we can't be sure if that is a reliable difference.

- Suppose me and you toss a coin 100 times. and find number of heads.

Your heads = 52, My heads = 49 times.

So this does not suggest, you will get heads more than me in future. This is only a chance.

→ **Descriptive statistics** - It does not guarantee that this are likely to happen again.

These are just stats such as mean that describe data you have, but can't generalized beyond that.

→ **Inferential statistics** - These are stats, such as t-test that allow us to make inference about population beyond our data. Example - Medicines made.

$$t = \frac{\text{variance between groups}}{\text{variance between within groups}}$$

A big t-value = different value groups.

A small t value = similar groups

- Each t-value has a p-value. The p-value tells us likelihood that there is real difference.

- Pvalue will tell if two group are really different or just a fluke. Pvalue is the probability that the pattern of data in the sample could be produced by random data.

$P = 0.10$ , there is 10% chance there is no real difference. (Rejected)

$P = 0.05$ , there is 5% chance there is no real difference.  $P \leq 0.05$

$P = 0.01$ , there is 1% chance there is no real difference (Accepted)

- Bigger samples make it easier to detect differences. with 2 group of 5,  $t = 2.0$ ,  $p = 0.04$

- A good guideline is to aim for 20 to 30 datapoint in each group. 2 group of 10,  $t = 2.0$ ,  $p = 0.03$

**Types of T-test** - i) Independent-sample ii) Paired Sample iii) One-sample

1) **Independent sample t test** → Testing the average quality of two different batches of beer. Also known as between-sample & unpaired-sample t test

2) **Paired sample t test** → Test the mean of one group twice. Eg- Testing balance before and after drinking. Also known as within-subject, repeated measures and dependent sample t test.

3) **One sample t test** → Test the mean of one group against known population mean. Eg- Testing if your co-workers IQ differ from average of 100.