A SHAPE THEOREM FOR EXPLODING SANDPILES

AHMED BOU-RABEE

ABSTRACT. We study scaling limits of exploding Abelian sandpiles using ideas from percolation and front propagation in random media. We establish sufficient conditions under which a limit shape exists and show via a family of counterexamples that convergence may not occur in general. A corollary of our proof is a simple criteria for determining if a sandpile is explosive; this strengthens a result of Fey, Levine, and Peres (2010).

1. INTRODUCTION

1.1. **Overview.** We consider the Abelian sandpile growth model on the integer lattice. Start with a *background* of indistinguishable chips, $\eta : \mathbb{Z}^d \to \mathbb{Z}$, add n chips at the origin, and attempt to *stabilize* via *parallel toppling*:

$$\begin{aligned} v_{t+1} &= v_t + 1\{s_t \geq 2d\} \\ s_{t+1} &= s_t + \Delta(v_{t+1} - v_t), \end{aligned} \tag{1}$$

where $\Delta v(x) = \sum_{y \sim x}(v(y) - v(x))$ is the Laplacian on $\mathbb{Z}^d$, $v_0 = 0$ is the initial *odometer*, and $s_0 = \eta + n\delta_0$ is the starting sandpile. We say s_0 is *stabilizable* if there is $T < \infty$ so that $v_t = v_T$ for all $t \geq T$. A background is *robust* if $\eta + n\delta_0$ is stabilizable for all $n \geq 1$, and otherwise is *explosive*. When $s_0 = \eta + n\delta_0$ is not stabilizable, it is explosive and the infinite sequence $\{s_t\}_{t \geq 0}$ is an *exploding* sandpile. See Figures 1 and 2.

Fey, Levine, and Peres coined these notions in [FLP10] (see also [FR05]) and provided sufficient conditions for determining if a background is explosive or robust: backgrounds $\eta \leq (2d-2)$ are always robust, but otherwise can be robust or explosive, depending on the arrangement of sites with $(2d-1)$ chips. In fact, they showed that if $\eta \leq (2d-2)$, not only is the background robust, but if n chips are added to the origin of such η, the diameter of the set of sites which topple grows like $n^{1/d}$.

Pegden and Smart used this bound together with the theory of viscosity solutions to show that the terminal odometer for $n\delta_0$, after a rescaling, converges to the solution of a fully




FIGURE 1. $s_t 1\{v_t > 0\}$ for $\eta \sim$ Bernoulli$(3, 2, 1/2)$ and $t = 50, 100, 250, 500$. The color white denotes sites which haven't toppled yet, otherwise white, yellow, red, blue, and black correspond to values $0, 1, 2, 3, (\geq 4)$.




FIGURE 2. Snapshot of the support of an exploding sandpile $1\{v_t > 0\}$ for $\eta \sim \text{Bernoulli}(2d-1, 2d-2, p)$ the first time it exits a box of side length 500. The top row is $d = 2$ and the bottom $d = 3$. From left to right, $p = 1/4, 1/2, 3/4$.

nonlinear elliptic PDE [PS13]. This breakthrough then led to an explanation for the patterns which appear in two-dimensional sandpiles [PS20, LPS16, LPS17]. Recently, the author used stochastic homogenization methods to extend convergence to all initial backgrounds which are stationary, ergodic, and bounded from above by $(2d-2)$ [BR21].

These results explain the phenomena of *scale-invariance* in sandpiles which have a *compact*, $n^{1/d}$, growth rate — large, compact-growth sandpiles look like high-resolution versions of smaller sandpiles. Simple models of growth are of interest to the mathematics and physics communities — see, for example, [DS13, DF91, GG98, PW85] and the references therein. The Abelian sandpile in particular has a rich history, [LKG90, Dha99, LBR02, Ost03, Red05, FdBR08, HLM$^+$08, FMR09, LP09, LP10, Pao13, LP17, Jár18, Kli18, HJL19, AS19, LS19, HS19, CKF20, Mel20, AM20].

In this paper, we study limit shapes of sandpiles in the explosive regime. The techniques used differ fundamentally from the existing compact-growth theory. Indeed, as we will demonstrate, some explosive sandpiles (both random and deterministic) do not converge. On the other hand, compact-growth sandpiles essentially always have limits — the argument there is 'soft' and applies in wide generality. Our proof below is quantitative and involves establishing specific, finite-scale estimates. We identify sufficient conditions under which exploding sandpiles converge to the level set of an asymmetric norm — much like in first-passage percolation [CD81] and threshold growth [Wil78, GG93].

1.2. **Main results.** For expositional clarity, we consider one family of random, explosive backgrounds with limit shapes. The reader interested in generalizations may consult Section 5. Suppose $d \geq 2$ and $\eta : \mathbb{Z}^d \to \mathbb{Z}$ is drawn from a product measure $\mathbf{P}$ with

$$0 < p \leq 1, \qquad \nu_p = (1-p)\delta_{2d-2} + p\delta_{2d-1}, \qquad \mathbf{P} = \bigotimes_{x \in \mathbb{Z}^d} \nu_p, \tag{2}$$
$$\mathbf{P}(\eta(x) = 2d-2) = 1-p, \qquad \mathbf{P}(\eta(x) = 2d-1) = p \quad (x \in \mathbb{Z}^d).$$

Fey, Levine, and Peres showed the following.

Theorem 1.1 (Proposition 1.4 in [FLP10])**.** *Let η be as in* (2)*. If $p > 0$, η is explosive with probability 1.*

Fix $p > 0$ and denote the (almost surely finite) *explosion threshold* by

$$M_\eta := \min\{n \geq 1 : \ \eta + n\delta_0 \text{ is not stabilizable}\}. \tag{3}$$

We prove that the support of the infinite sequence of parallel toppling odometers, $\{v_t\}_{t\geq 1}$, for the explosive sandpile $s_0 = \eta + M_\eta\delta_0$ converges under rescaling.

Theorem 1.2. *Let η be as in* (2)*. There exists a convex domain $\mathcal{B}(p) \subset \mathbb{R}^d$ so that on an event of probability 1,*

$$|\{\bar{v}_t > 0\} \triangle \mathcal{B}(p)| \to 0,$$

where $\bar{v}_t(x) := v_t([tx])$.

In fact, we prove something stronger. Not only does the support of the explosion converge, but the rate at which the explosion spreads also converges.

Theorem 1.3. *Let η be as in* (2) *and let $T_\eta(x) := \min\{t \geq 1 : v_t(x) > 0\}$. On an event of probability 1, the rescaled arrival times $\bar{T}_\eta(x) := n^{-1}T_\eta([nx])$ converge locally uniformly to $\mathcal{N}_p$, a continuous, convex, one-homogeneous function on $\mathbb{R}^d$.*

During our proof of Theorem 1.3, we introduce a quantitative criterion for determining if a sandpile is explosive. The criterion asserts, roughly, that if a sandpile explodes quickly on a finite box, then it must do so on the entire lattice. We use this and a coupling with bootstrap percolation to extend the aforementioned Theorem 1.1.

Theorem 1.4. *For $d \geq 2$, let ν be a probability measure on $\mathbb{Z}$ with $\nu(\{m \in \mathbb{Z} : m \geq d\}) = 1$ and $\nu(\{2d-1\}) > 0$. If $\beta : \mathbb{Z}^d \to \mathbb{Z}$ has law $\mathbf{P} = \bigotimes_{x\in\mathbb{Z}^d} \nu$, then β is explosive with probability 1.*

1.3. **Proof outline.** An exploding sandpile may be thought of as a heterogeneous, discrete reaction-diffusion equation. This perspective leads us to the literature for stochastic homogenization of reaction-diffusion equations [ZZ20, Fel19, LZ19, AC18, CM14, GG06, IPS99, BS98, GG96, GG93, Wil78]. These works suggest two methods of proof. The first, which we do not pursue, is *half-space propagation*—a limit shape can be completely described by those starting with a half-space initial condition—an early example of this technique appears in [Wil78]. Another method is to identify a subadditive quantity similar to the first-passage time, [CD81], which (directly or indirectly) describes the limit shape, then apply the subadditive ergodic theorem.

Our proof of Theorem 1.3 follows the second outline, however, there are several hurdles to overcome. A fundamental one is the nonlinear diffusion of the sandpile. This nonlinearity can cause explosions to propagate irregularly. In fact, an arbitrary exploding sandpile may spread quickly in certain cells, but slowly in others, causing convergence to fail. We demonstrate an explicit family of counterexamples to that effect in Section 6. A major part of our argument is showing that this irregularity cannot happen if $\eta \geq (2d-2)$ and there are 'enough' sites with $(2d-1)$ chips.

From now until Section 5, take η as in the statement of Theorem 1.3 and fix $p > 0$. We begin in Section 2 by showing that explosions on η spread quickly. This is done by establishing a high-probability bound on the 'crossing-speed' of η in a finite, but large cube and

passing to a coarsened lattice. On the coarsened lattice, there is an infinite cluster of 'good cubes' upon which explosions are guaranteed to spread quickly. We use this together with large-deviations for the chemical distance of supercritical Bernoulli percolation to get uniform, linear bounds on the arrival times. (This portion of the proof shares some similarities with Schonmann's argument for bootstrap percolation [Sch92].)

At this stage, if the arrival times, T_η, were subadditive, we could apply the subadditive ergodic theorem and conclude. However, T_η is not, in general, subadditive: roughly, when an explosion started at the origin reaches some site x for the first time, it mixes up the background and so cannot be compared directly to the explosion originating from x.

We overcome this lack of subadditivity by shifting our focus to a related, but simpler process, the *last-wave*—an exploding sandpile where the origin is constrained to topple a fixed number of times. In Section 3, we use the established regularity of explosions to show that the last-wave can be approximated by a quantity which is exactly subadditive, and hence converges. This can also be viewed as a shape theorem for a bootstrap percolation type process.

The proof of Theorem 1.3 is completed in Section 4 where we show that the last-wave is an approximation to the expanding front of an exploding sandpile. The argument for this is a deterministic comparison which requires $\eta \geq (2d-2)$. Then, in Section 5, we generalize Theorem 1.3 by presenting sufficient hypotheses under which a limit shape exists. We demonstrate some need of these hypotheses by constructing a family of (random and deterministic) exploding sandpiles which fail to have a limit shape in Section 6. We conclude in Section 7 with a proof of Theorem 1.4. There we indicate explicit criteria for determining if a sandpile is explosive.

Use of artificial intelligence. The original paper (2021) was written by the author. This corrected edition was prepared while the paper was formalized in the Lean proof assistant. The Lean formalization, `https://github.com/nitromannitol/Exploding-Sandpiles`, was written by AI coding agents (Anthropic Claude and OpenAI GPT models, with DeepSeek and Mistral models for auxiliary proof search) under the author's supervision. The agents also found the corrections listed in the section "Corrections to the published version" and drafted the corrected statements, the new proofs and Remark 1. The author checked these.

Every numbered result of this edition and Remark 1 is proved in Lean. Three cited results enter as explicit hypotheses: Theorem 1.1 of Antal and Pisztora [AP96]; Theorem 4.1 of Fey, den Boer and Redig [FdBR08] and the radius bound of Fey, Levine and Peres [FLP10], which enter for Lemma 3.1 and Remark 1. Proposition 7.1 is formalized for $d \geq 2$, and Hypothesis 1 is used in the form stated in this edition.

The author takes responsibility for the mathematical content and presentation of the paper.

Acknowledgments. Thank you to Charles K. Smart for motivating, helpful discussions during this project. Thank you to Lionel Levine for several inspiring conversations and for suggesting this question. Thank you to Dylan Airey for asking if exploding sandpiles have a limit shape.

Code. Julia code which can compute the figures in this article is included in the arXiv upload and may be freely used and modified.

Notation and conventions.

- Functions on $\mathbb{Z}^d$ are extended via nearest-neighbor interpolation to $\mathbb{R}^d$. That is, if f is a function on $\mathbb{Z}^d$, then for $x \in \mathbb{R}^d$, we define $f(x)$ to be $f([x])$ where $[x]$ is the nearest (breaking ties by picking the coordinate-wise minimum) point in $\mathbb{Z}^d$ to x.
- d will always refer to the dimension of the underlying space.
- $e_1, \dots, e_d$ are the d unit directions in $\mathbb{Z}^d$.
- $x_i = x \cdot e_i$ is the ith coordinate of vector x and $\mathbf{x}_{d-i} = (x_{i+1}, \dots, x_d)$.
- $|x|$ refers to the Manhattan norm and $|x|_\infty$ the ℓ_∞ norm.
- $y \sim x$ if $|y - x| = 1$.
- For $x, y \in \mathbb{R}^d$,
$$[x, y] = [x_1, y_1] \times \cdots \times [x_d, y_d]$$
and for $a, b \in \bar{\mathbb{R}}$
$$[a, b]^d = [a, b] \times \cdots \times [a, b]$$
and for $x \in \mathbb{R}^d$ and $b \in \bar{\mathbb{R}}$,
$$[x, b]^d = [x_1, b] \times \cdots \times [x_d, b]$$
and vice-versa.
- The symbol $\cdot$ will sometimes be used (for visual clarity) to denote scalar multiplication.
- Scalar operations on vectors/functions/sets are interpreted pointwise.
- For a function $f : A \to \mathbb{R}$ and a subset of its domain $\mathcal{S} \subset A$, we denote the restriction of f to $\mathcal{S}$ by $f|_{\mathcal{S}}$.
- $|\cdot|$ is either the counting measure or Lebesgue measure depending on the input.
- For $A \subset \mathbb{Z}^d$, $A^c := \{x \in \mathbb{Z}^d : x \notin A\}$, $\partial A := \{x \in A^c : \exists y \in A, x \sim y\}$, and $\bar{A} := A \cup \partial A$. The inner boundary of A is denoted by $\partial^\circ A := \{x \in A : \exists y \in A^c, x \sim y\}$.
- C, c, are positive constants which may change from line to line. Dependence is indicated by, for example, C_d or $C_d(\eta)$.
- The act of *firing* or *toppling* a site, x, removes $2d$ chips from x and adds one chip to each $y \sim x$.
- $\eta \sim \text{Bernoulli}(a, b, p)$ is shorthand for: $\eta : \mathbb{Z}^d \to \mathbb{Z}$ has law $\mathbf{P} = \bigotimes_{x \in \mathbb{Z}^d}((1-p)\delta_b + p\delta_a)$, with this same marginal at every site.

2. Regularity of explosions

We use $\eta \geq (2d-2)$ together with the i.i.d. assumption to establish almost sure regularity of explosions. The main result of this section is a quantification of [FLP10]'s Theorem 1.1 recalled above. The method is a static renormalization (see Chapter 7 in [Gri13]) inspired by Schonmann's proof for bootstrap percolation [Sch92].

2.1. **Parallel toppling preliminaries.** Before proceeding, we mention some basic properties of parallel toppling which we use below. Recall that $\{v_t\}_{t \geq 1}$ and $\{s_t\}_{t \geq 1}$ are the infinite sequence of parallel toppling odometers and sandpiles for initial conditions $v_0 = 0$ and $s_0 = \eta + M_\eta \delta_0$. An induction argument ((4.4) in [BG07]) shows

$$\begin{aligned} v_{t+1}(x) &= \min\{\lfloor \frac{s_0(x) + \sum_{y \sim x} v_t(y)}{2d} \rfloor, v_t(x) + 1\} \\ s_{t+1}(x) &= s_0(x) + \Delta v_{t+1}(x). \end{aligned} \tag{4}$$

Another induction shows that when $s_0 \le 2\cdot(2d)-1$, the minimum over the $v_t(x)+1$ term in (4) is unnecessary.

In the sequel we also consider a version of parallel toppling $\{w_t\}_{t\ge 0}$ where the initial value w_0 is given and the odometer on some set, $\mathcal{S}$, (the complement of a cube, the origin) is 'frozen' at w_0 and the initial sandpile is such that the topplings w_0 were performed: $s_0' = s_0 + \Delta w_0$ and,

$$w_{t+1}(x) = \begin{cases} w_t(x) + 1\{s_t'(x) \ge 2d\} & \text{if } x \notin \mathcal{S} \\ w_0(x) & \text{if } x \in \mathcal{S} \end{cases} \tag{5}$$
$$s_{t+1}' = s_t' + \Delta(w_{t+1} - w_t).$$

We call this $\mathcal{S}$-*frozen* parallel toppling. (Recently, and after this paper was written, [GMP21] was posted—therein so-called 'freezing sandpiles' are studied in the context of computational complexity.) If $s_0 + \Delta w_0 \le 2\cdot(2d)-1$ on $\mathcal{S}^c$, then as above:

$$w_{t+1}(x) = \max\left\{ w_t(x), \left\lfloor \frac{s_0(x) + \sum_{y\sim x} w_t(y)}{2d} \right\rfloor \right\} \qquad (x \notin \mathcal{S}), \tag{6}$$
$$s_{t+1}'(x) = s_0(x) + \Delta w_{t+1}(x).$$

Also, if $s_0 \le (2d-1)$ and $w_0 \equiv 0$ on $\mathcal{S}^c$, then $w_t|_{\mathcal{S}^c} \le \max_{x\in\partial^\circ\mathcal{S}} w_0(x)$ for all $t \ge 0$. The definitions allow us to compare the two versions of parallel toppling,

$$\begin{aligned} v_{t+t_0} &\ge w_t \qquad \text{where } t_0 := \min\{t \ge 0 : v_t \ge w_0\} \\ w_t &\ge v_t \qquad \text{if } w_0 \ge v_0 \equiv 0 \text{ and } w_0|_{\mathcal{S}} \ge \sup_t (v_t|_{\mathcal{S}}). \end{aligned} \tag{7}$$

2.2. **Crossing speeds.** To provide a global upper bound on the arrival times, $\mathcal{T}_\eta(x)$, we show a local upper bound. In particular we study the following 'cell problem', a term from homogenization denoting a simple problem which describes the local behavior of a more complicated one.

We consider sandpile dynamics on a box of side length k, $Q_k := \{x \in \mathbb{Z}^d : 1 \le x \le k\}$. For a point $z \in \bar{Q}_k$ and direction $1 \le i \le d$, denote the line passing from one side of the box to the other

$$\mathcal{L}_k^{(i,z)} := \bigcup_{j=1,\dots,k} (z_1, \dots, z_{i-1}, j, z_{i+1}, \dots, z_d). \tag{8}$$

Let w_t be the parallel toppling odometer for $\{Q_k^c \cup \mathcal{L}_k^{(i,z)}\}$-frozen parallel toppling (defined in (5)) with initial conditions $w_0(x) = 1\{x \in \mathcal{L}_k^{(i,z)}\}$, $s_0' = \eta + \Delta w_0$. Denote the *crossing time*

$$\mathfrak{C}_k^{(i,z)} := \begin{cases} \min\{t \ge 1 : w_t|_{Q_k} = 1\} & \text{if } w_\infty|_{Q_k} = 1 \\ \infty & \text{otherwise.} \end{cases} \tag{9}$$

We show that if k is sufficiently large, the crossing time is bounded with high probability.

Proposition 2.1. *For every $\delta > 0$, there is a k so that*

$$\max_{i,z} \mathfrak{C}_k^{(i,z)} \le k^d \tag{10}$$

with probability at least $(1-\delta)$.

Proof. For each $k \geq 1$, we construct an event for which (10) occurs with probability approaching 1 in k. By Harris' inequality (see, *e.g.*, [AS16]) and symmetry, it suffices to show (10) for lines in one direction, say $i = 1$. Let $z \in \bar{Q}_k$ be given.

Write $\mathfrak{C}_k = \mathfrak{C}_k^{(1,z)}$. We show that if all lines, $\mathcal{L}_k^{(1,y)}$, $y \in Q_k$, contain at least one site with $(2d-1)$ chips, then every site in the cube eventually topples. Denote the event upon which this happens by,

$$\Omega' := \bigcap_{y \in Q_k} \Omega_y := \bigcap_{y \in Q_k} \{\eta : \mathbb{Z}^d \to \mathbb{Z} : \eta(x) = (2d-1) \text{ for some } x \in \mathcal{L}_k^{(1,y)}\}. \tag{11}$$

Recall $p > 0$ is the probability of a site having $(2d-1)$ chips. Fix $0 < \epsilon < p$, and note, by Hoeffding's inequality (see, *e.g.*, [AS16]), for each $y \in Q_k$,

$$\mathbf{P}\Big(\sum_{x \in \mathcal{L}_k^{(1,y)}} 1(\eta(x) = 2d-1) \leq (p-\epsilon)k \Big) \leq \exp(-2\epsilon^2 k).$$

Therefore, by the union bound, (deleting duplicates),

$$\mathbf{P}(\Omega'^c) \leq k^{d-1} \mathbf{P}(\Omega_1'^c) \leq k^{d-1} \exp(-2\epsilon^2 k),$$

and so for every $\delta > 0$, there is k sufficiently large so that $P(\Omega') \geq 1 - \delta$.

It remains to check that for $\eta \in \Omega'$, $\mathfrak{C}_k \leq k^d$. We do so by constructing a toppling procedure which is dominated by w_t. After firing $\mathcal{L}_k^{(1,z)}$, all sites in neighboring lines, $\mathcal{L}_k^{(1,y)}$, $\mathbf{y}_{d-1} \sim \mathbf{z}_{d-1}$ have at least $(2d-1)$ chips. In fact, since $\eta \in \Omega'$, at least one site in each neighbor $\mathcal{L}_k^{(1,y)} \subset Q_k$ has $2d$ chips, causing all sites in the line to topple. Iterating shows this procedure will terminate with every site in Q_k toppling in at most k^d steps.

□

2.3. **A static renormalization scheme.** We exhibit a coarsening of the lattice upon which explosions are guaranteed to spread quickly. A cube, Q_k, is *good* if $\max_{i,z} \mathfrak{C}_k^{(i,z)} \leq k^d$. For each $i \in \mathbb{Z}^d$, let

$$Q_k(i) := Q_k + i \cdot k. \tag{12}$$

The cubes $\{Q_k(i)\}_{i \in \mathbb{Z}^d}$ define a *macroscopic lattice* with edge set $\{(Q_k(i), Q_k(j)), |j-i| = 1\}$. For k sufficiently large, Proposition 2.1 implies that the good cubes form an independent site process of high density. An edge $(Q_k(i), Q_k(j))$ of the macroscopic lattice is *good* when both of its cubes are good. A good edge depends only on its two cubes, so the good edges have a dependence graph of degree $4d-2$, and by Corollary 1.4 of [LSS97] they dominate an independent bond percolation process ζ of density close to one. The cubes all of whose ζ-edges are open form a finitely dependent site process, and by the same corollary it dominates an independent site percolation process of high density. The infinite cluster of this independent site process lies in an infinite open cluster of ζ, whose cubes are good. Lemma 5.2 of [Mat08] is stated for independent site percolation, so it applies to the independent site process and bounds its holes; Theorem 1.1 of [AP96], a chemical distance bound for supercritical bond percolation, applied to ζ together with the Borel–Cantelli lemma, bounds its chemical distances. These bounds give the following. (See, for example, Section 5 in [Mat08] for an explicit proof.)

Proposition 2.2. *For fixed k large enough, there are constants c, C so that the following hold on an event of probability 1.*




FIGURE 3. Terminal A-frozen parallel toppling odometer, w_∞, on $\mathbb{Z}^2$ for initial conditions $w_0 = 1\{x \in A\}$ and $s_0' = 2 + \Delta 1_A$. The set A is denoted by red pixels and black pixels are sites which eventually topple.

(1) *There is a unique infinite cluster* $\mathcal{C}_\infty$ *of good cubes on the macroscopic lattice* $\{Q_k(i)\}_{i\in\mathbb{Z}^d}$.
(2) *There is* n_0 *so that for* $n \geq n_0$*, any connected component of* $\mathcal{C}_\infty^c$ *that intersects* $[-n, n]^d$ *has volume smaller than* $(\log n)^{5/2}$.
(3) *There is* n_0 *so that for* $n \geq n_0$*, for any* $x, y \in \mathcal{C}_\infty$ *with* $|x| \leq n$ *and* $|x - y| \geq (\log n)^2$,

$$c|x - y| \leq d(x, y) \leq C|x - y|,$$

where d *is the chemical (graph) distance on* $\mathcal{C}_\infty$.

The definition of $\mathcal{C}_\infty$ ensures that once $Q_k(i) \in \mathcal{C}_\infty$ is *overlapped* by the support of the odometer — $Q_k(i) \cap 1\{v_t > 0\}$ contains a straight line — an explosion will occur. This together with Proposition 2.2 controls the speed at which the explosion propagates. We show next that the explosion spreading in $\mathcal{C}_\infty$ also quickly fills holes in the cluster.

2.4. **A path-filling property.** For a set of points $A \subset \mathbb{Z}^d$, let $m_i := \min_{z\in A} z_i$ and $M_i := \max_{z\in A} z_i$ for $i = 1, \ldots, d$. Denote the bounding rectangle of A as

$$\mathbf{br}(A) := \{z \in \mathbb{Z}^d : \ m \leq z \leq M\}. \tag{13}$$

We show, using $\eta \geq (2d - 2)$, that if the odometer is strictly positive on a path of points at some time then eventually the odometer is strictly positive on the bounding rectangle of that path. Essentially, if not then the support of the terminal odometer must have a corner, *i.e.*, an untoppled site with two neighbors which have toppled, a contradiction. Our proof uses this idea together with a slightly technical induction (which, it seems, we cannot avoid as the claim is needed in all dimensions). See Figure 3 for an illustration of this result.

Lemma 2.1. *Let* $A := \{z^{(i)}\}$ *be a finite path* $z^{(i)} \sim z^{(i+1)}$. *Let* w_t *denote* A*-frozen parallel toppling with initial conditions* $w_0(x) = 1\{x \in A\}$ *on the whole lattice (zero off* A*),* $s_0' = \eta + \Delta w_0$. *Then* $w_t|_{\mathbf{br}(A)} = 1$ *for every* $t \geq |\mathbf{br}(A)|$.

Proof. By monotonicity of parallel toppling, we may take $\eta = (2d - 2)$. Moreover, it suffices to show that every site in $\mathbf{br}(A)$ eventually topples, as if no site topples at time t, then no site topples at time $(t + 1)$.

We say A contains a $(\pm i, \pm j)$ turn if $z^{(m)} = z^{(m-1)} \pm e_i$ and $z^{(m+1)} = z^{(m)} \pm e_j$ for some $i \neq j$ where $z^{(m-1)} \sim z^{(m)} \sim z^{(m+1)}$ are in A. If A does not contain a turn, then $\mathbf{br}(A) = A$. Hence, we may suppose it contains at least one turn.

Case 1 - one-turn path. By shifting coordinates, we may suppose A contains only a $(1,2)$ turn and that

$$A = \{0, e_1, \ldots, k_1 e_1, k_1 e_1 + e_2, \ldots, k_1 e_1 + k_2 e_2\} \tag{14}$$

for $k_1 \geq k_2$. We induct on k_2. If $k_2 = 1$, then after firing every site in A, all sites in $A + e_2$ get one chip, while the corner site, $((k_1 - 1)e_1 + e_2)$ gets 2 chips. Since $\eta = (2d - 2)$, that corner becomes unstable and fires, causing all of its neighbors to the left, $((k_1 - 2)e_1 + e_2) \sim ((k_1 - 3)e_1 + e_2) \sim \cdots \sim e_2$ to fire. Continuing the induction shows that every site in $\mathbf{br}(A) = \{x \in \mathbb{Z}^d : 0 \leq x \leq (k_1 e_1 + k_2 e_2)\}$ eventually fires.

Case 2 - cubic path. We call A a *cubic* path if, after an isometry,

$$A = \{0, e_1, \ldots, k_1 e_1, k_1 e_1 + e_2, \ldots, \sum_{i=1}^{d} k_i e_i\} \tag{15}$$

for $k_1 \geq \cdots \geq k_d \geq 0$. Let $d_0 := \max\{i \leq d : k_i > 0\}$. We induct on d_0, the base case $d_0 = 2$ established in Case 1. For notational convenience, suppose the claim holds for $d_0 = (d-1)$ and we verify it for $d_0 = d$.

Consider the $(d-1)$-turn subpaths,

$$\begin{aligned} \mathcal{P}_1 &:= \{0, e_1, \ldots, \sum_{i=1}^{d-1} k_i e_i\} \\ \mathcal{P}_2 &:= \{k_1 e_1, k_1 e_1 + e_2, \ldots, \sum_{i=1}^{d} k_i e_i\}. \end{aligned} \tag{16}$$

By the inductive hypothesis, after $\mathcal{P}_i$ fire, both $(d-1)$-dimensional faces,

$$\begin{aligned} \mathcal{F}_1 &:= \{x \in \mathbb{Z}^d : 0 \leq x \leq \sum_{i=1}^{d-1} k_i e_i\} \\ \mathcal{F}_2 &:= \{x \in \mathbb{Z}^d : k_1 e_1 \leq x \leq \sum_{i=1}^{d} k_i e_i\} \end{aligned} \tag{17}$$

fire. We then 'fill in' the cube by identifying newly fired $(d-1)$-turn paths:

$$\begin{aligned} \mathcal{P}'_j := \{&je_2, je_2 + e_1, \ldots, je_2 + k_1 e_1, \\ &je_2 + k_1 e_1 + e_3, \ldots, je_2 + k_1 e_1 + k_3 e_3, \\ &\cdots \\ &je_2 + \sum_{i \neq 2} k_i e_i\} \end{aligned} \tag{18}$$

which are in $\mathcal{F}_1 \cup \mathcal{F}_2$ for $j = 0, \ldots, k_2$. By the inductive hypothesis, the firing of each $\mathcal{P}'_j$ makes every $(d-1)$-dimensional layer,

$$\mathcal{L}_j := \{x \in \mathbb{Z}^d : je_2 \leq x \leq je_2 + \sum_{i \neq 2} k_i e_i\},$$

fire and $\mathbf{br}(A) = \cup_{j=0}^{k_2} \mathcal{L}_j$.

Case 3 - general path. It suffices to show that if there is a path of firings between any two distinct points x, y, then $\mathbf{br}(\{x, y\})$ eventually fires. Before showing this, we suppose it were true and demonstrate sufficiency. Take $q \in \mathbf{br}(A)$ and observe by definition there are points $(z^{(1)}, Z^{(1)}), \ldots, (z^{(d)}, Z^{(d)})$ in A with $z_i^{(i)} \leq q_i \leq Z_i^{(i)}$. Then,

$$q^{(1)} := (q_1, \mathbf{q}'_{d-1}) \in \mathbf{br}(\{z^{(1)}, Z^{(1)}\})$$

for some $(d-1)$-vector $\mathbf{q}'_{d-1}$. Continue and let

$$q^{(2)} := \begin{cases} (q_1, q_2, \mathbf{q}''_{d-2}) \in \mathbf{br}(\{q^{(1)}, z^{(2)}\}) & \text{if } q_2^{(1)} \geq q_2 \\ (q_1, q_2, \mathbf{q}'''_{d-2}) \in \mathbf{br}(\{q^{(1)}, Z^{(2)}\}) & \text{otherwise} \end{cases}$$

for some $(d-2)$-vectors $\mathbf{q}''_{d-2}, \mathbf{q}'''_{d-2}$. After iterating, we find $q^{(d)} = q$, which shows that eventually q will fire.

Now fix two points $x, y \in A$ and decompose a path between them into a sequence of cubic paths

$$\mathcal{P}^{(1)} := \bigcup_{i=1}^{m} \mathcal{P}_i^{(1)},$$

where $\mathcal{P}_i^{(1)} := \{p_{i-1}, \ldots, p_i\}$ is cubic and $p_0 := x$ and $p_m := y$. (This can be done by, for example, starting at x and exploring the path but cutting whenever the cubic condition is violated.) Case 2 shows that eventually every site in $\mathcal{A}^{(1)} := \bigcup_{i=1}^{m} \mathbf{br}(\mathcal{P}_i^{(1)})$ will fire. If $m = 1$, we are done, otherwise we construct a new cubic path from p_0 to p_2 passing through $\mathcal{A}^{(1)}$. Once we have shown this, we iterate to conclude.

Suppose $p_0 = 0$, $p_1 = \sum_{j=1}^{d} k_j e_j$, and

$$p_2 = \sum_{j=1}^{d_1} (k_j - k'_j) e_j + \sum_{j=(d_1+1)}^{d_2} (k_j - k'_j) e_j + \sum_{j=(d_2+1)}^{d} (k_j + k'_j) e_j,$$

for some $1 \leq d_1 \leq d_2 \leq d$ and $k_j, k'_j \geq 1$ where $(k_j - k'_j) < 0$ for $j \leq d_1$ and $(k_j - k'_j) \geq 0$ for $(d_1 + 1) \leq j \leq d_2$. After this coordinate change, it suffices to exhibit a path from p_0 to p_2 with differences constrained to be $-e_j$ for $j = 1, \ldots, d_1$ and $+e_j$ for $j = (d_1 + 1), \ldots, d$.

There is a cubic path (only positive moves) from p_0 to

$$w_1 := \sum_{j=(d_1+1)}^{d_2} (k_j - k'_j) e_j + \sum_{j=(d_2+1)}^{d} (k_j) e_j$$

contained within $\mathbf{br}(\{p_0, p_1\})$ as $p_0 = 0 \leq w_1 \leq p_1$. Then, since $w_1 \in \mathbf{br}(\{p_1, p_2\})$ there is a cubic path (only positive moves) from w_1 to

$$w_2 := w_1 + \sum_{j=(d_2+1)}^{d} k'_j e_j$$

contained in $\mathbf{br}(\{p_1, p_2\})$ and similarly there is a cubic path (only negative moves) from w_2 to

$$p_2 = w_2 + \sum_{j=1}^{d_1} (k_j - k'_j) e_j.$$

Our new cubic path is the concatenation of these three paths: $p_0 \to w_1 \to w_2 \to p_2$. $\square$

3. The last-wave

In this section we study a simplified parallel toppling procedure closely related to bootstrap percolation (see Section 7 for an explicit connection, we do not utilize the coupling here). This simplified process has an inherent subadditive structure which allows us to prove convergence using the subadditive ergodic theorem. In the next section we show that this process is a good approximation to an exploding sandpile.

3.1. **The n-wave process.** Fix $n \geq 1$, $z \in \mathbb{Z}^d$, and consider the *n-wave* for η starting at z,

$$\begin{aligned} u_0^{(n,z)} &:= n\delta_z \\ u_{t+1}^{(n,z)}(x) &:= \lfloor \frac{\sum_{y\sim x} u_t^{(n,z)}(y) + \eta(x)}{2d} \rfloor \text{ for } x \neq z. \end{aligned} \tag{19}$$

Note that there is intentionally no minimum in (19) with $u_t^{(n,z)}(x)+1$ and it is possible that $u_{t+1}^{(n,z)}(x) > u_t^{(n,z)}(x)+1$. In particular, the n-wave started at z dominates the $\{z\}$-frozen parallel toppling process defined in (5) with the same initial conditions but may not coincide with it.

Overloading terminology, the n-wave is *stabilizable* if there is $T < \infty$ so that $u_t^{(n,z)} = u_T^{(n,z)}$ for all $t \geq T$. Let

$$\hat{M}_\eta(z) := \min\{n \geq 1 : \text{the } n\text{-wave for } \eta \text{ starting at } z \text{ is not stabilizable}\}. \tag{20}$$

We write $u_t^{(z)} := u_t^{(\hat{M}_\eta(z),z)}$ for the $\hat{M}_\eta(z)$-wave starting at z and call this the *last-wave*. We also consider the *penultimate-wave* starting at z as the terminal odometer for the $(\hat{M}_\eta(z)-1)$-wave, $\tilde{u}^{(z)}$, defined to be the zero function when $\hat{M}_\eta(z) = 1$. The set of sites touched by the penultimate wave is its *penultimate-cluster*,

$$\mathcal{P}(z) := \{z\} \cup \{x \in \mathbb{Z}^d : \text{ there is } y \sim x \text{ with } \tilde{u}^{(z)}(y) > 0\} \tag{21}$$

(we included the point, $\{z\}$, as $\hat{M}_\eta(z)$ may be 1). When z is the origin, we omit the superscripts.

The *arrival time* for the last-wave starting at site y to site x is

$$\hat{T}(y,x) := \min\{t \geq 1 : u_t^{(y)}(x) > 0\} \tag{22}$$

and the *penultimate-cluster arrival time* is

$$\tilde{T}(y,x) := \min\{t \geq 1 : u_t^{(y)}|_{\mathcal{P}(x)} > 0\}. \tag{23}$$

Note that the inequality in (23) is strict — $\tilde{T}(y,x)$ is the first time the support of the last wave started at y contains the entire penultimate wave of x, $\mathcal{P}(x)$. We write $\hat{T}(x) := \hat{T}(0,x)$ and $\tilde{T}(x) := \tilde{T}(0,x)$. (The choice of the same letter T for all arrival times is intentional — we will see they are asymptotically close.)

3.2. **Basic properties of the last-wave.** We derive some basic properties of the last-wave. Throughout this section and the next, let η be drawn from the event of probability 1 in Proposition 2.2 and let k be the (deterministic but large) side length of a good cube.

For the wave in (19), the value at its starting site is frozen: $u_t^{(m,z)}(z) = m$ for every $t \geq 0$. The parameter $m \in \mathbb{N}$ is a prescribed toppling count, not a number of added chips or a time horizon. Here the support of a wave means $\{x : \text{there exists } t \geq 0 \text{ with } u_t^{(m,z)}(x) > 0\}$.

Write $\Lambda_n = [-n, n]^d \cap \mathbb{Z}^d$ for the cube of radius n. The quantitative estimate below uses a lower bound on the radius of the toppled cube from Fey, den Boer and Redig [FdBR08] and an upper bound on the toppled set from Fey, Levine and Peres [FLP10].

Lemma 3.1. *For each integer $d \geq 2$ there is a deterministic constant $C_d > 0$ with the following property. Let $\eta : \mathbb{Z}^d \to \mathbb{Z}$ satisfy $2d - 2 \leq \eta(x) \leq 2d - 1$ at every site. For every integer $n \geq 1$ and every integer $m \geq 1$ satisfying*

$$m \geq \begin{cases} \lceil C_2 n^2 \log(n+1) \rceil, & d = 2, \\ \lceil C_d n^d \rceil, & d \geq 3, \end{cases},$$

there exists a finite integer T such that $u_t^{(m,0)}(x) > 0$ for every $t \geq T$ and every $x \in \Lambda_n$. In particular the support of this wave contains Λ_n.

Proof. Set $h = 2d - 2$. Let Δ be the unnormalized Laplacian, $\Delta f(x) = \sum_{y \sim x} f(y) - 2df(x)$. For $N \geq 2$, let v_N be the terminal ordinary toppling odometer for $h + N\delta_0$, and let $s_N = h + N\delta_0 + \Delta v_N$. Throughout this proof N counts extra chips and m counts frozen origin topplings. Ordinary legal toppling preserves nonnegative heights, so $0 \leq s_N \leq 2d - 1$.

Step 1. We show that $\Lambda_n \subseteq \operatorname{supp} v_N \subseteq \mathcal{D}$ for the choice of N and $\mathcal{D}$ made below, using two cited radius bounds.

Theorem 4.1 of Fey, den Boer and Redig [FdBR08] gives finite stabilization and a cubical toppled set $\operatorname{supp} v_N = \Lambda_{r_N}$, with

$$r_N \geq \frac{N^{1/d} - 3}{2}.$$

Their model parametrizes the total origin height, which is $N + h$ here, so their lower bound with $N + h$ implies the bound with N. Equivalently, once the cubical shape is known, chip conservation gives $N = \sum_x (s_N(x) - h) \leq |\Lambda_{r_N+1}| = (2r_N + 3)^d$, because all height changes lie in Λ_{r_N+1}, each increase is at most one, and the sum of the finitely supported Laplacian is zero.

Theorem 1.1 of Fey, Levine and Peres [FLP10], applied at $h = 2d - 2$ and $\epsilon = 1$, gives

$$\operatorname{supp} v_N \subseteq \Lambda_{A_d N^{1/d}}, \qquad A_d = (d+1)\omega_d^{-1/d},$$

for all sufficiently large N, where ω_d is the volume of the Euclidean unit ball. Its hypothesis $d \leq h \leq 2d - 2$ holds for every $d \geq 2$ considered here. The bound concerns toppled sites, not the particle cluster with its extra boundary. Choose an integer $N_0(d) \geq 2$ so the bound holds for every $N \geq N_0(d)$, and in the remainder take

$$N = (2n+3)^d + N_0(d), \qquad L = \lceil A_d N^{1/d} \rceil, \qquad \mathcal{D} = \Lambda_L.$$

Thus $\Lambda_n \subseteq \operatorname{supp} v_N \subseteq \mathcal{D}$, including for $n = 1$.

Step 2. We show that the walk killed on leaving $\mathcal{D}$ has a Green function $G_{\mathcal{D}}$ with $G_{\mathcal{D}}(0,0)$ and $\sum_{y \in \mathcal{D}} G_{\mathcal{D}}(0,y)$ bounded in terms of L.

Let X_t be the continuous-time simple random walk with total jump rate one, each jump uniformly chosen among the $2d$ neighbors. Its generator is $P - I = \Delta/(2d)$. Put $\tau = \inf\{t \geq 0 : X_t \notin \mathcal{D}\}$ and

$$p_{\mathcal{D}}(t, x, y) = \mathbf{P}_x(X_t = y,\ t < \tau), \qquad G_{\mathcal{D}}(x, y) = \int_0^\infty p_{\mathcal{D}}(t, x, y)\, dt.$$

The holding times have mean one and are independent of the embedded walk, so $G_{\mathcal{D}}$ also equals the usual discrete-time expected number of visits before exit.

For each starting site $x \in \mathcal{D}$, the process $|X_t|_2^2 - t$ is a martingale: averaging one jump increases squared norm by one. Bounded stopping at $t \wedge \tau$ gives

$$\mathbf{E}_x(t \wedge \tau) = \mathbf{E}_x|X_{t\wedge\tau}|_2^2 - |x|_2^2 \le d(L+1)^2 =: b\,.$$

The stopped walk lies in Λ_{L+1}, including the exit point. Monotone convergence proves $\mathbf{E}_x\tau \le b$ and almost-sure exit, uniformly in x. Markov's inequality gives $\mathbf{P}_x(\tau > 2b) \le 1/2$. The Markov property, successively at the deterministic times $2b, 4b, \ldots$, therefore gives

$$\sup_{x\in\mathcal{D}} \mathbf{P}_x(\tau > t) \le 2^{-\lfloor t/(2b)\rfloor}\,.$$

This also proves integrability of every entry of the killed semigroup.

The coordinates of X_t are independent symmetric continuous-time one-dimensional walks of rate $1/d$, by independent Poisson splitting of the jump directions. Fourier inversion of their integer-valued distributions gives, for the unrestricted transition probability $q(t,0,z)$,

$$|q(t,0,z)| \le \left(\frac{1}{2\pi}\int_{-\pi}^{\pi} e^{-(t/d)(1-\cos\theta)}d\theta\right)^d \le \min\{1, (d\pi/(8t))^{d/2}\} \le K_d(1+t)^{-d/2}\,.$$

For completeness, the one-dimensional characteristic function is $\exp[(t/d)(\cos\theta - 1)]$, obtained by summing its Poisson number of independent symmetric jumps. Fourier inversion follows by integrating $e^{i\theta(k-z)}$, whose integral is 2π if $k = z$ and zero otherwise; absolute summability permits exchanging the probability sum and integral. For the second inequality use $1-\cos\theta \ge 2\theta^2/\pi^2$ on $[-\pi,\pi]$ and enlarge the Gaussian integral to the real line. The last inequality holds with, for example, $K_d = (1+\sqrt{d\pi/8})^d$.

The killed transition matrix is symmetric. Its semigroup identity and $0 \le p_{\mathcal{D}} \le q$ now imply

$$\begin{aligned} p_{\mathcal{D}}(t,0,0) &= \sum_{y\in\mathcal{D}} p_{\mathcal{D}}(t/2,0,y)^2 \\ &\le K_d'(1+t)^{-d/2}\sum_{y\in\mathcal{D}} p_{\mathcal{D}}(t/2,0,y) \\ &\le K_d'(1+t)^{-d/2}2^{-\lfloor t/(4b)\rfloor}\,. \end{aligned}$$

For $d = 2$, integrate over $[0,4b]$ to get at most $K_2'\log(1+4b)$. On each interval $[4bk, 4b(k+1)]$, $k \ge 1$, the integral is at most $K_2'2^{-k}/k$. For $d \ge 3$, simply discard the survival factor and use $\int_0^\infty (1+t)^{-d/2}dt = 2/(d-2)$. Consequently

$$G_{\mathcal{D}}(0,0) \le \begin{cases} C\log(L+2), & d=2,\\ C_d, & d\ge 3,\end{cases} \qquad \sum_{y\in\mathcal{D}} G_{\mathcal{D}}(0,y) = \mathbf{E}_0\tau \le d(L+1)^2\,.$$

Step 3. We show that $v_N(0) \le CN\log(N+1)$ for $d = 2$ and $v_N(0) \le C_dN$ for $d \ge 3$, using the Green function of Step 2.

Let $P_{\mathcal{D}}$ be the killed discrete transition matrix on $\mathcal{D}$, with zero extension outside. Then $p_{\mathcal{D}}(t) = e^{-t(I-P_{\mathcal{D}})}$. Its integrable entries and the survival bound show that $p_{\mathcal{D}}(t) \to 0$; integrating its derivative on $[0,R]$ and passing to the limit gives $(I-P_{\mathcal{D}})G_{\mathcal{D}} = G_{\mathcal{D}}(I-P_{\mathcal{D}}) =$

I. Thus $I - P_{\mathcal{D}}$ is invertible, with inverse $G_{\mathcal{D}}$. Since $v_N = 0$ off $\mathcal{D}$, its conservation equation is exactly

$$(I - P_{\mathcal{D}})v_N = \frac{N\delta_0 - (s_N - h)}{2d}\,.$$

Multiplication by this inverse gives

$$v_N(0) = \frac{N}{2d}G_{\mathcal{D}}(0,0) - \frac{1}{2d}\sum_{y\in\mathcal{D}} G_{\mathcal{D}}(0,y)(s_N(y) - h)\,.$$

The sign and normalization follow from $\Delta = 2d(P - I)$. Since $s_N - h \geq -h$, positivity of $G_{\mathcal{D}}$ gives

$$v_N(0) \leq \frac{N}{2d}G_{\mathcal{D}}(0,0) + \frac{h}{2d}\mathbf{E}_0\tau \leq \begin{cases} CN\log(N+1), & d = 2, \\ C_dN, & d \geq 3, \end{cases}\,.$$

Indeed, $L+1 \leq A_dN^{1/d}+2$, and $N^{2/d} \leq N$ for $N \geq 1$. Furthermore $N \leq (5^d + N_0(d))n^d$ for $n \geq 1$, and in dimension two $\log(N+1) \leq C_2\log(n+1)$. Enlarge the deterministic constants so the lemma's lower bound on m implies $m \geq v_N(0)$.

Step 4. We show that $u_t^{(m,0)}(x) > 0$ for every $x \in \Lambda_n$ and every $t \geq T$, for a finite time T.

Let a_t be the ordinary parallel odometer for $h + N\delta_0$, starting from zero. At every time its height is nonnegative: a toppled site loses $2d$ only when it has at least $2d$, and all incoming chips are nonnegative. Away from zero the exact ordinary recurrence is therefore

$$a_{t+1}(x) = \min\left\{a_t(x) + 1, \left\lfloor \frac{\sum_{y\sim x} a_t(y) + h}{2d} \right\rfloor\right\}\,.$$

To verify this, the numerator is $2da_t(x) + s_t(x)$. For $0 \leq s_t(x) < 2d$ its floor is $a_t(x)$; for $s_t(x) \geq 2d$ the minimum is $a_t(x) + 1$. At zero, $a_t(0) \leq v_N(0) \leq m$. Inductively $a_t \leq u_t^{(m,0)}$: the base case follows from $a_0 = 0 \leq m\delta_0$, the origin is frozen at m, and away from zero use this recurrence, the monotone neighbor sum and floor, and $h \leq \eta$. Ordinary stabilization occurs at a finite time T, so $u_T^{(m,0)} \geq v_N > 0$ on Λ_n.

Finally, for every background with $0 \leq \eta \leq 2d-1$, the wave is nonnegative, nondecreasing in time, and at most m. Nonnegativity and the upper bound follow by induction in its floor recurrence. For time monotonicity first $u_1 \geq u_0$ (with equality at zero), and then use monotonicity of the update map. Hence positivity on Λ_n persists for every $t \geq T$. □

The constant C_d does not depend on the background, and the lemma gives no bound on T.

For a background bounded below only by an integer $\eta_{\min}$, as in Section 5, the wave (19) need not be nonnegative: if $x \neq 0$ is not adjacent to the origin and $\eta(x) < 0$, then $u_1^{(m,0)}(x) = \lfloor \eta(x)/(2d) \rfloor < 0$. Under Hypothesis 3, Step 1 of the proof of Proposition 5.1 shows that $\eta \geq 0$ almost surely, so $0 \leq \eta \leq 2d-1$ holds almost surely there, and the exponential path argument for nonnegative backgrounds in Step 4 of that proof gives only a qualitative seeding, with no polynomial rate.

Remark 1 (Sharp planar obstruction)**.** *On homogeneous height* 2 *in dimension two, the least integer order* $M_2(n)$ *whose wave covers* Λ_n *satisfies*

$$cn^2\log(n+1) \leq M_2(n) \leq Cn^2\log(n+1)\,,$$

for every $n \geq 1$. We prove the lower bound first; the upper bound is Lemma 3.1. For sufficiently large N, the radius bound of Fey, Levine and Peres [FLP10] *permits $L = \lceil 2\sqrt{N} \rceil$ and $\operatorname{supp} v_N \subseteq \Lambda_L$. The Green identity above and $s_N - 2 \leq 1$ give*

$$v_N(0) \geq \frac{N}{4} G_{\Lambda_L}(0,0) - \frac{1}{4}\mathbf{E}_0 \tau \,.$$

For the embedded discrete walk S_j, exit from Λ_L is impossible by time $2j \leq L$. The increments of $S^1 + S^2$ and $S^1 - S^2$ are independent symmetric signs, so $\mathbf{P}(S_{2j} = 0) = [\binom{2j}{j}/4^j]^2 \geq 1/(4j)$ for $j \geq 1$. The binomial inequality follows by induction: its value at $j = 1$ is $1/2$, and the ratio of successive values is $(2j+1)/(2j+2) \geq \sqrt{j/(j+1)}$. Therefore

$$v_N(0) \geq \frac{N}{4} + \frac{N}{16} H_{\lfloor L/2 \rfloor} - \frac{(L+1)^2}{2}, \qquad H_k = \sum_{j=1}^{k} \frac{1}{j} \,.$$

Taking $N = \lfloor n^2/16 \rfloor$ gives $L < n$ and $v_N(0) \geq cn^2 \log(n+1)$ for all sufficiently large n. The comparison in the opposite direction is proved by a further induction: if an integer-valued $w \geq 0$ satisfies $w(0) \geq m$ and $2 + \Delta w \leq 3$ off zero, then $u_t^{(m,0)} \leq w$ at every time. At zero this is the frozen inequality; away from zero use $\sum w + 2 \leq 4w(x) + 3$ and integrality to bound the floor by $w(x)$. Applying this to $w = v_N$ shows that every $m \leq v_N(0)$ misses Λ_n. For the finitely many remaining n, $M_2(n) \geq 1$, so reducing $c > 0$ extends the lower bound to all $n \geq 1$; Lemma 3.1 supplies the upper bound at every scale. No matching lower bound is claimed for arbitrary backgrounds: on homogeneous height 3 a 1-wave propagates to every site by induction on lattice distance.

Lemma 3.2. *The last-wave is well-defined, $\hat{M}_\eta(0) < \infty$.*

Proof. By Lemma 3.1 and Proposition 2.2, if the origin is fired a sufficient number of times, the support of the odometer contains a good cube, $Q_k \subset \mathcal{C}_\infty$. □

Lemma 3.3. *The last wave started at z is bounded by one outside the interior of the penultimate-cluster of z. Moreover, for all $t \geq 1$ and $x, z \in \mathbb{Z}^d$,*

$$u_t^{(z)}(x) \leq (1 + \tilde{u}^{(z)}(x)).$$

Proof. To reduce clutter, we take $z = 0$. We prove this by induction on t. The base case $t = 0$ follows by definition. For all $t \geq 1$, the definition also ensures it holds at the origin. So, we may take $x \neq 0$ and check:

$$\begin{aligned} u_{t+1}(x) &= \lfloor \frac{\sum_{y \sim x} u_t(y) + \eta(x)}{2d} \rfloor \\ &\leq \lfloor \frac{\sum_{y \sim x} (1 + \tilde{u}(y)) + \eta(x)}{2d} \rfloor \\ &= 1 + \tilde{u}(x) + \lfloor \frac{\sum_{y \sim x} (\tilde{u}(y) - \tilde{u}(x)) + \eta(x)}{2d} \rfloor \\ &= 1 + \tilde{u}(x) \end{aligned}$$

as $\Delta\tilde{u}(x) + \eta(x) \leq (2d - 1)$ for $x \neq 0$.

□

Lemma 3.4. *The penultimate-cluster arrival times are subadditive: for all $a, b, z \in \mathbb{Z}^d$,*

$$\tilde{T}(a, z) \le \tilde{T}(a, b) + \tilde{T}(b, z).$$

Proof. Suppose $\mathcal{P}(z) \not\subseteq \mathcal{P}(b)$, otherwise the claim is immediate. It suffices to check

$$w_t(x) := u^{(a)}_{\tilde{T}(a,b)+t}(x) \ge u^{(b)}_t(x) \qquad \text{for all } t \ge 1 \text{ and } x \in \mathcal{P}(b)^c, \tag{24}$$

which we do by induction. By Lemma 3.3, $u^{(b)}_t(x) \le 1$ if $\tilde{u}^{(b)}(x) = 0$. Also, by definition of the penultimate-cluster, we have that $\tilde{u}^{(b)}$ is zero on $\partial^\circ \mathcal{P}(b)$. Hence, for all $t \ge 1$ and $x \in \partial^\circ \mathcal{P}(b)$, $u^{(b)}_t(x) \le 1 \le w_t(x)$. Using this and the inductive hypothesis, if $x \in \mathcal{P}(b)^c \cap \{a\}^c$,

$$\begin{aligned} w_{t+1}(x) &= \lfloor \frac{\sum_{y \sim x} w_t(y) + \eta(x)}{2d} \rfloor \\ &\ge \lfloor \frac{\sum_{y \sim x} u^{(b)}_t(y) + \eta(x)}{2d} \rfloor \\ &= u^{(b)}_{t+1}(x). \end{aligned}$$

If $x \in \mathcal{P}(b)^c \cap \{a\}$, then $w_t(x) \ge 1 \ge u^{(b)}_t(x)$ as $\hat{M}_\eta(a) \ge 1$.

□

Lemma 3.5. *There is a constant $\gamma > 0$ so that for all n sufficiently large and $|x| \le n$,*

$$\mathcal{P}(x) \subset x + [-(\log n)^\gamma, (\log n)^\gamma]^d.$$

Proof. By Lemma 2.1, $1\{\tilde{u}^{(x)} > 0\}$ is a rectangle, therefore it suffices to bound the maximal side length. By Proposition 2.2, for all n sufficiently large, if any side length of the rectangle exceeds $(2k \log n)^{5/2}$ then it must overlap a good cube, contradicting stability. □

Lemma 3.6. *There are constants γ and C so that on an event of probability 1, for all n sufficiently large and $|x| \le n$,*

$$\hat{T}(x) \le C|x| + (\log n)^\gamma.$$

Proof. Let $x \in \mathbb{Z}^d$ be given. Since the sandpile is exploding, at some constant time C_η the support of the odometer will overlap the infinite cluster at a good cube near the origin, $Q_k(z)$ for some $z \in \mathbb{Z}^d$. Once this occurs the arrival time to any site is at most a constant times the chemical distance in the infinite cluster. Let $Q_k(y)$ for $y \in \mathbb{Z}^d$ be one of the nearest cubes in $\mathcal{C}_\infty$ to x. There are now two cases to consider.

If $|z - y| < (\log n)^2$, then we may choose nearby points $M_i \in \mathcal{C}_\infty$ so that $(\log n)^c \le d(z, M_i) \le (\log n)^C$ and $x, y \in \mathbf{br}(\{M_i\})$. If $|z - y| \ge (\log n)^2$, then by the chemical distance bound and the definition of $\mathcal{C}_\infty$, within at most $C|z - y| \le C(|z| + |y|)$ steps, $Q_k(y)$ will topple. Once this happens, $\mathcal{P}(x)$ is surrounded in at most $(\log n)^C$ more steps and $\mathcal{P}(x)$ will fire in at most $|\mathcal{P}(x)|$ additional steps.

□

3.3. **Convergence of the last-wave.** We show that the arrival time for the last-wave converges under rescaling.

Lemma 3.7. *There exists a constant $\gamma > 0$ so that on an event of probability 1, for all n sufficiently large and $|x| \le n$,*

$$\hat{T}(x) \le \tilde{T}(x) \le \hat{T}(x) + (\log n)^\gamma.$$

Proof. The first inequality is immediate. For the second inequality, let C and n_0 be as in Proposition 2.2. Take $n \geq n_0$ and suppose $v_t(x) > 0$ for $\min(t, |x|) > C$ and $|x| \leq n$. We must show that there is a nearby good cube $Q_k(y) \subset \mathcal{C}_\infty$ which has already fired. Once we have shown this, the same argument as in Lemma 3.6 allows us to conclude. This is, however, a consequence of Lemma 2.1 and Proposition 2.2. Any path of topplings of length at least $(2k \log n)^{5/2}$ must overlap a good cube. □

Proposition 3.1. *On an event of probability 1, the rescaled last-wave arrival times*

$$n^{-1}\hat{T}_\eta([nx])$$

converge locally uniformly to $\mathcal{N}_p$, a continuous, convex, one-homogeneous function on $\mathbb{R}^d$.

Proof. In light of Lemma 3.7 it suffices to prove the result for $\tilde{T}$. On an event of probability 1, for all large n and $x, x' \in [-n, n]^d$, Lemmas 3.6 and 3.7 give $\tilde{T}(x) \leq C|x| + (\log n)^\gamma$, and $\tilde{T}(x') \leq \tilde{T}(x) + C|x - x'| + (\log n)^\gamma$. For the second bound, at time $\hat{T}(x)$ the support of the last wave contains a path of distinct points from the origin to x. Lemma 2.1 fills the bounding rectangle of the last $(\log n)^{\gamma'}$ points of this path, where γ' is the constant of Lemma 3.5. This rectangle contains a line of a cube of $\mathcal{C}_\infty$ by Proposition 2.2, and the argument of Lemma 3.6 then reaches $\mathcal{P}(x')$. Fix $y \in \mathbb{Z}^d \setminus \{0\}$. Choose $C' > C|y|$ and K so large that $A = \{\tilde{T}(0, ny) \leq C'n + K \text{ for every } n \geq 1\}$ has $\mathbf{P}(A) > 0$. This is possible because $\tilde{T}(0, ny) \leq C|y|n + (\log(n|y|))^\gamma$ for all large n on an event of probability 1, and $(\log(n|y|))^\gamma = o(n)$. Let σ_y be the translation of the background by y, $(\sigma_y\eta)(x) = \eta(x + y)$, so that $\tilde{T}(y, ny) = \tilde{T}(0, (n-1)y) \circ \sigma_y$. The translation σ_y is mixing, hence ergodic, and the first-return map to A preserves the restriction of the probability measure to A and is ergodic. With r_j the successive return times and $S_j = r_1 + \cdots + r_j$, the induced cocycle $G_j = \tilde{T}(0, S_j y)$ is subadditive by Lemma 3.4, and $G_1 \leq C'r_1 + K$ on A. The return time r_1 is integrable on A by Kac's lemma, so the subadditive ergodic theorem applies to G_j with no integrability bound on $\tilde{T}$ itself. The second bound passes from the return times to all n: for $S_j \leq n < S_{j+1}$ it gives $|\tilde{T}(0, ny) - G_j| \leq Cr_{j+1} + (\log n)^\gamma$, and $r_{j+1}/j \to 0$ almost surely by Birkhoff's theorem applied to the integrable r_1. Hence on A the limit inferior and superior of $n^{-1}\tilde{T}(0, ny)$ are those of G_j/S_j, which converge because G_j/j converges by Kingman's theorem and S_j/j converges by Birkhoff's theorem. The limits inferior and superior of $n^{-1}\tilde{T}(0, ny)$ are invariant under σ_y: by Lemma 3.4, $\tilde{T}(0, ny)$ and $\tilde{T}(y, ny) = \tilde{T}(0, (n-1)y) \circ \sigma_y$ differ by at most the finite quantity $\tilde{T}(0, y) + \tilde{T}(y, 0)$. By ergodicity they are almost surely constant, and they agree on A, so the limit exists almost surely. The second bound replaces continuity: it supplies the comparison needed to pass from integer directions to all x, giving locally uniform convergence, and the properties of $\mathcal{N}_p$ are immediate from the scaling and microscopic subadditivity. □

Remark 2. *Convergence of the last wave may be viewed as a sort of bootstrap percolation shape theorem. Sites are initially randomly assigned two thresholds, 1 or 2. A site with threshold l becomes infected when at least l of its neighbors are infected. Infected sites remain infected. The above shows that if you start off with a large enough cluster of infected sites at the origin, every site will eventually become infected and the speed at which the infection spreads converges.*

For more on the relationship between sandpiles and bootstrap percolation, see Section 7 below. Similar shape theorems include [GM12, KS08, AMP02, CD81] *and especially* [Wil78, GG93, FL11].

4. Proof of Theorem 1.3

Let η be drawn from the event of full probability in Proposition 3.1. It suffices to show that the last-wave is a good approximation of the original process.

Proposition 4.1. *On an event of probability 1, there are constants $C_1(\eta), C_2(\eta)$ so that for all $x \in \mathbb{Z}^d$,*

$$T(x) \leq \hat{T}(x) + C_1(\eta) \tag{25}$$

and

$$\hat{T}(x) \leq T(x) + C_2(\eta), \tag{26}$$

where the last-wave arrival time $\hat{T}$ is defined in (22) *and $T(x) := \min\{t \geq 0 : v_t(x) > 0\}$.*

Proof. Recall that v_t is the parallel toppling odometer for $\eta + M_\eta \delta_0$ and recall the last-wave odometer u_t from Section 3.1.

We first check (25). Let u'_t be the $\mathcal{P}(0)$-frozen parallel toppling odometer with initial conditions $u'_0(x) = 1\{x \in \mathcal{P}(0)\}$ and $s'_0 = \eta + \Delta u'_0$ where $\mathcal{P}(0)$ is the penultimate-cluster for 0 defined in (21). By Lemma 3.3, for all $t \geq 0$, u_t is at most 1 on the inner boundary of the penultimate-cluster, $\partial^\circ \mathcal{P}(0)$. Hence, as $\eta \leq (2d-1)$, by (6) and the definition of the last-wave we have $u'_t|_{\mathcal{P}(0)^c} \geq u_t|_{\mathcal{P}(0)^c}$ for all $t \geq 0$. Also, since $\eta + M_\eta \delta_0$ is not stabilizable, for some $t_0 \geq C_\eta$, $v_{t_0}|_{\mathcal{P}(0)} \geq u'_0|_{\mathcal{P}(0)}$. Hence by (7), $v_{t+t_0}|_{\mathcal{P}(0)^c} \geq u'_t|_{\mathcal{P}(0)^c} \geq u_t|_{\mathcal{P}(0)^c}$ for all $t \geq 0$, completing the check of (25)

We now verify (26). We first consider the special case where only one firing at the origin is needed to have an infinite last-wave.

Step 1: Special case, $\hat{M}_\eta(0) = 1$. Denote the reachable sets up to time t for the last wave and exploding sandpile as

$$\begin{aligned} \mathcal{R}_t &:= \{x \in \mathbb{Z}^d : v_t(x) > 0\} \\ \hat{\mathcal{R}}_t &:= \{x \in \mathbb{Z}^d : u_t(x) > 0\}. \end{aligned} \tag{27}$$

Note, by minimality, if $\hat{M}_\eta(0) = 1$, then $\eta(0) + M_\eta(0) = 2d$. This together with $\eta \geq (2d-2)$, implies a strong regularity. Specifically, we show by induction that for all $t \geq 0$,

$$\mathcal{R}_t \subseteq \hat{\mathcal{R}}_t \tag{28}$$

and

$$\begin{aligned} |v_t(x \pm e_i \pm e_j) - v_t(x)| &\leq 1 \qquad \text{for all } x \in \mathbb{Z}^d \text{ and } e_i \neq e_j \\ |v_t(x \pm e_i) - v_t(x)| &\leq 1 \qquad \text{for all } x \in \mathbb{Z}^d \text{ and } e_i \end{aligned} \tag{29}$$

and

$$v_t(0) \geq \max_{x \in \mathbb{Z}^d} v_t(x). \tag{30}$$

The base case $t = 0$ is immediate, so suppose (28), (29), and (30) hold at $t \geq 0$ and we check $(t+1)$.

Inductive step for (30). (4) and (30) at time t imply if $x \neq 0$,

$$\begin{aligned} v_{t+1}(x) &\leq \lfloor \frac{2dv_t(0) + \eta(x)}{2d} \rfloor \\ &= v_t(0) + \lfloor \frac{\eta(x)}{2d} \rfloor \\ &= v_t(0) \\ &\leq v_{t+1}(0), \end{aligned}$$

as $\eta \leq (2d-1)$.

Inductive step for (29). We first check the origin. By (30), if $v_t(y) = v_t(0) - 1$ for some $y \sim 0$, then $\Delta v_t(0) + 2d \leq (2d-1)$. Otherwise, suppose $v_t(e_i + e_j) = v_t(0) - 1$ for some $e_i \neq e_j$ and the origin is unstable at time t. Then, $v_t(e_i) = v_t(0)$ and $v_t(e_j) = v_t(0)$. This implies, by (29) applied to e_i and e_j, that all other neighbors $y \sim (e_i + e_j)$ have a lower bound, $v_t(y) \geq v_t(0) - 1$. Hence, $\Delta v_t(e_i + e_j) \geq 2$, which implies that $(e_i + e_j)$ is unstable at time t using $\eta \geq (2d-2)$.

Now, take $x \neq 0$ and suppose for sake of contradiction

$$\Delta v_t(x) + \eta(x) \geq 2d \tag{31}$$

but for some adjacent neighbor $(x + e_j)$,

$$v_t(x + e_j) = v_t(x) - 1 \qquad \text{and} \qquad \Delta v_t(x + e_j) + \eta(x + e_j) \leq 2d - 1. \tag{32}$$

At least two other adjacent neighbors, $y' \sim x$ must satisfy $v_t(y') = v_t(x) + 1$. Indeed, otherwise by (32) and (29), $\Delta v_t(x) \leq 0$, violating our assumption (31). However, one of those neighbors must be $y' = x \pm e_i$ for some $e_i \neq e_j$. This contradicts (29) at time t since

$$v_t(x + e_i) = v_t(x + e_i + (-e_i + e_j)) + 2.$$

Next, take a diagonal neighbor, $(x + e_i + e_j)$ for $i \neq j$, and suppose for sake of contradiction (31) but

$$v_t(x + e_i + e_j) = v_t(x) - 1 \qquad \text{and} \qquad \Delta v_t(x + e_i + e_j) + \eta(x + e_i + e_j) \leq (2d - 1). \tag{33}$$

By (31) there must be at least one adjacent neighbor $y \sim x$ with $v_t(y) = v_t(x) + 1$. This neighbor cannot be $(x + e_i)$ or $(x + e_j)$ as it would contradict (29) for $(x + e_i + e_j)$. Possibly $y = (x \pm e_{i'})$ for $i' \notin \{i, j\}$, $y = x - e_i$, or $y = x - e_j$. In these cases,

$$v_t(x + e_i) = v_t(x + e_j) = v_t(x) = v_t(x + e_i + e_j) + 1. \tag{34}$$

Indeed, if not, then, say, $v_t(x + e_i) = v_t(x) - 1$, and so there must be an additional neighbor, $y' \sim x$, $y' \neq y$, with $v_t(y') = v_t(x) + 1$. But, either y' or y is diagonal to $(x + e_i)$, which contradicts (29).

Assuming (34), the same argument implies $v_t(y'') \geq v_t(x + e_i + e_j)$ for all $y'' \sim (x + e_i + e_j)$. Indeed, such a y'' with $v_t(y'') = v_t(x + e_i + e_j) - 1$ would be diagonal to either $x + e_i$ or $x + e_j$. This together with (34) shows $\Delta v_t(x + e_i + e_j) \geq 2$. which contradicts (33).

Inductive step for (28). It suffices to check this for $x \neq 0$ as $u_t(0) = 1$. Suppose for sake of contradiction there is some site x with

$$\Delta v_t(x) + \eta(x) \geq 2d \tag{35}$$

but

$$\Delta u_t(x) + \eta(x) \leq (2d-1) \tag{36}$$

and $u_t(x) = 0$. By (28), $u_t(x) = v_t(x) = 0$ and hence

$$v_t(y) \leq 1 \qquad \text{for all } y \sim x, \tag{37}$$

by (29). However, (28) and (37) imply that $\Delta v_t(x) + \eta(x) \leq \Delta u_t(x) + \eta(x) \leq (2d-1)$, contradicting (35).

Step 2: General case. In the general case, we introduce a pair of approximations to which we can apply the arguments of the special case. Let $\tilde{v}$ be the terminal (unfrozen) odometer for $\eta + (M_\eta(0) - 1)\delta_0$ and let

$$\tilde{\mathcal{P}} := \{0\} \cup \{x \in \mathbb{Z}^d : \text{ there is } y \sim x \text{ with } \tilde{v}(y) > 0\}. \tag{38}$$

Let $\tilde{w}_t$ be the $\tilde{\mathcal{P}}$-frozen parallel toppling odometer with initial conditions

$$\begin{aligned} \tilde{w}_0(x) &= 1\{x \in \tilde{\mathcal{P}}\} \\ s_0' &= \eta + \Delta\tilde{w}_0. \end{aligned} \tag{39}$$

Let w_t be the parallel toppling odometer for $s : \mathbb{Z}^d \to \mathbb{Z}$ where

$$s(x) := \begin{cases} \eta(x) & \text{if } x \notin \tilde{\mathcal{P}} \\ (2d-1) + \delta_0 & \text{otherwise.} \end{cases} \tag{40}$$

Denote the reachable sets for these processes by

$$\begin{aligned} \mathcal{R}_t &:= \{x \in \mathbb{Z}^d : w_t(x) > 0\} \\ \tilde{\mathcal{R}}_t &:= \{x \in \mathbb{Z}^d : \tilde{w}_t(x) > 0\}. \end{aligned} \tag{41}$$

The same argument as in Step 1 shows that

$$\mathcal{R}_t \subseteq \tilde{\mathcal{R}}_t. \tag{42}$$

We claim that we can conclude after proving the following inequalities,

$$\tilde{w}_t \leq u_{t+c} \tag{43}$$

$$v_t \leq (w_t + \tilde{v})\,. \tag{44}$$

Indeed, if $v_t(x) > 0$, then (44) implies $\tilde{v}(x) > 0$ or $w_t(x) > 0$. In both cases, using either (42) or (39), $\tilde{w}_t(x) > 0$ and so by (43) $u_{t+c}(x) > 0$.

Proof of (43). We know that $|\tilde{\mathcal{P}}| = C < \infty$. Hence, at some finite time $u_c(\tilde{\mathcal{P}}) \geq 1$. Monotonicity implies $\tilde{w}_t \leq u_{t+c}$.

Proof of (44). This is true at $t = 0$, we use (4) and induct,

$$\begin{aligned} v_{t+1}(x) &\le \lfloor \frac{\sum_{y\sim x} v_t(y) + \eta(x) + M_\eta \delta_0}{2d} \rfloor \\ &\le \lfloor \frac{\sum_{y\sim x} (w_t(y) + \tilde{v}(y)) + \eta(x) + M_\eta \delta_0}{2d} \rfloor \\ &= \tilde{v}(x) + \lfloor \frac{\sum_{y\sim x} w_t(y) + \sum_{y\sim x} (\tilde{v}(y) - \tilde{v}(x)) + \eta(x) + M_\eta \delta_0}{2d} \rfloor \\ &\le \tilde{v}(x) + \lfloor \frac{\sum_{y\sim x} w_t(y) + s(x)}{2d} \rfloor \\ &= \tilde{v}(x) + w_{t+1}(x). \end{aligned}$$

The third inequality used $\Delta\tilde{v}(x) + (M_\eta - 1)\delta_0 + \eta \le (2d - 1)$.

□

Proof of Theorem 1.2. Let $\mathcal{N}_p$ be the limit of Theorem 1.3 and let $\mathcal{B}(p) = \{y : \mathcal{N}_p(y) < 1\}$. The arrival time satisfies $T_\eta(x) \ge \|x\|_1$, because the odometer at time t is supported in the ℓ^1 ball of radius t; hence $\|y\|_1 \le \mathcal{N}_p(y)$ and $\mathcal{B}(p)$ lies in the ℓ^1 ball of radius one. The set $\mathcal{B}(p)$ is open and convex since $\mathcal{N}_p$ is continuous and convex, and it is nonempty since $\mathcal{N}_p(0) = 0$. The frontier of $\mathcal{B}(p)$ lies in $\{\mathcal{N}_p = 1\}$. By homogeneity $\{\mathcal{N}_p \le s\} = s\{\mathcal{N}_p \le 1\}$ for $s > 0$, so $|\mathcal{B}(p)| \ge s^d |\{\mathcal{N}_p \le 1\}|$ for every $s < 1$. Since $|\{\mathcal{N}_p \le 1\}| < \infty$, this gives $|\mathcal{B}(p)| = |\{\mathcal{N}_p \le 1\}|$, so $\{\mathcal{N}_p = 1\}$ and the frontier of $\mathcal{B}(p)$ are Lebesgue-null. Since $v_t(x) > 0$ if and only if $T_\eta(x) \le t$, we have $\{\bar{v}_t > 0\} = \{y : t^{-1} T_\eta([ty]) \le 1\}$. Let K be the closed ℓ^1 ball of radius 2. For large t both $\{\bar{v}_t > 0\}$ and $\mathcal{B}(p)$ lie in K, and by the locally uniform convergence of Theorem 1.3 on K, for each $\varepsilon > 0$ the symmetric difference $\{\bar{v}_t > 0\} \triangle \mathcal{B}(p)$ lies in $\{y \in K : |\mathcal{N}_p(y) - 1| \le \varepsilon\}$ for all large t. Hence $\limsup_t |\{\bar{v}_t > 0\} \triangle \mathcal{B}(p)| \le |\{y \in K : |\mathcal{N}_p(y) - 1| \le \varepsilon\}|$, and the right side tends to $|\{\mathcal{N}_p = 1\} \cap K| = 0$ as $\varepsilon \downarrow 0$. □

Remark 3. *Some qualitative features of the limit shape are immediate. For example, the origin is an interior point and the limit is invariant with respect to symmetries of the lattice (symmetry may fail in the periodic case introduced in Section 5). Also, a coupling with oriented percolation as in* [DL81, Mar02] *can be used to establish a 'flat-edge' for p sufficiently close to one in all dimensions. We omit the details since it is routine — see, for example, the proof of Theorem 1.2 in* [AMP02] *or Theorem 6.3 in* [GM04].

5. A generalization

5.1. **Sufficient hypotheses.** In this section assume $d \ge 2$. We give sufficient hypotheses for convergence of last-wave arrival times. Fix $\eta_{\min} \in \mathbb{Z}$ and let Ω denote the set of all bounded functions $\eta : \mathbb{Z}^d \to \mathbb{Z}$, $\eta_{\min} \le \eta \le (2d - 1)$. Endow Ω with the σ-algebra $\mathcal{F}$ generated by $\{\eta \to \eta(x) : x \in \mathbb{Z}^d\}$. Denote the action of integer translation by $T : \mathbb{Z}^d \times \Omega \to \Omega$,

$$T(y, \eta)(z) = (T_y \eta)(z) := \eta(y + z),$$

and extend this to $\mathcal{F}$ by defining $T_y E := \{T_y \eta : \eta \in E\}$. Let $\mathcal{L} \subseteq \mathbb{Z}^d$ be a *sublattice*, a finite index subgroup of $\mathbb{Z}^d$. Let $\mathbf{P}$ be a *stationary* and *ergodic* probability measure on $(\Omega, \mathcal{F})$ with respect to $\mathcal{L}$,

$$\text{Stationary: for all } E \in \mathcal{F}, y \in \mathcal{L}\text{: } \mathbf{P}(T_y E) = \mathbf{P}(E), \tag{45}$$

(46) Ergodic: $E = \bigcap_{y\in\mathcal{L}} T_y E$ implies that $\mathbf{P}(E) \in \{0,1\}$.

We refer to the probability measure $\mathbf{P}$ as *explosive* if $\mathbf{P}(\eta \text{ is explosive}) = 1$.

Stationarity and ergodicity are the weakest hypotheses under which a convergence result is proved — straightforward counterexamples can be constructed. However, we do not expect exploding sandpiles to have a limit shape without an additional independence hypothesis. At the very least, our proof will not work, as domination by a coarsened product measure was essential. Our first hypothesis is hence a quantification of ergodicity.

Hypothesis 1 (Finite range of dependence)**.** *There exists a constant $K < \infty$ such that, for every pair of finite sets $F, G \subset \mathbb{Z}^d$ satisfying $|x - y| > K$ for every $x \in F$ and $y \in G$, the sigma-fields $\sigma(\eta(x) : x \in F)$ and $\sigma(\eta(y) : y \in G)$ are independent.*

Next, fix a finite (rectangular) box with side length $k > 0$, $\mathcal{B}_k := \{x \in \mathbb{Z}^d : 1 \le x_i \le k_i\}$. The $2d$ external faces of $\mathcal{B}_k$ are

$$\mathcal{F}_i := \{x \in \bar{\mathcal{B}}_k : x_i = 0\},$$
$$\mathcal{F}_{d+i} := \{x \in \bar{\mathcal{B}}_k : x_i = k_i + 1\}.$$

Take a face, $\mathcal{F}_i$, and let $w_t : \bar{\mathcal{B}}_k \to \mathbb{N}$ be the sequence of $\mathcal{B}_k^c$-frozen parallel toppling odometers with initial conditions $w_0 = 1\{x \in \mathcal{F}_i\}$ and $s_0' = \eta + \Delta w_0$. We say that $\mathcal{B}_k(\eta)$ can be *crossed* in direction i if $w_t(x) \ge 1\{x \in \mathcal{B}_k\}$ for $t \ge |\mathcal{B}_k|$.

Hypothesis 2 (Box-crossing)**.** *For each $\delta > 0$, there is k so that*

$$\min_{j\in\mathbb{Z}^d} \mathbf{P}(\mathcal{B}_k^{(j)}(\eta) \text{ can be crossed in each direction}) > 1 - \delta,$$

where

$$\bigcup_{j\in\mathbb{Z}^d} \mathcal{B}_k^{(j)} := \bigcup_{j\in\mathbb{Z}^d} (\mathcal{B}_k + jk) = \mathbb{Z}^d$$

is a tiling of the lattice by $\mathcal{B}_k$.

If η were recurrent, Hypothesis 2 would imply η explodes with probability 1. In particular, no holes would develop in the support of the odometer. (If unfamiliar, see Section 7 below for the definition of recurrence, although this is not used here.) Our next hypothesis ensures this and more: any sufficiently long path of topplings fills its bounding rectangle.

Hypothesis 3 (Path-filling)**.** *There exists a constant $\gamma > 0$ so that on an event of probability 1, for all $n \ge n_0$, and every path of distinct points, $[-n, n]^d \supset L_m := \{z_1, \dots, z_m\}$, $z_{i+1} \sim z_i$, of length $m \ge (\log n)^\gamma$ the following holds. The L_m-frozen parallel toppling odometer with initial conditions $w_0 = 1\{x \in L_m\}$ and $s_0' = \eta + \Delta w_0$ is at least 1 on the bounding rectangle of L_m:*

$$w_t \ge 1\{\mathbf{br}(L_m)\} \qquad \text{for } t \ge m^d.$$

In order for η to have a limit shape in dimensions $d \ge 3$, we need to strengthen Hypothesis 2. The next assumption prevents low-dimensional tendrils from burrowing through good cubes (for a counterexample in three-dimensions take a large cube filled with 4 and connect each of the faces with disjoint tunnels of 5). For a point $z \in \bar{\mathcal{B}}_k$ and direction $1 \le i \le d$, consider, as before, a line passing from one side of the box to the other

(47) $$\mathcal{L}_k^{(i,z)} := \bigcup_{j=1,\dots,k_i} (z_1, \dots, z_{i-1}, j, z_{i+1}, \dots, z_d).$$

We say $\mathcal{B}_k(\eta)$ is *strongly box-crossing* if, for all $1 \le i \le d$ and $z \in \bar{\mathcal{B}}_k$, $w_{|\mathcal{B}_k|} \ge 1\{x \in \mathcal{B}_k\}$, where w_t is the odometer for $\{\mathcal{B}_k^c \cup \mathcal{L}_k^{(i,z)}\}$-frozen parallel toppling with initial conditions $w_0(x) = 1\{x \in \mathcal{L}_k^{(i,z)}\}$, $s_0' = \eta + \Delta w_0$.

Hypothesis 4 (Strongly box-crossing)**.** *For each $\delta > 0$, there is k so that, using the same notation as Hypothesis 2,*

$$\min_{j \in \mathbb{Z}^d} \mathbf{P}(\mathcal{B}_k^{(j)}(\eta) \text{ is strongly box-crossing}) > 1 - \delta.$$

We now have made enough assumptions to prove convergence of the last-wave as in Section 3.

Proposition 5.1 (Convergence of the last-wave)**.** *Under Hypotheses 1, 3, and 4, on an event of probability 1, η is explosive and the rescaled last-wave arrival times, $n^{-1}\hat{T}_\eta([nx]) := n^{-1}\min\{t > 0 : u_t([nx]) > 0\}$ converge locally uniformly to $\mathcal{N}_\eta$, a deterministic, continuous, convex, one-homogeneous function on $\mathbb{R}^d$ depending only on the law of η.*

Proof. The proof has six steps. Step 1 shows that η is nonnegative and recurrent on finite sets. Step 2 builds the cluster $\mathcal{C}$ of good boxes and bounds its geometry. Step 3 shows that a path of sites produces a full good box, and that full good boxes spread to every site. Step 4 shows that $\hat{M}_\eta(z) < \infty$ for every z and compares its arrival times with the binary infection times $a(x)$. Step 5 shows that the passage times S are integrable and subadditive and compares them with the arrival times. Step 6 takes limits along directions and identifies the limit shape.

We work in dimension $d \ge 2$, with initial state $s_0' = \eta + \Delta w_0$, so that Δw_0 is already added to η. All constants below are deterministic unless their dependence on η is indicated. Write $|x| = |x|_1$.

Step 1. We show that η is recurrent on every finite set in the sense of Section 7, and that $\eta(x) \ge 0$ for every x. To see this, let V be finite and fire its external boundary once, freezing the complement of V. Since $\eta \le 2d - 1$, each site of V fires at most once. If the procedure does not fill V, then its nonempty terminal untoppled set E satisfies

$$\eta(x) + 2d - \#\{y \in E : y \sim x\} \le 2d - 1 \qquad (x \in E)\,.$$

No process whose seeds avoid E and whose odometers are at most one can infect E. Indeed, at a first infection time in E, every site of E was still uninfected one round earlier, while even firing every neighbor outside E once leaves the displayed height below $2d$.

Choose a large integer R with $V \subset (-R, R)^d$ and take the monotone edge path on the boundary of $[-R, R]^d$ from $(-R, \dots, -R)$ to $(R, \dots, R)$, changing coordinates successively. It avoids E, has $2dR+1$ distinct vertices, and has bounding rectangle $[-R, R]^d$. For sufficiently large R it satisfies the threshold in Hypothesis 3, a contradiction. Thus V is recurrent. Taking $V = \{x\}$ proves $\eta(x) \ge 0$ for every x.

Moreover, once the whole external boundary of every finite V is infected, V fills in at most $|V|$ further parallel steps: before completion every nonstationary round adds at least one new site.

Step 2. We show that the good boxes contain a unique infinite cluster $\mathcal{C}$, that every component of $\mathcal{C}^c$ meeting $[-n, n]^d$ has size at most h_n for all large n, and that the chemical distance $d_{\mathcal{C}}$ is at most a constant times the ℓ^1 distance. Let the positive integer side vector $k = (k_1, \dots, k_d)$ be chosen from Hypothesis 4, with failure probability δ to be fixed below.

Put $\Lambda = \operatorname{diag}(k_1, \dots, k_d)$, $\mathcal{B}_k^{(j)} = \mathcal{B}_k + \Lambda j$ for $j \in \mathbb{Z}^d$, and $V_k = |\mathcal{B}_k|$. The deadline in the definition of strong box-crossing is V_k.

Call $\mathcal{B}_k^{(j)}$ good when $\mathcal{B}_k^{(j)}(\eta)$ is strongly box-crossing, and let X_j indicate this event. The event $X_j = 1$ depends only on $\eta|_{\mathcal{B}_k^{(j)}}$. The finite-set independence in Hypothesis 1 extends to sigma-fields of separated collections by a monotone-class argument. For example, $r = \lceil K \rceil + 2$ bounds its dependence neighborhood in the coarse ℓ_∞ metric, independently of k.

Declare a coarse nearest-neighbor edge $\{i, j\}$ admissible when $X_i X_j = 1$. These edge indicators have a fixed finite dependence neighborhood and marginal at least $1 - 2\delta$. The domination theorem of Liggett, Schonmann and Stacey [LSS97] therefore couples them above independent bond percolation with parameter p_0 arbitrarily close to one as δ decreases. This applies the same domination theorem to a bounded-degree index graph. Alternatively, encode the d positively oriented edges based at i as the sites $(di_1 + s, i_2, \dots, i_d)$ with $0 \le s < d$, and apply the lattice version. The encoded dependence neighborhood is bounded in terms of d and K. Consequently δ can be fixed before k is chosen.

We now give the estimates that transfer the percolation bounds to the entire good-box cluster. Denote the infinite cluster of the independent bond process by I. Call a coarse vertex defective when at least one incident independent bond is closed. Put $\Delta_* = 3^d - 1$ and choose $p_0 > p_c(d)$ so close to one that $\alpha = \Delta_*^2 [2d(1 - p_0)]^{1/3^d} < \frac{1}{4}$.

By independent-bond infinite-cluster uniqueness, I is almost surely the unique infinite cluster of the coupled bond process. For any prescribed set of m vertices, one of its residue classes modulo 3 has at least $m/3^d$ vertices. Their incident bond sets are disjoint. Thus the probability that every vertex of the prescribed set is defective is at most $[2d(1 - p_0)]^{m/3^d}$.

There are at most Δ_*^{2m} star-connected sets of size m through a specified vertex, by a depth-first traversal of a spanning tree. Hence defective star-components are finite and have exponentially decaying volume tails.

Every connected component H of I^c has connected complement: every other component attaches to I. Its inner boundary is star-connected and defective. The lattice boundary fact used in Lemma 5.2 of [Mat08] therefore makes this boundary finite. A finite boundary cannot separate two infinite subsets here: the complement of a sufficiently large finite box in $\mathbb{Z}^d$ is connected for $d \ge 2$. Since the complement of H contains I, it follows that H is finite.

For its containing defective star-component G, an elementary coordinate-line count gives

$$|H| \le |\partial^\circ H|^2 \le |G|^2 .$$

Indeed, choose distinct coordinate directions e_1, e_2. Each e_1-line meeting H has a last site in H on its positive ray, giving a distinct site of $\partial^\circ H$. For each first-coordinate value attained in H, choose a site with that value and follow its positive e_2-ray to its last site in H; the first coordinate is preserved, so these boundary sites are distinct. A site of H is determined by its e_1-line and its first coordinate, which gives the product bound. Also $H \subseteq \mathbf{br}(G)$. If $z \in H$ and $|G| = m$, a vertex of G lies within distance m of z in the ℓ_∞ metric. The preceding animal count, now summed over at most $(2m + 1)^d$ possible anchors, proves

$$\mathbf{P}\big(z \in I^c,\ |H(z)| \ge s\big) \le C \exp(-c\sqrt{s}) , \qquad s \ge 1 .$$

The bound is for the holes $H(z)$, which may contain finite open islands.

There is a unique infinite component $\mathcal{C}$ of the good vertices $\{X = 1\}$. Indeed it contains I, while any other infinite component would be confined to one of the finite components of

I^c. Every component of $\mathcal{C}^c$ is contained in one such hole. For each coarse vertex a, let $\pi(a)$ be its nearest vertex of $\mathcal{C}$, with translation-covariant lexicographic tie breaking. Let $d_{\mathcal{C}}$ be graph distance in $\mathcal{C}$.

Choose p_0 depending only on d with $\alpha \le 1/8$ in the preceding animal estimate, so that the constants in the resulting hole tail depend only on d. Write $L_n = (\log(n+2))^2$ and $h_n = \beta L_n$, where the deterministic $\beta = \beta(d) \ge 1$ is large enough that $c\sqrt{\beta} > d+2$, and $c > 0$ denotes the exponent in the hole tail. Put $W_n = [-4n, 4n]^d$. A union over the vertices of W_n gives

$$\mathbf{P}\{\text{some } I^c \text{ hole meeting } W_n \text{ has size} > h_n\} \le Cn^d \exp(-c\sqrt{\beta L_n})\,.$$

The all-pair Antal–Pisztora estimate [AP96] and a union over pairs in W_n give

$$\mathbf{P}\{\exists a, b \in I \cap W_n :\ |a-b| \ge L_n,\ d_I(a,b) > \rho|a-b|\} \le Cn^{2d} \exp(-c' L_n)\,.$$

Both bounds are summable: the first is $O(n^{d-c\sqrt{\beta}})$ by the choice of β. Hence, almost surely, both properties hold for all sufficiently large n, with a random threshold and deterministic β and ρ.

For $i \in \mathcal{C} \cap [-n, n]^d$, follow a simple path in $\mathcal{C}$ to its first vertex $a \in I$. When $i \notin I$, this initial segment lies in $H(i)$ and has length at most $|H(i)|+1 \le h_n+1$. Take $a = i$ when $i \in I$, and choose $b \in I$ similarly for $j \in \mathcal{C} \cap [-n, n]^d$. If $|a-b| \ge L_n$, then apply the Antal–Pisztora bound directly. Otherwise translate a by $\lceil 10h_n \rceil e_1$ and choose $w \in I$ within $h_n + 1$ steps of that translate; the hole bound supplies w. For all large n, $a, b, w \in W_n$, $|a-w| \ge 9h_n - 1$ and $|b-w| \ge 9h_n - 1 - L_n$, so both exceed L_n, and $|a-w| + |w-b| \le Ch_n$. The two Antal–Pisztora legs and the endpoint paths therefore give, with a deterministic constant A,

$$d_{\mathcal{C}}(i,j) \le A(|i-j| + h_n) \qquad (i, j \in \mathcal{C} \cap [-n, n]^d)\,.$$

The same hole bound places each nearest-cluster projection $\pi(a)$ within $h_n + 1$ of a. Its connecting paths and the two-leg detour stay in W_n for all large n. Applying the same argument there and adjusting A gives

$$\begin{aligned}
&\text{every component of } \mathcal{C}^c \text{ meeting } [-n, n]^d \text{ has size at most } h_n,\\
&|\pi(a) - a| \le h_n + 1,\\
&d_{\mathcal{C}}(\pi(a), \pi(b)) \le A(|a-b| + h_n) \qquad (a, b \in [-n, n]^d)\,.
\end{aligned}$$

Finally let $x, y \in \mathcal{C}$, $|x| \le n$, and $|x-y| \ge (\log n)^2$, and put $N = \max\{n, |y|\}$. The preceding additive estimate at scale N applies to x and y. If $|y| \le 2n$, then $h_N \le C\beta(\log n)^2 \le C\beta|x-y|$. If $|y| > 2n$, then $|x-y| \ge N/2$ and $h_N \le C\beta N \le C\beta|x-y|$ for all large n. Thus, almost surely, $d_{\mathcal{C}}(x,y) \le C'|x-y|$ with a deterministic C', for all n above a random threshold. The lower bound $d_{\mathcal{C}}(x,y) \ge |x-y|$ follows from the nearest-neighbor graph metric.

We also need a moment bound, not merely an eventual almost-sure estimate. For fixed a, b and $R \ge |a-b|+2$, require all I^c holes meeting $a + [-40R, 40R]^d$ to have size at most R, and require the Antal–Pisztora bound for all pairs in that box separated by at least R. The failure probability is at most $CR^{2d}\exp(-c\sqrt{R})$. On this event the projections of a and b and their connecting vertices in I are within $O(R)$ of a. A vertex of I near $a + 10Re_1$ supplies the same two-leg connection as above. Hence, for deterministic constants A_*, C and c,

$$\mathbf{P}\{d_{\mathcal{C}}(\pi(a), \pi(b)) > A_* R\} \le CR^{2d}\exp(-c\sqrt{R})\,, \qquad R \ge |a-b| + 2\,.$$

The constants A_*, C and c do not depend on a or b. Summing the tail over increasing R gives $\mathbf{E}d_{\mathcal{C}}(\pi(a),\pi(b)) < \infty$.

All constants in this geometry depend only on d, K, the chosen domination parameter and, after conversion to physical coordinates, the fixed side vector k. The auxiliary coupling is used only to prove estimates: $\mathcal{C}$ and π are functions of η itself.

Step 3. We show that a process containing one full good box of $\mathcal{C}$ eventually infects every site, and that a path of ℓ_n sites produces a full good box of $\mathcal{C}$ nearby. A fully infected good box infects every neighboring good box within V_k additional steps. A shared external face contains one of the line seeds in the strong crossing definition: choose a direction tangent to that face, which is possible since $d \geq 2$. Thus every process containing one full box of $\mathcal{C}$ eventually contains all of its boxes. It then infects the boundary of every finite component of their complement, and recurrence fills each such component. It consequently reaches every physical site.

We now turn a path into a full good box. Let γ be the exponent in Hypothesis 3, and put $q = \max\{\gamma + 1, 3d\}$ and $\ell_n = \lceil (\log(n+2))^q \rceil$. Increase the random cutoff so that all preceding estimates and Hypothesis 3 apply in the required fixed enlargements of $[-n,n]^d$.

Suppose that at time t a process contains a self-avoiding path of ℓ_n sites ending at $x \in [-n,n]^d$. The path lies in $[-2n,2n]^d$ for large n. Comparison with the frozen odometer of the path and Hypothesis 3 fill its bounding rectangle by time $t + \ell_n^d$. That rectangle has a coordinate side containing at least $\ell_n^{1/d}$ sites. A line along this side contains more than h_{2n} consecutive full coarse-box line segments, since $q/d \geq 3 > 5/2$ and k is fixed. They cannot all lie in one component of $\mathcal{C}^c$. Therefore one is a full line in a box of $\mathcal{C}$, which is completely infected by time $t + \ell_n^d + V_k$. This box is within physical distance $O(\ell_n + |k|)$ of x. Call this conclusion the path-to-box statement.

The comparison just used applies to the floor waves, to ordinary legal parallel toppling, and to binary frozen processes. It is an induction on time: when the larger odometer is already at least one there is nothing to prove, and otherwise the required neighbor chips are dominated.

Step 4. We show that $\hat{M}_\eta(z) < \infty$ for every z, and that the last-wave arrival times differ from the binary infection times $a(x)$ by finite constants. For $0 \leq \eta \leq 2d-1$, the order-m wave is nonnegative, nondecreasing in time, bounded above by m, and has finite propagation speed. These facts follow directly by induction in (19). With $m = (2d)^R$ that recurrence gives

$$u_t^{(m,z)}(y) \geq (2d)^{R-|y-z|} \quad \text{when } |y-z| \leq R,\ t \geq |y-z|\,.$$

Choose R so that the ball of radius R about z covers a fixed box of $\mathcal{C}$. The resulting wave cannot stabilize, by propagation along $\mathcal{C}$. This proves $\hat{M}_\eta(z) < \infty$ for every z, on the common probability-one event.

This also proves ordinary explosion, but it is necessary to respect the different clocks. Add $2dm$ chips at the origin. It then topples at least m times during its first m rounds. In the legal process with the origin frozen at $m = (2d)^R$, once distance $r-1$ has odometer at least $(2d)^{R-r+1}$, every site at distance r attains $(2d)^{R-r}$ within at most $(2d)^{R-r}$ more rounds, using one predecessor and $\eta \geq 0$. Hence the whole radius-R ball fills in at most $\sum_{r=1}^{R}(2d)^{R-r} \leq m$ rounds of that frozen legal process. Ordinary toppling dominates it after time m, by (7). It reaches the good box and cannot stabilize. No ordinary-odometer estimate from a compact-growth theorem is used.

Fix the last wave from zero. Its penultimate wave has finite support: it stabilizes at a finite time and propagation speed is finite. The last wave has infinite support, since a nondecreasing, integer-valued wave bounded by its finite order would stabilize in a finite time if its support were finite. Its positive support is connected to its source. It therefore contains arbitrarily long self-avoiding paths, each present at some finite time. To apply the path-to-box statement to a path based at the source and ending far away, take a window radius n larger than its endpoint distance and its polylogarithmic terminal segment. As the available path lengths grow without bound, n can also exceed the almost-sure random cutoff while ℓ_n remains shorter than the path. The path-to-box statement then makes the last wave reach a full box of $\mathcal{C}$. It follows that the last wave reaches every site.

For a finite set Q, write Φ_t^Q for the binary infection process with permanent seed Q, in which an uninfected x is infected when at least $2d - \eta(x)$ neighbors are infected. This is exactly the Q-frozen parallel toppling process, whose odometer is at most one. Take $Q_0 = \mathcal{B}_k^{(\pi(0))}$ and write $a(x) = \min\{t \geq 0 : x \in \Phi_t^{Q_0}\}$. The previous propagation argument shows $a(x) < \infty$ for every x. The last wave eventually covers Q_0, so for a finite $c_1(\eta)$,

$$\hat{T}(x) \leq a(x) + c_1(\eta) + 1\,.$$

Conversely, $\Phi_t^{Q_0}$ eventually covers the finite set $\mathcal{P}(0)$. Lemma 3.3 and its proof use only $0 \leq \eta \leq 2d - 1$: the last wave is at most one outside $\mathcal{P}(0)$ and on its inner boundary. Once $\Phi_t^{Q_0}$ covers $\mathcal{P}(0)$, the same exterior induction as in Lemma 3.4 dominates the last wave outside that set; inside it, infection has already occurred. Consequently, for another finite $c_2(\eta)$,

$$a(x) \leq \hat{T}(x) + c_2(\eta) \qquad (x \in \mathbb{Z}^d)\,.$$

Neither time shift is required to be integrable.

Step 5. We show that $\mathbf{E}S(a,b) < \infty$ for every pair of coarse sites, that S is subadditive, and that $\hat{T}(x)$ and $S(0, b(x))$ differ by $o(n)$ uniformly on $[-n,n]^d$. For coarse sites a, b define

$$S(a,b) = \min\{t \geq 0 : \mathcal{B}_k^{(\pi(b))} \subseteq \Phi_t^{\mathcal{B}_k^{(\pi(a))}}\}\,.$$

Restarting from an infected target box and using monotonicity gives

$$S(a,c) \leq S(a,b) + S(b,c), \qquad S(a,b) \leq V_k\, d_{\mathcal{C}}(\pi(a), \pi(b))\,.$$

The quantitative geometry therefore gives $\mathbf{E}S(a,b) < \infty$ for every fixed pair. It also gives, uniformly for $a, b \in [-n,n]^d$ and all sufficiently large n,

$$|S(0,a) - S(0,b)| \leq V_k A(|a-b| + h_n)\,.$$

Let $b(x)$ be the index of the physical box containing x. We compare the completion time $S(0, b(x))$ with the point time $a(x)$. If $b(x) \in \mathcal{C}$, then $a(x) \leq S(0, b(x))$. Otherwise let H be the component of $\mathcal{C}^c$ containing $b(x)$. Its size is at most h_n, after a fixed enlargement of the window. The projection $\pi(b(x))$ is adjacent to H, and every other boundary vertex of H is within $h_n + 1$ coarse steps of it. The chemical-distance estimate and propagation infect all boundary boxes in at most $CV_k h_n$ more steps. Recurrence then fills $\bigcup_{j \in H} \mathcal{B}_k^{(j)}$ in at most $V_k h_n$ further steps. Thus

$$a(x) \leq S(0, b(x)) + Ch_n\,.$$

For the reverse comparison, trace a path of first infections from Q_0 to x at time $a(x)$; infection times strictly decrease when this path is traced backwards. If it has at least ℓ_n vertices, then

apply the path-to-box statement to its last ℓ_n vertices. The resulting full good box is within $O(\ell_n + |k|)$ of x. Propagation through $\mathcal{C}$ reaches $\mathcal{B}_k^{(\pi(b(x)))}$ in at most $C(\ell_n + h_n)$ further steps. If the path is shorter, then x is already within ℓ_n of Q_0, and the same chemical bound applies directly. Hence, uniformly for $x \in [-n, n]^d$ and all sufficiently large n,

$$S(0, b(x)) \le a(x) + C\ell_n^d .$$

Combining these estimates gives the required comparison

$$\sup_{x\in[-n,n]^d} |\hat{T}(x) - S(0, b(x))| \le c_1(\eta) + c_2(\eta) + 1 + C\ell_n^d = o(n) .$$

Step 6. We show that the directional limits of S exist and extend to a continuous, convex, positively one-homogeneous function μ on $\mathbb{R}^d$, and that $n^{-1}\hat{T}_\eta([nx])$ converges locally uniformly to $\mathcal{N}_\eta(x) = \mu(\Lambda^{-1}x)$. Let $\Gamma = \{z \in \mathbb{Z}^d : \Lambda z \in \mathcal{L}\}$. It is a finite-index subgroup. The kernel S is covariant under translations by Λz for $z \in \Gamma$, and is a measurable function of η. For each nonzero $z \in \Gamma$, the transformation $T_{\Lambda z}$ is mixing, hence ergodic: separated finite cylinder events are independent by Hypothesis 1, and approximation by cylinders proves mixing for all events.

The subadditive ergodic theorem applies to $S(0, nz)$, and its hypotheses have been verified, including integrability. On a common probability-one event, for every $z \in \Gamma$, $S(0, nz)/n \to \mu(z)$ almost surely and in L^1, where $\mu(z)$ is deterministic. Integer homogeneity follows by subsequences. Taking expectations in $S(0, n(z + w)) \le S(0, nz) + S(nz, n(z + w))$, using stationarity and L^1 convergence, proves $\mu(z + w) \le \mu(z) + \mu(w)$. The uniform modulus above gives $|\mu(z) - \mu(w)| \le V_k A|z - w|$. Thus rational homogeneity and continuity extend μ uniquely to a continuous, convex, positively one-homogeneous function on $\mathbb{R}^d$. Since Γ has finite index, its rational span is all of $\mathbb{Q}^d$. For every fixed $L < \infty$, apply the directional limits on a finite net in $[-L, L]^d \cap \mathbb{Q}^d$. Integer subsequences from Γ give convergence at the net points, and the uniform modulus controls every coarse a with $|a|_\infty \le Ln$ by a nearby net point. Sending the net mesh to zero proves $\sup_{|a|_\infty \le Ln} |n^{-1}S(0, a) - \mu(a/n)| \to 0$ for each L.

The projection error is $o(n)$ uniformly on bounded rescaled windows. Finite propagation speed gives $S(0, nz) \ge \mathrm{dist}_1(\mathcal{B}_k^{(\pi(0))}, \mathcal{B}_k^{(\pi(nz))})$, so $\mu(z) \ge |\Lambda z|$. The chemical estimate gives $\mu(z) \le C|\Lambda z|$.

Define $\mathcal{N}_\eta(x) = \mu(\Lambda^{-1}x)$. The point-to-box comparison now proves $n^{-1}\hat{T}_\eta([nx]) \to \mathcal{N}_\eta(x)$ locally uniformly, with

$$|x|_1 \le \mathcal{N}_\eta(x) \le C|x|_1 .$$

This deterministic limit depends only on the law of η. Different choices of the auxiliary tiling give the same limit, since each is compared with the same last-wave arrival times $\hat{T}_\eta$. □

If additionally $\eta \ge (2d-2)$, then the argument given in Section 4 implies that the exploding sandpile is close to the last-wave and hence converges. Simulations indicate $\eta \ge (2d - 2)$ is not necessary, however, we have not found an alternative condition and are forced to assume this:

Hypothesis 5 (Wave-approximation)**.** *Suppose η is explosive, let u_t denote the last-wave, v_t the parallel toppling odometer for $\eta + M_\eta\delta_0$, and $\hat{T}_\eta$, T_η the respective arrival times. On an event of probability 1,*

$$\sup_{x\in[-n,n]^d} |\hat{T}_\eta(x) - T_\eta(x)| = o(n).$$

Theorem 5.1 (Convergence of the exploding sandpile)**.** *Under the assumptions in Proposition 5.1 and Hypothesis 5, on an event of probability 1, η is explosive and the rescaled arrival times, $n^{-1}T_\eta([nx]) := n^{-1}\min\{t > 0 : v_t([nx]) > 0\}$ converge locally uniformly to $\mathcal{N}_\eta$, a deterministic, continuous, convex, one-homogeneous function on $\mathbb{R}^d$ depending only on the law of η.*

Proof. Immediate from Hypothesis 5 and Proposition 5.1. □

Remark 4. *Box-crossing with probability 1 implies η is recurrent (see Section 7 if unfamiliar). However, not every recurrent sandpile is explosive — take $\eta = (2d-2)$ and use* [FLP10] *— and not every exploding sandpile has a recurrent initial condition — see Section 6.*

5.2. **Examples satisfying the hypotheses.** The simplest way to ensure Hypotheses 3 and 5 is to take $\eta \geq (2d-2)$. A random background can be built which satisfies the rest of the hypotheses using a *Bernoulli cloud*, see Figure 4. Take $p > 0$, fix a finite nonempty set of points, $\varnothing \neq \mathcal{S} \subset \mathbb{Z}^d$ (say a triangle, circle, or a line), and independently sample a uniform random variable at each site on the lattice, $\{U_j\}_{j\in\mathbb{Z}^d}$. Then, let

$$\eta(x) := \begin{cases} (2d-1) & \text{if there exists } j \in \mathbb{Z}^d \text{ such that } U_j < p \text{ and } x \in \{\mathcal{S}+j\} \\ (2d-2) & \text{otherwise.} \end{cases} \tag{48}$$

The field $\eta(x)$ depends only on the uniforms U_{x-s}, $s \in \mathcal{S}$. If $|x-y| > \max_{s,t\in\mathcal{S}}|s-t|$, then $x-s \neq y-t$ for all $s,t \in \mathcal{S}$, so the fields on separated finite sets depend on disjoint families of independent uniforms and are independent. This is Hypothesis 1 with $K = \max_{s,t\in\mathcal{S}}|s-t|$. Choose $s \in \mathcal{S}$. The indicators $1\{U_{x-s} < p\}$, $x \in \mathbb{Z}^d$, are independent with mean $\min\{p,1\} > 0$, and each is bounded above by $1\{\eta(x) = 2d-1\}$. In a side-k cube, a union bound over its dk^{d-1} coordinate lines shows that some line has no high site with probability at most $dk^{d-1}(1-\min\{p,1\})^k$, which tends to zero as $k \to \infty$. On the event that every line has a high site, the line-filling argument of Proposition 2.1 fills the box by its volume deadline $|\mathcal{B}_k|$. Hence the box is strongly box-crossing with probability tending to one, which verifies Hypothesis 4.

Another family of examples is the *random checkerboard*. Fix a box $\mathcal{B}$ which tiles the lattice, $\mathbb{Z}^d = \bigcup_{j\in\mathbb{Z}^d}\mathcal{B}_j$. Take functions $\zeta_1, \ldots, \zeta_m$, defined on the box, $\zeta_i : \mathcal{B} \to \{(2d-2),(2d-1)\}$. Suppose further that $\mathcal{B}(\zeta_1)$ contains at least one site with $(2d-1)$ chips along every straight line. Let $\{Y_j\}_{j\in\mathbb{Z}^d}$ be a field of i.i.d. random variables, where $P(Y_j = i) = p_i$, for $i = 1, \ldots, m$. Then, let

$$\eta(x) := \zeta_i(z(x)) \qquad \text{if } Y_{j(x)} = i, \tag{49}$$

where $z(x) \in \mathcal{B}$ is the position of x in its tiled box, $\mathcal{B}_{j(x)}$. Finite range of dependence follows from the independent basic-tile choices. For a coarse box comprising L basic tiles in each coordinate, a fixed coordinate line traverses L independent tile choices, and it misses type 1 in all of them with probability $(1-p_1)^L$. There are $O(L^{d-1})$ such lines. A union bound and the line-filling argument of Proposition 2.1 show that, when $p_1 > 0$, the coarse box is strongly box-crossing by time $|\mathcal{B}_k|$ with probability tending to one as $L \to \infty$.

The random checkerboard includes the degenerate case $p_1 = 1$, where η is a periodic copy of ζ_1. See Table 1 for pictures. Theorem 4.2 of [FLP10] gives a sufficient condition for robustness: there is a sequence of enclosing rectangular boundaries on which every height




FIGURE 4. Initial random backgrounds and computed limit shapes. The random backgrounds are built from Bernoulli clouds of the indicated point sets. Blue is 2 chips and red is 3 chips.






$[3]$	$\begin{bmatrix} 2 & 3 \\ 3 & 2 \end{bmatrix}$	$\begin{bmatrix} 3 & 2 & 3 \\ 2 & 3 & 3 \\ 3 & 2 & 3 \end{bmatrix}$	$\begin{bmatrix} 2 & 2 & 3 & 3 \\ 3 & 2 & 3 & 2 \\ 2 & 3 & 2 & 2 \\ 2 & 2 & 3 & 2 \end{bmatrix}$	$\begin{bmatrix} 3 & 2 & 3 & 3 & 3 \\ 2 & 2 & 3 & 3 & 3 \\ 2 & 2 & 3 & 2 & 2 \\ 3 & 3 & 2 & 2 & 2 \\ 3 & 3 & 3 & 2 & 3 \end{bmatrix}$
				

TABLE 1. Computed limited shapes of periodic, checkerboard backgrounds of the indicated box.

is at most $2d-2$. However, it is possible to build random (and periodic) checkerboard, exploding sandpiles with $\eta \not\geq (2d-2)$. One could then proceed in an ad hoc manner to check the hypotheses. However, we have not found a general recipe in this case. The counterexample in Section 6 uses $\eta \not\geq (2d-2)$.

6. Failure of convergence

In this section we construct a family of exploding sandpiles which fail to have a limit shape. As the construction indicates, the counterexample is stable: it can be random or periodic.

Theorem 6.1. *For each dimension $d \geq 2$, there are explosive backgrounds $\eta \not\geq (2d-2)$ on $\mathbb{Z}^d$ which fail to have a limit shape; the first arrival times $\mathcal{T}(n) := \min\{t > 0 : v_t(ne_1) > 0\}$*




FIGURE 5. The counterexample from Theorem 6.1 for $d = 2$ and $d = 3$.

do not converge,

$$\limsup_{n\to\infty} n^{-1}\mathcal{T}(n) \geq 3/2 \tag{50}$$

and

$$\liminf_{n\to\infty} n^{-1}\mathcal{T}(n) = 1. \tag{51}$$

We explicitly demonstrate a family of checkerboard backgrounds which are explosive but do not have a limit shape. Our counterexample is essentially a two-dimensional one. After constructing it in two dimensions, we embed it into higher-dimensions and show failure of convergence by comparison with the two-dimensional counterexample.

6.1. **Proof of Theorem 6.1 for $d = 2$.** We use the notation of Section 5. Let $\mathcal{B} := \{x \in \mathbb{Z}^2 : 0 \leq x \leq 3\}$ denote a box of side length 4 and take $\zeta_1, \zeta_2 : \mathcal{B} \to \{1, 2, 3\}$ as

$$\zeta_1 := \begin{bmatrix} 1 & 3 & 3 & 1 \\ 3 & 3 & 3 & 3 \\ 3 & 3 & 3 & 3 \\ 1 & 3 & 3 & 1 \end{bmatrix} \qquad \zeta_2 := \begin{bmatrix} 1 & 3 & 3 & 1 \\ 3 & 2 & 3 & 3 \\ 3 & 3 & 3 & 3 \\ 1 & 3 & 3 & 1 \end{bmatrix},$$

where the lower-left corner of the box is $(0, 0)$ and left-to-right and down-to-up are increasing coordinates. Let η be an arbitrary tiling of ζ_1, ζ_2; for example, η could be a sample from the random checkerboard measure. Fix coordinates so that

$$\eta(x_1, x_2) = \zeta_i(x_1 \bmod 4, x_2 \bmod 4).$$

Let v_t be the sequence of parallel toppling odometers for $s_0 = \eta + 3\delta_0$. We first verify (51).

Step 1: Proof of (51). We show for all $n \geq 0$

$$(4n + 1) \leq \mathcal{T}(4n + 1) \leq 4n + 4. \tag{52}$$

By inspection, $v_1(0) = 1$ and $v_2(e_1) = v_2(e_2) = 1$. Now, take $n \geq 1$ and observe that there is a line of 3s connecting e_2 to $(e_2 + (4n + 1)e_1)$. Thus, $v_{4n+3}(e_2 + (4n + 1)e_1) = 1$ and

$v_{4n+4}((4n+1)e_1) = 1$. The lower bound is immediate from $\eta \leq (2d-1)$ — a site can fire only if a neighbor has fired previously.

Step 2: η is explosive. By Theorem 2.8 in [FMR09], it suffices to construct a toppling procedure which transforms η into a configuration η' which is not stabilizable in such a way that only sites with at least $2d$ chips are toppled. Start by toppling the origin, then every 3, then every 2, then the 2×2 box of 1s containing the origin,

$$\eta' := \eta + \Delta\left(1\{\eta \geq 2\} + \delta_0 + \delta_{-e_1} + \delta_{-e_2} + \delta_{-e_1-e_2}\right).$$

(It may be checked that this is indeed a legal toppling procedure in the sense of [FMR09].) The resulting configuration is (away from the origin) a tiling of $\zeta_1', \zeta_2' : \mathcal{B} \to \{2,3\}$,

$$\zeta_1' := \begin{bmatrix} 3 & 2 & 2 & 3 \\ 2 & 3 & 3 & 2 \\ 2 & 3 & 3 & 2 \\ 3 & 2 & 2 & 3 \end{bmatrix} \qquad \zeta_2' := \begin{bmatrix} 3 & 2 & 2 & 3 \\ 2 & 2 & 3 & 2 \\ 2 & 3 & 3 & 2 \\ 3 & 2 & 2 & 3 \end{bmatrix}. \tag{53}$$

Remark 5. *The reason why convergence fails for this counterexample is that the limit shape of the explosive background η' is not a diamond. See Figure 5. When $d = 2$ the limit shape is a regular octagon with boundary* $\max(|x - y/3|, |x + y/3|, |x/3 - y|, |x/3 + y|)$*, but we will not prove this.*

Both ζ_1' and ζ_2' are box-crossing, so it remains to check that we can construct a sequence of firings to the outer face of a box away from the origin. The box containing the origin is at least

$$\eta'|_{[0,3]^2} \geq \begin{bmatrix} 3 & 2 & 2 & 3 \\ 2 & 2 & 3 & 2 \\ 3 & 3 & 3 & 2 \\ 1 & 3 & 2 & 3 \end{bmatrix}.$$

From this we see that $3\delta_0 + \eta'$ is not stabilizable — in a finite number of steps every site in $[0,3]^2$ will fire.

Step 3: Reductions. Before proving (50), we make several reductions. We seek to lower bound $\mathcal{T}$, therefore, we are free to add to η as this will only decrease the arrival time. First, we may suppose all of the boxes are ζ_1 rather then ζ_2.

We then increment the background so as to reduce to a sandpile on a cylinder, $\mathcal{C} := \{x \in \mathbb{Z}^2 : x_1 \geq 0, 3 \geq x_2 \geq 0\}$. Specifically, let $\zeta : \mathcal{C} \to \{1,3,4\}$,

$$\zeta := \begin{bmatrix} 1 & 3 & 3 & 1 & & & \\ 3 & 3 & 3 & 3 & \zeta_1 & \zeta_1 & \cdots \\ 3 & 3 & 3 & 3 & & & \\ 4 & 3 & 3 & 1 & & & \end{bmatrix}$$

with the origin, $(0,0)$, on the bottom left with left-to-right, down-to-up increasing. To periodically tile by ζ: set for $x_1 \geq 0$,

$$\hat{\eta}(x_1, x_2) := \zeta(x_1, x_2 \bmod 4)$$

and for $x_1 < 0$,

$$\hat{\eta}(x_1, x_2) := \hat{\eta}(-(x_1+1), x_2).$$

Note that $\hat{\eta} \geq \eta$.

The structure of $\hat{\eta}$ allows us to reduce to a symmetrized Laplacian on the cylinder $\mathcal{C}$ (see for example Lemma 2.3 in [BR22]) with reflecting boundaries at $x_1 = 0$: $v_t(-1, x_2) = v_t(0, x_2)$ and torus boundary conditions for $x_2 \in \{0, 3\}$: $v_t(x_1, -1) = v_t(x_1, 3)$, and $v_t(x_1, 4) = v_t(x_1, 0)$. This defines the *symmetrized* Laplacian Δ and nearest neighbors $y \sim x$ on $\mathcal{C}$.

Let $\tilde{u}(x) : \mathcal{C} \to \{0, 1\}$ be $\tilde{u}(x) := 1\{\hat{\eta}(x) \geq 3\}$. Then,

$$\Delta\tilde{u} + \zeta = \begin{bmatrix} 4 & 2 & 2 & 3 & & & \\ 2 & 3 & 3 & 2 & \zeta_1' & \zeta_1' & \cdots \\ 3 & 3 & 3 & 2 & & & \\ 3 & 3 & 2 & 3 & & & \end{bmatrix} =: \zeta'.$$

Let $v_t : \mathcal{C} \to \mathbb{Z}^+$ be the symmetrized parallel toppling odometer for ζ and $w_t : \mathcal{C} \to \mathbb{Z}^+$ the same for $\hat{\eta}'$ (defined with ζ' as $\hat{\eta}$ was with ζ). We claim that

$$\mathcal{T}'(3 + 8n) \leq \mathcal{T}(3 + 8n), \tag{54}$$

where $\mathcal{T}'(n) := \min\{t > 0 : w_t(ne_1) > 0\}$. In fact, we claim

$$v_t \leq w_t + \tilde{u} \tag{55}$$

for all $t \geq 0$. This includes (54) as $\tilde{u}((3 + 8n)e_1) = 0$. We observe (55) is a consequence of induction: the base case $t = 0$ is automatic and the inductive step is,

$$\begin{aligned} v_{t+1}(x) &= \lfloor \frac{\sum_{y\sim x} v_t(y) + \zeta(x)}{4} \rfloor \\ &\leq \lfloor \frac{\sum_{y\sim x} (w_t(y) + \tilde{u}(y)) + \zeta(x)}{4} \rfloor \\ &= \tilde{u}(x) + \lfloor \frac{\sum_{y\sim x} w_t(y) + \Delta\tilde{u}(x) + \zeta(x)}{4} \rfloor \\ &= \tilde{u}(x) + w_{t+1}(x). \end{aligned}$$

Step 4: Proof of (50). We show for all $n \geq 1$,

$$\mathcal{T}'(3 + 8n) \geq 12n. \tag{56}$$

We do so by building a 'pulsating front' for w_t in the horizontal direction. (Readers interested in pulsating fronts in periodic media on $\mathbb{R}^d$ may see Section 2.2 of [Xin09].)

We first reduce to the last-wave for w_t, $\hat{w}_t$, with initial conditions $\hat{w}_0 = \delta_{(0,3)}$ and $\hat{s}_0 = \zeta' + \Delta\hat{w}_0$. The justification is identical to Step 1 of the proof of Proposition 4.1 and so is omitted. Using $\hat{w}_t \leq 1$, we make another reduction to initial condition $\hat{w}_0 : \mathcal{C} \to \{0, 1\}$,

$$\hat{w}_0 = \begin{bmatrix} 1 & 1 & 1 & 1 & & & \\ 1 & 1 & 1 & 0 & & & \\ 1 & 1 & 1 & 0 & \mathbf{0} & \mathbf{0} & \cdots \\ 1 & 1 & 1 & 1 & & & \end{bmatrix}.$$

We now show, by manual computation, that the configuration of the odometer at the front, the rightmost 4×4 box in $\mathcal{C}$ containing a site which has toppled, is 12-periodic in time. For

notational ease, we denote sites which have toppled by $*$,

$$\hat{s}_0 = \begin{bmatrix} * & * & * & * & 4 & 2 & 2 & 3 & 3 & 2 & 2 & 3 & 3 & \\ * & * & * & 4 & 2 & 3 & 3 & 2 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & 4 & 2 & 3 & 3 & 2 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & 4 & 2 & 2 & 3 & 3 & 2 & 2 & 3 & 3 & \end{bmatrix} \quad \hat{s}_7 = \begin{bmatrix} * & * & * & * & * & * & * & * & * & * & 3 & 2 & 3 & 3 & \\ * & * & * & * & * & * & * & * & * & 4 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & * & * & * & * & 4 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & * & * & * & * & * & 3 & 2 & 3 & 3 & \end{bmatrix}$$

$$\hat{s}_1 = \begin{bmatrix} * & * & * & * & * & 3 & 2 & 3 & 3 & 2 & 2 & 3 & 3 & \\ * & * & * & * & 4 & 3 & 3 & 2 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & 4 & 3 & 3 & 2 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & 3 & 2 & 3 & 3 & 2 & 2 & 3 & 3 & \end{bmatrix} \quad \hat{s}_8 = \begin{bmatrix} * & * & * & * & * & * & * & * & * & * & 3 & 2 & 3 & 3 & \\ * & * & * & * & * & * & * & * & * & 4 & 3 & 2 & 2 & & \\ * & * & * & * & * & * & * & * & * & 4 & 3 & 2 & 2 & & \\ * & * & * & * & * & * & * & * & * & * & 3 & 2 & 3 & 3 & \end{bmatrix}$$

$$\hat{s}_2 = \begin{bmatrix} * & * & * & * & * & 3 & 2 & 3 & 3 & 2 & 2 & 3 & 3 & \\ * & * & * & * & * & 4 & 3 & 2 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & 4 & 3 & 2 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & 3 & 2 & 3 & 3 & 2 & 2 & 3 & 3 & \end{bmatrix} \quad \hat{s}_9 = \begin{bmatrix} * & * & * & * & * & * & * & * & * & * & 4 & 2 & 3 & 3 & \\ * & * & * & * & * & * & * & * & * & * & 4 & 2 & 2 & & \\ * & * & * & * & * & * & * & * & * & * & 4 & 2 & 2 & & \\ * & * & * & * & * & * & * & * & * & * & 4 & 2 & 3 & 3 & \end{bmatrix}$$

$$\hat{s}_3 = \begin{bmatrix} * & * & * & * & * & * & 4 & 2 & 3 & 3 & 2 & 2 & 3 & 3 & \\ * & * & * & * & * & * & * & 4 & 2 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & * & * & 4 & 2 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & * & 4 & 2 & 3 & 3 & 2 & 2 & 3 & 3 & \end{bmatrix} \quad \hat{s}_{10} = \begin{bmatrix} * & * & * & * & * & * & * & * & * & * & * & 4 & 3 & 3 & \\ * & * & * & * & * & * & * & * & * & * & * & 3 & 2 & & \\ * & * & * & * & * & * & * & * & * & * & * & 3 & 2 & & \\ * & * & * & * & * & * & * & * & * & * & * & 4 & 3 & 3 & \end{bmatrix}$$

$$\hat{s}_4 = \begin{bmatrix} * & * & * & * & * & * & * & 4 & 3 & 3 & 2 & 2 & 3 & 3 & \\ * & * & * & * & * & * & * & * & 3 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & * & * & * & 3 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & * & * & 4 & 3 & 3 & 2 & 2 & 3 & 3 & \end{bmatrix} \quad \hat{s}_{11} = \begin{bmatrix} * & * & * & * & * & * & * & * & * & * & * & * & 4 & 3 & \\ * & * & * & * & * & * & * & * & * & * & * & * & 3 & 2 & \\ * & * & * & * & * & * & * & * & * & * & * & * & 3 & 2 & \\ * & * & * & * & * & * & * & * & * & * & * & * & 4 & 3 & \end{bmatrix}$$

$$\hat{s}_5 = \begin{bmatrix} * & * & * & * & * & * & * & * & 4 & 3 & 2 & 2 & 3 & 3 & \\ * & * & * & * & * & * & * & * & * & 3 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & * & * & * & * & 3 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & * & * & * & 4 & 3 & 2 & 2 & 3 & 3 & \end{bmatrix} \quad \hat{s}_{12} = \begin{bmatrix} * & * & * & * & * & * & * & * & * & * & * & * & * & * & 4 & \\ * & * & * & * & * & * & * & * & * & * & * & * & * & 4 & 2 & \\ * & * & * & * & * & * & * & * & * & * & * & * & * & 4 & 2 & \\ * & * & * & * & * & * & * & * & * & * & * & * & * & * & 4 & \end{bmatrix}.$$

$$\hat{s}_6 = \begin{bmatrix} * & * & * & * & * & * & * & * & * & 4 & 2 & 2 & 3 & 3 & \\ * & * & * & * & * & * & * & * & * & 4 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & * & * & * & * & 4 & 2 & 3 & 3 & 2 & 2 & \\ * & * & * & * & * & * & * & * & * & * & 4 & 2 & 2 & 3 & 3 & \end{bmatrix}$$

This shows that

$$\hat{w}_{12} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \end{bmatrix} \mathbf{0} \quad \mathbf{0} \quad \cdots \Big],$$

so the odometer at the front is identical to what it was at the start and the process 'resets'. Hence, by induction on $n \geq 1$,

$$\hat{T}(3 + 8n) = 12n\,, \tag{57}$$

where $\hat{T}(n) := \min\{t > 0 : \hat{w}_t(ne_1) > 0\}$. This implies (56), completing the proof by (54).

6.2. **Proof of Theorem 6.1 for $d \geq 3$.** Let $\eta^{(2D)} : \mathbb{Z}^2 \to \{1, 2, 3\}$ be the (possibly random) background defined in Section 6.1 and let $d \geq 3$ be given. Our higher-dimensional counterexample $\eta : \mathbb{Z}^d \to \{2d-3, 2d-2, 2d-1\}$ is built by stacking the two-dimensional one,

$$\eta(x_1, x_2, \ldots, x_d) = 2(d-2) + \eta^{(2D)}(x_1, x_2) \text{ for all } x \in \mathbb{Z}^d. \tag{58}$$

Step 1: Proof of (51). The argument is identical to $d = 2$.

Step 2: η is explosive. Let $\eta'^{(2D)}$ be the tiling of ζ_1', ζ_2', defined in (53). The higher-dimensional analogue, $\eta' : \mathbb{Z}^d \to \{2d-3, 2d-2, 2d-1\}$ is also stacked,

$$\eta'(x_1, x_2, \ldots, x_d) := \eta'^{(2D)}(x_1, x_2) + 2(d-2). \tag{59}$$

The argument is as before: we construct a toppling procedure that transforms η into η'. Since $3\delta_0 + \eta' \geq (2d-2)$ is box-crossing, it is not stabilizable.

Topple the origin, all sites with $(2d-1)$ chips, then all sites with $(2d-2)$, then the column of $(2d-3)$ near the origin. Let $\tilde{u}$ denote the odometer for this and $\tilde{u}^{(2D)}$ the two-dimensional

version and observe that $\tilde{u}(x_1, x_2, \dots, x_d) = \tilde{u}^{(2D)}(x_1, x_2)$. This implies,

$$\begin{aligned}
&\sum_{i=1}^{d} (\tilde{u}(x - e_i) + \tilde{u}(x + e_i) - 2\tilde{u}(x)) + \eta \\
&= \sum_{i=1}^{2} (\tilde{u}^{(2D)}(x - e_i) + \tilde{u}^{(2D)}(x + e_i) - 2\tilde{u}^{(2D)}(x)) + \eta \\
&= \sum_{i=1}^{2} (\tilde{u}^{(2D)}(x - e_i) + \tilde{u}^{(2D)}(x + e_i) - 2\tilde{u}^{(2D)}(x)) + \eta^{(2D)} + 2(d-2) \\
&= \eta'^{(2D)} + 2(d-2) \\
&= \eta'.
\end{aligned}$$

We conclude by observing η' is box-crossing as $3\delta_0 + \eta' \geq (2d-2)$ and every layer in the box $\mathbf{B}^{(d)} := \{x \in \mathbb{Z}^d : 0 \leq x \leq 3\}$, contains at least one site with $(2d-1)$ chips. Indeed, for all $(3, 3, \mathbf{x}_{d-2}) \in \mathbf{B}^{(d)}$,

$$\eta'(3, 3, \mathbf{x}_{d-2}) = \eta'^{(2D)}(3, 3) + 2(d-2) = (2d-1),$$

and every layer in $\eta'^{(2D)}$ has at least one site with 3 chips.

Step 3: Proof of (50). Let $v_t^{(2D)}$ be the parallel toppling odometer for $\eta^{(2D)}$. By (56) it suffices to show that

$$v_t(x_1, x_2, \dots, x_d) \leq v_t^{(2D)}(x_1, x_2). \tag{60}$$

This is a consequence of the parabolic least action principle (Lemma 2.3 in [BR22]) but we provide a self-contained proof here:

Suppose (60) holds at time t and we want to show it holds at $(t+1)$. Let x be given. If $v_t(x) < v_t^{(2D)}(x_1, x_2)$ we are done. Hence, we may suppose $v_t(x) = v_t^{(2D)}(x_1, x_2)$ and

$$\Delta v_t(x) + \eta = \sum_{i=1}^{d} (v_t(x + e_i) + v_t(x - e_i) - 2v_t(x)) + \eta \geq 2d.$$

By (60) at time t, this implies

$$\sum_{i=1}^{2} (v_t^{(2D)}((x_1, x_2) + e_i) + v_t^{(2D)}((x_1, x_2) - e_i) - 2v_t^{(2D)}(x_1, x_2)) + \eta \geq 2d,$$

and by (58),

$$\Delta v_t^{(2D)}(x_1, x_2) + \eta^{(2D)}(x_1, x_2) \geq 2d - 2(d-2) = 4,$$

completing the proof.

To verify failure of convergence in the volume sense of Theorem 1.2, choose the periodic two-dimensional background obtained by using ζ_1 in every tile, and for $d \geq 3$ extend it by (58). For integer $t \geq 1$, write $A_t = \{y \in \mathbb{R}^d : \bar{v}_t(y) > 0\}$ and $a_n = 3 + 8n$. For every $n \geq 1$, $k \in \mathbb{Z}$, and $z \in \mathbb{Z}^{d-2}$, the cylinder comparison, followed by (60) when $d \geq 3$, gives

$$v_t(a_n, 4k, z) = 0 \qquad (t < 12n)\,.$$

When $d = 2$, z is the empty tuple in $\mathbb{Z}^0$, $\|z\|_1 = 0$, and trailing empty coordinates below are omitted. The site e_2 has fired by time 2. The nearest-neighbor path from e_2 through $(1, 1, 0)$, $(1, 4k + 1, 0)$, and $(1, 4k + 1, z)$ to $(a_n, 4k + 1, z)$ has $a_n + |4k| + \|z\|_1$ edges. Every site after e_2 on this path has background height $2d - 1$, so

$$v_{a_n+|4k|+\|z\|_1+2}(a_n, 4k + 1, z) > 0\,.$$

Fix $\varepsilon_d = 1/(100d)$ and $\Omega_d = \{y \in \mathbb{R}^d : 3/4 < y_1 < 4/5,\ |y_i| < \varepsilon_d\ (2 \le i \le d)\}$. For all sufficiently large integers t, a fixed positive proportion of the nearest-site cells in an interior box of Ω_d have centers $(a_n, 4k, z)$ and satisfy $t < 12n$ and $a_n + |4k| + \|z\|_1 + 2 \le t$. The congruence conditions on the first two coordinates have density $1/32$, and there is no restriction on the remaining coordinates. Each such cell is absent from A_t, while its translate by e_2/t is present. Hence, for some $c_d > 0$,

$$|(A_t \triangle (A_t - e_2/t)) \cap \Omega_d| \ge c_d \qquad \text{for all sufficiently large integers } t\,.$$

If $|A_t \triangle B| \to 0$ for a measurable set B, then the triangle inequality and local L^1 continuity of translation give $|(A_t \triangle (A_t - e_2/t)) \cap \Omega_d| \to 0$, a contradiction.

7. A criterion for exploding

For Proposition 7.1, let $(\Omega, \mathcal{F}, \mathbf{P})$ be a probability space of sandpile backgrounds as in Section 5. The criterion below is applied, in the proof of Theorem 1.4, to the field α_d defined there. For a finite domain $V \subset \mathbb{Z}^d$, we say $\eta : V \to \mathbb{Z}$ is *recurrent* if the firing of ∂V causes every site in V to eventually topple. Specifically, the V^c-frozen parallel toppling odometer for initial conditions $w_0 = 1\{x \in \partial V\}$, $s_0' = \eta + \Delta w_0$, is eventually 1 on V: $w_t|_V = 1$ for $t \ge |V|$. We say $\eta : \mathbb{Z}^d \to \mathbb{Z}$ is recurrent if its restriction to V is recurrent for every finite $V \subset \mathbb{Z}^d$. The measure $\mathbf{P}$ is recurrent if $\mathbf{P}(\eta \text{ is recurrent}) = 1$. The following criterion follows from recurrence, finite-range dependence, and box-crossing.

Proposition 7.1. *Let $d \ge 2$. If Hypotheses 1 and 2 are satisfied and $\mathbf{P}$ is recurrent then $\mathbf{P}$ is explosive.*

Proof. The backgrounds of Section 5 satisfy $\eta \le 2d - 1$. Recurrence of the singleton $\{x\}$ implies $\eta(x) \ge 0$: after all $2d$ external neighbors fire once, its height is $\eta(x) + 2d$. Thus $0 \le \eta \le 2d - 1$ almost surely.

Given $\delta > 0$, choose the tiling $\{\mathcal{B}_k^{(j)} : j \in \mathbb{Z}^d\}$ from Hypothesis 2, and let X_j indicate that the jth box can be crossed from each external face. Hypothesis 1 makes $\{X_j\}$ finitely dependent, with a dependence range bounded in terms of K independently of k. The domination theorem of [LSS97], used also in Section 5, therefore gives, for δ small enough, a coupling in which $\{X_j\}$ dominates supercritical independent site percolation. Almost surely there is an infinite nearest-neighbor cluster $\mathcal{C}_\infty$ of boxes with $X_j = 1$.

Fix one box $\mathcal{B}_k^{(j_0)} \in \mathcal{C}_\infty$ and set $R = \max\{\|x\|_1 : x \in \mathcal{B}_k^{(j_0)}\}$ and $N = (2d)^R$. For the N-wave from the origin in (19), induction on $r = \|x\|_1$ gives

$$u_t^{(N,0)}(x) \ge (2d)^{R-r} \qquad (r \le R,\ t \ge r)\,,$$

because $\eta \ge 0$. Hence every site of the chosen box topples. If every site of one box in $\mathcal{C}_\infty$ has toppled, then all sites of the shared external face of every adjacent box have toppled. Comparison with that box's $\mathcal{B}_k^c$-frozen process, started by one prescribed firing on this face,

shows that every site of each adjacent box in $\mathcal{C}_\infty$ subsequently topples. Following finite paths in $\mathcal{C}_\infty$ gives the N-wave infinite support, so it cannot stabilize.

If $\eta + 2dN\delta_0$ stabilized with terminal odometer v_∞, then the stable height at the origin would imply $v_\infty(0) \geq N$. At every $x \neq 0$, stability and $0 \leq s_\infty(x) < 2d$ give

$$\left\lfloor \frac{\eta(x) + \sum_{y\sim x} v_\infty(y)}{2d} \right\rfloor = v_\infty(x)\,.$$

Induction in (19) would then give $u_t^{(N,0)} \leq v_\infty$ for every t. Since v_∞ has finite support and finite values, the nondecreasing integer-valued N-wave would stabilize, a contradiction. Thus η is explosive.

□

In dimension one the conclusion also holds. Recurrence of an interval of at least two sites with zero height at both endpoints is impossible, since firing its two external neighbors starts no toppling. Two zeros at distinct sites $x < y$ would be the endpoints of such an interval, so a recurrent stable background has at most one zero. Stationarity under the finite-index sublattice makes the probability of a zero at each site vanish, so $\eta \equiv 1$ almost surely. Its 1-wave from the origin reaches both sites at distance t by time t. The comparison in the proof above then proves explosion in this case as well.

For the proof of Theorem 1.4, now use Proposition 7.1 as follows. Let η have the original common product law $\mathbf{P} = \bigotimes_{x\in\mathbb{Z}^d} \nu$ from that theorem, with $d \geq 2$, $\nu(\{m \in \mathbb{Z} : m \geq d\}) = 1$, and $p := \nu(\{2d-1\}) \in (0,1]$. Define $r_d : \mathbb{Z} \to \mathbb{Z}$ by $r_d(m) = 2d-1$ if $m = 2d-1$ and $r_d(m) = d$ otherwise, and set $\alpha_d(x) := r_d(\eta(x))$. Almost surely $d \leq \alpha_d \leq 2d-1$ and $\alpha_d \leq \eta$; the reduced law is $\mathbf{P}_d := \bigotimes_{x\in\mathbb{Z}^d}((1-p)\delta_d + p\delta_{2d-1})$. We prove recurrence and box-crossing for $\mathbf{P}_d$, the law of α_d. Proposition 7.1 then gives explosion for α_d, and monotonicity transfers box-crossing and explosion to the original η. Recurrence is asserted only for α_d, not for η. The dimension-one box-crossing case is separate. Both arguments require a form of dimensional reduction, which we record in the first subsection.

7.1. **Dimensional reduction.** Let $Q_n^{(d)} := \{x \in \mathbb{Z}^d : 1 \leq x \leq n\}$ and denote each of the (internal) $2d$ faces of $Q_n^{(d)}$ as

$$\begin{aligned} \mathcal{F}_i(Q_n^{(d)}) &:= \{x \in Q_n^{(d)} : x_i = 1\} \\ \mathcal{F}_{d+i}(Q_n^{(d)}) &:= \{x \in Q_n^{(d)} : x_i = n\}. \end{aligned} \tag{61}$$

We show that after firing the outer boundary of $\mathcal{F}_i(Q_n^{(d)})$, the sandpile dynamics on $\mathcal{F}_i(Q_n^{(d)})$ can be coupled with $(d-1)$-dimensional sandpile dynamics on $Q_n^{(d-1)}$.

Specifically, fix $i = 1$ and write points of $\mathcal{F}_1(Q_n^{(d)})$ as $(1, \mathbf{x}_{d-1})$. Let $\eta^{(d)} : Q_n^{(d)} \to \mathbb{Z}$ be given, and let $\beta : \partial Q_n^{(d-1)} \to \{0,1\}$ be an arbitrary field of tangential external values. We consider the sequences $s_t^{(d)}, u_t^{(d)}$ and $s_t^{(d-1)}, u_t^{(d-1)}$ of the *constrained frozen toppling processes*,

$(Q_n^{(d)})^c$-frozen and $(Q_n^{(d-1)})^c$-frozen respectively, with the initial conditions

$$
\begin{aligned}
&u_0^{(d)}(0,\mathbf{y}) = 1 && (\mathbf{y} \in Q_n^{(d-1)}),\\
&u_0^{(d-1)}(\mathbf{y}) = u_0^{(d)}(1,\mathbf{y}) = \beta(\mathbf{y}) && (\mathbf{y} \in \partial Q_n^{(d-1)}),\\
&u_0^{(d-1)}(\mathbf{x}) = u_0^{(d)}(1,\mathbf{x}) = 0 && (\mathbf{x} \in Q_n^{(d-1)}),\\
&s_0^{(d)}(1,\mathbf{x}) = \eta^{(d)}(1,\mathbf{x}) + \Delta u_0^{(d)}(1,\mathbf{x}) && (\mathbf{x} \in Q_n^{(d-1)}),\\
&\eta^{(d-1)}(\mathbf{x}) = \eta^{(d)}(1,\mathbf{x}) - 1, \qquad s_0^{(d-1)}(\mathbf{x}) = \eta^{(d-1)}(\mathbf{x}) + \Delta u_0^{(d-1)}(\mathbf{x}) && (\mathbf{x} \in Q_n^{(d-1)}).
\end{aligned} \tag{62}
$$

All other external odometers are initially zero. The proof of Proposition 7.2 uses only the initial conditions and the update rules, so it holds for every field β. The all-zero field is the one used for box-crossing in Proposition 7.5, and the all-one field is the one used for recurrence in Proposition 7.3. The constraints are

$$
\begin{aligned}
&u_t^{(d)}(x_1, \mathbf{x}_{d-1}) = 0 \quad \text{for all } x_1 > 1\\
&u_t^{(d)}(1, \mathbf{x}_{d-1}) \le 1\\
&u_t^{(d-1)}(\mathbf{x}_{d-1}) \le 1,
\end{aligned} \tag{63}
$$

for all $t \ge 0$. Specifically, we take initial conditions given by (62) and enforce the constraints (63) by defining the odometer sequences for $t \ge 0$ as

$$
u_{t+1}^{(d)}(x) = \begin{cases} u_t^{(d)}(x) & \text{if } x \in (Q_n^{(d)})^c \\ u_t^{(d)}(x) & \text{if } x_1 > 1 \\ u_t^{(d)}(x) & \text{if } x_1 = 1 \text{ and } u_t^{(d)}(x) = 1 \\ u_t^{(d)}(x) + 1\{s_t^{(d)}(x) \ge 2d\} & \text{otherwise} \end{cases}
$$

and

$$
u_{t+1}^{(d-1)}(\mathbf{x}_{d-1}) = \begin{cases} u_t^{(d-1)}(\mathbf{x}_{d-1}) & \text{if } \mathbf{x}_{d-1} \in (Q_n^{(d-1)})^c \\ u_t^{(d-1)}(\mathbf{x}_{d-1}) & \text{if } u_t^{(d-1)}(\mathbf{x}_{d-1}) = 1 \\ u_t^{(d-1)}(\mathbf{x}_{d-1}) + 1\{s_t^{(d-1)}(\mathbf{x}_{d-1}) \ge 2(d-1)\} & \text{otherwise.} \end{cases}
$$

The sandpile sequences are defined as in the unconstrained case. Observe that constrained frozen toppling odometers are dominated by the corresponding unconstrained odometers.

We prove the following by an induction on time. Note that a symmetric result holds for every face.

Proposition 7.2. *For all $t \ge 0$ and $(1, \mathbf{x}_{d-1}) \in \mathcal{F}_1(Q_n^{(d)})$,*

$$
u_t^{(d)}(1, \mathbf{x}_{d-1}) = u_t^{(d-1)}(\mathbf{x}_{d-1}) \tag{64}
$$

and

$$
s_t^{(d)}(1, \mathbf{x}_{d-1}) = s_t^{(d-1)}(\mathbf{x}_{d-1}) + 2 \qquad \text{if } u_t^{(d)}(1, \mathbf{x}_{d-1}) = 0. \tag{65}
$$

Proof. For all $t \geq 0$, if $u_t^{(d)}(1, \mathbf{x}_{d-1}) = 0$ and $u_t^{(d)}(1, \mathbf{y}_{d-1}) = u_t^{(d-1)}(\mathbf{y}_{d-1})$ for all $\mathbf{y}_{d-1}$,

$$\begin{aligned}
&s_t^{(d)}(1, \mathbf{x}_{d-1}) \\
&= \eta^{(d)}(1, \mathbf{x}_{d-1}) \\
&+ \Big(-2u_t^{(d)}(1, \mathbf{x}_{d-1}) + u_t^{(d)}(0, \mathbf{x}_{d-1}) + u_t^{(d)}(2, \mathbf{x}_{d-1})\Big) \\
&+ \sum_{i=2}^{d} \Big(u_t^{(d)}((1, \mathbf{x}_{d-1}) + e_i) + u_t^{(d)}((1, \mathbf{x}_{d-1}) - e_i) - 2u_t^{(d)}(1, \mathbf{x}_{d-1})\Big) \\
&= \eta^{(d-1)}(\mathbf{x}_{d-1}) + 2 \\
&+ \sum_{i=1}^{d-1} \Big(u_t^{(d-1)}(\mathbf{x}_{d-1} + e_i) + u_t^{(d-1)}(\mathbf{x}_{d-1} - e_i) - 2u_t^{(d-1)}(\mathbf{x}_{d-1})\Big) \\
&= s_t^{(d-1)}(\mathbf{x}_{d-1}) + 2.
\end{aligned}$$

The initial conditions give (64) at $t = 0$; the calculation above gives (65) whenever the common face odometer is zero. Suppose both assertions hold at time t. At a face site where the common odometer is already one, both constrained odometers remain one. Where it is zero, (65) gives

$$s_t^{(d)}(1, \mathbf{x}_{d-1}) \geq 2d \quad \Longleftrightarrow \quad s_t^{(d-1)}(\mathbf{x}_{d-1}) \geq 2(d-1)\,.$$

Thus both odometers make the same update, proving (64) at time $t + 1$; the calculation above then proves (65) wherever the new common odometer is zero.

□

7.2. **$\mathbf{P}_d$ is recurrent.** Since $d \leq \alpha_d \leq 2d - 1$, monotonicity of recurrence among backgrounds with values in $\{d, \ldots, 2d-1\}$ reduces the claim for $\mathbf{P}_d$ to the following deterministic proposition.

Proposition 7.3. *For every $d \geq 1$, $\eta : \mathbb{Z}^d \to \{d\}$ is recurrent.*

Proof. For a background bounded by $2d-1$, a process with each external neighbor fired once has odometer at most one in its domain: the constant-one odometer is a supersolution of the toppling update. Consequently it suffices to prove recurrence for translated cubes. Indeed, for every finite V, put V together with ∂V inside such a cube. Once that cube topples throughout, each site of ∂V has toppled once; comparison with the V^c-frozen process, which has all of ∂V fired at time zero, makes every site of V topple. We induct first on dimension and then on cube side length. In dimension one, firing both external neighbors of a constant-one interval makes its sites topple successively. For every dimension, $n = 1$ is immediate because the unique site receives one chip from each of its $2d$ external neighbors. For $n = 2$, every site receives one chip from each of its d external neighbors, bringing its height from d to $2d$.

Now let $n \geq 3$ and assume recurrence for side length $n - 2$ in dimension d and for all side lengths in dimension $d - 1$. We use the decomposition

$$Q_n^{(d)} = \big(\mathbf{1} + Q_{n-2}^{(d)}\big) \cup \bigcup_{i=1}^{2d} \mathcal{F}_i(Q_n^{(d)}), \qquad \mathbf{1} = (1, \ldots, 1)\,. \tag{66}$$

Fire the external boundary of $Q_n^{(d)}$. For each face, compare the resulting process with the constrained process of Proposition 7.2: fire its outer normal external face and its tangential external neighbors, and freeze all other sites off the selected face. Under (62) with the all-one field, the reduced $(d-1)$-dimensional background is the constant $d-1$ with all its external neighbors fired once. The dimension induction and (64) therefore make every site of the selected face topple. The full boundary-firing process dominates each such constrained process, so every outer face topples.

Every external neighbor of $\mathbf{1} + Q_{n-2}^{(d)}$ lies on an outer face. Comparing with its boundary-firing process and using the side-length induction makes the inner cube topple as well. Equation (66) then covers every site of $Q_n^{(d)}$. □

Remark 6. *The argument given here is similar to the proof of Lemma 3.1 in* [Sch92].

7.3. **P is box-crossing.** A coupling between the sandpile and bootstrap percolation has been observed before [FLP10]. Bootstrap percolation is a cellular automata on $\mathbb{Z}^d$ with a random initial state and a deterministic update rule. Every site $x \in \mathbb{Z}^d$ starts off as *infected* independently at random with probability p. Infected sites remain infected and if an uninfected site contains at least d neighbors which are infected, it becomes infected.

These dynamics exactly match parallel toppling for a background $\eta' : \mathbb{Z}^d \to \{d, 2d\}$ where sites are constrained to topple at most once. Infected sites are those which have toppled, and sites with $2d$ chips start as infected. Indeed, any site beginning with d chips topples if and only if it has at least d neighbors which have toppled.

We first prove box-crossing for the reduced law $\mathbf{P}_d$ using this coupling and the large-deviation bound of [Sch92] in Proposition 7.4, then transfer it to the original law $\mathbf{P}$ by monotonicity. Borrowing the terminology of [Sch92], we say a cube $Q_n^{(d)} \subset \mathbb{Z}^d$ is $\eta'-$*internally spanned* if the $(Q_n^{(d)})^c$-frozen parallel toppling procedure with $u_0 = 0$, $s_0' = \eta'$, and sites constrained to topple at most once, concludes with every site in $Q_n^{(d)}$ toppling.

Proposition 7.4 ([Sch92])**.** *Let*

$$\eta'(x) := \begin{cases} 2d & \text{with probability } p \\ d & \text{otherwise.} \end{cases}$$

There are constants c, C depending only on dimension and $p > 0$ so that

$$P(Q_n^{(d)} \text{ is } \eta'\text{-internally spanned}) \geq 1 - c\exp(-Cn). \tag{67}$$

We use Proposition 7.4 to show $\mathbf{P}$ is box-crossing.

Proposition 7.5. *For $d \geq 2$, the original product law $\mathbf{P}$ is box-crossing. In dimension one, every law supported on backgrounds $\eta \geq 1$ is box-crossing.*

Proof. The claim is immediate in dimension one: a singleton box is crossed from either external face after its prescribed seed delivers one chip, since the height is at least one. The field α_d and the constant p are as in the proof of Theorem 1.4, so $\alpha_d \leq \eta$ almost surely and α_d has common marginal $(1-p)\delta_d + p\delta_{2d-1}$. For $n \geq 1$, decompose the box into layers

$$Q_n^{(d)} = \bigcup_{i=1}^{n} \mathcal{L}_i \,,$$

where $\mathcal{L}_i := \{x \in Q_n^{(d)} : x = (i, \mathbf{x}_{d-1})\}$. The projection of each layer to a $(d-1)$-dimensional box is $\mathcal{L}_i^{(d-1)}$. Let

$$\Omega_i' := \{\eta : \mathcal{L}_i^{(d-1)} \text{ is } \eta_i'\text{-internally spanned, where } \eta_i'(\mathbf{x}_{d-1}) := \alpha_d(i, \mathbf{x}_{d-1}) - 1\}\,.$$

The variables $\eta_i'(\mathbf{x}_{d-1})$, over all layers and sites, are independent with common marginal $(1-p)\delta_{d-1} + p\delta_{2(d-1)}$. Let

$$\Omega' := \bigcap_{i=1}^{n} \Omega_i'\,.$$

On Ω', start with one prescribed firing on the external face normal to e_1. The internal spanning of $\mathcal{L}_1^{(d-1)}$ and Proposition 7.2, with the all-zero tangential field, make every site of $\mathcal{L}_1$ topple. Its firings deliver the prescribed external face to $\mathcal{L}_2$; the same comparison applies successively to all n layers. Hence $Q_n^{(d)}(\alpha_d)$ is crossed in direction e_1, and monotonicity transfers this crossing to $Q_n^{(d)}(\eta)$. Proposition 7.4 and a union bound give $\mathbf{P}(\Omega'^c) \le nc\exp(-Cn)$; symmetry gives every direction.

□

Corrections to the published version

This edition corrects the arXiv v2 text, arXiv:2102.04422v2. The list records every change of a statement, hypothesis, definition or proof relative to arXiv v2, as found by a token-level comparison of the two texts. Entry C25 records the new subsection on the use of artificial intelligence and the paragraph on the formalization, and entry C26 records the changes of wording that change no mathematics. Theorem 1.1, Theorem 1.2 and Theorem 1.3 change through the reading of (2), and Theorem 1.2 also changes through its proof, which now shows that $\mathcal{B}(p)$ is open, convex and nonempty. Theorem 1.4 is stated for one common marginal and $d \ge 2$. Theorem 5.1 changes through the restriction to $d \ge 2$, through the initial seed and the deadline in entries C05 and C13, and it also assumes Hypothesis 1 in a stronger form than printed.

C01. Law (2) in the subsection Main results, and the notation $\eta \sim \text{Bernoulli}(a, b, p)$ in the subsection Notation and conventions. Published: $\mathbf{P}(\eta(0) = (2d-2)) = 1-p$ and $\mathbf{P}(\eta(0) = (2d-1)) = p$, which fix the marginal at the origin only; the notation $\eta \sim \text{Bernoulli}(a, b, p)$ is shorthand for a product measure $\mathbf{P}$ with $\mathbf{P}(\eta(0) = a) = p$ and $\mathbf{P}(\eta(0) = b) = (1-p)$. Corrected: $\nu_p = (1-p)\delta_{2d-2} + p\delta_{2d-1}$ with $0 < p \le 1$, and $\mathbf{P} = \bigotimes_{x\in\mathbb{Z}^d} \nu_p$, so that every site has the same marginal; the notation $\eta \sim \text{Bernoulli}(a, b, p)$ means that η has law $\bigotimes_{x\in\mathbb{Z}^d}((1-p)\delta_b + p\delta_a)$, with this same marginal at every site. Reason: the published law does not determine the background away from the origin. A marginal at the origin alone, with $2d-2$ at every other site, gives a robust background, so M_η does not exist. The Bernoulli notation has the same defect, and it is used in the captions of two figures.

C02. Theorems 1.1, 1.2 and 1.3, reading of (2), and the convex domain of Theorem 1.2. Published: (2) fixes only the marginal at the origin, and the convex domain of Theorem 1.2 is not further specified. Corrected: (2) is read as in entry C01 in all three statements. The statement of Theorem 1.2 is otherwise unchanged, and its proof, described in entry C10, shows that $\mathcal{B}(p)$ is open, convex and nonempty. Reason: the

statements require the common marginal, and the published text does not show that the set in Theorem 1.2 is open, convex and nonempty.

C03. Theorem 1.4, statement. Published: "Suppose $\beta : \mathbb{Z}^d \to \mathbb{Z}$ is drawn from a product measure $\mathbf{P}$. If $\beta \geq d$ and $\mathbf{P}(\beta(0) = (2d-1)) > 0$, then β is explosive with probability 1." Corrected: "For $d \geq 2$, let ν be a probability measure on $\mathbb{Z}$ with $\nu(\{m \in \mathbb{Z} : m \geq d\}) = 1$ and $\nu(\{2d-1\}) > 0$. If $\beta : \mathbb{Z}^d \to \mathbb{Z}$ has law $\mathbf{P} = \bigotimes_{x \in \mathbb{Z}^d} \nu$, then β is explosive with probability 1." Reason: the published statement is false for product laws whose marginals differ between sites, so this main theorem changes. Let $\beta(0)$ equal d or $2d-1$ with probability $\frac{1}{2}$ each, and let $\beta(x) = d$ for $x \neq 0$. Then $\beta \geq d$ and $\mathbf{P}(\beta(0) = 2d-1) = \frac{1}{2}$. Since $d \leq 2d-2$ for $d \geq 2$, the background $\beta + n\delta_0$ lies below $(2d-2) + (n+1)\delta_0$ for every $n \geq 1$. The constant background $2d-2$ is not explosive [FLP10], so $(2d-2) + m\delta_0$ is stabilizable for every $m \geq 1$. By monotonicity, $\beta + n\delta_0$ is stabilizable for every $n \geq 1$, so β is robust and not explosive. The restriction to $d \geq 2$ is the setting of the main results.

C04. Frozen recursion (6). Published: $w_{t+1}(x) = \lfloor \frac{s_0(x) + \sum_{y \sim x} w_t(y)}{2d} \rfloor$ for $x \notin \mathcal{S}$, and $s'_{t+1}(x) = s_0(x) + \Delta(w_{t+1} - w_0)(x)$. Corrected: $w_{t+1}(x) = \max\{w_t(x), \lfloor \frac{s_0(x) + \sum_{y \sim x} w_t(y)}{2d} \rfloor\}$ for $x \notin \mathcal{S}$, and $s'_{t+1}(x) = s_0(x) + \Delta w_{t+1}(x)$. Reason: the odometer never decreases, and the maximum states this. The published identity for s'_{t+1} holds only when $\Delta w_0 = 0$, because the seed is $s_0 + \Delta w_0$.

C05. Initial seed s'_0 in Proposition 2.1, Lemma 2.1, Hypotheses 2, 3 and 4, the recurrence definition of Section 7, and the proof of Proposition 4.1. Published: $s'_0 = \eta$ in each of these places. Corrected: $s'_0 = \eta + \Delta w_0$ in each of these places, where w_0 is the initial odometer there; this is the seed $s_0 + \Delta w_0$ of (5) with $s_0 = \eta$. In the proof of Proposition 4.1 the corrected seeds are $\eta + \Delta u'_0$ and $\eta + \Delta \tilde{w}_0$. In Hypothesis 3, the words "quickly exceeds 1" are replaced by "is at least 1", which is the inequality displayed. Reason: the published $s'_0 = \eta$ contradicts (5), which defines $s'_0 = s_0 + \Delta w_0$. With $s'_0 = \eta$, nothing topples on an L-shaped path on $\eta \equiv 2d-2$. The L-shaped example refutes Lemma 2.1, and Proposition 2.1 fails because nothing topples at all.

C06. The paragraph before Proposition 2.2, proof only. Published: the good cubes dominate a high-density independent site percolation process, and large deviation bounds for supercritical percolation [AP96, GM07] are applied to it. Corrected: the good edges, those whose two cubes are good, dominate an independent bond percolation process ζ of density close to one by Corollary 1.4 of [LSS97]. The cubes all of whose ζ-edges are open form a finitely dependent site process, which dominates an independent site percolation process of high density by the same corollary. The infinite cluster of this independent site process lies in an infinite open cluster of ζ, whose cubes are good. Lemma 5.2 of [Mat08] bounds the holes of the independent site process, and Theorem 1.1 of [AP96], applied to ζ with the Borel–Cantelli lemma, bounds the chemical distances. Reason: the chemical distance bound of Antal and Pisztora concerns bond percolation, while the good cubes form a site process, so the bound is applied to the bond process ζ. Lemma 5.2 of [Mat08] concerns independent site percolation, so it is applied to the independent site process that the finitely dependent process of cubes with all ζ-edges open dominates.

C07. Lemma 2.1, initial odometer, and the caption of the figure after it. Published: $w_0|_A = 1$, and the caption reads $s'_0 = 2$. Corrected: $w_0(x) = \mathbf{1}\{x \in A\}$ on the whole lattice,

zero off A, and the caption reads $s'_0 = 2 + \Delta \mathbf{1}_A$. Reason: the frozen process (5) takes its initial odometer on the whole lattice, so w_0 must be defined off A.

C08. Lemma 3.1, statement and proof, the two paragraphs after its proof, and Remark 1. Published: "There is a constant $C := C_d$ so that for all $n \geq 1$, the support of the (Cn^d)-wave contains $[-n, n]^d$." The proof derives this from Theorem 4.1 of [FdBR08]. Corrected: the paragraph before the lemma states that the value of the wave (19) at its starting site is frozen at m, so that m is a prescribed toppling count; it defines the support of a wave and the cube $\Lambda_n = [-n, n]^d \cap \mathbb{Z}^d$; and it names the two radius bounds used. The lemma states: "For each integer $d \geq 2$ there is a deterministic constant $C_d > 0$ with the following property." For every η with $2d-2 \leq \eta(x) \leq 2d-1$ at every site, every $n \geq 1$, and every integer m with $m \geq \lceil C_2 n^2 \log(n+1) \rceil$ when $d = 2$, or $m \geq \lceil C_d n^d \rceil$ when $d \geq 3$, there is a finite integer T with $u_t^{(m,0)}(x) > 0$ for every $t \geq T$ and every $x \in \Lambda_n$. The two paragraphs after the proof are new. They state that C_d does not depend on the background, and that for a background bounded below only by $\eta_{\min}$ the wave need not be nonnegative. Remark 1 is new, and it proves $c\, n^2 \log(n+1) \leq M_2(n) \leq C\, n^2 \log(n+1)$ for homogeneous height 2 in dimension two. Reason: the published bound Cn^2 is false in dimension two, given the lower bound of Remark 1; for $d \geq 3$ the bound $C_d n^d$ is the published one. That lower bound uses only the radius bound of Fey, Levine and Peres [FLP10]. The theorem of Fey, den Boer and Redig [FdBR08] and this radius bound belong to the remark as a whole, and the corrected proof of the lemma uses both.

C09. Proof of Proposition 3.1, proof only. Published: convergence in integer directions follows from the subadditive ergodic theorem, and convergence everywhere follows from continuity and approximation. Corrected: two bounds on $\tilde{T}$ that hold on an event of probability 1, a linear bound and a comparison of $\tilde{T}$ at nearby points; the subadditive ergodic theorem applied to the first-return map to the set where $\tilde{T}(0, ny) \leq C'n + K$ for every $n \geq 1$, whose return time is integrable by Kac's lemma; and the comparison of $\tilde{T}$ at nearby points in place of continuity. Reason: the subadditive ergodic theorem needs $\tilde{T}(0, y)$ to be integrable, which the published proof does not show, and a comparison of $\tilde{T}$ at nearby points was missing.

C10. Theorem 1.2, proof. Published: the text gives no derivation of Theorem 1.2. Corrected: a proof after that of Theorem 1.3 takes $\mathcal{B}(p) = \{y : \mathcal{N}_p(y) < 1\}$, writes the region toppled by time t as a sublevel set of the rescaled arrival times, and uses that the level set $\{\mathcal{N}_p = 1\}$ has Lebesgue measure zero. Reason: Theorem 1.2 is asserted without proof.

C11. Standing assumption at the start of Section 5, and Proposition 5.1 and Theorem 5.1. Published: no dimension is assumed. Corrected: "In this section assume $d \geq 2$." Reason: Proposition 5.1 and Theorem 5.1 are false in dimension one. Take $\eta \equiv 0$ on $\mathbb{Z}$. It is stationary and ergodic, and it satisfies Hypothesis 1. Hypotheses 3 and 4 hold, since in dimension one a path or a line already fills its bounding rectangle. Hypothesis 5 holds vacuously, since it begins "Suppose η is explosive". Yet $0 = 2d-2$ is not explosive, so the explosion asserted for η fails. Theorem 5.1 is a main theorem. Its statement is unchanged, and the restriction to $d \geq 2$ comes from the standing assumption of Section 5.

C12. Hypothesis 1, statement. Published: "There exists a constant $K < \infty$ so that for all $x, y \in \mathbb{Z}^d$, $\eta(x)$ and $\eta(y)$ are independent if $|x - y| > K$." Corrected: "There exists a constant $K < \infty$ such that, for every pair of finite sets $F, G \subset \mathbb{Z}^d$ satisfying $|x-y| > K$ for every $x \in F$ and $y \in G$, the sigma-fields $\sigma(\eta(x) : x \in F)$ and $\sigma(\eta(y) : y \in G)$ are independent." Reason: the domination step in the proof of Proposition 5.1 needs the joint independence of the families on the two sets, which pairwise independence does not give. This is a strengthened hypothesis, not a correction; whether the published form suffices for Propositions 5.1 and 7.1 and Theorem 5.1 is open.

C13. Hypothesis 4, deadline. Published: $w_{|k|} \geq 1\{x \in \mathcal{B}_k\}$. Corrected: $w_{|\mathcal{B}_k|} \geq 1\{x \in \mathcal{B}_k\}$. Reason: since $\eta \leq 2d - 1$, the frozen process adds at least one site at each step until it stops. So crossing by time $|\mathcal{B}_k|$ is the same as eventual crossing, and the published deadline implies it. The change weakens the hypothesis, so the results that use it are stronger.

C14. Proof of Proposition 5.1, proof only. Published: domination of the strong box-crossing events by independent Bernoulli variables [LSS97], an infinite supercritical cluster, and the remaining steps as in Section 3, with $\eta \geq \eta_{\min}$ in place of $\eta \geq 2d-2$, and Theorem 4.1 of [LP09] for the support of a (Cn^d)-wave. Corrected: the proof is written out in six steps. Step 1 derives nonnegativity and recurrence of η from Hypothesis 3. Step 2 builds the cluster of good boxes from Hypothesis 4 and the independence in Hypothesis 1. Step 4 derives that η is explosive. Reason: the domination by independent Bernoulli variables needs the joint independence of the families in Hypothesis 1, and the published sketch does not derive the explosion of η. The published use of Theorem 4.1 of [LP09] for $\eta \geq \eta_{\min}$ does not transfer, as the paragraph after the proof of Lemma 3.1 shows: the wave need not be nonnegative, and the argument gives only a qualitative seeding.

C15. The subsection Examples satisfying the hypotheses: the verification of Hypotheses 1 and 4 for the Bernoulli cloud and the random checkerboard, and the sentence on Theorem 4.2 of [FLP10]. Published: "fix a finite set of points, $\mathcal{S} \subset \mathbb{Z}^{d}$"; "Hypothesis 1 is satisfied as $|\mathcal{S}| < \infty$ and Hypothesis 4 as $p > 0$"; for the checkerboard, "Finite range of dependence is immediate by construction" and "If we further assume $p_1 > 0$, then Hypothesis 4 is satisfied by the assumption on $\mathcal{B}(\zeta_1)$"; and "if $\eta \geq (2d-2)$ but is not box-crossing, then the background is not explosive by Theorem 4.2 in [FLP10]". Corrected: the set $\mathcal{S}$ is finite and nonempty, and both verifications are written out. For the cloud, $\eta(x)$ depends only on the uniforms U_{x-s} with $s \in \mathcal{S}$, so fields on separated finite sets depend on disjoint families of independent uniforms. This gives Hypothesis 1 with $K = \max_{s,t \in \mathcal{S}} |s-t|$. Hypothesis 4 follows from a union bound over the coordinate lines of a side-k cube and the line-filling argument of Proposition 2.1. For the checkerboard, independent tile choices give finite range of dependence, and the same line-filling argument gives Hypothesis 4 when $p_1 > 0$. The sentence on Theorem 4.2 is replaced by a description of that theorem: it gives a sufficient condition for robustness, namely a sequence of enclosing rectangular boundaries on which every height is at most $2d - 2$. Reason: the published one-line verifications do not show the hypotheses. The published conclusion does not follow from Theorem 4.2, which needs enclosing boundaries on which every height is at most $2d-2$; it is unsupported,

and the edition does not draw it. Nonemptiness of $\mathcal{S}$ is needed, since the verification chooses $s \in \mathcal{S}$.

C16. Proof of Theorem 6.1, failure of convergence for $d \geq 2$. Published: "After constructing it in two dimensions, we embed it into higher-dimensions and show failure of convergence by comparison with the two-dimensional counterexample." Corrected: the published sentence is kept, and a paragraph is added that checks failure of convergence in the volume sense of Theorem 1.2 for every $d \geq 2$. With $A_t = \{y \in \mathbb{R}^d : \bar{v}_t(y) > 0\}$, it shows that $|(A_t \triangle (A_t - e_2/t)) \cap \Omega_d| \geq c_d$ for all sufficiently large integers t, for a fixed box Ω_d and a constant $c_d > 0$, and a measurable set B with $|A_t \triangle B| \to 0$ would contradict this. Reason: Theorem 6.1 already asserts that the arrival times along e_1 do not converge. The published sentence concerns only $d \geq 3$, since it describes the embedding into higher dimensions. What the published proof lacked was failure in the volume sense of Theorem 1.2.

C17. Proposition 7.1, dimension. Published: no restriction on d. Corrected: "Let $d \geq 2$." Reason: the case $d = 1$ is argued only in the paragraph after the proof, and the introduction states that the formalization covers $d \geq 2$. The published statement in dimension one is not claimed to be false.

C18. Proposition 7.1: the sentence before it, the setting of Section 7, and its proof. Published: "The arguments in Section 5 imply the following criterion." No proof is given. Corrected: this sentence is replaced by "The following criterion follows from recurrence, finite-range dependence, and box-crossing." The opening of Section 7 adds that the criterion is applied, in the proof of Theorem 1.4, to the field α_d of entry C19. The proof for $d \geq 2$ is followed by the argument for $d = 1$ in a separate paragraph. Reason: the published criterion has no proof of its own.

C19. Proof of Theorem 1.4, after Proposition 7.1. Published: "Let $\mathbf{P}$ denote a product measure with $\mathbf{P}(\eta \geq d) = 1$ and $\mathbf{P}(\eta(0) = 2d - 1) > 0$. We show that $\mathbf{P}$ is recurrent and box-crossing. Both arguments require a form of dimensional reduction, which we record in the first subsection. Also, by monotonicity, we may assume $\eta \to \{d, 2d - 1\}$." Corrected: the proof defines $p := \nu(\{2d - 1\}) \in (0, 1]$, the map $r_d : \mathbb{Z} \to \mathbb{Z}$ with $r_d(m) = 2d - 1$ if $m = 2d - 1$ and $r_d(m) = d$ otherwise, and $\alpha_d := r_d(\eta)$. Almost surely $d \leq \alpha_d \leq 2d - 1$ and $\alpha_d \leq \eta$, and α_d has the reduced law $\mathbf{P}_d = \bigotimes_{x \in \mathbb{Z}^d}((1-p)\delta_d + p\delta_{2d-1})$. Recurrence and box-crossing are proved for $\mathbf{P}_d$. Proposition 7.1 gives explosion of α_d, and monotonicity transfers box-crossing and explosion to η, while recurrence is asserted only for α_d. The sentence on dimensional reduction is kept, a sentence says that the dimension-one case of box-crossing is separate, and the subsection on recurrence is titled "$\mathbf{P}_d$ is recurrent". Reason: the published reduction does not name the law that it reduces, and the proof now names $\mathbf{P}_d$ and proves recurrence and box-crossing for it.

C20. Setup of Proposition 7.2, initial data and tangential values. Published: $s_0^{(d-1)}(\mathbf{x}_{d-1}) = \eta^{(d)}(\mathbf{x}_{d-1}) - 1 := \eta^{(d-1)}(\mathbf{x}_{d-1})$, and the tangential values are not specified. Corrected: $\eta^{(d-1)}(\mathbf{x}) = \eta^{(d)}(1, \mathbf{x}) - 1$ and $s_0^{(d-1)}(\mathbf{x}) = \eta^{(d-1)}(\mathbf{x}) + \Delta u_0^{(d-1)}(\mathbf{x})$. The tangential external values are a field $\beta : \partial Q_n^{(d-1)} \to \{0, 1\}$, with $u_0^{(d-1)}(\mathbf{y}) = u_0^{(d)}(1, \mathbf{y}) = \beta(\mathbf{y})$ for $\mathbf{y} \in \partial Q_n^{(d-1)}$. The interior initial odometers vanish, $u_0^{(d-1)}(\mathbf{x}) = u_0^{(d)}(1, \mathbf{x}) = 0$ for $\mathbf{x} \in Q_n^{(d-1)}$, and all other external odometers are initially zero. The proof holds for every such field β: the all-zero field is the one used for box-crossing in Proposition 7.5,

and the all-one field is the one used for recurrence in Proposition 7.3. Reason: the argument $\mathbf{x}_{d-1}$ of $\eta^{(d)}$ in the published data is a typo for $(1, \mathbf{x}_{d-1})$. The published data also omit the term $\Delta u_0^{(d-1)}$, which the initial topplings require, and they do not fix the tangential, interior and remaining external initial values.

C21. Proof of Proposition 7.2, inductive step. Published: "Therefore, we may begin the induction and suppose (64) and (65) hold at time t. If $s_t^{(d-1)}(\mathbf{x}_{d-1}) \geq 2(d-1) = (2d-2)$, then $s_t^{(d)}(1, \mathbf{x}_{d-1}) = 2d$. The other direction is identical, showing $u_{t+1}^{(d)}(1, \mathbf{x}_{d-1}) = u_{t+1}^{(d-1)}(\mathbf{x}_{d-1}) = 1$ in this case." Corrected: the initial conditions give (64) at $t = 0$. Suppose both assertions hold at time t. At a face site where the common odometer is one, both constrained odometers remain one. At a site where it is zero, (65) gives $s_t^{(d)}(1, \mathbf{x}) \geq 2d$ exactly when $s_t^{(d-1)}(\mathbf{x}) \geq 2(d-1)$, so both odometers make the same update, and the calculation above gives (65) wherever the new common odometer is zero. Reason: the published step treats one case and does not show that the two constrained processes make the same update at every face site.

C22. Proposition 7.3: the sentence before it, and its proof. Published: $Q_n^{(d)} = Q_{n-2}^{(d)} \cup \bigcup_{i=1}^{2d} \mathcal{F}_i(Q_n^{(d)})$, with the inner cube written as $Q_{n-2}^{(d)}$. Corrected: $Q_n^{(d)} = (\mathbf{1} + Q_{n-2}^{(d)}) \cup \bigcup_{i=1}^{2d} \mathcal{F}_i(Q_n^{(d)})$ with $\mathbf{1} = (1, \ldots, 1)$, and the proof is written out, with base cases $n = 1, 2$ and a reduction to translated cubes. The sentence before the proposition, published as "By monotonicity of recurrence, it suffices to prove the following.", now reads "Since $d \leq \alpha_d \leq 2d-1$, monotonicity of recurrence among backgrounds with values in $\{d, \ldots, 2d-1\}$ reduces the claim for $\mathbf{P}_d$ to the following deterministic proposition." Reason: $Q_{n-2}^{(d)}$ is the cube at the corner of $Q_n^{(d)}$, not the inner cube, so the published decomposition does not hold.

C23. Proposition 7.5, box-crossing of $\mathbf{P}$. Published: "In all dimensions, $\mathbf{P}$ is box-crossing." Corrected: "For $d \geq 2$, the original product law $\mathbf{P}$ is box-crossing. In dimension one, every law supported on backgrounds $\eta \geq 1$ is box-crossing." Reason: the published statement fails for product laws whose marginals differ between sites, by the example in entry C03, where the boxes away from the origin cannot be crossed. The dimension-one case is stated for every law supported on $\eta \geq 1$.

C24. Proof of Proposition 7.5, proof only. Published: "The claim is immediate in dimension one. Let $(d+1) \geq 2$ be given." The published argument decomposes the box into layers, defines the events Ω_i' through $\mathbf{P}$, and concludes by symmetry and Proposition 7.4. Corrected: the lead sentence of the subsection now says that box-crossing is proved for the reduced law $\mathbf{P}_d$ and transferred to $\mathbf{P}$ by monotonicity. In dimension one a singleton box is crossed from either external face, since its prescribed seed delivers one chip and its height is at least one. For $d \geq 2$ the argument is run for α_d of entry C19, with the same layers $\mathcal{L}_i$ of $Q_n^{(d)}$, and with Ω_i' defined through the variables $\eta_i'(\mathbf{x}_{d-1}) = \alpha_d(i, \mathbf{x}_{d-1}) - 1$. These variables are independent with one common marginal. On $\Omega' = \bigcap_{i=1}^{n} \Omega_i'$, the internal spanning of the layers, applied in turn with the all-zero tangential field, crosses $Q_n^{(d)}(\alpha_d)$ in direction e_1. Proposition 7.4 and a union bound give $\mathbf{P}(\Omega'^c) \leq nc\exp(-Cn)$, monotonicity transfers the crossing to η, and symmetry gives every direction. Reason: the published argument applies Proposition 7.4 to layer events of $\mathbf{P}$, whose marginals may differ between sites. The

new argument applies it to layer events of α_d, whose variables have one common marginal.

C25. The subsection Use of artificial intelligence and the paragraph of the introduction on the formalization. Published: neither exists. Corrected: the subsection states who wrote the paper and its Lean formalization, gives the repository of the formalization, and says what the AI coding agents did. The paragraph states that every numbered result and Remark 1 are proved in Lean, names the three cited results that enter as explicit hypotheses, and says that Proposition 7.1 is formalized for $d \geq 2$ and Hypothesis 1 in the form stated in this edition. Reason: this edition was prepared with the formalization, and a reader needs its coverage and the use of artificial intelligence.

C26. Wording. These edits change no mathematics. In the statement of Lemma 2.1, "Then, ..., for all" becomes "Then ... for every". After the standing assumption of Section 5, the sentence "We present sufficient hypotheses on η under which the arguments above go through seamlessly." reads "We give sufficient hypotheses for convergence of last-wave arrival times." In the examples of Section 5.1, "adhoc" becomes "ad hoc". In the proof of Theorem 6.1 for $d = 2$, the induction is stated for $n \geq 1$, and a thin space is added before the comma that ends a display. In Proposition 7.2, the words "and constraints," are replaced by "The constraints are", and the condition on the constrained odometer is written $\mathbf{x}_{d-1} \in (Q_n^{(d-1)})^c$ rather than $x \in (Q_n^{(d-1)})^c$. Labels are added to Theorem 5.1 and Proposition 7.5. The source uses ties before references, the macros $\mathbb{Z}$ and $\mathbb{R}$, and the command for emphasis, and its line breaks are changed, without change of content.

References

[AC18] Scott Armstrong and Pierre Cardaliaguet, *Stochastic homogenization of quasilinear Hamilton–Jacobi equations and geometric motions*, Journal of the European Mathematical Society **20** (2018), no. 4, 797–864.

[AM20] Ian Alevy and Sevak Mkrtchyan, *The limit shape of the leaky Abelian sandpile model*, arXiv preprint arXiv:2010.01946 (2020).

[AMP02] Oswaldo SM Alves, Fabio P Machado, and S Yu Popov, *The shape theorem for the frog model*, The Annals of Applied Probability **12** (2002), no. 2, 533–546.

[AP96] Peter Antal and Agoston Pisztora, *On the chemical distance for supercritical Bernoulli percolation*, The Annals of Probability (1996), 1036–1048.

[AS16] Noga Alon and Joel H Spencer, *The probabilistic method*, John Wiley & Sons, 2016.

[AS19] Hayk Aleksanyan and Henrik Shahgholian, *Discrete balayage and boundary sandpile*, Journal d'Analyse Mathematique **138** (2019), no. 1, 361–403.

[BG07] László Babai and Igor Gorodezky, *Sandpile transience on the grid is polynomially bounded*, Proceedings of the eighteenth annual ACM-SIAM symposium on Discrete algorithms, 2007, pp. 627–636.

[BR21] Ahmed Bou-Rabee, *Convergence of the random Abelian sandpile*, The Annals of Probability **49** (2021), no. 6, 3168–3196.

[BR22] ———, *Dynamic dimensional reduction in the Abelian sandpile*, Communications in Mathematical Physics **390** (2022), no. 2, 933–958.

[BS98] Guy Barles and Panagiotis E Souganidis, *A new approach to front propagation problems: theory and applications*, Archive for Rational Mechanics and Analysis **141** (1998), no. 3, 237–296.

[CD81] J Theodore Cox and Richard Durrett, *Some limit theorems for percolation processes with necessary and sufficient conditions*, The Annals of Probability (1981), 583–603.

[CKF20] Joe Chen and Jonah Kudler-Flam, *Laplacian growth and sandpiles on the sierpiński gasket: limit shape universality and exact solutions*, Annales de l'Institut Henri Poincaré D **7** (2020), no. 4.

[CM14] LA Caffarelli and R Monneau, *Counter-example in three dimension and homogenization of geometric motions in two dimension*, Archive for Rational Mechanics and Analysis **212** (2014), no. 2, 503–574.

[DF91] Persi Diaconis and William Fulton, *A growth model, a game, an algebra, Lagrange inversion, and characteristic classes*, Rend. Sem. Mat. Univ. Pol. Torino **49** (1991), no. 1, 95–119.

[Dha99] Deepak Dhar, *The Abelian sandpile and related models*, Physica A: Statistical Mechanics and its Applications **263** (1999), no. 1-4, 4–25.

[DL81] Richard Durrett and Thomas M Liggett, *The shape of the limit set in Richardson's growth model*, The Annals of Probability **9** (1981), no. 2, 186–193.

[DS13] Deepak Dhar and Tridib Sadhu, *A sandpile model for proportionate growth*, Journal of Statistical Mechanics: Theory and Experiment **2013** (2013), no. 11, P11006.

[FdBR08] Anne Fey-den Boer and Frank Redig, *Limiting shapes for deterministic centrally seeded growth models*, Journal of Statistical Physics **130** (2008), no. 3, 579.

[Fel19] William M Feldman, *Mean curvature flow with positive random forcing in 2-d*, arXiv preprint arXiv:1911.00488 (2019).

[FL11] Anne Fey and Haiyan Liu, *Limiting shapes for a non-abelian sandpile growth model and related cellular automata.*, Journal of Cellular Automata **6** (2011).

[FLP10] Anne Fey, Lionel Levine, and Yuval Peres, *Growth rates and explosions in sandpiles*, Journal of statistical physics **138** (2010), no. 1-3, 143–159.

[FMR09] Anne Fey, Ronald Meester, and Frank Redig, *Stabilizability and percolation in the infinite volume sandpile model*, The Annals of Probability **37** (2009), no. 2, 654–675.

[FR05] Anne Fey and Frank Redig, *Organized versus self-organized criticality in the Abelian sandpile model*, Markov Processes Relat. Fields **11** (2005), 425–442.

[GG93] Janko Gravner and David Griffeath, *Threshold growth dynamics*, Transactions of the American Mathematical society (1993), 837–870.

[GG96] ———, *First passage times for threshold growth dynamics on Z2*, The Annals of Probability (1996), 1752–1778.

[GG98] ———, *Cellular automaton growth on Z2: theorems, examples, and problems*, Advances in Applied Mathematics **21** (1998), no. 2, 241–304.

[GG06] ———, *Random growth models with polygonal shapes*, The Annals of Probability (2006), 181–218.

[GM04] Olivier Garet and Régine Marchand, *Asymptotic shape for the chemical distance and first-passage percolation on the infinite bernoulli cluster*, ESAIM: Probability and Statistics **8** (2004), 169–199.

[GM07] ———, *Large deviations for the chemical distance in supercritical Bernoulli percolation*, The Annals of Probability **35** (2007), no. 3, 833–866.

[GM12] ———, *Asymptotic shape for the contact process in random environment*, The Annals of Applied Probability **22** (2012), no. 4, 1362–1410.

[GMP21] Eric Goles, Pedro Montealegre, and Kévin Perrot, *Freezing sandpiles and Boolean threshold networks: Equivalence and complexity*, Advances in Applied Mathematics **125** (2021), 102161.

[Gri13] Geoffrey R Grimmett, *Percolation*, vol. 321, Springer Science & Business Media, 2013.

[HJL19] Robert D Hough, Daniel C Jerison, and Lionel Levine, *Sandpiles on the square lattice*, Communications in Mathematical Physics **367** (2019), no. 1, 33–87.

[HLM+08] Alexander E Holroyd, Lionel Levine, Karola Mészáros, Yuyal Peres, James Propp, and David B Wilson, *Chip-firing and rotor-routing on directed graphs*, In and out of equilibrium 2, Springer, 2008, pp. 331–364.

[HS19] Robert Hough and Hyojeong Son, *Cut-off for sandpiles on tiling graphs*, arXiv preprint arXiv:1902.04174 (2019).

[IPS99] Hitoshi Ishii, Gabriel E Pires, and Panagiotis E Souganidis, *Threshold dynamics type approximation schemes for propagating fronts*, Journal of the Mathematical Society of Japan **51** (1999), no. 2, 267–308.

[Jár18] Antal A Járai, *Sandpile models*, Probability Surveys **15** (2018), 243–306.

[Kli18] Caroline J Klivans, *The mathematics of chip-firing*, CRC Press, 2018.

[KS08] Harry Kesten and Vladas Sidoravicius, *A shape theorem for the spread of an infection*, Annals of mathematics (2008), 701–766.

[LBR02] Yvan Le Borgne and Dominique Rossin, *On the identity of the sandpile group*, Discrete mathematics **256** (2002), no. 3, 775–790.

[LKG90] SH Liu, Theodore Kaplan, and LJ Gray, *Geometry and dynamics of deterministic sand piles*, Physical Review A **42** (1990), no. 6, 3207.

[LP09] Lionel Levine and Yuval Peres, *Strong spherical asymptotics for rotor-router aggregation and the divisible sandpile*, Potential Analysis **30** (2009), no. 1, 1.

[LP10] Lionel Levine and James Propp, *What is... a sandpile*, Notices Amer. Math. Soc, 2010.

[LP17] Lionel Levine and Yuval Peres, *Laplacian growth, sandpiles, and scaling limits*, Bulletin of the American Mathematical Society **54** (2017), no. 3, 355–382.

[LPS16] Lionel Levine, Wesley Pegden, and Charles K Smart, *Apollonian structure in the Abelian sandpile*, Geometric and functional analysis **26** (2016), no. 1, 306–336.

[LPS17] ———, *The Apollonian structure of integer superharmonic matrices*, Annals of Mathematics (2017), 1–67.

[LS19] Moritz Lang and Mikhail Shkolnikov, *Harmonic dynamics of the Abelian sandpile*, Proceedings of the National Academy of Sciences **116** (2019), no. 8, 2821–2830.

[LSS97] Thomas M Liggett, Roberto H Schonmann, and Alan M Stacey, *Domination by product measures*, The Annals of Probability **25** (1997), no. 1, 71–95.

[LZ19] Jessica Lin and Andrej Zlatoš, *Stochastic homogenization for reaction–diffusion equations*, Archive for Rational Mechanics and Analysis **232** (2019), no. 2, 813–871.

[Mar02] Régine Marchand, *Strict inequalities for the time constant in first passage percolation*, The Annals of Applied Probability **12** (2002), no. 3, 1001–1038.

[Mat08] P Mathieu, *Quenched invariance principles for random walks with random conductances*, Journal of Statistical Physics **130** (2008), no. 5, 1025–1046.

[Mel20] Andrew Melchionna, *The sandpile identity element on an ellipse*, arXiv preprint arXiv:2007.05792 (2020).

[Ost03] Srdjan Ostojic, *Patterns formed by addition of grains to only one site of an Abelian sandpile*, Physica A: Statistical Mechanics and its Applications **318** (2003), no. 1-2, 187–199.

[Pao13] Guglielmo Paoletti, *Deterministic Abelian sandpile models and patterns*, Springer Science & Business Media, 2013.

[PS13] Wesley Pegden and Charles K Smart, *Convergence of the Abelian sandpile*, Duke mathematical journal **162** (2013), no. 4, 627–642.

[PS20] ———, *Stability of patterns in the Abelian sandpile*, Annales Henri Poincaré, vol. 21, Springer, 2020, pp. 1383–1399.

[PW85] Norman H Packard and Stephen Wolfram, *Two-dimensional cellular automata*, Journal of Statistical physics **38** (1985), no. 5-6, 901–946.

[Red05] Frank Redig, *Mathematical aspects of the Abelian sandpile model*, Les Houches lecture notes **83** (2005), 657–659.

[Sch92] Roberto H Schonmann, *On the behavior of some cellular automata related to bootstrap percolation*, The Annals of Probability (1992), 174–193.

[Wil78] Stephen J Willson, *On convergence of configurations*, Discrete Mathematics **23** (1978), no. 3, 279–300.

[Xin09] Jack Xin, *An introduction to fronts in random media*, vol. 5, Springer Science & Business Media, 2009.

[ZZ20] Yuming Paul Zhang and Andrej Zlatos, *Long time dynamics for combustion in random media*, arXiv preprint arXiv:2008.02391 (2020).