

# Superdiffusion for Brownian motion with random drift


Ahmed Bou-Rabee (Courant Institute)

Joint work with Scott Armstrong (Courant Institute) and Tuomo Kuusi (University of Helsinki)

Probabilistic Field Theories, Aalto University, Finland

June 17, 2024

# Long time behavior of Brownian motion with isotropic random drift



Brownian motion  $X_t$  with an isotropic “random” drift  $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$

$$dX_t = f(X_t) dt + dW_t$$

## Physics predictions

- ▶ isotropic random drift = quenched Gaussian field
- ▶ predictions: (Fisher et. al (1984, 1985)) (Kravtsov et. al (1985))  
(Bouchaud et. al (1987))

## Physics predictions

- ▶ isotropic random drift = quenched Gaussian field
- ▶ predictions: (Fisher et. al (1984, 1985)) (Kravtsov et. al (1985))  
(Bouchaud et. al (1987))
  - $d \geq 3$ : if drift is small, universally *diffusive*,

$$\mathbb{E}^0[|X_t|^2] \sim t$$

proved in (Bricmont-Kupiainen (1991)), (Snitzman-Zeitouni (2006))

## Physics predictions

- ▶ isotropic random drift = quenched Gaussian field
- ▶ predictions: (Fisher et. al (1984, 1985)) (Kravtsov et. al (1985))  
(Bouchaud et. al (1987))
  - $d \geq 3$ : if drift is small, universally *diffusive*,

$$\mathbb{E}^0[|X_t|^2] \sim t$$

proved in (Bricmont-Kupiainen (1991)), (Snitzman-Zeitouni (2006))

- $d = 2$ : depends on the type of vector field

## Physics predictions

- ▶ isotropic random drift = quenched Gaussian field
- ▶ predictions: (Fisher et. al (1984, 1985)) (Kravtsov et. al (1985))  
(Bouchaud et. al (1987))
  - $d \geq 3$ : if drift is small, universally *diffusive*,

$$\mathbb{E}^0[|X_t|^2] \sim t$$

proved in (Bricmont-Kupiainen (1991)), (Snitzman-Zeitouni (2006))

- $d = 2$ : depends on the type of vector field
  - ▶ potential:  $f = \nabla P$ , *subdiffusive*

$$\mathbb{E}^0[|X_t|^2] \sim t^{1-\gamma}$$

Gaussian multiplicative chaos, progress in (Biskup, Ding, Goswami (2016))

## Physics predictions

- ▶ isotropic random drift = quenched Gaussian field
- ▶ predictions: (Fisher et. al (1984, 1985)) (Kravtsov et. al (1985))  
(Bouchaud et. al (1987))
  - $d \geq 3$ : if drift is small, universally *diffusive*,

$$\mathbb{E}^0[|X_t|^2] \sim t$$

proved in (Bricmont-Kupiainen (1991)), (Snitzman-Zeitouni (2006))

- $d = 2$ : depends on the type of vector field
  - ▶ potential:  $\mathbf{f} = \nabla P$ , *subdiffusive*

$$\mathbb{E}^0[|X_t|^2] \sim t^{1-\gamma}$$

Gaussian multiplicative chaos, progress in (Biskup, Ding, Goswami (2016))


- ▶ solenoidal:  $\operatorname{div} \mathbf{f} = 0$ , *superdiffusive*

$$\mathbb{E}^0[|X_t|^2] \sim t\sqrt{\log t}$$


**Critical dimension  $d = 2$**

## Critical dimension $d = 2$

Potential





## Critical dimension $d = 2$




## Critical dimension $d = 2$

Unconstrained







## Divergence-drift — passive tracer

$$dX_t = f(X_t)dt + \sqrt{2\nu}dW_t$$

- ▶ a divergence-free drift:  $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$  enhances the diffusivity of a Brownian particle

## Divergence-drift — passive tracer

$$dX_t = f(X_t)dt + \sqrt{2\nu}dW_t$$

- ▶ a divergence-free drift:  $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$  enhances the diffusivity of a Brownian particle
  - subcritical — **diffusive**,  $\nu \rightarrow \nu + \delta$

$$\text{cov}[f(x), f(y)] \simeq |x - y|^{-a} \quad \text{for } a > 2$$

classical: (Osada (1983)), (Oelschläger (1988)), (Fannjiang, Papanicolaou (1996))

## Divergence-drift — passive tracer

$$dX_t = f(X_t)dt + \sqrt{2\nu}dW_t$$

- ▶ a divergence-free drift:  $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$  enhances the diffusivity of a Brownian particle

- subcritical — **diffusive**,  $\nu \rightarrow \nu + \delta$

$$\text{cov}[f(x), f(y)] \simeq |x - y|^{-a} \quad \text{for } a > 2$$

classical: (Osada (1983)), (Oelschläger (1988)), (Fannjiang, Papanicolaou (1996))

- supercritical — superdiffusive,  $E^0[|X_t|^2] \sim t^{1+\delta}$

$$\text{cov}[f(x), f(y)] \simeq |x - y|^{-a} \quad \text{for } a < 2$$

(Komorowski, Olla (2002)) (annealed estimate)

## Divergence-drift — passive tracer

$$dX_t = f(X_t)dt + \sqrt{2\nu}dW_t$$

- ▶ a divergence-free drift:  $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$  enhances the diffusivity of a Brownian particle

- subcritical — **diffusive**,  $\nu \rightarrow \nu + \delta$

$$\text{cov}[f(x), f(y)] \simeq |x - y|^{-a} \quad \text{for } a > 2$$

classical: (Osada (1983)), (Oelschläger (1988)), (Fannjiang, Papanicolaou (1996))

- supercritical — superdiffusive,  $E^0[|X_t|^2] \sim t^{1+\delta}$

$$\text{cov}[f(x), f(y)] \simeq |x - y|^{-a} \quad \text{for } a < 2$$

(Komorowski, Olla (2002)) (annealed estimate)

- critical — **barely superdiffusive**,  $E^0[|X_t|^2] \sim t\sqrt{\log t}$

$$\text{cov}[f(x), f(y)] \simeq |x - y|^{-2}$$

*this talk*

## Critical divergence-free drift in $d = 2$ — past work

- ▶ consider a drift given by  $f = \nabla^\perp(H * \eta)$ , where  $H$  is a standard Gaussian free field in  $d = 2$  and  $\eta$  is a smooth, radial mollifier

## Critical divergence-free drift in $d = 2$ — past work

- ▶ consider a drift given by  $f = \nabla^\perp(H * \eta)$ , where  $H$  is a standard Gaussian free field in  $d = 2$  and  $\eta$  is a smooth, radial mollifier
- ▶ (Tóth, Valkó (2010))

$$C^{-1} \log \log t \leq \frac{1}{t} \mathbb{E}^0[|X_t|^2] \leq C \log t$$

## Critical divergence-free drift in $d = 2$ — past work

- ▶ consider a drift given by  $f = \nabla^\perp(H * \eta)$ , where  $H$  is a standard Gaussian free field in  $d = 2$  and  $\eta$  is a smooth, radial mollifier
- ▶ (Tóth, Valkó (2010))

$$C^{-1} \log \log t \leq \frac{1}{t} \mathbb{E}^0[|X_t|^2] \leq C \log t$$

- ▶ (Cannizzaro, Haunschmid-Sibitz, Toninelli (2021)), for every  $\varepsilon, \nu > 0$ , as  $t \rightarrow \infty$

$$\frac{1}{C_{\nu, \varepsilon} (\log \log t)^{1+\varepsilon}} \leq \frac{\mathbb{E}^0[|X_t|^2]}{t (\log t)^{1/2}} \leq C_{\nu, \varepsilon} (\log \log t)^{1+\varepsilon}$$

## Critical divergence-free drift in $d = 2$ — past work

- ▶ consider a drift given by  $f = \nabla^\perp(H * \eta)$ , where  $H$  is a standard Gaussian free field in  $d = 2$  and  $\eta$  is a smooth, radial mollifier
- ▶ (Tóth, Valkó (2010))

$$C^{-1} \log \log t \leq \frac{1}{t} \mathbb{E}[\mathbb{E}^0[|X_t|^2]] \leq C \log t$$

- ▶ (Cannizzaro, Haunschmid-Sibitz, Toninelli (2021)), for every  $\varepsilon, \nu > 0$ , as  $t \rightarrow \infty$

$$\frac{1}{C_{\nu, \varepsilon}(\log \log t)^{1+\varepsilon}} \leq \frac{\mathbb{E}[\mathbb{E}^0[|X_t|^2]]}{t(\log t)^{1/2}} \leq C_{\nu, \varepsilon}(\log \log t)^{1+\varepsilon}$$

- ▶ (Chatzigeorgiou, Morfe, Otto, Wang (2023)), for every  $\nu > 0$ , as  $t \rightarrow \infty$

$$C_\nu^{-1} \leq \frac{\mathbb{E}[\mathbb{E}^0[|X_t|^2]]}{t(\log t)^{1/2}} \leq C_\nu$$

## Critical divergence-free drift in $d = 2$ — past work

- ▶ consider a drift given by  $f = \nabla^\perp(H * \eta)$ , where  $H$  is a standard Gaussian free field in  $d = 2$  and  $\eta$  is a smooth, radial mollifier
- ▶ (Tóth, Valkó (2010))

$$C^{-1} \log \log t \leq \frac{1}{t} \mathbb{E}[\mathbb{E}^0[|X_t|^2]] \leq C \log t$$

- ▶ (Cannizzaro, Haunschmid-Sibitz, Toninelli (2021)), for every  $\varepsilon, \nu > 0$ , as  $t \rightarrow \infty$

$$\frac{1}{C_{\nu, \varepsilon}(\log \log t)^{1+\varepsilon}} \leq \frac{\mathbb{E}[\mathbb{E}^0[|X_t|^2]]}{t(\log t)^{1/2}} \leq C_{\nu, \varepsilon}(\log \log t)^{1+\varepsilon}$$

- ▶ (Chatzigeorgiou, Morfe, Otto, Wang (2023)), for every  $\nu > 0$ , as  $t \rightarrow \infty$

$$C_\nu^{-1} \leq \frac{\mathbb{E}[\mathbb{E}^0[|X_t|^2]]}{t(\log t)^{1/2}} \leq C_\nu$$

- ▶ above results are annealed:  $\mathbb{E}[\mathbb{E}^0[\cdot]]$

# Quenched superdiffusive invariance principle in two dimensions

Theorem (Armstrong, B., Kuusi 2024)

$\mathbb{P}$ -almost surely,

$$|\log \varepsilon^2|^{-1/4} \varepsilon X_{t/\varepsilon^2} \Rightarrow (2\pi)^{-1/2} W_t \quad \text{as } \varepsilon \rightarrow 0,$$

where  $\{W_t\}$  is a standard, two-dimensional Brownian motion and

$$\lim_{t \rightarrow \infty} \frac{\mathsf{E}^0[|X_t|^2]}{t(\log t)^{1/2}} = \frac{1}{\pi}.$$

# Quenched superdiffusive invariance principle in two dimensions

Theorem (Armstrong, B., Kuusi 2024)

$\mathbb{P}$ -almost surely,

$$|\log \varepsilon^2|^{-1/4} \varepsilon X_{t/\varepsilon^2} \Rightarrow (2\pi)^{-1/2} W_t \quad \text{as } \varepsilon \rightarrow 0,$$

where  $\{W_t\}$  is a standard, two-dimensional Brownian motion and

$$\lim_{t \rightarrow \infty} \frac{\mathbb{E}^0[|X_t|^2]}{t(\log t)^{1/2}} = \frac{1}{\pi}.$$

- ▶ first result establishing quenched (even qualitative) superdiffusivity

# Quenched superdiffusive invariance principle in two dimensions

Theorem (Armstrong, B., Kuusi 2024)

$\mathbb{P}$ -almost surely,

$$|\log \varepsilon^2|^{-1/4} \varepsilon X_{t/\varepsilon^2} \Rightarrow (2\pi)^{-1/2} W_t \quad \text{as } \varepsilon \rightarrow 0,$$

where  $\{W_t\}$  is a standard, two-dimensional Brownian motion and

$$\lim_{t \rightarrow \infty} \frac{\mathbb{E}^0[|X_t|^2]}{t(\log t)^{1/2}} = \frac{1}{\pi}.$$

- ▶ first result establishing quenched (even qualitative) superdiffusivity
- ▶ rate of superdiffusivity independent of molecular diffusivity  $\nu$  (universality)

# Quenched superdiffusive invariance principle in two dimensions

Theorem (Armstrong, B., Kuusi 2024)

$\mathbb{P}$ -almost surely,

$$|\log \varepsilon^2|^{-1/4} \varepsilon X_{t/\varepsilon^2} \Rightarrow (2\pi)^{-1/2} W_t \quad \text{as } \varepsilon \rightarrow 0,$$

where  $\{W_t\}$  is a standard, two-dimensional Brownian motion and

$$\lim_{t \rightarrow \infty} \frac{\mathbb{E}^0[|X_t|^2]}{t(\log t)^{1/2}} = \frac{1}{\pi}.$$

- ▶ first result establishing quenched (even qualitative) superdiffusivity
- ▶ rate of superdiffusivity independent of molecular diffusivity  $\nu$  (universality)
- ▶ special case of a much more general result in  $\mathbb{R}^d$

## Generator and stream matrices

$$dX_t = f(X_t)dt + \sqrt{2\nu}dW_t$$

## Generator and stream matrices

$$dX_t = f(X_t)dt + \sqrt{2\nu}dW_t$$

- ▶ the infinitesimal generator of  $\{X_t\}$  is the elliptic operator  $\mathcal{L}$  given by

$$\mathcal{L}u = \nu\Delta u + f \cdot \nabla u$$

## Generator and stream matrices

$$dX_t = f(X_t)dt + \sqrt{2\nu}dW_t$$

- ▶ the infinitesimal generator of  $\{X_t\}$  is the elliptic operator  $\mathfrak{L}$  given by

$$\mathfrak{L}u = \nu\Delta u + f \cdot \nabla u$$

- ▶ any divergence free vector field  $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$  can be written in terms of a *stream matrix*,

$$k : \mathbb{R}^d \rightarrow \{\text{skew symmetric } d \times d \text{ matrices}\}$$

as

$$\nabla \cdot k = f$$

## Generator and stream matrices

$$dX_t = f(X_t)dt + \sqrt{2\nu}dW_t$$

- ▶ the infinitesimal generator of  $\{X_t\}$  is the elliptic operator  $\mathcal{L}$  given by

$$\mathcal{L}u = \nu \Delta u + f \cdot \nabla u$$

- ▶ any divergence free vector field  $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$  can be written in terms of a *stream matrix*,

$$k : \mathbb{R}^d \rightarrow \{\text{skew symmetric } d \times d \text{ matrices}\}$$

as

$$\nabla \cdot k = f$$

- ▶ thus,

$$\mathcal{L}u = \nabla \cdot (\nu I_d + k) \nabla u$$

is a pure divergence form operator, and, in our case  $k$  is the two-dimensional GFF mollified at the unit scale (curl of the GFF)

## Convergence in law and convergence of generators

## Convergence in law and convergence of generators

- ▶ for Feller processes,

$$X_t^\varepsilon \Rightarrow X_t \iff \langle \mathfrak{L}^\varepsilon \rightarrow \mathfrak{L}'' \rangle$$

## Convergence in law and convergence of generators

- ▶ for Feller processes,

$$X_t^\varepsilon \Rightarrow X_t \iff \langle \mathcal{L}^\varepsilon \rightarrow \mathcal{L}'' \rangle$$

- ▶ in our case:

$$2\sqrt{\pi} |\log \varepsilon^2|^{-1/4} \varepsilon X_{t/\varepsilon^2} \Rightarrow \sqrt{2} W_t$$

is equivalent to

$$\langle \mathcal{L}^\varepsilon \rightarrow \Delta'' \rangle$$

where

$$\mathcal{L}^\varepsilon = ((4\sqrt{2}\pi)^{-2} |\log \varepsilon|)^{-1/2} \nabla \cdot (\nu I_d + k^\varepsilon) \nabla, \quad \text{for } k^\varepsilon(x) := k(x/\varepsilon)$$

Heuristic: study the operator scale-by-scale

## Heuristic: study the operator scale-by-scale

- ▶ can express the GFF  $k$  as a (formal) infinite sum of waves

$$k = \sum_{\ell=0}^{\infty} j_{\ell}$$

where each  $j_{\ell}$  is an independent Gaussian wave with wavelength  $3^{\ell}$

## Heuristic: study the operator scale-by-scale

- ▶ can express the GFF  $k$  as a (formal) infinite sum of waves

$$k = \sum_{\ell=0}^{\infty} j_{\ell}$$

where each  $j_{\ell}$  is an independent Gaussian wave with wavelength  $3^{\ell}$

- ▶ below scale  $3^n$ , the waves  $j_{\ell}$  for  $\ell \geq n$  are essentially constant, and so

$$\nabla \cdot (\nu I_d + k) \nabla u \approx \mathcal{L}_n u := \nabla \cdot (\nu I_d + \sum_{\ell=0}^n j_{\ell}) \nabla u \quad \text{on scales } \leq 3^n$$

## Heuristic: study the operator scale-by-scale

- ▶ can express the GFF  $k$  as a (formal) infinite sum of waves

$$k = \sum_{\ell=0}^{\infty} j_{\ell}$$

where each  $j_{\ell}$  is an independent Gaussian wave with wavelength  $3^{\ell}$

- ▶ below scale  $3^n$ , the waves  $j_{\ell}$  for  $\ell \geq n$  are essentially constant, and so

$$\nabla \cdot (\nu I_d + k) \nabla u \approx \mathcal{L}_n u := \nabla \cdot (\nu I_d + \sum_{\ell=0}^n j_{\ell}) \nabla u \quad \text{on scales } \leq 3^n$$

- ▶ the operator  $\mathcal{L}_n$  is (essentially) uniformly elliptic and so, by classical homogenization

$$\mathcal{L}_n u \approx \bar{s}_n \Delta u$$

for a deterministic constant  $\bar{s}_n > 0$ .

## Heuristic: study the operator scale-by-scale

- ▶ can express the GFF  $k$  as a (formal) infinite sum of waves

$$k = \sum_{\ell=0}^{\infty} j_{\ell}$$

where each  $j_{\ell}$  is an independent Gaussian wave with wavelength  $3^{\ell}$

- ▶ below scale  $3^n$ , the waves  $j_{\ell}$  for  $\ell \geq n$  are essentially constant, and so

$$\nabla \cdot (\nu I_d + k) \nabla u \approx \mathcal{L}_n u := \nabla \cdot (\nu I_d + \sum_{\ell=0}^n j_{\ell}) \nabla u \quad \text{on scales } \leq 3^n$$

- ▶ the operator  $\mathcal{L}_n$  is (essentially) uniformly elliptic and so, by classical homogenization

$$\mathcal{L}_n u \approx \bar{s}_n \Delta u$$

for a deterministic constant  $\bar{s}_n > 0$ .

- ▶ goal: show convergence of the effective diffusivities,  $\bar{s}_n$

## A recurrence for the effective diffusivities

- ▶ assuming no interaction between the waves,

$$\begin{aligned}\mathfrak{L}_{n+k} &\approx \nabla \cdot \left( \bar{s}_n \mathbf{I}_d + (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla \\ &= \bar{s}_n \nabla \cdot \left( \mathbf{I}_d + \bar{s}_n^{-1} (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla\end{aligned}$$

## A recurrence for the effective diffusivities

- ▶ assuming no interaction between the waves,

$$\begin{aligned}\mathfrak{L}_{n+k} &\approx \nabla \cdot \left( \bar{s}_n \mathbf{I}_d + (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla \\ &= \bar{s}_n \nabla \cdot \left( \mathbf{I}_d + \bar{s}_n^{-1} (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla\end{aligned}$$

- ▶ if  $\bar{s}_n$  is large, then this shows

$$\mathfrak{L}_{n+k} \approx \bar{s}_n \nabla \cdot \left( \mathbf{I}_d + \boldsymbol{\delta} \right) \nabla$$

for a small perturbation  $\boldsymbol{\delta}$

## A recurrence for the effective diffusivities

- ▶ assuming no interaction between the waves,

$$\begin{aligned}\mathfrak{L}_{n+k} &\approx \nabla \cdot \left( \bar{s}_n \mathbf{I}_d + (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla \\ &= \bar{s}_n \nabla \cdot \left( \mathbf{I}_d + \bar{s}_n^{-1} (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla\end{aligned}$$

- ▶ if  $\bar{s}_n$  is large, then this shows

$$\mathfrak{L}_{n+k} \approx \bar{s}_n \nabla \cdot \left( \mathbf{I}_d + \boldsymbol{\delta} \right) \nabla$$

for a small perturbation  $\boldsymbol{\delta}$

- ▶ a direct argument implies

$$\nabla \cdot (\mathbf{I}_d + \boldsymbol{\delta}) \nabla \approx \bar{s} \Delta, \quad \text{where } \bar{s} = 1 + \underbrace{\mathbb{E}[|\nabla \Delta^{-1} \nabla \cdot \boldsymbol{\delta} e_1|^2(0)]}_{k c_* \log 3} + O(\|\boldsymbol{\delta}\|^4)$$

## A recurrence for the effective diffusivities

- ▶ assuming no interaction between the waves,

$$\begin{aligned}\mathcal{L}_{n+k} &\approx \nabla \cdot \left( \bar{s}_n \mathbf{I}_d + (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla \\ &= \bar{s}_n \nabla \cdot \left( \mathbf{I}_d + \bar{s}_n^{-1} (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla\end{aligned}$$

- ▶ if  $\bar{s}_n$  is large, then this shows

$$\mathcal{L}_{n+k} \approx \bar{s}_n \nabla \cdot \left( \mathbf{I}_d + \boldsymbol{\delta} \right) \nabla$$

for a small perturbation  $\boldsymbol{\delta}$

- ▶ a direct argument implies

$$\nabla \cdot (\mathbf{I}_d + \boldsymbol{\delta}) \nabla \approx \bar{s} \Delta, \quad \text{where } \bar{s} = \underbrace{1 + \mathbb{E}[|\nabla \Delta^{-1} \nabla \cdot \boldsymbol{\delta} e_1|^2(0)]}_{k c_* \log 3} + O(\|\boldsymbol{\delta}\|^4)$$

- ▶ we deduce that

$$\bar{s}_{n+k} = \bar{s}_n \left( 1 + (k c_* \log 3) \bar{s}_n^{-2} \right) + O(\bar{s}_n^{-1}),$$


and so  $\frac{\bar{s}_n}{\sqrt{2 c_* \log 3^n}} \rightarrow 1$

## Why this recurrence?


$$\bar{s}_{n+1} \approx \bar{s}_n + \bar{s}_n^{-1}$$

## Why this recurrence?


$$\bar{s}_{n+1} \approx \bar{s}_n + \bar{s}_n^{-1}$$




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?




## Why this recurrence?



## Why this recurrence?



## Why this recurrence?



## Making the heuristic rigorous

- ▶ two challenges:

## Making the heuristic rigorous

- ▶ two challenges:
  - a priori lower bound on  $\bar{s}_n$ 
    - ▶ use a pigeonhole argument to see that it must grow by at least  $n^{1/2} \log^{-7}(n)$

## Making the heuristic rigorous

- ▶ two challenges:
  - a priori lower bound on  $\bar{s}_n$ 
    - ▶ use a pigeonhole argument to see that it must grow by at least  $n^{1/2} \log^{-7}(n)$
  - homogenization and scale separation

$$\mathfrak{L}_{n+k} \approx \nabla \cdot \left( \bar{s}_n \mathbf{I}_d + (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla$$

## Making the heuristic rigorous

- ▶ two challenges:
  - a priori lower bound on  $\bar{s}_n$ 
    - ▶ use a pigeonhole argument to see that it must grow by at least  $n^{1/2} \log^{-7}(n)$
  - homogenization and scale separation

$$\mathfrak{L}_{n+k} \approx \nabla \cdot \left( \bar{s}_n \mathbf{I}_d + (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla$$

- ▶ requires precise bounds on the approximation

$$\mathfrak{L}_n \approx \bar{s}_n \Delta$$

out of reach of prior techniques

## Making the heuristic rigorous

- ▶ two challenges:
  - a priori lower bound on  $\bar{s}_n$ 
    - ▶ use a pigeonhole argument to see that it must grow by at least  $n^{1/2} \log^{-7}(n)$
  - homogenization and scale separation

$$\mathfrak{L}_{n+k} \approx \nabla \cdot \left( \bar{s}_n \mathbf{I}_d + (\mathbf{j}_{n+1} + \cdots + \mathbf{j}_{n+k}) \right) \nabla$$

- ▶ requires precise bounds on the approximation

$$\mathfrak{L}_n \approx \bar{s}_n \Delta$$

- out of reach of prior techniques
- ▶ also, need to argue that the nearby waves do not affect the homogenization

## General case: LGF

assume the stream matrix is given by the formal sum

$$k(x) = \sum_{n=0}^{\infty} j_n(x),$$

where  $\{j_0, j_1, j_2, \dots\}$  is a sequence of independent  $\mathbb{Z}^d$ -stationary random fields satisfying roughly

## General case: LGF

assume the stream matrix is given by the formal sum

$$k(x) = \sum_{n=0}^{\infty} j_n(x),$$

where  $\{j_0, j_1, j_2, \dots\}$  is a sequence of independent  $\mathbb{Z}^d$ -stationary random fields satisfying **roughly**

- (J1) Independent pieces:  $j_n$  has range of dependence  $3^n$ .

## General case: LGF

assume the stream matrix is given by the formal sum

$$k(x) = \sum_{n=0}^{\infty} j_n(x),$$

where  $\{j_0, j_1, j_2, \dots\}$  is a sequence of independent  $\mathbb{Z}^d$ -stationary random fields satisfying roughly

- (J1) Independent pieces:  $j_n$  has range of dependence  $3^n$ .
- (J2) Local regularity: the entries of  $j_n$  are smooth in space on scale  $3^n$

## General case: LGF

assume the stream matrix is given by the formal sum

$$k(x) = \sum_{n=0}^{\infty} j_n(x),$$

where  $\{j_0, j_1, j_2, \dots\}$  is a sequence of independent  $\mathbb{Z}^d$ -stationary random fields satisfying roughly

- (J1) Independent pieces:  $j_n$  has range of dependence  $3^n$ .
- (J2) Local regularity: the entries of  $j_n$  are smooth in space on scale  $3^n$
- (J3) Dihedral symmetry: the joint law of the sequence  $\{j_n\}_{n \in \mathbb{N}}$  is invariant under coordinate rotations and negation

## General case: LGF

assume the stream matrix is given by the formal sum

$$k(x) = \sum_{n=0}^{\infty} j_n(x),$$

where  $\{j_0, j_1, j_2, \dots\}$  is a sequence of independent  $\mathbb{Z}^d$ -stationary random fields satisfying roughly

- (J1) Independent pieces:  $j_n$  has range of dependence  $3^n$ .
- (J2) Local regularity: the entries of  $j_n$  are smooth in space on scale  $3^n$
- (J3) Dihedral symmetry: the joint law of the sequence  $\{j_n\}_{n \in \mathbb{N}}$  is invariant under coordinate rotations and negation
- (J4) Non-degeneracy: there exists a constant  $c_*$  such that

$$\mathbb{E} \left[ \left| \nabla \Delta^{-1} \left( \nabla \cdot \sum_{\ell=n+1}^m j_\ell e \right)(0) \right|^2 \right] = c_* \log(3^{m-n}) + O(1)$$

## General case: LGF

assume the stream matrix is given by the formal sum

$$k(x) = \sum_{n=0}^{\infty} j_n(x),$$

where  $\{j_0, j_1, j_2, \dots\}$  is a sequence of independent  $\mathbb{Z}^d$ -stationary random fields satisfying roughly

- (J1) Independent pieces:  $j_n$  has range of dependence  $3^n$ .
- (J2) Local regularity: the entries of  $j_n$  are smooth in space on scale  $3^n$
- (J3) Dihedral symmetry: the joint law of the sequence  $\{j_n\}_{n \in \mathbb{N}}$  is invariant under coordinate rotations and negation
- (J4) Non-degeneracy: there exists a constant  $c_*$  such that

$$\mathbb{E} \left[ \left| \nabla \Delta^{-1} \left( \nabla \cdot \sum_{\ell=n+1}^m j_\ell e \right)(0) \right|^2 \right] = c_* \log(3^{m-n}) + O(1)$$

- ▶ canonical example: entries given by independent copies of mollified log-correlated Gaussian field

## Quenched superinvariance principle

Theorem (Armstrong, B., Kuusi 2024)

$\mathbb{P}$ -almost surely,

$$|\log \varepsilon|^{-1/4} \varepsilon X_{t/\varepsilon^2} \Rightarrow \sqrt{2c_*} W_t \quad \text{as } \varepsilon \rightarrow 0,$$

where  $\{W_t\}$  is a standard Brownian motion in  $\mathbb{R}^d$  and

$$\lim_{t \rightarrow \infty} \left| \frac{1}{2c_* t (\log t)^{1/2}} \mathbb{E}^0 [|X_t|^2] - d \right| = 0,$$

and, consequently, for every  $p \in [1, \infty)$

$$\lim_{t \rightarrow \infty} \mathbb{E} \left[ \left| \frac{1}{2c_* t (\log t)^{1/2}} [\mathbb{E}^0 [|X_t|^2]] - d \right|^p \right]^{1/p} = 0.$$

## Quenched superinvariance principle

Theorem (Armstrong, B., Kuusi 2024)

$\mathbb{P}$ -almost surely,

$$|\log \varepsilon|^{\frac{1}{4}} \varepsilon X_{t/\varepsilon^2} \Rightarrow \sqrt{2c_*} W_t \quad \text{as } \varepsilon \rightarrow 0,$$

where  $\{W_t\}$  is a standard Brownian motion in  $\mathbb{R}^d$  and

$$\lim_{t \rightarrow \infty} \left| \frac{1}{2c_* t (\log t)^{1/2}} \mathbb{E}^0 [|X_t|^2] - d \right| = 0,$$

and, consequently, for every  $p \in [1, \infty)$

$$\lim_{t \rightarrow \infty} \mathbb{E} \left[ \left| \frac{1}{2c_* t (\log t)^{1/2}} [\mathbb{E}^0 [|X_t|^2]] - d \right|^p \right]^{1/p} = 0.$$

- corollary of analogous result for the generators ([homogenization](#))

## Quenched superinvariance principle

Theorem (Armstrong, B., Kuusi 2024)

$\mathbb{P}$ -almost surely,

$$|\log \varepsilon|^{\frac{1}{4}} \varepsilon X_{t/\varepsilon^2} \Rightarrow \sqrt{2c_*} W_t \quad \text{as } \varepsilon \rightarrow 0,$$

where  $\{W_t\}$  is a standard Brownian motion in  $\mathbb{R}^d$  and

$$\lim_{t \rightarrow \infty} \left| \frac{1}{2c_* t (\log t)^{1/2}} \mathbb{E}^0 [|X_t|^2] - d \right| = 0,$$

and, consequently, for every  $p \in [1, \infty)$

$$\lim_{t \rightarrow \infty} \mathbb{E} \left[ \left| \frac{1}{2c_* t (\log t)^{1/2}} [\mathbb{E}^0 [|X_t|^2]] - d \right|^p \right]^{1/p} = 0.$$

- ▶ corollary of analogous result for the generators ([homogenization](#))
- ▶ actual result is quantitative — has explicit rates

## Qualitative homogenization

Theorem (Armstrong, B., Kuusi 2024)

For every Lipschitz domain  $U \subset \mathbb{R}^d$ ,  $f \in L^\infty(U)$  and  $g \in \text{Lip}(U)$  if we let  $u^\varepsilon, u_{\text{hom}} \in H^1(U)$  denote the solutions of the boundary value problems

$$\begin{cases} -\mathfrak{L}^\varepsilon u^\varepsilon = f & \text{in } U, \\ u^\varepsilon = g & \text{on } \partial U, \end{cases} \quad \text{and} \quad \begin{cases} -\Delta u_{\text{hom}} = f & \text{in } U, \\ u_{\text{hom}} = g & \text{on } \partial U, \end{cases}$$

then

$$\lim_{\varepsilon \rightarrow 0} \|u^\varepsilon - u_{\text{hom}}\|_{L^\infty(U)} = 0.$$

## Qualitative homogenization

Theorem (Armstrong, B., Kuusi 2024)

For every Lipschitz domain  $U \subset \mathbb{R}^d$ ,  $f \in L^\infty(U)$  and  $g \in \text{Lip}(U)$  if we let  $u^\varepsilon, u_{\text{hom}} \in H^1(U)$  denote the solutions of the boundary value problems

$$\begin{cases} -\mathfrak{L}^\varepsilon u^\varepsilon = f & \text{in } U, \\ u^\varepsilon = g & \text{on } \partial U, \end{cases} \quad \text{and} \quad \begin{cases} -\Delta u_{\text{hom}} = f & \text{in } U, \\ u_{\text{hom}} = g & \text{on } \partial U, \end{cases}$$

then

$$\lim_{\varepsilon \rightarrow 0} \|u^\varepsilon - u_{\text{hom}}\|_{L^\infty(U)} = 0.$$

- ▶ qualitative version of our result

## Quantitative homogenization

Theorem (Armstrong, B., Kuusi 2024)

For every Lipschitz domain  $U \subset \mathbb{R}^d$  and  $\alpha \in (0, 1/4)$ , there exists a finite random variable  $\mathcal{Z}$  such that for every  $f \in L^\infty(U)$  and  $g \in Lip(U)$  and  $\varepsilon^{-1} \geq \mathcal{Z}$

$$\begin{aligned} & \|u^\varepsilon - u_{\text{hom}}\|_{L^\infty(U)} + [\nabla u^\varepsilon - \nabla u_{\text{hom}}]_{H^{-1}(U)} \\ & + \left[ \frac{\nu I_d + k^\varepsilon - (k^\varepsilon)_U}{(2c_*^2 |\log \varepsilon|)^{1/2}} \nabla u^\varepsilon - \nabla u_{\text{hom}} \right]_{H^{-1}(U)} \\ & \leq |\log \varepsilon|^{-1/2+\alpha} (\|\nabla g\|_{L^\infty(U)} + \|f\|_{L^\infty(U)}). \end{aligned}$$

## Quantitative homogenization

Theorem (Armstrong, B., Kuusi 2024)

For every Lipschitz domain  $U \subset \mathbb{R}^d$  and  $\alpha \in (0, 1/4)$ , there exists a finite random variable  $\mathcal{Z}$  such that for every  $f \in L^\infty(U)$  and  $g \in Lip(U)$  and  $\varepsilon^{-1} \geq \mathcal{Z}$

$$\begin{aligned} & \|u^\varepsilon - u_{\text{hom}}\|_{L^\infty(U)} + [\nabla u^\varepsilon - \nabla u_{\text{hom}}]_{H^{-1}(U)} \\ & + \left[ \frac{\nu I_d + k^\varepsilon - (k^\varepsilon)_U}{(2c_*^2 |\log \varepsilon|)^{1/2}} \nabla u^\varepsilon - \nabla u_{\text{hom}} \right]_{H^{-1}(U)} \\ & \leq |\log \varepsilon|^{-1/2+\alpha} (\|\nabla g\|_{L^\infty(U)} + \|f\|_{L^\infty(U)}). \end{aligned}$$

- ▶ actual statement is slightly more quantitative

## Homogenization and coarse-graining

- ▶ instead of analyzing the generators  $\mathfrak{L}$  directly, study auxillary quantities, the *coarse-grained diffusion matrices*,  $s(\square)$ ,  $s_*(\square)$  and  $k(\square)$

## Homogenization and coarse-graining

- ▶ instead of analyzing the generators  $\mathfrak{L}$  directly, study auxillary quantities, the *coarse-grained diffusion matrices*,  $s(\square)$ ,  $s_*(\square)$  and  $k(\square)$
- ▶ defined in terms of the solution to a variational problem involving the equation restricted to a cube

## Homogenization and coarse-graining

- ▶ instead of analyzing the generators  $\mathfrak{L}$  directly, study auxillary quantities, the *coarse-grained diffusion matrices*,  $s(\square)$ ,  $s_*(\square)$  and  $k(\square)$
- ▶ defined in terms of the solution to a variational problem involving the equation restricted to a cube
- ▶ the matrix  $k(\square)$  represents the renormalized antisymmetric part of the diffusion

## Homogenization and coarse-graining

- ▶ instead of analyzing the generators  $\mathfrak{L}$  directly, study auxillary quantities, the *coarse-grained diffusion matrices*,  $s(\square)$ ,  $s_*(\square)$  and  $k(\square)$
- ▶ defined in terms of the solution to a variational problem involving the equation restricted to a cube
- ▶ the matrix  $k(\square)$  represents the renormalized antisymmetric part of the diffusion
- ▶ the matrices  $s(\square)$  and  $s_*(\square)$  give a confidence interval for the symmetric part of the diffusion

## Homogenization and coarse-graining

- ▶ instead of analyzing the generators  $\mathfrak{L}$  directly, study auxillary quantities, the *coarse-grained diffusion matrices*,  $s(\square)$ ,  $s_*(\square)$  and  $k(\square)$
- ▶ defined in terms of the solution to a variational problem involving the equation restricted to a cube
- ▶ the matrix  $k(\square)$  represents the renormalized antisymmetric part of the diffusion
- ▶ the matrices  $s(\square)$  and  $s_*(\square)$  give a confidence interval for the symmetric part of the diffusion
- ▶ quantities are central to the theory of homogenization,

## Homogenization and coarse-graining

- ▶ instead of analyzing the generators  $\mathfrak{L}$  directly, study auxillary quantities, the *coarse-grained diffusion matrices*,  $s(\square)$ ,  $s_*(\square)$  and  $k(\square)$
- ▶ defined in terms of the solution to a variational problem involving the equation restricted to a cube
- ▶ the matrix  $k(\square)$  represents the renormalized antisymmetric part of the diffusion
- ▶ the matrices  $s(\square)$  and  $s_*(\square)$  give a confidence interval for the symmetric part of the diffusion
- ▶ quantities are central to the theory of homogenization,
  - first introduced by (Armstrong, Smart (2014))

## Homogenization and coarse-graining

- ▶ instead of analyzing the generators  $\mathfrak{L}$  directly, study auxillary quantities, the *coarse-grained diffusion matrices*,  $s(\square)$ ,  $s_*(\square)$  and  $k(\square)$
- ▶ defined in terms of the solution to a variational problem involving the equation restricted to a cube
- ▶ the matrix  $k(\square)$  represents the renormalized antisymmetric part of the diffusion
- ▶ the matrices  $s(\square)$  and  $s_*(\square)$  give a confidence interval for the symmetric part of the diffusion
- ▶ quantities are central to the theory of homogenization,
  - first introduced by (Armstrong, Smart (2014))
  - a special case is classcial and goes back to (De Giorgi (1972))

Regularity improves homogenization

## Regularity improves homogenization

- ▶ proof proceeds by showing convergence of the coarse-grained matrices

## Regularity improves homogenization

- ▶ proof proceeds by showing convergence of the coarse-grained matrices
- ▶ the coarse-grained matrices describe solutions which in turn describe the matrices

## Regularity improves homogenization

- ▶ proof proceeds by showing convergence of the coarse-grained matrices
- ▶ the coarse-grained matrices describe solutions which in turn describe the matrices
- ▶ in order to close the bootstrap, we need precise analytic estimates on solutions

weak homogenization  $\Rightarrow$  regularity  $\Rightarrow$  stronger homogenization

## Regularity improves homogenization

- ▶ proof proceeds by showing convergence of the coarse-grained matrices
- ▶ the coarse-grained matrices describe solutions which in turn describe the matrices
- ▶ in order to close the bootstrap, we need precise analytic estimates on solutions
  - weak homogenization  $\Rightarrow$  regularity  $\Rightarrow$  stronger homogenization
- ▶ technical heart: *coarse-grained elliptic regularity estimates* and high contrast homogenization estimates established in the companion work (**Armstrong, Kuusi 2024**)

# Superdiffusive elliptic estimates

## Superdiffusive elliptic estimates

- ▶ vanilla Poincaré, for  $h \in H^1(B_r)$

$$r^{-1} \|h - (h)_{B_r}\|_{\underline{L}^2(B_r)} \leq \|\nabla h\|_{\underline{L}^2(B_r)}$$

and vanilla Caccioppoli, solutions to  $Ah = 0$  when  $A$  uniformly elliptic satisfy

$$\|\nabla h\|_{\underline{L}^2(B_r)} \leq \frac{C}{r} \|h - (h)_{B_{2r}}\|_{\underline{L}^2(B_{2r})}$$

## Superdiffusive elliptic estimates

- ▶ vanilla Poincaré, for  $h \in H^1(B_r)$

$$r^{-1} \|h - (h)_{B_r}\|_{\underline{L}^2(B_r)} \leq \|\nabla h\|_{\underline{L}^2(B_r)}$$

and vanilla Caccioppoli, solutions to  $Ah = 0$  when  $A$  uniformly elliptic satisfy

$$\|\nabla h\|_{\underline{L}^2(B_r)} \leq \frac{C}{r} \|h - (h)_{B_{2r}}\|_{\underline{L}^2(B_{2r})}$$

- ▶ key ingredient: superdiffusive version of these basic elliptic estimates for solutions  $\mathcal{L}u = 0$

## Superdiffusive elliptic estimates

- ▶ vanilla Poincaré, for  $h \in H^1(B_r)$

$$r^{-1} \|h - (h)_{B_r}\|_{\underline{L}^2(B_r)} \leq \|\nabla h\|_{\underline{L}^2(B_r)}$$

and vanilla Caccioppoli, solutions to  $Ah = 0$  when  $A$  uniformly elliptic satisfy

$$\|\nabla h\|_{\underline{L}^2(B_r)} \leq \frac{C}{r} \|h - (h)_{B_{2r}}\|_{\underline{L}^2(B_{2r})}$$

- ▶ key ingredient: superdiffusive version of these basic elliptic estimates for solutions  $\mathcal{L}u = 0$
- ▶ **superdiffusive Caccioppoli**

$$\|\nabla u\|_{\underline{L}^2(B_r)} \lesssim \frac{(\log r)^{1/4}}{r} \|u - (u)_{B_{2r}}\|_{\underline{L}^2(B_{2r})}$$

and **superdiffusive Poincaré**

$$\frac{(\log r)^{1/4}}{r} \|u - (u)_{B_r}\|_{\underline{L}^2(B_r)} \lesssim \|\nabla u\|_{\underline{L}^2(B_r)}$$

## Quantitative zeroth-order Liouville theorem

Theorem (Armstrong, B., Kuusi 2024)

For each  $\alpha \in (0, 1)$ , there exists  $C < \infty$  such that for every  $R \geq \mathcal{X}$ ,  $f \in L^\infty(B_R)$  and solution  $u \in H^1(B_R)$  of the equation

$$-\mathfrak{L}u = f \quad \text{in } B_R,$$

we have, for every  $r \in [\mathcal{X}, R]$ , the estimate

$$\begin{aligned} \|u - (u)_{B_r}\|_{\underline{L}^2(B_r)} &\leq C \left( \frac{r}{R} \right)^\alpha \left( \|u - (u)_{B_R}\|_{\underline{L}^2(B_R)} \right. \\ &\quad \left. + (\log R)^{-1/2} R^2 \|f\|_{L^\infty(B_R)} \right). \end{aligned}$$

## Quantitative zeroth-order Liouville theorem

Theorem (Armstrong, B., Kuusi 2024)

For each  $\alpha \in (0, 1)$ , there exists  $C < \infty$  such that for every  $R \geq \mathcal{X}$ ,  $f \in L^\infty(B_R)$  and solution  $u \in H^1(B_R)$  of the equation

$$-\mathfrak{L}u = f \quad \text{in } B_R,$$

we have, for every  $r \in [\mathcal{X}, R]$ , the estimate

$$\begin{aligned} \|u - (u)_{B_r}\|_{\underline{L}^2(B_r)} &\leq C \left( \frac{r}{R} \right)^\alpha \left( \|u - (u)_{B_R}\|_{\underline{L}^2(B_R)} \right. \\ &\quad \left. + (\log R)^{-1/2} R^2 \|f\|_{L^\infty(B_R)} \right). \end{aligned}$$

- ▶ any solution growing slower than  $x^{1-\alpha}$  is constant

## Quantitative zeroth-order Liouville theorem

Theorem (Armstrong, B., Kuusi 2024)

For each  $\alpha \in (0, 1)$ , there exists  $C < \infty$  such that for every  $R \geq \mathcal{X}$ ,  $f \in L^\infty(B_R)$  and solution  $u \in H^1(B_R)$  of the equation

$$-\mathfrak{L}u = f \quad \text{in } B_R,$$

we have, for every  $r \in [\mathcal{X}, R]$ , the estimate

$$\begin{aligned} \|u - (u)_{B_r}\|_{\underline{L}^2(B_r)} &\leq C \left( \frac{r}{R} \right)^\alpha \left( \|u - (u)_{B_R}\|_{\underline{L}^2(B_R)} \right. \\ &\quad \left. + (\log R)^{-1/2} R^2 \|f\|_{L^\infty(B_R)} \right). \end{aligned}$$

- ▶ any solution growing slower than  $x^{1-\alpha}$  is constant
- ▶ states that if you mollify at scale  $R(\log R)^{-1000}$ , the solution is Hölder

## Subquadratic solutions

- ▶ recall the classical result

$$\begin{aligned} \{u : \mathbb{R}^d \rightarrow \mathbb{R} : \Delta u = 0 \text{ and } u \text{ grows slower than a quadratic}\} \\ = \{\text{space of linear harmonic functions}\} \end{aligned}$$

## Subquadratic solutions

- ▶ recall the classical result

$$\begin{aligned} \{u : \mathbb{R}^d \rightarrow \mathbb{R} : \Delta u = 0 \text{ and } u \text{ grows slower than a quadratic}\} \\ = \{\text{space of linear harmonic functions}\} \end{aligned}$$

- ▶ this is also true for operators of the form  $\nabla \cdot a \nabla$  where

## Subquadratic solutions

- ▶ recall the classical result

$$\begin{aligned} & \{u : \mathbb{R}^d \rightarrow \mathbb{R} : \Delta u = 0 \text{ and } u \text{ grows slower than a quadratic}\} \\ &= \{\text{space of linear harmonic functions}\} \end{aligned}$$

- ▶ this is also true for operators of the form  $\nabla \cdot a \nabla$  where
  - $a$  is periodic (Avellaneda, Lin (1989))

## Subquadratic solutions

- ▶ recall the classical result

$$\begin{aligned} & \{u : \mathbb{R}^d \rightarrow \mathbb{R} : \Delta u = 0 \text{ and } u \text{ grows slower than a quadratic}\} \\ &= \{\text{space of linear harmonic functions}\} \end{aligned}$$

- ▶ this is also true for operators of the form  $\nabla \cdot a \nabla$  where
  - $a$  is periodic (Avellaneda, Lin (1989))
  - $a$  is random, but bounded uniformly (Armstrong, Smart (2014))  
(Gloria, Neukamm, Otto (2014))

## Subquadratic solutions

- ▶ recall the classical result

$$\begin{aligned} \{u : \mathbb{R}^d \rightarrow \mathbb{R} : \Delta u = 0 \text{ and } u \text{ grows slower than a quadratic}\} \\ = \{\text{space of linear harmonic functions}\} \end{aligned}$$

- ▶ this is also true for operators of the form  $\nabla \cdot a \nabla$  where
  - $a$  is periodic (Avellaneda, Lin (1989))
  - $a$  is random, but bounded uniformly (Armstrong, Smart (2014))  
(Gloria, Neukamm, Otto (2014))
- ▶ false for us, since  $a := \nu Id + k$  is *not uniformly bounded*

## Quantitative first-order Liouville theorem

$$\mathcal{A}^1(\mathbb{R}^d) = \{u : \mathbb{R}^d \rightarrow \mathbb{R} : \nabla \cdot a \nabla u = 0 \text{ and } u = O(|x|^{2-\delta})\}$$

Theorem (Armstrong, B., Kuusi 2024)

1. Finite dimensionality. We have  
that  $\mathbb{P}[\dim \mathcal{A}^1(\mathbb{R}^d) = d + 1] = 1$ .
2. Flatness at every scale. There exists  $C < \infty$  such that, for  
every  $\phi \in \mathcal{A}^1(\mathbb{R}^d)$  and  $r \geq \mathcal{X}$ , we have

$$\inf_{e \in \mathbb{R}^d} \|\phi - \ell_e\|_{L^2(B_r)} \leq C(\log r)^{-1/2+\delta} \|\phi\|_{L^2(B_r)}.$$

3. Large-scale  $C^{1,\gamma}$  estimate. For each  $R \in [\mathcal{X}, \infty)$  and  $u$   
solving  $\nabla \cdot a \nabla u = 0$  in  $B_R$  there exists  $\phi \in \mathcal{A}^1(\mathbb{R}^d)$  such that

$$\|\nabla u - \nabla \phi\|_{L^2(B_r)} \leq C \left( \frac{r}{R} \right)^\gamma \|\nabla u\|_{L^2(B_R)}, \quad \forall r \in [\mathcal{X}, R).$$

## Recap and outlook

$$dX_t = f(X_t) dt + dW_t$$

- ▶ we proved a superdiffusive quenched invariance principle, convergence of the Dirichlet problem, and Liouville theorems

## Recap and outlook

$$dX_t = f(X_t) dt + dW_t$$

- ▶ we proved a superdiffusive quenched invariance principle, convergence of the Dirichlet problem, and Liouville theorems
- ▶ phenomena occurs because of scale-by-scale *reverse cascade of homogenization*

## Recap and outlook


$$dX_t = f(X_t) dt + dW_t$$

- ▶ we proved a superdiffusive quenched invariance principle, convergence of the Dirichlet problem, and Liouville theorems
- ▶ phenomena occurs because of scale-by-scale *reverse cascade of homogenization*
- ▶ technical engine driving our proof is the regularity theory of  $\mathfrak{L}$ -harmonic functions

## Recap and outlook

$$dX_t = f(X_t) dt + dW_t$$

- ▶ we proved a superdiffusive quenched invariance principle, convergence of the Dirichlet problem, and Liouville theorems
- ▶ phenomena occurs because of scale-by-scale *reverse cascade of homogenization*
- ▶ technical engine driving our proof is the regularity theory of  $\mathfrak{L}$ -harmonic functions
- ▶ approach seems flexible and could work for other types of drift (and other problems which are renormalizable)



“Superdiffusive central limit theorem for a Brownian particle in a  
critically-correlated incompressible random drift”

(Scott Armstrong, Ahmed Bou-Rabee, and Tuomo Kuusi.)  
[arXiv:2404.01115]