APPENDIX TO : INTEGER SUPERHARMONIC MATRICES ON THE F-LATTICE

AHMED BOU-RABEE

The table below displays a Farey quadruple $(p_0, q_0, p_1, q_1) = (\mathcal{C}(p_1, q_1), p_1, q_1)$ and the Laplacian of the odd child's standard and alternate tile odometers. We only draw the Laplacian of p_0 since the Laplacian of any odd $(\frac{n}{d})$ is the rotated Laplacian of even $(\frac{d-n}{d+n})$. All quadruples with $14 \leq \det(L'(p_0)) \leq 10000$ are displayed.

| (p_0, q_0, p_1, q_1) | standard tile odometer | alternate tile odometer |
|------------------------|------------------------|--|
| (1/2, 1/3, 0/1, 1/1) | | 74 |
| (2/3, 3/5, 1/2, 1/1) | | 类 |
| (1/4, 1/5, 0/1, 1/3) | vg3k. | |
| (3/4, 5/7, 2/3, 1/1) | # | # |
| (2/5, 3/7, 1/2, 1/3) | | |
| (1/6, 1/7, 0/1, 1/5) | 18/3p | A STATE OF THE PARTY OF THE PAR |
| (4/5, 7/9, 3/4, 1/1) | # | |















