

PHYS UN1601 Recitation Worksheet 1

TA: Nitya Nigam

September 11, 2024

Problem 1

- a) In unit-vector notation, what is $\vec{r} = \vec{a} - \vec{b} + \vec{c}$ if $\vec{a} = 5.0\hat{i} + 4.0\hat{j} - 6.0\hat{k}$, $\vec{b} = -2.0\hat{i} + 2.0\hat{j} + 3.0\hat{k}$, and $\vec{c} = 4.0\hat{i} + 3.0\hat{j} + 2.0\hat{k}$?

$$\begin{aligned}\vec{a} - \vec{b} + \vec{c} &= (5 - (-2) + 4)\hat{i} + (4 - 2 + 3)\hat{j} + (-6 - 3 + 2)\hat{k} \\ &= 11\hat{i} + 5\hat{j} - 7\hat{k}\end{aligned}$$

- b) Calculate the angle between \vec{r} and the positive z axis.

$$\hat{z} = 0\hat{i} + 0\hat{j} + 1\hat{k} \Rightarrow \vec{r} \cdot \hat{z} = 11 \cdot 0 + 5 \cdot 0 + -7 \cdot 1 = -7$$


$$\vec{r} \cdot \hat{z} = |\vec{r}| |\hat{z}| \cos \theta \Rightarrow \cos \theta = \frac{-7}{\sqrt{11^2 + 5^2 + (-7)^2}} = \frac{-7}{\sqrt{145}} \approx -0.501$$

$$\Rightarrow \theta = \cos^{-1}(-0.501) = 2.10 \text{ rad} \approx 120^\circ$$

- c) What is the component of \vec{a} along the direction of \vec{b} ?

$$\vec{a} \cdot \vec{b} = 5 \cdot (-2) + 4 \cdot 2 + (-6) \cdot 3 = -10 + 8 - 18 = -20 = |\vec{a}| |\vec{b}| \cos \theta$$

$$\text{we need } |\vec{a}| \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{-20}{\sqrt{2^2 + 2^2 + 7^2}} = \frac{-20}{\sqrt{17}} = -4.85$$



- d) What is the component of \vec{a} perpendicular to the direction of \vec{b} but in the plane of \vec{a} and \vec{b} ?

we need this

subtract \vec{b}_{\parallel} component : $\vec{a} - \left(-\frac{20}{\sqrt{17}} \vec{b} \right) = \vec{a} + \frac{20}{\sqrt{17}} (-2\hat{i} + 2\hat{j} + 3\hat{k})$

$$\begin{aligned}&= 5\hat{i} + 4\hat{j} - 6\hat{k} + \left(-\frac{40}{\sqrt{17}} \hat{i} + \frac{40}{\sqrt{17}} \hat{j} + \frac{60}{\sqrt{17}} \hat{k} \right) \\ &= \frac{45}{\sqrt{17}} \hat{i} + \frac{104}{\sqrt{17}} \hat{j} - \frac{42}{\sqrt{17}} \hat{k}\end{aligned}$$

$$|\vec{a} - |\vec{a}| \cos \theta \vec{b}| = \sqrt{2.65^2 + 6.35^2 + 2.49^2} \approx 7.31$$