

# PHYS UN1602 Recitation Week 11 Worksheet

TA: Nitya Nigam

April 23-24, 2025

## Problem 1

The star Gliese 832 is located 16 lightyears away from the Earth. Suppose we send a spacecraft to investigate the recently discovered exoplanet in that star system with velocity  $v = \frac{4}{5}c$  and  $\gamma = \frac{5}{3}$ .

- a) According to the occupants of the spacecraft, how long does the trip take?
- b) According to the occupants of the spacecraft, how far did they travel?
- c) Once they arrive, they send a light-based signal back to Earth. According to the people on Earth, how much time passed between the spacecraft leaving and the signal being received?

*Solution.*

- a) According to someone in the Earth's rest frame, traveling 16 lightyears at a velocity of  $v = \frac{4}{5}c$   $\text{ms}^{-1}$  will take:

$$\Delta t_E = \frac{16 \text{ ly}}{\frac{4}{5} \text{ ly} / \text{yr}} = \left( \frac{5}{4} \text{ yr} / \text{ly} \right) (16 \text{ ly}) = 20 \text{ yr}$$

Someone on the spacecraft moving at a relative velocity  $v$  will experience time dilation compared to the observer on Earth and measure an elapsed time of:

$$\Delta t_{SC} = \frac{1}{\gamma} \Delta t_E = \frac{3}{5} (20 \text{ yr}) = 12 \text{ yr}$$

- b) According to a stationary observer in Earth's rest frame, the spacecraft traveled 16 lightyears. However, due to length contraction the occupants of the spacecraft measure the distance to be:

$$\Delta L_{SC} = \frac{1}{\gamma} \Delta L_E = \frac{3}{5} (16 \text{ ly}) = 9.6 \text{ ly}$$

- c) According to the Earth-based observer, it took them 20 yr to get to Gliese 832, and then 16 yr for the light to return. So 36 years in total.

## Problem 2

High-energy photons propagating through space can convert into electron-positron pairs by scattering with cosmic microwave background (CMB) photons. Taking the average CMB temperature of 2.8 K, a typical CMB photon will have an energy of roughly  $7 \times 10^{-4}$  eV. Calculate the minimum energy required for the high-energy photon to produce an electron-positron pair ( $m_e = 511$  keV) if:

- The CMB photon momentum is perpendicular to that of the high-energy photon.
- The CMB photon propagates in the direction opposite the high-energy photon.
- Suppose the CMB photon propagates in the same direction as the high-energy photon. Is it ever possible for the two photons to collide and produce an electron-positron pair?

*Solution.*

- The ability to produce an  $e^-e^+$  pair is constrained by the system mass of the scattering, which is given by:

$$m_{\text{total}}^2 = \mathbf{p}_{\text{total}}^2 = \mathbf{p}_\gamma^2 + \mathbf{p}_{\text{CMB}}^2 + 2\mathbf{p}_\gamma \cdot \mathbf{p}_{\text{CMB}} = 2\mathbf{p}_\gamma \cdot \mathbf{p}_{\text{CMB}}$$

where we used the fact that photons have zero mass. Further simplifying:

$$m_{\text{total}}^2 = 2(E_\gamma E_{\text{CMB}} - \vec{p}_\gamma \cdot \vec{p}_{\text{CMB}}) = 2E_\gamma E_{\text{CMB}}(1 - \cos \theta)$$

Solving for  $E_\gamma$  gives:

$$E_\gamma = \frac{m_{\text{total}}^2}{2E_{\text{CMB}}(1 - \cos \theta)}$$

So at an angle of  $\theta$  between the high-energy photon and the CMB photon, the minimum energy required to produce an electron-positron pair is:

$$E_\gamma(\theta) = \frac{(2 \cdot 511 \times 10^3)^2}{2 \cdot 7 \times 10^{-4} \cdot (1 - \cos \theta)}$$

Using this formula:

$$E_\gamma\left(\theta = \frac{\pi}{2}\right) = 746 \text{ TeV}$$

- 

$$E_\gamma(\theta = \pi) = 373 \text{ TeV}$$

- At  $\theta = 0$ , the formula for  $E_\gamma$  becomes singular. Looking again at the expression for the system mass, we see that this is because in this case  $m_{\text{total}}$  is always 0. This tells us that there is no energy available for interactions, so the photons cannot collide to produce an electron-positron pair. This is in line with our intuition, as both photons must always move colinearly at the speed of light in all inertial frames.