



Rapid Solution Exchange to Neuronal Cultures Grown on Multi Electrode Arrays

by Nitzan Herzog

Thesis submitted to The University of Nottingham
for the degree of Doctor of Philosophy, January 2017

Contents

List of Figures	5
Abstract	8
Acknowledgements	9
1 Introduction	10
1.1 The enterprise of experimental neuroscience	10
1.2 The MEA culture model for network activity in neuronal ensembles	12
1.2.1 Notable achievements and advantages of neuronal cultures on MEAs	15
1.3 Volume transmission in neuronal systems	17
1.3.1 Neuromodulator transmission and plasticity	19
1.3.2 The Izhikevic thought experiment	21
1.4 Proposed system and how it addresses present experimental gaps	24
1.5 Rapid solution exchange with microfluidics	25
1.5.1 Microfluidics in neuroscience	27
1.5.2 Neurons and flow	28
1.6 Ph.D objectives	29
2 Methods	30
2.1 Basic fabrication elements	30
2.1.1 PDMS preparation	30
2.1.2 Thin film spinning	30
2.1.3 Soft lithography	31
2.1.4 PDMS extraction	31
2.2 Bonding techniques	31
2.2.1 Plasma bonding	31
2.2.2 Double sided silicone tape	31
2.3 Surface coating	32
2.3.1 Open surface	32
2.3.2 Devices - Bond-then-PLL	32
2.3.3 Devices - PEI-then-bond	32
2.4 Seeding and maintaining of Neuronal cultures	32

2.5	MEA recording and stimulation	32
2.5.1	Spike detection and noise removal	32
2.5.1.1	Automatic quantification of active channels	32
2.5.2	Electrical stimulation	33
2.5.2.1	Automatic detection of responsive channels	33
2.5.3	Correlation maps	33
2.5.4	Burst detection	33
2.5.5	Functional connectivity analysis	33
2.6	Plasticity protocol	33
2.7	Conditioned media production	33
2.8	Flow experiments	34
2.8.1	Heated chamber	34
2.8.2	Steady flow	34
2.8.3	Pulsing	34
2.9	Immunohistochemistry	34
3	Establishment of a culture model for network activity in neuronal ensembles	40
3.1	Introduction	40
3.2	Development of spontaneous activity in Mouse cultures	41
3.2.1	Statistics of activity and synchronicity measures	44
3.2.2	Comparison between mouse and rat cultures	48
3.3	Evoked activity	49
3.4	Plasticity induction in the presence of dopamine	52
3.4.1	Examining changes in response to stimulation	54
3.4.2	Examining changes in functional connectivity	57
3.5	Chapter conclusion	60
4	Viability of neuronal cultures in microfluidic devices in static conditions and under steady flow	62
4.1	Introduction	62
4.2	Long term neuronal cultures in microfluidic devices	64
4.2.1	Development of protocol	64
4.2.1.1	Evaporation and surface chemistry considerations	66
4.2.1.2	Considerations of factor circulation	66
4.2.1.3	Alternative bonding methods	70
4.2.1.4	Extraction of PDMS	71
4.2.2	Growing microcultures in plasma bonded devices	74
4.3	Viability of neuronal cultures under steady microfluidic flow	78
4.3.1	Pilot flow study	78

4.3.2	Quantitative viability analysis	80
4.4	Chapter conclusion	83
5	Activity under steady microfluidic flow	86
5.1	Introduction	86
5.2	Neuronal cultures in cross flow devices on MEAs	87
5.3	Activity under flow for young cultures	88
5.3.1	Effect of flow rate	90
5.3.2	The semi-permeable membrane approach for shear reduction	92
5.3.3	Considerations of diffusive flux	94
5.4	Activity under flow for old cultures	97
5.4.1	The effect of the media source	100
5.4.2	How old conditioned media performs on young cultures	102
5.5	Interpretation of the activity under flow results	104
5.6	Chapter conclusion	109
6	Rapid programmatic agonist delivery to a neuronal microculture	110
6.1	Introduction	110
6.2	Fabrication and establishment of long term Neuronal microcultures	111
6.2.1	PEI-then-all-tape devices	112
6.2.2	PEI-then-PDMS-tape device	113
6.2.3	Network Activity in microcultures	116
6.3	Pulsing performance in microculture devices	118
6.3.1	Analysis of pulsing visualized by fluorescein	119
6.3.2	Numerical simulation of drug pulsing	120
6.3.3	Shortening of the transient time scales by changing microfluidic parameters	125
6.4	Glutamate pulsing	127
6.5	Dopamine pulses	129
6.6	Chapter conclusion	137
7	Discussion	140
7.1	The utility of the culture model	140
7.2	The shear protection vs. conditioning protection issue	140
7.2.1	Limitation uncovered by the conducted experiments	140
7.2.2	Avenues for further expansion of the model	140
7.3	Importance of extrasynaptic environment	140
7.4	Prospects of the <i>in vitro</i> volume transmission model	141

A Appendix	142
A.1 MEA Data sheets	142
A.2 Parameters of Comsol model	142
Bibliography	142

List of Figures

1.1	Illustration of the volume and wired transmission concepts	18
1.2	Volume transmission view of dopamine signalling	21
1.3	The Izhikevich proposal for a plasticity rule involving dopamine	23
1.4	Generating a pulse with the interface shifting method	27
2.1	profiles	30
2.2	microwell profile	31
3.1	Representative images of a cortical mouse culture developing on a planar multi electrode array	42
3.2	Development of synchrony in the spontaneous activity of a representative mouse culture	43
3.3	Averaged statistics of development of activity measures in mouse cultures . .	45
3.4	Averaged statistics of development of bursting measures in mouse cultures . .	47
3.5	Comparison between spontaneous activity in mouse and rat based cultures . .	49
3.6	Example of responses to test stimuli applied at 2 different electrodes	51
3.7	Outline of the combined dopamine-and-tetanus-induced open bath plasticity experiments	54
3.8	Example response rasters from the combined dopamine and tetanus plasticity induction experiment	56
3.9	Example stimulation response maps for the combined dopamine and tetanus plasticity induction experiment	57
3.10	Statistics of changes to evoked responses in the combined dopamine and tetanus plasticity induction experiment	58
3.11	Statistics of Change to functional connectivity and average unit firing rate in the combined dopamine and tetanus plasticity induction experiment	59
4.1	Schematics of the standard single layer microfluidic devices	65
4.2	Effects of osmolarity drift in early protocol for long term culturing of neurons in microfluidic devices	67
4.3	The immersion maintenance configuration	68
4.4	Demonstration of the limitations of circulation in planar microfluidic devices	69

4.5	Effect of the pre-polymer bonding approach on the development of neuronal cultures in microfluidic devices	70
4.6	Comparison between cultures growing in plasma bonded devices and tape based devices	72
4.7	Demonstration of PDMS related contaminations and the extraction procedure	73
4.8	Schematics of the 2-layered microfluidic devices with microwells	75
4.9	Neuronal microcultures growing without a support culture	75
4.10	Development of neuronal microcultures	76
4.11	Statistics of microculture viability over development	77
4.12	Demonstration of the microculture isolation issue with the bond-then-surface approach	78
4.13	Time lapse of neuronal culture under steady microfluidic flow	79
4.14	Effect of media conditioning and flow rate on viability of neuronal cultures under steady microfluidic flow	81
4.15	Example of viability curves from individual steady flow experiments	82
4.16	Statistics of death rates for various steady flow conditions	84
5.1	Illustration of the cross flow devices used for measuring culture activity under flow	88
5.2	Images of a neuronal culture growing in cross flow devices	89
5.3	Example for the Effect of flow rate on the activity of a young culture under flow	91
5.4	Averaged time course of activity measures in young cultures placed under flow at different rates	93
5.5	Bright field and staining images of a culture in a cross flow device inclusive of a semi-permeable membrane	95
5.6	Averaged time course of activity measures in young cultures separated from the fast flow by means of a semi-permeable membrane	96
5.7	Example for the effect of culture age on the network activity under flow	98
5.8	Averaged time course of activity measures in young versus old cultures under flow	99
5.9	Averaged time course of activity measures in old cultures under flow with different media types	101
5.10	Averaged time course of activity measures in young cultures under flow with old media	103
5.11	Illustration of a synaptic cleft with flow running through it	107
6.1	Illustration of the new microwell devices used for dopamine pulsing on a microculture	112
6.2	Microcultures grown in all-tape devices exhibit bad surface adhesion	113

6.3	Development of microcultures in hybrid PDMS-tape devices	114
6.4	Immunohistochemistry of microcultures in hybrid PDMS-tape devices	115
6.5	Effectiveness of surface-then-bond in maintaining an isolated microculture . .	116
6.6	Effectiveness of surface-then-bond in maintaining an isolated microculture - immunostaining evidence	117
6.7	Spontaneous and evoked activity in microcultures	119
6.8	Fluorescence visualization of agonist pulsing	120
6.9	Analysis pipeline of fluorescence pulsing data	121
6.10	Measured and model flow rate switches during an agonist pulse	122
6.11	Comparison between model predictions and measured agonist time course . .	123
6.12	Comparison between model predictions and measured agonist time course including variability between devices	124
6.13	Spatial distributions of agonist concentration in the microwell during a pulse	126
6.14	Shortening of the agonist transient time scales by changing the microfluidic parameters	128
6.15	Microculture electrical responses to glutamate pulses	129
6.16	Illustration of the dopamine pulsing experimental protocol	130
6.17	Example response rasters from the dopamine pulsing experiment	131
6.18	Channel PSTHs before and after administration of pulsing protocol with dopamine or control solutions - example of global increase in response intensity	132
6.19	Channel PSTHs before and after administration of pulsing protocol with dopamine or control solutions - example of global decrease in response in- tensity	133
6.20	Timing of dopamine transient relative to reverberative stimulation response .	135
6.21	Global stimulation response and functional connectivity analysis of the dopamine pulsing experiments	136
6.22	Statistics of the change in evoked responses following dopamine or blank pulsing	139