## CONCORDIA UNIVERSITY

## Department of Mathematics & Statistics

1		
Course	Number	Sections
Mathematics	205	All
Examination	Date	Pages
Final	December 2016	2
Instructors:	M. Bertola, J. Brody,	Course Examiner
	A. Conti , K. Lagota	A. Atoyan and H. Proppe
Special	Only approved calculators are allowed.	
Instructions:	Show all your work for full marks.	

 $\overline{MARKS}$ 

[10] 1. a. Sketch the graph of the function

$$f(x) = \begin{cases} 1 - 2x & \text{if } x \le 1\\ \frac{|1 - x|}{1 - x} & \text{if } 1 < x \le 2\\ x - 3 & \text{if } x > 2 \end{cases}$$

and find the definite integral  $\int_{0}^{3} f(x) dx$  in terms of area (do not antidifferentiate).

**b** Use the Fundamental Theorem of Calculus to calculate the derivative of  $F(x) = \int\limits_{1-x}^{x^2} 2^{-t^2} \, \mathrm{d}t \; .$ 

[16] **2.** Find the following indefinite integrals:

(a) 
$$\int \frac{(x^{2/3} + x^{-2/3})^2}{x} dx$$
 (b)  $\int \arcsin(x) dx$  (c)  $\int \frac{x^3 + 4}{x^3 - 4x} dx$ 

[12] **3.** Evaluate the following definite integrals (give the exact answers):

(a) 
$$\int_{0}^{\pi/2} x^{2} \cos(2x) dx$$
 (b)  $\int_{0}^{4} x \sqrt{16 - x^{2}} dx$ 

[6] 4. Find F(t) such that  $F'(t) = \sin^3(t) \cos^5(t)$  and  $F\left(\frac{\pi}{2}\right) = 0$ .

[8] 5. Evaluate the given improper integral or show that it diverges:

(a) 
$$\int_{0}^{1/e} \frac{\mathrm{d}x}{x \ln^{2}(x)}$$
 (b) 
$$\int_{0}^{\infty} \frac{e^{x}}{e^{x} + 4} dx$$

- Sketch the curves  $y = x + \frac{3}{x}$  and y = 4, and find the area enclosed. [16] **6.** 
  - Find the volume of a solid obtained by rotating the region bounded by the curve  $y = \sin(x)$  and the x-axis on the interval  $[0, \pi]$  about the axis y = -1
  - Find the exact average value of  $f(x) = \tan^2 x$  on the interval  $[0, \pi/4]$ .
- Find the limit of the sequence  $\{a_n\}$  at  $n \to \infty$  or prove that it does not exist:

(a) 
$$a_n = \frac{(3^n + 1)^2}{6^n}$$
 (b)  $a_n$ 

(b) 
$$a_n = \ln(1+2n^2) - 2\ln(1+n)$$

Determine whether the series is divergent or convergent, and if convergent, whether absolutely or conditionally:

(a) 
$$\sum_{n=1}^{\infty} \frac{(-1)^n \sqrt{1+n^3}}{n^2}$$
 (b)  $\sum_{n=0}^{\infty} \frac{(-3)^n}{5+e^n}$  (c)  $\sum_{n=2}^{\infty} \frac{\sin(n)}{n^2}$ 

(b) 
$$\sum_{n=0}^{\infty} \frac{(-3)^n}{5+e^n}$$

(c) 
$$\sum_{n=2}^{\infty} \frac{\sin(n)}{n^2}$$

- 9. Find (a) the radius of convergence, and (b) the interval convergence of the series  $\sum_{n=0}^{\infty} \frac{(x+1)^n}{(n+1) 2^n}$ .
- [8] 10. (a) Use the integrability of the power series to express the function

$$F(x) = \int_{0}^{x} \left( \sum_{n=1}^{\infty} n t^{2n-1} \right) dt$$
 as an elementary function

(i.e. sum the series for F(x) within the radius of its convergence).

- (b) Find the MacLaurin series for the function  $x e^{-x^2}$ . (Hint: start with the series for  $e^z$  then replace z by  $-x^2$ )
- [5] **Bonus Question.** A solid is generated by rotating about the x-axis the region under the curve y = f(x), where f is a positive function and x > 0. The volume generated by the part of the curve from x = 0 to x = b is  $\pi b^2$  for all  $b \ge 0$ . Find f(x).

The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.