

CONCORDIA UNIVERSITY
Department of Mathematics & Statistics

Course	Number	Sections
Mathematics	205	All
Examination	Date	Pages
Final	April 2019	2
Instructors:	R. Bairakdar, B. Falahat, H. Greenspan, T. Hughes, S. Ky, A. Panait, S. Zaman	Course Examiner A. Atoyan, H. Proppe
Special Instructions:	Only approved calculators are allowed Show all your work for full marks	

MARKS

- [10] **1.** (a) Sketch the graph of $f(x) = 2x + x^2$ on the interval $[-1, 2]$, and write in sigma notation the formula for the right Riemann sum R_n for $f(x)$ with partitioning of the interval $[-1, 2]$ into n subintervals of equal length. Then calculate $\int_{-1}^2 f(x) dx$ as the limit of R_n at $n \rightarrow \infty$

NOTE: you may need the formulas $\sum_{k=1}^n k = \frac{n(n+1)}{2}$, $\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$.

- (b) Use the Fundamental Theorem of Calculus to calculate the derivative of $F(x) = \int_{-x^4}^0 \sin(t^2) dt$. Is the function F increasing or decreasing at $x = -1$?

- [15] **2.** Calculate the following indefinite integrals:

$$(a) \int \frac{x^2 + 3}{x^2 - 9} dx \quad (b) \int \sqrt{x} \ln x dx \quad (c) \int \frac{x^3}{\sqrt{x^2 + 1}} dx$$

- [6] **3.** Find the antiderivative $F(t)$ of the function $f(t) = \frac{e^t}{4 + e^{2t}}$ such that $F(\ln 2) = \frac{\pi}{8}$

- [12] **4.** Evaluate the following definite integrals (give the **exact answers**):

$$(a) \int_0^{\pi} \cos^2(x) \sin^3(x) dx \quad (b) \int_0^4 x^2 \sqrt{1 + 2x} dx$$

- [8] **5.** Evaluate the given improper integral or show that it diverges:

$$(a) \int_1^{\infty} \frac{1}{x(1 + \ln x)^3} dx \quad (b) \int_0^1 \frac{1}{(1 - x)^{2/3}} dx$$

- [17] **6.** (a) Sketch the curves $x = y^2 - 4y$ and $x = 2y - y^2$, find their points of intersection, and find the area enclosed by the curves.
- (b) Sketch the region between $y = \sin(x)$ and the x -axis on the interval $x \in [0, \pi]$, and find the volume of the solid generated by rotating this region about the line $y = -1$.
- (c) Suppose $g(x) = \int_0^x f(t) dt$ where $f'(t) > 0$ for all t and $f(1) = 0$. Investigate whether $g(x)$ has (i) a local maximum or local minimum, (ii) an inflection point, or (iii) none of the above at $x = 1$. **Explain.**

- [6] **7.** Find the limit of the sequence $\{a_n\}$ or prove that the limit does not exist:

$$(a) \quad a_n = \frac{2n^2 \cos(\pi n)}{\sqrt{100 + 4n^4}} \quad (b) \quad a_n = \sqrt{n+1} - \sqrt{n}$$

- [12] **8.** Determine whether the series is divergent or convergent, and if convergent, whether absolutely or conditionally:

$$(a) \quad \sum_{n=1}^{\infty} \frac{(-1)^n n}{1 + n^2} \quad (b) \quad \sum_{n=1}^{\infty} \frac{(-2)^{10n}}{n!} \quad (c) \quad \sum_{n=2}^{\infty} \frac{\sin(n)}{n^2}$$

- [6] **9.** Find (a) the interval of convergence, and (b) the radius of convergence of the series

$$\sum_{n=1}^{\infty} \frac{(6x+3)^n}{n+1}$$

- [8] **10.** (a) Derive the Maclaurin series of $f(x) = x^2 \ln(1+2x)$

(HINT: start with the series for $\ln(1+z)$ where $z = 2x$).

- (b) Use the differentiability of power series to find the sum $S(x) = \sum_{n=1}^{\infty} \frac{x^{2n}}{n}$ in the form of an elementary function within the radius of convergence of $S(x)$.

(HINT: find first the sum for the derivative of $S(x)$, then antidifferentiate).

- [5] **Bonus question.** If f is continuous, prove that

$$\int_0^{\pi/2} f(\cos x) dx = \int_0^{\pi/2} f(\sin x) dx$$

The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. No part of the present document may be sold, reproduced or re-disseminated in any manner or form without the written permission of its owner and copyright holder.