

## RESEARCH ARTICLE

# Hydrocarbon biodegradation by Arctic sea-ice and sub-ice microbial communities during microcosm experiments, Northwest Passage (Nunavut, Canada)

Marie-Ève Garneau<sup>1,\*</sup>, Christine Michel<sup>1</sup>, Guillaume Meisterhans<sup>1</sup>, Nathalie Fortin<sup>2</sup>, Thomas L. King<sup>3</sup>, Charles W. Greer<sup>2</sup> and Kenneth Lee<sup>4</sup>

<sup>1</sup>Fisheries and Oceans Canada, Marine Productivity Laboratory, 501 University Crescent, Winnipeg, MB R3T 2N6, Canada, <sup>2</sup>Energy, Mining and Environment, National Research Council of Canada, 6100 Royalmount Avenue, Montreal, QC H4P 2R2, Canada, <sup>3</sup>Fisheries and Oceans Canada, Bedford Institute of Oceanography, PO Box 1006, Dartmouth, NS B2Y 4A2, Canada and <sup>4</sup>Oceans and Atmosphere National Research Flagship, Australian Resources Research Centre, CSIRO, 26 Dick Perry Avenue, Kensington, WA 6151, Australia

\*Corresponding author: Fisheries and Oceans Canada Central and Arctic Region, Marine Productivity Laboratory, 501 University Crescent, Winnipeg, MB R3T 2N6, Canada. Tel: 1-204-984-8726; E-mail: [me.garneau@gmail.com](mailto:me.garneau@gmail.com)

One sentence summary: Ice-associated microorganisms of the Arctic Ocean can degrade petroleum at  $-1.7^{\circ}\text{C}$  within 15 days, and their response to oil varied whether they lived in ice or in underlying waters.

Editor: Max Hägglom

## ABSTRACT

The increasing accessibility to navigation and offshore oil exploration brings risks of hydrocarbon releases in Arctic waters. Bioremediation of hydrocarbons is a promising mitigation strategy but challenges remain, particularly due to low microbial metabolic rates in cold, ice-covered seas. Hydrocarbon degradation potential of ice-associated microbes collected from the Northwest Passage was investigated. Microcosm incubations were run for 15 days at  $-1.7^{\circ}\text{C}$  with and without oil to determine the effects of hydrocarbon exposure on microbial abundance, diversity and activity, and to estimate component-specific hydrocarbon loss. Diversity was assessed with automated ribosomal intergenic spacer analysis and Ion Torrent 16S rRNA gene sequencing. Bacterial activity was measured by  $^{3}\text{H}$ -leucine uptake rates. After incubation, sub-ice and sea-ice communities degraded 94% and 48% of the initial hydrocarbons, respectively. Hydrocarbon exposure changed the composition of sea-ice and sub-ice communities; in sea-ice microcosms, Bacteroidetes (mainly *Polaribacter*) dominated whereas in sub-ice microcosms, the contribution of *Epsilonproteobacteria* increased, and that of *Alphaproteobacteria* and *Bacteroidetes* decreased. Sequencing data revealed a decline in diversity and increases in *Colwellia* and *Moritella* in oil-treated microcosms. Low concentration of dissolved organic matter (DOM) in sub-ice seawater may explain higher hydrocarbon degradation when compared to sea ice, where DOM was abundant and composed of labile exopolysaccharides.

**Keywords:** Arctic; sea ice; hydrocarbon biodegradation; bioremediation; 16S ribosomal RNA gene sequencing; microbial community fingerprinting

## INTRODUCTION

The warming of the Arctic drives an accelerated decline in sea-ice cover extent and duration, which opens navigation routes and improves access to natural resources (Meier et al. 2014; Kwok and Cunningham 2015). Extensive offshore oil and gas activities have been taking place in the Arctic since the 1940s (AMAP 2010), but increased accessibility made possible by a longer open water season has created new opportunities for the petroleum industry (AMAP 2012; Meier et al. 2014). Higher risks of accidental hydrocarbon releases in the Arctic, either as oil spills or discharges from ships, are anticipated. One of the major transportation routes, the Northwest Passage in the Canadian Arctic Archipelago, has seen an increase in vessel transits in recent years (AMAP 2012; Michel 2013), thereby making the area particularly vulnerable to accidental leaks of hydrocarbons. Hydrocarbon pollution poses a major threat to Arctic marine ecosystems, particularly to the survival of seabirds and marine mammals (Meltofte et al. 2013). Oil may persist in the Arctic marine environment longer than in southern regions in part because of the challenges associated with oil spill clean-up in ice-infested waters (Meltofte et al. 2013).

While mechanical and chemical methods used to remove hydrocarbons from contaminated sites have limited effectiveness and can be expensive, microbial degradation of hydrocarbons appears as a non-invasive and relatively cost-effective bioremediation technique (Das and Chandran 2011). This natural process is responsible for the removal of the last traces of oil along polluted shores (ITOPF 2011). In marine environments, oil biodegradation is mostly attributable to bacteria, with an efficiency ranging from 0.003% to 100%, whereas degradation by fungi and algae (Das and Chandran 2011) and archaea (Brakstad and Bonaunet 2006) are considered negligible. Hydrocarbon-degrading bacteria exist in very low abundance (usually less than 1%) in the world's marine environments (Atlas and Cerniglia 1995), but may reach 10% of the total bacterial population in contaminated sites, especially where the pollution is chronic such as in harbors (Hassanshahian and Cappello 2013). The complete degradation of hydrocarbon components requires a large consortium of indigenous bacteria with overall broad enzymatic capabilities (Leahy and Colwell 1990) and the presence of microorganisms possessing adequate metabolic potential (Das and Chandran 2011).

Rates of biodegradation depend on oil characteristics, temperature and the availability of oxygen and nutrients (mainly nitrogen in marine systems) (Das and Chandran 2011; ITOPF 2011). Temperature also directly affects oil contaminant solubility and viscosity, with degradation efficiency decreasing at low temperatures (Atlas 1981; Das and Chandran 2011). Nonetheless, cold-adapted bacteria that possess hydrocarbon-degrading genes have been found in Arctic soils (Whyte et al. 1999; Yergeau et al. 2012) and seawater (McFarlin et al. 2014). Sea ice is another distinctive Arctic ecosystem, characterized by very cold temperature and high brine content, and yet it harbors abundant and diverse microbial assemblages thriving in several distinct microhabitats (Bowman et al. 2012). Some studies have addressed the hydrocarbon degradation potential of ice-associated microbes (Atlas, Horowitz and Busdosh 1978; Deppe et al. 2005; Gerdes et al. 2005; Brakstad et al. 2008), but overall this area of research is rather unexplored, especially regarding first-year sea ice. The identification of key oil-degrading microorganisms that live in and below the sea ice is essential for the development of bioremediation strategies suitable for the ice-covered Arctic Ocean.

The Canadian Arctic Archipelago is a region of high biological production that supports large populations of seabirds and marine mammals (McLaughlin et al. 2004; Michel, Ingram

and Harris 2006). Bottom ice maintains the highest ice-algal biomass reported for the Arctic (Arrigo, Mock and Lizotte 2010) and contains elevated sea-ice bacterial abundances (Smith et al. 1989; Maranger, Bird and Juniper 1994). We studied the potential for hydrocarbon degradation of natural microbial communities thriving in first-year sea ice and sub-ice seawater, in the eastern part of the Northwest Passage during spring 2012. The objectives were to evaluate (i) the diversity of bacterial communities and the presence of oil-degrading taxa; (ii) the hydrocarbon degradation potential, including preferential degradation of certain components; and (iii) changes in communities induced by hydrocarbon exposure in terms of abundance, activity and diversity. We hypothesized that (i) the hydrocarbon degradation rates would be different in sea ice and sub-ice seawater because of variations in microbial abundance, activity and environmental conditions and that (ii) exposure to hydrocarbons would favor oil-degrading taxa. These hypotheses were tested in microcosms, in which natural microbial communities collected during the ice-algae bloom period were exposed to persistent hydrocarbons. Our study is one of the few that has combined data on hydrocarbon components and bacterial metabolic rates, as well as phylogenetic analysis and community fingerprinting, to investigate oil degradation by ice-associated bacteria.

## MATERIALS AND METHODS

### Study site and sample collection

Samples were collected on 1 May 2012 at a station located on the landfast ice of the Resolute Passage (74.7°N, 95.5°W), a subsection of the Northwest Passage in the Canadian Arctic Archipelago (Fig. 1A and B). The Northwest Passage is the main channel across the archipelago, linking Baffin Bay in the east to Beaufort Sea in the west (Fig. 1A). The region is extensively covered by first-year ice during most of the year, i.e. from mid/late November to the end of June. The Northwest Passage is influenced by different water masses, mainly Pacific waters to the west, Atlantic waters to the east and modified Arctic waters flowing from the northern channels (McLaughlin et al. 2004).

Two types of samples were collected: first-year sea-ice cores and samples of the seawater immediately under the ice cover, hereafter referred to as sub-ice seawater. Sub-ice seawater receives brine rejections during sea-ice formation from autumn to spring, at times making it more saline than the surface seawater found in ice-free areas (Petrich and Eicken 2009). Salinity in the sub-ice samples was 32.0 (Table 1), which is comparable to what is typically found at surface in Barrow Strait during open-water periods (McLaughlin et al. 2004). A total of 40 l of sub-ice seawater was collected with a pump submerged at the ice-water interface; 20 l was used for ancillary measurements and 20 l was used in microcosms (see below). The water was stored in acid-washed containers in the dark at -1°C until further processing (see below). A total of 13 sea-ice cores were collected using a manual ice corer (Mark II coring system, 9 cm internal diameter, Kovacs Enterprises, Roseburg, OR, USA). The ice cores were cut with a stainless steel saw to collect the bottom 3 cm, where most of the microbial biomass is found (e.g. Michel et al. 1996). Sterile gloves were worn at all times when handling the cores. All cores were kept in sterile Whirl-Pak® bags or in an acid-washed isothermal container in cold, dark conditions until further processing and analysis.

Initial conditions (T0) in sea ice and sub-ice seawater were characterized in terms of microbial variables, i.e. autotroph abundance, prokaryote abundance and heterotrophic activity (herein referred to as bacterial abundance and bacterial




Figure 1. Canadian Arctic Archipelago (A) showing the sampling region in Resolute Passage, in the eastern part of the Northwest Passage (B). The star identifies the location of the sampling station.

**Table 1.** Environmental conditions and microbial biomass and production in sea ice and sub-ice water at a landfast ice station in Resolute Passage (Nunavut, Canada) on 1 May 2012. Values are average of duplicate measurements ( $\pm$  standard error), except for salinity, nitrate and total bacterial abundance ( $N = 1$ ) and bacterial production (BP;  $N = 3$ ).  $\text{NO}_3^-$ : nitrate;  $\text{PO}_4^{3-}$ : phosphate;  $\text{Si(OH)}_4$ : silicic acid; DOC: dissolved organic carbon; DN: dissolved nitrogen.

	Sea ice	Sub-ice seawater
Salinity	8.0	32.0
$\text{NO}_3^-$ (mmol m $^{-3}$ )	$12.58 \pm 0.2$	$7.08 \pm 0.04$
$\text{PO}_4^{3-}$ (mmol m $^{-3}$ )	$8.53 \pm 0.09$	$1.07 \pm 0.00$
$\text{Si(OH)}_4$ (mmol m $^{-3}$ )	$6.83 \pm 0.23$	$12.00 \pm 0.06$
DOC (mmol m $^{-3}$ )	$2030 \pm 358$	$59.5 \pm 2.8$
DN (mmol m $^{-3}$ )	$126 \pm 29$	$10.4 \pm 0.3$
Chlorophyll <i>a</i> (mg m $^{-3}$ )	$4563 \pm 391$	$0.18 \pm 0.05$
Total protists (cells ml $^{-1}$ )	$2.1 \times 10^6$	$1.0 \times 10^2$
Bacteria (cells ml $^{-1}$ )	$16 \times 10^6$	$2.3 \times 10^5$
BP (pmol leu L $^{-1}$ h $^{-1}$ )	$441 \pm 18$	$7.83 \pm 0.72$

production, respectively). Biochemical analyses were performed to determine concentrations of nutrients, dissolved organic carbon (DOC), dissolved nitrogen (DN) and salinity. Microbial variables in sea ice were measured on three ice cores that were previously combined in an acid-washed isothermal container and melted in filtered surface seawater (0.22  $\mu\text{m}$ , 142 mm diameter GV Durapore® filter, Millipore, Darmstadt, Germany). The addition of 500 ml filtered seawater per core gave an average dilution factor of 3.7. One core was kept in a sterile Whirl-Pak bag for salinity, nutrients, DOC and DN analyses without the addition of filtered seawater to avoid a dilution effect of dissolved materials.

Salinity was measured with a salinometer (Portasal 8410A, Guildline Instruments Limited, Smiths Falls, Ontario, Canada) according to Müller (1999). Inorganic nutrients, i.e. nitrate ( $\text{NO}_3^-$ ), phosphate ( $\text{PO}_4^{3-}$ ) and silicic acid ( $\text{Si(OH)}_4$ ), were analyzed on a SmartChem discrete analyzer (Westco Scientific Instruments Inc., Brookfield, CT, USA) according to Hansen and Koroleff (1999). DOC and DN concentrations were determined by high-temperature catalytic combustion on a Shimadzu TOC-VCPh analyzer with an ASI-V auto sampler and TNM-1 Total Nitrogen module (Knap et al. 1996). Chlorophyll *a* (chl *a*), a proxy for autotroph biomass, was measured by fluorometry using 10AU Turner Designs (San Jose, CA, USA) after filtration onto Whatman GF/F 25mm filters, according to Parsons, Maita and Lalli

(1984). Protist and bacterial cell numbers, as well as high nucleic acid content (HNA) bacteria, were stained with SYBR Green I and counted on an Epics Altra flow cytometer (Beckman Coulter, Brea, CA, USA) according to Belzile et al. (2008).

### Microcosm experiments

The remaining 20 l of sub-ice seawater and nine ice cores were used in microcosm experiments, which started the day following sample collection. The ice cores were combined with 4.5 l of 0.2  $\mu\text{m}$  filtered surface seawater (total volume of 6.2 l) into a 9 l acid-washed carboy (Nalgene, Rochester, NY, USA), and were left to melt slowly overnight in a cold room in the dark. The dilution factor was 3.6. Melted and diluted sea ice samples are convenient and routinely used because microbes in the liquid phase can be reached by stable isotope additions while at the same time be protected from osmotic shock and cell lysis caused by the fresher salinities of melt water (Deming 2010). The use of melted sea ice in the microcosms also ensured that the entire sea-ice bacterial community was exposed to hydrocarbons. This approach was based on previous experimental results showing that upward infiltration of hydrocarbons in sea-ice channels was very low in May (Karlsson, Petrich and Eicken 2011).

Control microcosms without oil were set up in duplicate for both sub-ice seawater and diluted ice samples. Triplicate microcosms of sub-ice seawater and diluted sea-ice were spiked with 25 mg L $^{-1}$  of Arabian Light oil. The oil was previously weathered to remove volatiles. Arabian Light oil is a moderately waxy crude oil (McKetta 1997) and has a pour point of -40°C (ITOPF 2011). The API gravity (i.e. oil density in relation to pure water, a standard established by the American Petroleum Institute) of this oil varies between 31.8° and 33.4° (Environment Canada 1991), which indicates that it is lighter than water (which has an API gravity of 10°). The oil is classified in Group 3 (°API 17.5–35) and is considered persistent (ITOPF 2011).

Samples were incubated for 15 days in oil-treated and control microcosms and those were gently mixed by a custom-made rotating shaker. Incubations were run in sterile transparent Nalgene polycarbonate bottles at near *in situ* temperature (-1.7  $\pm$  0.02°C; average  $\pm$  standard error) and under light conditions comparable to those under the sea ice (7.8  $\pm$  0.8  $\mu\text{E m}^{-2} \text{s}^{-1}$ ). Temperature was recorded every 10 min during the incubation period, and light was measured on 10 occasions. Average under-ice PAR at sampling stations visited was 9.4  $\mu\text{E m}^{-2} \text{s}^{-1}$ . Water was collected from each microcosm and control at five time points (T), i.e. days 0, 3, 7, 11 and 15, using 250 ml Nalgene

bottles for sea ice and 250 ml or 1 l Nalgene bottles for sub-ice seawater, depending on the analytical volume required. Bacterial production and hydrocarbon composition were measured at each time point. Bacterial abundance was analyzed at the beginning (T0) and at the end (T15) of the experiment by flow cytometry (as mentioned above). Community composition was evaluated using automated ribosomal intergenic spacer analysis (ARISA) fingerprinting and 16S rRNA gene sequencing at T0 and T15.

### Bacterial production

Bacterial production was measured from the incorporation rates of the titrated ( $^3\text{H}$ -) amino acid leucine (leu) according to the centrifugation protocol (Smith and Azam 1992). This approach measures the production rate of proteins (Kirchman and Ducklow 1993) in both bacteria and archaea (Herndl et al. 2005), although we assume a minor contribution of archaea in our samples, as observed on the Beaufort Sea shelf (< 1% to 7% of archaea in total DAPI counts; Garneau et al. 2006) and in the Northwest Passage (2% to 13%; Wells and Deming 2003). Triplicate subsamples of 1.2 ml were spiked with  $^3\text{H}$ -leucine (specific activity, 60 Ci mmol $^{-1}$ ) to give a final concentration of 10 nM. Duplicate controls were immediately spiked with 50% trichloroacetic acid (TCA; 5% final concentration) to kill the bacteria. Vials for uptake measurement and killed control vials were incubated in the dark at  $-1.7^\circ\text{C}$  for 4 h. At the end of the incubation, TCA (5% final concentration) was immediately added to the vials. All samples were frozen at  $-80^\circ\text{C}$  until final analysis in our main laboratory. Samples were then thawed at  $4^\circ\text{C}$  and centrifuged at 14 000 rpm for 10 min. The supernatant was removed and pellets were rinsed by adding 1 ml of TCA (5% final concentration). This rinsing TCA was removed after a second centrifugation. Scintillation cocktail (Ecolume, MP Biomedicals, Santa Ana, CA, USA) was added to the vials, and bacterial cells were resuspended by vortex mixing.  $^3\text{H}$ -Leucine incorporation was measured using a liquid scintillation counter (TRI-CARB 2100 TR, Packard Bio-Science, Meriden, CT, USA) after 48 h of incubation in the dark at  $4^\circ\text{C}$ . Data are given in pmol leu L $^{-1}$  h $^{-1}$ .

### Hydrocarbon composition

Gas chromatography-mass spectroscopy (GC-MS) was used to quantify the different petroleum components at each experimental time point. Alkanes, polycyclic aromatic hydrocarbons and their alkylated homologs represent > 80% weight of the total oil in Arabian Light oil (Environment Canada 2001). Thus, these three major groups of hydrocarbon compounds were analyzed: (i) alkanes, including 28 linear *n*-alkanes ranging in carbon number from C<sub>10</sub> to C<sub>35</sub>, the branched alkanes pristane and phytane, and the cyclic alkane hopane; (ii) polycyclic aromatic hydrocarbons (PAHs) consisting of the 2-, 3- and 4-ring aromatics (anthracene, chrysenes, dibenzothiophenes, fluorenes, fluoranthenes, naphthalenes, naphthobenzothiophenes, perylenes, phenanthrenes, pyrenes); and (iii) the methylated homologs of these aromatics (C<sub>1</sub>- to C<sub>4</sub>-methylated PAHs). These analytes were selected because they are most likely to contribute to toxicity in fish exposed to an oil spill, as the high-molecular-weight (HMW) resins and asphaltenes are not expected to cross fish tissue membranes. In parallel, we also analyzed 9 internal standards (5 $\alpha$ -androstane, d<sub>10</sub>-anthracene, d<sub>12</sub>-chrysene, d<sub>12</sub>-perylene, d<sub>22</sub>-n-decane, d<sub>34</sub>-hexadecane, d<sub>42</sub>-n-eicosane, d<sub>52</sub>-n-tricontane and d<sub>8</sub>-naphthalene) and 10 surrogate standards (5 $\beta$ -cholestane, d<sub>10</sub>-phenanthrene,

d<sub>10</sub>-pyrene, d<sub>12</sub>-benzo[a]pyrene, d<sub>12</sub>-benzo[b]fluoranthene, d<sub>14</sub>-dibenz[a,h]anthracene, d<sub>26</sub>-dodecane, d<sub>36</sub>-heptadecane, d<sub>50</sub>-n-tetracosane and d<sub>66</sub>-n-dotriacontane). A GC-MS detection of <100 ng L $^{-1}$  was applied.

Sub-ice seawater and sea-ice samples (100 ml) were placed in clean amber glass bottles sealed with Teflon tape after adding dichloromethane (5% final concentration). The samples were refrigerated until analysis, which was performed within 4 months of collection. Hydrocarbons were extracted in dichloromethane and quantified using an Agilent 6890 Series GC with a 5973N MSD operated in the selected ion-monitoring mode. The column was a 30 m  $\times$  0.25 mm internal diameter, with 0.2  $\mu\text{m}$  film thickness MS-5 or equivalent column (Supelco Pack (Supelco, Bellefonte, PA, USA)). Concentrations of the analytes at a given time point were divided by the concentration of the internal marker 17 $\alpha$ (H),21 $\beta$ (H)-hopane to avoid any bias caused by physico-chemical losses of hydrocarbons (Prince et al. 1994). Hopane-normalized concentrations allow for the measurement of oil biodegradation as the internal marker biodegrade very slowly. The coefficients of variation on the measured concentrations (expressed in ng L $^{-1}$ ) between the triplicate filters confirmed that the precision and repeatability of the assay were good. The average coefficient of variation was 1% (N = 81) for the internal standards, 9% (N = 90) for the surrogate standards and 19% for the analytes (N = 429). The hydrocarbon loss (in %) was calculated as HC<sub>loss</sub> = (HC<sub>T0</sub> - HC<sub>Tx</sub>)  $\div$  HC<sub>T0</sub>  $\times$  100, where HC<sub>T0</sub> is the hydrocarbon concentration at T0 (initial concentration) and HC<sub>Tx</sub> is the hydrocarbon concentration at sampling time x.

Hydrocarbon compound concentrations in seawater and melted ice cores were measured before the addition of Arabian Light oil. Methyl-PAH and PAH concentrations were below detection. Alkanes were detected but were, on average, at < 4% of initial concentrations in spiked bottles ( $5.3 \pm 3.5 \times 10^4$  ng L $^{-1}$ ). Unfortunately, contamination occurred in T15 sub-ice samples during the preservation and results were excluded from the data analysis.

### Bacterial community fingerprinting

ARISA fingerprinting (Fisher and Triplett 1999) was used to determine bacterial community structure under the experimental conditions. Sub-ice seawater (1400 ml) and bottom sea-ice (75 ml) samples that were taken at T0 and T15 were filtered onto 0.2  $\mu\text{m}$  sterile filters (Millipore, 47 mm diameter). DNA was extracted by bead-beating and purified using a PowerWater DNA isolation kit (MO BIO Laboratories, Carlsbad, CA, USA). The amount of DNA was then determined by spectrofluorometry (Tecan Safire, Tecan Group Ltd, Männedorf, Switzerland) using Quant-iT PicoGreen dsDNA assay (Invitrogen, Carlsbad, CA, USA). The PCR amplification of the 16S–23S rRNA intergenic transcribed spacer (ITS) region was conducted in triplicate for each DNA extract and according to Ranjard et al. (2000). We used the universal bacterial primers SDBact (eubacterial rRNA gene small subunit, 5'-TGC GGC TGG ATC CCC TCC TT-3') and LDBact (eubacterial rRNA gene large subunit, 5'-CCG GGT TTC CCC ATT CGG-3') (Normand et al. 1996). The primers were synthesized by Molecular Probes Invitrogen. The primer SDBact was labeled at the 5' end with the phosphoramidite dye 5-FAM fluorochrome for subsequent analysis by ARISA. The three-step thermal PCR amplification protocol was as follows: (i) 94 $^\circ\text{C}$  for 5 min; (ii) 35 cycles of 94 $^\circ\text{C}$  for 1 min, 55 $^\circ\text{C}$  for 1 min, 72 $^\circ\text{C}$  for 1 min; and (iii) 72 $^\circ\text{C}$  for 5 min. The 25  $\mu\text{l}$  reaction mixtures contained 1  $\times$  PCR buffer, 1.5 mM MgCl<sub>2</sub>, 0.2 mM of each deoxynucleoside triphosphate (dNTP), 0.1  $\mu\text{M}$  of each primer, 1 U of rTaq polymerase

(GE Healthcare, Little Chalfont, UK) and 10 ng of DNA template. Triplicate amplified ITS fragments were pooled together and then purified using a QIAquick PCR Purification Kit (QIAGEN, Hilden, Germany) to eliminate primer surplus that could interfere during electrophoresis. The bacterial ITS fragments were quantified by spectrofluorometry.

Bacterial ITS products were denatured using Hi Di formamide followed by a heat treatment at 95°C for 5 min. An internal fluorescent size marker (LIZ1200) was added to each sample and 10 ng of bacterial product were run on a 3130XL Capillary Genetic Analyzer (Applied Biosystems, Foster City, CA, USA) using a 50 cm capillary. ARISA electrophoresis profiles were analyzed with Genemapper software (Applied Biosystems). Peak lengths and absolute areas that characterized the different operational taxonomic units and their abundances were used in data analysis. Peaks with a length inferior to 200 bp and superior to 1200 bp were considered noise and were thus excluded from further analysis. The area of each peak was divided by the total area of the sample to obtain relative abundance (%). Background and electrophoresis internal deviation were eliminated using the variable percentage threshold method (Osborne et al. 2006) and the binning procedure under R environment (interactive and automatic binning algorithms; Ramette 2009). Data were transformed to abundance matrices, from which Bray–Curtis dissimilarities ( $BC_{diss}$ ) between samples were calculated using the Primer software (Clarke and Warwick 2001). The complement of the Bray–Curtis measure was used to calculate the similarities.

### **Ion Torrent 16S rRNA gene sequencing**

At the beginning of the experiment (T0), triplicate samples of sub-ice seawater (1400 ml) and melted sea ice (75 ml) were filtered onto 0.2 µm sterile filters (Millipore, 47 mm diameter) for DNA extraction using hexadecyltrimethylammonium bromide (CTAB), following a slightly modified version of the method published by Ausubel et al. (2002). A phenol/chloroform extraction was first performed and was followed by a chloroform/isoamyl alcohol (24:1) extraction. This extraction method gives superior DNA purity than the bead-beating method we used for DNA fingerprinting, but it produces toxic phenol–chloroform wastes. The DNA was precipitated overnight at -20°C. The DNA pellet was resuspended in 100 µl of sterile deionized water and quantified using the Quant-iT PicoGreen dsDNA assay (Invitrogen). The water from triplicate oiled microcosms was harvested at T15. The control treatments at T0 and T15 were in duplicate. These samples were filtered and extracted as described above for the initial sub-ice seawater and melted sea-ice samples.

Partial 16S ribosomal (rRNA) gene amplicons were produced using the universal primers F343 (5'-TAC GGR AGG CAG CAG-3') and R534 (5'-ATT ACC GCG GCT GCT GGC-3') as in Yergeau et al. (2014). The primers contained 10 bp multiplex identifiers (MIDs) and adaptor sequences for Ion Torrent sequencing (Yergeau et al. 2012; Bell et al. 2013). Reactions were performed in 25 µl volumes containing 10 ng of DNA, 0.3 µM of each primer, 0.4 mg ml<sup>-1</sup> of bovine serum albumin, 0.2 µM of dNTPs and 0.05 U µl<sup>-1</sup> of rTaq DNA polymerase (GE Healthcare). The PCR cycling conditions involved an initial denaturation step at 95°C for 5 min, followed by (i) 30 cycles of 30 s at 95°C, (ii) 30 s at 55°C, (iii) 45 s at 72°C and (iv) a final elongation step of 3 min at 72°C. The PCR products were purified on agarose gel using the QIAquick gel extraction kit from QIAGEN and quantified with the Quant-iT PicoGreen dsDNA assay (Invitrogen). All of the 20 purified products were pooled in equimolar ratios. A total of  $39.15 \times 10^7$  molecules was used in the

emulsion PCR using the Ion PGM Template OT2 200 and the One-Touch instrument (Life Technologies, Carlsbad, CA, USA). Community composition assessment was performed using the RDP pyrosequencing pipeline (<http://pyro.cme.msu.edu/>); sequences were separated based on their adapter, as well as deconvoluted and binned according to their respective MIDs.

Sequencing was performed on an Ion Torrent personal genome machine using the Ion 314 chip and the Ion PGM Sequencing 200 kitV2 (Life Technologies) following the manufacturer's instructions. Sequences with an average Phred quality score below 20 or a length of less than 150 were filtered out of the data set. Taxonomic identities were assigned by classifying sequences in parallel with a bootstrap value of 80% using the MultiClassifier tool from the Ribosomal Database Project Pipeline Classifier (Release 9; Wang et al. 2007). Number of reads before and after quality control generated during the 16Sr RNA gene sequencing analysis are presented in Table S1 (Supporting Information). The results of the replicate sequences belonging to a given taxonomic level were averaged and expressed as a percentage of all sequences. Raw sequence data produced in this study was deposited in NCBI under the Bioproject accession PRJNA321190.

### **Statistical analysis**

Differences between two groups (i.e. control versus oil-treated microcosms; ice versus sub-ice seawater) were tested with the R software (R Core Team 2013). Student's t-test was used since data distributions were not far from a normal distribution and variances were equal. Differences between incubation days were tested with the Kruskal–Wallis test (KW) and multiple comparison post-hoc tests were run using the kruskalmc function in the pgirmess package of R software (Giraudoux 2014). Mean values are presented with standard error (SE).

Bray–Curtis similarities were illustrated using non-metric multidimensional scaling (MDS), where each data point represents a bacterial assemblage, and the distance between data points represents the level of similarity in bacterial assemblages; the longer the distance, the lower the similarity. Bacterial community composition was assessed using ANalysis Of SIMilarity (ANOSIM), based on a random Monte-Carlo test of 100 000 permutations (Primer V06 software; Clarke and Warwick 2001). The test compares ranks of distances between groups with ranks of distances within groups. Resulting R values > 0.75 are interpreted as well distinct, R > 0.5 as different but overlapping and R < 0.25 as barely different (Clarke and Gorley 2001). The Mann–Withney U test (MW) was performed to test difference in assemblage similarity between (i) sub-ice seawater and sea-ice microcosms, (ii) controls and oil-treated microcosms and (iii) T0 and T15.

## **RESULTS**

### **Environmental parameters and initial experimental conditions**

Environmental conditions in sub-ice seawater and in sea ice at the time of the sampling were typical of the spring period (Table 1). Nutrient inventories were high in sub-ice seawater but bacterial ( $2.3 \times 10^5$  cells ml<sup>-1</sup>) and phytoplankton biomass (< 0.2 mg chla m<sup>-3</sup>) were low, indicating pre-bloom conditions. Pre-bloom nitrate concentrations in the surface waters of the Canadian Archipelago are typically ca. 10 mmol m<sup>-3</sup> (Codispoti

**Table 2.** Initial (T0) bacterial abundances in microcosms. Values are averages of duplicate control microcosms ( $\pm$  standard error). The asterisks indicate significant difference (t-test,  $P < 0.01$ ) in sea ice when compared to sub-ice seawater. HNA: high nucleic acid.


	Sea ice	Sub-ice seawater
Bacteria ( $\times 10^5$ cells $\text{ml}^{-1}$ )	4.2* $\pm$ 0.7	2.1 $\pm$ 0.0
HNA bacteria (% cells)	92* $\pm$ 1	64 $\pm$ 1
Bacterial production (pmol leu $\text{L}^{-1} \text{h}^{-1}$ )	21* $\pm$ 2	12.7 $\pm$ 2.3

et al. 2013), which is concordant with our nitrate concentrations (Table 1). Comparatively, sea ice had more nutrients and dissolved matter, and had higher bacterial ( $16 \times 10^6$  cells  $\text{ml}^{-1}$ ) and algal ( $> 4000$  mg chl  $a$   $\text{m}^{-3}$ ) biomasses (Table 1). Concentrations of DOC and chl  $a$ , and protist abundance were three to five orders of magnitude higher in the ice than in sub-ice seawater (Table 1), indicating that the ice-algal bloom was well underway.

For microcosm experiments, sea ice was diluted to reach a bacterial abundance comparable to that measured in sub-ice seawater (Table 2). Initial (T0) bacterial abundance and production in the microcosms, as well as the proportion of HNA bacteria, were significantly higher in the sea ice than in sub-ice seawater (t-test,  $P < 0.01$ ; Table 2). Nonetheless, values were of the same order of magnitude (Table 2).

### Bacterial production

Bacterial production during the incubation reached values that were about 10 times higher in sea ice (Fig. 2). Bacterial production in sub-ice microcosms at T0 was significantly higher in the controls than in the oil-treated microcosms (13 versus 2 pmol leu  $\text{L}^{-1} \text{h}^{-1}$ ; t-test,  $P < 0.01$ ; Fig. 2B). This situation was reversed during the remainder of the incubation (i.e. excluding T0) with a mean of  $6.5 \pm 1.0$  pmol leu  $\text{L}^{-1} \text{h}^{-1}$  in oil treatment compared to a mean of  $3.7 \pm 0.8$  pmol leu  $\text{L}^{-1} \text{h}^{-1}$  in controls (t-test,  $P < 0.05$ ; Fig. 2B).



**Figure 2.** Time series of bacterial production in sea ice (A) and sub-ice seawater (B) during microcosm experiments. Bars and vertical lines represent averages and standard errors, respectively, for oil-exposed ( $N = 3$ ) or control ( $N = 2$ ) microcosms. Note different scales.




Figure 3. Time series of total hydrocarbon concentrations normalized to hopanes (A) and cumulative percent loss of total hydrocarbons (B) in sea ice and sub-ice seawater during microcosm experiments. Squares and vertical lines represent averages and standard error, respectively, for oil-exposed microcosms ( $N = 3$ ). nd: no data.

In the experiments with sea ice, however, there was no significant difference in bacterial production between the oil-treated and control microcosms ( $t$ -test,  $P > 0.05$ ; Fig. 2A). The overall bacterial production in the sea-ice microcosms increased significantly over the course of the experiment, with a 4-fold increase from T0 to T7 ( $19 \pm 1$  versus  $248 \pm 4$  pmol leu L $^{-1}$  h $^{-1}$ ; KW,  $P < 0.05$ , Fig. 2A).

#### Hydrocarbon degradation in oil-treated microcosms

Total hydrocarbon concentrations at the beginning of the incubation (T0) were not significantly different between sea ice and sub-ice seawater, averaging  $1819 \pm 141$  hopane-normalized units ( $t$ -test,  $P > 0.05$ ; Fig. 3A). The 28 alkanes analyzed were de-

tected in both microcosms at T0 (data not shown). Three low-molecular-weight (LMW) PAHs (i.e. two or three rings; Canadian Council of Ministers of the Environment 1999) were also detected. They were phenanthrene, naphthalene and dibenzothiophene, but there was no PAH compound of HMW (four to seven rings; Canadian Council of Ministers of the Environment 1999). A total of 26 methyl-PAHs were detected in the microcosms at T0 (data not shown).

Total concentrations decreased in both sea-ice and sub-ice seawater microcosms during the experiment (Fig. 3). Concentration in sea-ice microcosms was  $972 \pm 92$  hopane-normalized after 15 days, but there was no significant difference between T0 and T15 ( $t$ -test,  $P > 0.05$ ). In sub-ice seawater, there was a significant difference between T0 and T11, at which time the

concentration was  $115 \pm 8$  hopane-normalized units remaining after 11 days (*t*-test,  $P < 0.005$ ).


The hydrocarbon loss rate in sea-ice microcosms was constant over the duration of the experiment, as shown by the consistent slope of percent hydrocarbon loss by incubation days (Fig. 3B). In these microcosms,  $1635 \pm 195$  hopane-normalized units remained at T3 (Fig. 3A), corresponding to a loss of  $12 \pm 11\%$  (Fig. 3B). Comparatively, in sub-ice seawater microcosms most of the total hydrocarbon loss ( $63 \pm 5\%$  of initial concentrations) occurred during the first days, with  $660 \pm 82$  hopane-normalized units remaining at T3 (Fig. 3A). The estimated loss rate of hydrocarbons between T0 and T3 was significantly higher in sub-ice seawater than in the sea-ice microcosms ( $371$  versus  $76$  hopane-normalized unit  $d^{-1}$ , respectively; test on slopes,  $P < 0.01$ ). On average, the loss of hydrocarbon components by the end of the incubation was 94% in sub-ice seawater and 48% in sea ice (Fig. 3B). Based on the time series of hydrocarbon percent loss, the overall half-life of crude oil in Resolute Passage would be ca. 3 days in seawater under ice cover, but at least 2 weeks when in contact with sea-ice bacteria under similar environmental conditions.

There was no significant difference in the initial (T0) amounts of the major groups of hydrocarbon compounds between sea-ice and sub-ice seawater microcosms (Fig. S1, Supporting Information), which consisted mainly of alkanes ( $85 \pm 1\%$ ), followed by methylated PAHs ( $15 \pm 1\%$ ) and PAHs ( $0.50 \pm 0.03\%$ ). These proportions remained unchanged from T0 to T15

in sea-ice (Fig. S1A, Supporting Information) microcosms (KW,  $P > 0.05$ ), but slightly changed in sub-ice seawater microcosms during the incubation although the difference was not statistically significant (Fig. S1B, Supporting Information). The final hydrocarbon composition in sub-ice seawater consisted of  $65 \pm 4\%$  alkanes,  $34 \pm 4\%$  methylated PAHs and  $1.8 \pm 0.3\%$  PAHs (T11).

The three major groups of hydrocarbon compounds showed different rates of decrease during the incubation period (Fig. 4). Alkanes showed the highest loss in terms of absolute concentrations, followed by methyl-PAHs and PAHs in both sea ice and sub-ice seawater, and this trend is especially evident in sub-ice microcosms (Fig. 4A and C). This also applied to the relative loss values (in %) measured in sub-ice water microcosms (Fig. 4D), but not to sea-ice microcosms where the PAHs experienced higher loss ( $43 \pm 5\%$ ) than methyl-PAHs ( $36 \pm 9\%$ ) throughout the experiment (Fig. 4B). Loss rates (in % per day) of alkanes and methyl-PAHs were significantly higher in sub-ice seawater than in sea ice (*t*-test,  $P < 0.01$ , Fig. 4).

Few aromatics were completely biodegraded in sea-ice microcosms (Fig. 5A); only  $C_1$ -methylated fluorene, phenanthrene and  $C_2$ -methylated pyrene were fully degraded. The loss of alkanes varied between 50% and 56%, whereas the loss of other compounds was below the average total hydrocarbon loss of 48% for sea ice (Fig. 5A). Comparatively, many aromatics were completely biodegraded (i.e. 100% loss) in sub-ice seawater:  $C_1$ - to  $C_3$ -methylated fluorenes, phenanthrene and its  $C_1$ - to



**Figure 4.** Time series of hopane-normalized concentrations of hydrocarbons and percent loss of main hydrocarbon compounds in sea ice (A and B) and sub-ice seawater (C and D) during microcosm experiments. Bars (or symbols) and vertical lines represent averages and standard deviations, respectively, for oil-exposed microcosms ( $N = 3$ ). Note logarithm scale on y-axis in (A) and (C). nd: no data.





Figure 5. Loss of hydrocarbon compounds in sea ice (A) and sub-ice seawater (B) after 15 and 11 days in microcosms, respectively.

C<sub>4</sub>-methylated forms, C<sub>1</sub>- to C<sub>4</sub>-methylated naphthobenzothiophenes, as well as C<sub>4</sub>-methylated naphthalenes, and the dibenzothiophene and its C<sub>3</sub>- and C<sub>4</sub>-methylated forms (Fig. 5B). The loss of linear *n*-alkanes (C<sub>10</sub> to C<sub>35</sub>) was close to the average total hydrocarbon loss of 94%, whereas the loss of the small methylated aromatics C<sub>0</sub>-C<sub>2</sub> naphthalenes and C<sub>1</sub>-C<sub>2</sub> dibenzothiophenes was below that average (Fig. 5B).

#### Bacterial community composition

Differences in bacterial community composition were observed between sub-ice seawater and sea-ice microcosms (ANOSIM, R = 0.77, P < 0.01; Fig. 6). The non-metric MDS plot had a 2D stress of 0.12, indicating good representations of the similarity matrices (Fig. 6). The initial (T0) bacterial community structure was similar in oil-enriched and control microcosms for a given



**Figure 6.** 2D MDS configuration of microcosms based on Bray–Curtis similarities of bacterial community composition in sub-ice seawater (black) and sea ice (gray) at the beginning (T0, circles) and end (T15, squares) of the incubation. Full symbols represent controls, and open symbols the oil-exposed microcosms. The 2D stress of 0.12 indicated good representations of the Bray–Curtis similarity matrices. Data points positioned closer together are similar. Microbial community structure was different in sub-ice and sea-ice microcosms (ANOSIM,  $R = 0.77$ ,  $P < 0.01$ ).

sample type; similarity was  $75 \pm 6\%$  (MW,  $P < 0.01$ ) in sea ice and  $69 \pm 5\%$  (MW,  $P < 0.01$ ) in sub-ice seawater. There was a significant difference in bacterial community between T0 and T15 in sea ice (ANOSIM,  $R = 1$ ,  $P < 0.05$ ), but at T15 the control and oil-treated microcosms were similar (MW,  $P > 0.05$ ; Fig. 6). Significant difference between T0 and T15 was also observed in oil-treated microcosms for the sub-ice seawater (ANOSIM,  $R = 0.61$ ,  $P < 0.05$ ). The similarity in community structure was significantly higher between T0 and T15 in the controls ( $38 \pm 6\%$ , MW,  $P < 0.01$ ) when compared to the similarity in oil-treated microcosms over the same period ( $26 \pm 7\%$ , MW,  $P < 0.01$ ). In other terms, the bacterial community composition in sub-ice seawater changed more in oil-enriched microcosms than in controls after the incubation time.

In sea ice at the beginning of the experiment, an average of  $5474 \pm 970$  16S rRNA gene sequences were obtained from each replicate sample (Table S1, Supporting Information), and chloroplasts (a proxy for algal sequences) represented  $72 \pm 2\%$  of the total sequences. After 2 weeks of incubation, less than 0.1% of the sequences were chloroplast for both the oiled-enriched microcosms and the controls (data not shown). In sub-ice seawater microcosms, an average of  $5964 \pm 540$  16S rRNA gene sequences were obtained (Table S1, Supporting Information). Of these,  $1.9\% \pm 0.1$  were affiliated to chloroplasts in both treatments at T0 and T15 (not shown). This is consistent with the higher algal biomass (chl  $a$ ) measured in our sea-ice samples when compared to sub-ice seawater (Table 1). Sea-ice bacterial community composition at T0 consisted of, on average,  $77 \pm 1\%$  of Proteobacteria (alpha: 0.3%; gamma: 49% and others: 28%),  $23 \pm 1\%$  of Bacteroidetes (Flavobacteria: 9%) and >1% of Actinobacteria, Acidobacteria and Firmicutes (data not shown). In sub-ice seawater, the bacterial community at T0 was composed of, on average,  $72 \pm 1\%$  of Proteobacteria (alpha: 7%; epsilon: 0.1%; gamma: 58% and others: 6%),  $28 \pm$

$1\%$  of Bacteroidetes (Flavobacteria: 17%) and >2% of Actinobacteria, Verrucomicrobia, Fusobacteria, Firmicutes and Planctomycetales (not shown).

Amplicon sequencing analyses of the 16S rRNA gene also revealed that the bacterial community composition changed significantly over the course of the experiment (i.e. between T0 and T15) in the control and oil-enriched microcosms for both sea ice and sub-ice seawater (Fig. 7). All sea-ice microcosms experienced an increase in the relative contribution of Flavobacteria/Bacteroidetes and a decrease in Proteobacteria (Fig. 7A). The reduction in Proteobacteria was less important in oil-enriched microcosms. Community composition evolved differently in the sub-ice seawater microcosms; the relative contribution of Bacteroidetes decreased and relative contribution of Proteobacteria increased, especially the Gammaproteobacteria (Fig. 7B). Notably, in the oil-enriched microcosms, the contribution of Epsilonproteobacteria increased (T15:  $4 \pm 2\%$  versus T0:  $0.1 \pm 0.0\%$ ; t-test,  $P < 0.01$ ) whereas the contribution of Alphaproteobacteria decreased significantly (T15:  $3 \pm 1\%$  versus T0:  $7 \pm 1\%$ ; t-test,  $P < 0.01$ ).

Resolution at lower taxonomic ranks revealed finer changes in sea ice (Fig. 8). Increases in the relative contribution of the genera Polaribacter and Moritella were observed at T15 in control and oil-treated microcosms, as well as a decrease in the relative contribution of Glaciecola (Fig. 8C and D). Polaribacter contributed for  $99 \pm 1\%$  of the Bacteroidetes, whereas Moritella and Glaciecola contributed  $69 \pm 7\%$  and  $1.2 \pm 0.4\%$  of the Proteobacteria, respectively. Moritella was the only genus that was more abundant in oil-treated microcosms (41%) than in the controls (24%). The most striking change was in oil-treated microcosms where the community diversity decreased because minor taxa (i.e. Arenicella, Octadecabacter, Lewinella, Ulvibacter and Perscivirga) were below detection (Fig. 8D). Sequences from the genus Algicola were only detected after incubation with crude oil (Fig. 8D).




Figure 7. Bacterial community taxonomic composition based on 16S rRNA gene sequences in sea ice (A) and sub-ice seawater (B) at the beginning (T0) and end (T15) of the incubation in control and experimental microcosms.

Interestingly, the sub-ice seawater at T0 in both the oil-treated and controls microcosms consisted of ~25% unclassified bacteria at the class level, which increased to 97% unclassified bacteria at the genus level (Fig. 9A and B). The presence of oil stimulated the growth of *Colwellia*, which constituted 33% of the population at the end of the incubation period (Fig. 9D). Three genera were only obtained after 2 weeks in the oil-treated microcosms; *Alcanivorax*, *Sulfitobacter* and *Oleispira* contributed to 0.52%, 0.20% and 0.05%, respectively (Fig. 9D). Even though new genera were detected after 2 weeks in the oil-treated microcosms, the overall diversity had decreased; the contribution of unclassified bacteria was to 61% in oil-treated microcosms at T15, whereas it was 96% in controls (Fig. 9C and D).

## DISCUSSION

### Contrasting hydrocarbon degradation within and under first-year sea ice

Ice-associated microorganisms are able to substantially degrade typical crude oil components at temperatures as low as  $-1.7^{\circ}\text{C}$ , without additional nutrients or oil dispersant. The biodegradation capacity of the indigenous ice-associated microbes of the eastern Canadian Arctic Archipelago was markedly distinct whether they lived in the ice matrix or in the underlying waters. Bacterial communities in sub-ice seawater were much more efficient in degrading hydrocarbons than those living in sea ice, as the former degraded the bulk (94%) of added hydrocarbons within 11 days (Fig. 3B). Sea ice bacteria removed less than 40% of the hydrocarbons over a similar period (Fig. 3B), although these cells were two times more abundant at the beginning of the experiment (Table 2).


Sub-ice seawater bacteria depleted almost all the resolvable alkanes ( $\text{C}_9$  to  $\text{C}_{40}$ , pristane and phytane; sensu Atlas and Bragg 2009) within 11 days at  $-1.7^{\circ}\text{C}$  (Fig. 5B). Alkanes are major components of crude oil, and are typically utilized as carbon and energy sources by bacteria (Rojo 2009), including Arctic populations growing at  $4^{\circ}\text{C}$  (Deppe et al. 2005) down to  $1^{\circ}\text{C}$  (Gerdes et al. 2005),

and as low as  $-1^{\circ}\text{C}$  (McFarlin et al. 2014). Interestingly, the methylated forms of PAHs were degraded almost as much as the alkanes by the sub-ice seawater bacteria (Fig. 4B). This contrasts with recent experimental results obtained with Arctic seawater bacteria where parent PAHs were degraded before their methylated homologs (McFarlin et al. 2014). These compounds are considerably toxic and bioaccumulate more than the non-methyl forms because the addition of a methyl group decreases water solubility (Seo, Keum and Li 2009). Methylated PAHs are nonetheless biodegradable under aerobic conditions, but few studies have addressed this topic (Seo, Keum and Li 2009). Sea-ice bacteria were not as efficient in degrading methyl-PAHs; only 2 compounds out of the 26 detected were fully metabolized (Fig. 5A), whereas the remaining ones were partially degraded (24% to 64%).

LMW PAHs are highly toxic to most aquatic organisms (Peng et al. 2008), naphthalenes being the most noxious (International Association of Oil and Gas Producers 2002), but they can be metabolized by microorganisms (van Agteren, Keuning and Oosterveld 1998). Phenanthrene, a biodegradable and relatively water-soluble PAH (Head, Jones and Röling 2006), was completely degraded by all ice-associated microbes, whereas naphthalene and dibenzothiophene were degraded to a lesser extent (Fig. 5). Nonetheless, naphthalene-degrader microbes in cold Arctic ecosystems are as efficient as their temperate counterparts (Bagi et al. 2014). Assuming the degradation rate observed during our experiment would persist, 15 more days would be needed for the complete degradation of these components. Our results indicate that sub-ice microbial communities have the physiological capabilities to degrade crude oil and could possibly biodegrade naphthalenes from an accidental spill within a month, even under seasonal sea-ice cover.

### Microcosms to study hydrocarbon degradation

The bacterial communities studied in microcosms are representative, in terms of abundance, production and diversity, of



**Figure 8.** Bacterial community taxonomic composition (in %) based on 16S rRNA gene sequences in sea ice at the beginning (T0) and end (T15) of the incubation period in control and experimental (hydrocarbons) microcosms. Note logarithm scale on y-axis.

natural communities found in the Northwest Passage in spring. In May–June, the area is typically covered with first-year ice that supports high ice algal production (Cota et al. 1991), which was reflected by the very high bottom-ice chl *a* concentrations at the time of our experiment (Table 1). Bacterial abundances in sub-ice seawater and in sea ice (Table 1) were comparable to previous records for the same region;  $1.0 \times 10^5$  cells ml<sup>-1</sup> were found at the ice-water interface, and from 1.5 up to  $100 \times 10^5$  cells ml<sup>-1</sup> in sea ice of Resolute Passage during April–May (Maranger, Bird and Juniper 1994). When applying the conservative conversion factor of  $1.5 \times 10^6$  mg C mol<sup>-1</sup> leucine (Ducklow 2003) to our leucine uptake rates at T0, daily bacterial production was  $0.77$  mg C m<sup>-3</sup> d<sup>-1</sup> in sea ice and  $0.46$  mg C m<sup>-3</sup> d<sup>-1</sup> in sub-ice seawater. These rates are consistent with rates in sea ice of  $0.48$  to  $3$  mg C m<sup>-3</sup> d<sup>-1</sup>, and

a mean of  $0.72$  mg C m<sup>-3</sup> d<sup>-1</sup> in sub-ice waters of the Resolute Passage (Maranger, Bird and Juniper 1994).

*Gammaproteobacteria* is typically the predominant group in Arctic sea ice (Junge et al. 2002; Brinkmeyer et al. 2003; Bowman et al. 2012) followed by *Bacteroidetes* (Deming 2010), whereas *Alphaproteobacteria* dominates in Arctic surface waters (Garneau et al. 2006; Alonso-Sáez et al. 2008). Accordingly, our results showed a dominance of *Gammaproteobacteria* and *Bacteroidetes* in the marine communities on the underside of landfast ice but no *Alphaproteobacteria* were detected (Fig. 7A), which is concordant with observed low abundance (ca. 4% of 16S rRNA gene sequence tags) in Arctic multiyear ice (Bowman et al. 2012). Bacterial composition in sub-ice seawater at the beginning of our experiment showed the presence of the same three dominant groups,




Figure 9. Bacterial community taxonomic composition (in %) based on 16S rRNA gene sequences in sub-ice seawater at the beginning (T0) and end (T15) of the incubation period in control and experimental (hydrocarbons) microcosms. Note logarithm scale on y-axis.

but with a predominance of the Gammaproteobacteria over the Alphaproteobacteria (Fig. 7B). This could be related to the phytoplankton pre-bloom conditions, sub-ice conditions or exchange of ice bacteria with sub-ice seawater. Given that our experiment was run in oxic conditions, as would be expected in the diffusive interfacial sea-ice layer, sequencing results did not reveal the presence of anaerobic bacteria. However, anoxic microsites have been found in sea-ice brine channels in the Arctic (Rysgaard and Glud 2004), where anaerobic bacteria were detected (Petri and Imhoff 2001; Bowman et al. 2012). Biodegradation of hydrocarbon is also possible in anoxic pockets when  $\text{NO}_3^-$ ,  $\text{Fe}^{3+}$  and  $\text{SO}_4^{2-}$  are used as electron acceptors (Widdel and Rabus 2001). The extent of anaerobic oil biodegradation in ice could not be estimated in our oil-treated microcosms, and would require further investigation.

One cannot rule out that part of naphthalene losses may be due to volatilization in the headspace of the incubation microcosms. Irwin and colleagues (1997) showed that volatiliza-

tion in soils accounted for about 20% and 30% of the loss of C<sub>1</sub>-methylnaphthalene and naphthalene, respectively, but for the 12 remaining HMW PAHs, volatilization was not an important loss mechanism. Given that at low temperature the volatilization of short-chain alkanes is reduced (Yang et al. 2009), volatilization of naphthalenes in our microcosms was probably smaller than what was observed in soils and therefore most losses were likely due to biodegradation.

The overall half-life of low-pour point oil spilled in Resolute Passage would be ca. 3 days in seawater under ice cover. This is faster than other recent experimental results, which suggest that the half-life of crude oil would be ca. 60 days in the Chukchi Sea at sub-zero temperatures (McFarlin et al. 2014). No bacterial activity was measured during that experiment, but nutrient concentrations were below the detection limit in the Chukchi Sea, suggesting that bacterial growth may have been limited and explaining the comparatively shorter half-life measured in our microcosms. In contrast, a half-life of 14 days was suggested for

crude oil in temperate ( $8^{\circ}\text{C}$ ) seawater offshore of New Jersey, USA (Prince et al. 2013).

### Microbial response to oil exposure

Initial bacterial abundances in microcosms did not have much impact on the overall degradation of oil because even though cell abundance in sub-ice seawater microcosms was half the count in ice (Table 2), sub-ice microbes degraded three times more hydrocarbons in 3 days (Fig. 3). Results rather support the idea that difference in the starting community composition and/or environmental conditions (e.g. DOC concentrations) were key factors. The relative contribution of Gammaproteobacteria to the total bacteria was higher in sub-ice seawater than in the ice matrix, where the contribution of other Proteobacteria was also important (Fig. 7). Gammaproteobacteria were confirmed to be the main players of oil biodegradation in cold marine environments (Yakimov et al. 2004; Gerdes et al. 2005; Brakstad and Bonaunet 2006; Brakstad et al. 2008). Bacterial diversity also changed in the presence of hydrocarbons (Figs 7–9) and shifted toward a higher contribution of Gammaproteobacteria in both sub-ice seawater and sea ice, further supporting the contention that oil-degrading bacteria are affiliated to this class. Interestingly, the Epsilonproteobacteria in sub-ice seawater also seemed to have benefited from the presence of oil, as their contribution increased in oil-enriched microcosms but not in controls (Fig. 7B). Several genera of this class that are able to oxidize sulfur compounds dominated the microbial biomass in groundwater contaminated by crude-oil extraction activities (Kodama and Watanabe 2003; Hubert et al. 2011). This may explain why dibenzothiophene, a sulfur-containing compound, and its methylated forms were highly oxidized (average of 97% loss) in sub-ice waters but not in sea ice (< 40% loss; Fig. 5).

Bacteroidetes increased the most after 15 days of incubation in sea-ice microcosms, regardless of the presence of hydrocarbons (Fig. 7A) and following bacterial production rates (Fig. 2), i.e. Bacteroidetes increased in sea ice, but decreased in seawater. These changes agree with previous observations of a Bacteroidetes predominance associated with high bacterial biomass and production (Teira et al. 2008). Bacteroidetes are the main degraders of complex polymeric organic compounds (Cottrell and Kirchman 2000; Fernández-Gómez et al. 2013); therefore, the presence of extracellular polymeric substances (EPS) in the sea ice of the Canadian Arctic Archipelago (Aslam et al. 2016) could have provided a competitive advantage for resources to the Bacteroidetes over the Gammaproteobacteria. Conversely, the low organic matter content in sub-ice seawater (Table 1) could have been more advantageous for the Gammaproteobacteria.

The characterization at the genus level indicated that crude-oil addition reduced microbial diversity (Figs 8 and 9), as previously shown for oil-polluted fjord ice (Brakstad et al. 2008) and Arctic multiyear ice (Gerdes et al. 2005). Three genera benefited from the oil enrichment in sea ice, the Gammaproteobacteria genera *Moritella*, *Algicola* and *Colwellia* (Fig. 8). *Moritella* was recently reported to be the second most abundant genus in naphthalene-spiked Arctic seawater (Bagi et al. 2014), whereas the genus *Algicola* has not been previously described as oil degrader. It is unclear if *Algicola* strains metabolized the added hydrocarbons or if they were rather secondary consumers that profited from other compounds released by oil-degrading bacteria.

Members of the genus *Colwellia*, which are oil-degrading Gammaproteobacteria (Yakimov et al. 2004; Redmond and Valentine 2012), were also stimulated by the presence of oil in sub-

ice seawater (Fig. 9). Like *Polaribacter* (Bacteroidetes), *Colwellia* are known psychrophiles and may have benefited from the low incubation temperature ( $< -1^{\circ}\text{C}$ ), as previously observed in cold marine environments (Yakimov et al. 2004; Brakstad et al. 2008). Three typical marine hydrocarbonoclastic bacteria, i.e. *Alcanivorax* and *Oleispira* (Yakimov et al. 2007), only grew in oil-treated sub-ice seawater microcosms, as well as *Sulfitobacter*, a sulfite-oxidizing Alphaproteobacteria bacterium. This genera was also obtained in beach sands in the Gulf of Mexico after the Deepwater Horizon oil spill (Kostka et al. 2011). The very high percentage of unclassified bacteria in the sub-ice seawater (97%) at the genus level indicates that the majority of bacteria in cold Arctic waters have yet to be discovered.

The oil treatment in sea-ice microcosms did not induce an increase in bacterial production as it did in sub-ice microcosms after 3 days and until the end of incubation (Fig. 2). This lag period was similar to the 5 to 8 day lag observed during other carbon enrichment experiments on polar bacterial communities (Dyda et al. 2009; Ducklow et al. 2011). Bacterial communities usually need time to activate the pathway of enzymes required for hydrocarbon degradation (Rojo 2009). The fast response to hydrocarbon exposure observed in our study may be due to a sub-population of alkane degraders that is maintained by alkanes, which are ubiquitous in the marine environment, being continuously produced by many organisms (plants, green algae, bacteria and animals; Rojo 2009). The sub-ice carbon:nitrogen (DOC:DN, mol:mol) ratio of 6:1 (Table 1) suggests a carbon-limited community because it is below the ratio of 10:1 for optimal hydrocarbon degradation (Alexander 1999). The addition of hydrocarbons very likely stimulated the activity of the carbon-limited sub-ice bacteria. Likewise, the addition of  $^{3}\text{H}$ -leucine may have induced an artificial stimulation of the uptake rates, explaining why bacterial production in control microcosms at T0 were much higher than at any other time point (Fig. 2). Toxic stress on bacteria after oil addition (e.g. Siron et al. 1993) may have impeded such stimulation in the oil-treated microcosms.

In sea ice, the DOC:DN ratio of 16:1 (Table 1) was above the optimal ratio, indicating that the biodegradation of spilled hydrocarbons would rather be limited by nutrients. Accordingly, the addition of hydrocarbons did not lead to a sharp increase in bacterial production (Fig. 2A). Nutrient limitation after oil spills is common in marine environments because hydrocarbons have high organic carbon content and no bioavailable nitrogen (Das and Chandran 2011). The limited biodegradation in sea ice may also be explained by the availability of highly labile dissolved organic matter produced by sea-ice algae (Amon, Fitznar and Benner 2001). Algal-derived carbon supports substantial bacterial growth (Riedel et al. 2008), and most alkane-degrading bacteria were shown to prefer such labile compounds than alkanes when available (Rojo 2009). The very high DOC and chl a concentrations found in sea ice as well as the apparent absence of (or a short) lag period in sea-ice bacterial activity would agree with a bacterial production supported by sea-ice organic carbon.

### Factors governing hydrocarbon degradation in ice-covered Arctic waters

Among the 59 crude oils classified in ITOPF's technical paper (2011), about 61% have a pour point  $< -2^{\circ}\text{C}$  (including the Arabian Light used in this study), implying that a majority of oils would still behave like fluids and spread in droplets under wave action even if the Arctic seawater is at its coldest. Arabian Light oil was

efficiently degraded in this study in part because the oil was still fluid and accessible to microorganisms, even at sub-zero temperatures. High degradation rates could be found in the field for several other low pour-point oils, although in some cases oil composition could lead to divergent results. In ice-covered areas, however, where temperature in first-year ice varies between  $-2^{\circ}\text{C}$  and  $-20^{\circ}\text{C}$  (Junge, Eicken and Deming 2004), degradation rates would potentially be much lower than what was observed in ice microcosms because only 39% of the ITOFP-classified oils would exhibit liquid characteristics below  $-15^{\circ}\text{C}$ . At temperature as low as  $-25^{\circ}\text{C}$  in multiyear ice (Perovich and Elder 2001), biodegradation rates would be negligible, if not null, because most oils will solidify (ITOPF 2011) and because very cold ice has little brine volume fraction and thus, fewer microhabitats to support an active bacterial community (Deming 2010).

The 'law of five' states that sea ice that has a brine volume fraction of 5% and the typical salinity of 5 ppt will be impermeable at temperatures below  $-5^{\circ}\text{C}$  (Golden, Ackley and Lytle 1998). Conversely, at temperatures warmer than  $-5^{\circ}\text{C}$ , the sea ice will be permeable because brine inclusions become connected, allowing heat, brine, nutrients and seawater to move through ice (Golden, Ackley and Lytle 1998). Thus, whether an oil spill occurs in winter or during the warmer ice melting season has important implications for the biodegradation of hydrocarbons. In winter, chances are that sea ice would be impermeable because the temperature can drop below  $-5^{\circ}\text{C}$  (Junge, Eicken and Deming 2004). This means that oil spilled on the ice cover would not percolate and be in contact with bottom-ice bacterial communities, thereby limiting biodegradation. However, if some oil was floating at the ice-water interface, it could reach the lowermost part of the ice where the temperature remains near to the seawater freezing point of  $-1.8^{\circ}\text{C}$  (Notz and Worster 2009).

Field observations indicate that oil spilled in calm seas spreads out under the ice where it gradually fills depressions (Fingas and Hollebone 2003). Our results imply that bacteria in these cavities and growing close to the ice-water interface should be capable of biodegradation during winter. Bacteria were shown to be active in the upper water column even in winter, but rates were low at  $< 0.05 \text{ mg C m}^{-3} \text{ d}^{-1}$  (Garneau et al. 2008). Assuming the oil reaches the ice-water interface, biodegradation rates could be higher than the ones measured in the microcosms because bottom-ice DOC and EPS concentrations are very low in winter (Riedel et al. 2008), as well as the primary producer biomass ( $< 0.04 \text{ mg chl a m}^{-3}$ ; Garneau et al. 2008). In such carbon-limiting conditions, oil additions could enhance bacterial production.

It is unclear if under-ice biodegradation during cold seasons could be significant. Limited oil biodegradation was found under a winter ice cover in the St. Lawrence River, a temperate marine system, and was attributed to the low productivity of the microbial communities (Siron et al. 1993). Another study in the Beaufort Sea investigated the movement of oil under first-year ice and showed that oil spilled in October was encapsulated in ice within days, and remained in the bottom 5–10 cm of the first-year ice until February, without apparent degradation (NORCOR Engineering Research Limited 1975). Recent results show that full-depth connectivity between the top and bottom of first-year ice can happen in spring, well before summer melt (Jardon et al. 2013), potentially favoring bacterial degradation in the event of oil release during spring. Therefore, remediation strategies for oil mitigation in the Arctic likely need to vary according to seasonal conditions. Finally, the biodegradation potential of microbial communities in melt ponds formed at the ice-cover surface remains to be determined.

## CONCLUDING REMARKS

The experimental results demonstrate great potential for bioremediation techniques in the Arctic Ocean, whether in sea ice or in the water beneath it. Nutrient concentrations should not be a limiting factor if a small amount of contamination occurs during the pre-bloom period, considering that 94% of the initial hydrocarbons were lost within 2 weeks in our oil-treated microcosms. So far, biostimulation (i.e. supply of limiting resources; Brakstad 2008) has been more successful than bioaugmentation in cold environments because the added microorganisms were stressed by the polar conditions and were outcompeted by indigenous species (Yang et al. 2009). Biostimulation in winter or early spring, when nutrient concentrations in surface water are elevated, will likely have little effect. However, such a technique could be beneficial if used after the phytoplankton bloom period, when the surface water is depleted of nutrients. Likewise, biostimulation could be efficiently used in case of an accidental release of large amounts of hydrocarbons, which would result in rapid depletion of the available nutrients.

Our results highlight the influence of the natural, ice-algal carbon resources on the activity of oil-degrading bacteria. Hydrocarbon degradation by sub-ice planktonic bacteria under high DOC concentration (i.e. during the phytoplankton bloom), as well as by sea-ice bacteria under low DOC concentration (before the ice-algal bloom), could provide a better understanding of bioremediation during the spring-summer transition. Further studies could also target the oil-degradation functional genes of ice-associated bacteria, as was previously reported for Arctic soils (Yergeau et al. 2012). All this information will provide a finer comprehension of natural oil degradation and help in developing bioremediation strategies for the Arctic.

## SUPPLEMENTARY DATA

Supplementary data are available at FEMSEC online.

## ACKNOWLEDGEMENTS

The authors sincerely thank Anke Reppchen, Michel Poulin, Matthew Dunbar and Julie Champagne for their assistance in the field and laboratory, Gary Wohlgeschaffen, and three anonymous referees for constructive comments on the manuscript. The authors are also grateful to the Hunters and Trappers Association of Resolute Bay for their continued support, and appreciated the excellent logistical support of the Polar Continental Shelf Program in Resolute, Nunavut.

## FUNDING

This research was supported by: the Program of Energy Research and Development (Portfolio Frontier Oil and Gas, Program 1.2.4); Natural Sciences and Research Council of Canada (Discovery Grant no. 327416 to Christine Michel); and the Natural Resources Canada Polar Continental Shelf Program (Project 107–12 to Christine Michel).

**Conflict of interest.** None declared.

## REFERENCES

- Alexander M. *Biodegradation and Bioremediation*. Ithaca: Gulf Professional Publishing, 1999.

- Alonso-Sáez L, Sánchez O, Gasol JM et al. Winter-to-summer changes in the composition and single-cell activity of near-surface Arctic prokaryotes. *Environ Microbiol* 2008;19:2444–54.
- AMAP. Assessment 2007: Oil and Gas Activities in the Arctic - Effects and Potential Effects, Vol 1. Oslo: Arctic Monitoring and Assessment Programme (AMAP), 2010.
- AMAP. Arctic Climate Issues 2011: Changes in Arctic Snow, Water, Ice and Permafrost. Oslo: Arctic Monitoring and Assessment Programme (AMAP), 2012.
- Amon RMW, Fitznar H-P, Benner R. Linkages among the bioreactivity, chemical composition, and diagenetic state of marine dissolved organic matter. *Limnol Oceanogr* 2001;46:287–97.
- Arrigo KR, Mock T, Lizotte MP. Primary producers and sea ice. In: Thomas DN, Dieckmann GS (eds). *Sea Ice*, 2nd edn. Oxford: Blackwell Publishing, 2010, 283–325.
- Aslam SN, Michel C, Niemi A et al. Patterns and drivers of carbohydrate budgets in ice algal assemblages from first year Arctic sea ice. *Limnol Oceanogr* 2016;61:919–37.
- Atlas RM. Microbial degradation of petroleum hydrocarbons: an environmental perspective. *Microbiol Rev* 1981;45:180–209.
- Atlas RM, Bragg J. Bioremediation of marine oil spills: when and when not—the Exxon Valdez experience. *Microb Biotechnol* 2009;2:213–21.
- Atlas RM, Cerniglia CE. Bioremediation of petroleum pollutants. *BioScience* 1995;45:332–8.
- Atlas RM, Horowitz A, Busdosh M. Prudhoe crude oil in Arctic marine ice, water, and sediment ecosystems: degradation and interactions with microbial and benthic communities. *J Fish Res Board Can* 1978;35:585–90.
- Ausubel FM, Brent R, Kingston RE et al. (eds). *Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology*. New York: John Wiley & Sons, 2002.
- Bagi A, Pampanin DM, Lanzén A et al. Naphthalene biodegradation in temperate and arctic marine microcosms. *Biodegradation* 2014;25:111–25.
- Bell TH, Yergeau É, Maynard C et al. Predictable bacterial composition and hydrocarbon degradation in Arctic soils following diesel and nutrient disturbance. *ISME J* 2013;7:1200–10.
- Belzile C, Brugel S, Nozaïs C et al. Variations of the abundance and nucleic acid content of heterotrophic bacteria in Beaufort Shelf waters during winter and spring. *J Mar Syst* 2008;74:946–56.
- Bowman JS, Rasmussen S, Blom N et al. Microbial community structure of Arctic multiyear sea ice and surface seawater by 454 sequencing of the 16S RNA gene. *ISME J* 2012;6:11–20.
- Brakstad OG. Natural and stimulated biodegradation of petroleum in cold marine environments. In: Margesin R, Schinner F, Marx J-C et al. (eds). *Psychrophiles: From Biodiversity to Biotechnology*. Berlin Heidelberg: Springer, 2008, 389–407.
- Brakstad OG, Bonaunet K. Biodegradation of petroleum hydrocarbons in seawater at low temperatures (0–5°C) and bacterial communities associated with degradation. *Biodegradation* 2006;17:71–82.
- Brakstad OG, Nonstad I, Faksness L-G et al. Responses of microbial communities in Arctic sea ice after contamination by crude petroleum oil. *Microb Ecol* 2008;55:540–52.
- Brinkmeyer R, Knittel K, Jurgens J et al. Diversity and structure of bacterial communities in Arctic versus Antarctic pack ice. *Appl Environ Microb* 2003;69:6610–9.
- Canadian Council of Ministers of the Environment. Canadian sediment quality guidelines for the protection of aquatic life: Polycyclic aromatic hydrocarbons (PAHs). In: *Canadian Environmental Quality Guidelines*. Winnipeg: Canadian Council of Ministers of the Environment, 1999.
- Clarke KR, Gorley RN. PRIMER V5: User Manual/Tutorial. Plymouth UK: PRIMER-E Limited, 2001.
- Clarke KR, Warwick RM. *Change in Marine Communities: An Approach to Statistical Analysis and Interpretation*. Plymouth: PRIMER-E Limited, 2001.
- Codispoti LA, Kelly V, Thessen A et al. Synthesis of primary production in the Arctic Ocean: III. Nitrate and phosphate based estimates of net community production. *Prog Oceanogr* 2013;110:126–50.
- Cota GF, Legendre L, Gosselin M et al. Ecology of bottom ice algae: I. Environmental controls and variability. *J Mar Syst* 1991;2:257–77.
- Cottrell MT, Kirchman DL. Natural assemblages of marine proteobacteria and members of the Cytophaga-Flavobacter cluster consuming low- and high-molecular-weight dissolved organic matter. *Appl Environ Microb* 2000;66:1692–7.
- Das N, Chandran P. Microbial degradation of petroleum hydrocarbon contaminants: an overview. *Biotechnol Res Int* 2011;2011:1–13.
- Deming JW. Sea ice bacteria and viruses. In: Thomas DN, Dieckmann GS (eds). *Sea Ice*, 2nd edn. Oxford: Blackwell Publishing, 2010, 247–82.
- Deppe U, Richnow H-H, Michaelis W et al. Degradation of crude oil by an arctic microbial consortium. *Extremophiles* 2005;9:461–70.
- Ducklow HW. Seasonal production and bacterial utilization of DOC in the Ross Sea, Antarctica. In: DiTullio GR, Dunbar RB (eds). *Biogeochemistry of the Ross Sea*, Vol 78. Washington, DC: American Geophysical Union, 2003, 143–57.
- Ducklow HW, Myers K, Erickson M et al. Response of a summertime Antarctic marine bacterial community to glucose and ammonium enrichment. *Aquat Microb Ecol* 2011;64:205–20.
- Dyda RY, Suzuki MT, Yoshinaga MY et al. The response of microbial communities to diverse organic matter sources in the Arctic Ocean. *Deep-Sea Res II* 2009;56:1249–63.
- Environment Canada. Arabian Light. Emergencies Science and Technology Division, 1991.
- Environment Canada. Oil Properties. 2001, <http://www/etc-cte.ec.gc.ca/databases/oilproperties/> (11 July 2016, date last accessed).
- Fernández-Gómez B, Richter M, Schüler M et al. Ecology of marine Bacteroidetes: a comparative genomics approach. *ISME J* 2013;7:1026–37.
- Fingas MF, Hollebone BP. Review of behaviour of oil in freezing environments. *Mar Pollut Bull* 2003;47:333–40.
- Fisher MM, Triplett EW. Automated approach for ribosomal intergenic spacer analysis of microbial diversity and its application to freshwater bacterial communities. *Appl Environ Microb* 1999;65:4630–6.
- Garnier M-È, Roy S, Lovejoy C et al. Seasonal dynamics of bacterial biomass and production in a coastal arctic ecosystem: Franklin Bay, western Canadian Arctic. *J Geophys Res-Oceans* 2008;113:C07S91.
- Garnier M-È, Vincent WF, Alonso-Sáez L et al. Prokaryotic community structure and heterotrophic production in a river-influenced coastal arctic ecosystem. *Aquat Microb Ecol* 2006;42:27–40.
- Gerdes B, Brinkmeyer R, Dieckmann GS et al. Influence of crude oil on changes of bacterial communities in Arctic sea-ice. *FEMS Microbiol Ecol* 2005;53:129–39.
- Giraudoux P. Pgirmess: Data Analysis in Ecology. R package version 1.5.9. 2014, <http://cran.r-project.org/web/packages/pgirmess/index.html> (11 July 2016, date last accessed).

- Golden KM, Ackley SF, Lytle VI. The percolation phase transition in sea ice. *Science* 1998;282:2238–41.
- Hansen HP, Koroleff F. Determination of nutrients. In: Grasshoff K, Kremling K, Ehrhardt M (eds). *Methods of Seawater Analysis*. Weinheim: Wiley-VCH, 1999, 159–228.
- Hassanshahian M, Cappello S. Crude oil biodegradation in the marine environments. In: Chamy R, Rosenkranz F (eds). *Biodegradation Engineering and Technology*. Rijeka, Croatia: In-TechOpen, 2013, 101–35.
- Head IM, Jones DM, Röling WFM. Marine microorganisms make a meal of oil. *Nat Rev Microbiol* 2006;4:173–82.
- Herndl GJ, Reinhäler T, Teira E et al. Contribution of Archaea to total prokaryotic production in the deep Atlantic Ocean. *Appl Environ Microb* 2005;71:2303–9.
- Hubert CRJ, Oldenburg TBP, Fustic M et al. Massive dominance of *Epsilonproteobacteria* in formation waters from a Canadian oil sands reservoir containing severely biodegraded oil. *Environ Microb* 2011;14:387–404.
- International Association of Oil & Gas Producers. *Aromatics in Produced Water: Occurrence, Fate and Effects, and Treatment*. London: International Association of Oil & Gas Producers, 2002.
- Irwin RJ, Van Mouwerik M, Stevens L et al. *Environmental Contaminants Encyclopedia C1-Naphthalene Entry*. Fort Collins, Colorado: National Park Service, 1997:1–50.
- ITOPF. *Fate of Marine Oil Spills. Technical Information Paper*. London: The International Tanker Owners Pollution Federation Limited (ITOPF), 2011.
- Jardon FP, Vivier F, Vancoppenolle M et al. Full-depth desalination of warm sea ice. *J Geophys Res-Oceans* 2013;118:435–47.
- Junge K, Eicken H, Deming JW. Bacterial activity at –2 to –20°C in Arctic wintertime sea ice. *Appl Environ Microb* 2004;70:550–7.
- Junge K, Imhoff F, Staley T et al. Phylogenetic diversity of numerically important arctic sea-ice bacteria cultured at subzero temperature. *Microb Ecol* 2002;43:315–28.
- Karlsson J, Petrich C, Eicken H. Oil entrainment and migration in laboratory-grown saltwater ice. *Proceedings of the 21st International Conference on Port and Ocean Engineering under Arctic Conditions*. Boca Raton, Florida, 2011.
- Kirchman DL, Ducklow HW. Estimating conversion factors for the thymidine and leucine methods for measuring bacterial production. In: Kemp PF, Sherr BF, Sherr EB et al. (eds). *Handbook of Methods in Aquatic Microbial Ecology*. Boca Raton: Lewis Publications, 1993, 513–7.
- Knap A, Michaels A, Close A et al. Protocols for the Joint Global Ocean Flux Study (JGOFS) Core Measurements. JGOFS Report No 29. Paris: UNESCO, 1996.
- Kodama Y, Watanabe K. Isolation and characterization of a sulfur-oxidizing chemolithotroph growing on crude oil under anaerobic conditions. *Appl Environ Microb* 2003;69:107–12.
- Kostka JE, Prakash O, Overholst WA et al. Hydrocarbon-degrading bacteria and the bacterial community response in Gulf of Mexico beach sands impacted by the Deepwater Horizon oil spill. *Appl Environ Microb* 2011;77:7962–74.
- Kwok R, Cunningham GF. Variability of Arctic sea ice thickness and volume from CryoSat-2. *Philos T Roy Soc A* 2015;373:20140157.
- Leahy JG, Colwell RR. Microbial degradation of hydrocarbons in the environment. *Microbiol Rev* 1990;54:305–15.
- McFarlin KM, Prince RC, Perkins R et al. Biodegradation of dispersed oil in Arctic seawater at –1°C. *PLoS One* 2014;9:e84297.
- McKetta JJ, Jr. *Encyclopedia of Chemical Processing and Design*. New York: CRC Press, 1997.
- McLaughlin FA, Carmack EC, Ingram RG et al. Oceanography of the Northwest Passage. In: Robinson AR, Brink KH (eds). *The Sea, Vol. 14: the Global Coastal ocean, Interdisciplinary Regional Studies and Syntheses*. New York: Wiley, 2004, 1211–42.
- Maranger R, Bird DF, Juniper SK. Viral and bacterial dynamics in Arctic sea ice during the spring algal bloom near Resolute, N.W.T., Canada. *Mar Ecol Prog Ser* 1994;111:121–7.
- Meier WN, Hovelsrud GK, Oort BEH et al. Arctic sea ice in transformation: a review of recent observed changes and impacts on biology and human activity. *Rev Geophys* 2014;52:185–217.
- Meltofte H, Barry T, Berteaux D et al. Synthesis: implications for conservation. In: Meltofte H (ed.). *Arctic Biodiversity Assessment. Status and Trends in Arctic Biodiversity*. Oslo: Conservation of Arctic Flora and Fauna (CAFF), 2013, 21–65.
- Michel C. Marine ecosystems. In: Meltofte H (ed.). *Arctic Biodiversity Assessment. Status and Trends in Arctic Biodiversity*. Oslo: Conservation of Arctic Flora and Fauna, 2013, 379–419.
- Michel C, Ingram RG, Harris LR. Variability in oceanographic and ecological processes in the Canadian Arctic Archipelago. *Prog Oceanogr* 2006;71:379–401.
- Michel C, Legendre L, Ingram RG et al. Carbon budget of sea-ice algae in spring: evidence of a significant transfer to zooplankton grazers. *J Geophys Res-Oceans* 1996;101:18.345–60.
- Müller TJ. Determination of salinity. In: Grasshoff K, Kremling K, Ehrhardt M (eds). *Methods of Seawater Analysis*. 3rd edn. Weinheim: Wiley-VCH, 1999, 41–73.
- NORCOR Engineering Research Limited. *The Interaction of Crude Oil with Arctic Sea Ice*. Victoria: Beaufort Sea Project, Department of the Environment, 1975:1–213.
- Normand P, Ponsonnet C, Nesme X et al. ITS analysis of prokaryotes. In: Akkermans DL, van Elsas JD, de Brujin FJ (eds). *Molecular Microbial Ecology Manual*, Vol 3. Amsterdam: Kluwer Academic Publishers, 1996, 1–12.
- Notz D, Worster MG. Desalination processes of sea ice revisited. *J Geophys Res-Oceans* 2009;114:C05006.
- Osborne CA, Rees GN, Bernstein Y et al. New threshold and confidence estimates for terminal restriction fragment length polymorphism analysis of complex bacterial communities. *Appl Environ Microb* 2006;72:1270–8.
- Parsons TR, Maita Y, Lalli CM. *A Manual of Chemical and Biological Methods for Seawater Analysis*. Oxford: Pergamon Press, 1984.
- Peng RH, Xiong AS, Xue Y et al. Microbial biodegradation of polycyclic aromatic hydrocarbons. *FEMS Microbiol Rev* 2008;32:927–55.
- Perovich DK, Elder BC. Temporal evolution of Arctic sea-ice temperature. *Ann Glaciol* 2001;33:207–11.
- Petri R, Imhoff JF. Genetic analysis of sea-ice bacterial communities of the Western Baltic Sea using an improved double gradient method. *Polar Biol* 2001;24:252–7.
- Petrich C, Eicken H. Growth, structure and properties of sea ice. In: Thomas DN, Dieckmann GS (eds). *Sea Ice*, 2nd edn. Oxford: Wiley-Blackwell, 2009, 23–77.
- Prince RC, Elmendorf DL, Lute JR et al.  $17\alpha(H),21\beta(H)$ -Hopane as a conserved internal marker for estimating the biodegradation of crude oil. *Environ Sci Technol* 1994;28:142–5.
- Prince RC, McFarlin KM, Butler JD et al. The primary biodegradation of dispersed crude oil in the sea. *Chemosphere* 2013;90:521–6.
- Ramette A. Quantitative community fingerprinting methods for estimating the abundance of operational taxonomic units in natural microbial communities. *Appl Environ Microb* 2009;75:2495–505.
- Ranjard L, Poly F, Combrisson J et al. Heterogeneous cell density and genetic structure of bacterial pools associated with various soil microenvironments as determined by

- enumeration and DNA fingerprinting approach (RISA). *Microb Ecol* 2000;49:263–72.
- R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing. 2013, <http://www.R-project.org/> (11 July 2016, date last accessed).
- Redmond MC, Valentine DL. Natural gas and temperature structured a microbial community response to the Deepwater Horizon oil spill. *P Natl Acad Sci USA* 2012;109:20292–7.
- Riedel A, Michel C, Gosselin M et al. Winter-spring dynamics in sea-ice carbon cycling in the coastal Arctic Ocean. *J Mar Syst* 2008;74:918–32.
- Rojo F. Degradation of alkanes by bacteria. *Environ Microbiol* 2009;11:2477–90.
- Rysgaard S, Glud RN. Anaerobic N<sub>2</sub> production in Arctic sea ice. *Limnol Oceanogr* 2004;49:86–94.
- Seo J-S, Keum Y-S, Li QX. Bacterial degradation of aromatic compounds. *Int J Environ Res Public Health* 2009;6:278–309.
- Siron R, Pelletier E, Delille D et al. Fate and effects of dispersed crude oil under icy conditions simulated in mesocosms. *Mar Environ Res* 1993;35:273–302.
- Smith DC, Azam F. A simple, economical method for measuring bacterial protein synthesis rates in seawater using <sup>3</sup>H-leucine. *Mar Microb Food Webs* 1992;6:107–14.
- Smith REH, Clement P, Cota GF. Population dynamics of bacteria in Arctic sea ice. *Microb Ecol* 1989;17:63–76.
- Teira E, Gasol JM, Aranguren-Gassis M et al. Linkages between bacterioplankton community composition, heterotrophic carbon cycling and environmental conditions in a highly dynamic coastal ecosystem. *Environ Microbiol* 2008;10:906–17.
- van Agteren MH, Keuning S, Oosterhaven J. Handbook on Biodegradation and Biological Treatment of Hazardous Organic Compounds. Dordrecht, the Netherlands: Springer, 1998.
- Wang Q, Garrity GM, Tiedje JM et al. Naïve Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microb* 2007;73:5261–7.
- Wells LE, Deming JW. Abundance of Bacteria, the Cytophaga-Flavobacterium cluster and Archaea in cold oligotrophic waters and nepheloid layers of the Northwest Passage, Canadian Archipelago. *Aquat Microb Ecol* 2003;31:19–31.
- Whyte LG, Bourbonnière L, Bellerose C et al. Bioremediation assessment of hydrocarbon-contaminated soils from the High Arctic. *Bioremed J* 1999;3:69–80.
- Widdel F, Rabus R. Anaerobic biodegradation of saturated and aromatic hydrocarbons. *Curr Opin Biotech* 2001;12:259–76.
- Yakimov MM, Gentile G, Bruni V et al. Crude oil-induced structural shift of coastal bacterial communities of rod bay (Terra Nova Bay, Ross Sea, Antarctica) and characterization of cultured cold-adapted hydrocarbonoclastic bacteria. *FEMS Microbiol Ecol* 2004;49:419–32.
- Yakimov MM, Timmis KN, Golyshin PN et al. Obligate oil-degrading marine bacteria. *Curr Opin Biotechnol* 2007;18:257–66.
- Yang S-Z, Jin H-J, Wei Z et al. Bioremediation of oil spills in cold environments: a review. *Pedosphere* 2009;19:371–81.
- Yergeau É, Sanschagrin S, Beaumier D et al. Metagenomic analysis of the bioremediation of diesel-contaminated Canadian High Arctic soils. *PLoS One* 2012;7:e30058.
- Yergeau É, Sanschagrin S, Maynard C et al. Microbial expression profiles in the rhizosphere of willows depend on soil contamination. *ISME J* 2014;8:344–58.