

Introduction to Probability, Statistics and Random Processes

Chapter 6: Multiple Random Variables

Hossein Pishro-Nik
University of Massachusetts, Amherst
Email: pishro@ecs.umass.edu

Methods for More Than Two Random Variables

Joint Distributions
and Independence
Sums of Random
Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Joint Distributions and Independence

- ▶ The joint PMF of n discrete random variables X_1, X_2, \dots, X_n is

$$P_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n).$$

- ▶ The joint PDF of n continuous random variables is a function $f_{X_1 X_2 \dots X_n}(x_1, x_2, \dots, x_n)$ such that probability of $A \subset \mathbb{R}^n$ is given by

$$P\left((X_1, X_2, \dots, X_n) \in A\right) = \int_A f_{X_1 X_2 \dots X_n}(x_1, x_2, \dots, x_n) dx_1 dx_2 \dots dx_n.$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

- ▶ Marginal PDF of X_1 can be obtained from the joint PDF by

$$f_{X_1}(x_1) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f_{X_1 X_2 \dots X_n}(x_1, x_2, \dots, x_n) dx_2 \cdots dx_n.$$

- ▶ Joint CDF of n random variables X_1, X_2, \dots, X_n is given by

$$\begin{aligned} F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) \\ = P(X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n). \end{aligned}$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Independence

- Random variables X_1, X_2, \dots, X_n are independent, if for all $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$,

$$F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = F_{X_1}(x_1)F_{X_2}(x_2) \cdots F_{X_n}(x_n).$$

- If X_1, X_2, \dots, X_n are discrete, then they are independent if for all $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, we have

$$P_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = P_{X_1}(x_1)P_{X_2}(x_2) \cdots P_{X_n}(x_n).$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

- ▶ If X_1, X_2, \dots, X_n are continuous, then they are independent if for all $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, we have

$$f_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = f_{X_1}(x_1)f_{X_2}(x_2) \cdots f_{X_n}(x_n).$$

- ▶ If random variables X_1, X_2, \dots, X_n are independent, then we have

$$E[X_1 X_2 \cdots X_n] = E[X_1]E[X_2] \cdots E[X_n].$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Random variables X_1, X_2, \dots, X_n are said to be **independent and identically distributed (i.i.d.)** if they are *independent*, and they have the *same marginal distributions*:

$$F_{X_1}(x) = F_{X_2}(x) = \dots = F_{X_n}(x), \text{ for all } x \in \mathbb{R}.$$

Methods for More Than Two Random Variables

Joint Distributions and Independence

Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Sums of Random Variables

In general

$$Var\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n Var(X_i) + 2 \sum_{i < j} Cov(X_i, X_j)$$

- ▶ The MGF is very useful because of the following reasons:
 - ▶ The MGF of X gives us all the moments of X .
 - ▶ The MGF, if it exists uniquely determines the distribution.
- ▶ With the MGF of a random variable, we have determined it's distribution. This method is very useful when we work on sums of several independent random variables.

Methods for More
Than Two
Random Variables

Joint Distributions
and Independence
Sums of Random
Variables

Moment Generating
Functions

Characteristic
Functions

Random Vectors

Probability Bounds

The Union Bounds
and its Extensions

Markov and
Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz
Inequality

Jensen's Inequality

Finding Moments from MGF

- Using the Taylor series for e^x , we can write

$$e^{sX} = \sum_{k=0}^{\infty} \frac{(sX)^k}{k!} = \sum_{k=0}^{\infty} \frac{X^k s^k}{k!}.$$

We can obtain all moments of X^k from its MGF:

$$M_X(s) = \sum_{k=0}^{\infty} E[X^k] \frac{s^k}{k!}$$

$$E[X^k] = \left. \frac{d^k}{ds^k} M_X(s) \right|_{s=0}$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

- **Theorem:** Consider two random variables X and Y . Suppose that there exists a positive constant c such that MGFs of X and Y are finite and identical for all values of s in $[-c, c]$. Then,

$$F_X(t) = F_Y(t), \text{ for all } t \in \mathbb{R}.$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Sum on Independent Random Variables

If X_1, X_2, \dots, X_n are n independent random variables, then

$$M_{X_1+X_2+\dots+X_n}(s) = M_{X_1}(s)M_{X_2}(s)\cdots M_{X_n}(s).$$

Methods for More Than Two Random Variables

Joint Distributions and Independence

Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Characteristic Functions

- ▶ The MGF does not exist for all random variables.
- ▶ If a random variable does have a well-defined MGF, we can use the characteristic function defined as

$$\phi_X(\omega) = E[e^{j\omega X}]$$

- ▶ $j = \sqrt{-1}$ and ω is a real number.
- ▶ The characteristic function is defined for all real-valued random variables.
- ▶ $|\phi_X(\omega)| \leq 1$.
- ▶ It has similar properties to the MGF. If X_1, X_2, \dots, X_n are n independent random variables, then

$$\phi_{X_1+X_2+\dots+X_n}(\omega) = \phi_{X_1}(\omega)\phi_{X_2}(\omega)\cdots\phi_{X_n}(\omega).$$

Methods for More
Than Two
Random Variables

Joint Distributions
and Independence
Sums of Random
Variables

Moment Generating
Functions

Characteristic
Functions

Random Vectors

Probability Bounds

The Union Bounds
and its Extensions

Markov and
Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz
Inequality

Jensen's Inequality

Random Vectors

- ▶ When we have n random variables X_1, X_2, \dots, X_n we can put them in a (column) vector \mathbf{X} :

$$\mathbf{X} = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix}.$$

- ▶ \mathbf{X} is a n -dimensional random vector.
- ▶ The CDF of the random vector \mathbf{X} is

$$\begin{aligned} F_{\mathbf{X}}(\mathbf{x}) &= F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) \\ &= P(X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n). \end{aligned}$$

- ▶ If the X_i 's are jointly continuous, the PDF of \mathbf{X} can be written as

$$f_{\mathbf{X}}(\mathbf{x}) = f_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n).$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Expectation

- ▶ **Expected value vector** or **mean vector** of random vector \mathbf{X} is defined as

$$E\mathbf{X} = \begin{bmatrix} EX_1 \\ EX_2 \\ \vdots \\ EX_n \end{bmatrix}.$$

- ▶ A **random matrix** is a matrix whose elements are random variables. We have an m by n random matrix \mathbf{M} as

$$\mathbf{M} = \begin{bmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ X_{m1} & X_{m2} & \dots & X_{mn} \end{bmatrix}.$$

- ▶ Sometimes it is written as $\mathbf{M} = [X_{ij}]$.

Expectation

- ▶ The mean matrix of \mathbf{M} is given as

$$E\mathbf{M} = \begin{bmatrix} EX_{11} & EX_{12} & \dots & EX_{1n} \\ EX_{21} & EX_{22} & \dots & EX_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ EX_{m1} & EX_{m2} & \dots & EX_{mn} \end{bmatrix}.$$

- ▶ Linearity of expectation is valid for random vectors and matrices.
- ▶ For the random vector $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$ with \mathbf{A} , a fixed (non-random) m by n matrix and \mathbf{b} a fixed m -dimensional vector, we have

$$E\mathbf{Y} = \mathbf{A}E\mathbf{X} + \mathbf{b}.$$

- ▶ If $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_k$ are n -dimensional random vectors, then we have

$$E[\mathbf{X}_1 + \mathbf{X}_2 + \dots + \mathbf{X}_k] = E\mathbf{X}_1 + E\mathbf{X}_2 + \dots + E\mathbf{X}_k.$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

- For a random vector \mathbf{X} , we define the **correlation** and **covariance** matrix as

Correlation matrix of \mathbf{X} :

$$\mathbf{R}_X = E[\mathbf{X}\mathbf{X}^T]$$

Covariance matrix of \mathbf{X} :

$$\mathbf{C}_X = E[(\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T] = \mathbf{R}_X - E\mathbf{X}E\mathbf{X}^T$$

Methods for More
Than Two
Random Variables

Joint Distributions
and Independence
Sums of Random
Variables

Moment Generating
Functions

Characteristic
Functions

Random Vectors

Probability Bounds

The Union Bounds
and its Extensions

Markov and
Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz
Inequality

Jensen's Inequality

$$\mathbf{R}_X = E[\mathbf{X}\mathbf{X}^T] = \begin{bmatrix} EX_1^2 & E[X_1X_2] & \dots & E[X_1X_n] \\ EX_2X_1 & E[X_2^2] & \dots & E[X_2X_n] \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ E[X_nX_1] & E[X_nX_2] & \dots & E[X_n^2] \end{bmatrix},$$

$$\mathbf{C}_X = E[(\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T]$$

$$= \begin{bmatrix} \text{Var}(X_1) & \text{Cov}(X_1, X_2) & \dots & \text{Cov}(X_1, X_n) \\ \text{Cov}(X_2, X_1) & \text{Var}(X_2) & \dots & \text{Cov}(X_2, X_n) \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ \text{Cov}(X_n, X_1) & \text{Cov}(X_n, X_2) & \dots & \text{Var}(X_n) \end{bmatrix}.$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Properties of the Covariance Matrix

- ▶ The covariance matrix $\mathbf{C}_\mathbf{X}$ is a symmetric matrix.

$$c_{ij} = \text{Cov}(X_i, X_j) = \text{Cov}(X_j, X_i) = c_{ji}.$$

- ▶ $\mathbf{C}_\mathbf{X}$ can be diagonalized and all the eigenvalues of $\mathbf{C}_\mathbf{X}$ are real.
- ▶ A symmetric matrix \mathbf{M} is
 - ▶ **positive semi-definite**(PSD) if, for all vectors \mathbf{b} , $\mathbf{b}^T \mathbf{M} \mathbf{b} \geq 0$.
 - ▶ **positive definite** (PD), if for all vectors $\mathbf{b} \neq 0$, $\mathbf{b}^T \mathbf{M} \mathbf{b} > 0$.
- ▶ The covariance matrix is positive semi-definite.

Methods for More
Than Two
Random Variables

Joint Distributions
and Independence
Sums of Random
Variables

Moment Generating
Functions

Characteristic
Functions

Random Vectors

Probability Bounds

The Union Bounds
and its Extensions

Markov and
Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz
Inequality

Jensen's Inequality

Properties of the Covariance Matrix

- ▶ **Theorem:** Let X be a random vector with n elements. Then, its covariance matrix \mathbf{C}_X is positive semi-definite(PSD).
- ▶ **Theorem:** Let X be a random vector with n elements. Then its covariance matrix \mathbf{C}_X is positive definite (PD), if and only if all its eigenvalues are larger than zero. Equivalently, \mathbf{C}_X is positive definite (PD), if and only if $\det(\mathbf{C}_X) > 0$.
- ▶ For random vectors \mathbf{X} and \mathbf{Y} , the **cross correlation matrix** is given by

$$\mathbf{R}_{\mathbf{XY}} = E[\mathbf{XY}^T]$$

- ▶ The **cross covariance matrix** of \mathbf{X} and \mathbf{Y} is

$$\mathbf{C}_{\mathbf{XY}} = E[(\mathbf{X} - E\mathbf{X})(\mathbf{Y} - E\mathbf{Y})^t]$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Functions of Random Vectors: The Method of Transformations

- ▶ A function of a random vector is a random vector. We can use the method of transformations to find distributions of random vectors.
- ▶ If \mathbf{X} is an n -dimensional random vector with joint PDF $f_{\mathbf{X}}(\mathbf{x})$. Let $G : \mathbb{R}^n \mapsto \mathbb{R}^n$ be a continuous and invertible function with continuous partial derivatives and let $H = G^{-1}$.
- ▶ Let the random vector \mathbf{Y} be given by $\mathbf{Y} = G(\mathbf{X})$ and thus $\mathbf{X} = G^{-1}(\mathbf{Y}) = H(\mathbf{Y})$.

$$\mathbf{X} = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix} = \begin{bmatrix} H_1(Y_1, Y_2, \dots, Y_n) \\ H_2(Y_1, Y_2, \dots, Y_n) \\ \vdots \\ H_n(Y_1, Y_2, \dots, Y_n) \end{bmatrix}.$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Functions of Random Vectors: The Method of Transformations

- The PDF of \mathbf{Y} , $f_{Y_1, Y_2, \dots, Y_n}(y_1, y_2, \dots, y_n)$, is given by

$$f_{\mathbf{Y}}(\mathbf{y}) = f_{\mathbf{X}}(H(\mathbf{y}))|J|$$

where J is the Jacobian of H defined by

$$J = \det \begin{bmatrix} \frac{\partial H_1}{\partial y_1} & \frac{\partial H_1}{\partial y_2} & \cdots & \frac{\partial H_1}{\partial y_n} \\ \frac{\partial H_2}{\partial y_1} & \frac{\partial H_2}{\partial y_2} & \cdots & \frac{\partial H_2}{\partial y_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial H_n}{\partial y_1} & \frac{\partial H_n}{\partial y_2} & \cdots & \frac{\partial H_n}{\partial y_n} \end{bmatrix},$$

and evaluated at (y_1, y_2, \dots, y_n) .

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Normal (Gaussian) Random Vectors

Random variables X_1, X_2, \dots, X_n are said to be **jointly normal** if, for all $a_1, a_2, \dots, a_n \in \mathbb{R}$, the random variable

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n$$

is a normal random variable.

A random vector

$$\mathbf{X} = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix}$$

is said to be **normal** or **Gaussian** if the random variables X_1, X_2, \dots, X_n are jointly normal.

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

For a normal random vector \mathbf{X} with mean \mathbf{m} and covariance matrix \mathbf{C} , the PDF is given by

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{(2\pi)^{\frac{n}{2}} \sqrt{\det \mathbf{C}}} \exp \left\{ -\frac{1}{2} (\mathbf{x} - \mathbf{m})^T \mathbf{C}^{-1} (\mathbf{x} - \mathbf{m}) \right\}$$

Methods for More
Than Two
Random Variables

Joint Distributions
and Independence

Sums of Random
Variables

Moment Generating
Functions

Characteristic
Functions

Random Vectors

Probability Bounds

The Union Bounds
and its Extensions

Markov and
Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz
Inequality

Jensen's Inequality

Normal (Gaussian) Random Vectors

If $\mathbf{X} = [X_1, X_2, \dots, X_n]^T$ is a normal random vector, and we know $\text{Cov}(X_i, X_j) = 0$ for all $i \neq j$, then X_1, X_2, \dots, X_n are independent.

If $\mathbf{X} = [X_1, X_2, \dots, X_n]^T$ is a normal random vector, $\mathbf{X} \sim N(\mathbf{m}, \mathbf{C})$, \mathbf{A} is an m by n fixed matrix, and \mathbf{b} is an m -dimensional fixed vector, then the random vector $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$ is a normal random vector with mean $\mathbf{A}\mathbf{E}\mathbf{X} + \mathbf{b}$ and covariance matrix $\mathbf{A}\mathbf{C}\mathbf{A}^T$.

$$\mathbf{Y} \sim N(\mathbf{A}\mathbf{E}\mathbf{X} + \mathbf{b}, \mathbf{A}\mathbf{C}\mathbf{A}^T)$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

- ▶ **Union bound** or **Boole's inequality** is applicable when you need to show that the probability of union of some events is less than some value.

The Union Bound

For any events A_1, A_2, \dots, A_n , we have

$$P\left(\bigcup_{i=1}^n A_i\right) \leq \sum_{i=1}^n P(A_i)$$

- ▶ It is widely used in the area of *random graphs*.

Methods for More
Than Two
Random Variables

Joint Distributions
and Independence
Sums of Random
Variables

Moment Generating
Functions

Characteristic
Functions

Random Vectors

Probability Bounds

The Union Bounds
and its Extensions

Markov and
Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz
Inequality

Jensen's Inequality

Union Bound and its Extensions

- ▶ We can extend the union bounds and obtain the lower and upper bound on probability of union of events.
- ▶ These bounds are known as **Bonferroni inequalities**.
- ▶ Start writing the inclusion-exclusion formula. If you stop at the first term, you obtain an upper bound on the probability of union. If you stop at the second term, you obtain a lower bound. If you stop at the third term, you obtain an upper bound, etc.
- ▶ In general, if you write an odd number of terms, you get an upper bound and if you write an even number of terms, you get a lower bound.

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Union Bound and its Extensions

Generalization of the Union Bound: Bonferroni inequalities For any events A_1, A_2, \dots, A_n , we have

$$P\left(\bigcup_{i=1}^n A_i\right) \leq \sum_{i=1}^n P(A_i)$$

$$P\left(\bigcup_{i=1}^n A_i\right) \geq \sum_{i=1}^n P(A_i) - \sum_{i < j} P(A_i \cap A_j)$$

$$P\left(\bigcup_{i=1}^n A_i\right) \leq \sum_{i=1}^n P(A_i) - \sum_{i < j} P(A_i \cap A_j) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k)$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Markov's Inequality

If X is any nonnegative random variable, then

$$P(X \geq a) \leq \frac{EX}{a}, \text{ for any } a > 0.$$

Chebyshev's Inequality

If X is any random variable, then for any $b > 0$ we have

$$P(|X - EX| \geq b) \leq \frac{\text{Var}(X)}{b^2}.$$

Methods for More Than Two Random Variables

Joint Distributions and Independence

Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Chernoff Bounds:

$$P(X \geq a) \leq e^{-sa} M_X(s), \text{ for all } s > 0,$$

$$P(X \leq a) \leq e^{-sa} M_X(s), \text{ for all } s < 0$$

Methods for More
Than Two
Random Variables

Joint Distributions
and Independence

Sums of Random
Variables

Moment Generating
Functions

Characteristic
Functions

Random Vectors

Probability Bounds

The Union Bounds
and its Extensions

Markov and
Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz
Inequality

Jensen's Inequality

Comparison between Markov, Chebyshev, and Chernoff Bounds

- ▶ We found upper bounds on $P(X \geq \alpha n)$ for $X \sim \text{Binomial}(n, p)$ with $p = \frac{1}{4}$ and $\alpha = \frac{3}{4}$:

$$P(X \geq \frac{3n}{4}) \leq \frac{2}{3} \quad \text{Markov,}$$

$$P(X \geq \frac{3n}{4}) \leq \frac{4}{n} \quad \text{Chebyshev,}$$

$$P(X \geq \frac{3n}{4}) \leq \left(\frac{16}{27}\right)^{\frac{n}{4}} \quad \text{Chernoff.}$$

- ▶ The bound given by Markov is the “weakest”.
- ▶ The bound given by Chebyshev’s inequality is “stronger” than the one given by Markov’s inequality. Note that $\frac{4}{n}$ goes to zero as n goes to infinity.
- ▶ The strongest bound is the Chernoff bound. It goes to zero exponentially fast.

Methods for More Than Two Random Variables

Joint Distributions and Independence

Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Cauchy-Schwarz Inequality

Cauchy-Schwarz Inequality

For any two random variables X and Y , we have

$$|EXY| \leq \sqrt{E[X^2]E[Y^2]},$$

where equality holds if and only if $X = \alpha Y$, for some constant $\alpha \in \mathbb{R}$.

Methods for More
Than Two
Random Variables

Joint Distributions
and Independence
Sums of Random
Variables

Moment Generating
Functions

Characteristic
Functions

Random Vectors

Probability Bounds

The Union Bounds
and its Extensions

Markov and
Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz
Inequality

Jensen's Inequality

Definition: Consider a function $g : I \rightarrow \mathbb{R}$, where I is an interval in \mathbb{R} . We say that g is a **convex** function if for any two points x and y in I and any $\alpha \in [0, 1]$, we have

$$g(\alpha x + (1 - \alpha)y) \leq \alpha g(x) + (1 - \alpha)g(y).$$

We say that g is **concave** if

$$g(\alpha x + (1 - \alpha)y) \geq \alpha g(x) + (1 - \alpha)g(y).$$

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Jensen's Inequality




Figure: Pictorial representation of a convex function and a concave function.

Methods for More Than Two Random Variables

Joint Distributions and Independence

Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality

Jensen's Inequality:

If $g(x)$ is a convex function on R_X , and $E[g(X)]$ and $g(E[X])$ are finite, then

$$E[g(X)] \geq g(E[X]).$$

A twice-differentiable function $g : I \rightarrow \mathbb{R}$ is convex if and only if $g''(x) \geq 0$ for all $x \in I$.

Methods for More Than Two Random Variables

Joint Distributions and Independence
Sums of Random Variables

Moment Generating Functions

Characteristic Functions

Random Vectors

Probability Bounds

The Union Bounds and its Extensions

Markov and Chebyshev Inequalities

Chernoff Bounds

Cauchy-Schwarz Inequality

Jensen's Inequality