


Problem 1



(a)



(b)
$$P = \begin{bmatrix} P_{00} & P_{01} \\ P_{10} & P_{11} \end{bmatrix} \begin{matrix} q \\ 1 \end{matrix}$$

$$\begin{bmatrix} 1-p & p \\ q & 1-q \end{bmatrix} = I$$

$$(c) \quad P^{(2)} = \begin{bmatrix} P_{00}^{(2)} & P_{01}^{(2)} \\ P_{10}^{(2)} & P_{11}^{(2)} \end{bmatrix}$$

$$= P^2 = \begin{bmatrix} 1-p & p \\ q & 1-q \end{bmatrix} \begin{bmatrix} 1-p & p \\ q & 1-q \end{bmatrix}$$

$$P^{(n)} = P^n$$


✓

$$\approx \begin{bmatrix} (1-p)^2 + pq & p^2 + (1-q)^2 \\ q(1-p) + (1-q)q & pq + (1-q)^2 \end{bmatrix}$$

Problem 2 10 ms

10 ms

10 ms



talking
silent

$$P = \begin{bmatrix} 139 \\ 140 \\ 1 \\ 100 \end{bmatrix}$$

$$\begin{matrix} 140 \\ 99 \\ 100 \end{matrix}$$

$$P_{00} = ?$$

$$1 - \frac{1}{140} = \frac{139}{140}$$

$$P_{01} = ? = \frac{1}{140}$$

$$P_{11} = ?$$

$$1 - \frac{1}{100} = \frac{99}{100}$$


$$P_{10} = ? = \frac{1}{100}$$

silent


talking

Problem 3


discrete random walk \rightarrow a kind of Markov chain



$$S = \{ \dots, -2, -1, 0, 1, 2, \dots \}$$



Problem 4



$$P = \begin{bmatrix} 0.9 & 0.1 \\ 0.01 & 0.99 \end{bmatrix} = I \quad \checkmark$$




106%



Share

Problem 5:

Consider the Markov chain with three states $S = 1, 2, 3$, that has the following state transition diagram. Suppose $P[X_1 = 1] = 1/2$ and $P[X_1 = 2] = 1/4$.



- (a) Find the state transition matrix for this chain.
- (b) Find $P[X_1 = 3, X_2 = 2, X_3 = 1]$.
- (c) Find $P(X_1 = 3, X_3 = 1)$.

Problem 6:

Search tools

Export PDF

Adobe Export PDF

Convert PDF Files to Word or Excel Online

[Select PDF File](#)

Problem_Set_10.pdf x

Convert to

Microsoft Word (*.docx) v

Document Language:
English (U.S.) [Change](#)

Convert


Convert and edit PDFs
with Acrobat Pro DC

[Start Free Trial](#)

ENG

3:33 PM






Share

Problem 5:Consider the Markov chain with three states $S = \{1, 2, 3\}$, that has the following state transition diagramSuppose $P[X_1 = 1] = 1/2$ and $P[X_1 = 2] = 1/4$.

$$P[X_1 = 3] = ?$$

$$1/2 - 1/4$$



$$= 1/2 - 1/4$$

$$= 1/4$$

$$\pi^{(n)} = \pi^{(n-1)} P$$

$$\pi^{(n)} = [P[X_n = 1] \ P[X_n = 2] \ P[X_n = 3]]$$

(a) Find the state transition matrix for this chain.

(b) Find $P[X_1 = 3, X_2 = 2, X_3 = 1]$.(c) Find $P(X_1 = 3, X_3 = 1)$.**Problem 6:**

Search tools

Export PDF

Adobe Export PDF

Convert PDF Files to Word or Excel Online

Select PDF File

Problem_Set_10.pdf x

Convert to

Microsoft Word (*.docx) v

Document Language:
English (U.S.) Change

Convert

Convert and edit PDFs
with Acrobat Pro DC

Start Free Trial

$$P = \begin{bmatrix} \frac{1}{4} & 0 & \frac{3}{4} \\ \frac{1}{2} & 0 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{4} & \frac{1}{4} \end{bmatrix}$$

$$C = \{X_1 = 3\}$$

$$B = \{X_2 = 2\}$$

$$A = \{X_3 = 1\}$$

$$A = \{X_1 = 3\}$$

$$B = \{X_2 = 2\}$$

$$C = \{X_3 = 1\}$$

$$P[X_1=3, X_2=2, X_3=1] = P[X_1=3] \cdot P[X_2=2|X_1=3] \cdot P[X_3=1|X_1=3, X_2=2]$$

$$(b) P[X_1=3, X_2=2, X_3=1] = P[X_3=1 | X_2=2, X_1=3] \cdot P[X_2=2 | X_1=3] \cdot P[X_1=3]$$

$$P[A, B] = P[A|B]P[B]$$

$$P[A, B, C] = P[A|B, C]P[B, C]$$

$$= P[A|B, C]P[B|C]P[C]$$

$$= P[X_3=1 | X_2=2, X_1=3] \cdot P[X_2=2 | X_1=3] \cdot P[X_1=3]$$

$$= \left(\frac{1}{2}\right) \cdot \left(\frac{1}{4}\right) \cdot \left(\frac{1}{4}\right) = \frac{1}{32}$$

(c) $P[X_1=3, X_3=1] = \sum_{k=1}^3 P[X_1=3, X_2=k, X_3=1]$


2 steps

$$= \sum_{k=1}^3 P[X_1=3] P_{3k} P_{k1}$$

X_2

Low & total probability

$$P[A] = P[A \cap B_2] + P[A \cap B_3] + P[A \cap B_4]$$



(a)

$$P[X_1=3, X_2=2, X_3=1] = P[X_1=3] P_{32} P_{21}$$

$$P[X_3=j | X_1=i] = \sum_{k=1}^3 p_{ik} p_{kj} \quad \begin{matrix} n=1 \\ m=1 \end{matrix}$$

•

$$P[X_3=j, X_1=i]$$

$$= P[X_1=i] P[X_3=j | X_1=i]$$

$$= P[X_1=i] \sum_{k=1}^3 p_{ik} p_{kj}$$

Problem 6




106%



Share

Problem 6:

Consider the Markov chain shown in the figure below. It is assumed that when there is an arrow from state i to state j , then $p_{ij} > 0$. Find the communicating classes (also called equivalence classes) for this Markov chain.

**Problem 7:**

Consider the Markov chain in Problem 6.

- Is Class 1 = {state 1, state 2} aperiodic?
- Is Class 2 = {state 3, state 4} aperiodic?

Search tools

Export PDF

Adobe Export PDF

Convert PDF Files to Word or Excel Online

Select PDF File

Problem_Set_10.pdf

Convert to

Microsoft Word (*.docx)

Document Language:
English (U.S.) [Change](#)

Convert

Convert and edit PDFs
with Acrobat Pro DC[Start Free Trial](#)




106%



Share

(c) Find $\lim_{n \rightarrow \infty} (x_1 - 0, x_3 - 1)$.**Problem 6:**

Consider the Markov chain shown in the figure below. It is assumed that when there is an arrow from state i to state j , then $p_{ij} > 0$. Find the communicating classes (also called equivalence classes) for this Markov chain.

**Problem 7:**

Consider the Markov chain in Problem 6.

- (a) Is Class 1 = {state 1, state 2} aperiodic?
- (b) Is Class 2 = {state 3, state 4} aperiodic?

Search tools

Export PDF

Adobe Export PDF

Convert PDF Files to Word or Excel Online

Select PDF File

Problem_Set_10.pdf

Convert to

Microsoft Word (*.docx)

Document Language:
English (U.S.) [Change](#)

Convert

Convert and edit PDFs
with Acrobat Pro DC

[Start Free Trial](#)

ENG

4:14 PM








106%



Share

Markov chain.

**Problem 7:**

Consider the Markov chain in Problem 6.

- (a) Is Class 1 = {state 1, state 2} aperiodic?
- (b) Is Class 2 = {state 3, state 4} aperiodic?

Search tools

Export PDF

Adobe Export PDF

Convert PDF Files to Word or Excel Online

Select PDF File

Problem_Set_10.pdf

Convert to

Microsoft Word (*.docx)


Document Language:
English (U.S.) [Change](#)

Convert

Convert and edit PDFs
with Acrobat Pro DC[Start Free Trial](#)

**Problem 6:**

Consider the Markov chain shown in the figure below. It is assumed that when there is an arrow from state i to state j , then $p_{ij} > 0$. Find the communicating classes (also called equivalence classes) for this Markov chain.



$P_{88}^{(1)} = 0$
 $P_{88}^{(2)} = 0$
 $P_{88}^{(3)} > 0$
 $P_{88}^{(4)} = 0$
 $P_{88}^{(5)} = 0$
 $P_{88}^{(6)} > 0$

$P_{66}^{(1)} = 0$
 $P_{66}^{(2)} > 0$
 $P_{66}^{(3)} > 0$

$\gcd\{2, 3\} = 1$

Problem 7:

Consider the Markov chain in Problem 6.

- Is Class 1 = {state 1, state 2} aperiodic?
- Is Class 2 = {state 3, state 4} aperiodic?

Search tools

Export PDF

Adobe Export PDF

Convert PDF Files to Word or Excel Online

Select PDF File

Problem_Set_10.pdf

Convert to

Microsoft Word (*.docx)

Document Language:
English (U.S.) Change


Convert

Convert and edit PDFs
with Acrobat Pro DC

Start Free Trial

Problem 8

$$S = \{0, 1\}$$



where a and b are in $[0, 1]$


such that

$0 \leq a+b \leq 2$

$a \neq 0, b \neq 0$

$a \neq 1, b \neq 1$

$\pi^{(0)}$ = $\begin{bmatrix} P[X_0=0] & P[X_0=1] \end{bmatrix} = \begin{bmatrix} a & 1-a \end{bmatrix}$



To have a limiting distribution for a finite markov chain we need

→ irreducible (= 1 class that is recurrent)

→ aperiodic

$$P^{(n)} = P^{(n)} \xrightarrow{1 \text{ step}}$$

(b) $P^{(n)} = \frac{1}{a+b} \begin{bmatrix} b-a & a \\ b & a \end{bmatrix} + \frac{(1-a-b)^n}{2+b} \begin{bmatrix} a & -a \\ -b & b \end{bmatrix}$

$$\begin{bmatrix} P_{00}^{(n)} & P_{01}^{(n)} \\ P_{10}^{(n)} & P_{11}^{(n)} \end{bmatrix}$$

$$\lim_{n \rightarrow \infty} P^{(n)}$$

$$\lim_{n \rightarrow \infty} \underline{\underline{\pi}}^{(n)}$$

$$= \pi^{(0)} \lim_{n \rightarrow \infty} P^{(n)}$$

$$\pi^{(n)} = \frac{\pi^{(0)} P^{(n)}}{\sum [\alpha \quad 1-\alpha]}$$

$$\frac{1}{2+b} [\alpha \quad 1-\alpha] \begin{bmatrix} b & a \\ b & a \end{bmatrix} = \frac{1}{2+b} \begin{bmatrix} b & a \end{bmatrix}$$

$$[P[X_n=0] \quad P[X_n=1]]$$

$$\lim_{n \rightarrow \infty} \pi^{(n)} = \lim_{n \rightarrow \infty} \frac{\pi^{(0)} P^{(n)}}{\sum}$$

$$= \frac{\pi^{(0)}}{\sum} \lim_{n \rightarrow \infty} P^{(n)}$$

$n \rightarrow \infty$



where a and b are two real numbers in the interval $[0, 1]$ such that $0 < a + b < 2$. Suppose that the system is in state 0 at time $n = 0$ with probability α , i.e., $\pi^{(0)} = [P(X_0 = 0) \ P(X_0 = 1)] = [\alpha \ 1 - \alpha]$, where $\alpha \in [0, 1]$.

- (a) Sketch the Markov chain. How many classes are in the chain, are they recurrent/transient, and are they periodic/aperiodic?
 (b) Given that

show that

$\lim_{n \rightarrow \infty}$

$$P^n = \frac{1}{a+b} \begin{bmatrix} b & a \\ b & a \end{bmatrix} + \frac{(1-a-b)^n}{a+b} \begin{bmatrix} a & -a \\ -b & b \end{bmatrix}$$

$$\lim_{n \rightarrow \infty} P^n = \frac{1}{a+b} \begin{bmatrix} b & a \\ b & a \end{bmatrix}$$

and that the Markov chain has a limiting distribution given by

$$\lim_{n \rightarrow \infty} \pi^{(n)} = \left[\frac{b}{a+b} \quad \frac{a}{a+b} \right]$$

- (c) Find the stationary distribution for this chain by solving the balance equations. Is this also the limiting distribution for this chain? Why or why not?
 (d) Consider the case when $a = b = 1$. Sketch the Markov chain and comment on the existence of a limiting distribution for this Markov chain.
 (e) Consider the case when $a = b = 0$. Sketch the Markov chain and comment on the existence of a limiting distribution for this Markov chain.

Search tools

Export PDF

Adobe Export PDF

Convert PDF Files to Word or Excel Online

Select PDF File

Problem_Set_10.pdf

Convert to

Microsoft Word (*.docx)

Document Language:
English (U.S.) [Change](#)

Convert

Convert and edit PDFs
with Acrobat Pro DC

[Start Free Trial](#)



ENG

4:59 PM



(c)

$$\underline{\pi} = \underline{\pi} P$$

$$\sum_{j=0}^1 \underline{\pi}_j = 1$$

$$\underline{\pi}^{(n)} = \pi^{(0)} P^n$$

$$\text{rel} \hookrightarrow \underline{\pi} = \underline{\pi} P$$

$$\underline{\pi} = \begin{bmatrix} \pi_0 & \pi_1 \end{bmatrix}$$

$$\lim_{n \rightarrow \infty} P[\underline{x}_n = 0] \quad \lim_{n \rightarrow \infty} P[x_n = 1]$$



Search tools

Export PDF

Adobe Export PDF

Convert PDF Files to Word
or Excel Online

Select PDF File

Problem_Set_10.pdf x

Convert to

Microsoft Word (*.docx)

Document Language:
English (U.S.) [Change](#)

Convert

Convert and edit PDFs
with Acrobat Pro DC[Start Free Trial](#)

(c) Is Class 4 = {state 6, state 7, state 8} aperiodic?

Problem 8:

Consider a Markov chain with two possible states, $S = \{0, 1\}$. In particular, suppose that the transition matrix is given by

$$P = \begin{bmatrix} 1-a & a \\ b & 1-b \end{bmatrix},$$

where a and b are two real numbers in the interval $[0, 1]$ such that $0 < a + b < 2$. Suppose that the system is in state 0 at time $n = 0$ with probability α , i.e., $\pi^{(0)} = [P(X_0 = 0) \ P(X_0 = 1)] = [\alpha \ 1 - \alpha]$, where $\alpha \in [0, 1]$.

(a) Sketch the Markov chain. How many classes are in the chain, are they recurrent/transient, and are they periodic/aperiodic?

(b) Given that

$$P^n = \frac{1}{a+b} \begin{bmatrix} b & a \\ b & a \end{bmatrix} + \frac{(1-a-b)^n}{a+b} \begin{bmatrix} a & -a \\ -b & b \end{bmatrix}.$$

show that

$$\lim_{n \rightarrow \infty} P^n = \frac{1}{a+b} \begin{bmatrix} b & a \\ b & a \end{bmatrix}.$$

$a \neq 1, b \neq 1$

$$\pi = \pi P \Rightarrow \begin{bmatrix} \pi_0 & \pi_1 \end{bmatrix} = \begin{bmatrix} \pi_0 & \pi_1 \end{bmatrix} \begin{bmatrix} 1-a & a \\ b & 1-b \end{bmatrix}$$

Final answer: $\pi = \begin{bmatrix} \frac{b}{a+b} & \frac{a}{a+b} \end{bmatrix}$

$$\pi_0 + \pi_1 = 1$$

$$\begin{bmatrix} \pi_0 & \pi_1 \end{bmatrix} = \begin{bmatrix} (1-a)\pi_0 + b\pi_1 & a\pi_0 + (1-b)\pi_1 \end{bmatrix}$$

(1)

$$\pi_0 = (1-a)\pi_0 + b\pi_1$$

~~$$\pi_1 = a\pi_0 + (1-b)\pi_1$$~~

$$\pi_0 + \pi_1 = 1$$

(2)

From (2)

$$\pi_0 = 1 - \pi_1 \Rightarrow \pi_1 = 1 - \pi_0$$

plug in to (1)

$$\pi_0 = (1-a)\pi_0 + b(1-\pi_0)$$

~~π_0~~

$$a\pi_0 = b - b\pi_0 \Rightarrow \pi_0 = \frac{b}{a+b}$$

$$\pi_1 = \frac{a}{a+b}$$

$$\pi = \pi P$$

$$\sum_j \pi_j = 1$$

→ find stationary
dist.

because my chain
is irreducible
and aperiodic

↓
this is the limiting dist
 $\lim_{n \rightarrow \infty} \pi(n)$


(d) Is the stationary distribution a limiting distribution for the chain?

Problem 10:
Consider the following Markov chain

$P =$

$\frac{1}{2}$	$\frac{1}{2}$	0
0	$\frac{1}{3}$	$\frac{2}{3}$
$\frac{1}{2}$	$\frac{1}{2}$	0

$= 1$



$$\underline{\pi} = \underline{\pi} P$$
$$\sum \pi_j = 1$$

π is
stationary
 \downarrow
limiting.

- (a) Is this chain irreducible?
- (b) Is this chain aperiodic?
- (c) Find the stationary distribution for this chain.
- (d) Is the stationary distribution a limiting distribution for the chain?

Search tools

Export PDF

Adobe Export PDF

Convert PDF Files to Word or Excel Online

Select PDF File

Problem_Set_10.pdf

Convert to

Microsoft Word (*.docx)

Document Language: English (U.S.)

Convert

Convert and edit PDFs with Acrobat Pro DC

Start Free Trial

$$P = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & \frac{1}{3} & \frac{2}{3} \\ \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix}$$

$$\pi < \pi P, \sum \pi_i = 1$$

$$\begin{bmatrix} \pi_1 & \pi_2 & \pi_3 \end{bmatrix} = \begin{bmatrix} \pi_1 & \pi_2 & \pi_3 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & \frac{1}{3} & \frac{2}{3} \\ \frac{1}{2} & \frac{1}{2} & 0 \end{bmatrix}$$

$$\begin{bmatrix} \pi_1 & \pi_2 & \pi_3 \end{bmatrix} = \begin{bmatrix} \frac{1}{2}\pi_1 + \frac{1}{2}\pi_3 & \frac{1}{2}\pi_1 + \frac{1}{3}\pi_2 + \frac{1}{2}\pi_3 & \frac{2}{3}\pi_2 \end{bmatrix}$$

$$\pi_1 = \frac{1}{2}\pi_1 + \frac{1}{2}\pi_3$$

$$\pi_2 = \frac{1}{2}\pi_1 + \frac{1}{3}\pi_2 + \frac{1}{2}\pi_3$$

$$\pi_3 = \frac{2}{3}\pi_2$$

$$\pi_1 + \pi_2 + \pi_3 = 1$$

**Problem 8:**

Consider a Markov chain with two possible states, $S = \{0, 1\}$. In particular, suppose that the transition matrix is given by

$$P = \begin{bmatrix} 1-a & a \\ b & 1-b \end{bmatrix},$$

where a and b are two real numbers in the interval $[0, 1]$ such that $0 < a + b < 2$. Suppose that the system is in state 0 at time $n = 0$ with probability α , i.e., $\pi^{(0)} = [P(X_0 = 0) \ P(X_0 = 1)] = [\alpha \ 1 - \alpha]$, where $\alpha \in [0, 1]$.

- (a) Sketch the Markov chain. How many classes are in the chain, are they recurrent/transient, and are they periodic/aperiodic?
- (b) Given that

$$P^n = \frac{1}{a+b} \begin{bmatrix} b & a \\ b & a \end{bmatrix} + \frac{(1-a-b)^n}{a+b} \begin{bmatrix} a & -a \\ -b & b \end{bmatrix}.$$

show that

$$\lim_{n \rightarrow \infty} P^n = \frac{1}{a+b} \begin{bmatrix} b & a \\ b & a \end{bmatrix}.$$

and that the Markov chain has a limiting distribution given by

$$\lim_{n \rightarrow \infty} \pi^{(n)} = \left[\frac{b}{a+b} \quad \frac{a}{a+b} \right].$$

- (c) Find the stationary distribution for this chain by solving the balance equations. Is this also the limiting distribution for this chain? Why or why not?

Search tools

Export PDF

Adobe Export PDF

Convert PDF Files to Word or Excel Online

Select PDF File

Problem_Set_10.pdf

Convert to

Microsoft Word (*.docx)

Document Language:

English (U.S.) [Change](#)

Convert

Convert and edit PDFs with Acrobat Pro DC

[Start Free Trial](#)

ENG

5:01 PM

