

Evaluation of Pitch-Class Set Similarity Measures for Tonal Analysis (Literature Review)

Nicholas Harley - Masters Thesis UPF

June 10, 2014

Contents

| | | |
|----------|---|----------|
| 1 | Introduction | 3 |
| 2 | Tonality | 3 |
| 2.1 | Defining Tonality | 3 |
| 2.1.1 | General Definition | 3 |
| 2.1.2 | Babbitt's Domains | 3 |
| 2.2 | Modelling Tonality | 3 |
| 2.2.1 | Tonality as Context | 3 |
| 2.2.2 | Tonality in MIR | 4 |
| 2.2.3 | Similarity | 4 |
| 2.3 | The Semantic Gap | 4 |
| 2.3.1 | Acoustic Domain | 4 |
| 2.3.2 | Graphemic Domain | 4 |
| 2.3.3 | Problems | 4 |
| 2.4 | Systematicity | 5 |
| 2.4.1 | The Musical Surface | 5 |
| 2.4.2 | Systematic Description | 5 |
| 3 | Pitch-Class Set Theory | 5 |
| 3.1 | Pitch Class | 5 |
| 3.2 | Set Classification | 5 |
| 3.3 | Vector Analysis | 6 |
| 3.3.1 | Membership and Inclusion | 6 |
| 3.3.2 | Embedding Number | 7 |
| 3.3.3 | Subset Vectors | 7 |
| 3.3.4 | Notation | 8 |
| 3.4 | Set-Class Similarity | 8 |
| 3.4.1 | Similarity Relations | 8 |
| 3.4.2 | Similarity Measures | 8 |
| 3.4.3 | Castren's Criteria | 9 |
| 3.5 | Perceptual Relevance | 10 |
| 3.5.1 | Octave Equivalence | 10 |
| 3.5.2 | Set-Class Equivalence | 10 |
| 3.5.3 | Perceived vs Theoretical Similarity | 10 |
| 3.6 | PC-set Theory for Analysis | 11 |

| | | |
|-----------|---|-----------|
| 4 | Multidimensional Scaling | 11 |
| 4.1 | Non-Metric MDS | 11 |
| 4.2 | Cluster Analysis | 11 |
| 4.3 | MDS with Similarity Measures | 12 |
| 5 | Spatial Models of Tonality | 12 |
| 5.1 | Similarity and Distance | 12 |
| 5.2 | Spatial Representations | 12 |
| 5.3 | Cognitive Psychology | 12 |
| 5.4 | Set-Class Spaces | 12 |
| 6 | Similarity Measure Selection | 13 |
| 6.1 | Criteria | 13 |
| 6.2 | Cardinality | 13 |
| 6.3 | SC-type | 13 |
| 6.4 | Measure Type | 14 |
| 6.5 | Total Measures | 14 |
| 6.5.1 | Trivial Forms | 14 |
| 6.5.2 | Rahn: ATMEMB | 15 |
| 6.5.3 | Lewin: REL | 15 |
| 6.5.4 | Buchler: AvgSATSIM and TSATSIM | 15 |
| 6.5.5 | Castren: T%REL and RECREL | 15 |
| 6.5.6 | Scale of Values | 15 |
| 7 | Practical Analysis | 16 |
| 7.1 | Chords as PC-Sets | 16 |
| 7.2 | Repertoire | 16 |
| 7.2.1 | Billboard Chords | 16 |
| 7.2.2 | Beatles Chords | 16 |
| 7.2.3 | Diatoniscm in Bach and Victoria | 16 |
| 7.2.4 | Josquin? | 16 |
| 7.2.5 | Essen Collection | 17 |
| 7.3 | Comparison of Chords | 17 |
| 7.4 | Conclusions | 18 |
| 7.5 | Measure discrimination | 18 |
| 7.6 | Structure | 18 |
| 8 | Class Scape | 18 |
| 8.1 | replicate | 18 |
| 8.2 | genere specific analysis | 18 |
| 8.3 | structure analysis | 18 |
| 9 | MDS | 18 |
| 10 | Latex Stuff | 19 |
| A | SC Similarity Measures | 20 |
| A.1 | MORRIS | 20 |
| A.1.1 | K | 20 |
| A.1.2 | SIM | 20 |
| A.1.3 | ASIM | 21 |
| A.2 | LORD | 21 |
| A.2.1 | sf | 21 |
| A.3 | TEITELBAUM | 22 |
| A.3.1 | s.i. | 22 |

| | | |
|-------|-----------------------------|----|
| A.4 | ROGERS | 22 |
| A.4.1 | IcVD ₁ | 22 |
| A.4.2 | IcVD ₂ | 22 |
| A.4.3 | Cos(θ) | 23 |
| A.5 | RAHN | 23 |
| A.5.1 | AK | 23 |
| A.5.2 | MEMB _n | 24 |
| A.5.3 | TMEMB | 24 |
| A.5.4 | ATMEMB | 24 |
| A.6 | ISAACSON | 25 |
| A.6.1 | AMEMB2 | 25 |
| A.6.2 | IcVSIM | 25 |
| A.6.3 | ISIM2 | 25 |
| A.6.4 | ANGLE (Isaacson & Scott) | 26 |
| A.7 | LEWIN | 26 |
| A.7.1 | REL | 26 |
| A.7.2 | REL ₂ | 27 |
| A.8 | CASTREN | 27 |
| A.8.1 | Castren's Difference Vector | 27 |
| A.8.2 | nC%V | 27 |
| A.8.3 | %REL _n | 27 |
| A.8.4 | T%REL | 28 |
| A.8.5 | RECREL | 28 |
| A.9 | BUCHLER | 28 |
| A.9.1 | nSATV | 28 |
| A.9.2 | SATSIM _n | 28 |
| A.9.3 | AvgSATSIM | 29 |
| A.9.4 | TSATSIM | 29 |

```
%makeindex -s doc.ist -o doc.gls doc.glo
```

1 Introduction

The objectives of the present work are principally concerned with descriptive modelling of tonality using PC-set theory. In particular, the analytical potential of set-class similarity measures will be assessed and evaluated through comparison with similar work. The tentative goal, therefore, is to present this approach in such a way as to make it relevant to the wider community of music researchers, such as those in MIR.

Chapters 2-5 comprise a literature review containing background information and some basic theory. Chapter 2 addresses the challenges and problems involved in descriptive modelling of tonality as a means of justification for the proposed approach. Chapter 3 introduces the basic concepts of PC-set theory including set-class similarity measures. Chapter 4 gives a basic description of MDS techniques and how they can play a role in set-class analysis. Chapter 5 contains a review of the relevant tonal models that exist so as to give context work that follows.

2 Tonality

2.1 Defining Tonality

2.1.1 General Definition

Tonality is a notoriously complex musical phenomenon and numerous definitions have been proposed from a variety of viewpoints. Perhaps the most general definition is that provided by Hyer (Hyer2013): "... refers to the systematic arrangements of pitch phenomena and relations between them." Explanations of tonality have been provided through many different disciplines (acoustics, music theory, linguistics, cognitive psychology) and a detailed discussion of these areas is certainly beyond the scope of this work. However, it is generally agreed that tonality is an abstract cultural and cognitive construct that can have many different physical representations.

2.1.2 Babbitt's Domains

? proposed three domains to categorise different types of representation of music: acoustic (physical), auditory (perceived), graphemic (notated). Western music theory provides a lexicon for describing abstract tonal objects with terms such as note, chord and key. These objects have a hierarchical relationship and the meaning of these labels is highly dependent on musical context and the scale of observation. Musicological descriptions, which constitute the majority of reasoning about tonality, reside mainly in the Babbitt's graphemic domain, although arguably they reflect some aspects of the other two. Each domain, whilst connected to every other, provides only a projection of the musical whole and examination of tonality from just one will most likely result in an incomplete picture. However, these three domains provide a convenient framework for the discussion that follows.

2.2 Modelling Tonality

The challenge of mathematically modelling aspects of tonality has been approached in numerous ways and from different domains. In the graphemic domain, musicologists and composers have proposed theoretical models, attempting to rethink tonal theory from a mathematical perspective. These models employ different branches of mathematics such as geometry (?) or group theory (?) to describe harmonic structure. From the auditory domain, cognitive psychologists have built models of tonal induction based on perceptual ratings of tonal stimuli (?).

2.2.1 Tonality as Context

Many models approach the concept of tonality as a context, within which the relations and hierarchies of tonal phenomena can be understood. A sense of tonality can be induced when musical stimuli resemble some a priori contextual category. For western music of the major-minor period, key signatures comprise a collection of categories that give context to the tonal components of music. ? identifies three important aspects of tonality as context: dimensionality (the relatedness of categories), ambiguity (reference to two or

more categories simultaneously) and timing (the dynamics of tonal context). He highlights the importance of a model's capability to describe these aspects.

2.2.2 Tonality in MIR

The MIR community is primarily concerned with the extraction of tonal descriptors from audio signals such as chord and key estimates. Most systems use chroma features as a preliminary step, obtained by mapping STFT or CQ transform energies to chroma bins. Template matching is used to compare the chroma vectors to a tonal model (contextual category) using some distance measure. A commonly used tonal model for key estimation are the KK-profiles (Krumholtz, 1990) (e.g. in (Krumholtz, 1990)). Distance measures such as inner product (e.g. in (Krumholtz, 1990)) and fuzzy distance (e.g. in (Krumholtz, 1990)) are used to compare vectors. Statistical methods, such as HMMs, have been used for chord and key tracking (Krumholtz, 1990). Of additional interest in the field is the concept of musical similarity (for music recommendation, structure analysis, cover detection etc.). (Krumholtz, 1990) computed self-similarity matrices for visualisation of structure by correlating the MFCC feature vector time-series. (Krumholtz, 1990) proposed the application of this method to tonal feature vectors.

2.2.3 Similarity

The importance of defining the similarity or closeness between musical phenomena, be it theoretical, physical or perceptual, is central to almost every model of tonality and often leads to a geometric configuration of tonal objects. The concepts of similarity and distance is discussed further in Chapter 5 where a review of spatial models of tonality is given.

2.3 The Semantic Gap

2.3.1 Acoustic Domain

(Krumholtz, 1990) discusses, what is referred to in MIR as, the “Semantic Gap”: the inability of systems to achieve success rates beyond a conspicuous boundary. He examines the fundamental methodological groundings of MIR in terms of Babbitt's three domains, discussing the limits of each representation and regarding the incarnate nature of music. He concludes that the audio signal (acoustic domain) simply cannot contain all of the information that systems seek to retrieve. He points towards the auditory domain as the chief residence of music information and urges for it not to be overlooked in MIR and wider music research.

2.3.2 Graphemic Domain

Furthermore, Wiggins criticises the purely graphemic approach and the tendency of music research to presuppose musicological axioms. (Wiggins, 1992) argues that music (tonal) theory is, rather than a theory in the scientific sense, a highly developed folk psychology (internal human theory for explaining common behaviour). Thus, the rules of music theory are not like scientific laws but rather abstract descriptions of a specific musical behaviour. This idea challenges the validity of formalising such rules in mathematics and prompts the question, “What is actually being modelled?” He concludes that to apply mathematical models to musical output alone (scales or chords) without consideration of the musical mind is a scientific failure.

2.3.3 Problems

The two assertions of Wiggins sit contrary to a number of the aspects of the tonal models discussed in 2.2. Firstly, the major-minor paradigm, upon which so many approaches are based, whilst certainly possessing cognitive significance, is still a musicological concept and therefore a misleading basis for both mathematical and cognitive approaches. A second problem is that of the numerical methods used by some MIR systems, in particular, distance measures. As will be discussed in Chapter 5, similarity (and by extension distance) is a central part of the auditory domain. MIR systems often use distance measures from mathematics such as Mahalanobis (Mahalanobis, 1936) or Cosine (Cosine, 1997) with little consideration of their perceptual or musical significance.

2.4 Systematicity

2.4.1 The Musical Surface

Having cautioned against a purely musicological approach, ? , pp. 481 proposes a compromise: to adopt a bottom-up approach to music theory, exploring the concepts through systematic mid-level representations. He states that “methods starting at, for example, the musical surface of notes is a useful way of proceeding” The concept of musical surface is illustrated by ? , pp. 159 with a metaphor: “...to approach a musical landscape not by drawing a map, which necessarily confines itself to a limited set of structurally important features, but by presenting a bird’s-eye view of the musical surface – an aerial photograph, as it were, which details the position of every pitched component.”

2.4.2 Systematic Description

? also advocates this mid-level approach, observing that surface description influences analytical observation and that, for an unbiased view, the researcher must be provided with the adequate raw materials with which to make more in-depth observation. Such a systematic, descriptive model would be fundamentally independent of high level concepts such as chords and key but, at the same time, capable of capturing them. ? also discusses the importance of systematicity in terms of dimensionality, ambiguity and timing. He finds that models based on the major-minor paradigm are incapable of adequately describing tonal ambiguity even in some Western music (?, chap. 3).

With a systematic description of the musical surface, theories and models from different domains can be gathered and evaluated together in the same analytical arena, thus helping to bridge the gap between traditional musicology, cognitive psychology and MIR.

3 Pitch-Class Set Theory

One such method available for systematic description of the musical surface is Pitch class set theory. PC-set theory is a system for analysing the pitch content of music. It uses class equivalence relations to reduce the amount of data required to describe any sequence of pitches. This chapter will outline the basic principles.

3.1 Pitch Class

Pitch-class set theory uses octave equivalence. In Western equal temperament (TET), a pitch-class (PC) is an integer representing the residue class modulo 12 of a pitch (Babbitt 1955) and indicates the position of a note within the octave. A PC-set is a collection of PCs ignoring any repetitions and the order in which they occur. PC-sets are notated as follows $\{0,1,2,3,4\}$ with PCs ordered from lowest to highest as a convention (Example 1). The cardinality of a set, denoted $\#S$, is the number of PCs it contains (Example 2). There are 4096 (2^{12}) unique PC-sets with which any segment of music can be represented.

Table 1: Notes and corresponding pitch-classes

| | | | | | | | | | | | | |
|------|---|----|---|----|---|---|----|---|----|---|----|----|
| Note | C | C# | D | D# | E | F | F# | G | G# | A | A# | B |
| PC | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 |

| | | | |
|------------|-------------|-----------|------------------------------------|
| Example 1: | PC-set | Pitch-set | $S = \{A4, C5, E5, A5\}$ (A minor) |
| | | PC-set | $S = \{9, 0, 4, 9\} = \{0, 4, 9\}$ |
| Example 2: | Cardinality | | $\#S = 3$ |

3.2 Set Classification

Defining equivalence classes of PC-sets further reduces the total number of tonal objects. A set-class (SC) is a group of PC-sets related by a transformation or group of transformations. The two types of transformation

commonly used are transposition and inversion. A transposition, $T_n(S)$, transposes the set, S , by the interval, n , (by adding n to all PCs, Example 3). An inversion, $I(S)$, inverts the set S , replacing all PCs with their inverse (12-PC, Example 4). From these two transformations it is possible to define three types of SC: T_n , T_nI and I .

Example 3: Transposition $S = \{0,4,9\}$, $T_3(S) = \{3,7,0\} = \{0,3,7\}$
Example 4: Inversion $S = \{0,4,9\}$, $I(S) = \{11,7,2\} = \{2,7,11\}$

T_n All PC-sets that can be transformed to each by transposition belong to the same class. There are 348 distinct T_n types.
 I All PC-sets that can be transformed to each other by inversion belong to the same SC. There are 197 distinct I types.
 T_nI All PC-sets that can be transformed to each other by transposition, inversion or both belong to the same SC. There are 220 distinct T_nI types.

Transpositional (T_n): All PC-sets that can be transformed to each by transposition belong to the same class. There are 348 distinct T_n types.
Inversional (I): All PC-sets that can be transformed to each other by inversion belong to the same SC. There are 197 distinct I types.
Transpositional/Inversional (T_nI): All PC-sets that can be transformed to each other by transposition, inversion or both belong to the same SC. There are 220 distinct T_nI types.

The Prime Form of a PC-set is a convention for denoting the SC it belongs to. The convention was introduced by Allan Forte (?) for T_nI types and has since been adopted by the majority of theorists. In addition, he devised a system for ordering T_nI -type SCs and assigning to each one a cardinality-ordinal number. For example, the Forte number 3-11 refers to the 11th SC of cardinality 3. This convention has been modified for use with T_n types by adding A and B to the names of inversionally related SCs.

One additional concept is that of cardinality-class (nC), which refers to all the SCs of cardinality n . Cardinality-class 2 is commonly referred to as interval-class (IC) and there are 6 distinct interval-classes.

Table 2: Forte's Prime form and numbering convention

| | |
|-----------------------|-------------|
| PC-set | $\{1,4,9\}$ |
| Prime Form (T_nI) | $\{0,3,7\}$ |
| Prime Form (T_n) | $\{0,4,7\}$ |
| Forte Name (T_nI) | 3-11 |
| Forte Name (T_n) | 3-11B |

3.3 Vector Analysis

3.3.1 Membership and Inclusion

Two concepts that are crucial in PC-set theory are membership and inclusion. Membership of a set is denoted $p \in S$ and means that PC p is a member of set S (Example 5). Inclusion in a set is denoted $Q \subset S$ and means that all members of set Q are also members of set S (Example 6). Q is said to be a subset of S .

Example 5: Membership $4 \in \{0,4,9\}$
Example 6: Inclusion $\{0,4,9\} \subset \{0,1,4,5,9\}$

3.3.4 Notation

Some additional vector notation is required for the comparison procedures of the similarity measures described in 3.4.

Difference Vector is the absolute difference between corresponding terms in the nCVs of two SCs, X and Y:

$$DV(nCV(X), nCV(Y)) = |nCV(X) - nCV(Y)|$$

Vector Magnitude is the length of the nCV in euclidean space:

$$\|nCV(X)\| = \sqrt{\sum_{i=1}^{\#nCV} (nCV(X)_i)^2}$$

Unit Vector is the normalised nCV (unit length):

$$nCV\hat{V}(X) = \frac{nCV(X)}{\|nCV(X)\|}$$

Euclidean Distance is the distance between the points defined by two nCVs in n-dimensional Euclidean space:

$$d(X, Y) = \sqrt{\sum_{i=1}^n (X_i - Y_i)^2} = \|DV(X, Y)\|$$

3.4 Set-Class Similarity

3.4.1 Similarity Relations

The assessment of similarity between two SCs has been discussed in the literature for decades and a large number theoretical models have been proposed. Different models approach the problem from different conceptual standpoints and theorists have different opinions about the contributing factors. All these models are described under the blanket term “similarity relations”. Despite the perennial fascination with the concept, little or no consensus exists as to what constitutes a good similarity relation.

? provides a comprehensive and in-depth review of a large number of similarity relations and categorises them according to some fundamental principles. Firstly, he distinguishes between methods that produce binary outcomes and those that produce a range of values. The former category, termed “plain relations”, include Forte’s R-relations (?) and indicate whether the two SCs are related in a specific way, which in turn may give some indication of whether they are similar. The latter category, termed “similarity measures”, indicate a degree of similarity, returning a value from a known range. This property appears to be more inline with the perceptual notion of similarity and therefore the focus of this work shall be exclusively on similarity measures.

3.4.2 Similarity Measures

The vast number and diversity of the different approaches to similarity measures can only be approached by narrowing the focus to a specific type. Here we will focus on measures that use the Tn and TnI-type SCs (3.2), and furthermore we will only consider those methods based on vector analysis (3.3). These measure usually involve the comparison of the SC’s nCVs. Of this (still sizeable) subset, ? identifies two main categories.

| | |
|-----------------|--|
| Single nC: | Single nC measures compare the nCVs of the two SCs for one particular value of n. Many of the relations in this category compare ICVs (2CVs). |
| Total Measures: | Total Measures consider the subsets of all cardinalities contained within in two SCs. All the relevant nCVs are compared to produce a final value. |

Table 4 shows the majority of the Tn and TnI-Type, vector based similarity measures from the PC-set theoretical literature. Vector Type indicates whether the measure compares ICVs or nCVs. Card (Cardinality) indicates whether the measure is capable of comparing SCs of different cardinalities while the Measure Type indicates which of Castren’s categories it belongs to. nC indicates it is a Single nC measure and TOTAL indicates it is a Total Measure. All these measure are described more thoroughly in A.

Table 5: Comparison table of similarity measures

| THEORIST | SIMILARITY MEASURE | VECTOR TYPE | CARD | MEASURE TYPE |
|------------|--------------------|-------------|------|--------------|
| MORRIS | K | ICV | SAME | nC |
| | SIM | ICV | SAME | nC |
| | ASIM | ICV | ANY | nC |
| LORD | sf | ICV | SAME | nC |
| TEITELBAUM | s.i. | ICV | SAME | nC |
| ROGERS | IcVD1 | ICV | ANY | nC |
| | IcVD2 | ICV | ANY | nC |
| | COS | ICV | ANY | nC |
| ISAACSON | AMEMB2 | ICV | ANY | nC |
| | IcVSIM | ICV | ANY | nC |
| | ISIM2 | ICV | ANY | nC |
| | ANGLE | ICV | ANY | nC |
| RAHN | AK | ICV | ANY | nC |
| | MEMB _n | nCV | ANY | nC |
| | TMEMB | nCV | ANY | TOTAL |
| | ATMEMB | nCV | ANY | TOTAL |
| LEWIN | REL2 | ICV | ANY | nC |
| | REL | nCV | ANY | TOTAL |
| CASTREN | %REL _n | nC%V | ANY | nC |
| | T%REL | nC%V | ANY | TOTAL |
| | RECREL | nC%V | ANY | TOTAL |
| BUCHLER | SATSIM | nSATV | ANY | nC |
| | CSATSIM | CSATV | ANY | nC |
| | TSATSIM | nSATV | ANY | TOTAL |
| | AvgSATSIM | nSATV | ANY | TOTAL |

3.4.3 Castren’s Criteria

In addition to his categorisation, ? proposes several criteria which a good similarity relation should meet. Later, these criteria will be used in assessing the specific capabilities of various similarity measures.

Castren says that a similarity measure should:

- C1: allow comparisons between SCs of different cardinalities
- C2: provide a distinct value for every pair of SCs
- C3: provide a comprehensible scale of values such that
 - C3.1: All values are commensurable
 - C3.2: the end points are not just some extreme values but can be meaningfully associated with maximal and minimal similarity.
 - C3.3: The values are integers or other easily manageable numbers
 - C3.4: the degree of discrimination is not too coarse and not unrealistically fine
- C4: produce a uniform value for all comparable cases
- C5: observe mutually embeddable subset-classes of all meaningful cardinalities

- C6: observe also the mutual embeddable subset-classes not in common between the SCs being compared.

3.5 Perceptual Relevance

The many equivalence relations used in PC-set theory give rise to a highly abstract description of musical objects. Thus, an important question to be asked is whether these theoretical assumptions and models of similarity reflect perceptual equivalence. This chapter contains a summary and discussion of some relevant studies.

3.5.1 Octave Equivalence

Pitch is a percept that derives from a particular harmonic structure and is roughly proportional to the logarithm of the fundamental frequency. This allows pitch to be modelled as a straight line. Music psychologists have observed a strong perceptual similarity between pitches with fundamental frequencies in the ratio of 2:1. This property of octave similarity leads the straight line model of pitch to be bent into a helix. Division of the octave into a number of categories is thought to offer a more efficient cognitive representation in memory and thus confers evolutionary advantage. The resulting pitch equivalence classes are implicitly learned through exposure at an early age. TET has 12 pitch equivalence classes which, in PC-set theory, are modelled as a circular projection of the pitch helix. Thus the two most fundamental components of PC-set theory, i.e. octave equivalence and pitch-class labelling, would appear to have a solid basis in perception.

? investigated the perceived similarity of pairs of chords with varying numbers of octave related pitches. He found that in general chords with identical PC contents were perceived as more similar than chords with near identical PC contents, regardless of the octave. However, in further studies he his findings suggest that there are other factors that play a significant role (?).

3.5.2 Set-Class Equivalence

Some researchers have attempted to examine whether there is perceived equivalence between different manifestations of a PC-set. ? presented subjects with sequences of tones derived by transforming two different PC-sets. They noted that subjects were able to distinguish between the different sets both in neutral and musical contexts.

? investigated the perceptual similarity of different PC-sets derived from the same set class under TnI classification. Subjects were presented with three-note melodies and asked to judge which was equivalent to a reference melody. Some melodies preserved the SC identity whilst others did not. She found transpositions to be perceived more similar than inversions and in addition she discovered that the order of the notes and melodic contour was a strong factor in perceived similarity.

Some authors have questioned the perceptual relevance of using TnI equivalence as a basis for set classification. ? seems unconvinced by evidence for the perceptual similarity of inverted intervals. This can be illustrated by the example of major and minor triads which, while perceptually distinct, are equivalent under TnI.

3.5.3 Perceived vs Theoretical Similarity

A number of studies have been done to ascertain the connection between perceptual similarity ratings and the theoretical values obtained from some PC-set similarity measures. A large number of relevant studies are summarised by ? and the most significant ones are mentioned here.

? used multidimensional scaling on subjects' similarity ratings between trichords and tetrachords and on the similarity values obtained from SIM. She compared the 2-dimensional solutions and found there to be little correlation.

? investigated non-traditional chords. He compared subjects' ratings with similarity assessments calculated from Forte's R-relations and Lord's similarity function. He also concluded there was little correspondence between the two.

? compared subjects' ratings of 4 note melodies with the theoretical values obtained from SIM. She found the ratings of subjects with more musical training to be more correlated with the SIM values.

? compared subjects' ratings of pitch sequences with corresponding values of seven ICV-based similarity measures: ASIM, MEMB2, REL2, s.i., IcVSIM and AMEMB2 and concluded there to be a strong relation.

? compared subjects' ratings of pentachords with the values obtained from 9 similarity measures. He found there to be a connection between aurally estimated ratings and the theoretical values and concluded that the abstract properties of set-classes do have some perceptual relevance. He also comments on the way in which this kind of study is conducted, suggesting that the way in which subjects are presented with the stimuli has a significant effect on the outcome.

3.6 PC-set Theory for Analysis

PC-set theory as means for descriptive modelling of tonality is not widely known outside of highly theoretical circles and the use of PC-set similarity measures seems mainly restricted to the theorists who proposed them (for example, ?). The basic premise is simple: a musical piece is segmented and each segment described by its SC. Similarity measures can be used to assess the similarity between segments or between a segment and some reference SC.

? used a pentachordal tail segmentation policy (each successive note defines a segment that includes the preceding four notes) and compared these segments to comparison sets 7-1 (chromaticism) and 7-35 (diatonicism) using the REL distance (A.7.1). They claim that the visual results of their analysis "reflect pertinent aspects of our listening experience" (?, pp. 204).

?, chap. 5.3 uses a more systematic approach to segmentation using multiple time scales. He proposes the class-scape, a two-dimensional visualisation of a piece of music with time on the x-axis and segmentation time-scale on the y-axis. A single SC can be represented by highlighting the segment or alternatively each segment can be shaded according to its REL distance from a comparison SC. He emphasises that class-scape is an exploratory tool rather than an automated analysis system.

4 Multidimensional Scaling

Multidimensional scaling (MDS) is a numerical visualisation technique that, given a matrix of pairwise distances between objects, provides a geometric configuration of the objects in some abstract space. It provides an efficient means of observing relationships in large, complex data sets and the resulting dimensions often give valuable insight into the data as a whole.

4.1 Non-Metric MDS

Non-Metric MDS was described by ? and it assumes that the distance matrix values are related to points in an abstract N-dimensional Euclidean space. An important consideration is that of the dimensionality of the solution. For comprehension and visualisation it is important to minimise the number of dimensions however, there is a trade-off between the number of dimensions and the accuracy of the model. For a given dimensionality, we obtain two values: Stress and r^2 .

| | |
|--------|---|
| Stress | Stress is a "goodness of fit" measure which characterises the distortion that occurs in a given number of dimensions. |
| | As the number of dimensions increases the stress decreases. |
| r^2 | r^2 is the percentage variability of the data being explained by the solution. |

By plotting stress against r^2 for a number of dimensionalities is possible to observe the point at which additional dimensions do not significantly improve the solution (the "elbow"). Ultimately, the choice of dimensions should be based on interpretation.

4.2 Cluster Analysis

Cluster analysis (CA) is method for dealing with dimensions that are highly separable. First, the most similar pair of objects are selected and grouped together in a cluster. The process is repeated, creating a

binary tree structure. The distance between objects is then related to their separation along the branches of the tree.

4.3 MDS with Similarity Measures

Using MDS on the values produced by similarity measures is one way to approach an understanding of the constructs they are measuring. There are two potentially interesting issues to consider. Firstly, a measure may be inconsistent with itself, meaning that the geometries it produces are not “robust” (changing the set of objects changes the distances between the original set). This kind of problem cannot be observed through inspection of the values alone. The second issue is that two different measures that are both self-consistent may produce very different geometries from the same group of SCs. The question then is, what exactly do the measures measure?

5 Spatial Models of Tonality

5.1 Similarity and Distance

Judgements of similarity form the basis of many cognitive processes including the perception of tonality. Similarity between two objects is often conceived as being inversely related to distance between them in geometric space. For example, some tonal objects (chords, for example) are perceived as close to one another whereas others are further apart. In addition, the number of dimensions of the geometric space is in connection with the number of independent properties that are relevant for similarity comparisons. ? suggests that humans are naturally predisposed to create spatial cognitive representations of perceptual stimuli due to the geometric nature of the world we have evolved to inhabit. Therefore spatial modelling of tonality, as well as helping to visualise the complex multidimensional relationships between tonal phenomena, has the potential to reflect cognitive aspects of the way they are perceived.

5.2 Spatial Representations

Throughout history theorists have proposed many spatial representations of tonality from different domains. From the graphemic domain, ? and ? both proposed simple 2-dimensional charts to display the proximity between keys. For representation of chords, ? models major and minor triads as regions in a 2-dimensional space whilst ? proposes a variety high dimensional, non-euclidean chord spaces that reflect the theoretical principles of voice leading. From the acoustic domain, ? proposes a five-dimensional model to represent interval relations between pitches. Some theorists have attempted to incorporate relations between several levels of tonal hierarchy into one configuration. The “spiral array” of ? is a three-dimensional mathematical model which simultaneously captures the relations between pitches, chords and keys. The “chordal-regional space” of ? models the relations between chords within a certain key.

5.3 Cognitive Psychology

The auditory domain has been addressed through cognitive psychology by ? who used the probe-tone methodology (?) to establish major and minor key profiles (12-dimensional vectors containing the perceptual stability ratings of each of the 12 pitch classes within a major or minor context). These profiles, known as Krumhansl-Kessler profiles (KK-profiles), show the hierarchy of pitches in major and minor keys. Correlating each of the 24 major and minor profiles produced a matrix of pairwise distances which was fed to a dimensional scaling algorithm. The resulting geometrical solution was found to have a double circular property (circle of fifths and relative-parallel relations) which can be modelled as the surface of a 3D torus. Many spatial models of tonality have this double circular property whether it is implicit (??) or stated explicitly (?).

5.4 Set-Class Spaces

Most of these models are limited to description of music in the major-minor paradigm and are not capable of generalising beyond the “western common practice”. PC-set theory, once again, provides a possible means

to generalise to any kind of pitch-based music. By considering a collection of tonal objects described by SCs, a geometric space can be constructed to model their relations based on some theoretical principle. Some PC-set theorists have proposed explicit geometric spaces to model relations between SCs. The distances in these spaces are expressed by models of similarity based on voice leading (??) or ICVs and the Fourier transform (??). However, these models are only designed to represent SCs of one cardinality-class at a time and cannot model the relations between arbitrary collections of pitches.

Alternative spatial models are provided by the implicit geometries of the values produced by the SC similarity measures discussed in ???. As mentioned in 4.3, MDS can be used on values produced by similarity measure to create a geometric space. ? and ? both applied MDS to the values produced from a variety of similarity measures. Samplaski used TnI-type SCs while Kuusi used Tn-type. They both found reasonably low-dimensional solutions and attempted to interpret each of the dimensions. Kuusi interpreted three dimensions as corresponding to chromaticism, wholetoneness and pentatonicism. Samplaski made similar observations but found some dimensions in the higher-dimensional spaces difficult to interpret. Nevertheless, he concluded that values from similarity measure tend to agree (with some exceptions) and that they measure constructs relating to familiar scales (diatonic, hexatonic, octatonic, etc.).

6 Similarity Measure Selection

So far, PC-set theory has been presented as viable means for systematic descriptive modelling of tonality and brief reference has been made to the extensive existing literature on SC similarity measures (see 3.4). In this section, the large number of measures will be discussed in relation to Castren’s criteria (see 3.4.3) in order to gauge their suitability for use in systematic surface description models. The most suitable measures will be subjected to a more thorough analysis over the course of the work.

6.1 Criteria

Castren’s criteria (see 3.4.3) for similarity measures provide a basis for assessment of similarity measures for our purposes. A detailed descriptions and justification for the criteria can be found in ?, chap. 2, however here we will focus on one or two specific aspects. The table below shows the list of similarity measures with marks indicating whether each of the criteria is met. In sections 6.2 to 6.4 specific criteria are used to exclude measures from further consideration with justification in terms of systematicity and perceptual relevance.

6.2 Cardinality

Measures which fail to meet criteria C1, i.e. that cannot compare SCs of different cardinalities, are clearly inadequate for systematic analysis of music, which might require the comparison of any two arbitrary segments regardless of how many PCs they contain. Both s.i. (A.3.1) and sf (A.2.1) were proposed specifically for SCs of the same cardinality and so will be excluded from further discussion. Some other measures which were intended to compare SCs of different cardinalities nonetheless have problems. Measures such as SIM (A.1.2) and K (A.1.1) give unintuitive values when the cardinalities of the SCs being compared differ greatly and, in addition, the range of values produced depends on the cardinality of the sets (failure to meet criteria C3.1). Measures of this type will also be excluded.

6.3 SC-type

An important consideration when using similarity measures is the type of SC being compared. Many of the measures are designed for comparison of TnI-type SCs, however, owing to issues raised in 3.5 regarding the perceptual relevance of inversionally related sets, here, measures will be selected for use with Tn-type SCs. This means that the measure should be able to discriminate between inversionally related sets. All the measures which exclusively consider interval content in the comparison procedure can automatically be discounted, as inversionally related sets have identical ICVs.

Table 6: Castren’s Criteria

| SIMILARITY MEASURE | C1 | C2 | C3.1 | C3.2 | C3.3 | C3.4 | C4 | C5 | C6 |
|-----------------------|----|----|------|------|------|------|----|----|----|
| s.i. | | | | | X | X | | | |
| sf | | | | | X | X | X | | |
| IcVSIM | X | X | | | | X | | | |
| ISIM2 | X | X | | | | X | | | |
| K | X | X | | | X | X | X | | |
| SIM | X | X | | | X | X | X | | |
| MEMB _n | X | X | | | X | X | X | | |
| AMEMB2 | X | X | X | | | | | | |
| ASIM | X | X | X | X | | X | X | | |
| IcVD1 | X | X | X | X | | X | X | | |
| IcVD2 | X | X | X | X | | X | | | |
| COS | X | X | X | X | | X | | | |
| ANGLE | X | X | X | X | | X | | | |
| AK | X | X | X | X | | X | X | | |
| SATSIM | X | X | X | | | | | | |
| CSATSIM | X | X | X | | | | | | |
| REL2 | | | | | | | | | |
| %REL _n | X | X | X | X | X | X | X | | |
| TMEMB | X | X | | | X | | X | X | |
| ATMEMB | X | X | X | X | | X | X | X | |
| TSATSIM | X | X | X | X | | X | | X | |
| AvgSATSIM | X | X | X | X | | X | | X | |
| REL | X | X | X | X | | X | X | X | |
| T%REL | X | X | X | X | X | X | X | X | |
| RECREL | X | X | X | X | X | X | X | X | X |

6.4 Measure Type

Although many theorists have supposed that interval-class subsets are of paramount importance in similarity judgments, no thorough investigation has been carried out as to the exact perceptual significance of subset cardinality. Single nC measures presuppose that subsets of one particular cardinality contribute to similarity above all others. In the interest of systematicity, we will not make this assumption instead assuming that subsets of all cardinalities are equally relevant and should be considered. Similarity measures that exhaustively consider all subset cardinalities meet criteria C5 and are total measures (see 3.4.2). The six total measures from 3.4.2 will be investigated further.

6.5 Total Measures

In this section each of the six total measures are examined in more detail and assessed in terms of their particular abilities and discriminatory power. Section ?? describes the concept of trivial forms and for each measure the method for handling these cases is described.

6.5.1 Trivial Forms

Three of the 351 Tn -type SCs are known as trivial forms: 1-1, 11-1 and 12-1. Due to their lack of musical or harmonic significance, these SCs are usually excluded from the work of SC-theorists. However, it is important that they be included in any systematic description and that their similarity to other sets be given a meaningful value. The next chapter will discuss each of the total measures and specify how each of the trivial forms is to be dealt with if it was not made explicit by the theorist.

The total measures which will be discussed make comparisons based on the subset content of a set. SC 1-1, which has no subsets, is rarely accounted for in such measures and in these cases a simple method will be used: Comparisons involving $X = 1-1$ and Y will be given the value $1/\#Y$. Thus, the value will be the ratios of the cardinalities with 1 indicating maximum similarity.

Table 7: Trivial Forms

| | |
|------|---------------------------------|
| 1-1 | $\{0\}$ |
| 11-1 | $\{0,1,2,3,4,5,6,7,8,9,10\}$ |
| 12-1 | $\{0,1,2,3,4,5,6,7,8,9,10,11\}$ |

6.5.2 Rahn: ATMEMB

Details on how to calculate ATMEMB are give in A.5.4. In his analysis of the measure, Castren concludes that “divisor term is flawed, resulting in values suggesting suspiciously high degrees of dissimilarity between SCs of clearly different cardinalities. The general reliability and usefulness of the measure is difficulty to determine” (?, pp. 89). The trivial forms 11-1 and 12-1 are accommodated explicitly by the fomulation of ?, however SC 1-1 is not and thus values will be obtained using the method specified in 6.5.1.

6.5.3 Lewin: REL

Details on how to calculate REL are given in A.7.1. From the basic equation it is possible to define three different formulations depending on the exact nature of $SUB(X)$. In each formulation the trivial forms 11-1 and 12-1 are accomodated. The three formulations are as follows:

1. $SUB(X)$ consists of the concatenated nCVs from 2 to 12. Here comparisons involving SC 1-1 will be evaluated with the method specified in 6.5.1.
2. $SUB(X)$ consists of the concatenated nCVs from 1 to 12 ($1CV(X) = \#X$). This formulation accommodates SC 1-1.
3. ? specifies an alternative formulation where $SUB(X)$ begins with the ICV (2CV) followed by the concatenated nCVs from 1 to 12. This formulation accommodates SC 1-1.

6.5.4 Buchler: AvgSATSIM and TSATSIM

Details on how to calculate AvgSATSIM and TSATSIM are given in A.9.3 and A.9.4 respectively. Comparisons involving SC 1-1 are not accommodated and thus the method specified in 6.5.1 will be used to provide values. Comparisons involving SCs 11-1 and 12-1 are accommodated except for the single comparison that involves both. This is because their $MAX_n(\#X)$ and $MIN_n(\#X)$ vectors are equal and thus all terms of the nSATVs are 0. The value for this comparison will be set to 0 (indicating maximal similarity). For comparisons involving ICs the value will be given by $SATSIM_2(X,Y)$ (see A.9.2).

6.5.5 Castren: T%REL and RECREL

Details on how to calculate T%REL and RECREL are given in A.8.4 and A.8.5 respectively. Comparisons involving SCs 11-1 and 12-1 are accommodated in both by Castren’s formulation. Comparisons involving SC 1-1 will be given values by the method specified in 6.5.1. Castren comments that some T%REL values are too high to be intuitively plausible. Finally it should be noted that the basic algorithm provided by Castren for calculating RECREL is not feasible for large sets. Compaisons of such sets require tables of pre-computed branch values.

6.5.6 Scale of Values

The values of each measure will be adjusted to the same scale for comparability by the same method as ?, pp. 48). This scale is from 0 to 100 with with 0 indicating maximum similarity. The modified values will be signalled by adding the symbol “prime” to the name.

- $\text{ATMEMB-prime}(X,Y) = (1-\text{ATMEMB}(X,Y))*100$
- $\text{REL-prime}(X,Y) = (1-\text{REL}(X,Y))*100$
- $\text{AvgSATSIM-prime}(X,Y) = \text{AvgSATSIM}(X,Y)*100$
- $\text{TSATSIM-prime}(X,Y) = \text{TSATSIM}(X,Y)*100$
- $\text{T\%REL-prime}(X,Y) = \text{T\%REL}(X,Y)$
- $\text{RECREL-prime}(X,Y) = \text{RECREL}(X,Y)$

7 Practical Analysis

7.1 Chords as PC-Sets

So far in the previous chapters, PC-set theory and the similarity measures have been discussed with only the broadest reference to familiar musical or musicological concepts. In order to use these techniques for real music description and analysis some effort should be made to link SC theoretical concepts to musicological ones.

The table below shows a collection of familiar chord types, cadence types and scale types with their corresponding Tn-type prime-form SC, Forte Name and index number (the position in an ordered list of all 351 Tn-type SCs). The table is divided into five vertical segments: three-note chords, four-note chords, five-note chords, cadences and scales.

In PC-set theory it is only the chord type that is relevant seeing as transpositionally related (Tn-type) SCs are considered equivalent. For example all major chords are considered equivalent regardless of the root.

7.2 Repertoire

In addition it is also useful to understand the tonal or chordal content of some specific music. This section details existing analysis and annotations of the tonal content of some music collections.

7.2.1 Billboard Chords

The McGill Billboard data set constitutes an expert ground truth for chord recognition algorithms in MIR. It contains detailed transcriptions of the chords for more than one thousand songs selected at random from Billboard magazine's "Hot 100" charts. ?, pp. 637 provides the frequency distribution of the 50 most common chord types in the data set. Major chords alone account for more than half of the data set, followed by minor chords and the basic 7th chords. No information is given as to the scales or modes used.

7.2.2 Beatles Chords

?, chap. 6 provides a collection chord transcriptions for 180 Beatles songs and statistical analysis of the collection. Again, over 70% by both occurrence and time are major and minor chords.

7.2.3 Diatonism in Bach and Victoria

A Matorell. MIDI format

- Bach WTC
 - Major mode preludes
- Victoria
 - Ionian mode

7.2.4 Josquin?

-

Table 8: Chord types and their SCs

| Chord | Tn-Type SC (Prime Form) | Forte Name (Tn-type) | index |
|------------|----------------------------|-------------------------|-------|
| maj | {0,4,7} | 3-11B | 25 |
| min | {0,3,7} | 3-11A | 24 |
| dim | {0,3,6} | 3-10 | 23 |
| aug | {0,4,8} | 3-12 | 26 |
| sus4 | {0,2,7} | 3-9 | 22 |
| sus2 | {0,2,7} | 3-9 | 22 |
| maj7 | {0,1,5,8} | 4-20 | 57 |
| min7 | {0,3,5,8} | 4-26 | 64 |
| hdim7 | {0,2,5,8} | 4-27A | 65 |
| 7 | {0,3,6,8} | 4-27B | 66 |
| dim7 | {0,3,6,9} | 4-28 | 67 |
| min(7) | {0,1,4,8} | 4-19A | 55 |
| aug(7) | {0,3,4,8} | 4-19B | 56 |
| maj(9) | {0,2,4,7} | 4-22A | 59 |
| min(9) | {0,2,3,7} | 4-14A | 46 |
| maj6 | {0,3,5,8} | 4-26 | 64 |
| min6 | {0,1,5,8} | 4-20 | 57 |
| sus4(7) | {0,2,6,7} | 4-16B | 51 |
| sus4(b7) | {0,2,5,7} | 4-23 | 61 |
| 9 | {0,2,4,6,9} | 5-34 | 129 |
| maj9 | {0,1,3,5,8} | 5-27A | 116 |
| min9 | {0,3,5,7,8} | 5-27B | 117 |
| V-I/IV-I | {0,1,3,5,8} | 5-27A | 116 |
| V7-I | {0,1,3,5,6,8} | 6-Z25A | 176 |
| V-IV | {0,2,4,6,7,9} | 6-33B | 189 |
| Pentatonic | {0,2,4,7,9} | 5-35 | 130 |
| Wholetone | {0,2,4,6,8,10} | 6-35 | 192 |
| Diatonic | {0,1,3,5,6,8,10} | 7-35 | 276 |
| Octatonic | {0,1,3,4,6,7,9,10} | 8-28 | 322 |

7.2.5 Essen Collection

- 10000 folk songs
- MIDI format

7.3 Comparison of Chords

For a preliminary idea of the utility of the total measures it is useful to visualise the vlaues produced for comparisons involving the common tonal objects described in 7.1. This information can be visualised as 2D grids with each square corresponding to the comparison between two tonal objects and coloured according the distance between them i.e. the value of MEASURE-prime (see 6.5.6).

%#+CAPTION: ATMEMB distance between chord types

- Six such plots are shown below, one for each total measure.
- What can be seen?
- which measures discriminate what?
- eliminate some measures?
- Comparison - plot difference between plots
- which measures disagree and agree most?

- To what extent do the measures agree?
- correlate plots - plot overall correlation
- discard TSATSIM
- which rel?

7.4 Conclusions

- not suitable for common chord recognition or differentiation
- segmentation?!
- subclass vectors

7.5 Measure descrimination

- which measures best descriminate between different SC dictionaries?
- what sets could be best for comparison sets in class scapes?
- compare/look at graphs from ??

7.6 Structure

- autocorrelation of class scape?
- cadential sets mark ends of sections

8 Class Scape

- from martorell
- what?
- segementation
- class scape
 - 3d sparse matrix
- class matrix
 - remove timescalee information
- class vector
 - %time of class matrix

8.1 replicate

- list his examples
- replicate with different measures

8.2 genere specific analysis

- visualise examples of corpora using class-scape

8.3 structure analysis

- structure analysis with autocorrelation and cadential sets
- self similarity matrix with the user able to vary the segment size.

9 MDS

- Visualisation of values using MDS
- r2 against stress for different measures
- comparison
- visualisation

10 Latex Stuff

A SC Similarity Measures

This chapter contains a concise summary of the SC similarity measures from the literature organised by theorist. Each section specifies the publication in which the measure was proposed and brief description of the theoretical approach adopted by the theorist. A mathematical formula is given where possible using standard notation. A reference for notation can be found in 3.3.4 and commonly used symbols are defined in the glossary. Where a mathematical formula does suffice, the comparison procedure is described in words. In addition, each section contains a table specifying important statistics:

- SC-Type: the type of SC the measure compares (Tn or TnI)
- Cardinality: whether the measure can compare SCs of different cardinalities.
- Vector Type: the type of vector used in the comparison procedure (see 3.3).
- Max Similarity: the value that indicates maximum similarity from all possible comparisons.
- Min Similarity: the value that indicates minimum similarity from all possible comparisons.
- Average Value: the average value from all possible comparisons.
- No. Values: the number of distinct values produced from all possible comparisons.
- Criteria Met: a list of Castren's criteria which the measure meets.
- I-related: whether the measure discriminates between inversionally related sets.
- Z-related: whether the measure discriminates between Z-related sets.

A.1 MORRIS

A.1.1 K

Presented in ?, pp. 448, the K measure gives the number of intervals-classes (dyad-classes) shared by two SCs, X and Y.

$$K(X, Y) = \sum_{i=1}^6 MIN(x_i, y_i)$$

| | |
|-----------------|--------------------|
| SC Type: | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | 55 |
| Min Similarity: | 0 |
| Average Value: | 10 |
| No. Values: | 35 |
| Criteria Met: | C1,C2,C3.3,C3.4,C4 |
| I-related: | No |
| Z-related: | No |

- Problems: scale of values not the same for all value groups.

A.1.2 SIM

Presented in ?, pp. 446, SIM compares the ICVs of two SCs (the value is the cardinality of the DV).

$$SIM(X, Y) = \sum_{i=1}^6 |x_i - y_i|$$

or

$$SIM(X, Y) = \#DV(ICV(X), ICV(Y))$$

SIM is a function of K:

$$SIM(X, Y) = \#ICV(X) + \#ICV(Y) - 2.K(X, Y)$$

| | |
|-----------------|--------------------|
| SC Type: | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | 0 |
| Min Similarity: | 65 |
| Average Value: | 13 |
| No. Values: | 44 |
| Criteria Met: | C1,C2,C3.3,C3.4,C4 |
| I-related: | No |
| Z-related: | No |

- Problems: scale not the same for all value groups. course resolution when cardinalities differ greatly

A.1.3 ASIM

Presented in ?, pp. 450, ASIM (Absolute SIM) is a scaled version of SIM to address criteria C3.1.

$$ASIM(X, Y) = \frac{SIM(X, Y)}{\#ICV(X) + \#ICV(Y)}$$

| | |
|-----------------|-------------------------|
| SC Type: | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | 0 |
| Min Similarity: | 1 |
| Average Value: | 0.42 |
| No. Values: | 79 |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.4,C4 |
| I-related: | No |
| Z-related: | No |

Problems: Fixed the scale of values, but still coarse resolution when cardinalities differ greatly. Scaling is done as the last step.

A.2 LORD

A.2.1 sf

Presented in (?, pp. 93), sf (Similarity Function) is similar to SIM but developed independently. sf is a subset of SIM:

$$sf(X, Y) = \frac{\#DV(ICV(X), ICV(Y))}{2} = \frac{SIM(X, Y)}{2}$$

| | |
|-----------------|--------------|
| SC Type: | TnI |
| Cardinality: | Same |
| Vector Type: | ICV |
| Max Similarity: | 0 |
| Min Similarity: | 9 |
| Average Value: | 3 |
| No. Values: | 10 |
| Criteria Met: | C3.3,C3.4,C4 |
| I-related: | No |
| Z-related: | No |

A.3 TEITELBAUM

A.3.1 s.i.

Presented in ?, pp. 88, s.i. (Similarity Index) is the Euclidean distance between the cartesian coordinates defined by the ICVs of two SCs. This is equivalent to the magnitude of the difference vector.

$$s.i.(X, Y) = \sqrt{\sum_{i=1}^6 (x_i - y_i)^2} = \|DV(ICV(X), ICV(Y))\|$$

| | |
|-----------------|-----------|
| SC Type: | TnI |
| Cardinality: | Same |
| Vector Type: | ICV |
| Max Similarity: | 1.41 |
| Min Similarity: | 8.49 |
| Average Value: | 2.85 |
| No. Values: | 31 |
| Criteria Met: | C3.3,C3.4 |
| I-related: | No |
| Z-related: | No |

- Same cardinality only
- Z-related sets not compared

A.4 ROGERS

A.4.1 IcVD₁

Presented in ?, IcVD₁ (Distance Formula 1) is a modification of SIM (A.1.2). The ICV components are scaled before being summed. IcVD₁ is related to Castren's %REL₂ (A.8.3): %REL₂(X,Y) = IcVD₁(X,Y) × 50.

$$IcVD_1(X, Y) = \#DV \left(\frac{ICV(X)}{\#ICV(X)}, \frac{ICV(Y)}{\#ICV(Y)} \right)$$

| | |
|-----------------|-------------------------|
| SC Type: | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | 0 |
| Min Similarity: | 2 |
| Average Value: | 0.59 |
| No. Values: | 140 |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.4,C4 |
| I-related: | No |
| Z-related: | No |

A.4.2 IcVD₂

Presented in ?, IcVD₂ (Distance Formula 2) is similar to s.i. (A.3.1), but instead returns the Euclidean distance between the ends of the normalised ICVs.

$$IcVD_2(X, Y) = \|DV(IC\hat{V}(X), IC\hat{V}(Y))\|$$

| | |
|-----------------|----------------------|
| SC Type: | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | 0 |
| Min Similarity: | 1.41 |
| Average Value: | 0.54 |
| No. Values: | 133 |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.4 |
| I-related: | No |
| Z-related: | No |

- Problems: does not produce uniform values for comparable cases

A.4.3 Cos(θ)

Presented in ?, $\text{Cos}\theta$, gives the cosine of the angle between the ICVs in six-dimensional euclidean space. As the angle decreases the similarity approaches 1.

$$\text{Cos}\theta(X, Y) = \frac{ICV(X) \cdot ICV(Y)}{\|ICV(X)\| \times \|ICV(Y)\|}$$

| | |
|-----------------|----------------------|
| SC Type: | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | 1 |
| Min Similarity: | 0 |
| Average Value: | 0.81 |
| No. Values: | 92 |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.4 |
| I-related: | No |
| Z-related: | No |

- Problems: C4

A.5 RAHN

A.5.1 AK

Presented in /citet[pp. 489]{Rahn1979}, AK is an absolute or adjusted version of Morris' K (A.1.1), addressing the C3.1 criteria. AK is related to Morris' ASIM: $\text{AK}(X,Y)=1-\text{ASIM}(X,Y)$.

$$\text{AK}(X, Y) = \frac{2K(X, Y)}{\#ICV(X) + \#ICV(Y)}$$

| | |
|-----------------|-------------------------|
| SC Type: | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | 1 |
| Min Similarity: | 0 |
| Average Value: | 0.58 |
| No. Values: | 78 |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.4,C4 |
| I-related: | No |
| Z-related: | No |

- Problems: single scale of values (C4), but poor discrimination for some value groups.

A.5.2 MEMB_n

Presented in ?, pp. 492, MEMB_n (Mutual Embedding Number) compares the nCVs of two SCs for one nC at a time. It measures the mutual embedding of subsets such that only non-zero components of the nCVs contribute. By setting n = 2 (MEMB₂) it compares ICVs.

$$MEMB_n(X, Y) = \sum_{i=1}^{\#nC} nCV(X)_i + nCV(Y)_i$$

such that $nCV(X)_i > 0$ and $nCV(Y)_i > 0$.

| | |
|-----------------|--------------------|
| SC Type: | TnI or Tn |
| Cardinality: | Any |
| Vector Type: | nCV |
| Max Similarity: | 121 |
| Min Similarity: | 0 |
| Average Value: | 30 |
| No. Values: | 79 |
| Criteria Met: | C1,C2,C3.3,C3.4,C4 |
| I-related: | Yes* |
| Z-related: | Yes* |

- Problems: does not produce uniform scale of values for all value groups.

A.5.3 TMEMB

Presented in ?, pp. 492, TMEMB (Total Mutual Embedding Number) counts the mutually embedded subsets of every cardinality. TMEMB is a total measure.

$$TMEMB(X, Y) = \sum_{n=2}^{12} MEMB_n(X, Y)$$

| | |
|-----------------|------------------|
| SC Type: | TnI or Tn |
| Cardinality: | Any |
| Vector Type: | nCV |
| Max Similarity: | 6118 |
| Min Similarity: | 0 |
| Average Value: | 131 |
| No. Values: | 877 |
| Criteria Met: | C1,C2,C3.3,C4,C5 |
| I-related: | Yes |
| Z-related: | Yes |

- Problems: Different value scales for different value groups

A.5.4 ATMEMB

Presented in ?, pp. 494, ATMEMB (Adjusted Total Mutual Embedding Number) is a scaled version of TMEMB to address criteria C3.1 (like SIM and ASIM; A and AK). ATMEMB is a total measure.

$$ATMEMB(X, Y) = \frac{TMEMB(X, Y)}{2^{\#X} + 2^{\#Y} - (\#X + \#Y + 2)}$$

| | |
|-----------------|----------------------------|
| SC Type: | TnI or Tn |
| Cardinality: | Any |
| Vector Type: | nCV |
| Max Similarity: | 1 |
| Min Similarity: | 0 |
| Average Value: | 0.45 |
| No. Values: | 101 |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.4,C4,C5 |
| I-related: | Yes |
| Z-related: | Yes |

A.6 ISAACSON

A.6.1 AMEMB₂

Proposed by ?, pp. 8, AMEMB₂ (Adjusted MEMB₂) is a scaled version MEMB₂ (A.5.2), measuring the mutual embedding of ICs.

$$AMEMB_2 = \frac{2 \times MEMB_2(X, Y)}{(\#X(\#X - 1) + \#Y(\#Y - 1))}$$

| | |
|-----------------|-----|
| SC Type: | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | 1 |
| Min Similarity: | 0 |
| Average Value: | |
| No. Values: | |
| Criteria Met: | |

A.6.2 IcVSIM

Presented in ?, pp. 18, IcVSIM (Interval-Class Vector Similarity Relation) is the standard deviation of the entries in the ICVs of two SCs. IcVSIM is a scaled version of s.i. (A.3.1). IdV_i is the i th term in the vector defined by ICV(X)-ICV(Y) and \overline{dV} is the average (mean) of its entries.

$$IcVSIM(X, Y) = \sqrt{\frac{\sum (IdV_i - \overline{dV})^2}{6}}$$

| | |
|-----------------|------------|
| SC Type | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | 0 |
| Min Similarity: | 3.64 |
| Average Value: | 1.2 |
| No. Values: | 121 |
| Criteria Met: | C1,C2,C3.4 |
| I-related: | No |
| Z-related: | No |

A.6.3 ISIM2

Presented in ?, ISIM2 is a scaled version of IcVSIM (A.6.2). The square root is taken of each term in the ICVs. Isaacson argues that each additional instance of an IC contributes less to similitude. However, ? found ISIM2 to be inconsistent with itself when applying MDS to the values produced.

| | |
|-----------------|------------|
| SC Type | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | |
| Min Similarity: | |
| Average Value: | |
| No. Values: | |
| Criteria Met: | C1,C2,C3.4 |

A.6.4 ANGLE (Isaacson & Scott)

? propose a geometric method which is identical to that of Cos/theta (A.4.3) but instead gives the size of the angle in degrees.

$$ANGLE(X, Y) = \arccos Cos\theta(X, Y)$$

| | |
|-----------------|----------------------|
| SC Type | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | |
| Min Similarity: | |
| Average Value: | |
| No. Values: | |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.4 |
| I-related: | No |
| Z-related: | No |

A.7 LEWIN

A.7.1 REL

Presented in ?, REL compares the nCVs of two SCs for all the nCs. Like MEMB_n (A.5.2), REL only considers non-zero entries however, this is achieved by multiplication (taking the geometric mean) of corresponding nCV terms.

$$REL(X, Y) = \frac{\sum_{i=1}^p \sqrt{SUB(X)_i \times SUB(Y)_i}}{\sqrt{\#SUB(X) \times \#SUB(Y)}}$$

$$REL(X, Y) = \frac{\#GMV(SUB(X), SUB(Y))}{GM(\#SUB(X), \#SUB(Y))}$$

where SUB(X) consists of concatenated nCVs and has a length p.

| | |
|-----------------|----------------------------|
| SC Type: | TnI or Tn |
| Cardinality: | Any |
| Vector Type: | nCV |
| Max Similarity: | 1 |
| Min Similarity: | 0 |
| Average Value: | 0.57 |
| No. Values: | 91 |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.4,C4,C5 |
| I-related: | Yes |
| Z-related: | Yes |

A.7.2 REL₂

? suggested a number of manifestations of the basic REL concept including REL₂ which measures only intervallic similarity.

$$REL_2(X, Y) = \frac{2 \times \sum \sqrt{(x_i y_i)}}{\sqrt{(\#X(\#X - 1)\#Y(\#Y - 1))}}$$

| | |
|-----------------|----------------------|
| SC Type: | TnI |
| Cardinality: | Any |
| Vector Type: | ICV |
| Max Similarity: | 1 |
| Min Similarity: | 0 |
| Average Value: | |
| No. Values: | |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.4 |

A.8 CASTREN

A.8.1 Castren's Difference Vector

Castren specifies a different type of DV, which we shall call cDV to distinguish it from the regular DV. It consists of two rows, $cDV_x(X, Y) = X - Y$ and $cDV_y(X, Y) = Y - X$. Any negative values in either of the rows are set to zero. In addition Castren defines the weighted difference vector (wcDV) of two vectors X and Y as:

$$wcDV = \frac{cDV(X, Y)}{\#cDV(X, Y)} \times 100$$

A.8.2 nC%V

Presented in ? for use in %REL_n, nC%V(X) (n-class subset percentage vector) gives the percentage subset-class contents of an SC, X. The 2C%V is the Interval percentage vector.

$$nC\%V(X) = \frac{nCV(X)}{\#nCV(X)} \times 100$$

A.8.3 %REL_n

Presented in ?, %REL_n (Percentage Relation) is a modification of sf (A.2.1) using the nC%Vs (A.8.2) instead of ICVs. %REL_n can be used as a stand-alone measure, however it is primarily intended as an intermediate step in T%REL and RECREL (A.8.4 and A.8.5).

$$\%REL_n(X, Y) = \frac{\#DV(nC\%V(X), nC\%V(Y))}{2}$$

| | |
|-----------------|------------------------------|
| SC Type | TnI or Tn |
| Cardinality: | Any |
| Measure Type: | Single nC |
| Vector Type: | nC%V |
| Max Similarity: | 0 |
| Min Similarity: | 100 |
| Average Value: | 30 |
| No. Values: | 85 |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.3,C3.4,C4 |
| I-related: | Sometimes |
| Z-related: | Sometimes |

A.8.4 T%REL

Presented in ?, T%REL (Total Percentage Relation) is the mean average of the vlaues of %REL_n for all values of n from 2 to m where, if $\#X \neq \#Y$, $m = \text{MIN}(\#X, \#Y)$ else $m = \#X - 1$.

$$T\%REL(X, Y) = \frac{\sum_{n=2}^m \%REL_n(X, Y)}{m - 1}$$

| | |
|-----------------|---------------------------------|
| SC Type: | TnI or Tn |
| Cardinality: | Any |
| Measure Type: | Total |
| Vector Type: | nC%V |
| Max Similarity: | 0 |
| Min Similarity: | 100 |
| Average Value: | 63 |
| No. Values: | 79 |
| Criteria Met: | C1,C2,C3.1,C3.2,C3.3,C3.4,C4,C5 |
| I-related: | Yes |
| Z-related: | Yes |

A.8.5 RECREL

Presented in ?, RECREL (Recursive Relation) recursively compares the subsets and subsets of subsets of two SCs using %REL_n (A.8.3). The comparison procedure is quite complicated and potentially involves evaluating %REL_n thousands of times.

| | |
|-----------------|-----------|
| SC Type: | TnI or Tn |
| Cardinality: | Any |
| Measure Type: | Total |
| Vector Type: | nC%V |
| Max Similarity: | 0 |
| Min Similarity: | 100 |
| Average Value: | |
| No. Values: | 89 |
| Criteria Met: | All |
| I-related: | Yes |
| Z-related: | Yes |

A.9 BUCHLER

A.9.1 nSATV

Presented in ?, chap. 2.3 nSATV(X) (Saturation Vector) is a dual vector consisting of two rows, nSATV_A(X) and nSATV_B(X). It shows extent to which an SC is saturated with subclasses of cardinality n. The steps for computing nSATV(X) are as follows:

1. Compute the nCVs for all SCs of cardinality $\#X$.
2. Find the minimum and maximum values for each vector position. These values form vectors $Max_n(\#X)$ and $Min_n(\#X)$.
3. Compute the following two vectors: $MaxMinus = DV(nCV(X), Max_n(\#X))$ and $MinPlus = DV(nCV(X), Min_n(\#X))$
4. $nSATV_A(X)_i = \text{MIN}(MaxMinus_i, MinPlus_i)$ and $nSATV_B(X)_i = \text{MAX}(MaxMinus_i, MinPlus_i)$
5. If $MaxMinus_i = MinPlus_i$, $nSATV_A(X)_i = MaxMinus_i$ and $nSATV_B(X)_i = MinPlus_i$

A.9.2 SATSIM_n

Presented in ?, chap. 2.4, SATSIM_n (Saturation Similarity index) compares the nSATVs of two SCs and involves the following steps:

1. Calculate $nSATV(X)$ and $nSATV(Y)$
2. Calculate the vectors $nSATV_{row}(X)$ and $nSATV_{row}(Y)$.
3. The function “row” maps the MaxMinus values of one $nSATV$ to the MaxMinus values of the other. If $nSATV_A(X)_i$ is a MaxMinus value and $nSATV_A(X)_i$ is also a MaxMinus value, $row = A$ ($nSATV_{row}(X)_i = nSATV_A(X)_i$), otherwise $row = B$.
4. Finally $SATSIM_n(X, Y)$ is given by the formula:

$$SATSIM_n(X, Y) = \frac{\#DV(nSATV_A(X), nSATV_{row}(Y)) + \#DV(nSATV_A(Y), SATV_{row}(X))}{\#DV(nSATV_A(X), SATV_B(X)) + \#DV(SATV_A(Y), SATV_B(Y))}$$

A.9.3 AvgSATSIM

Presented in ?, chap. 2.10, AvgSATSIM (Average Saturation Similarity index) is the mean of $SATSIM_n$ values where $m = MIN(\#X, \#Y)$.

$$AvgSATSIM(X, Y) = \frac{\sum_{n=2}^{m-1} SATSIM_n(X, Y)}{m - 2}$$

A.9.4 TSATSIM

Presented in ?, chap. 2.10, TSATSIM (Total Saturartion Vector Similarity index) is an extension of $SATSIM_n$. TSATSIM is the quotient of the sum of all $SATSIM_n$ numerators and denominators for all values of n from 2 to $m-1$ where $m = MIN(\#X, \#Y)$.