

Green Bonds EDA

Noah Leibowitz & Siya Motwani

2026-07-08

```
# Load in all libraries
library(dplyr)
library(tidyverse)
library(readxl)
library(ggplot2)
library(maps)
library(tidyr)

# Solar Data
solar_2024 <- read_excel("Solar.xlsx", sheet = "2024")
solar_2023 <- read_excel("Solar.xlsx", sheet = "2023")
solar_2022 <- read_excel("Solar.xlsx", sheet = "2022")
solar_2021 <- read_excel("Solar.xlsx", sheet = "2021")
solar_2020 <- read_excel("Solar.xlsx", sheet = "2020")
solar_2019 <- read_excel("Solar.xlsx", sheet = "2019")
solar_2018 <- read_excel("Solar.xlsx", sheet = "2018")
solar_2017 <- read_excel("Solar.xlsx", sheet = "2017")
solar_2016 <- read_excel("Solar.xlsx", sheet = "2016")
solar_2015 <- read_excel("Solar.xlsx", sheet = "2015")
solar_2014 <- read_excel("Solar.xlsx", sheet = "2014")
solar_2013 <- read_excel("Solar.xlsx", sheet = "2013")

# Wind Data
wind_2024 <- read_excel("Wind.xlsx", sheet = "2024")
wind_2023 <- read_excel("Wind.xlsx", sheet = "2023")
wind_2022 <- read_excel("Wind.xlsx", sheet = "2022")
wind_2021 <- read_excel("Wind.xlsx", sheet = "2021")
wind_2020 <- read_excel("Wind.xlsx", sheet = "2020")
wind_2019 <- read_excel("Wind.xlsx", sheet = "2019")
wind_2018 <- read_excel("Wind.xlsx", sheet = "2018")
wind_2017 <- read_excel("Wind.xlsx", sheet = "2017")
wind_2016 <- read_excel("Wind.xlsx", sheet = "2016")
wind_2015 <- read_excel("Wind.xlsx", sheet = "2015")
wind_2014 <- read_excel("Wind.xlsx", sheet = "2014")
wind_2013 <- read_excel("Wind.xlsx", sheet = "2013")

# Green Bond Issuance and RPS Data
GBI <- read_excel("GB_RPS Hard Code.xlsx")

# Pre-Trends Wind Data
pre_wind <- read_csv("Wind Pre.csv")
```

```

# Pre-Trends Solar Data
pre_solar <- read_excel("Solar 2010 - 2012.xlsx")

# Real GDP Data
gdp_raw <- read_csv("SAGDP1_ALL_AREAS_1997_2025.csv")

# State Population Data (2010 - 2020)
intercensal_raw <- read_excel(
  "nst-est2020int-pop.xlsx",
  sheet = 1,
  skip = 3
)

# State Population Data (2020 - 2024)
vintage2024_raw <- read_csv("NST-EST2024-POPCHG2020-2024.csv", stringsAsFactors = FALSE)

```

```

# Cleaning Pre-Wind Data

clean_wind_panel <- pre_wind %>%

# Remove any rows where the operational year is missing
filter(!is.na(p_year)) %>%

# Select only the state, year, and capacity columns
select(t_state, p_year, t_cap) %>%

# Group by state and year, and convert individual turbine kW to total MW added
group_by(t_state, p_year) %>%
summarize(mw_added = sum(t_cap, na.rm = TRUE) / 1000, .groups = 'drop') %>%

# Open 50-state grid for pre-trends window
complete(t_state, p_year = 2000:2026, fill = list(mw_added = 0)) %>%
arrange(t_state, p_year) %>%
group_by(t_state) %>%

# Cumulative sum holds capacity at 0 until first project,
# then holds total capacity flat forward through the years
mutate(total_wind_mw = cumsum(mw_added)) %>%
rename(state = t_state, year = p_year) %>%

# Filter the outcome variable to strictly include 2010-2012
filter(year >= 2010 & year <= 2013)

# Split by each year and pivot year column to display state and capacity by year
clean_wind_panel <- clean_wind_panel %>%
  select(state, year, total_wind_mw) %>%
  group_by(year) %>%
  group_split(.keep = FALSE)
wind_2010 <- clean_wind_panel[[1]]
wind_2011 <- clean_wind_panel[[2]]
wind_2012 <- clean_wind_panel[[3]]

# Used to compare discrepancies between two different data sources for wind capacity

```

```
wind_2013_test <- clean_wind_panel[[4]]
```

```
# Cleaning Pre-Solar Data
```

```
# 2. Extract state, and 2010 - 2013 capacity data
```

```
solar_wide_cumulative <- pre_solar %>%
```

```
  select(
    State = 1,
    raw_2010 = 2,
    raw_2011 = 3,
    raw_2012 = 4,
    raw_2013 = 5
  )
```

```
# Convert from solar capacity added to cumulative capacity
```

```
solar_wide_cumulative <- solar_wide_cumulative %>%
```

```
  mutate(
    # 2010 is just the base year
    Year_2010 = replace_na(raw_2010, 0),
```

```
    # 2011 = 2010 + 2011
```

```
    Year_2011 =
      replace_na(raw_2010, 0) +
      replace_na(raw_2011, 0),
```

```
    # 2012 = 2010 + 2011 + 2012
```

```
    Year_2012 =
      replace_na(raw_2010, 0) +
      replace_na(raw_2011, 0) +
      replace_na(raw_2012, 0),
```

```
    # 2013 = 2010 + 2011 + 2012 + 2013
```

```
    # Used to compare discrepancies between two different data sources for solar capacity
```

```
    Year_2013_test =
      replace_na(raw_2010, 0) +
      replace_na(raw_2011, 0) +
      replace_na(raw_2012, 0) +
      replace_na(raw_2013, 0)
  ) %>%
```

```
# 5. Output clean state name and cumulative columns for each year
```

```
select(State, Year_2010, Year_2011, Year_2012, Year_2013_test)
```

```
# Cleaning Post-Wind Data
```

```
# Add year columns
```

```
wind_2010 <- wind_2010 %>%
```

```
  rename(State = `state`, wind_capacity = `total_wind_mw`) %>%
  mutate(year = 2010) %>%
  arrange(State)
```

```
wind_2011 <- wind_2011 %>%
```

```
  rename(State = `state`, wind_capacity = `total_wind_mw`) %>%
```

```

mutate(year = 2011) %>%
  arrange(State)

wind_2012 <- wind_2012 %>%
  rename(State = `state`, wind_capacity = `total_wind_mw`) %>%
  mutate(year = 2012) %>%
  arrange(State)

# Loop through every year from 2013 to 2024
for (y in 2013:2024) {

  # Generate the variable name ("wind_year")
  var_name <- paste0("wind_", y)

  # Temporary data frame to do cleaning
  temp_df <- get(var_name)
  cleaned_df <- temp_df %>%

    # Adds the current loop's year
    mutate(year = y) %>%
    rename(wind_capacity = `Nameplate Capacity (MW)`) %>%
    group_by(State, year) %>%
    summarize(
      wind_capacity = sum(wind_capacity, na.rm = TRUE), .groups = "drop"
    ) %>%
    arrange(State)

  # 4. Save the cleaned data frame back to its original variable name
  assign(var_name, cleaned_df)
}

# Combine all years (2010 - 2024)
post_wind_all = bind_rows(
  wind_2010, wind_2011, wind_2012, wind_2013, wind_2014,
  wind_2015, wind_2016, wind_2017, wind_2018, wind_2019,
  wind_2020, wind_2021, wind_2022, wind_2023, wind_2024
)

# To account for discrepancies between pre and post data sources
# Scale pre-trend data
wind_2013_uswtodb <- wind_2013_test %>%
  rename(State = state, wind_uswtodb = total_wind_mw)

wind_ratios <- wind_2013 %>%
  rename(wind_eia = wind_capacity) %>%
  left_join(wind_2013_uswtodb, by = "State") %>%
  mutate(

    # Ratio = EIA Capacity / USWTDB Capacity
    scale_factor = wind_eia / wind_uswtodb
  ) %>%
  mutate(scale_factor = if_else(
    is.nan(scale_factor) | is.infinite(scale_factor), 1, scale_factor)

```

```

    ) %>%
    select(State, scale_factor)

# Scale down 2010-2012 wind capacities to meet the EIA baseline
post_wind_all <- post_wind_all %>%
  left_join(wind_ratios, by = "State") %>%
  mutate(
    wind_capacity = if_else(
      year <= 2012, wind_capacity * scale_factor, wind_capacity
    )
  ) %>%
  select(-scale_factor)

# Create a master data frame containing all 48 contiguous states
# Excludes Hawaii and Alaska
contiguous_48 <- state.abb[!state.abb %in% c("AK", "HI")]
master_states <- data.frame(State = contiguous_48)

# Pivot wind data from long format to wide format
wind_wide <- post_wind_all %>%
  pivot_wider(
    names_from = year,
    values_from = wind_capacity,
    names_prefix = "yr_"
  )

# Left join the master list with wind data and sort states alphabetically
final_wind_panel <- master_states %>%
  left_join(wind_wide, by = "State") %>%
  arrange(State)

final_wind_panel_clean <- final_wind_panel %>%

  # Temporarily bring columns back to a long format to work row-by-row per state
  pivot_longer(
    cols = starts_with("yr_"),
    names_to = "year",
    values_to = "wind_capacity"
  ) %>%
  # 2. Group by State and sort by year to ensure sequential order
  group_by(State) %>%
  arrange(State, year) %>%
  # Fill NA values with the previous year's value
  fill(wind_capacity, .direction = "down") %>%
  # If it's still NA (meaning it never had any capacity), replace with 0
  mutate(wind_capacity = replace_na(wind_capacity, 0)) %>%
  # Bring it back to the wide layout
  pivot_wider(
    names_from = year,
    values_from = wind_capacity
  ) %>%
  ungroup()

```

```

# Cleaning Post-Solar Data

# Add year columns
solar_2010 <- solar_wide_cumulative %>%
  select(State, solar_capacity = Year_2010) %>%
  mutate(
    year = 2010,
    State = state.abb[match(State, state.name)]
  ) %>%
  arrange(State)

solar_2011 <- solar_wide_cumulative %>%
  select(State, solar_capacity = Year_2011) %>%
  mutate(
    year = 2011,
    State = state.abb[match(State, state.name)]) %>%
  arrange(State)

solar_2012 <- solar_wide_cumulative %>%
  select(State, solar_capacity = Year_2012) %>%
  mutate(
    year = 2012,
    State = state.abb[match(State, state.name)]) %>%
  arrange(State)

solar_2013_test <- solar_wide_cumulative %>%
  select(State, solar_capacity = Year_2013_test) %>%
  mutate(
    year = 2013,
    State = state.abb[match(State, state.name)]) %>%
  arrange(State)

# Loop through every year from 2013 to 2024
# Same way we did for the post wind data
for (y in 2013:2024) {
  var_name <- paste0("solar_", y)
  temp_df <- get(var_name)
  cleaned_df <- temp_df %>%
    mutate(year = y) %>%
    rename(solar_capacity = `Nameplate Capacity (MW)`) %>%
    group_by(State, year) %>%
    summarize(
      solar_capacity = sum(solar_capacity, na.rm = TRUE), .groups = "drop"
    ) %>%
    arrange(State)
  assign(var_name, cleaned_df)
}

post_solar_all = bind_rows(
  solar_2010, solar_2011, solar_2012, solar_2013,
  solar_2014, solar_2015, solar_2016, solar_2017,
  solar_2018, solar_2019, solar_2020, solar_2021,
  solar_2022, solar_2023, solar_2024
)

```

```

)

# Isolate the 2013 Berkeley Lab data and convert names to abbreviations
solar_2013_lbnl_check <- solar_wide_cumulative %>%
  select(State, solar_lbnl = Year_2013_test) %>%
  mutate(
    State = state.abb[match(State, state.name)]
  )

# Join the EIA 2013 data with the Berkeley Lab data
solar_ratios <- solar_2013 %>%
  rename(solar_eia = solar_capacity) %>%
  left_join(solar_2013_lbnl_check, by = "State") %>%
  mutate(

    # Create a conditional scale factor to prevent division by zero or Na errors
    # If there's missing or 0 data, don't scale by ratio to avoid dividing by 0
    # Otherwise, scale by the ratio of the two data source values
    scale_factor = if_else(
      is.na(solar_eia) | is.na(solar_lbnl) | solar_eia == 0 | solar_lbnl == 0,
      1,
      solar_eia / solar_lbnl
    )
  ) %>%
  select(State, scale_factor)

# Apply the ratio to the pre-trends years (2010-2012)
post_solar_all <- post_solar_all %>%
  left_join(solar_ratios, by = "State") %>%
  mutate(
    # Double check protection for any lingering NAs
    scale_factor = replace_na(scale_factor, 1),
    solar_capacity = if_else(
      year <= 2012, solar_capacity * scale_factor, solar_capacity
    )
  ) %>%
  select(-scale_factor)

contiguous_48 <- state.abb[!state.abb %in% c("AK", "HI")]
master_states <- data.frame(State = contiguous_48)
solar_wide <- post_solar_all %>%
  group_by(State, year) %>%
  summarize(solar_capacity = sum(solar_capacity, na.rm = TRUE), .groups = "drop") %>%
  pivot_wider(
    names_from = year,
    values_from = solar_capacity,
    names_prefix = "yr_"
  )

final_solar_panel <- master_states %>%
  left_join(solar_wide, by = "State") %>%
  arrange(State)

```

```

# Interpolate missing values
final_solar_panel_clean <- final_solar_panel %>%
  pivot_longer(
    cols = starts_with("yr_"),
    names_to = "year",
    values_to = "solar_capacity"
  ) %>%
  group_by(State) %>%
  arrange(State, year) %>%
  fill(solar_capacity, .direction = "down") %>%
  mutate(solar_capacity = replace_na(solar_capacity, 0)) %>%
  pivot_wider(
    names_from = year,
    values_from = solar_capacity
  ) %>%
  ungroup()

```

```

# Merge state population data: 2010-2019 (Intercensal)
# + 2020-2024 (Vintage 2024)

# Rename columns: first column is geography, next is 2010 base, then 2011-2019,
# then 2020 census
colnames(intercensal_raw) <- c(
  "Geography", "Base2010", paste0("Y", 2010:2019), "Census2020"
)

intercensal_states <- intercensal_raw %>%
  filter(grepl("^\\.", Geography)) %>%
  mutate(State = sub("^\\.", "", Geography)) %>%
  filter(State != "Puerto Rico") %>%

# filter(State != "District of Columbia") %>%
select(State, paste0("Y", 2010:2019)) %>%
pivot_longer(
  cols = paste0("Y", 2010:2019),
  names_to = "Year",
  values_to = "Population"
) %>%
mutate(Year = as.integer(sub("Y", "", Year)))

vintage2024_states <- vintage2024_raw %>%
  filter(trimws(as.character(SUMLEV)) %in% c("40", "040")) %>%
  filter(NAME != "Puerto Rico") %>%
  select(NAME, POPESTIMATE2020, POPESTIMATE2021, POPESTIMATE2022,
    POPESTIMATE2023, POPESTIMATE2024) %>%
  rename(State = NAME) %>%
  pivot_longer(
    cols = starts_with("POPESTIMATE"),
    names_to = "Year",
    values_to = "Population"
  ) %>%
  mutate(Year = as.integer(sub("POPESTIMATE", "", Year)))

```



```

# Merge into one long-format table: 2010-2024, states only
state_population_2010_2024 <- dplyr::bind_rows(
  intercensal_states, vintage2024_states
) %>%
  arrange(State, Year)

# Reshape to wide format: one row per state, one column per year
state_population_wide <- state_population_2010_2024 %>%
  pivot_wider(
    names_from = Year, values_from = Population, names_prefix = "Y"
  )

# Map full state names to 2-letter postal codes
population_long <- state_population_2010_2024 %>%
  filter(State %in% c(state.name, "District of Columbia")) %>%
  mutate(
    year = as.numeric(Year),
    Population = as.numeric(Population),
    State = case_when(
      State == "District of Columbia" ~ "DC",
      TRUE ~ state.abb[match(State, state.name)]
    )
  ) %>%
  select(State, year, Population)

# Save outputs
write.csv(
  state_population_2010_2024,
  "state_population_2010_2024_long.csv",
  row.names = FALSE
)

write.csv(
  state_population_wide,
  "state_population_2010_2024_wide.csv",
  row.names = FALSE
)

```

```

# Cleaning Real GDP data

gdp_long <- gdp_raw %>%
  # Keep only the Real GDP rows (LineCode is 1)
  filter(LineCode == 1) %>%

  # Drop the "United States" row and any regional summaries
  filter(GeoName %in% c(state.name, "District of Columbia")) %>%

  # 4. Keep only the State name column and years 2010 - 2024
  select(State = GeoName, `2010`:`2024`) %>%

  # Reshape from wide (columns) to long (rows) to match population data
  pivot_longer(
    cols = `2010`:`2024`,

```

```

    names_to = "year",
    values_to = "real_gdp"
  ) %>%

  # Fix data types, scale by 1 million,
  # and convert full state names to 2-letter postal codes
  mutate(
    year = as.numeric(year),
    real_gdp = as.numeric(real_gdp) * 1000000,
    State = case_when(
      State == "District of Columbia" ~ "DC",
      TRUE ~ state.abb[match(State, state.name)]
    )
  )

# Building the Panel data set

# Create a dedicated long-format version for regressions and plotting
final_wind_panel_long <- final_wind_panel %>%

# Remove any rows where the operational year is missing/blank
# Pivot to long format
pivot_longer(
  cols = starts_with("yr_"),
  names_to = "year",
  values_to = "wind_capacity"
) %>%

# Strip "yr_" so the year column becomes a clean, parse-able number
mutate(year = as.numeric(gsub("yr_", "", year))) %>%

# Sort sequentially by state and year to prepare for the carry-forward step
group_by(State) %>%
arrange(State, year) %>%

# Handle missing data gaps using Last Observation Carried Forward (LOCF)
fill(wind_capacity, .direction = "down") %>%

# Switch any remaining early NAs to 0
mutate(wind_capacity = replace_na(wind_capacity, 0)) %>%

# Ungroup so it behaves like a standard data frame
ungroup()

# Create a dedicated long-format version for solar regressions and plotting
final_solar_panel_long <- final_solar_panel %>%
pivot_longer(
  cols = starts_with("yr_"),
  names_to = "year",
  values_to = "solar_capacity"
) %>%
mutate(year = as.numeric(gsub("yr_", "", year))) %>%
group_by(State) %>%

```

```

arrange(State, year) %>%
fill(solar_capacity, .direction = "down") %>%
mutate(solar_capacity = replace_na(solar_capacity, 0)) %>%
ungroup()

# One-hot-encoding of corporate & municipal issuance, and RPS issuance
# Corporate: 1 if current year >= corporate year, otherwise 0
# Municipal: 1 if current year >= municipal year, otherwise 0
# RPS: 1 if current year >= RPS enacted AND current year < RPS expired
# If never expired, remains 1
# GB: 1 if a state issued either type of green bond,
# 0 if a state issued neither
policy_clean <- GBI %>%

# 1. Standardize state names to postal abbreviations
mutate(State = state.abb[match(State, state.name)]) %>%
filter(!is.na(State)) %>%

# 2. Convert character strings/NA values to proper numeric years or infinity
mutate(
  corp_start = as.numeric(gsub("N/A", NA, `Corporate Year`)),
  muni_start = as.numeric(gsub("N/A", NA, `Municipal Year`)),
  rps_start = as.numeric(gsub("N/A", NA, `RPS Year`)),
  rps_end = as.numeric(gsub("N/A", NA, `RPS Repealed/Expired`)),

  # If RPS was never repealed, set its expiration to infinity
  rps_end = if_else(is.na(rps_end), Inf, rps_end)
) %>%
select(State, corp_start, muni_start, rps_start, rps_end)

# Merge Wind and Solar long panels into a single joint panel
joint_energy_panel <- final_wind_panel_long %>%
full_join(final_solar_panel_long, by = c("State", "year"))

# Combine panels and compute the logical thresholds
final_joint_regression_panel <- final_wind_panel_long %>%

# Join wind and solar together side-by-side
full_join(final_solar_panel_long, by = c("State", "year")) %>%

# Join cleaned timeline parameters
left_join(policy_clean, by = "State") %>%

# Join population metrics sequentially
left_join(population_long, by = c("State", "year")) %>%

# Join BEA real GDP metrics sequentially
left_join(gdp_long, by = c("State", "year")) %>%

# Build the binary metrics dynamically
mutate(
  is_corporate = if_else(!is.na(corp_start) & year >= corp_start, 1, 0),
  is_public = if_else(!is.na(muni_start) & year >= muni_start, 1, 0),

```

```

gb = if_else(is_corporate == 1 | is_public == 1, 1, 0),

# RPS Policy Dummy
is_rps = if_else(!is.na(rps_start) & year >= rps_start & year < rps_end, 1, 0)
) %>%

# Calculate all per capita and economic density metrics
mutate(
  real_gdp_per_capita      = real_gdp / Population,
  wind_capacity_per_capita = wind_capacity / Population,
  solar_capacity_per_capita = solar_capacity / Population
) %>%

# Drop the following columns to only include the one-hot-encodings in the panel
select(-corp_start, -muni_start, -rps_start, -rps_end)

# Export final panel data set to a CSV file
# This is the data set to be used in Stata for our econometric analysis
write.csv(final_joint_regression_panel, "panel.csv", row.names = FALSE)

# Wind Line Chart


# 1. Pull data directly from the panel for plotting
plot_data <- final_joint_regression_panel %>%

  # Keep only the state, year, and wind capacity per capita columns
  select(State, year, wind_capacity_per_capita)

# Create the line plot showing per capita distribution
ggplot(plot_data, aes(x = year, y = wind_capacity_per_capita, group = State, color = State)) +
  geom_line(size = 0.8, alpha = 0.7) +
  geom_point(size = 1.2, alpha = 0.5) +
  scale_x_continuous(breaks = seq(2010, 2024, by = 2)) +
  labs(
    title = "Wind Capacity Per Capita Over Time For Lower 48 States (2010 - 2024)",
    x = "Year",
    y = "Wind Capacity Per Capita (MW / Population)"
  ) +
  theme_minimal() +
  theme(
    legend.position = "none",
    plot.title = element_text(hjust = 0.5, face = "bold"),
    plot.subtitle = element_text(hjust = 0.5)
  )

```

Wind Capacity Per Capita Over Time For Lower 48 States (2010 – 2024)



```
ggsave(
  filename = "wind_capacity_per_capita_timeline.png",
  width = 10,
  height = 6,
  dpi = 300
)
```


Solar Line Chart

```
solar_plot_data_all <- final_joint_regression_panel %>%
  select(State, year, solar_capacity_per_capita)

ggplot(solar_plot_data_all, aes(x = year, y = solar_capacity_per_capita, group = State, color = State))
  geom_line(size = 0.8, alpha = 0.7) +
  geom_point(size = 1.2, alpha = 0.6) +
  scale_x_continuous(breaks = seq(2010, 2024, by = 2)) +
  labs(
    title = "Solar Capacity Per Capita Over Time For Lower 48 States (2010 - 2024)",
    x = "Year",
    y = "Solar Capacity Per Capita (MW / Population)"
  ) +
  theme_minimal() +
  theme(
    legend.position = "none",
    plot.title = element_text(hjust = 0.5, face = "bold"),
    plot.subtitle = element_text(hjust = 0.5)
  )
```

)

Solar Capacity Per Capita Over Time For Lower 48 States (2010 – 2024)



```
ggsave(
  filename = "solar_capacity_per_capita_timeline.png",
  width = 10,
  height = 6,
  dpi = 300
)
```

```
# Wind map

# Determine which states are never treated (gb == 0)
never_treated_states <- final_joint_regression_panel %>%
  group_by(State) %>%
  summarize(ever_issued = max(gb, na.rm = TRUE), .groups = "drop") %>%
  filter(ever_issued == 0) %>%
  pull(State)

# Extract 2024 data
wind_map_2024 <- final_joint_regression_panel %>%
  filter(year == 2024) %>%
  select(State, capacity = wind_capacity_per_capita) %>%
  mutate(region = tolower(state.name[match(State, state.abb)]))

us_states <- map_data("state")
```

```

wind_geo_data <- us_states %>%
  left_join(wind_map_2024, by = "region")

# Create a label data set containing only the never-treated states
never_treated_labels <- data.frame(
  State = state.abb,
  long = state.center$x,
  lat = state.center$y
) %>%

# Filter out AK and HI, and isolate the never-treated control states
filter(!State %in% c("AK", "HI"), State %in% never_treated_states)

# Generate Wind Map
ggplot() +
  # Base map layer showing capacities
  geom_polygon(data = wind_geo_data, aes(x = long, y = lat, group = group, fill = capacity),
    color = "white", size = 0.2) +


  # Overlay text labels for the never-treated states
  geom_text(data = never_treated_labels, aes(x = long, y = lat, label = State),
    color = "black", fontface = "bold", size = 3.5, check_overlap = TRUE) +

  # Use Journal of Economics appropriate coloring scheme
  scale_fill_gradientn(
    name = "MW / Capita",
    colors = c("#E6D3B3", "#D49A73", "#C07A5A"),
    labels = scales::comma_format(),
    na.value = "grey90"
  ) +
  coord_map("albers", lat0 = 39, lat1 = 45) +
  labs(
    title = "Cumulative US Wind Capacity Per Capita by State (2024)",
    subtitle = "Labeled states represent the never-treated control group"
  ) +
  theme_void() +
  theme(
    legend.position = "right",
    legend.box.margin = margin(t = 0, r = 20, b = 0, l = 0),
    legend.margin = margin(t = 0, r = 0, b = 0, l = 0),
    plot.title = element_text(hjust = 0.5, face = "bold"),
    plot.subtitle = element_text(hjust = 0.5, size = 10, color = "gray30")
  )

```

Cumulative US Wind Capacity Per Capita by State (2024)

Labeled states represent the never-treated control group



```
ggsave(  
  filename = "wind_map_per_capita.png",  
  width = 10,  
  height = 6,  
  dpi = 300  
)  
  
# Solar map  
  
never_treated_states <- final_joint_regression_panel %>%  
  group_by(State) %>%  
  summarize(ever_issued = max(gb, na.rm = TRUE), .groups = "drop") %>%  
  filter(ever_issued == 0) %>%  
  pull(State)  
  
solar_map_2024 <- final_joint_regression_panel %>%  
  filter(year == 2024) %>%  
  select(State, capacity = solar_capacity_per_capita) %>%  
  mutate(region = tolower(state.name[match(State, state.abb)]))  
  
us_states <- map_data("state")  
  
solar_geo_data <- us_states %>%  
  left_join(solar_map_2024, by = "region")  
  
never_treated_labels <- data.frame(  
  State = never_treated_states,  
  capacity = 0,  
  region = "never_treated"
```



```


State = state.abb,
long = state.center$x,
lat = state.center$y
) %>%
  filter(!State %in% c("AK", "HI"), State %in% never_treated_states)

ggplot() +
  geom_polygon(data = solar_geo_data, aes(x = long, y = lat, group = group, fill = capacity),
    color = "white", size = 0.2) +
  geom_text(data = never_treated_labels, aes(x = long, y = lat, label = State),
    color = "black", fontface = "bold", size = 3.5, check_overlap = TRUE) +
  scale_fill_gradientn(
    name = "MW / Capita",
    colors = c("#E6D3B3", "#D49A73", "#C07A5A"),
    labels = scales::comma_format(),
    na.value = "grey90"
  ) +
  coord_map("albers", lat0 = 39, lat1 = 45) +
  labs(
    title = "Cumulative US Solar Capacity Per Capita by State (2024)",
    subtitle = "Labeled states represent the never-treated control group"
  ) +
  theme_void() +
  theme(
    legend.position = "right",
    legend.box.margin = margin(t = 0, r = 20, b = 0, l = 0),
    legend.margin = margin(t = 0, r = 0, b = 0, l = 0),
    plot.title = element_text(hjust = 0.5, face = "bold"),
    plot.subtitle = element_text(hjust = 0.5, size = 10, color = "gray30")
  )

```

Cumulative US Solar Capacity Per Capita by State (2024)

Labeled states represent the never-treated control group



```
ggsave(
  filename = "solar_map_per_capita.png",
  width = 10,
  height = 6,
  dpi = 300
)
```

Corporate Green Bond Dot Plot

Filter out NAs and calculate a stacking index for overlapping years

```
corp_plot_data <- policy_clean %>%
  filter(!is.na(corp_start)) %>%
  select(State, year = corp_start) %>%
  group_by(year) %>%
  mutate(stack_index = row_number()) %>%
  ungroup()
```

Generate Corporate timeline

```
ggplot(corp_plot_data, aes(x = year, y = stack_index, label = State)) +
```

Draw a clean baseline

```
geom_hline(yintercept = 0, color = "gray50", size = 0.5) +
```

Replace dots/digits with actual state abbreviations

```
geom_text(color = "#C07A5A", fontface = "bold", size = 4.5) +
```

```

# Force x-axis to show clear integer years (year interval is 1)
scale_x_continuous(breaks = seq(min(corp_plot_data$year), max(corp_plot_data$year), by = 1)) +


# Pad the vertical axis slightly so text doesn't cut off
scale_y_continuous(expand = expansion(mult = c(0.1, 0.2))) +
labs(
  title = "Corporate Green Bond Issuance Timeline by State",
  x = "Year of Initial Issuance",
  y = ""
) +
theme_minimal() +
theme(
  plot.title = element_text(hjust = 0.5, face = "bold", size = 14),

  # Clear horizontal lines to keep it looking like a stem plot
  panel.grid.major.y = element_blank(),
  panel.grid.minor.y = element_blank(),

  # Hide y-axis numbers since height just represents stacking
  axis.text.y = element_blank(),
  axis.ticks.y = element_blank()
)

```

Corporate Green Bond Issuance Timeline by State



```

ggsave(
  filename = "corporate_issuance_timeline.png",

```

```

width = 10,
height = 5,
dpi = 300
)


# Public/Municipal Green Bond Stem and Leaf Plot

muni_plot_data <- policy_clean %>%
  filter(!is.na(muni_start)) %>%
  select(State, year = muni_start) %>%
  group_by(year) %>%
  mutate(stack_index = row_number()) %>%
  ungroup()

ggplot(muni_plot_data, aes(x = year, y = stack_index, label = State)) +
  geom_hline(yintercept = 0, color = "gray50", size = 0.5) +
  geom_text(color = "#D49A73", fontface = "bold", size = 4.5) +
  scale_x_continuous(breaks = seq(min(muni_plot_data$year), max(muni_plot_data$year), by = 1)) +
  scale_y_continuous(expand = expansion(mult = c(0.1, 0.2))) +
  labs(
    title = "Public/Municipal Green Bond Issuance Timeline by State",
    x = "Year of Initial Issuance",
    y = ""
  ) +
  theme_minimal() +
  theme(
    plot.title = element_text(hjust = 0.5, face = "bold", size = 14),
    panel.grid.major.y = element_blank(),
    panel.grid.minor.y = element_blank(),
    axis.text.y = element_blank(),
    axis.ticks.y = element_blank()
  )

```

Public/Municipal Green Bond Issuance Timeline by State



```
ggsave(  
  filename = "municipal_issuance_timeline.png",  
  width = 10,  
  height = 5,  
  dpi = 300  
)
```