How do entropy and transport control Langevin convergence?

1 How do entropy and transport control Langevin convergence?

This handout takes one question from the Sampling unit and studies it in continuous time: how fast does the overdamped Langevin law approach equilibrium, and which geometric assumptions control that rate? The diffusion itself is

$$dX_t = -\nabla U(X_t)\,dt + \sqrt{2}\,dB_t,$$

with stationary law

$$\pi(x) = Z^{-1}e^{-U(x)}.$$

Here $Z = \int_{\mathbb{R}^d} e^{-U(x)}\,dx$ is assumed finite, and the evolving object is the law $\rho_t = \mathcal{L}(X_t)$ when X_0 does not start from equilibrium. The main quantities will be the free-energy gap, its dissipation along the Langevin flow, and the transport inequalities that turn dissipation into quantitative convergence.

This viewpoint extends the Sampling unit's gradient-based MCMC subsection, reuses the Optimization unit's language of objectives, proximal structure, and conditioning in measure space, and feeds directly into the Augmented-state sampling unit. There the same local geometry drives HMC and NUTS through momentum-carrying trajectories rather than one-step Langevin corrections.

There is also a concrete geometric reason to care. For a quadratic target

$$U(x) = \frac{1}{2}x^\top Hx,$$

every mode in an eigenbasis of H relaxes at its own exponential rate. If H is poorly conditioned, then one slow direction controls the last stage of convergence even when the other directions have already equilibrated. So even in the friendliest Gaussian case, convergence is already controlled by geometry. The rest of the note develops the general version of that picture: we identify a free-energy gap to equilibrium, compute its dissipation along the Langevin flow, convert that dissipation into explicit entropy and transport rates, and then read off how preconditioning changes the governing matrix in quadratic and locally quadratic models.

Throughout we work in the smooth, rapidly decaying setting when we differentiate under the integral sign or integrate by parts. Those formal calculations are the right ones, and the cited theorems extend them to broader classes of densities.

2 A notion of distance to equilibrium

The first task is to identify the right notion of distance to equilibrium. For Langevin, the natural candidate is not Euclidean distance between points but a functional on densities. The standard object here is the free-energy gap, and the corresponding density evolution is governed by the Fokker–Planck equation.

2.1 Relative entropy is a free-energy gap

Fix a potential $U : \mathbb{R}^d \to \mathbb{R}$ and define the target density

$$\pi(x) = Z^{-1} e^{-U(x)}, \qquad Z = \int_{\mathbb{R}^d} e^{-U(x)}\, dx.$$

For a probability density ρ we define the free energy

$$\mathcal{F}(\rho) := \int_{\mathbb{R}^d} U(x)\rho(x)\, dx + \int_{\mathbb{R}^d} \rho(x) \log \rho(x)\, dx. \tag{1}$$

The first term is potential energy. The second is minus the Shannon entropy, so $\mathcal{F}$ is the natural energy–entropy tradeoff for Langevin diffusion.

Proposition 2.1 (Relative entropy equals the free-energy gap)**.** *For every probability density ρ with finite free energy,*

$$D_{\mathrm{KL}}(\rho \| \pi) = \mathcal{F}(\rho) - \mathcal{F}(\pi). \tag{2}$$

In particular,

$$\mathcal{F}(\pi) = -\log Z,$$

so minimizing free energy is the same as minimizing relative entropy to π.

Proof. Since $\log \pi(x) = -U(x) - \log Z$,

$$\begin{aligned} D_{\mathrm{KL}}(\rho \| \pi) &= \int \rho(x) \log \frac{\rho(x)}{\pi(x)}\, dx \\ &= \int \rho(x) \log \rho(x)\, dx + \int U(x)\rho(x)\, dx + \log Z \\ &= \mathcal{F}(\rho) + \log Z. \end{aligned}$$

Applying the same identity with $\rho = \pi$ gives

$$0 = D_{\mathrm{KL}}(\pi \| \pi) = \mathcal{F}(\pi) + \log Z,$$

so $\mathcal{F}(\pi) = -\log Z$. Substituting this into the first formula yields (2). ■

Proposition 2.1 is the basic dictionary for the rest of the note. The object that decreases along Langevin is not an ad hoc Lyapunov function. It is exactly the distance to equilibrium measured in relative entropy.

Exercise 2.2 (Deriving the free-energy identity)**.** Starting from Equation (1) and $\pi(x) = Z^{-1} e^{-U(x)}$, derive Equation (2). Then show directly that π minimizes $\mathcal{F}$ over all densities.

Workspace

2.2 Langevin as a continuity equation

The overdamped Langevin diffusion is

$$dX_t = -\nabla U(X_t)\,dt + \sqrt{2}\,dB_t. \tag{3}$$

If ρ_t denotes the density of X_t, then ρ_t solves the Fokker–Planck equation

$$\partial_t \rho_t = \nabla \cdot (\rho_t \nabla U) + \Delta \rho_t. \tag{4}$$

The key rewrite is that the diffusion term can be absorbed into the logarithmic derivative of ρ_t.

Proposition 2.3 (Continuity-equation form of Fokker–Planck)**.** *Assume ρ_t is smooth and strictly positive. Then (4) can be written as*

$$\partial_t \rho_t = \nabla \cdot \big(\rho_t \nabla (U + \log \rho_t)\big) = -\nabla \cdot (\rho_t v_t), \tag{5}$$

with velocity field

$$v_t = -\nabla\big(U + \log \rho_t\big). \tag{6}$$

Proof. Because $\rho_t > 0$,

$$\Delta \rho_t = \nabla \cdot (\nabla \rho_t) = \nabla \cdot \big(\rho_t \nabla \log \rho_t\big).$$

Substituting this into (4) gives

$$\partial_t \rho_t = \nabla \cdot \big(\rho_t \nabla U\big) + \nabla \cdot \big(\rho_t \nabla \log \rho_t\big) = \nabla \cdot \big(\rho_t \nabla (U + \log \rho_t)\big).$$

Defining v_t by (6) yields the continuity equation in (5). ■

The field $U + \log \rho_t$ is the first variation of free energy, up to an irrelevant additive constant. So Proposition 2.3 says that mass moves downhill in free energy. What it does not yet give is a one-step rule. If we freeze time and ask which nearby density should count as the next descent step, we need to say how expensive it is to move one probability law to another. That is the role of the JKO scheme.



Figure 1: One JKO step balances two competing effects. Moving farther toward equilibrium lowers the free-energy term, but it also increases the Wasserstein movement penalty, so the minimizer of the total objective stops at an interior compromise rather than jumping all the way to π.

3 A variational one-step descent rule

Free energy tells us what should decrease, but we still need a rule for how to choose the next density from the current one. Starting from ρ_k, we want a new law that lowers $\mathcal{F}$ without moving mass arbitrarily far in a single step. The Jordan–Kinderlehrer–Otto (JKO) construction is the variational rule that makes that tradeoff precise.

The analogy from Euclidean optimization is the proximal form of implicit Euler. For the ODE $\dot{x} = -\nabla F(x)$,

$$x_{k+1} \in \arg\min_x \frac{1}{2h}\|x - x_k\|_2^2 + F(x).$$

The JKO step does the same thing for densities. It replaces the Euclidean quadratic penalty by a transport penalty between probability measures. This is the point where the optimization and sampling viewpoints literally meet. The same implicit-step logic used to stabilize descent methods in optimization becomes a rule for evolving *laws* in sampling, with Wasserstein distance replacing Euclidean distance and free energy replacing the objective function.

For a time step $h > 0$, the JKO update is

$$\rho_{k+1} \in \arg\min_{\rho \in \mathcal{P}_2(\mathbb{R}^d)} \left\{ \frac{1}{2h} W_2^2(\rho, \rho_k) + \mathcal{F}(\rho) \right\}. \tag{7}$$

The first term penalizes moving too far from the current density. The second rewards lower free energy. Figure 1 is the picture to keep in mind: JKO is not "jump to the minimizer of $\mathcal{F}$"; it is a one-step compromise between transport cost and dissipation.

3.1 A Gaussian JKO calculation

Take a quadratic target

$$U(x) = \frac{1}{2}x^\top Hx, \qquad H \succ 0,$$

so the equilibrium law is $\pi = \mathcal{N}(0, H^{-1})$. To see what (7) is doing, first restrict attention to the translated Gaussian family

$$\rho_m := \mathcal{N}(m, H^{-1}), \qquad m \in \mathbb{R}^d.$$

The covariance already matches the target, so only the mean is moving.

Proposition 3.1 (Gaussian JKO mean update)**.** *Fix $m_k \in \mathbb{R}^d$ and let $\rho_k = \rho_{m_k}$. If we minimize the JKO objective (7) over translated Gaussians ρ_m, then*

$$m_{k+1} = (I_d + hH)^{-1} m_k. \tag{8}$$

Equivalently, the mean follows implicit Euler for the ODE $\dot{m} = -Hm$.

Proof. All densities ρ_m have the same covariance, so their entropy terms are equal. By Proposition 2.1,

$$\mathcal{F}(\rho_m) - \mathcal{F}(\pi) = D_{\mathrm{KL}}(\rho_m \| \pi).$$

For Gaussians with common covariance H^{-1},

$$D_{\mathrm{KL}}(\rho_m \| \pi) = \frac{1}{2} m^\top Hm, \qquad W_2^2(\rho_m, \rho_{m_k}) = \|m - m_k\|_2^2.$$

Therefore minimizing (7) over the translated family is exactly

$$m_{k+1} \in \arg\min_{m \in \mathbb{R}^d} \left\{ \frac{1}{2h}\|m - m_k\|_2^2 + \frac{1}{2} m^\top Hm \right\}.$$

The first-order condition is

$$\frac{1}{h}(m_{k+1} - m_k) + Hm_{k+1} = 0,$$

which rearranges to (8). ■

This is already a real calculation, not a slogan. JKO reproduces the correct mean evolution and it does so through an *implicit* step. That implicit character matters. If $H = Q\Lambda Q^\top$ with $\Lambda = \mathrm{diag}(\lambda_1, \dots, \lambda_d)$ and $u_k := Q^\top m_k$, then

$$u_{k+1,i} = \frac{1}{1 + h\lambda_i} u_{k,i}, \qquad i = 1, \dots, d. \tag{9}$$

Every mode contracts, and stiffer modes contract more aggressively. There is no explicit-Euler stability restriction of the form $h < 2/\lambda_{\max}$ in this mean calculation. That is one reason minimizing movements are such a natural analytic tool.

Figure 2 makes the same point visually. The left panel shows the mean locations of those translated-Gaussian minimizers. Because the covariance is fixed inside this restricted family, tracking the means is enough to see the entire update. The right panel shows the mode-by-mode damping from (9).



Figure 2: For a quadratic target, restricting JKO to translated Gaussians keeps the covariance fixed and moves only the mean. That mean update is exactly implicit Euler, and in diagonal coordinates each mode contracts by $(1 + h\lambda_i)^{-1}$, so stiff directions settle faster.

Exercise 3.2 (Computing the Gaussian JKO mean step)**.** Work in the translated Gaussian family from Proposition 3.1.

1. Show that $W_2^2(\rho_m, \rho_{m_k}) = \|m - m_k\|_2^2$.
2. Show that $\mathcal{F}(\rho_m) - \mathcal{F}(\pi) = \frac{1}{2} m^\top H m$.
3. Derive (8) and the mode-by-mode contraction formula (9).

Workspace

3.2 The implicit velocity law

The mean calculation shows the basic mechanism in one tractable family, but the full measure-space update has an equally important first-order condition. If ρ_{k+1} is a smooth positive minimizer of (7) and T_{k+1} is the optimal transport map from ρ_{k+1} to ρ_k, then the Euler–Lagrange condition can be written as

$$\frac{\mathrm{id} - T_{k+1}}{h} = -\nabla\big(U + \log \rho_{k+1}\big) \qquad \rho_{k+1}\text{-a.e.} \tag{10}$$

see Santambrogio (2016, §4.5, Prop. 4.16). This is exactly the measure-space analogue of the implicit Euler equation

$$\frac{x_{k+1} - x_k}{h} = -\nabla F(x_{k+1}).$$

The transport displacement per unit time equals the free-energy velocity, evaluated at the *new* density.

3.3 The general JKO theorem

The full theorem is much deeper than the Gaussian example, and we do not try to reproduce the compactness proof here. For our purposes, the conclusion is the key point.

Theorem 3.3 (JKO scheme for Fokker–Planck, cited)**.** *In the regular setting treated in Santambrogio (2016, §4.5, Eq. (4.13), Prop. 4.14, Prop. 4.19, pp. 37–42), each minimization problem in (7) is well posed, and as $h \downarrow 0$ the recursive minimizers admit piecewise constant or geodesic interpolants that converge to the solution of the corresponding Fokker–Planck equation with the prescribed initial density. In this sense, the Fokker–Planck equation is the W_2-gradient flow of free energy.*

The cited proof is written on a compact domain with a C^1 confinement potential. For the Langevin setting on $\mathbb{R}^d$, one imposes the corresponding confinement and integrability assumptions to recover the same minimizing-movement picture. That theorem is the mathematical heart of the note. It upgrades the formal continuity-equation picture from Proposition 2.3 into a precise variational statement: Langevin is not

merely a diffusion with invariant law π; it is the steepest descent of $\mathcal{F}$ in Wasserstein geometry. The right way to read the theorem is as an infinite-dimensional implicit-Euler convergence statement. For each fixed h, the JKO rule produces a discrete descent path of densities. As the step size shrinks, those variational steps converge to the actual Fokker–Planck evolution. So the proximal picture is not just an analogy borrowed from optimization language; it is a literal construction of the PDE.

4 An exact decay identity

JKO identifies the geometry of the flow. The next step is to compute an exact time derivative. That is what turns the variational picture into a quantitative convergence argument.

4.1 The instantaneous decay rate

The quantity that records this instantaneous decay rate is the *relative Fisher information.* It is defined by

$$I(\rho\|\pi) := \int_{\mathbb{R}^d} \|\nabla \log \frac{\rho}{\pi}\|_2^2 \, \rho \, dx. \tag{11}$$

Because $\log \pi = -U - \log Z$, this can also be written as

$$I(\rho\|\pi) = \int \|\nabla(U + \log \rho)\|_2^2 \, \rho \, dx.$$

It measures the squared size of the free-energy slope.

Proposition 4.1 (Entropy dissipation identity)**.** *Let $(\rho_t)_{t\geq 0}$ solve the Langevin Fokker–Planck equation. Under the smoothness and decay assumptions needed for integration by parts,*

$$\frac{d}{dt} D_{\mathrm{KL}}(\rho_t\|\pi) = -I(\rho_t\|\pi). \tag{12}$$

Proof. Using mass conservation, $\int \partial_t \rho_t \, dx = 0$, we may differentiate the entropy functional as

$$\frac{d}{dt} D_{\mathrm{KL}}(\rho_t\|\pi) = \int \log \frac{\rho_t}{\pi} \, \partial_t \rho_t \, dx.$$

From Proposition 2.3,

$$\partial_t \rho_t = \nabla \cdot \left(\rho_t \nabla \log \frac{\rho_t}{\pi} \right).$$

Therefore

$$\begin{aligned} \frac{d}{dt} D_{\mathrm{KL}}(\rho_t\|\pi) &= \int \log \frac{\rho_t}{\pi} \nabla \cdot \left(\rho_t \nabla \log \frac{\rho_t}{\pi} \right) dx \\ &= -\int \|\nabla \log \frac{\rho_t}{\pi}\|_2^2 \rho_t \, dx \\ &= -I(\rho_t\|\pi), \end{aligned}$$

which is (12). ∎

Proposition 4.1 is the exact dynamic counterpart of Proposition 2.1. Relative entropy is the gap to equilibrium, and Fisher information is the instantaneous rate at which Langevin closes that gap.

Exercise 4.2 (Deriving entropy dissipation)**.** Starting from Equation (5), prove Equation (12). Then explain why $I(\rho_t\|\pi) = 0$ forces $\rho_t = \pi$ under mild connectedness assumptions.

Workspace

4.2 An exact Ornstein–Uhlenbeck example

The entropy dissipation identity is completely explicit for Gaussian targets. Let

$$U(x) = \frac{1}{2}x^\top Hx, \qquad H \succ 0, \qquad \pi = \mathcal{N}(0, H^{-1}).$$

Suppose the initial law is

$$\rho_0 = \mathcal{N}(m_0, H^{-1}),$$

so the covariance already matches equilibrium. Then the Langevin diffusion preserves Gaussianity and only the mean moves:

$$\rho_t = \mathcal{N}(m_t, H^{-1}), \qquad m_t = e^{-Ht}m_0.$$

Writing $m_{0,i}$ in an eigenbasis of H with eigenvalues λ_i gives

$$D_{\mathrm{KL}}(\rho_t\|\pi) = \frac{1}{2}\sum_{i=1}^{d}\lambda_i m_{0,i}^2 e^{-2\lambda_i t}, \qquad I(\rho_t\|\pi) = \sum_{i=1}^{d}\lambda_i^2 m_{0,i}^2 e^{-2\lambda_i t}. \tag{13}$$

Differentiating the first display immediately yields

$$\frac{d}{dt}D_{\mathrm{KL}}(\rho_t\|\pi) = -I(\rho_t\|\pi).$$

This example is worth keeping in mind because every later result here collapses to an exact mode-by-mode statement in this quadratic setting.

5 When decay becomes a uniform rate

Entropy dissipation by itself is only an identity. It tells us the instantaneous decay rate,

$$\frac{d}{dt}D_{\mathrm{KL}}(\rho_t\|\pi) = -I(\rho_t\|\pi),$$

but not whether that rate is uniformly large when the chain is still far from equilibrium. In principle the Fisher information could be small even while the entropy gap is still substantial. To turn dissipation into a quantitative rate, we need an inequality that forces a large entropy gap to come with a large slope. The standard tool is the log-Sobolev inequality, and the Bakry–Émery criterion is the curvature condition that often guarantees it.

5.1 A functional inequality that closes the estimate

A log-Sobolev inequality compares the entropy gap directly to its dissipation. Because

$$I(\rho\|\pi) = \int \|\nabla \log(\rho/\pi)(x)\|_2^2 \, \rho(x)\, dx,$$

the Fisher information is the squared $L^2(\rho)$ size of the relative log-density gradient. Geometrically, it measures how sharply the density ratio ρ/π is still tilted across space. The log-Sobolev inequality says that a distribution cannot keep a large free-energy gap while that relative tilt is uniformly tiny. Equivalently, a large mismatch from equilibrium must leave enough slope for Langevin to keep pushing downhill.

Definition 5.1 (Log-Sobolev inequality)**.** We say that π satisfies a log-Sobolev inequality with constant $\alpha > 0$ if

$$D_{\mathrm{KL}}(\rho\|\pi) \le \frac{1}{2\alpha} I(\rho\|\pi) \qquad \text{for every smooth density } \rho. \tag{14}$$

Proposition 5.2 (LSI implies exponential KL decay)**.** *If π satisfies (14) with constant $\alpha > 0$, then every sufficiently regular Langevin solution satisfies*

$$D_{\mathrm{KL}}(\rho_t\|\pi) \le e^{-2\alpha t} D_{\mathrm{KL}}(\rho_0\|\pi). \tag{15}$$

Proof. By Proposition 4.1,

$$\frac{d}{dt} D_{\mathrm{KL}}(\rho_t\|\pi) = -I(\rho_t\|\pi).$$

Applying the log-Sobolev inequality at time t gives

$$I(\rho_t\|\pi) \ge 2\alpha D_{\mathrm{KL}}(\rho_t\|\pi),$$

hence

$$\frac{d}{dt} D_{\mathrm{KL}}(\rho_t\|\pi) \le -2\alpha D_{\mathrm{KL}}(\rho_t\|\pi).$$

Gronwall's inequality yields (15). ■

This is the precise place where "strong log-concavity implies quantitative mixing" becomes a theorem. Once Fisher information dominates entropy, the free-energy gap decays exponentially. The geometric message is that under an LSI the target has no extended almost-flat directions in which a nonequilibrium density can hide. If ρ_t still differs substantially from π, then the relative log-density must still have noticeable slope, so the Fokker–Planck flow cannot stall prematurely.

5.2 The Gaussian benchmark

For the Gaussian target

$$\gamma_m(dx) \propto e^{-\frac{m}{2}\|x\|_2^2}\,dx,$$

the correct LSI constant is $\alpha = m$. One way to see that is to invoke the curvature criterion in the next subsection, since $\nabla^2 U = mI_d$. Inside the translated Gaussian family, the constant is visible exactly: if $\rho = \mathcal{N}(a, m^{-1}I_d)$, then

$$D_{\mathrm{KL}}(\rho\|\gamma_m) = \frac{m}{2}\|a\|_2^2, \qquad I(\rho\|\gamma_m) = m^2\|a\|_2^2,$$

so equality holds in (14). That is a useful sanity check. The Gaussian measure is not merely an example where LSI is true; it is the model case where the constants are exact.

5.3 Curvature gives LSI

The next question is where a bound like (14) comes from. For strongly log-concave targets, the answer is curvature. If U bends upward everywhere by at least a fixed quadratic amount, then the measure has no broad flat directions in which entropy can stay large while the free-energy slope becomes tiny. The Bakry–Émery criterion turns that idea into a global log-Sobolev constant.

Theorem 5.3 (Bakry–Émery criterion, cited). *Let $\pi(dx) = Z^{-1}e^{-U(x)}\,dx$ on $\mathbb{R}^d$, where $U \in C^2(\mathbb{R}^d)$ and $Z < \infty$. If*

$$\nabla^2 U(x) \succeq mI_d \qquad \text{for every } x \in \mathbb{R}^d \tag{16}$$

for some $m > 0$, then π satisfies a log-Sobolev inequality with constant $\alpha = m$.

We cite Otto and Villani (2000, Thm. 2) for this form of the theorem. Historically it is the Bakry–Émery curvature condition. The proof uses Γ_2-calculus rather than the transport arguments used in the surrounding development, so we do not reproduce it.

The geometric meaning of (16) is that the ambient space always looks at least as confining as a quadratic well. If U has a uniform quadratic lower bound on curvature, then the target has no flat directions that allow entropy to linger and no broad corridors in which mass can drift with only a weak restoring force. Misplaced mass must create a visible change in the relative log-density, so the free-energy slope cannot be tiny unless the entropy gap is already small. That is exactly what (14) formalizes. What makes the theorem powerful is that it is global. The lower Hessian bound is required at every point, not just near the mode, and the conclusion controls every smooth density ρ. One can think of m as a universal restoring-force constant: no matter where mass moves, the landscape is always at least as curved as a quadratic well of curvature m. Bakry–Émery says that this geometric fact is enough to force a uniform entropy-dissipation estimate.

6 From entropy to transport

We now have an exponential decay theorem in KL divergence. For proving convergence, that is already enough. The reason to keep going is that KL and Wasserstein distance answer different questions. The KL divergence measures excess free energy, while W_2 measures how far the mass of ρ_t still has to move in space to match π. That geometric viewpoint is closer to the coupling language of the Sampling unit and the conditioning language of the Optimization unit, so it is worth translating entropy control into transport control as well.

Definition 6.1 (Talagrand's T_2 inequality)**.** We say that π satisfies $T_2(\alpha)$ if

$$W_2^2(\rho, \pi) \leq \frac{2}{\alpha} D_{\mathrm{KL}}(\rho \| \pi) \qquad \text{for every density } \rho. \tag{17}$$

Theorem 6.2 (Otto–Villani, cited)**.** *Let $\pi(dx) = Z^{-1} e^{-U(x)}\, dx$ be a probability measure on $\mathbb{R}^d$ with finite second moment, where $U \in C^2(\mathbb{R}^d)$ and*

$$\nabla^2 U(x) \succeq C I_d \qquad \text{for every } x \in \mathbb{R}^d$$

for some $C \in \mathbb{R}$. If π satisfies a log-Sobolev inequality with constant $\alpha > 0$, then π also satisfies $T_2(\alpha)$.

This is Otto and Villani (2000, Thm. 1). It says that once entropy is controlled, the amount of spatial transport still needed to reach equilibrium is controlled automatically. That is not obvious a priori because the two quantities measure different kinds of mismatch. The KL divergence compares density values pointwise, while W_2 measures the cost of rearranging mass in space. The theorem says that an LSI is strong enough to connect them: under sufficient curvature, any large remaining transport cost would itself force a large entropy gap. So transport control is not an extra ingredient added on top of entropy control; it is a geometric consequence of the same inequality.

Corollary 6.3 (Wasserstein decay from LSI)**.** *Under the hypotheses of Theorem 6.2, every sufficiently regular Langevin solution satisfies*

$$W_2(\rho_t, \pi) \leq \sqrt{\frac{2}{\alpha}}\, e^{-\alpha t} \sqrt{D_{\mathrm{KL}}(\rho_0 \| \pi)}. \tag{18}$$

Proof. Combine Theorem 6.2 with Proposition 5.2:

$$W_2^2(\rho_t, \pi) \leq \frac{2}{\alpha} D_{\mathrm{KL}}(\rho_t \| \pi) \leq \frac{2}{\alpha} e^{-2\alpha t} D_{\mathrm{KL}}(\rho_0 \| \pi).$$

Taking square roots gives (18). ■

For the Gaussian family from Section 5.2, the bridge is exact on translated states: if $\rho = \mathcal{N}(a, m^{-1} I_d)$ and $\gamma_m = \mathcal{N}(0, m^{-1} I_d)$, then

$$W_2^2(\rho, \gamma_m) = \|a\|_2^2 = \frac{2}{m} D_{\mathrm{KL}}(\rho \| \gamma_m).$$

So the LSI-to-T_2 story is not merely qualitative. Its constants are already sharp in the Gaussian model.

Exercise 6.4 (From LSI to a Wasserstein bound)**.** Assume π satisfies LSI(α).

1. Use Proposition 4.1 to derive Equation (15).
2. Combine that with Equation (17) to prove Equation (18).
3. Verify directly in the translated Gaussian family that $T_2(\alpha)$ holds with equality.

Workspace

7 Preconditioning and a real posterior

The functional-inequality chain explains why Langevin converges. The final question is where the constants come from in practice. For quadratic or locally quadratic targets, the answer is geometric: the relevant rates are the eigenvalues of the preconditioned Hessian.

7.1 Quadratic targets and the rate matrix

Now return to geometry. Consider the preconditioned overdamped Langevin diffusion

$$dX_t = -M\nabla U(X_t)\,dt + \sqrt{2M}\,dB_t, \tag{19}$$

where $M \succ 0$ is a fixed symmetric positive-definite matrix. For a quadratic potential centered at $x_\star$,

$$U(x) = \frac{1}{2}(x - x_\star)^\top H(x - x_\star), \qquad H \succ 0,$$

the geometry is again exact.

Proposition 7.1 (The relevant rate matrix is $M^{1/2}HM^{1/2}$)**.** *Let $Y_t := M^{-1/2}(X_t - x_\star)$. Then (19) becomes*

$$dY_t = -AY_t\,dt + \sqrt{2}\,dB_t, \qquad A := M^{1/2}HM^{1/2}. \tag{20}$$

Hence the mode-by-mode decay rates are the eigenvalues of A, not of H alone. In particular, if $M = H^{-1}$, then $A = I_d$ and all local modes are equalized.

Proof. Write $X_t = x_\star + M^{1/2}Y_t$. Then

$$dX_t = M^{1/2}dY_t, \qquad \nabla U(X_t) = H(X_t - x_\star) = HM^{1/2}Y_t.$$

Substituting into (19),

$$M^{1/2}dY_t = -MHM^{1/2}Y_t\,dt + \sqrt{2M}\,dB_t.$$



Figure 3: For an anisotropic Gaussian target, KL, Fisher information, and W_2^2 decay exactly under the local Ornstein–Uhlenbeck dynamics. Whitening turns the rate matrix into the identity and removes the slow mode.

Multiplying by $M^{-1/2}$ gives

$$dY_t = -M^{1/2}HM^{1/2}Y_t\,dt + \sqrt{2}\,dB_t,$$

which is (20). ■

The local conclusion is immediate. Whitening is not mysterious. It is exactly the choice of M that turns the local quadratic surrogate into an isotropic Ornstein–Uhlenbeck process.

The entropy and transport quantities from earlier sections are also explicit here. If the transformed process starts from

$$Y_0 \sim \mathcal{N}(m_0, A^{-1}),$$

then

$$Y_t \sim \mathcal{N}(e^{-At}m_0, A^{-1}),$$

so in an eigenbasis of A with eigenvalues ν_i and coordinates c_i of m_0,

$$D_{\mathrm{KL}}(\rho_t\|\pi) = \frac{1}{2}\sum_i \nu_i c_i^2 e^{-2\nu_i t}, \qquad I(\rho_t\|\pi) = \sum_i \nu_i^2 c_i^2 e^{-2\nu_i t}, \qquad W_2^2(\rho_t,\pi) = \sum_i c_i^2 e^{-2\nu_i t}. \tag{21}$$

When $M = H^{-1}$, all ν_i become 1, so every mode decays on the same time scale.

Figure 3 shows (21) for a simple two-dimensional quadratic target. The visual simplification reflects an analytic one. Whitening changes the exact exponents appearing in every convergence quantity derived above.

7.2 Bayesian logistic regression on the Problem Set 2 data

To keep the real-data example aligned with the notes, we return to the regularized logistic-regression model that runs through the Optimization unit and reappears in Problem Set 2. The dataset has five predictors and a binary response. In the optimization material, this model supplied explicit gradients, Hessians, and



Figure 4: The Problem Set 2 Bayesian logistic posterior has a nontrivial MAP Hessian spectrum. Whitening with $H_{\mathrm{MAP}}^{-1/2}$ collapses the local spectrum to 1.

the MAP/Laplace calculation. Here the same local quadratic surrogate serves a different purpose: it reveals the local rate matrix seen by overdamped Langevin near the MAP.

Write

$$\beta = (\beta_0, \beta_1, \dots, \beta_5)^\top, \qquad \tilde{x}_i = (1, x_{i1}, \dots, x_{i5})^\top.$$

We also keep the same isotropic Gaussian prior as in Problem Set 2,

$$\beta \sim \mathcal{N}(0, 100 I_6),$$

so the prior precision is $\lambda = 0.01$. The negative log posterior is

$$U(\beta) = \sum_{i=1}^{n} \left[\log\left(1 + e^{\tilde{x}_i^\top \beta}\right) - y_i \tilde{x}_i^\top \beta \right] + \frac{\lambda}{2} \|\beta\|_2^2. \tag{22}$$

If $p_i(\beta) = \sigma(\tilde{x}_i^\top \beta)$ with $\sigma(t) = 1/(1 + e^{-t})$, then

$$\nabla U(\beta) = X^\top (p(\beta) - y) + \lambda \beta, \qquad \nabla^2 U(\beta) = X^\top W(\beta) X + \lambda I_6,$$

where $W(\beta)$ is diagonal with entries $p_i(\beta)(1 - p_i(\beta))$.

We compute the MAP estimate β_{MAP} numerically and use the Hessian

$$H_{\mathrm{MAP}} := \nabla^2 U(\beta_{\mathrm{MAP}})$$

as a local Gaussian surrogate. This is exactly the Laplace approximation, and we use it only locally. The claim is not that the posterior is globally Gaussian. The claim is that the Hessian near the MAP reveals which local preconditioner equalizes the quadratic surrogate.

Figure 4 shows the full Hessian spectrum. The raw local condition number is about 5.8. That is not extreme, but it is large enough to make the soft/stiff splitting visible in a familiar course example. That spectrum is



Figure 5: The softest/stiffest eigenspace of the Laplace surrogate around the MAP. Without preconditioning, local mean paths pinch first along the stiff direction and only later move along the soft direction; local whitening equalizes the contraction rates.

the continuous-time Langevin analogue of the conditioning language from the Optimization unit and the Sampling unit's gradient-based MCMC subsection.

Figure 5 turns the spectrum into a local picture. Because this model has only six parameters, the figure uses the softest and stiffest eigendirections directly. The message is the same in either case. Without preconditioning, the local mean paths collapse first in the stiff direction while progress in the soft direction lags behind. With $M = H_{\mathrm{MAP}}^{-1}$, the local quadratic surrogate has balanced rates.

This is the right closing connection to the course's conditioning language. For overdamped Langevin, preconditioning changes the exact rate matrix controlling entropy dissipation and transport contraction. The same local Hessian picture is what later makes mass-matrix choices matter in the Augmented-state sampling unit's HMC/NUTS development.

Exercise 7.2 (Reading the local Hessian geometry)**.** Use Proposition 7.1, Figure 4, and Figure 5.

1. Explain why a spread in the eigenvalues of H_{MAP} produces multiple local mixing time scales for isotropic Langevin.
2. Show that choosing $M = H_{\mathrm{MAP}}^{-1}$ makes the local rate matrix equal to the identity.
3. Why is H_{MAP}^{-1} the natural quadratic preconditioner, up to an overall scalar factor?

Workspace

There is also a discrete-time algorithmic payoff. Once one has an LSI and enough smoothness, these continuous-time decay arguments feed directly into nonasymptotic ULA theory; see Vempala and Wibisono (2019, §3.1, Thm. 1). The point of that cited result is not that Euler discretization exactly preserves the diffusion analysis. Rather, good continuous-time geometry survives discretization up to a controlled error, so the same inequalities lead to quantitative bounds for an actual algorithm. The course-level synthesis is now clear. Free energy is the Lyapunov functional, JKO is the proximal structure, Fisher information supplies dissipation, LSI turns dissipation into a uniform rate, Talagrand converts that rate into transport control, and preconditioning changes the constants by changing the local geometry. Carried back to the main notes, this is the optimization story returning in measure space: descent becomes entropy dissipation, proximal structure becomes JKO, and conditioning still determines which directions are easy and which are slow.

References

Felix Otto and Cédric Villani. Generalization of an inequality by talagrand and links with the logarithmic sobolev inequality. *Journal of Functional Analysis*, 173(2):361–400, 2000.

Filippo Santambrogio. Euclidean, metric, and wasserstein gradient flows: An overview, 2016.

Santosh S. Vempala and Andre Wibisono. Rapid convergence of the unadjusted langevin algorithm: Isoperimetry suffices, 2019.