

# Usage for **genulens**

Naoki Koshimoto

Email: koshimoto.work@gmail.com

GitHub: <https://github.com/nkoshimoto/genulens>

Paper for the Galactic model: Koshimoto, Baba, and Bennett (2021), arXiv: 2104.03306

May 25, 2021

To use **genulens**, you need to specify input parameter values or options in the arguments.

A simplest example is

(Example 1)

```
$/genulens Nsimu 1e+5 l 2.32 b -2.38 PA 60.31 vEarthlb 1.10 -6.71 VERBOSITY 2 > tmp.dat
```

This generates  $10^5$  ( $=$  Nsimu) artificial microlensing events toward the sky direction of  $(l, b) = (2.32^\circ, -2.38^\circ)$ . It uses the E+E<sub>X</sub> model in Koshimoto, Baba, and Bennett (2021) [1] as the Galactic model by default. There are other three Galactic models developed in the paper, and you can use each of the E, G and G+G<sub>X</sub> models by adding “E\_fg0 1”, “G\_fg0 1”, “GXG\_fg0 1” to the arguments, respectively.

The output format can be specified by the parameter VERBOSITY, and Fig. 1 shows part of output lines of example 1. With “VERBOSITY 2”, the output format is (from left to right):

$wt_j$  – Weight for  $j$ th line. **You need to weight each line by  $wt_j$**  when you combine them to make a probability distribution.  $wt_j$  takes mostly 1 and sometimes  $> 1$  unless you use the “SMALLGAMMA 1” option.

$t_E$  – The Einstein radius crossing time in days.

$\theta_E$  – The angular Einstein radius in mas.

$\pi_{E,N}$  – Equatorial north component of the microlens parallax vector.

$\pi_{E,E}$  – Equatorial east component of the microlens parallax vector.

$D_S$  – Distance to the source system in pc.

$\mu_{S,l,hel}$  – Galactic longitudinal component of the source proper motion in the heliocentric frame in mas/yr.

$\mu_{S,b,hel}$  – Galactic latitudinal component of the source proper motion in the heliocentric frame in mas/yr.

$i_S$  – Flag to indicate which component the source system belongs to. The components are:

- 0: Thin disk (0.00 - 0.15 Gyr)
- 1: Thin disk (0.15 - 1.00 Gyr)
- 2: Thin disk (1.00 - 2.00 Gyr)
- 3: Thin disk (2.00 - 3.00 Gyr)
- 4: Thin disk (3.00 - 5.00 Gyr)
- 5: Thin disk (5.00 - 7.00 Gyr)
- 6: Thin disk (7.00 - 10.00 Gyr)
- 7: Thick disk (12.0 Gyr)
- 8: Bulge ( $9 \pm 1$  Gyr)

$i_L$  – Flag to indicate which component the lens system belongs to.

$f_{REM}$  – Flag to indicate evolutionary stage of the lens.

- 0: Main sequence
- 1: White dwarf
- 2: Neutron star
- 3: Black hole

Note that  $f_{REM}$  can be  $> 0$  only when the “REMNANT 1” or “onlyWD 1” option is added to the argument, and can be  $> 1$  only when the “REMNANT 1” option is added.

```

#----- Output of Monte Carlo simulation w/ VERBOSITY= 2 and seed= 12304357 -----
#      wtj      tE      thetaE      piEN      piEE      D_S      muSl      muSb      iS      iL      fREM
1.789982e+00 1.026014e+01 2.893354e-01 -2.670026e-02 4.126398e-02 8159 -6.23573e+00 4.13179e+00 8 8 0
1.000000e+00 1.449606e+01 2.533772e-01 2.800766e-01 2.697542e-01 13858 -6.68344e+00 -1.68848e-01 6 3 0
1.000000e+00 6.392391e+01 2.665412e-01 -2.151892e-02 -4.385056e-02 7157 -2.54691e+00 -1.72749e+00 8 8 0
1.000000e+00 4.701406e+00 1.194560e-01 4.380464e-02 3.561815e-02 8184 -1.55973e+01 2.67340e+00 8 8 0
1.000000e+00 1.739493e+01 2.549459e-01 9.644269e-02 -4.377408e-03 8763 -6.70109e+00 -1.46376e+00 8 8 0
1.000000e+00 5.888771e+00 1.094510e-01 6.294772e-02 -2.766520e-02 7904 -8.48463e+00 -1.29959e+00 8 8 0

```




Figure 1: Part of the output lines of example 1 (top) and the probability distributions for each output parameter (bottom).

In example 1, PA is the position angle in degree, which is the angle between the equatorial east direction and the galactic longitudinal direction, on the sky plane. The two values after `vEarthlb` are the galactic longitudinal and latitudinal components of the Earth velocity projected onto the sky plane in km/sec. PA and `vEarthlb` do not have to be specified if you are just interested in the magnitude of the microlens parallax vector, and not interested in its direction. Technically, `vEarthlb` affects  $t_E$  value because it is used to convert  $t_{E, \text{hel}} (= \theta_E / \mu_{\text{rel, hel}})$  into  $t_E (= \theta_E / \mu_{\text{rel, geo}})$ . However, in most cases, the difference in the resulted probability distributions of parameters other than the direction of the microlens parallax vector is very small between when you specify `vEarthlb` and when you do not specify it.

The output parameters with “`VERBOSITY 2`” are sufficient to calculate other (perhaps more interesting) microlensing parameters such as the lens mass and distance. Appendix A provides the formulae for the conversion.

## 1 Posterior Calculation

For an individual event analysis,  $t_E$  is commonly measured, and  $\theta_E$  is mostly measured for a caustic-crossing event. A frequent situation is that you want to calculate the posterior probability distributions for the lens mass and distance by incorporating these constraints from observed parameters into the Galactic prior.

Let me use OGLE-2008-BLG-355 (Koshimoto et al. 2014 [2]) to show how to calculate a posterior distribution with `genulens`. You have measured  $t_E = 34.0 \pm 2.2$  days and  $\theta_E = 0.28 \pm 0.03$  mas for this event occurred at  $(l, b) = (-0.08^\circ, -3.45^\circ)$ .

You can calculate the posterior probability distribution by

(Example 2)

```

$./genulens NlikeMIN 1e+5 1 -0.08 b -3.45 tE 34.0 2.2 thetaE 0.28 0.03 VERBOSITY 3 > tmp.dat

```

and Fig. 2 shows part of the output lines (with `VERBOSITY = 3`) and the distribution for some parameters. The “`VERBOSITY 3`” option is more redundant than “`VERBOSITY 2`”, and it outputs  $M_L$ ,  $D_L$ ,  $\pi_E$ ,  $\mu_{\text{rel, geo}}$ , and  $I_L$ , in addition to the output parameters with `VERBOSITY 2`. In example 2, I do not specify PA or `vEarthlb` assuming that we are not interested in the direction of  $\pi_E$  vector now.

The “`tE [val] [err]`” option multiplies a Gaussian probability of

$$\mathcal{L}_{t_E} = \exp \left( -0.5 \left( \frac{\text{val} - t_E}{\text{err}} \right)^2 \right) \quad (1)$$

```

#----- Output of Monte Carlo simulation w/ VERBOSITY= 3 and seed= 12304357 -----
#      wtj      M_L      D_L      D_S      t_E      theta_E      pi_E      pi_EN      pi_EE      mu_rel      mu_Sl      mu_Sb      I_L  iS  iL  fREM
1.090589e+00 6.073099e-01 9798 11270 3.337356e+01 2.567712e-01 5.191626e-02 1.268154e-02 5.034359e-02 2.810178e+00 -8.36728e+00 1.97348e+00 99.000 7 8 0
1.000000e+00 8.924341e-01 7467 8010 3.675385e+01 2.568369e-01 3.533856e-02 -2.433739e-02 -2.562236e-02 2.552377e+00 -4.00786e+00 -2.02992e-01 99.000 8 8 0
1.000000e+00 5.172451e-01 7701 9233 3.254831e+01 3.012192e-01 7.150788e-02 4.492107e-02 5.563698e-02 3.380217e+00 -7.02563e+00 -3.33934e-01 99.000 4 8 0
1.000000e+00 4.967772e-01 8126 9339 3.581116e+01 2.542376e-01 6.284139e-02 -1.548968e-02 -6.090247e-02 2.593054e+00 -4.58248e+00 -6.41778e-01 99.000 8 6 0
1.000000e+00 5.501188e-01 6733 7838 3.407724e+01 3.062447e-01 6.835647e-02 -4.535272e-02 -4.831792e-02 3.282422e+00 -3.57722e+00 -4.30568e+00 99.000 8 8 0
1.000000e+00 7.642087e-01 7927 8963 3.233027e+01 3.013273e-01 4.841657e-02 -3.061996e-02 -3.750443e-02 3.404233e+00 -6.64013e+00 1.77290e+00 99.000 8 8 0

```




Figure 2: Part of the output lines of example 2 (top) and the probability distributions for some parameters (bottom). Note that the lens apparent magnitude in  $I$ -band,  $I_L$ , only takes 99 unless the extinction value for red clump centroid  $A_{I,rc}$  is specified by `AIRC`.

by the Galactic prior by default, while it multiplies a uniform probability of

$$\mathcal{L}_{t_E} = \begin{cases} 1 & \text{when } |\text{val} - t_E| \leq \text{err} \\ 0 & \text{when } |\text{val} - t_E| > \text{err} \end{cases} \quad (2)$$

by the Galactic prior when “UNIFORM 1” is added to the arguments. The same is true for the “`thetaE [val] [err]`” option which multiplies another probability of  $\mathcal{L}_{\theta_E}$ .

In practice, when you add the `tE` and `thetaE` constraints, `genulens` generates many microlensing events following the Galactic model, and outputs its parameters only if the condition “ $P_{\text{ran}} < \mathcal{L}_{t_E} \mathcal{L}_{\theta_E}$ ” is true, where  $P_{\text{ran}}$  is a random number generated from the uniform distribution between 0 and 1.

Example 2 outputs  $\sim 10^5$  ( $= \text{NlikeMIN}$ ) lines that passed the condition “ $P_{\text{ran}} < \mathcal{L}_{t_E} \mathcal{L}_{\theta_E}$ ”, which means the number of generated events are much larger than `NlikeMIN`. This is the difference between `NlikeMIN` and `Nsimu` used in example 1. If you use “`Nsimu 1e+5`” instead of “`NlikeMIN 1e+5`” in example 2, it would generate  $10^5$  microlensing events, and would output parameters for small fraction of them that passed the condition.

The “`NlikeMIN`” option is useful, but you need to be careful when using it because you cannot easily tell the amount of time it will take unlike the “`Nsimu`” option. I recommend that you always check the computational time with a small `NlikeMIN` value before you try a calculation with a large `NlikeMIN` value, because the computational time with a certain `NlikeMIN` value totally depends on the constraints you used for `tE`, `thetaE`, or other parameters. In the case of example 2, it takes  $\sim 80$  secs with my laptop to generate  $1.4 \times 10^7$  events and accept  $\sim 10^5$  events out of them.

In general, if you find that your desired calculation would take too long, you can divide the calculation into multiple runs with different random seed values using the `seed` option. For example, you can run

```
./genulens [options for calculation] seed 1 > tmp1.dat
```

```
./genulens [options for calculation] seed 2 > tmp2.dat
```

```
.
```

```
.
```

```
./genulens [options for calculation] seed 50 > tmp50.dat
```

in parallel if you have sufficient cores in your machine(s), and combine the output files to make a posterior probability distribution.

## 1.1 How to incorporate the parameter distribution from the MCMC calculation

In example 2, I incorporated the measured  $t_E$  and  $\theta_E$  values assuming they are independent Gaussian distributions. Ideally, we want to convolve the Galactic prior with a posterior distribution from a MCMC calculation for the light curve data to include correlations among parameters and non-Gaussian shape of parameter distributions.

Currently, `genulens` incorporates the observed constraints in either a Gaussian probability form (Eq. 1) or a uniform probability form (Eq. 2), and cannot directly incorporate the MCMC chain distribution. An alternative way, in the case of OGLE-2008-BLG-355, is to conduct a similar calculation to example

2, but with the “UNIFORM 1” option and larger error bars for  $t_E$  and  $\theta_E$ , e.g.,

(Example 3)

```
./genulens NlikeMIN 1e+6 1 -0.08 b -3.45 tE 34.0 9.0 thetaE 0.28 0.12 VERBOSITY 2
```

```
UNIFORM 1 > tmp.dat
```

Separately, you should be able to calculate a joint probability distribution of  $t_E$  and  $\theta_E$ ,  $P_{LC}(t_E, \theta_E)$ , from your MCMC calculation for the light curve data. By multiplying  $P_{LC}(t_E, \theta_E)$  by  $wt_j$  in each line of the output of example 3, you can calculate a posterior probability distribution for  $t_E$  and  $\theta_E$  (and other output parameters) including the correlation between  $t_E$  and  $\theta_E$ , and non-Gaussian shape of them.

## 2 Input Parameters

Coming soon.

## A Formulae for Conversion

The following summarizes formulae to convert the output parameters with `VERBOSITY = 2` into other common parameters used in microlensing studies.

$$\pi_E = \sqrt{\pi_{E,N}^2 + \pi_{E,E}^2} \quad (3)$$

$$M_L = \frac{\theta_E}{\kappa \pi_E} \quad (\kappa = 8.144 \text{ mas } M_\odot^{-1}) \quad (4)$$

$$D_L = \frac{\text{AU}}{\pi_E \theta_E + \text{AU}/D_S} \quad (5)$$

$$\boldsymbol{\mu}_{\text{rel,geo}} = \frac{\theta_E}{t_E} \frac{\boldsymbol{\pi}_E}{\pi_E} \quad (6)$$

$$\text{or } \begin{pmatrix} \mu_{\text{rel,geo,N}} \\ \mu_{\text{rel,geo,E}} \end{pmatrix} = \frac{\theta_E}{t_E} \frac{1}{\pi_E} \begin{pmatrix} \pi_{E,N} \\ \pi_{E,E} \end{pmatrix} \quad (7)$$

$$\text{or } \begin{pmatrix} \mu_{\text{rel,geo,b}} \\ \mu_{\text{rel,geo,l}} \end{pmatrix} = \begin{pmatrix} \cos \text{PA} & -\sin \text{PA} \\ \sin \text{PA} & \cos \text{PA} \end{pmatrix} \begin{pmatrix} \mu_{\text{rel,geo,N}} \\ \mu_{\text{rel,geo,E}} \end{pmatrix} \quad (8)$$

$$\boldsymbol{\mu}_{\text{rel,hel}} = \boldsymbol{\mu}_{\text{rel,geo}} + \frac{\pi_E \theta_E}{\text{AU}} \mathbf{v}_{\oplus,\perp} \quad (9)$$

$$\boldsymbol{\mu}_{L,\text{hel}} = \boldsymbol{\mu}_{\text{rel,hel}} + \boldsymbol{\mu}_{S,\text{hel}} \quad (10)$$

$$\mathbf{v}_{L,\perp} = D_L \boldsymbol{\mu}_{\text{rel,hel}} + \mathbf{v}_{\odot,\perp} \quad (11)$$

## References

- [1] Koshimoto, N., Baba, J., & Bennett, D. P. 2021, arXiv:2104.03306
- [2] Koshimoto, N., Udalski, A., Sumi, T., et al. 2014, ApJ, 788, 128
- [3] Nataf, D. M., Gould, A., Fouqué, P., et al. 2013, ApJ, 769, 88. doi:10.1088/0004-637X/769/2/88