

TREE APPROXIMATORS OF A NORMAL

Why

What is the optimal tree approximator of a multivariate normal density?

Result

Prop. 1. Let $g: \mathbb{R}^n \to \mathbb{R}$ be a normal density with mean $\mu \in \mathbb{R}^d$ and covariance $\Sigma \in \mathbb{S}_{++}^d$. The normal density $f_T^*: \mathbb{R}^d \to \mathbb{R}$ with mean μ and precision matrix P defined by

•
$$P_{11} = \Sigma_{11}^{-1} + \sum_{\text{pa}j=1} \Sigma_{j1}^2 \Sigma_{11}^{-2} \Sigma_{j|1}^{-1}$$

• for
$$i = 2, ..., d$$
, $P_{ii} = \sum_{i \mid \text{pa } i}^{-1} + \sum_{\text{pa } j = i}^{2} \sum_{j \mid i}^{2} \sum_{i \mid i}^{-2} \sum_{j \mid i}^{-1}$

•
$$i, j = 1, \dots d$$
 and $i = \text{pa } j, P_{ij} = P_{ji} = -\sum_{ji} \sum_{jj}^{-1} \sum_{j|i}^{-1}$

where pai is the parent of i in an optimal approximator tree T (i = 2, ..., n) is an optimal tree approximator of g.

Proof. Using Proposition 1 of Best Tree Density Approximators, express an optimal tree approximator of g by

$$(1/c) \exp \left(-\frac{1}{2} \left(\Sigma_{11}^{-1} \bar{x}_1^2 + \sum_{i \neq 1} \left(\bar{x}_i - \Sigma_{i, \text{pa } i} \Sigma_{\text{pa } i, \text{pa } i}^{-1} \bar{x}_{\text{pa } i} \right)^2 \Sigma_{i \mid \text{pa } i}^{-1} \right) \right)$$

where
$$\bar{x}_i = x_i - \mu_i$$
 and $c = \sqrt{(2\pi)^d \sum_{11} \prod_{i \neq 1} \sum_{i \mid \text{pa } i}}$.

Second, express the quadratic in the exponential as

$$\Sigma_{11}^{-1} \bar{x}_{1}^{2} + \sum_{i \neq 1} \left[\Sigma_{i \mid \text{pa}\, i}^{-1} \bar{x}_{i}^{2} - 2 \Sigma_{i, \text{pa}\, i} \Sigma_{\text{pa}\, i, \text{pa}\, i}^{-1} \Sigma_{i \mid \text{pa}\, i}^{-1} \bar{x}_{i} \bar{x}_{\text{pa}\, i} + \Sigma_{i, \text{pa}\, i}^{2} \Sigma_{\text{pa}\, i, \text{pa}\, i}^{-2} \Sigma_{i \mid \text{pa}\, i}^{-1} \bar{x}_{\text{pa}\, i}^{2} \right]$$

With P defined as earlier, we can express the above as $\bar{x}^{\top}P\bar{x}$.

Third, note that c is $\sqrt{(2\pi)^d \det P^{-1}}$ since f_T^* is a density and so integrates to one.

Notice that f_T^* is a tree normal density.

Empirical Normal

In particular, notice that we can approximate the empirical normal density of a dataset with a density that factors according to a tree.

