### Why

[1]

## 1 Definition

Let $A \subset \mathbf{R}$ open. Let $f : A \to \mathbf{R}$ be differentiable with derivative $f' : \mathbf{R} \to \mathbf{R}$. We call $f$ *twice differentiable* (or *two times differentiable*) if its derivative $f'$ is differentiable. In this case, we call the derivative of $f'$ the *second derivative* of $f$.

### Notation

Let $A \subset R$. The second derivative of the twice-differentiable function $f : A \to \mathbf{R}$ is sometimes denoted $f''(x) : A \to \mathbf{R}$

[1] Future editions will include.


Second Derivatives
Differentiable Functions
Real Open Sets
N-Dimensional Space
Real Functions
Distance
Space Distance
Plane Distance
Interval Length
Intervals
Real Line
Real Space
Real Plane
Integral Line
Real Order
Real Numbers
Geometry
Rational Numbers
Integer Arithmetic
Sequences
Natural Multiplicative Identity
Natural Additive Identity
Set Numbers
Direct Products
Arithmetic
Identity Elements
Finite Sets
Natural Powers
Integer Products
Operations
Equivalent Sets
Natural Products
Integer Sums
Function Inverses
Family Unions and Intersections
Natural Sums
Integer Numbers
Function Composites
Function Images
Families
Recursion Theorem
Natural Order
Equivalence Relations
Functions
Peano Axioms
Relations
Natural Induction
Set Products
Natural Numbers
Set Powers
Generalized Set Dualities
Partitions
Intersection of Empty Set
Successor Sets
Unordered Triples
Set Dualities
Set Unions and Intersections
Ordered Pairs
Set Intersections
Pair Unions
Set Complements
Set Unions
Set Differences
Pair Intersections
Unordered Pairs
Empty Set
Set Specification
Set Inclusion
Standardized Accounts
Accounts
Deductions
Quantified Statements
Logical Statements
Statements
Identities
Sets
Names
Letters
Objects