

# ENHANCING BRAIN CONNECTIVITY MAPPING: A RADIOMICS APPROACH TO DIFFUSION MRI TRACTOGRAPHY

LEVENTE ZSOLT NAGY

**Thesis supervisor**

ALFREDO VELLIDO ALCACENA (Department of Computer Science)

**Thesis co-supervisor**

ESTELA CAMARA MANCHA (Hospital Universitari de Bellvitge)

**Degree**

Master's Degree in Artificial Intelligence

**Master's thesis**

**School of Engineering**

Universitat Rovira i Virgili (URV)

**Faculty of Mathematics**

Universitat de Barcelona (UB)

**Barcelona School of Informatics (FIB)**

Universitat Politècnica de Catalunya (UPC) - BarcelonaTech

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>Fulfillment of the Objectives</b>	<b>5</b>
2.1	Objectives . . . . .	5
2.2	Motivation . . . . .	5
2.3	Experiments . . . . .	5
<b>3</b>	<b>Adjustments Made</b>	<b>8</b>
<b>4</b>	<b>Work Plan for Completion</b>	<b>9</b>

# List of Notations & Abbreviations

**MRI** magnetic resonance imaging ..... 3

**dMRI** diffusion magnetic resonance imaging..... 3

**FA** fractional anisotropy ..... 3

**MD** mean diffusivity ..... 3

**RD** radial diffusivity ..... 3

**ROI** region of interest ..... 3

**FMRIB** functional magnetic resonance imaging of the brain

**FNIRT** FMRIB’s nonlinear image registration tool..... 3

# Introduction

Basal ganglia is a part of the human brain which is group of subcortical nuclei responsible primarily for motor control, as well as other roles such as motor learning, executive functions and behaviors, and emotions. [1] Huntington's disease is a disorder that causes the progressive degeneration of the basal nuclei. [2]

Hospital de Bellvitge provided an excellent dataset of magnetic resonance imaging (MRI) and diffusion magnetic resonance imaging (dMRI) records of 32 control and 37 Huntington patient records of T1 and T1/T2 MRI images with isotropic voxels of 1 millimeter resolution and dMRI fractional anisotropy (FA), mean diffusivity (MD) and radial diffusivity (RD) images with isotropic voxels of 2 millimeter resolution and 1 second temporal resolution. Furthermore this dataset also contains the mask for the basal ganglia, which will also be referenced as the region of interest (ROI). Masks for the 7 main cortical regions of the brain, which will also be referenced as the target regions: Limbic, Executive, Rostral-Motor, Caudal-Motor, Parietal, Occipital and Temporal are also included in the dataset. Tractography was performed on the dMRI images to figure out which parts of the ROI are connected to which cortical target, in a similar manner to how it was done in this paper [3]; where the relative connectivity maps are representing the ratio of the number of streamlines to each cortical target. Furthermore, the raw streamline images are also available, where there are a maximum of 5000 streamlines from each voxel in the ROI. The subcortical segmentation of the Basal Ganglia is also available, for the Caudate ,Putamen and Accumbens on the control patients. And lastly FMRIB's nonlinear image registration tool (FNIRT) warp fields were also provided for converting the records into normalized space.




Figure 1.1: Basal Ganglia (ROI) & Cortical Targets

Color	Region
□ White	Basal Ganglia (ROI)
■ Green	Limbic
■ Brown	Executive
■ Light Blue	Rostral-Motor
■ Purple	Caudal-Motor
■ Red	Parietal
■ Blue	Occipital
■ Yellow	Temporal

Table 1.1: Regions Legend

Furthermore, for both the ROI and cortical targets, the dataset distinguishes between the right and left halves of the brain. Thus there are actually 2 ROIs and  $2 \cdot 7 = 14$  target regions.




Figure 1.2: Connectivity Maps

# Fulfillment of the Objectives

## 2.1 Objectives

The end goal is to predict the relative connectivity of the Basal Ganglia to the cortical targets, from the radiomics features of the T1 and T1/T2 images.

This being a very complex problem, there is the possibility that the correlation between the connectivity of the brain and the T1, T1/T2 images are too weak to be mapped on this dataset. As from a datascience perspective, 69 datapoints are not much. But from a medical perspective it is substantial as it is very hard to collect uniform, clean data, with permissions to use it for research.

A simpler task leading up to the complex end goal, is a model for the simple segmentation of the Basal Ganglia for the subcortical regions Caudate Putamen and Accumbens. In order to confirm that the radiomics texture of the T1 and T1/T2 images of this dataset are correlated to the segmentation of the Basal Ganglia. This problem is inherently connected to the main goal, as the relative connectivity does obey certain anatomical restrictions, and the subcortical segmentation of the Basal Ganglia is confirmed to be related to the relative connectivity. Thus if this simpler prediction fails, there is a good chance that the complex end goal will fail as well.

Another intermediate task, is a model for predicting FA and MD images. This is also related to the main goal, as these images are computed from the dMRI images, the same image that the relative connectivity was computed from. But it is inherently simpler, not needing to perform complex algorithms like tractography.

The biggest obstacle of this project is the preprocessing of the data, as there are many variations and hyperparameters that can be tuned. An exhaustive search definitely will not be viable, thus the preprocessing and model will need to be tuned in a waterfall like manner, making educated guesses and comparing model performances across different tries. The main metric to measure model performance, will be the accuracy of the label prediction across voxels, as it should be comparable between all approaches. The accuracy metric will be used for the subcortical label prediction problem as well, and pearson correlation will be used as the metric to evaluate the FA and MD predictions.

## 2.2 Motivation

The motivation for predicting the connectivity maps from the T1 and T1/T2 MRI images, is skipping the time and resource consuming process performing dMRI and tractography.

## 2.3 Experiments

The following contributing factors are needed to be explored and experimented with:

- Experiment Type:
  - Basal Ganglia subcortical segmentation (classification)
  - Brain / Basal Ganglia diffusion FA and MD prediction (regression)
  - Basal Ganglia relative connectivity segmentation (classification)
  - Basal Ganglia number of streamlines prediction (regression)
- Input Space:
  - Native
  - Normalized
- Input Data:
  - T1
  - T1/T2
  - Mixed (T1 and T1/T2)
- Radiomics Extraction Parameters:
  - Kernel size
  - Bin width
  - Relative bin width (fixed number of bins)
- Additional Non-Voxel Based Radiomics Inputs:
  - Basal Ganglia
  - Entire Brain
  - Target Regions
- Left and Right Hemispheres:
  - Left only
  - Right only
  - Left and Right with concatenated label information (e.g. no differentiation between left and right target regions)
  - Left and Right with NOT concatenated label information (e.g. differentiation between left and right target regions)
- Control and Patient Datapoints:
  - Control only
  - Patient only
  - Mixed (Control and Patient)
- Additional Clinical Data Input
- Sequential Backward Feature Selection

- Feature Scaling and/or Normalization
- Relative Connectivity Thresholding
- Model Properties
  - General Architecture
    - \* Single FNN Model
    - \* Dual FNN Model (more on this later)
    - \* Mixture of Experts FNN Model (more on this later)
  - Model Size
    - \* Number of Layers
    - \* Layer Width
  - Activation Function
  - Batch Size
  - Early Stopping
  - Loss Function
  - Learning Rate
  - Regularization

As mentioned, an exhaustive search of the hyperparameter space is not feasible, thus it must be explored in a linear way. The following experiments are already done, or partially done:

- Basal Ganglia subcortical segmentation
  - Sanity Check: tried to predict it from T1 to make sure there is no direct correlation
  - Additional Non-Voxel Based Radiomics Inputs: experimented with additional non-voxel based radiomics features
  - Radiomics Kernel Sizes: experimented with including voxel based features with different kernel sizes
  - Short Conclusion: including multiple different kernel sizes are the best, non-voxel based features made no improvement, final test accuracy 91%
- Brain / Basal Ganglia diffusion FA and MD prediction
  - FA: predicted FA in Basal Ganglia from many different kernel sizes (did not use any other features besides voxel based radiomics), yielded test pearson correlation of 0.92
  - MD: predicted MD in Basal Ganglia from many different kernel sizes (did not use any other features besides voxel based radiomics), yielded test pearson correlation of 0.83
  - TODO: whole brain prediction
- Basal Ganglia relative connectivity segmentation
  -



# Adjustments Made

# Work Plan for Completion