



Working Title
Natasha Hopkins

Stage 3 Project for Master of Biology (MBiol) Degree
Univeristy of York, UK

Project Director
Dr. Richard Maguire

Examination Date
18 April, 2022

Contents

Introduction	2
Flow	2
Flow During Development	2
Developmental Proteins / mechanosensors	2
Endothelial	2
Atherosclerosis	2
Pathway	2
Hypothesis	2
Methods	2
Orbital Shaker	2
mRNA Isolation and qPCR	2
Results	3
Discussion	3
Atheroprotective gene expression	3
Future	4
Acknowledgements	4
References	4

List of Figures

- 1 Relative mRNA expression of angiopoietin-2, axin-2, and thrombospondin-1 from HUVECs exposed to high and low oscillatory shear stress (OSS). Normalised to the HPRT control and relative to the static control. 3

List of Tables

- 1 Oligonucleotide qPCR primers from Ensembl. 3

With diam quis enim lobortis scelerisque fermentum dui faucibus in ornare quam viverra orci sagittis eu volutpat odio facilisis mauris sit amet massa vitae tortor condimentum lacinia quis vel eros donec ac odio tempor orci dapibus ultrices in iaculis nunc sed augue lacus viverra vitae congue eu consequat ac felis donec et odio pellentesque diam volutpat commodo sed egestas egestas fringilla phasellus faucibus scelerisque eleifend donec pretium vulputate sapien nec sagittis aliquam malesuada bibendum arcu vitae elementum curabitur vitae nunc sed velit dignissim sodales ut eu sem integer vitae justo eget magna fermentum iaculis eu non diam phasellus vestibulum lorem sed risus ultricies tristique nulla aliquet enim tortor at auctor urna nunc id cursus metus aliquam eleifend mi in nulla posuere sollicitudin aliquam ultrices sagittis orci a scelerisque purus semper eget dui at tellus at urna condimentum mattis pellentesque id nibh tortor id aliquet lectus proin nibh nisl condimentum id venenatis a condimentum vitae sapien pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas sed tempus urna et pharetra pharetra massa massa ultricies mi quis hendrerit dolor magna eget est lorem ipsum dolor sit amet consectetur adipiscing elit pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas integer eget aliquet nibh praesent tristique magna sit amet purus gravida quis blandit turpis cursus in hac habitasse platea dictumst quisque sagittis purus sit amet volutpat consequat mauris nunc congue nisi vitae suscipit tellus mauris a diam maecenas sed enim ut sem viverra aliquet eget sit.

Key Words: Atherosclerosis • Wnt Signalling Pathway • β catenin • Shear Stress
• Human Umbilical Vein Endothelial Cells (HUVECs) • Angiopoietin-2 •
Thrombospondin-1

(250 Words)

Introduction

Flow

Flow During Development

Developmental Proteins / mechanosensors

Endothelial

Atherosclerosis

WSS

Pathway

Does Axin, Angp2, Thrombosin-2 change if Wnt is inhibited?

Hypothesis

XAV-939 Wnt/Beta Catenin inhibitor, acts by inhibiting tankyrase

Methods

Orbital Shaker

HUVECs were cultured in M199 complete growth medium until ~80% confluent. Cells were then washed with warmed PBS and incubated with 1ml of trypsin until cells thoroughly detached. Cells were transferred to a falcon tube with M199 and spun for 5 minutes at 400g. The supernatant was discarded, and cells were re-suspended in M199 media and transferred to 10mm radius 6 well plates. Once confluent, 3ml of 1% DMSO in M199 was added to one half of the plates, and 3ml of 1% XAV939 Wnt inhibitor in M199 to the other half. Cells were subjected to flow using a orbital shaker at 210 rpm for 72 hours, with the exception of a static controls.

mRNA Isolation and qPCR


Media was removed from the plates and cells were washed with cold PBS. Cells were isolated from the periphery and centre of the plates with PBS and centrifuged for 5 minutes at 400g to remove the supernatant. Total mRNA was extracted using the

RNEasy Mini Kit (Qiagen). The amount of isolated mRNA was determined spectrophotometrically. cDNA synthesis was performed using the Verso cDNA Synthesis Kit (Thermo Scientific) with 5.5µl of 0.64% of mRNA. ANGPT2, AXIN2, THSB1, and HPRT1 mRNA was quantified using StepOne qPCR (Thermo Scientific) using oligonucleotide qPCR primers from Ensembl (Howe et al., 2020) (Table 1).

Table 1. Oligonucleotide qPCR primers from Ensembl.

Gene	Direction	Sequence
ANGPT2	L	CGGCTGTGATGATAGAAATAGGGA
	R	GTTCCAAGAGCTGAAGTTCAAGTC
AXIN1	L	TGTCACCTACTTTTTCTGTGGGGA
	R	TGTCACCTACTTTTTCTGTGGGGA
HPRT1	L	TTGGTCAGGCAGTATAATCC
	R	GGGCATATCCTACAACAAC
THSB1	L	AAAGATGGAGAATGCTGAGTTGGA
	R	GGTCCAAGACAAACCTCACATT

Results



Discussion

Atheroprotective gene expression

Limitations of orbital shaker = improve method

64 **Future**

65 epigenetics

66 look at proliferation, apoptosis, senescence, inflammation = PERP, p53

67 look at vascular repair = wound scratch assay?

68 look at emt = slug/snail?

69 **Acknowledgements**

70 (291 Words) remove headers !!!

71 **References**

72 Howe, K. L. et al. (2020). Ensembl 2021. *Nucleic Acids Research*, 49 (D1), pp.D884–
73 D891. [Online]. Available at: doi:[10.1093/nar/gkaa942](https://doi.org/10.1093/nar/gkaa942).