

TITLE

by Natasha Louise Hopkins

Master of Biology (Honours), Molecular Cell Biology

University of York, UK

Project Director

Prof. Robert J White

Examination Date

17 April, 2023

Word Count

Abstract:

Main:



UNIVERSITY
of York

Contents

Introduction	1
FOXA1 Expression and ERα+ Breast Cancer	1
tRNAs and Gene Expression	1
Materials & Methods	1
Acquisition of Public ChIP-seq Datasets	1
EaSeq for Chip-seq Peak Quantification	2
Motif Analysis	2
Statistics	3
Results	3
FOXA1 and H3k27ac Binding Events	3
Background	3
-	3
Why? What?	3
Heatmaps	3
Average Tracks	3
Suggests	5
Boxplots	5
Why?	5
What?	5
Boxplot	5
Volcano	5
Localisation of FOXA1 at individual tRNA genes in MCF-7 cells	5
Discussion	7
Future	7
References	7

TITLE

Natasha L. Hopkins

Abstract

1 Words

Introduction

FOXA1 Expression and ER α + Breast Cancer

tRNAs and Gene Expression

Materials & Methods

Acquisition of Public ChIP-seq Datasets

ChIP-seq was performed on genetically modified MCF7L cells (*insertion, using a lentiviral cDNA delivery system to express Dox-inducible FOXA1*)^[1]. Datasets were deposited into the National Centre for Biotechnology Information (NCBI) Sequence Read Archive (SRA)^[2] under accession no. PRJNA512997 (Table 1). Using “Genetic Manipulation Tools” within the Galaxy^[3] environment (v 23.0.rc1), SRAs were converted to FastQ files. FastQ files were then aligned to the human genome assembly GRCh37 (hg19) using Bowtie2 (v 2.5.0)^[4] to output BAM files.

Table 1. Publicly available ChIP-seq SRA files acquired from the NCBI SRA database (accession no. PRJNA512997).

Experiment	SRA	Factor	Tissue	Assembly
PRJNA512997	SRR8393424	FOXA1	MCF-7LP	GRCh37 (Hg19)
	SRR8393425			
	SRR8393426			
	SRR8393427	H3K27ac		
	SRR8393428			
	SRR8393431	None (input)		
	SRR8393432			

EaSeq for Chip-seq Peak Quantification

BAM files were uploaded into EaSeq (v1.111) as “Datasets” using the standard settings for Chip-seq data. GRCh37 (hg19) tRNA sequences (n = 606) were downloaded as a “Geneset” from the UCSC Table Browser^[5], (available at <https://genome.ucsc.edu>). High-confidence tRNAs (n = 416) identified in the GtRNAdb^[6] were extracted as a “Regionset”.

Signal peak intensities surrounding tRNAs were quantified using the EaSeq “quantify” tool. Here the default settings “Normalize to reads per million” and “Normalize counts to DNA-fragments” were left checked. The default setting “Normalise to a signal of 1000 bp” was unchecked. The window size was offset ± 500 bp from the start of each tRNA gene. Outputs are referred to as “Q-values”.

To quantify upstream and downstream signals, the “quantify” tool was used with adjusted window sizes. The upstream region was defined as 500 bp preceding and the first nucleotide of tRNA loci. Thus, the start position was offset to 0 bp, and the end position was offset to -500 bp. The downstream region constitutes the 500 bp region beginning with the first nucleotide of tRNA gene body. The start position was offset to 1 bp, and the end position was offset to 500 bp.

Following quantification, tRNA binding events were arranged in ascending order -DOX Q-value and visualised as heatmaps. Data was also visualised with “average”, and “overlay” EaSeq tools.

EaSeq^[7] is available at <http://easeq.net>.

Motif Analysis

Multiple EM for Motif Elicitation ChIP (MEME) Suite

Statistics

Statistical tests and graphs were generated with R^[8] (v 4.2.3), R Studio^[9] (v 2023.03.0.386) and the tidyverse^[10] package.

Results

FOXA1 and H3k27ac Binding Events

Background

-

Why? What?

To investigate the impact of FOXA1 on tRNA enhancers in ER+ MCF-7 cells, public ChIP-Seq datasets from Fu et al. (2019)^[1] were interrogated. In this paper, a doxycycline (Dox) inducible OE system was used to achieve FOXA1 OE akin to tamoxifen-resistant (TamR) MCF-7 cells^[1].

Heatmaps

Quantification of 416 high-confidence tRNAs revealed that both FOXA1 and H3k27ac interact with the majority of tRNA genes, relative to ± 10 kb flanking regions. Whereas input reads generated minimal peak enrichment (Supplementary Figures). Upon FOXA1 OE, average FOXA1 binding at tRNAs increases tRNA genes. On the other hand, H3K27ac binding decreases.

Average Tracks

This was confirmed by average signal intensity tracks 4/1.




Figure 1. (B) Ratiometric heatmaps of the log₂ ratio between the binding of FOXA1 or H3k27ac with endogenous FOXA1 expression vs. the binding of FOXA1 or H3k27ac with FOXA1 OE.

Suggests

Boxplots

Why?

What?

Mann Whitney U to compare means

Boxplot

Volcano

X2 = x number of significantly upregulated and downregulated genes

more Up than down (FOX)

about the same for H3

Localisation of FOXA1 at individual tRNA genes in MCF-7 cells

Table 2. .

Group	Function
ALOXE	Insulator Function ^[11,12]
Ebersole	Insulator Function ^[12,13]
HES7	
Per1	
TMEM107	Insulator Function ^[11,12]
Arg-CCG	Implicated in Cancer Progression ^[14]
Glu-TTC	Implicated in Cancer Progression ^[14]
iMET	Proliferation of Breast Cancer
Met	iMet Control
SeC	Involved in REDOX ^[15]




Figure 2. (B) Ratiometric heatmaps of the log2 ratio between the binding of FOXA1 or H3k27ac with endogenous FOXA1 expression vs. the binding of FOXA1 or H3k27ac with FOXA1 OE.

Discussion

- not compared mcf-7 to treatment responsive cells

Future

- FOXA1 alone not efficient to increase activity
 - p300
- FOXA1 moves nucleosomes to make other TF accessible
- Loses fox = weak binding?
- Dynamic and stable marks
- perturbations
- ATAC-seq

574 Words

References

1. Fu X, Pereira R, De Angelis C, Veeraraghavan J, Nanda S, Qin L, et al. FOXA1 upregulation promotes enhancer and transcriptional reprogramming in endocrine-resistant breast cancer. *Proceedings of the National Academy of Sciences* [Internet]. 2019 Dec 11;116(52):26823–34. Available from: <http://dx.doi.org/10.1073/pnas.1911584116>
2. Leinonen R, Sugawara H, Shumway M. The Sequence Read Archive. *Nucleic Acids Research* [Internet]. 2010 Nov 9;39(Database):D19–21. Available from: <http://dx.doi.org/10.1093/nar/gkq1019>
3. Afgan E, Nekrutenko A, Grünig BA, Blankenberg D, Goecks J, Schatz MC, et al. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2022 update. *Nucleic Acids Research* [Internet]. 2022 Apr 21;50(W1):W345–51. Available from: <http://dx.doi.org/10.1093/nar/gkac247>
4. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nature Methods* [Internet]. 2012 Mar 4;9(4):357–9. Available from: <http://dx.doi.org/10.1038/nmeth.1923>

5. Karolchik D. The UCSC Table Browser data retrieval tool. *Nucleic Acids Research* [Internet]. 2004 Jan 1;32(90001):493D–496. Available from: <http://dx.doi.org/10.1093/nar/gkh103>
6. Chan PP, Lowe TM. GtRNAdb 2.0: an expanded database of transfer RNA genes identified in complete and draft genomes. *Nucleic Acids Research* [Internet]. 2015 Dec 15;44(D1):D184–9. Available from: <http://dx.doi.org/10.1093/nar/gkv1309>
7. Lerdrup M, Johansen JV, Agrawal-Singh S, Hansen K. An interactive environment for agile analysis and visualization of ChIP-sequencing data. *Nature Structural & Molecular Biology* [Internet]. 2016 Feb 29;23(4):349–57. Available from: <http://dx.doi.org/10.1038/nsmb.3180>
8. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2023. Available from: <https://www.R-project.org/>
9. Posit Team. RStudio: Integrated development environment for r [Internet]. Boston, MA: Posit Software, PBC; 2023. Available from: <http://www.posit.co/>
10. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the tidyverse. *Journal of Open Source Software* [Internet]. 2019 Nov 21;4(43):1686. Available from: <http://dx.doi.org/10.21105/joss.01686>
11. Raab JR, Chiu J, Zhu J, Katzman S, Kurukuti S, Wade PA, et al. Human tRNA genes function as chromatin insulators. *The EMBO Journal* [Internet]. 2011 Nov 15;31(2):330–50. Available from: <http://dx.doi.org/10.1038/emboj.2011.406>
12. Sizer RE, Chahid N, Butterfield SP, Donze D, Bryant NJ, White RJ. TFIIC-based chromatin insulators through eukaryotic evolution. *Gene* [Internet]. 2022 Aug;835:146533. Available from: <http://dx.doi.org/10.1016/j.gene.2022.146533>
13. Ebersole T, Kim J-H, Samoshkin A, Kouprina N, Pavlicek A, White RJ, et al. tRNA genes protect a reporter gene from epigenetic silencing in mouse cells. *Cell Cycle* [Internet]. 2011 Aug 15;10(16):2779–91. Available from: <http://dx.doi.org/10.4161/cc.10.16.17092>
14. Goodarzi H, Nguyen HCB, Zhang S, Dill BD, Molina H, Tavazoie SF. Modulated Expression of Specific tRNAs Drives Gene Expression and Cancer Progression. *Cell* [Internet]. 2016 Jun;165(6):1416–27. Available from: <http://dx.doi.org/10.1016/j.cell.2016.05.046>
15. Sangha AK, Kantidakis T. The Aminoacyl-tRNA Synthetase and tRNA Expression Levels Are Deregulated in Cancer and Correlate Independently with Patient Survival. *Current Issues in Molecular Biology* [Internet]. 2022 Jul 2;44(7):3001–19. Available from: <http://dx.doi.org/10.3390/cimb44070207>