



Pairs ★

- Problem
- Submissions
- Leaderboard
- Editorial
- Topics

Given an array of integers and a target value, determine the number of pairs of array elements that have a difference equal to the target value.

Example

$k = 1$
 $arr = [1, 2, 3, 4]$

There are three values that differ by $k = 1$: $2 - 1 = 1$, $3 - 2 = 1$, and $4 - 3 = 1$. Return **3**.

Function Description

Complete the pairs function below.

pairs has the following parameter(s):

- int k: an integer, the target difference
- int arr[n]: an array of integers

Returns

- int: the number of pairs that satisfy the criterion

Input Format

The first line contains two space-separated integers n and k , the size of arr and the target value.

The second line contains n space-separated integers of the array arr .

Constraints

- $2 \leq n \leq 10^5$
- $0 < k < 10^9$
- $0 < arr[i] < 2^{31} - 1$
- each integer $arr[i]$ will be unique

Sample Input

STDIN	Function
5 2	arr[] size n = 5, k =2
1 5 3 4 2	arr = [1, 5, 3, 4, 2]

Sample Output

3

Explanation

There are 3 pairs of integers in the set with a difference of 2: [5,3], [4,2] and [3,1]. .

```
20     return 0;
21   });
22
23   function readLine() {
```

Line: 31 Col: 2

Submit Code

You have passed the sample test cases. Click the submit button to run your code against all the test cases.

Download

1 3