

SoftwareX

duet-tools: A Python package for interacting with the DUET program

--Manuscript Draft--

Manuscript Number:	SOFTX-D-25-00776R1
Article Type:	Original software publication
Section/Category:	Environmental Sciences
Keywords:	Python; fuel modeling; understory; fire modeling
Corresponding Author:	Niko Tutland New Mexico Consortium UNITED STATES OF AMERICA
First Author:	Niko Tutland
Order of Authors:	Niko Tutland
Abstract:	Fuel modeling is a key avenue for understanding dynamics of fire-vegetation interactions across landscapes. Three-dimensional (3D) fuel models are used as inputs for fire behavior models to develop strategies for prescribed fire application and wildfire risk assessment and mitigation. Distribution of Understory using Elliptical Transport (DUET) is a recently developed program for creating surface fuel inputs for 3D fuel models. DUET simulates litter fall from 3D tree canopy inputs and grass growth, producing spatially heterogeneous estimates of fine fuel distribution and characteristics in forested domains. Here we introduce duet-tools, a Python package that streamlines the process for creating DUET simulation input files and calibrating the values in DUET output files. The package handles two primary aspects of the DUET workflow: (1) programmatic creation and management of the DUET input file with validation and documentation, and (2) calibration of fine fuel outputs towards values provided by online data sources or directly from the user. By simplifying these tasks, duet-tools allows modelers and managers to more easily interact with the DUET program while incorporating locally accurate fuels data.

Reviewer #1:

Dear Authors,

Greetings, and thank you for your interesting submission to SoftwareX. Your work on duet-tools addresses a relevant and practical problem in wildland fire science by facilitating DUET input/output management and calibration of 3D fuel fields for models such as QUIC-Fire and FIRETEC. Overall, the software appears scientifically sound, useful to the prescribed fire and fire-modeling community, and well aligned with the aims of the journal.

Many thanks for the helpful comments and suggestions. They have improved the manuscript.

However, I recommend major revisions before the manuscript can be considered for publication. My main concerns are not about the core scientific contribution, but about template compliance, structure, and completeness of the required SoftwareX elements. The most important issues are summarized below:

1. Template and structure issues

- The submission does not fully follow the SoftwareX template. In particular, the PDF appears to contain two versions of the article text, with duplicated sections and metadata. Please ensure that only a single, clean and final version of the manuscript remains in the file.

Thanks for pointing this out. The duplicate article was a result of a misunderstanding of the submission guidelines, where I submitted a rendered PDF as well as a LaTeX file. The first copy is the PDF rendered in Overleaf, which was the version of the article I meant to be the final submission. The second copy was rendered by SoftwareX, and contains the formatting errors you're referring to. In my resubmission I will ensure that only one copy of the manuscript is submitted, and that it contains the proper formatting.

- In the second copy of the text, there are multiple unresolved placeholders such as "Listing ??", "Figure ??" and citation markers like "[?]". These must be fully resolved and checked so that all cross-references and references are correctly formatted.

See my response to the above comment.

2. Missing "Software Metadata" table

- The article currently includes a Code metadata table (repository, license, dependencies, documentation, etc.), which is very good.

- However, SoftwareX requires two specification tables: a Software metadata table and a Code metadata table.

- Please add a complete Software metadata table in accordance with the Guide for Authors and ensure that both tables are fully and correctly filled in.

You are correct that the Software Metadata table was omitted, however, according to the article template linked in the Author Guidelines, this table is optional. To the best of my knowledge, it is intended for executable softwares, which duet-tools is not. Though the DUET program is an executable software, this manuscript only presents a Python package that interacts with the outputs of DUET. Other papers published by SoftwareX that describe Python or R packages do not include a Software Metadata table, so I felt it was appropriate to omit.

3. Length and conciseness

- Because of the duplicated version of the manuscript within the same file, the effective length is longer than needed and may exceed what is appropriate for this article type.

I have removed the duplicated article. All other conciseness considerations are addressed in the comment below.

- After removing the duplicate sections, I encourage you to review the text for redundancies and streamline overlapping explanations so that the final version is concise and easy to follow.

Thanks for drawing my attention to redundancies in the text. I have resolved a few overlapping explanations, summarized here:

L243 Previously: The continued development of \texttt{duet-tools} can help it join the ranks as way to facilitate this goal. As fuel structures and fire dynamics become more uncertain in a changing climate \cite{davis_fire-catalyzed_2020, coop_wildfire-driven_2020, wion_multidecadal_2024}, the combination of DUET and \texttt{duet-tools} can help capture the fine-scale, local fuel dynamics that are essential for wildland fire modeling.

L243 Now: The continued development of \texttt{duet-tools} can help it join the ranks as a way to facilitate this goal, especially as fuel structures and fire dynamics become more uncertain in a changing climate \cite{davis_fire-catalyzed_2020, coop_wildfire-driven_2020, wion_multidecadal_2024}.

L251 Previously: Accurate descriptions of fuel and vegetation are essential for wildland fire modeling \cite{loudermilk_vegetations_2022}, which is an integral aspect of land management planning \cite{drury_interagency_2016}, disaster mitigation \cite{staley_updated_2016}, and species conservation \cite{lucash_managing_2022}. Prescribed burning is a key management practice for forests across the U.S. \cite{ryan_prescribed_2013} that utilizes fuel and fire modeling for prioritization, implementation and incident planning, and practitioner training. For all of these purposes, fire modeling that accounts for 3D fuel structures, topography, and dynamic winds is poised to become an increasingly vital component for planning prescribed fire in high-priority landscapes \cite{parsons_going_2023, hiers_prescribed_2020}.

L251 Now: Accurate descriptions of fuel and vegetation are essential for wildland fire modeling \cite{loudermilk_vegetations_2022}, which is an integral aspect of prescribed fire planning and training \cite{parsons_going_2023, hiers_prescribed_2020} that supports land management \cite{drury_interagency_2016}, disaster mitigation \cite{staley_updated_2016}, and species conservation \cite{lucash_managing_2022}.

4. Empirical evaluation and examples

- The current example (code listing plus figure showing the effect of calibration on fuel load distributions) is useful and illustrative.

- To strengthen the paper, you might consider adding one or two brief additional examples or small benchmarks (e.g., different landscapes or calibration settings, or a short performance note). This would make the empirical evaluation clearer and more convincing, although I see this as a secondary priority compared to the structural issues above.

Thanks for the encouragement to expand the demonstration of the package capabilities. At the behest of another reviewer, I have added an example of DUET output calibration using LANDFIRE data, supported by a second figure illustrating the process (Listing 2 and Fig. 2).

5. Minor editorial and clarity comments

- Please run a careful language and formatting check, correcting minor typos and ensuring consistent terminology (e.g., "landscapes" versus any misspellings).

I appreciate the attention to detail. I have corrected misspellings and typos throughout the document.

- Verify the reference list for any truncated entries, inconsistent styles, or missing information (years, page numbers, DOIs) and bring it fully into line with the journal's reference format.

Thanks for your attention to the references. I prepared the bibliography file according to the article template's specifications, but it looks like the Overleaf rendering (the first copy of the article in the submission) did not render the formatting as it appears in the Author Guidelines, and the SoftwareX rendering (second copy of article in submission) did not detect the bibtex file I submitted. I will work with the journal editor to ensure that all references are formatted correctly in the next submission.

- Where you refer to online documentation or examples (e.g., GitHub pages), it can help the reader if you briefly indicate what a new user will find there (tutorials, API reference, example scripts).

Thanks for the suggestion. I have listed the materials available online where applicable, which consist of code documentation, how-to guides, and example scripts.

Metadata Table, Nr. C8: "Full documentation, how-to guides, example scripts, and issue tracking: <https://nmc-cafes.github.io/duet-tools/>"

Section 3, Paragraph 1: "Full documentation, how-to guides, example scripts, and issue tracking are hosted on GitHub."

I believe that duet-tools is a valuable contribution and has clear potential impact for the wildland fire modeling community. Once the manuscript fully follows the SoftwareX template, includes the missing Software metadata table, removes duplicated/placeholder material, and slightly strengthens the empirical presentation, it will be much closer to being ready for publication.

Best Regards

Thank you for your time.

Reviewer #2: The manuscript presents a valuable Python package that enhances the usability and calibration workflow of DUET for surface fuel modeling. The contribution is well aligned with SoftwareX objectives, and the technical explanations are clear. However, several formatting, citation, and structural issues need attention before the manuscript can be considered for publication.

I appreciate the thoughtful and organized comments. They have improved the manuscript.

Comments for the Authors

1. Correction of Citation Placeholders

Numerous instances of unresolved citation markers such as "[?]", "??", and broken cross-references appear throughout the manuscript. These must be corrected to ensure proper academic referencing and readability.

Thanks for pointing out these issues. All unresolved citation markers are present in the second copy of the manuscript, which was included in error. Please see comment 2 below for more details.

2. Removal of Duplicate Content

Large sections including the abstract, metadata table, introduction, and software description appear duplicated in the later pages. These duplicated blocks seem accidental and should be removed to avoid confusion and redundancy.

A duplicate of the manuscript was included in error due to the submission of both a PDF and a .tex file. I am in contact with SoftwareX to ensure that my resubmission will consist of the proper file types, cross-references and bibliography links.

3. Typographical and Formatting Errors

Several typographical issues (e.g., "foresteddomains," "lanscapes") and spacing inconsistencies are noticeable. A few lines appear merged or improperly broken. A thorough proofreading is recommended to correct these errors.

Thanks for identifying these issues. The two typographical issues you highlighted have been rectified. Merged or improperly broken lines have been resolved through proper file submission procedures.

4. Suggested "Limitations & Future Work" Section

While the structure broadly fits SoftwareX guidelines, adding a brief subsection on "Limitations and Future Work" would strengthen the completeness of the manuscript by acknowledging current boundaries and outlining ongoing developments.

I appreciate this suggestion. Please see Section 4.1, which acknowledges the simplicity of the current calibration functionality and describes a few areas where future development is aimed, specifically in regards to the incorporation of more broad-scale fuels data.

5. Clarification of DUET Input Requirements

The calibration workflow and algorithms are well explained. However, including a short description of DUET input expectations such as grid dimensions, units, and file structure would provide valuable context for new users.

Good idea. We have added the following sentence when describing the Inputs module in the Software Functionalities section:

L115: Input parameters for the DUET program include the 3D fuel array dimensions and resolution, the number of years to simulate, and information on prevailing winds: speed, direction, and variability.

6. Impact Section Requires Reference Fixes

The Impact section is conceptually strong, but it contains unresolved citation placeholders similar to earlier sections. Please revise this section to include proper citation links and ensure consistency.


These issues were resolved by deleting the duplicate copy of the manuscript. References now match the journal guidelines and are cross-referenced correctly in the text.

7. Enhancement of Example Section

Listing 1 provides a clear demonstration of the calibration workflow. To better showcase the full capabilities of the package, including an example that uses the LANDFIRE calibration module would be highly beneficial.

1
2
3
4 *Thanks for the suggestion. I have added an example of DUET output calibration using LANDFIRE data,*
5 *supported by a figure illustrating the process (See Listing 2, Fig. 2, and associated text).*
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



[Click here to view linked References](#)



[Click here to access/download](#)

LaTeX Source File

duet-tools_LaTeX_source_files.zip



duet-tools: A Python package for interacting with the DUET program

Niko J. Tutland^{a,*}

^a*New Mexico Consortium, Center for Applied Fire and Ecosystem Science, Los Alamos,
NM, USA*

Abstract

Fuel modeling is a key avenue for understanding dynamics of fire-vegetation interactions across landscapes. Three-dimensional (3D) fuel models are used as inputs for fire behavior models to develop strategies for prescribed fire application and wildfire risk assessment and mitigation. Distribution of Understory using Elliptical Transport (DUET) is a recently developed program for creating surface fuel inputs for 3D fuel models. DUET simulates litter fall from 3D tree canopy inputs and grass growth, producing spatially heterogeneous estimates of fine fuel distribution and characteristics in forested domains. Here we introduce **duet-tools**, a Python package that streamlines the process for creating DUET simulation input files and calibrating the values in DUET output files. The package handles two primary aspects of the DUET workflow: (1) programmatic creation and management of the DUET input file with validation and documentation, and (2) calibration of fine fuel outputs towards values provided by online data sources or directly from the user. By simplifying these tasks, **duet-tools** allows modelers and managers to more easily interact with the DUET program while incorporating locally accurate fuels data.

Keywords: Python, fuel modeling, understory, fire modeling, simulation, fire behavior

Metadata

1. Motivation and significance

In landscapes across North America, wildland fire is a critical driver of ecosystem processes that can both maintain and disturb habitats [1]. Modeling

*Corresponding author

Email address: ntutland@gmail.com (Niko J. Tutland)

Nr.	Code metadata description	Metadata
C1	Current code version	v1.0.1
C2	Permanent link to code/repository used for this code version	https://github.com/nmc-cafes/duet-tools
C3	Permanent link to Reproducible Capsule	—
C4	Legal Code License	MIT License
C5	Code versioning system used	git
C6	Software code languages, tools, and services used	Python (NumPy, pandas, rasterio, landfire)
C7	Compilation requirements, operating environments & dependencies	Microsoft Windows, Linux; Python ≥ 3.10 ; dependencies installed via pip
C8	Link to developer documentation/-manual	Full documentation, how-to guides, example scripts, and issue tracking: https://nmc-cafes.github.io/duet-tools/
C9	Support email for questions	ntutland@gmail.com

Table 1: Code metadata

wildland fire behavior is a key tool for planning land management practices like prescribed burning, where complexities in fire-atmospheric interactions are mediated by heterogeneous, three-dimensional (3D) fuel structures. Physics-based fire behavior models, such as FIRETEC [2] or QUIC-Fire [3], simulate fire behavior in 3D, but take in relatively fine-scale ($<4 \text{ m}^3$) inputs describing fine fuel density, size, moisture, and height. In forested landscapes, modeling the distribution of fine fuels on the ground surface (e.g., leaf litter, herbaceous fuels) is challenging at these scales because of their heterogeneity and the difficulty of observing them with survey tools such as airborne LiDAR.

The Distribution of Understory using Elliptical Transport (DUET) program [4] presents a solution by simulating the spatial arrangement of leaf or needle drop based on tree crown dimensions and prevailing wind direction, placing grass where litter levels would not limit its growth. However, these simulations may not capture some essential elements of understory dynamics that vary geographically, such as decay rates, herbaceous fuel loading, or fuel moisture content. Thus, users of DUET may wish to modify outputs of the simulations, differentially shifting the magnitudes of loading, height, and/or moisture values based on field-measured values, expert opinions, or broad-scale datasets, without altering the spatial distributions predicted by DUET.

Despite the potential for DUET to be used to improve 3D fuel modeling in forested landscapes, no standardized tool yet exists for managing the inputs and outputs of DUET simulations to help parameterize realistic, place-based 3D fire behavior simulations.

Here, we introduce the `duet-tools` Python package, which addresses these needs by providing: (1) a simple, validated interface for creating, reading, and modifying DUET input files; (2) intuitive management of DUET output arrays; (3) built-in functions for calibration of fuel elements; and (4) comprehensive documentation with how-to guides and example scripts.

2. Software description

2.1. Software architecture

`duet-tools` is implemented in Python using a modular design with a combination of functional and object-oriented features. Functions are applied to two main classes representing the package’s functionalities: (1) the `InputFile` class, which manages DUET input files by loading, creating, and writing them with validation; and (2) the `DuetRun` class, which organizes DUET outputs, enabling calibration, format conversion, and export to FIRETEC or QUIC-Fire formats. Typically, a DUET simulation will be run locally in a directory containing the DUET executable file and a text file specifying simulation parameters called `duet.in`. In `duet-tools`, the `InputFile` class represents the parameters within `duet.in` for easy programmatic interaction, and additionally provides the information for automatically importing the data outputs of a completed DUET simulation. These data outputs, which are stored in idiosyncratic data structures, can then be read and represented by a `DuetRun` object.

Once the DUET simulation outputs are imported, `duet-tools` offers flexible methods for modifying or "calibrating" the outputs. Target values and calibration methods for the magnitudes of DUET data can be defined, assigned to various fuel types, and applied differentially based on the specific needs of the user. By following the general workflow of assigning targets, setting those targets for fuel parameters, then calibrating a `DuetRun` object, a user can incorporate local data or expert knowledge into their DUET simulations. The resulting data arrays are represented in standard NumPy [5] formats, which can then be processed manually by the user or conveniently exported to data formats accepted by QUIC-Fire or FIRETEC.

2.2. Software functionalities

The package consists of several modules, three of which we highlight here: (1) **inputs**, (2) **calibration**, and (3) **landfire**. The Inputs module facilitates loading, modifying, and writing validated DUET input files. Input parameters for the DUET program include the 3D fuel array dimensions and resolution, the number of years to simulate, and information on prevailing winds: speed, direction, and variability. The `from_directory()` classmethod automatically creates a representation of `duet.in` which can be conveniently modified and written to a file programmatically. Every parameter is validated upon write, meaning that any errors will be caught before the DUET simulation takes place. This involves type checking and ensuring inputs fall within accepted data ranges.

The Calibration module comprises the bulk of the current functionalities of `duet-tools`, providing methods for shifting and/or scaling DUET output values based on target ranges or distributions. Field data on fuel properties such as fuel loading may be collected and summarized as a range of values or as a distribution with a center and spread. The Calibration module can perform shifting and scaling based on either type of target, and can differentially apply those methods for different fuel parameters and/or fuel types. Targets are assigned using the `assign_targets()` function, which creates an object of class `Targets`. To assign targets based on a range of values, the "maxmin" method applies the following algorithm:

$$x'_i = \begin{cases} \text{min_target} + \frac{x_i - \min(x^+)}{\max(x^+) - \min(x^+)} \cdot (\text{max_target} - \text{min_target}), & x_i > 0 \\ x_i, & x_i \leq 0 \end{cases}$$

where $x^+ = \{x_i \in x \mid x_i > 0\}$.

This shifts, then stretches or squeezes the values in the DUET output data array to match the provided range, without modifying the relative relationships between the data. Similarly, calibration based on mean and standard deviation is applied using the "meansd" argument and the following algorithm:

$$x'_i = \begin{cases} \max \left(0, \text{mean_target} + \frac{(x_i - \mu_x) \cdot \text{sd_target}}{\sigma_x} \right), & x_i > 0 \\ x_i, & x_i \leq 0 \end{cases}$$

where $x^+ = \{x_i \in x \mid x_i > 0\}$, $\mu_x = \text{mean}(x^+)$, $\sigma_x = \text{std}(x^+)$

The user may create one or multiple **Targets** objects which can then be set to different fuel parameters and fuel types. Briefly, a fuel parameter is one of the three required fuel inputs for the QUIC-Fire model: (1) fuel loading, (2) fuel moisture, or (3) fuel depth. A fuel type, on the other hand, corresponds a category of fuel simulated by DUET that each parameter may describe. DUET simulates litter fall from any species of tree represented in the DUET fuel grid input as well as growth of grass or other herbaceous plants. Upon creation of a **DuetRun** object, **duet-tools** groups together the litter from coniferous species, then the litter from deciduous species. This creates three main fuel types to which a **Targets** object may be set: (1) grass, (2) conifer litter, and (3) deciduous litter. Additionally, a target may be set for a combination of both litter types, or to all fuel types together.

The targets corresponding to a single fuel parameter can be set for one or more fuel types using **set_fuel_parameter()**, which creates an object of class **FuelParameter**. Once each desired fuel parameter has been set using targets for each desired fuel type, the **DuetRun** object can be calibrated using the **calibrate()** function. Crucially, any combination of fuel type and fuel parameter can be assigned targets using either calibration method. In practice, this means there is a single function, **calibrate()**, that handles every possible calibration specification. The flexibility to mix and match methods, fuel types, and fuel parameters is facilitated by the use of a small set of keyword arguments in the **assign_targets()** and **set_fuel_parameter()** functions. Running **calibrate()** on a **DuetRun** object produces a new **DuetRun** instance that stores the calibrated arrays. These arrays can be accessed or saved via the classmethods **to_numpy()** and **to_quicfire()**, the latter of which saves the calibrated arrays to a specified directory in the format expected by the QUIC-Fire model.

The **landfire** module supports calibration by offering a convenient source for calibration targets. The main functionality of the module is the execution of a data query of from the LANDFIRE database [6] to be used to calibrate DUET outputs. The module leverages the *landfire* python package [7] to download fuels data, then converts those data to calibration targets. Fuels data themselves are derived from the Scott and Burgan 40 Fire Behavior Fuel Models [8]. Future package development is poised incorporate additional data sources from LANDFIRE, such as the Fuel Characteristics and Classification System (FCCS) [9].

3. Illustrative examples

The main functionalities of `duet-tools` package are compatible with Python version 3.10+ and can be installed using PyPI. Since the `landfire` module requires dependencies that are only compatible with Python version 3.10, it is included as an optional package extra. Full documentation, how-to guides, example scripts, and issue tracking are hosted on GitHub.

3.1. DUET calibration workflow

The use of `duet-tools` is contingent upon the user possessing the results of a DUET simulation. Most often they will exist alongside a DUET input file. A common workflow will begin by creating a `DuetRun` object using `import_duet()`, which derives DUET data dimensions from the `duet.in` input file. If the input file is not present, but the data dimensions are known, the user can alternatively specify `import_duet_manual()`. Next, targets may be assigned using either of the two available calibration methods, and subsequently set to the desired fuel parameter(s). Once targets for each desired fuel parameter have been specified, the `calibrate()` function applies the calibration methods accordingly. A simple example of this process is shown in Listing 1 below.

```
1  # Import DUET outputs
2  duet_run = import_duet(directory=duet_path)
3
4  # Assign targets for each fuel type and fuel parameter
5  coniferous_loading = assign_targets(method="maxmin", max
6      =5.0, min=0)
7  deciduous_loading = assign_targets(method="meansd", mean
8      =0.5, sd=0.1)
9  grass_loading = assign_targets(method="meansd", mean=0.5,
10     sd=0.25)
11
12 # Bring together fuel types for each parameter
13 loading_targets = set_fuel_parameter(
14     parameter="loading",
15     grass=grass_loading,
16     deciduous=deciduous_loading,
17     coniferous=coniferous_loading,
18 )
19
20 # Calibrate the DUET run
```

```

18 calibrated_duet = calibrate(duet_run=duet_run,
19                               fuel_parameter_targets=loading_targets)
20 # Export to a QUIC-Fire input file
21 calibrated_duet.to_quicfire(quicfire_path)

```

Listing 1: Example of a typical workflow using `duet-tools`, including importing a DUET run, setting calibration targets, calibrating fuel parameter values, and exporting to QUIC-Fire fuel arrays.

In Listing 1, the fuel loading parameter is calibrated using separately defined targets for each fuel type. Grass and deciduous litter loading are calibrated using their respective target means and standard deviations, whereas coniferous loading is calibrated based on a target value range. Figure 1 shows how each calibration shifts the magnitudes of fuel loading values for each fuel type, while retaining the relative relationships between values. In this example, coniferous litter and grass fuel loading have been calibrated to reflect lower loading than was predicted by DUET, whereas deciduous litter loading was increased. After these calibrations, overall loading was decreased, but with patterns corresponding to the modifications of the individual fuel types. The total litter loading is available to be exported as an input file to QUIC-Fire, or as a NumPy array for other applications.




Figure 1: Example of DUET calibration. The top row shows original fuel loading outputs from DUET for each fuel type; the bottom row shows calibrated results based on user-specified targets.

Fuel parameter targets may be assigned from LANDFIRE data in a similar fashion. Listing 2 shows the process of querying the LANDFIRE database to retrieve Scott and Burgan 40 Fuel Model data [8] for an area of interest. Spatial data defining an area of interest can be supplied as either a geojson or a polygon. For each fuel type (grass and/or litter; Fig. 2B), the values of the desired fuel parameters (Fig. 2C) are automatically assigned to a **Targets** object, which can then be used to differentially adjust DUET output values for those values (Fig. 2D-E).

```

1  # Import DUET outputs
2  duet_run = import_duet(directory=duet_path)
3
4  # Query the LANDFIRE database
5  landfire_query = query_landfire(
6      area_of_interest=aoi_geojson, year=2019, directory=
7      duet_path, input_epsg=4326
8  )
9
10 # Assign fuel targets from LANDFIRE query
11 grass_loading_targets = assign_targets_from_sb40(
12     landfire_query, "grass", "loading")
13 litter_loading_targets = assign_targets_from_sb40(
14     landfire_query, "litter", "loading")
15 landfire_loading = set_loading(
16     grass=grass_loading_targets, litter=
17     litter_loading_targets
18 )
19
20 # Calibrate the DUET run
21 calibrated_duet_landfire = calibrate(duet_run,
22     landfire_loading)

```

Listing 2: Example of DUET calibration using data from LANDFIRE [6]

4. Impact

Accurate descriptions of fuel and vegetation are essential for wildland fire modeling [10], which is an integral aspect of prescribed fire planning and training [11, 12] that supports land management [13], disaster mitigation [14], and species conservation [15]. Simulating realistic fire behavior using tools like QUIC-Fire and FIRETEC requires an understanding of the heterogeneity of fuel structures on a landscape [16, 17], but that heterogene-




Figure 2: Example of DUET calibration using Scott and Burgan 40 Fuel Model [8] data from the LANDFIRE database [6]. Original DUET output values (A) are assigned fuel parameter targets from a provided area of interest based on the mapped fuel type (B-C). Values for grass (D) and tree litter (E) may be calibrated separately and combined to produce a calibrated DUET output (F).

ity is difficult to characterize. Data fusion approaches that combine local data, broad-scale mapping, and ecological modeling can provide more realistic characterizations wildland fuel heterogeneity [18, 19]. By facilitating the fusion of modeled ecological processes from DUET, satellite-based gridded data from LANDFIRE, and direct input from users, `duet-tools` offers one of the first such approaches for surface fuel modeling.

Though there is utility in fine-scale, data-driven fire behavior modeling, user access to the models, their computational weight, and accurate fuel parameterization stand as barriers to operational use [20]. `duet-tools` helps ease the burden of fuel parameterization by providing a simple and convenient tool for harnessing a fuel modeling software that may otherwise be under-utilized. By enabling the programmatic management and modification of DUET inputs and outputs, `duet-tools` has already allowed it to be incorporated into the most robust 3D fuel modeling platform to date, FastFuels [18]. As a national-scale data platform, FastFuels’ coverage includes a wide variety of ecosystems, fuel types, and climate zones which can have vastly

different litter decay rates [21] and herbaceous growth patterns [22], or may experience a variety of management actions that may modify surface fuels. `duet-tools` provides a key mechanism for accounting for these differences by scaling DUET’s predictions to local data inputs, without compromising the spatial heterogeneity predictions that are central to DUET’s utility.

As the science of wildland fire management continues to expand its toolkit of computing technologies, tools and frameworks for harmonizing fuel data across scales are becoming more widespread [23, 24, 19]. The continued development of `duet-tools` can help it join the ranks as a way to facilitate this goal, especially as fuel structures and fire dynamics become more uncertain in a changing climate [25, 26, 27].

4.1. Limitations and Future Work

This article aims to introduce `duet-tools`, which is still in the early stages of its development. Currently, manipulation of DUET output files is limited to two simple calibration algorithms and 2-3 fuel types. Future work should incorporate more complex and place-based calibration parameters, such as seasonality [28], topography [29], and the addition of other types of surface fuel such as shrubs. Because local fuel data may not be available, further package development should focus on forging more connections to large geospatial datasets, including the expansion of the `landfire` module to access additional fuel layers, such as FCCS data [9]. Since these layers are generally compiled at a resolution of 20-30 m, the variability they describe is currently applied wholesale when calibrating the fine-scale variability simulated by DUET. Future package development could add the option to incorporate the broad-scale fuel heterogeneity described by broad-scale data while maintaining the fine-scale patterns predicted by DUET simulations. These improvements will help `duet-tools` play an integral role in streamlining the integration of DUET with fuels data platforms such as FastFuels [18].

5. Conclusions

`duet-tools` bridges a key usability gap between DUET and downstream fire behavior models. It simplifies DUET file management, enables automated calibration against data sources, and promotes transparent and reproducible workflows for surface fuel modeling.

Acknowledgements

Many thanks to Jenna McDanold for the introduction and continued development of the DUET program. I am also grateful for the guidance and men-

torship from Anthony Marcozzi, as well as the support from Scott Pokswinski and the rest of the NMC CAFES team. Special thanks to Rachel Loehman for encouraging and supporting the development of this tool. Funding for this project is provided by the Robert T. Stafford Disaster Relief and Emergency Assistance Act (42 U.S.C. 5121 et seq.) and supplemental funding acts for Federal disaster relief activities via the U.S. Geological Survey.

References

- [1] J. E. Keeley, Ecological foundations for fire management in North American forest and shrubland ecosystems, Vol. 779, US Department of Agriculture, Forest Service, Pacific Northwest Research Station, 2009.
URL https://books.google.com/books?hl=en&lr=&id=4_Y4IT6YBX8C&oi=fnd&pg=PA1&dq=role+of+fire+in+north+american+forests&ots=80ARlcD01P&sig=lvNYsm1xxHeTZr0RC-xBgMsgfLU
- [2] R. Linn, J. Reisner, J. J. Colman, J. Winterkamp, Studying wild-fire behavior using FIRETEC, International Journal of Wildland Fire 11 (4) (2002) 233–246, publisher: CSIRO PUBLISHING. doi:10.1071/wf02007.
URL <https://www.publish.csiro.au/wf/wf02007>
- [3] R. R. Linn, S. L. Goodrick, S. Brambilla, M. J. Brown, R. S. Middleton, J. J. O'Brien, J. K. Hiers, QUIC-fire: A fast-running simulation tool for prescribed fire planning, Environmental Modelling & Software 125 (2020) 104616, publisher: Elsevier.
URL <https://www.sciencedirect.com/science/article/pii/S1364815219307388>
- [4] J. S. McDanold, R. R. Linn, A. K. Jonko, A. L. Atchley, S. L. Goodrick, J. K. Hiers, C. M. Hoffman, E. L. Loudermilk, J. J. O'Brien, R. A. Parsons, C. H. Sieg, J. A. Oliveto, DUET - Distribution of Understory using Elliptical Transport: A mechanistic model of leaf litter and herbaceous spatial distribution based on tree canopy structure, Ecological Modelling 483 (2023) 110425. doi:10.1016/j.ecolmodel.2023.110425.
URL <https://www.sciencedirect.com/science/article/pii/S0304380023001564>
- [5] C. R. Harris, K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau, E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van Kerkwijk, M. Brett, A. Haldane,

- J. F. del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant, K. Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke, T. E. Oliphant, Array programming with NumPy, *Nature* 585 (7825) (2020) 357–362. doi:10.1038/s41586-020-2649-2.
URL <https://doi.org/10.1038/s41586-020-2649-2>
- [6] I. P. La Puma, LANDFIRE technical documentation, Tech. Rep. 2023-1045, Reston, VA (2023). doi:10.3133/ofr20231045.
URL <https://pubs.usgs.gov/publication/ofr20231045>
- [7] J. Clark, landfire-python (2023).
URL <https://landfire-python.firesci.io/en/latest/index.html#>
- [8] J. H. Scott, R. E. Burgan, Standard Fire Behavior Fuel Models: A Comprehensive Set for Use with Rothermel’s Surface Fire Spread Model, Tech. rep. (2005).
URL <http://www.fs.fed.us/rm>
- [9] R. D. Ottmar, D. V. Sandberg, C. L. Riccardi, S. J. Prichard, An overview of the Fuel Characteristic Classification System — Quantifying, classifying, and creating fuelbeds for resource planning This article is one of a selection of papers published in the Special Forum on the Fuel Characteristic Classification System., *Canadian Journal of Forest Research* 37 (12) (2007) 2383–2393. doi:10.1139/X07-077.
URL <http://www.nrcresearchpress.com/doi/10.1139/X07-077>
- [10] E. L. Loudermilk, J. J. O’Brien, S. L. Goodrick, R. R. Linn, N. S. Skowronski, J. K. Hiers, Vegetation’s influence on fire behavior goes beyond just being fuel, *Fire Ecology* 18 (1) (2022) 9, s42408-022-00132-9. doi:10.1186/s42408-022-00132-9.
URL <https://fireecology.springeropen.com/articles/10.1186/s42408-022-00132-9>
- [11] R. Parsons, L. Wells, A. Marcozzi, R. R. Linn, J. K. Hiers, F. Pimont, K. Riley, I. Altintas, S. Flanary, Going 3D With Fuel and Fire Modeling: FastFuels and QUIC-Fire, *Fire Management Today* 81 (1) (2023) 49–51.
URL <https://www.fs.usda.gov/sites/default/files/fire-management-today/fmt-vol-81-n1.pdf>
- [12] J. K. Hiers, J. J. O’Brien, J. M. Varner, B. W. Butler, M. Dickinson, J. Furman, M. Gallagher, D. Godwin, S. L. Goodrick, S. M.

- Hood, A. Hudak, L. N. Kobziar, R. Linn, E. L. Loudermilk, S. McCaffrey, K. Robertson, E. M. Rowell, N. Skowronski, A. C. Watts, K. M. Yedinak, Prescribed fire science: the case for a refined research agenda, *Fire Ecology* 16 (1) (2020) 11, s42408-020-0070-8. doi:10.1186/s42408-020-0070-8.
URL <https://fireecology.springeropen.com/articles/10.1186/s42408-020-0070-8>
- [13] S. A. Drury, H. M. Rauscher, E. M. Banwell, S. Huang, T. L. Lavezzo, The Interagency Fuels Treatment Decision Support System: Functionality for Fuels Treatment Planning, *Fire Ecology* 12 (1) (2016) 103–123. doi:10.4996/fireecology.1201103.
URL <https://fireecology.springeropen.com/articles/10.4996/fireecology.1201103>
- [14] D. M. Staley, J. A. Negri, J. W. Kean, J. L. Laber, A. C. Tillery, A. M. Youberg, Updated logistic regression equations for the calculation of post-fire debris-flow likelihood in the western United States, Tech. rep., US Geological Survey (2016).
URL <https://pubs.usgs.gov/publication/ofr20161106>
- [15] M. S. Lucash, S. Weiss, M. J. Duveneck, R. M. Scheller, Managing for red-cockaded woodpeckers is more complicated under climate change, *The Journal of Wildlife Management* 86 (8) (2022) e22309, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/jwmg.22309>. doi:10.1002/jwmg.22309.
URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/jwmg.22309>
- [16] S. R. Bonner, C. M. Hoffman, R. R. Linn, W. T. Tinkham, A. L. Atchley, C. H. Sieg, J. M. Varner, J. J. O'Brien, J. K. Hiers, Forest structural complexity and ignition pattern influence simulated prescribed fire effects, *Fire Ecology* 20 (1) (2024) 82. doi:10.1186/s42408-024-00314-7.
URL <https://fireecology.springeropen.com/articles/10.1186/s42408-024-00314-7>
- [17] R. A. Parsons, W. E. Mell, P. McCauley, Linking 3D spatial models of fuels and fire: effects of spatial heterogeneity on fire behavior, *Ecological Modelling* 222 (3) (2011) 679–691, publisher: Elsevier.
URL <https://www.sciencedirect.com/science/article/pii/S0304380010005831>

- [18] A. Marcozzi, L. Wells, R. Parsons, E. Mueller, R. Linn, J. K. Hiers, FastFuels: Advancing wildland fire modeling with high-resolution 3D fuel data and data assimilation, *Environmental Modelling & Software* 183 (2025) 106214. doi:10.1016/j.envsoft.2024.106214.
URL <https://www.sciencedirect.com/science/article/pii/S1364815224002755>
- [19] N. J. Tutland, A. P. Wion, C. J. May, G. C. Hutchings, H. A. Nowak, J. R. Gattiker, J. K. Hiers, R. R. Linn, S. M. Pokswinski, E. Q. Margolis, Representing 3-dimensional fuels for physics-based fire behavior models: a general framework and case study in a type-converted post-fire shrubfield, *Fire Ecology* 21 (1) (2025) 43. doi:10.1186/s42408-025-00383-2.
URL <https://fireecology.springeropen.com/articles/10.1186/s42408-025-00383-2>
- [20] W. Mell, R. Linn, FIRETEC and WFDS Modeling of Fire Behavior and Smoke in Support of FASMEE JFSP PROJECT ID: 16-4-05-1 (2017).
URL https://www.academia.edu/download/76406618/16-4-05-1_final_report.pdf
- [21] J. M. Blair, R. W. Parmelee, M. H. Beare, Decay Rates, Nitrogen Fluxes, and Decomposer Communities of Single- and Mixed-Species Foliar Litter, *Ecology* 71 (5) (1990) 1976–1985, eprint: <https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.2307/1937606>. doi:10.2307/1937606.
URL <https://onlinelibrary.wiley.com/doi/abs/10.2307/1937606>
- [22] P. B. Reich, D. W. Peterson, D. A. Wedin, K. Wragge, FIRE AND VEGETATION EFFECTS ON PRODUCTIVITY AND NITROGEN CYCLING ACROSS A FOREST–GRASSLAND CONTINUUM, *Ecology* 82 (6) (2001) 1703–1719. doi:10.1890/0012-9658(2001)082[1703:FAVEOP]2.0.CO;2.
URL [http://doi.wiley.com/10.1890/0012-9658\(2001\)082\[1703:FAVEOP\]2.0.CO;2](http://doi.wiley.com/10.1890/0012-9658(2001)082[1703:FAVEOP]2.0.CO;2)
- [23] D. Roten, L. Wells, D. Crawl, R. A. Parsons, A. Marcozzi, R. R. Linn, K. Hiers, I. Altintas, TrueTrees: A Scalable Workflow for the Integration of Airborne LiDAR Scanning Data into Fuel Models for Prescribed Fire Simulations, in: 2023 IEEE 19th International Conference on e-Science (e-Science), IEEE, 2023, pp. 1–10.

URL <https://ieeexplore.ieee.org/abstract/document/10254923/>

- [24] I. G. Moreno, I. Nealey, D. Roten, M. H. Nguyen, D. Crawl, K. O’Laughlin, M. Floca, S. Pokswinski, I. Altintas, Visualization and Labeling of Terrestrial LiDAR Data for Three-Dimensional Fuel Classification, in: 2023 IEEE 19th International Conference on e-Science (e-Science), IEEE, Limassol, Cyprus, 2023, pp. 1–2.

URL <https://ieeexplore.ieee.org/abstract/document/10254841/>

- [25] K. T. Davis, P. E. Higuera, S. Z. Dobrowski, S. A. Parks, J. T. Abatzoglou, M. T. Rother, T. T. Veblen, Fire-catalyzed vegetation shifts in ponderosa pine and Douglas-fir forests of the western United States, *Environmental Research Letters* 15 (10) (2020) 1040b8, publisher: IOP Publishing. doi:10.1088/1748-9326/abb9df.

URL <https://dx.doi.org/10.1088/1748-9326/abb9df>

- [26] J. D. Coop, S. A. Parks, C. S. Stevens-Rumann, S. D. Crausbay, P. E. Higuera, M. D. Hurteau, A. Tepley, E. Whitman, T. Assal, B. M. Collins, K. T. Davis, S. Dobrowski, D. A. Falk, P. J. Fornwalt, P. Z. Fulé, B. J. Harvey, V. R. Kane, C. E. Littlefield, E. Q. Margolis, M. North, M.-A. Parisien, S. Prichard, K. C. Rodman, Wildfire-Driven Forest Conversion in Western North American Landscapes, *BioScience* 70 (8) (2020) 659–673. doi:10.1093/biosci/biaa061.

URL <https://doi.org/10.1093/biosci/biaa061>

- [27] A. P. Wion, J. T. Stevens, K. Beeley, R. Oertel, E. Q. Margolis, C. D. Allen, Multidecadal vegetation transformations of a New Mexico ponderosa pine landscape after severe fires and aerial seeding, *Ecological Applications* 34 (6) (2024) e3008, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/eap.3008>. doi:10.1002/eap.3008.

URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/eap.3008>

- [28] Y. Li, Y. Zhang, X. Quan, B. He, S. Veraverbeke, Z. Liao, T. A. J. Janssen, Estimating forest litter fuel load by integrating remotely sensed foliage phenology and modeled litter decomposition, *Remote Sensing of Environment* 317 (2025) 114526. doi:10.1016/j.rse.2024.114526.

URL <https://www.sciencedirect.com/science/article/pii/S0034425724005522>

- 1
2
3
4
5 [29] P. Nyman, D. Metzen, P. J. Noske, P. N. Lane, G. J. Sheridan, Quantify-
6 ing the effects of topographic aspect on water content and temperature
7 in fine surface fuel, International Journal of Wildland Fire 24 (8) (2015)
8 1129–1142, publisher: CSIRO Publishing.
9 URL <https://www.publish.csiro.au/wf/wf14195>
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Declaration of interests

☒The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

duet-tools: A Python package for interacting with the DUET program

Niko J. Tutland^{a,*}

^a*New Mexico Consortium, Center for Applied Fire and Ecosystem Science, Los Alamos, NM, USA*

Abstract

Fuel modeling is a key avenue for understanding dynamics of fire-vegetation interactions across landscapes. Three-dimensional (3D) fuel models are used as inputs for fire behavior models to develop strategies for prescribed fire application and wildfire risk assessment and mitigation. Distribution of Understory using Elliptical Transport (DUET) is a recently developed program for creating surface fuel inputs for 3D fuel models. DUET simulates litter fall from 3D tree canopy inputs and grass growth, producing spatially heterogeneous estimates of fine fuel distribution and characteristics in forested domains. Here we introduce **duet-tools**, a Python package that streamlines the process for creating DUET simulation input files and calibrating the values in DUET output files. The package handles two primary aspects of the DUET workflow: (1) programmatic creation and management of the DUET input file with validation and documentation, and (2) calibration of fine fuel outputs towards values provided by online data sources or directly from the user. By simplifying these tasks, **duet-tools** allows modelers and managers to more easily interact with the DUET program while incorporating locally accurate fuels data.

Keywords: Python, fuel modeling, understory, fire modeling, simulation, fire behavior

Metadata

1. Motivation and significance

In landscapes across North America, wildland fire is a critical driver of ecosystem processes that can both maintain and disturb habitats [1]. Modeling

*Corresponding author

Email address: ntutland@gmail.com (Niko J. Tutland)

Nr.	Code metadata description	Metadata
C1	Current code version	v1.0.1
C2	Permanent link to code/repository used for this code version	https://github.com/nmc-cafes/duet-tools
C3	Permanent link to Reproducible Capsule	—
C4	Legal Code License	MIT License
C5	Code versioning system used	git
C6	Software code languages, tools, and services used	Python (NumPy, pandas, rasterio, landfire)
C7	Compilation requirements, operating environments & dependencies	Microsoft Windows, Linux; Python ≥ 3.10 ; dependencies installed via pip
C8	Link to developer documentation/-manual	Full documentation, how-to guides, example scripts, and issue tracking: https://nmc-cafes.github.io/duet-tools/
C9	Support email for questions	ntutland@gmail.com

Table 1: Code metadata

wildland fire behavior is a key tool for planning land management practices like prescribed burning, where complexities in fire-atmospheric interactions are mediated by heterogeneous, three-dimensional (3D) fuel structures. Physics-based fire behavior models, such as FIRETEC [2] or QUIC-Fire [3], simulate fire behavior in 3D, but take in relatively fine-scale ($<4 \text{ m}^3$) inputs describing fine fuel density, size, moisture, and height. In forested landscapes, modeling the distribution of fine fuels on the ground surface (e.g., leaf litter, herbaceous fuels) is challenging at these scales because of their heterogeneity and the difficulty of observing them with survey tools such as airborne LiDAR.

The Distribution of Understory using Elliptical Transport (DUET) program [4] presents a solution by simulating the spatial arrangement of leaf or needle drop based on tree crown dimensions and prevailing wind direction, placing grass where litter levels would not limit its growth. However, these simulations may not capture some essential elements of understory dynamics that vary geographically, such as decay rates, herbaceous fuel loading, or fuel moisture content. Thus, users of DUET may wish to modify outputs of the simulations, differentially shifting the magnitudes of loading, height, and/or moisture values based on field-measured values, expert opinions, or broad-scale datasets, without altering the spatial distributions predicted by DUET.

Despite the potential for DUET to be used to improve 3D fuel modeling in forested landscapes, no standardized tool yet exists for managing the inputs and outputs of DUET simulations to help parameterize realistic, place-based 3D fire behavior simulations.

Here, we introduce the **duet-tools** Python package, which addresses these needs by providing: (1) a simple, validated interface for creating, reading, and modifying DUET input files; (2) intuitive management of DUET output arrays; (3) built-in functions for calibration of fuel elements; and (4) comprehensive documentation with how-to guides and example scripts.

2. Software description

2.1. Software architecture

duet-tools is implemented in Python using a modular design with a combination of functional and object-oriented features. Functions are applied to two main classes representing the package’s functionalities: (1) the **InputFile** class, which manages DUET input files by loading, creating, and writing them with validation; and (2) the **DuetRun** class, which organizes DUET outputs, enabling calibration, format conversion, and export to FIRETEC or QUIC-Fire formats. Typically, a DUET simulation will be run locally in a directory containing the DUET executable file and a text file specifying simulation parameters called `duet.in`. In **duet-tools**, the **InputFile** class represents the parameters within `duet.in` for easy programmatic interaction, and additionally provides the information for automatically importing the data outputs of a completed DUET simulation. These data outputs, which are stored in idiosyncratic data structures, can then be read and represented by a **DuetRun** object.

Once the DUET simulation outputs are imported, **duet-tools** offers flexible methods for modifying or "calibrating" the outputs. Target values and calibration methods for the magnitudes of DUET data can be defined, assigned to various fuel types, and applied differentially based on the specific needs of the user. By following the general workflow of assigning targets, setting those targets for fuel parameters, then calibrating a **DuetRun** object, a user can incorporate local data or expert knowledge into their DUET simulations. The resulting data arrays are represented in standard NumPy [5] formats, which can then be processed manually by the user or conveniently exported to data formats accepted by QUIC-Fire or FIRETEC.

2.2. Software functionalities

The package consists of several modules, three of which we highlight here: (1) **inputs**, (2) **calibration**, and (3) **landfire**. The Inputs module facilitates loading, modifying, and writing validated DUET input files. Input parameters for the DUET program include the 3D fuel array dimensions and resolution, the number of years to simulate, and information on prevailing winds: speed, direction, and variability. The `from_directory()` classmethod automatically creates a representation of `duet.in` which can be conveniently modified and written to a file programmatically. Every parameter is validated upon write, meaning that any errors will be caught before the DUET simulation takes place. This involves type checking and ensuring inputs fall within accepted data ranges.

The Calibration module comprises the bulk of the current functionalities of `duet-tools`, providing methods for shifting and/or scaling DUET output values based on target ranges or distributions. Field data on fuel properties such as fuel loading may be collected and summarized as a range of values or as a distribution with a center and spread. The Calibration module can perform shifting and scaling based on either type of target, and can differentially apply those methods for different fuel parameters and/or fuel types. Targets are assigned using the `assign_targets()` function, which creates an object of class `Targets`. To assign targets based on a range of values, the "maxmin" method applies the following algorithm:

$$x'_i = \begin{cases} \min_target + \frac{x_i - \min(x^+)}{\max(x^+) - \min(x^+)} \cdot (\max_target - \min_target), & x_i > 0 \\ x_i, & x_i \leq 0 \end{cases}$$

where $x^+ = \{x_i \in x \mid x_i > 0\}$.

This shifts, then stretches or squeezes the values in the DUET output data array to match the provided range, without modifying the relative relationships between the data. Similarly, calibration based on mean and standard deviation is applied using the "meansd" argument and the following algorithm:

$$x'_i = \begin{cases} \max \left(0, \min_target + \frac{(x_i - \mu_x) \cdot sd_target}{\sigma_x} \right), & x_i > 0 \\ x_i, & x_i \leq 0 \end{cases}$$

where $x^+ = \{x_i \in x \mid x_i > 0\}$, $\mu_x = \text{mean}(x^+)$, $\sigma_x = \text{std}(x^+)$

The user may create one or multiple **Targets** objects which can then be set to different fuel parameters and fuel types. Briefly, a fuel parameter is one of the three required fuel inputs for the QUIC-Fire model: (1) fuel loading, (2) fuel moisture, or (3) fuel depth. A fuel type, on the other hand, corresponds a category of fuel simulated by DUET that each parameter may describe. DUET simulates litter fall from any species of tree represented in the DUET fuel grid input as well as growth of grass or other herbaceous plants. Upon creation of a **DuetRun** object, **duet-tools** groups together the litter from coniferous species, then the litter from deciduous species. This creates three main fuel types to which a **Targets** object may be set: (1) grass, (2) conifer litter, and (3) deciduous litter. Additionally, a target may be set for a combination of both litter types, or to all fuel types together.

The targets corresponding to a single fuel parameter can be set for one or more fuel types using **set_fuel_parameter()**, which creates an object of class **FuelParameter**. Once each desired fuel parameter has been set using targets for each desired fuel type, the **DuetRun** object can be calibrated using the **calibrate()** function. Crucially, any combination of fuel type and fuel parameter can be assigned targets using either calibration method. In practice, this means there is a single function, **calibrate()**, that handles every possible calibration specification. The flexibility to mix and match methods, fuel types, and fuel parameters is facilitated by the use of a small set of keyword arguments in the **assign_targets()** and **set_fuel_parameter()** functions. Running **calibrate()** on a **DuetRun** object produces a new **DuetRun** instance that stores the calibrated arrays. These arrays can be accessed or saved via the classmethods **to_numpy()** and **to_quicfire()**, the latter of which saves the calibrated arrays to a specified directory in the format expected by the QUIC-Fire model.

The **landfire** module supports calibration by offering a convenient source for calibration targets. The main functionality of the module is the execution of a data query of from the LANDFIRE database [6] to be used to calibrate DUET outputs. The module leverages the *landfire* python package [7] to download fuels data, then converts those data to calibration targets. Fuels data themselves are derived from the Scott and Burgan 40 Fire Behavior Fuel Models [8]. Future package development is poised incorporate additional data sources from LANDFIRE, such as the Fuel Characteristics and Classification System (FCCS) [9].

3. Illustrative examples

The main functionalities of `duet-tools` package are compatible with Python version 3.10+ and can be installed using PyPI. Since the `landfire` module requires dependencies that are only compatible with Python version 3.10, it is included as an optional package extra. Full documentation, how-to guides, example scripts, and issue tracking are hosted on GitHub.

3.1. DUET calibration workflow

The use of `duet-tools` is contingent upon the user possessing the results of a DUET simulation. Most often they will exist alongside a DUET input file. A common workflow will begin by creating a `DuetRun` object using `import_duet()`, which derives DUET data dimensions from the `duet.in` input file. If the input file is not present, but the data dimensions are known, the user can alternatively specify `import_duet_manual()`. Next, targets may be assigned using either of the two available calibration methods, and subsequently set to the desired fuel parameter(s). Once targets for each desired fuel parameter have been specified, the `calibrate()` function applies the calibration methods accordingly. A simple example of this process is shown in Listing 1 below.

```
1 # Import DUET outputs
2 duet_run = import_duet(directory=duet_path)
3
4 # Assign targets for each fuel type and fuel parameter
5 coniferous_loading = assign_targets(method="maxmin", max
   =5.0, min=0)
6 deciduous_loading = assign_targets(method="meansd", mean
   =0.5, sd=0.1)
7 grass_loading = assign_targets(method="meansd", mean=0.5,
   sd=0.25)
8
9 # Bring together fuel types for each parameter
10 loading_targets = set_fuel_parameter(
11     parameter="loading",
12     grass=grass_loading,
13     deciduous=deciduous_loading,
14     coniferous=coniferous_loading,
15 )
16
17 # Calibrate the DUET run
```

```

18 calibrated_duet = calibrate(duet_run=duet_run,
    fuel_parameter_targets=loading_targets)
19
20 # Export to a QUIC-Fire input file
21 calibrated_duet.to_quicfire(quicfire_path)

```

Listing 1: Example of a typical workflow using `duet-tools`, including importing a DUET run, setting calibration targets, calibrating fuel parameter values, and exporting to QUIC-Fire fuel arrays.

In Listing 1, the fuel loading parameter is calibrated using separately defined targets for each fuel type. Grass and deciduous litter loading are calibrated using their respective target means and standard deviations, whereas coniferous loading is calibrated based on a target value range. Figure 1 shows how each calibration shifts the magnitudes of fuel loading values for each fuel type, while retaining the relative relationships between values. In this example, coniferous litter and grass fuel loading have been calibrated to reflect lower loading than was predicted by DUET, whereas deciduous litter loading was increased. After these calibrations, overall loading was decreased, but with patterns corresponding to the modifications of the individual fuel types. The total litter loading is available to be exported as an input file to QUIC-Fire, or as a NumPy array for other applications.




Figure 1: Example of DUET calibration. The top row shows original fuel loading outputs from DUET for each fuel type; the bottom row shows calibrated results based on user-specified targets.

Fuel parameter targets may be assigned from LANDFIRE data in a similar fashion. Listing 2 shows the process of querying the LANDFIRE database to retrieve Scott and Burgan 40 Fuel Model data [8] for an area of interest. Spatial data defining an area of interest can be supplied as either a geojson or a polygon. For each fuel type (grass and/or litter; Fig. 2B), the values of the desired fuel parameters (Fig. 2C) are automatically assigned to a `Targets` object, which can then be used to differentially adjust DUET output values for those values (Fig. 2D-E).

```

1 # Import DUET outputs
2 duet_run = import_duet(directory=duet_path)
3
4 # Query the LANDFIRE database
5 landfire_query = query_landfire(
6     area_of_interest=aoi_geojson, year=2019, directory=
7     duet_path, input_epsg=4326
8 )
9
10 # Assign fuel targets from LANDFIRE query
11 grass_loading_targets = assign_targets_from_sb40(
12     landfire_query, "grass", "loading")
13 litter_loading_targets = assign_targets_from_sb40(
14     landfire_query, "litter", "loading")
15 landfire_loading = set_loading(
16     grass=grass_loading_targets, litter=
17     litter_loading_targets
18 )
19
20 # Calibrate the DUET run
21 calibrated_duet_landfire = calibrate(duet_run,
22     landfire_loading)

```

Listing 2: Example of DUET calibration using data from LANDFIRE [6]

4. Impact

Accurate descriptions of fuel and vegetation are essential for wildland fire modeling [10], which is an integral aspect of prescribed fire planning and training [11, 12] that supports land management [13], disaster mitigation [14], and species conservation [15]. Simulating realistic fire behavior using tools like QUIC-Fire and FIRETEC requires an understanding of the heterogeneity of fuel structures on a landscape [16, 17], but that heterogene-




Figure 2: Example of DUET calibration using Scott and Burgan 40 Fuel Model [8] data from the LANDFIRE database [6]. Original DUET output values (A) are assigned fuel parameter targets from a provided area of interest based on the mapped fuel type (B-C). Values for grass (D) and tree litter (E) may be calibrated separately and combined to produce a calibrated DUET output (F).

ity is difficult to characterize. Data fusion approaches that combine local data, broad-scale mapping, and ecological modeling can provide more realistic characterizations wildland fuel heterogeneity [18, 19]. By facilitating the fusion of modeled ecological processes from DUET, satellite-based gridded data from LANDFIRE, and direct input from users, `duet-tools` offers one of the first such approaches for surface fuel modeling.

Though there is utility in fine-scale, data-driven fire behavior modeling, user access to the models, their computational weight, and accurate fuel parameterization stand as barriers to operational use [20]. `duet-tools` helps ease the burden of fuel parameterization by providing a simple and convenient tool for harnessing a fuel modeling software that may otherwise be under-utilized. By enabling the programmatic management and modification of DUET inputs and outputs, `duet-tools` has already allowed it to be incorporated into the most robust 3D fuel modeling platform to date, FastFuels [18]. As a national-scale data platform, FastFuels’ coverage includes a wide variety of ecosystems, fuel types, and climate zones which can have vastly

different litter decay rates [21] and herbaceous growth patterns [22], or may experience a variety of management actions that may modify surface fuels. `duet-tools` provides a key mechanism for accounting for these differences by scaling DUET’s predictions to local data inputs, without compromising the spatial heterogeneity predictions that are central to DUET’s utility.

As the science of wildland fire management continues to expand its toolkit of computing technologies, tools and frameworks for harmonizing fuel data across scales are becoming more widespread [23, 24, 19]. The continued development of `duet-tools` can help it join the ranks as a way to facilitate this goal, especially as fuel structures and fire dynamics become more uncertain in a changing climate [25, 26, 27].

4.1. Limitations and Future Work

This article aims to introduce `duet-tools`, which is still in the early stages of its development. Currently, manipulation of DUET output files is limited to two simple calibration algorithms and 2-3 fuel types. Future work should incorporate more complex and place-based calibration parameters, such as seasonality [28], topography [29], and the addition of other types of surface fuel such as shrubs. Because local fuel data may not be available, further package development should focus on forging more connections to large geospatial datasets, including the expansion of the `landfire` module to access additional fuel layers, such as FCCS data [9]. Since these layers are generally compiled at a resolution of 20-30 m, the variability they describe is currently applied wholesale when calibrating the fine-scale variability simulated by DUET. Future package development could add the option to incorporate the broad-scale fuel heterogeneity described by broad-scale data while maintaining the fine-scale patterns predicted by DUET simulations. These improvements will help `duet-tools` play an integral role in streamlining the integration of DUET with fuels data platforms such as FastFuels [18].

5. Conclusions

`duet-tools` bridges a key usability gap between DUET and downstream fire behavior models. It simplifies DUET file management, enables automated calibration against data sources, and promotes transparent and reproducible workflows for surface fuel modeling.

Acknowledgements

Many thanks to Jenna McDanold for the introduction and continued development of the DUET program. I am also grateful for the guidance and men-

torship from Anthony Marcozzi, as well as the support from Scott Pokswinski and the rest of the NMC CAFES team. Special thanks to Rachel Loehman for encouraging and supporting the development of this tool. Funding for this project is provided by the Robert T. Stafford Disaster Relief and Emergency Assistance Act (42 U.S.C. 5121 et seq.) and supplemental funding acts for Federal disaster relief activities via the U.S. Geological Survey.

References

- [1] J. E. Keeley, Ecological foundations for fire management in North American forest and shrubland ecosystems, Vol. 779, US Department of Agriculture, Forest Service, Pacific Northwest Research Station, 2009.
URL https://books.google.com/books?hl=en&lr=&id=4_Y4IT6YBX8C&oi=fnd&pg=PA1&dq=role+of+fire+in+north+american+forests&ots=80ARlcD01P&sig=1VNYsm1xxHeTZr0RC-xBgMsgfLU
- [2] R. Linn, J. Reisner, J. J. Colman, J. Winterkamp, Studying wild-fire behavior using FIRETEC, *International Journal of Wildland Fire* 11 (4) (2002) 233–246, publisher: CSIRO PUBLISHING. doi:10.1071/wf02007.
URL <https://www.publish.csiro.au/wf/wf02007>
- [3] R. R. Linn, S. L. Goodrick, S. Brambilla, M. J. Brown, R. S. Middleton, J. J. O’Brien, J. K. Hiers, QUIC-fire: A fast-running simulation tool for prescribed fire planning, *Environmental Modelling & Software* 125 (2020) 104616, publisher: Elsevier.
URL <https://www.sciencedirect.com/science/article/pii/S1364815219307388>
- [4] J. S. McDanold, R. R. Linn, A. K. Jonko, A. L. Atchley, S. L. Goodrick, J. K. Hiers, C. M. Hoffman, E. L. Loudermilk, J. J. O’Brien, R. A. Parsons, C. H. Sieg, J. A. Oliveto, DUET - Distribution of Understory using Elliptical Transport: A mechanistic model of leaf litter and herbaceous spatial distribution based on tree canopy structure, *Ecological Modelling* 483 (2023) 110425. doi:10.1016/j.ecolmodel.2023.110425.
URL <https://www.sciencedirect.com/science/article/pii/S0304380023001564>
- [5] C. R. Harris, K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau, E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van Kerkwijk, M. Brett, A. Haldane,

- J. F. del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant, K. Shepard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke, T. E. Oliphant, Array programming with NumPy, *Nature* 585 (7825) (2020) 357–362. doi:10.1038/s41586-020-2649-2.
URL <https://doi.org/10.1038/s41586-020-2649-2>
- [6] I. P. La Puma, LANDFIRE technical documentation, Tech. Rep. 2023-1045, Reston, VA (2023). doi:10.3133/ofr20231045.
URL <https://pubs.usgs.gov/publication/ofr20231045>
- [7] J. Clark, landfire-python (2023).
URL <https://landfire-python.firesci.io/en/latest/index.html#>
- [8] J. H. Scott, R. E. Burgan, Standard Fire Behavior Fuel Models: A Comprehensive Set for Use with Rothermel’s Surface Fire Spread Model, Tech. rep. (2005).
URL <http://www.fs.fed.us/rm>
- [9] R. D. Ottmar, D. V. Sandberg, C. L. Riccardi, S. J. Prichard, An overview of the Fuel Characteristic Classification System — Quantifying, classifying, and creating fuelbeds for resource planning This article is one of a selection of papers published in the Special Forum on the Fuel Characteristic Classification System., *Canadian Journal of Forest Research* 37 (12) (2007) 2383–2393. doi:10.1139/X07-077.
URL <http://www.nrcresearchpress.com/doi/10.1139/X07-077>
- [10] E. L. Loudermilk, J. J. O’Brien, S. L. Goodrick, R. R. Linn, N. S. Skowronski, J. K. Hiers, Vegetation’s influence on fire behavior goes beyond just being fuel, *Fire Ecology* 18 (1) (2022) 9, s42408-022-00132-9. doi:10.1186/s42408-022-00132-9.
URL <https://fireecology.springeropen.com/articles/10.1186/s42408-022-00132-9>
- [11] R. Parsons, L. Wells, A. Marcozzi, R. R. Linn, J. K. Hiers, F. Pimont, K. Riley, I. Altintas, S. Flanary, Going 3D With Fuel and Fire Modeling: FastFuels and QUIC-Fire, *Fire Management Today* 81 (1) (2023) 49–51.
URL <https://www.fs.usda.gov/sites/default/files/fire-management-today/fmt-vol-81-n1.pdf>
- [12] J. K. Hiers, J. J. O’Brien, J. M. Varner, B. W. Butler, M. Dickinson, J. Furman, M. Gallagher, D. Godwin, S. L. Goodrick, S. M.

- Hood, A. Hudak, L. N. Kobziar, R. Linn, E. L. Loudermilk, S. McCaffrey, K. Robertson, E. M. Rowell, N. Skowronski, A. C. Watts, K. M. Yedinak, Prescribed fire science: the case for a refined research agenda, *Fire Ecology* 16 (1) (2020) 11, s42408-020-0070-8. doi:10.1186/s42408-020-0070-8.
URL <https://fireecology.springeropen.com/articles/10.1186/s42408-020-0070-8>
- [13] S. A. Drury, H. M. Rauscher, E. M. Banwell, S. Huang, T. L. Lavezzo, The Interagency Fuels Treatment Decision Support System: Functionality for Fuels Treatment Planning, *Fire Ecology* 12 (1) (2016) 103–123. doi:10.4996/fireecology.1201103.
URL <https://fireecology.springeropen.com/articles/10.4996/fireecology.1201103>
- [14] D. M. Staley, J. A. Negri, J. W. Kean, J. L. Laber, A. C. Tillery, A. M. Youberg, Updated logistic regression equations for the calculation of post-fire debris-flow likelihood in the western United States, Tech. rep., US Geological Survey (2016).
URL <https://pubs.usgs.gov/publication/ofr20161106>
- [15] M. S. Lucash, S. Weiss, M. J. Duveneck, R. M. Scheller, Managing for red-cockaded woodpeckers is more complicated under climate change, *The Journal of Wildlife Management* 86 (8) (2022) e22309, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/jwmg.22309>. doi:10.1002/jwmg.22309.
URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/jwmg.22309>
- [16] S. R. Bonner, C. M. Hoffman, R. R. Linn, W. T. Tinkham, A. L. Atchley, C. H. Sieg, J. M. Varner, J. J. O'Brien, J. K. Hiers, Forest structural complexity and ignition pattern influence simulated prescribed fire effects, *Fire Ecology* 20 (1) (2024) 82. doi:10.1186/s42408-024-00314-7.
URL <https://fireecology.springeropen.com/articles/10.1186/s42408-024-00314-7>
- [17] R. A. Parsons, W. E. Mell, P. McCauley, Linking 3D spatial models of fuels and fire: effects of spatial heterogeneity on fire behavior, *Ecological Modelling* 222 (3) (2011) 679–691, publisher: Elsevier.
URL <https://www.sciencedirect.com/science/article/pii/S0304380010005831>

- [18] A. Marcozzi, L. Wells, R. Parsons, E. Mueller, R. Linn, J. K. Hiers, FastFuels: Advancing wildland fire modeling with high-resolution 3D fuel data and data assimilation, *Environmental Modelling & Software* 183 (2025) 106214. doi:10.1016/j.envsoft.2024.106214.
URL <https://www.sciencedirect.com/science/article/pii/S1364815224002755>
- [19] N. J. Tutland, A. P. Wion, C. J. May, G. C. Hutchings, H. A. Nowak, J. R. Gattiker, J. K. Hiers, R. R. Linn, S. M. Pokswinski, E. Q. Margolis, Representing 3-dimensional fuels for physics-based fire behavior models: a general framework and case study in a type-converted post-fire shrubfield, *Fire Ecology* 21 (1) (2025) 43. doi:10.1186/s42408-025-00383-2.
URL <https://fireecology.springeropen.com/articles/10.1186/s42408-025-00383-2>
- [20] W. Mell, R. Linn, FIRETEC and WFDS Modeling of Fire Behavior and Smoke in Support of FASMEE JFSP PROJECT ID: 16-4-05-1 (2017).
URL https://www.academia.edu/download/76406618/16-4-05-1_final_report.pdf
- [21] J. M. Blair, R. W. Parmelee, M. H. Beare, Decay Rates, Nitrogen Fluxes, and Decomposer Communities of Single- and Mixed-Species Foliar Litter, *Ecology* 71 (5) (1990) 1976–1985, eprint: <https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.2307/1937606>. doi:10.2307/1937606.
URL <https://onlinelibrary.wiley.com/doi/abs/10.2307/1937606>
- [22] P. B. Reich, D. W. Peterson, D. A. Wedin, K. Wragge, FIRE AND VEGETATION EFFECTS ON PRODUCTIVITY AND NITROGEN CYCLING ACROSS A FOREST–GRASSLAND CONTINUUM, *Ecology* 82 (6) (2001) 1703–1719. doi:10.1890/0012-9658(2001)082[1703:FAVEOP]2.0.CO;2.
URL [http://doi.wiley.com/10.1890/0012-9658\(2001\)082\[1703:FAVEOP\]2.0.CO;2](http://doi.wiley.com/10.1890/0012-9658(2001)082[1703:FAVEOP]2.0.CO;2)
- [23] D. Roten, L. Wells, D. Crawl, R. A. Parsons, A. Marcozzi, R. R. Linn, K. Hiers, I. Altintas, TrueTrees: A Scalable Workflow for the Integration of Airborne LiDAR Scanning Data into Fuel Models for Prescribed Fire Simulations, in: 2023 IEEE 19th International Conference on e-Science (e-Science), IEEE, 2023, pp. 1–10.

- URL <https://ieeexplore.ieee.org/abstract/document/10254923/>
- [24] I. G. Moreno, I. Nealey, D. Roten, M. H. Nguyen, D. Crawl, K. O’Laughlin, M. Floca, S. Pokswinski, I. Altintas, Visualization and Labeling of Terrestrial LiDAR Data for Three-Dimensional Fuel Classification, in: 2023 IEEE 19th International Conference on e-Science (e-Science), IEEE, Limassol, Cyprus, 2023, pp. 1–2.
URL <https://ieeexplore.ieee.org/abstract/document/10254841/>
- [25] K. T. Davis, P. E. Higuera, S. Z. Dobrowski, S. A. Parks, J. T. Abatzoglou, M. T. Rother, T. T. Veblen, Fire-catalyzed vegetation shifts in ponderosa pine and Douglas-fir forests of the western United States, *Environmental Research Letters* 15 (10) (2020) 1040b8, publisher: IOP Publishing. doi:10.1088/1748-9326/abb9df.
URL <https://dx.doi.org/10.1088/1748-9326/abb9df>
- [26] J. D. Coop, S. A. Parks, C. S. Stevens-Rumann, S. D. Crausbay, P. E. Higuera, M. D. Hurteau, A. Tepley, E. Whitman, T. Assal, B. M. Collins, K. T. Davis, S. Dobrowski, D. A. Falk, P. J. Fornwalt, P. Z. Fulé, B. J. Harvey, V. R. Kane, C. E. Littlefield, E. Q. Margolis, M. North, M.-A. Parisien, S. Prichard, K. C. Rodman, Wildfire-Driven Forest Conversion in Western North American Landscapes, *BioScience* 70 (8) (2020) 659–673. doi:10.1093/biosci/biaa061.
URL <https://doi.org/10.1093/biosci/biaa061>
- [27] A. P. Wion, J. T. Stevens, K. Beeley, R. Oertel, E. Q. Margolis, C. D. Allen, Multidecadal vegetation transformations of a New Mexico ponderosa pine landscape after severe fires and aerial seeding, *Ecological Applications* 34 (6) (2024) e3008, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/eap.3008>. doi:10.1002/eap.3008.
URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/eap.3008>
- [28] Y. Li, Y. Zhang, X. Quan, B. He, S. Veraverbeke, Z. Liao, T. A. J. Janssen, Estimating forest litter fuel load by integrating remotely sensed foliage phenology and modeled litter decomposition, *Remote Sensing of Environment* 317 (2025) 114526. doi:10.1016/j.rse.2024.114526.
URL <https://www.sciencedirect.com/science/article/pii/S0034425724005522>

- [29] P. Nyman, D. Metzen, P. J. Noske, P. N. Lane, G. J. Sheridan, Quantifying the effects of topographic aspect on water content and temperature in fine surface fuel, *International Journal of Wildland Fire* 24 (8) (2015) 1129–1142, publisher: CSIRO Publishing.
URL <https://www.publish.csiro.au/wf/wf14195>