

¹ Mixing cover crops suppresses weeds and roto-till
² improves urban soil compaction and infiltration

³ Naim Edwards ^{a*} Nicholas Medina ^b Elizabeth Asker ^a

⁴ ^a Agriculture and Natural Resources, Michigan State University, 16745 Lamphere St, Detroit, MI, USA 48219; ^b Ecology and
⁵ Evolutionary Biology, University of Michigan, 1105 N University Ave, Ann Arbor, MI, USA

⁶

⁷

⁸

⁹ **Keywords:** urban agriculture; soil compaction; weed suppression; roto-till, cover crop mix;
¹⁰ soil infiltration

¹¹

¹²

¹³

¹⁴ ***Corresponding author:** Naim Edwards, edwar649@msu.edu

¹⁵ Author emails: edwar649@msu.edu, nmedina@umich.edu, askereli@msu.edu

¹⁶ **Highlights:**

- ¹⁷ • Roto-till improves urban soil compaction and infiltration vs. no-till
- ¹⁸ • Tractor-till improves compaction but not infiltration and also increases weeds
- ¹⁹ • Cover crop mixes suppress weeds
- ²⁰ • Forage radish yield not affected by till or cover crop mixes
- ²¹ • Roto-till and cover crop mixes help improve soils for urban agriculture

²² Abstract

²³ Urban soils have been degraded by decades of industrial activities, but they also represent
²⁴ opportunities to improve food sovereignty for urban residents practicing urban agriculture.
²⁵ Urban growers often use varying practices of compost, tillage, and cover cropping, yet fur-
²⁶ ther integrated approaches could facilitate by model analyses of how different practices
²⁷ may compare or complement each other. This study examined how tillage methods repre-
²⁸ senting various intensities and cover crop mixes targeting different functions affected agri-
²⁹ cultural variables including soil compaction, water infiltration rate, herbaceous weedy plant
³⁰ pressure, and crop yield. Results showed that both roto- and tractor-till significantly af-
³¹ fected compaction but not yield compared to no-till, and roto-till also improved infiltration,
³² while tractor-till reached deeper soils but allowed denser weed growth. Mixing sorghum-
³³ sudangrass, buckwheat, and cowpea cover crops significantly reduced weed pressure com-
³⁴ pared to other mixtures, and perennials affected compaction but not soil water infiltration
³⁵ under no-till. These results reveal that medium-intensity tillage may offer more balanced
³⁶ trade-offs for initial management, and that cover crops can help reduce weeds under low-till
³⁷ strategies. Overall this study offers evidence detailing effects of various tillage and cover
³⁸ crop styles that can be of use for smallholder urban growers.

³⁹ **1 Introduction**

⁴⁰ Urban soils could improve the livelihoods of most of the world (*Acuto et al., 2018*) by helping
⁴¹ adapt to climate change, slowing erosion and storm-water runoff management, and promot-
⁴² ing local forestry (*Pavao-Zuckerman, 2008*), however, many urban soils are degraded for
⁴³ agriculture, after decades of industrial use, including sealing and structural engineering (*Lal*
⁴⁴ *et al., 2015*). Urban soil issues are notable in post-industrial cities of the mid-western USA,
⁴⁵ where thousands of vacant lots still show high compaction, pH, and chemical contamination
⁴⁶ (*Beniston et al., 2016*). These degraded urban soils also have low organic matter, but also
⁴⁷ being far from carbon saturation (*Stewart et al., 2007*) can increase potential responses to
⁴⁸ intervention (*Kumar and Hundal, 2016; Kuzyakov and Zamanian, 2019*). Single strategies
⁴⁹ like adding compost are popular, and indeed are beneficial for various physical, chemical, and
⁵⁰ biological properties (*Cogger, 2005*), but they also can have limiting side effects like excess
⁵¹ phosphorus (*Small et al., 2019*), which in turn highlights the benefits of simultaneous strate-
⁵² gies, like cover cropping and occasional tillage, that could better target multi-functionality
⁵³ (*Blesh, 2017; Garbach et al., 2017; O'Riordan et al., 2021; Sircely and Naeem, 2012; Tresch*
⁵⁴ *et al., 2018*). Urban agriculture has spread as a response to diverse community needs (*Lon-*
⁵⁵ *don et al., 2021*), from systemic food insecurity to schooling access and labor imbalances,
⁵⁶ and also widely engages non-profits, politicians, and individuals in environmental steward-
⁵⁷ ship addressing public health issues like pollution (*Block et al., 2012; Clendenning et al.,*
⁵⁸ *2016; García-Sempere et al., 2019; Siebert, 2020*). Community-led infrastructure governing
⁵⁹ vacant land additionally means that urban growers invest much of their personal money,
⁶⁰ time, and other limited resources into lot preparation for initial cultivation (*Daftary-Steel et*
⁶¹ *al., 2015*), but often need to move ahead with varying models of holistic approaches (*Gross-*
⁶² *man, 2003*) to jump-starting cultivation in urban soils with industrial legacy effects (*Wade*
⁶³ *et al., 2021*), jeopardizing regionally high yields (*McDougall et al., 2019*).
⁶⁴ Mechanized tilling is one strategy that can offer short-term benefits, but also at the cost

of long-term soil health, especially as mechanical intensity increases. In the short term, tilling can improve soil porosity to lower soil bulk density (*Badalíková, 2010; Hill et al., 1985*), improve nutrient availability (*Wolkowski, 1990*), and control weeds (*Barberi and Lo Cascio, 2001; Cordeau et al., 2020*), thereby also likely improving water infiltration and drainage that can facilitate faster seeding and early crop establishment (*Monti et al., 2001*). However, in the long term (i.e. over five years), soil aggregates can weaken (*Catania et al., 2018; Six et al., 2002*) lead to faster soil erosion (*Richter, 2021*), eventually increasing grower dependency on intense tillage to maintain previous yields (*de Cárcer et al., 2019*), all of which resemble causes of the USA Dust Bowl and even the fall of ancient civilizations (*Amundson et al., 2015; Lal, 2007; Montgomery, 2007*). To combat degradation, no-till and minimal-till have been supported as sustainable alternatives to industrial agri-business farming (*Roger-Estrade et al., 2010; Wang et al., 2006*), although, continuing research is still needed to address different challenges like more weed pressure (*Anderson, 2007*). Since urban growers already have limited access to machinery (*Daniel, 2007*), yet given the short-term benefits of tillage for quick initial productivity, community sharing systems have been set up for tractors and rotary implements, which can lead to mixed or variable management strategies being adopted for urban soil cultivation, which are in need to further study (*Bazzoffi, 1998; Materechera, 2009*).

Cover cropping is another regenerative agriculture practice with old origins, but whose lasting benefits are increasingly recognized (*Perez, 2021; Richter, 2021*), however, more studies could go beyond single species to complementary species mixtures. Cover crops are named so because they cover fallow soils, also continuing root activity and limiting erosion (*García-González et al., 2018*), but benefits can vary by species used. For example, legumes like cowpea (or black-eyed peas, *Vigna unguiculata* subsp. *unguiculata*), clovers (*Trifolium sp.*), and hairy vetch (*Vicia villosa*) have symbiotic root bacteria that fix nitrogen from the air into soil pores where it becomes bioavailable to plants (*Grossman et al., 2005*). Somewhat similarly, buckwheat (*Fagopyrum esculentum*) helps scavenge soil phosphorus (*Possinger et*

92 *al., 2013*), often a limiting macro-nutrient in clay soils (*Mori et al., 2022*), which could also be
93 combined with phosphorus-rich compost to alleviate recurring soil phosphorus deficiencies.
94 Other plants, including grasses like sorghum (*Sorghum bicolor*) can grow deep roots with
95 chemical defenses, called allelopathy, that harm other weed roots (*Weston et al., 1989*).
96 Overall, cover cropping may also increase soil organic matter through complex processes
97 (*King, 2020*), though few studies show direct correlations between soil organic matter and
98 yield (*Oldfield et al., 2019*). Organic yet industrial farms can benefit from specific cover
99 crops, but their mechanization also limits their use to monoculture, where as small urban
100 agriculture can make use of labor that replaces machinery to study new cover crop mixture
101 designs that could accelerate early cultivation efforts. Cover crop mixtures generally remain
102 understudied empirically in agriculture (*Baraibar et al., 2020; Bedoussac et al., 2015; Bourke
103 et al., 2021; Mead and Riley, 1981*), but it could be hypothesized that combining sorghum,
104 cowpea, and buckwheat together would improve soil nitrogen, phosphorus, and weed control,
105 via their root symbioses and chemical defenses. In general, integrated approaches to small-
106 scale urban agriculture could be useful internationally (*Stewart et al., 2013*), but tailored
107 research that informs grower decision-making remains diffuse.


108 In this study, we investigated how different tillage techniques and cover crop species mixes,
109 representing various possible integrated management strategies, affect urban soil functions
110 for agriculture. Tillage methods studied ranged from low intensity using a broadfork to
111 high intensity using a tractor and implements. Additionally, cover crop species mixes were
112 chosen based on target functions including reducing compaction, suppressing weeds, and
113 perenniability (i.e. potential for sustainable re-growth). We hypothesized that both tillage and
114 cover crop mixes would confer similar benefits to soil functions, which would also translate
115 to affect weed pressure and yield. Accordingly, we predicted that roto-till, a moderate-
116 intensity option, would best balance compaction and weed pressure benefits, deepening where
117 soil hardpan layers occur that limit root penetration, and thereby also increase soil water
118 infiltration rates, along with reducing weed cover, density, and diversity. We also expected

¹¹⁹ that the cover crop mix designed against soil compaction would have the deepest depth to
¹²⁰ soil harpan, along with the fastest water infiltration rates compared to other mixes, mostly
¹²¹ due to the deep rooting potential of forage radish (*Raphanus sativus var. longipinnatus*) and
¹²² ryegrass (*Secale cereale*). Finally, we expected that the cover crop mix designed for weed
¹²³ suppression would experience the lowest local weed cover, density, and diversity, due to
¹²⁴ allelopathic chemical defense traits from buckwheat (*Fagopyrum esculentum*) and surghum-
¹²⁵ sudangrass (*Sorghum bicolor x Sorghum bicolor var. sudanese*).

¹²⁶ 2 Methods

¹²⁷ 2.1 Site

¹²⁸ The study site was located at the Michigan State University (MSU) - Detroit Partnership for
¹²⁹ Food, Learning, and Innovation (DPFLI) (42.4, -83.3), a 1.6-ha (4 acres) extension facility
¹³⁰ dedicated to urban agriculture and engaging with local small-scale growers in Detroit, MI,
¹³¹ USA. The climate is temperate with four seasons, with mean annual temperature of ~9.5 C
¹³² (49.1 F) and precipitation at ~787 mm (31 in) (ncdc.noaa.gov). The site was formerly a school
¹³³ building and associated playground until 2016 when it was demolished after closing due to
¹³⁴ low funding and the land became vacant. The habitat is ~1.2 km (~0.8 mi) away from a small
¹³⁵ river, conferring some wetland ecosystem properties like denser soils. It is also surrounded
¹³⁶ by sealed sidewalk and small roads on all four sides, which likely affects runoff and drainage



137 patterns (Fig ??a).

138 Site soils can be classified as Technosols (Fig ??b), given that large metal artifacts can be
 139 found throughout various profiles (*FAO, 2014*), from when the area was filled in with nearby
 140 soils during highway road construction, as was common in mid-western USA industrial man-
 141 ufacturing cities many decades ago in the *1960s* (*Beniston et al., 2016*). Accordingly, the
 142 growing area has both a finer- and coarser-textured side (Fig ??a), and this study was done
 143 on the side with consistent clay of ~37%. Topsoil A horizons are 1-2" (<5 cm) deep, and
 144 subsoil B horizons can be >30.5 cm (1 ft) deep, with a muted yellow color 10YR 8/4 (Fig
 145 ??b). A baseline site-level soil lab assessment determined that the top *4 in* (10 cm) of soils
 146 around the site together have relatively good organic matter at ~2.5 ± 0.3% and nutrient
 147 levels, including concentrations of heavy metals like lead and arsenic which were present
 148 below harmful government human-contact standards (cfpub.epa.gov/ecotox). Site soils were

¹⁴⁹ also assessed to have decent but sub-optimal CO_2 respiration rates of 0.2 ± 0.04 mg per day
¹⁵⁰ (Table ??). Initial main concerns limiting productivity include high alkaline pH of $8.1 \pm$
¹⁵¹ 0.1, lowering availability of existing nutrients, as well as weak aggregate stability of $19 \pm$
¹⁵² 4.4, leading to concerns with aeration, infiltration, rooting, crusting when dry, erosion, and
¹⁵³ runoff (Table ??).

Table 1: Baseline Soil Health Assessment (Cornell,
 Ithaca, NY, USA)

Kind	Variable	Median (n=10)	Deviation	Descriptor
Biological	Organic Matter (%)	2.5	0.3	Very Low
	Respiration (mg per day)	0.2	0.0	Medium
Physical	Aggregate Stability (%)	19.0	4.4	Very Low
Chemical	pH	8.1	0.1	Poor
	Phosphorus (ppm)	2.2	1.0	Medium
	Potassium (ppm)	103.8	36.3	Optimal
	Iron (ppm)	6.0	4.4	Optimal
	Magnesium (ppm)	463.6	24.9	Optimal
	Manganese (ppm)	42.1	4.9	Optimal
	Zinc (ppm)	3.8	2.9	Optimal
	Heavy metals (Pb, Al, As, Cu)	-	-	Safe

¹⁵⁴ 2.2 Design

¹⁵⁵ The study area was a $278 m^2$ ($2992.4 ft^2$) section on the East side of the site under the
¹⁵⁶ former school building that was divided into 36 separate $4.6 m^2$ ($49.5 ft^2$) plots in nine rows
¹⁵⁷ and four columns (Fig ??c). Tillage groups spanned the nine columns in adjacent groups of
¹⁵⁸ three, while cover crop mix treatments spanned the rows with one row per cover crop mix,

¹⁵⁹ totaling 36 plots, or 12 plots per tillage group and nine plots per cover crop mix. Before
¹⁶⁰ applying treatments, approximately 0.2 m^3 (8.5 ft^3) of compost was incorporated into each
¹⁶¹ plot.

¹⁶² Tillage treatments represented methods of increasing intensity available for small scale agri-
¹⁶³ culture, also varying in cost, machinery needed, and sometimes grower preferences (*Drugova*
¹⁶⁴ *et al.*, 2022). Specifically, treatments included no-till with a broadfork (*NT*), roto-tiller (*RT*),
¹⁶⁵ and tractor-till (*TT*) with implements. Tractor-till plots were worked with a subsoiler, mold-
¹⁶⁶ board plow, and roto-tiller attached to a tractor up to 30.5 cm (1 ft) deep. Roto-till plots
¹⁶⁷ were treated with a rototiller implement up to 20 cm (7.9 in) deep. Lastly, no-till plots were
¹⁶⁸ worked with only a broadfork up to 10 cm (3.9 in) deep. All tilling was done once early in
¹⁶⁹ the season after one typical compost application and before planting cover crops.

¹⁷⁰ Cover crop mixes were designed primarily based on plants associated with targeted bene-
¹⁷¹ fits, and as possible, relative simplicity of re-seeding and winter-kill (e.g. more heat tolerant)
¹⁷² (*Clark*, 2007). Three mixes were designed to target three functions, with each mix containing
¹⁷³ three different plant species (Table ??). The mix specifically designed to alleviate compaction
¹⁷⁴ generally focused on plants with roots that tend to penetrate and loosen soil well, and ulti-
¹⁷⁵ mately included crimson clover (*Trifolium incarnatum*), forage radish (*Raphanus sativus var.*
¹⁷⁶ *longipinnatus*), and cereal ryegrass (*Secale cereale*). The mix targeting weed suppression
¹⁷⁷ included heat- and drought-tolerant crops that tend to grow rapidly, allowing them to out-
¹⁷⁸ compete other plants—the taxa chosen were sorghum-sudangrass (*Sorghum bicolor x Sorghum*
¹⁷⁹ *bicolor var. sudanese*), cowpea/black-eyed pea (*Vigna unguiculata subsp. *unguiculata**), and
¹⁸⁰ buckwheat (*Fagopyrum esculentum*). Lastly, a mix was dedicated to perennial cover crops,
¹⁸¹ which in contrast to annuals can survive the winter and thus tend to accumulate biomass
¹⁸² and establish before spring weeds—this mix included hairy vetch (*Vicia villosa*), red clover
¹⁸³ (*Trifolium pratense*), and wheat (*Triticum aestivum*). We also had a null control group
¹⁸⁴ consisting of established vegetation within the plot, where no additional seeds were sown.

Table 2: Cover crop mixes

Function	Plants	Binomial
Weed Suppression	Sorghum-Sudangrass	Sorghum bicolor x S. bicolor var. sudanese
	Cowpea/Black-Eyed Pea	Vigna unguiculata subsp. unguiculata
	Buckwheat	Fagopyrum esculentum
Perennial	Hairy Vetch	Vicia villosa
	Red Clover	Trifolium pratense
	Wheat	Triticum aestivum
Compaction	Forage Radish	Raphanus sativus var. longipinnatus
	Crimson Clover	Trifolium incarnatum
	Cereal Ryegrass	Secale cereale
Null	Existing vegetation (no manipulation)	-

¹⁸⁵ 2.3 Sampling

¹⁸⁶ Soil compaction was measured with a penetrometer (*AgraTronix #08180*) in four randomly
¹⁸⁷ selected spots within each quarter of every plot, as the depth where resistance was 2 MPa
¹⁸⁸ (290.1 psi, *lbs in⁻²*), which is considered hardpan that roots typically cannot penetrate (*Cor-*
¹⁸⁹ *rea et al., 2019*). Measurements were recorded to the nearest 2.5 cm (1 inch) on dry days.

¹⁹⁰ Soil water infiltration down to 10 cm (8.75 in) depth was measured using a 16.5 (9.5 in)
¹⁹¹ wide aluminum cylinder, set away from dense vegetation and any impeding large roots, and
¹⁹² recording the time up to 160 sec for 1 L (32 fl oz) to pass through, representing a typical
¹⁹³ local rainfall onto ~0.10 m² (~1 ft²) of soil area (waterdata.usgs.gov).

¹⁹⁴ Weed pressure was measured using percent cover, richness, and density, following similar
¹⁹⁵ studies (*Storkey and Neve, 2018*). Weed cover was estimated as the total proportion of plot

area covered by any weed biomass, descretized into intervals of ten. Weed richness, a measure of diversity, was recorded by counting the number of unique morphospecies observed in each plot. Finally, weed density was measured as the number of stems of either of the two most abundant weed taxa, pigweed (*Amaranthus viridis*) and velvetleaf (*Abutilon theophrasti*), also descretized into intervals of ten up to 50 stems per plot.

Five forage radish (*Brassica Raphanus sativus var. longipinnatus*) roots were randomly selected from each plot in the compaction treatment and measured for length, individually, and wet weight, as a cluster. The length of a radish root was measured from the hypocotyl, or root cap, to where the root became ~6.3 mm (~1/4 in) wide.

Sampling was done in July and October 2019 and the following Spring.

2.4 Statistics

Field space limited strict plot replication for treatment combinations ($n=3$), and thus inference from advanced nested mixed models (*Silk et al., 2020*), so analysis focused on specific hypotheses tested using simpler, more conservative non-parametric tests that make few underlying assumptions about data and thus appropriate for data with lower replication. Kruskal-Wallis tests were run for tillage and cover crop treatments separately, with alpha corrections from 0.05 to 0.01 under multiple comparisons to descriptively parse any treatment interactions, and overall significant treatment effects were followed up by post-hoc Wilcoxon pairwise tests with Holm-corrected p-value adjustments. All data were centered at plot-level medians, often more robust than means, and where applicable pooled across sampling times given no preliminary significant variation along this axis, together with minimal relevance to focal hypotheses in field studies (*Davies and Gray, 2015*), and was a general solution to uneven sampling across response variables, minimally increasing statistical power ($n>3-6$). For clarity, results figures were designed to reflect statistical models and grouping transparently. Significant treatment effect sizes were estimated with η^2 (*Tomczak and Tomczak,*

221 2014) and raw median differences at finer pairwise levels. All calculations and analyzes
222 were done in R version 4.2.0 (2022-04-22) (*R Core Team, 2022*) with useful functions from
223 the packages *tidyverse* 1.3.1 (*Wickham et al., 2019*), *rstatix* 0.7.0 and *ggpubr* 0.4.0 (*Kassam-*
224 *bara, 2021*). Data and code are stored at github.com/nmedina17/must, documented using R
225 packages *here* 1.0.1 (*Müller, 2020*), *bookdown* 0.26 (*Xie, 2022a*), *measurements* 1.4.0 (*Birk,*
226 *2019*), *taxize* 0.9.100 (*Chamberlain et al., 2020*), *knitr* 1.39 (*Xie, 2022b*), and *rmarkdown*
227 2.14 (*Allaire et al., 2022*).

228 3 Results

229 3.1 Compaction

230 Compaction was affected significantly overall by tillage treatments ($H = 38.2$, $df = 2$, n
231 = 72, $p = <0.0001$) by ~52.4% across cover crop treatments (Fig 1a). Tractor-till had
232 the largest significant effect on depth to hardpan compared to no-till ($p_{adj} = <0.0001$),
233 deepening the depth to hardpan by ~9.4 cm (3.7 in, or ~83.3%) compared to no-till, down
234 to $\sim 20.6 \pm 4.6$ cm (8.1 ± 1.8 in) across all cover crop mixes. Roto-till also had a marginally
235 significant effect on depth to hardpan compared to no-till ($p_{adj} = 0.1$), deepening the depth
236 to hardpan by ~9.4 cm (3.7 in, or ~83.3%) compared to no-till, down to $\sim 13.8 \pm 1.9$ cm
237 (5.4 ± 0.7 in). The overall effect from tillage stemmed from significant effects among the
238 perennial ($p_{adj} = <0.01$) and weed suppression ($p_{adj} = <0.01$) mixes (Fig 1a). The
239 effect of roto-till was more pronounced in the perennial mix ($p_{adj} = <0.01$), where depth
240 to hardpan was about twice as deep as in no-till plots (Fig 1a). There was also a significant
241 difference of ~6.9 cm (2.7 in, or ~50%) between tractor- and roto-till among all cover crop
242 mixes (Fig 1a).

243 Compaction was not affected by cover crops among tillage groups overall ($H = 2$, $df = 3$,
244 $n = 72$, $p = 0.57$), but was significantly affected by cover crops specifically under no-till




Figure 1: Compaction data (a) by tillage, and (b) cover crop mix. Gray dots show plot medians and black point ranges show group mean \pm 1 std error and may be small. Significant pairwise post-hoc Wilcoxon test outcomes shown (***) $p < 0.0001$, ** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, *, $p < 0.1$, ' $p > 0.1$ or ns)

²⁴⁵ conditions ($p_{adj} = <0.01$) (Fig 1b). Under no-till, the perennial mix had significantly
²⁴⁶ shallower depth to hardpan compared to both null ($p_{adj} = <0.01$) and weed suppression
²⁴⁷ mixes ($p_{adj} = <0.01$), raising the depth to hardpan by ~ 2.5 cm (1 in, or $\sim 16.7\%$) compared
²⁴⁸ to each mix, up to $\sim 12.5 \pm 7.4$ cm (4.9 ± 2.9 in) below the soil surface (Fig 1b).

²⁴⁹ 3.2 Infiltration

²⁵⁰ Soil infiltration was significantly affected by tillage ($H = 8.5$, $df = 2$, $n = 48$, $p = 0.01$)
²⁵¹ and marginally significantly by cover crop mix ($H = 5.9$, $df = 3$, $n = 48$, $p = 0.1$) (Fig
²⁵² 2). Roto-till had significantly faster infiltration compared to no-till ($p_{adj} = 0.027$) and
²⁵³ marginally significantly compared to tractor-till ($p_{adj} = 0.1$), speeding up infiltration by
²⁵⁴ $\sim 14.5\%$ compared to each tillage groups, up to $\sim 13.4 \pm 10.7$ mL per sec (0.2 ± 0.2 gal per
²⁵⁵ min) (Fig 2a).

²⁵⁶ 3.3 Weed pressure

²⁵⁷ Weed density was overall significantly affected by tillage ($H = 6.5$, $df = 2$, $n = 72$, $p =$
²⁵⁸ 0.039) by $\sim 25.1\%$, although weed cover ($H = 1.6$, $df = 2$, $n = 72$, $p = 0.44$) and richness
²⁵⁹ ($H = 0.2$, $df = 2$, $n = 36$, $p = 0.92$) were not (Fig 3a). Weeds under tractor-till were
²⁶⁰ significantly denser compared to no-till ($p_{adj} = 0.06$) and marginally significantly compared
²⁶¹ to roto-till ($p_{adj} = 0.1$), denser by $\sim 6.5\%$ compared to each tillage group, up to $\sim 8 \pm 2$
²⁶² *stems per m⁻²*.

²⁶³ All measured weed variables were affected significantly by cover crop mix, including weed
²⁶⁴ density ($H = 20.1$, $df = 3$, $n = 72$, $p = 0.00016$) changing overall by $\sim 6.5\%$, weed cover
²⁶⁵ ($H = 31$, $df = 3$, $n = 72$, $p = <0.0001$) lowering overall by $\sim 0.5\%$, and weed richness ($H = 10$,
²⁶⁶ $df = 3$, $n = 36$, $p = 0.019$) also lowering overall by $\sim 5.5\%$ (Fig 3b). The weed
²⁶⁷ suppression mix had the most detectable effects on both weed density and cover. The weed
²⁶⁸ suppression mix significantly lowered weed density compared to all other cover crop mix




Figure 2: Infiltration data (a) by tillage, and (b) cover crop mix. Gray dots show plot medians and black point ranges show group mean ± 1 std error and may be small. Significant pairwise post-hoc Wilcoxon test outcomes shown (**** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, *, $p < 0.1$, ' $p > 0.1$)




Figure 3: Weeds data (a) by tillage, and (b) cover crop mix. Gray dots show plot medians and black point ranges show group mean \pm 1 std error and may be small. Significant pairwise post-hoc Wilcoxon test outcomes shown (***) $p < 0.0001$, ** $p < 0.001$, * $p < 0.01$, * $p < 0.05$, * $p < 0.1$, ' $p > 0.1$ or ns)

treatments, namely the null ($p_{adj} = <0.001$), perennial ($p_{adj} = 0.017$), and compaction ($p_{adj} = 0.025$) mixes, by ~ 4 stems m^{-2} ($\sim 50\%$), down to ~ 4 stems per m^{-2} . The weed suppression mix also significantly lowered weed cover compared to all other cover crop mix treatments, namely the null ($p_{adj} = <0.0001$), perennial ($p_{adj} = <0.0001$), and compaction ($p_{adj} = 0.00093$) mixes, by ~ 20 stems m^{-2} ($\sim 33.3\%$), down to $\sim 40 \pm 15\%$. Finally, the null mix showed significantly higher richness compared to the weed suppression mix ($p_{adj} = 0.03$) and marginally significantly compared to perennial ($p_{adj} = 0.1$) and compaction ($p_{adj} = 0.2$) mixes, up to ~ 4 taxa.

3.4 Yield

Radish yield was not significantly affected by tillage ($H = 1.4$, $df = 2$, $n = 8$, $p = 0.5$), and centered at ~ 67.8 g m^{-2} (0.1 lbs m^{-2}) and ~ 13.2 cm (5.2 in) long (Fig 4).




Figure 4: Yield data (a) from no-till, (b) tractor-till, and (c) all tillage groups. Gray dots show plot medians and black point ranges show group mean ± 1 std error and may be small. Photo credits: Naim Edwards.

280 **4 Discussion**

281 Overall this study informs urban soil management by supporting the use of tillage to address
282 compaction issues and improve infiltration, together with cover crops to also reduce weed
283 pressure. We hypothesized that cover crop use would be comparable to tillage effects, which
284 was in part supported, because overall tillage significantly deepened the depth to hardpan by
285 ~0.5 (Fig 1a), which was within the range of effect sizes measured among the various cover
286 crop mixes within the no-till treatment (Fig 1b). Additionally, infiltration was significantly
287 affected by tillage, with roto-till showing the fastest rates (Fig 2a), which agreed with our
288 predictions. Furthermore, weed pressure was significantly affected by both cover crop mixes
289 and tillage (Fig 3); although effects from cover crop mixes, especially the weed suppression
290 mix, were more widespread among multiple measured variables (Fig 3b). Despite these
291 significant effects on soils, infiltration, and weeds, yields did not respond to tillage treatments
292 in this study.

293 Short-term soil compaction issues are commonly alleviated by annual tilling (*Badalíková,*
294 *2010; Salem et al., 2015*), and in addition to validating this practice, this study showed
295 that cover cropping can also be used to manage compaction under no-till, although effects
296 vary by mixture of taxa used. Under tillage, this study validates that tillage intensity
297 corresponds negatively with compaction (measured as depth to hardpan), and additionally
298 clarifies that tractor-till can alleviate compaction in slightly deeper soils below main root
299 zones under $\sim 20.6 \pm 4.6$ cm (8.1 ± 1.8 in), as well as that roto-till can be useful under
300 perennial crops, although under annuals, no-till can be just as effective as roto-till, saving
301 grower time, energy, and cost for areas with crops harvested before rooting surpasses ~ 10
302 cm (4 in) (*Krause and Black, 1995*). For urban Technosol soils, however, it is worth noting
303 that some initial tillage may help remove large metal artifacts and legacy construction debris
304 like rebar, wires, cables, bricks, cinder blocks, and pipes, that could limit root growth under
305 stricter no-till management. Additionally, results suggest that when used together, tillage

may obscure varying but notable effects of cover crops on compaction, however, cover crops would still provide separate benefits to soils, like available macro-nutrients (*Chapagain et al., 2020*). Under no-till, this study found that perennial crop mixes can have significant effects on compaction, but rather than deep roots loosening soils, in some cases depth to hardpan can instead become shallower. This may be due to dense root mats that can form under grasses (*Douglas et al., 1992*) like sorghum-sudangrass, which could further fill already limited pore space in densely-structured clay soils, helping water to pool under the soil surface (*Hoogmoed and Bouma, 1980*). Other studies have generally found similar results (*Ozpinar and Cay, 2006*), suggesting short-term benefits of tillage to soil functions, while acknowledging long-term costs.

Water infiltration is a key function of wide interest for urban environmental management, needed to not only increase available root water, but also reducing erosion and potentially contaminated storm-water runoff and flooding (*Masoner et al., 2019*) after even short heavy rains, due to soil sealing by concrete near hillslopes (*Dreelin et al., 2006*). This study found that roto-till resulted in significantly faster infiltration compared to no-till, unlike tractor-till, suggesting that roto-till management can generally be effective for improving infiltration and drainage. This result could be explained by medium intensity roto-till increasing soil macro-porosity, which compared to micro-pores bind water less tightly, allowing soil water to flow faster (*Gerke, 2006*). In contrast, the tractor diffused tillage energy across deeper soil volume, lowering the density of any added soil macro-pores and thereby making it easier for soil particles to settle back together, and whereas no-till may have needed more time to improve macro-porosity via organic matter effects on soil structure (*King, 2020*). It is also possible that this result could be explained by compost incorporation, where tractor-till similarly incorporated compost more diffusely throughout the soil profile, diluting compost benefits to infiltration. Overall, this study supported the use of roto-till, but not no-till or tractor-till, against a one inch rain event, since both others showed rates of only $\sim 6.2 \pm 0$ mL per sec (0.1 ± 0 gal per min), which would likely be associated with more rain water

333 runoff and soil erosion, worse field drainage, and pooling or flooding into roads. Regarding
334 cover crops, this study suggests that cover crop mixes can generally affect infiltration, though
335 specifically perennials may not have notable significant effects on infiltration rates, despite
336 detectable effects on compaction. Based on these findings, roto-till (alongside compost) can
337 be an effective practice to specifically improve urban soil water infiltration, at least in the
338 short-term, after which no-till may prevail (*Cusser et al., 2020*).

339 In urban settings, weed suppression not just alleviates competition with crops that may
340 already be stressed, but also lowers human health risks, including asthma and other respira-
341 tory issues stemming from allergens like pollen (*Katz and Carey, 2014*), and this study shows
342 evidence that cover crops may be better at this than tilling (*Barberi and Lo Cascio, 2001;*
343 *Cordeau et al., 2020*). Tractor-till alleviated the deepest soil compaction but at the cost of
344 showing the highest density of the two most common weeds, velvet leaf (*Abutilon theophrasti*)
345 and pigweed (*Palmer amaranth*). This may have been due to their fast-growing weed life
346 histories, which can grow denser roots in looser soil with varying microbes (*Korneykova et*
347 *al., 2021*), possibly helping explain slower infiltration, with roots that could re-sprout more,
348 clonally and/or from seed banks (*Hesse et al., 2007*). Most notably for weed suppression, the
349 targeted mix consisting of sorghum-sudangrass, buckwheat, and cowpea indeed significantly
350 reduced both weed density and richness by about half compared to the other cover crop
351 mixes. This result agrees with other studies pairing buckwheat and sorghum-sudangrass
352 (*Smith et al., 2015*), and may have occurred due to any of several reasons—competitive ex-
353 clusion of other weeds by either taxon, such via allelopathic chemical root defenses (*Weston*
354 *et al., 1989*), competition for light (*Liu et al., 2009*), better phosphorus mining and use by
355 buckwheat (*Zhu et al., 2002*), facilitation or amplification of these listed effects by cowpea’s
356 added nitrogen supply (*Martins et al., 2003; Sanginga et al., 2000*), and/or existing adapta-
357 tions to poor dry soils (*Bàrberi et al., 2018*) allowing high biomass accumulation. Given both
358 effectiveness and relative ease of re-seeding and winter-kill, this weed suppression mix could
359 serve well to frame crop beds, keeping out encroaching weeds, or to reduce weed pressure in

³⁶⁰ an area that might be planted in the fall or following season.

³⁶¹ Despite overall significant effects by tillage on compaction, infiltration, and weeds, tillage
³⁶² did not significantly affect radish yield, which in fact agrees with other similar studies, in
³⁶³ contrast to common hypotheses. As is, this study does not rule out more complex rela-
³⁶⁴ tionships between soil compaction, infiltration, and crop yield, as suggested by emerging
³⁶⁵ ideas (*Ryan et al., 2007; Vandermeer and Perfecto, 2017*). With further replication, fu-
³⁶⁶ ture similar studies no-till might be expected to show slightly higher yields (*Nunes et al.,*
³⁶⁷ *2018*), due to resulting longer-term reservoirs of water and nutrients, like from mulched com-
³⁶⁸ post, less reliance on transient influxes from infiltration (*Schlegel et al., 2015; Schlegel and*
³⁶⁹ *Havlin, 1995*), and better soil structure (*Du et al., 2015; Sheehy et al., 2015*). However,
³⁷⁰ despite these reasonable hypotheses, recent studies appear to converge with results shown
³⁷¹ here, namely that benefits to soil from no-till may not scale up to detectably affect yields
³⁷² (*Martínez et al., 2016; Pittelkow et al., 2015; VandenBygaart, 2016*). While forage radish
³⁷³ may not itself respond to management, it may still confer benefits to surrounding soils, even-
³⁷⁴ tually reducing compaction and building soil structure over time, such as with minimal or
³⁷⁵ no mechanical tillage (*Chen and Weil, 2010; Lawley et al., 2011*). Together with others, this
³⁷⁶ study suggests a need for future studies to tie yield to land management strategies, includ-
³⁷⁷ ing in urban clay soils, to aid small-scale growers in addressing legacy compaction and pH
³⁷⁸ issues, potentially acknowledging short-term benefits of occasional tillage (*Blanco-Canqui*
³⁷⁹ *and Wortmann, 2020; Ekboir, 2001*).

³⁸⁰ Taken together, this study presents findings that, in addition to validating previous stud-
³⁸¹ ies supporting general tillage for short-term soil fertility, also supports the targeted use of
³⁸² medium-intensity roto-till and cover crop mixtures (*Chapagain et al., 2020*) specifically for
³⁸³ weed suppression. This study serves as a model demonstration of both widely-accessible and
³⁸⁴ effective strategies for growing on re-purposed urban soils after industrial land-use turnover.
³⁸⁵ Overall, we advocate for the maximal use of cover crop mixes for various target functions,
³⁸⁶ with medium-intensity tillage to jump-start urban cultivation.

³⁸⁷ **Funding**

³⁸⁸ *This research did not receive any specific grant from funding agencies in the public, commer-*
³⁸⁹ *cial, or not-for-profit sectors.*

³⁹⁰ **Declaration of interests**

³⁹¹ Authors declare no conflicts of interest.

³⁹² **Acknowledgements**

³⁹³ Thanks to previous anonymous reviewers and peers KS, ZHF, and JK for discussion of initial
³⁹⁴ drafts.

³⁹⁵ **Author contributions**

³⁹⁶ NE conceived, designed, and performed the study; NE and NM helped collect data; NM
³⁹⁷ analyzed data; NE wrote the initial report. All authors wrote and revised second draft; NM
³⁹⁸ wrote the third draft; all authors revised later drafts.

³⁹⁹ **Data statement**

⁴⁰⁰ Data and code stored at github.com/nmedina17/must.

401 **References**

- 402 Acuto, M., Parnell, S., Seto, K.C., 2018. Building a global urban science. *Nature Sustainability* 1, 2–4. <https://doi.org/10.1038/s41893-017-0013-9>
- 403
- 404 Allaire, J., Xie, Y., McPherson, J., Luraschi, J., Ushey, K., Atkins, A., Wickham, H., Cheng, J., Chang, W., Iannone, R., 2022. Rmarkdown: Dynamic documents for r.
- 405
- 406 Amundson, R., Berhe, A.A., Hopmans, J.W., Olson, C., Sztein, A.E., Sparks, D.L., 2015. Soil science. *Soil and human security in the 21st century*. *Science* (New York, N.Y.) 348, 1261071. <https://doi.org/10.1126/science.1261071>
- 407
- 408
- 409 Anderson, R.L., 2007. Managing weeds with a dualistic approach of prevention and control. A review. *Agronomy for Sustainable Development* 27, 13–18. <https://doi.org/10.1051/agro:2006027>
- 410
- 411
- 412 Badalíková, B., 2010. Influence of Soil Tillage on Soil Compaction, in: Dedousis, A.P., Bartzanas, T. (Eds.), *Soil Engineering*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 19–30. https://doi.org/10.1007/978-3-642-03681-1_2
- 413
- 414
- 415 Baraibar, B., Murrell, E.G., Bradley, B.A., Barbercheck, M.E., Mortensen, D.A., Kaye, J.P., White, C.M., 2020. Cover crop mixture expression is influenced by nitrogen availability and growing degree days. *PLOS ONE* 15, e0235868. <https://doi.org/10.1371/journal.pone.0235868>
- 416
- 417
- 418
- 419 Barberi, P., Lo Cascio, B., 2001. Long-term tillage and crop rotation effects on weed seedbank size and composition. *Weed Research* 41, 325–340. <https://doi.org/10.1046/j.1365-3180.2001.00241.x>
- 420
- 421
- 422 Bärberi, P., Bocci, G., Carlesi, S., Armengot, L., Blanco-Moreno, J.M., Sans, F.X., 2018. Linking species traits to agroecosystem services: A functional analysis of weed communities. *Weed Research* 58, 76–88. <https://doi.org/10.1111/wre.12283>
- 423
- 424
- 425 Bazzoffi, P., 1998. The effect of urban refuse compost and different tractors tyres on soil physical properties, soil erosion and maize yield. *Soil and Tillage Research* 48, 275–286. [https://doi.org/10.1016/S0167-1987\(98\)00133-0](https://doi.org/10.1016/S0167-1987(98)00133-0)
- 426
- 427

- 428 Bedoussac, L., Journet, E.-P., Hauggaard-Nielsen, H., Naudin, C., Corre-Hellou, G., Jensen,
429 E.S., Prieur, L., Justes, E., 2015. Ecological principles underlying the increase of produc-
430 tivity achieved by cereal-grain legume intercrops in organic farming. A review. *Agronomy*
431 for Sustainable Development
- 35, 911–935. <https://doi.org/10.1007/s13593-014-0277-7>
- 432 Beniston, J.W., Lal, R., Mercer, K.L., 2016. Assessing and Managing Soil Quality for
433 Urban Agriculture in a Degraded Vacant Lot Soil: ASSESSING AND MANAGING
434 SOIL QUALITY FOR URBAN AGRICULTURE. *Land Degradation & Development*
435 27, 996–1006. <https://doi.org/10.1002/ldr.2342>
- 436 Birk, M.A., 2019. Measurements: Tools for units of measurement.
- 437 Blanco-Canqui, H., Wortmann, C.S., 2020. Does occasional tillage undo the ecosystem
438 services gained with no-till? A review. *Soil and Tillage Research* 198, 104534. <https://doi.org/10.1016/j.still.2019.104534>
- 440 Blesh, J., 2017. Functional traits in cover crop mixtures: Biological nitrogen fixation and
441 multifunctionality. *Journal of Applied Ecology* 55, 38–48. <https://doi.org/10.1111/1365-2664.13011>
- 443 Block, D.R., Chávez, N., Allen, E., Ramirez, D., 2012. Food sovereignty, urban food ac-
444 cess, and food activism: Contemplating the connections through examples from Chicago.
445 *Agriculture and Human Values* 29, 203–215. <https://doi.org/10.1007/s10460-011-9336-8>
- 446 Bourke, P.M., Evers, J.B., Bijma, P., van Apeldoorn, D.F., Smulders, M.J.M., Kuyper,
447 T.W., Mommer, L., Bonnema, G., 2021. Breeding Beyond Monoculture: Putting the
448 “Intercrop” Into Crops. *Frontiers in Plant Science* 12, 734167. <https://doi.org/10.3389/fpls.2021.734167>
- 450 Catania, P., Badalucco, L., Laudicina, V.A., Vallone, M., 2018. Effects of tilling methods
451 on soil penetration resistance, organic carbon and water stable aggregates in a vineyard
452 of semiarid Mediterranean environment. *Environmental Earth Sciences* 77, 348. <https://doi.org/10.1007/s12665-018-7520-5>
- 454 Chamberlain, S., Szoechs, E., Foster, Z., Arendsee, Z., Boettiger, C., Ram, K., Bartomeus,

- 455 I., Baumgartner, J., O'Donnell, J., Oksanen, J., Tzovaras, B.G., Marchand, P., Tran, V.,
456 Salmon, M., Li, G., Grenié, M., 2020. Taxize: Taxonomic information from around the
457 web.
- 458 Chapagain, T., Lee, E.A., Raizada, M.N., 2020. The Potential of Multi-Species Mixtures
459 to Diversify Cover Crop Benefits. *Sustainability* 12, 2058. <https://doi.org/10.3390/su12052058>
- 460 Chen, G., Weil, R.R., 2010. Penetration of cover crop roots through compacted soils. *Plant
461 and Soil* 331, 31–43. <https://doi.org/10.1007/s11104-009-0223-7>
- 462 Clark, A. (Ed.), 2007. Managing cover crops profitably, 3rd ed. ed, Handbook series.
463 Sustainable Agriculture Research & Education (SARE), College Park, MD.
- 464 Clendenning, J., Dressler, W.H., Richards, C., 2016. Food justice or food sovereignty?
465 Understanding the rise of urban food movements in the USA. *Agriculture and Human
466 Values* 33, 165–177. <https://doi.org/10.1007/s10460-015-9625-8>
- 467 Cogger, C.G., 2005. Potential Compost Benefits for Restoration Of Soils Disturbed by
468 Urban Development. *Compost Science & Utilization* 13, 243–251. <https://doi.org/10.1080/1065657X.2005.10702248>
- 469 Cordeau, S., Baudron, A., Adeux, G., 2020. Is Tillage a Suitable Option for Weed Man-
470 agement in Conservation Agriculture? *Agronomy* 10, 1746. <https://doi.org/10.3390/agronomy10111746>
- 471 Correa, J., Postma, J.A., Watt, M., Wojciechowski, T., 2019. Soil compaction and the
472 architectural plasticity of root systems. *Journal of Experimental Botany* 70, 6019–6034.
473 <https://doi.org/10.1093/jxb/erz383>
- 474 Cusser, S., Bahlai, C., Swinton, S.M., Robertson, G.P., Haddad, N.M., 2020. Long-term re-
475 search avoids spurious and misleading trends in sustainability attributes of no-till. *Global
476 Change Biology* 26, 3715–3725. <https://doi.org/10.1111/gcb.15080>
- 477 Daftary-Steel, S., Herrera, H., Porter, C., 2015. The Unattainable Trifecta of Urban Agri-
478 culture. *Journal of Agriculture, Food Systems, and Community Development* 19–32.
- 479

- 482 https://doi.org/10.5304/jafscd.2015.061.014
- 483 Daniel, P., 2007. African American Farmers and Civil Rights 37.
- 484 Davies, G.M., Gray, A., 2015. Don't let spurious accusations of pseudoreplication limit our
485 ability to learn from natural experiments (and other messy kinds of ecological monitor-
486 ing). *Ecology and Evolution* 5, 5295–5304. https://doi.org/10.1002/ece3.1782
- 487 de Cárcer, P.S., Sinaj, S., Santonja, M., Fossati, D., Jeangros, B., 2019. Long-term effects
488 of crop succession, soil tillage and climate on wheat yield and soil properties. *Soil and*
489 *Tillage Research* 190, 209–219. https://doi.org/10.1016/j.still.2019.01.012
- 490 Douglas, J.T., Koppi, A.J., Moran, C.J., 1992. Alteration of the structural attributes of a
491 compact clay loam soil by growth of a perennial grass crop. *Plant and Soil* 139, 195–202.
492 https://doi.org/10.1007/BF00009310
- 493 Dreelin, E.A., Fowler, L., Ronald Carroll, C., 2006. A test of porous pavement effectiveness
494 on clay soils during natural storm events. *Water Research* 40, 799–805. https://doi.org/
495 10.1016/j.watres.2005.12.002
- 496 Drugova, T., Curtis, K.R., Ward, R.A., 2022. Producer preferences for drought management
497 strategies in the arid west. *Renewable Agriculture and Food Systems* 37, 14–23. https:
498 //doi.org/10.1017/S1742170521000259
- 499 Du, Z., Ren, T., Hu, C., Zhang, Q., 2015. Transition from intensive tillage to no-till enhances
500 carbon sequestration in microaggregates of surface soil in the North China Plain. *Soil*
501 and *Tillage Research* 146, 26–31. https://doi.org/10.1016/j.still.2014.08.012
- 502 Ekboir, J., 2001. Developing No-Till Packages for Small-Scale Farmers.
- 503 FAO, 2014. World reference base for soil resources 2014: International soil classification
504 system for naming soils and creating legends for soil maps. FAO, Rome.
- 505 Garbach, K., Milder, J.C., DeClerck, F.A.J., Montenegro de Wit, M., Driscoll, L., Gemmill-
506 Herren, B., 2017. Examining multi-functionality for crop yield and ecosystem services
507 in five systems of agroecological intensification. *International Journal of Agricultural*
508 *Sustainability* 15, 11–28. https://doi.org/10.1080/14735903.2016.1174810

- 509 García-González, I., Hontoria, C., Gabriel, J.L., Alonso-Ayuso, M., Quemada, M., 2018.
510 Cover crops to mitigate soil degradation and enhance soil functionality in irrigated land.
511 Geoderma 322, 81–88. <https://doi.org/10.1016/j.geoderma.2018.02.024>
- 512 García-Sempere, A., Morales, H., Hidalgo, M., Ferguson, B.G., Rosset, P., Nazar-
513 Beutelspacher, A., 2019. Food Sovereignty in the city?: A methodological proposal
514 for evaluating food sovereignty in urban settings. Agroecology and Sustainable Food
515 Systems 43, 1145–1173. <https://doi.org/10.1080/21683565.2019.1578719>
- 516 Gerke, H.H., 2006. Preferential flow descriptions for structured soils. Journal of Plant
517 Nutrition and Soil Science 169, 382–400. <https://doi.org/10.1002/jpln.200521955>
- 518 Grossman, J.M., 2003. Exploring farmer knowledge of soil processes in organic coffee systems
519 of Chiapas, Mexico. Geoderma 111, 267–287. [https://doi.org/10.1016/S0016-7061\(02\)00268-9](https://doi.org/10.1016/S0016-7061(02)00268-9)
- 520
- 521 Grossman, J.M., Sheaffer, C., Wyse, D., Graham, P.H., 2005. Characterization of slow-
522 growing root nodule bacteria from *Inga oerstediana* in organic coffee agroecosystems in
523 Chiapas, Mexico. Applied Soil Ecology 29, 236–251. <https://doi.org/10.1016/j.apsoil.2004.12.008>
- 524
- 525 Hesse, E., Rees, M., Müller-Schärer, H., 2007. Seed bank persistence of clonal weeds in
526 contrasting habitats: Implications for control. Plant Ecology 190, 233–243. <https://doi.org/10.1007/s11258-006-9203-7>
- 527
- 528 Hill, R.L., Horton, R., Cruse, R.M., 1985. Tillage Effects on Soil Water Retention and
529 Pore Size Distribution of Two Mollisols. Soil Science Society of America Journal 49,
530 1264–1270. <https://doi.org/10.2136/sssaj1985.03615995004900050039x>
- 531
- 532 Hoogmoed, W.B., Bouma, J., 1980. A Simulation Model for Predicting Infiltration into
Cracked Clay Soil. Soil Science Society of America Journal 44, 458–461. <https://doi.org/10.2136/sssaj1980.03615995004400030003x>
- 533
- 534 Kassambara, A., 2021. Rstatix: Pipe-friendly framework for basic statistical tests.
- 535 Katz, D.S.W., Carey, T.S., 2014. Heterogeneity in ragweed pollen exposure is determined

- 536 by plant composition at small spatial scales. *Science of the Total Environment* 485–486,
537 435–440. <https://doi.org/10.1016/j.scitotenv.2014.03.099>
- 538 King, A.E., 2020. Soil Organic Matter as Catalyst of Crop Resource Capture. *Frontiers in*
539 *Environmental Science* 8, 8.
- 540 Korneykova, M.V., Vasenev, V.I., Nikitin, D.A., Soshina, A.S., Dolgikh, A.V., Sotnikova,
541 Y.L., 2021. Urbanization Affects Soil Microbiome Profile Distribution in the Russian
542 Arctic Region. *International Journal of Environmental Research and Public Health* 18,
543 11665. <https://doi.org/10.3390/ijerph182111665>
- 544 Krause, M.A., Black, J.R., 1995. Optimal Adoption Strategies for No-till Technology in
545 Michigan. *Review of Agricultural Economics* 17, 299. <https://doi.org/10.2307/1349575>
- 546 Kumar, K., Hundal, L.S., 2016. Soil in the City: Sustainably Improving Urban Soils. *Journal*
547 *of Environmental Quality* 45, 2–8. <https://doi.org/10.2134/jeq2015.11.0589>
- 548 Kuzyakov, Y., Zamanian, K., 2019. Reviews and syntheses: Agropedogenesis – humankind
549 as the sixth soil-forming factor and attractors of agricultural soil degradation. *Biogeosciences*
550 16, 4783–4803. <https://doi.org/10.5194/bg-16-4783-2019>
- 551 Lal, R., 2007. Soil Science and the Carbon Civilization 71, 1425–1437. <https://doi.org/10.2136/sssaj2007.0001>
- 553 Lal, R., Negassa, W., Lorenz, K., 2015. Carbon sequestration in soil. *Current Opinion in*
554 *Environmental Sustainability* 15, 79–86. <https://doi.org/10.1016/j.cosust.2015.09.002>
- 555 Lawley, Y.E., Weil, R.R., Teasdale, J.R., 2011. Forage Radish Cover Crop Suppresses Winter
556 Annual Weeds in Fall and Before Corn Planting. *Agronomy Journal* 103, 137–144. <https://doi.org/10.2134/agronj2010.0187>
- 558 Liu, J.G., Mahoney, K.J., Sikkema, P.H., Swanton, C.J., 2009. The importance of light
559 quality in crop-weed competition: Light quality and crop competition. *Weed Research*
560 49, 217–224. <https://doi.org/10.1111/j.1365-3180.2008.00687.x>
- 561 London, J.K., Cutts, B.B., Schwarz, K., Schmidt, L., Cadenasso, M.L., 2021. Unearthing
562 the entangled roots of urban agriculture. *Agriculture and Human Values* 38, 205–220.

- 563 <https://doi.org/10.1007/s10460-020-10158-x>
- 564 Martínez, I., Chervet, A., Weisskopf, P., Sturny, W.G., Etana, A., Stettler, M., Forkman,
565 J., Keller, T., 2016. Two decades of no-till in the Oberacker long-term field experiment:
566 Part I. Crop yield, soil organic carbon and nutrient distribution in the soil profile. *Soil
567 and Tillage Research* 163, 141–151. <https://doi.org/10.1016/j.still.2016.05.021>
- 568 Martins, L.M.V., Xavier, G.R., Rangel, F.W., Ribeiro, J.R.A., Neves, M.C.P., Morgado,
569 L.B., Rumjanek, N.G., 2003. Contribution of biological nitrogen fixation to cowpea: A
570 strategy for improving grain yield in the semi-arid region of Brazil. *Biology and Fertility
571 of Soils* 38, 333–339. <https://doi.org/10.1007/s00374-003-0668-4>
- 572 Masoner, J.R., Kolpin, D.W., Cozzarelli, I.M., Barber, L.B., Burden, D.S., Foreman, W.T.,
573 Forshay, K.J., Furlong, E.T., Groves, J.F., Hladik, M.L., Hopton, M.E., Jaeschke, J.B.,
574 Keefe, S.H., Krabbenhoft, D.P., Lowrance, R., Romanok, K.M., Rus, D.L., Selbig, W.R.,
575 Williams, B.H., Bradley, P.M., 2019. Urban Stormwater: An Overlooked Pathway of Ex-
576 tensive Mixed Contaminants to Surface and Groundwaters in the United States. *Environ-
577 mental Science & Technology* 53, 10070–10081. <https://doi.org/10.1021/acs.est.9b02867>
- 578 Materechera, S.A., 2009. Tillage and tractor traffic effects on soil compaction in horticul-
579 tural fields used for peri-urban agriculture in a semi-arid environment of the North West
580 Province, South Africa. *Soil and Tillage Research* 103, 11–15. <https://doi.org/10.1016/j.still.2008.09.001>
- 582 McDougall, R., Kristiansen, P., Rader, R., 2019. Small-scale urban agriculture results in high
583 yields but requires judicious management of inputs to achieve sustainability. *Proceedings
584 of the National Academy of Sciences of the United States of America* 116, 129–134.
585 <https://doi.org/10.1073/pnas.1809707115>
- 586 Mead, R., Riley, J., 1981. A Review of Statistical Ideas Relevant to Intercropping Research.
587 Journal of the Royal Statistical Society. Series A (General) 144, 462. <https://doi.org/10.2307/2981827>
- 589 Montgomery, D.R., 2007. Soil erosion and agricultural sustainability 104, 13268–13272.

- 590 Monti, A., Venturi, P., Elbersen, H.W., 2001. Evaluation of the establishment of lowland and
591 upland switchgrass (*Panicum virgatum* L.) Varieties under different tillage and seedbed
592 conditions in northern Italy. *Soil and Tillage Research* 63, 75–83. [https://doi.org/10.1016/S0167-1987\(01\)00238-0](https://doi.org/10.1016/S0167-1987(01)00238-0)
- 593
- 594 Mori, T., Wang, S., Zhang, W., Mo, J., 2022. Microbial assembly adapted to low-P soils in
595 three subtropical forests by increasing the maximum rate of substrate conversion of acid
596 phosphatases but not by decreasing the half-saturation constant. *European Journal of
597 Soil Biology* 108, 103377. <https://doi.org/10.1016/j.ejsobi.2021.103377>
- 598 Müller, K., 2020. Here: A simpler way to find your files.
- 599 Nunes, M.R., van Es, H.M., Schindelbeck, R., Ristow, A.J., Ryan, M., 2018. No-till and
600 cropping system diversification improve soil health and crop yield. *Geoderma* 328, 30–43.
601 <https://doi.org/10.1016/j.geoderma.2018.04.031>
- 602 O'Riordan, R., Davies, J., Stevens, C., Quinton, J.N., Boyko, C., 2021. The ecosystem
603 services of urban soils: A review. *Geoderma* 395, 115076. <https://doi.org/10.1016/j.geoderma.2021.115076>
- 604
- 605 Oldfield, E.E., Bradford, M.A., Wood, S.A., 2019. Global meta-analysis of the relationship
606 between soil organic matter and crop yields 15–32.
- 607 Ozpinar, S., Cay, A., 2006. Effect of different tillage systems on the quality and crop
608 productivity of a clay-loam soil in semi-arid north-western Turkey. *Soil and Tillage
609 Research* 88, 95–106. <https://doi.org/10.1016/j.still.2005.04.009>
- 610 Pavao-Zuckerman, M.A., 2008. The Nature of Urban Soils and Their Role in Ecological
611 Restoration in Cities. *Restoration Ecology* 16, 642–649. [https://doi.org/10.1111/j.1526-100X.2008.00486.x](https://doi.org/10.1111/j.1526-
612 100X.2008.00486.x)
- 613 Perez, R., 2021. Freedom Farmers: Agricultural Resistance and the Black Freedom Move-
614 ment, by Monica M.White, Chapel Hill: University of North Carolina Press, 2018.
615 208 pp. \$14.99 (e-book). ISBN: 978-1-4696-4370-0. *Rural Sociology* 86, 974–977.
616 <https://doi.org/10.1111/ruso.12423>

- 617 Pittelkow, C.M., Linquist, B.A., Lundy, M.E., Liang, X., van Groenigen, K.J., Lee, J., van
618 Gestel, N., Six, J., Venterea, R.T., van Kessel, C., 2015. When does no-till yield more?
619 A global meta-analysis. *Field Crops Research* 183, 156–168. <https://doi.org/10.1016/j.fcr.2015.07.020>
- 621 Possinger, A.R., Byrne, L.B., Breen, N.E., 2013. Effect of buckwheat (*Fagopyrum Esculentum*) on soil-phosphorus availability and organic acids. *Journal of Plant Nutrition and*
622 *Soil Science* 176, 16–18. <https://doi.org/10.1002/jpln.201200337>
- 624 R Core Team, 2022. R: A language and environment for statistical computing. R Foundation
625 for Statistical Computing, Vienna, Austria.
- 626 Richter, D.D., 2021. Searching for solutions to our soil woes—A World Without Soil: The
627 Past, Present, and Precarious Future of the Earth Beneath Our Feet, Jo Handelsman,
628 Yale University Press, 2021, 272 pp. *Science* 374, 1452–1452. <https://doi.org/10.1126/science.abm4765>
- 630 Roger-Estrade, J., Anger, C., Bertrand, M., Richard, G., 2010. Tillage and soil ecology:
631 Partners for sustainable agriculture. *Soil and Tillage Research* 111, 33–40. <https://doi.org/10.1016/j.still.2010.08.010>
- 633 Ryan, J.G., Ludwig, J.A., Mcalpine, C.A., 2007. Complex adaptive landscapes (CAL): A
634 conceptual framework of multi-functional, non-linear ecohydrological feedback systems.
635 *Ecological Complexity* 4, 113–127. <https://doi.org/10.1016/j.ecocom.2007.03.004>
- 636 Salem, H.M., Valero, C., Muñoz, M.Á., Rodríguez, M.G., Silva, L.L., 2015. Short-term
637 effects of four tillage practices on soil physical properties, soil water potential, and maize
638 yield. *Geoderma* 237–238, 60–70. <https://doi.org/10.1016/j.geoderma.2014.08.014>
- 639 Sanginga, N., Lyasse, O., Singh, B.B., 2000. Phosphorus use efficiency and nitrogen balance
640 of cowpea breeding lines in a low P soil of the derived savanna zone in West Africa. *Plant*
641 *and Soil* 10.
- 642 Schlegel, A.J., Assefa, Y., Bond, H.D., Wetter, S.M., Stone, L.R., 2015. Soil Physicochemical
643 Properties after 10 Years of Animal Waste Application. *Soil Science Society of America*

- 644 Journal 79, 711–719. <https://doi.org/10.2136/sssaj2014.11.0461>
- 645 Schlegel, A.J., Havlin, J.L., 1995. Corn Response to Long-Term Nitrogen and Phosphorus
646 Fertilization. Journal of Production Agriculture 8, 181–185. <https://doi.org/10.2134/jpa1995.0181>
- 648 Sheehy, J., Regina, K., Alakukku, L., Six, J., 2015. Impact of no-till and reduced tillage
649 on aggregation and aggregate-associated carbon in Northern European agroecosystems.
650 Soil and Tillage Research 150, 107–113. <https://doi.org/10.1016/j.still.2015.01.015>
- 651 Siebert, A., 2020. Transforming urban food systems in South Africa: Unfolding food
652 sovereignty in the city. The Journal of Peasant Studies 47, 401–419. <https://doi.org/10.1080/03066150.2018.1543275>
- 654 Silk, M.J., Harrison, X.A., Hodgson, D.J., 2020. Perils and pitfalls of mixed-effects regression
655 models in biology. PeerJ 8, e9522. <https://doi.org/10.7717/peerj.9522>
- 656 Sircely, J., Naeem, S., 2012. Biodiversity and Ecosystem Multi-Functionality: Observed
657 Relationships in Smallholder Fallows in Western Kenya. PLoS ONE 7, e50152. <https://doi.org/10.1371/journal.pone.0050152>
- 659 Six, J., Feller, C., Denef, K., Ogle, S., De Moraes, J.C., Albrecht, A., 2002. Soil organic
660 matter, biota and aggregation in temperate and tropical soils - Effects of no-tillage.
661 <https://doi.org/10.1051/agro>
- 662 Small, G., Shrestha, P., Metson, G.S., Polsky, K., Jimenez, I., Kay, A., 2019. Excess phosphorus
663 from compost applications in urban gardens creates potential pollution hotspots
664 Excess phosphorus from compost applications in urban gardens creates potential pollution
665 hotspots.
- 666 Smith, A.P., Marín-Spiotta, E., Balser, T., 2015. Successional and seasonal variations in soil
667 and litter microbial community structure and function during tropical postagricultural
668 forest regeneration: A multiyear study. Global Change Biology 21, 3532–3547. <https://doi.org/10.1111/gcb.12947>
- 670 Stewart, C.E., Paustian, K., Conant, R.T., Plante, A.F., Six, J., 2007. Soil carbon satura-

- 671 tion: Concept, evidence and evaluation. *Biogeochemistry* 86, 19–31. <https://doi.org/10.1007/s10533-007-9140-0>
- 672
- 673 Stewart, R., Korth, M., Langer, L., Rafferty, S., Da Silva, N.R., van Rooyen, C., 2013. What
674 are the impacts of urban agriculture programs on food security in low and middle-income
675 countries? *Environmental Evidence* 2, 7. <https://doi.org/10.1186/2047-2382-2-7>
- 676 Storkey, J., Neve, P., 2018. What good is weed diversity? *Weed Research* 58, 239–243.
677 <https://doi.org/10.1111/wre.12310>
- 678 Tomczak, M., Tomczak, E., 2014. The need to report effect size estimates revisited. An
679 overview of some recommended measures of effect size 1, 7.
- 680 Tresch, S., Moretti, M., Bayon, R.C.L., Mäder, P., Zanetta, A., Frey, D., Fliessbach, A.,
681 2018. A gardener's influence on urban soil quality. *Frontiers in Environmental Science*
682 6. <https://doi.org/10.3389/fenvs.2018.00025>
- 683 VandenBygaart, A.J., 2016. The myth that no-till can mitigate global climate change. *Agricri-
684 culture, Ecosystems & Environment* 216, 98–99. <https://doi.org/10.1016/j.agee.2015.09.013>
- 685
- 686 Vandermeer, J., Perfecto, I., 2017. Ecological complexity and agroecosystems : Seven themes
687 from theory. *Agroecology and Sustainable Food Systems* 41, 697–722. <https://doi.org/10.1080/21683565.2017.1322166>
- 688
- 689 Wade, A.M., Richter, D.D., Craft, C.B., Bao, N.Y., Heine, P.R., Osteen, M.C., Tan,
690 K.G., 2021. Urban-Soil Pedogenesis Drives Contrasting Legacies of Lead from Paint
691 and Gasoline in City Soil. *Environmental Science & Technology* 55, 7981–7989. <https://doi.org/10.1021/acs.est.1c00546>
- 692
- 693 Wang, X.-B., Cai, D.-X., Hoogmoed, W.B., Oenema, O., Perdok, U.D., 2006. Potential
694 Effect of Conservation Tillage on Sustainable Land Use: A Review of Global Long-Term
695 Studies. *Pedosphere* 16, 587–595. [https://doi.org/10.1016/S1002-0160\(06\)60092-1](https://doi.org/10.1016/S1002-0160(06)60092-1)
- 696
- 697 Weston, L.A., Harmon, R., Mueller, S., 1989. Allelopathic potential of sorghum-sudangrass
hybrid (sudex). *Journal of Chemical Ecology* 15, 1855–1865. <https://doi.org/10.1007/>

698 BF01012272

- 699 Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemund,
700 G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Bache, S.M.,
701 Müller, K., Ooms, J., Robinson, D., Seidel, D.P., Spinu, V., Takahashi, K., Vaughan, D.,
702 Wilke, C., Woo, K., Yutani, H., 2019. Welcome to the tidyverse. Journal of Open Source
703 Software 4, 1686. <https://doi.org/10.21105/joss.01686>
- 704 Wolkowski, R.P., 1990. Relationship between Wheel-Traffic-Induced Soil Compaction, Nu-
705 trient Availability, and Crop Growth: A Review. Journal of Production Agriculture 3,
706 460–469. <https://doi.org/10.2134/jpa1990.0460>
- 707 Xie, Y., 2022a. Bookdown: Authoring books and technical documents with r markdown.
- 708 Xie, Y., 2022b. Knitr: A general-purpose package for dynamic report generation in r.
- 709 Zhu, Y.G., He, Y.Q., Smith, S.E., Smith, F.A., 2002. Buckwheat (*Fagopyrum esculentum*
710 Moench) has high capacity to take up phosphorus (P) from a calcium (Ca)-bound source.
711 Plant and Soil 239, 1–8. <https://doi.org/10.1023/A:1014958029905>