

1. Let  $u = \sin x$ , so that  $du = \cos x \, dx$ . Then  $\int \cos x(1 + \sin^2 x) \, dx = \int (1 + u^2) \, du = u + \frac{1}{3}u^3 + C = \sin x + \frac{1}{3}\sin^3 x + C$ .

2. 
$$\int \frac{\sin^3 x}{\cos x} \, dx = \int \frac{\sin^2 x \sin x}{\cos x} \, dx = \int \frac{(1 - \cos^2 x) \sin x}{\cos x} \, dx = \int \frac{1 - u^2}{u} (-du) \quad \left[ \begin{array}{l} u = \cos x \\ du = -\sin x \, dx \end{array} \right]$$
$$= \int \left(u - \frac{1}{u}\right) du = \frac{1}{2}u^2 - \ln|u| + C = \frac{1}{2}\cos^2 x - \ln|\cos x| + C$$

3. 
$$\int \frac{\sin x + \sec x}{\tan x} \, dx = \int \left( \frac{\sin x}{\tan x} + \frac{\sec x}{\tan x} \right) dx = \int (\cos x + \csc x) \, dx = \sin x + \ln|\csc x - \cot x| + C$$

4. 
$$\int \tan^3 \theta \, d\theta = \int (\sec^2 \theta - 1) \tan \theta \, d\theta = \int \tan \theta \sec^2 \theta \, d\theta - \int \frac{\sin \theta}{\cos \theta} \, d\theta$$
$$= \int u \, du + \int \frac{dv}{v} \quad \left[ \begin{array}{ll} u = \tan \theta, & v = \cos \theta, \\ du = \sec^2 \theta \, d\theta & dv = -\sin \theta \, d\theta \end{array} \right]$$
$$= \frac{1}{2}u^2 + \ln|v| + C = \frac{1}{2}\tan^2 \theta + \ln|\cos \theta| + C$$

5. 
$$\int_0^2 \frac{2t}{(t-3)^2} \, dt = \int_{-3}^{-1} \frac{2(u+3)}{u^2} \, du \quad \left[ \begin{array}{l} u = t-3, \\ du = dt \end{array} \right] = \int_{-3}^{-1} \left( \frac{2}{u} + \frac{6}{u^2} \right) du = \left[ 2 \ln|u| - \frac{6}{u} \right]_{-3}^{-1}$$
$$= (2 \ln 1 + 6) - (2 \ln 3 + 2) = 4 - 2 \ln 3 \text{ or } 4 - \ln 9$$

6. Let  $u = x^2$ . Then  $du = 2x \, dx \Rightarrow \int \frac{x \, dx}{\sqrt{3-x^4}} = \frac{1}{2} \int \frac{du}{\sqrt{3-u^2}} = \frac{1}{2} \sin^{-1} \frac{u}{\sqrt{3}} + C = \frac{1}{2} \sin^{-1} \frac{x^2}{\sqrt{3}} + C$ .

7. Let  $u = \arctan y$ . Then  $du = \frac{dy}{1+y^2} \Rightarrow \int_{-1}^1 \frac{e^{\arctan y}}{1+y^2} \, dy = \int_{-\pi/4}^{\pi/4} e^u \, du = [e^u]_{-\pi/4}^{\pi/4} = e^{\pi/4} - e^{-\pi/4}$ .

8. 
$$\int x \csc x \cot x \, dx \quad \left[ \begin{array}{ll} u = x, & dv = \csc x \cot x \, dx, \\ du = dx & v = -\csc x \end{array} \right] = -x \csc x - \int (-\csc x) \, dx = -x \csc x + \ln|\csc x - \cot x| + C$$

9. 
$$\int_1^3 r^4 \ln r \, dr \quad \left[ \begin{array}{ll} u = \ln r, & dv = r^4 \, dr, \\ du = \frac{dr}{r} & v = \frac{1}{5}r^5 \end{array} \right] = \left[ \frac{1}{5}r^5 \ln r \right]_1^3 - \int_1^3 \frac{1}{5}r^4 \, dr = \frac{243}{5} \ln 3 - 0 - \left[ \frac{1}{25}r^5 \right]_1^3$$
$$= \frac{243}{5} \ln 3 - \left( \frac{243}{25} - \frac{1}{25} \right) = \frac{243}{5} \ln 3 - \frac{242}{25}$$

10. 
$$\frac{x-1}{x^2-4x-5} = \frac{x-1}{(x-5)(x+1)} = \frac{A}{x-5} + \frac{B}{x+1} \Rightarrow x-1 = A(x+1) + B(x-5). \text{ Setting } x = -1 \text{ gives}$$
$$-2 = -6B, \text{ so } B = \frac{1}{3}. \text{ Setting } x = 5 \text{ gives } 4 = 6A, \text{ so } A = \frac{2}{3}. \text{ Now}$$

$$\int_0^4 \frac{x-1}{x^2-4x-5} \, dx = \int_0^4 \left( \frac{2/3}{x-5} + \frac{1/3}{x+1} \right) dx = \left[ \frac{2}{3} \ln|x-5| + \frac{1}{3} \ln|x+1| \right]_0^4$$
$$= \frac{2}{3} \ln 1 + \frac{1}{3} \ln 5 - \frac{2}{3} \ln 5 - \frac{1}{3} \ln 1 = -\frac{1}{3} \ln 5$$

$$11. \int \frac{x-1}{x^2-4x+5} dx = \int \frac{(x-2)+1}{(x-2)^2+1} dx = \int \left( \frac{u}{u^2+1} + \frac{1}{u^2+1} \right) du \quad [u = x-2, du = dx]$$

$$= \frac{1}{2} \ln(u^2+1) + \tan^{-1} u + C = \frac{1}{2} \ln(x^2-4x+5) + \tan^{-1}(x-2) + C$$

$$12. \int \frac{x}{x^4+x^2+1} dx = \int \frac{\frac{1}{2} du}{u^2+u+1} \quad \left[ \begin{array}{l} u = x^2, \\ du = 2x dx \end{array} \right] = \frac{1}{2} \int \frac{du}{(u+\frac{1}{2})^2+\frac{3}{4}}$$

$$= \frac{1}{2} \int \frac{\frac{\sqrt{3}}{2} dv}{\frac{3}{4}(v^2+1)} \quad \left[ \begin{array}{l} u+\frac{1}{2} = \frac{\sqrt{3}}{2}v, \\ du = \frac{\sqrt{3}}{2} dv \end{array} \right] = \frac{\sqrt{3}}{4} \cdot \frac{4}{3} \int \frac{dv}{v^2+1}$$

$$= \frac{1}{\sqrt{3}} \tan^{-1} v + C = \frac{1}{\sqrt{3}} \tan^{-1} \left( \frac{2}{\sqrt{3}} \left( x^2 + \frac{1}{2} \right) \right) + C$$

$$13. \int \sin^3 \theta \cos^5 \theta d\theta = \int \cos^5 \theta \sin^2 \theta \sin \theta d\theta = - \int \cos^5 \theta (1 - \cos^2 \theta)(-\sin \theta) d\theta$$

$$= - \int u^5(1-u^2) du \quad \left[ \begin{array}{l} u = \cos \theta, \\ du = -\sin \theta d\theta \end{array} \right]$$

$$= \int (u^7 - u^5) du = \frac{1}{8}u^8 - \frac{1}{6}u^6 + C = \frac{1}{8}\cos^8 \theta - \frac{1}{6}\cos^6 \theta + C$$

Another solution:

$$\int \sin^3 \theta \cos^5 \theta d\theta = \int \sin^3 \theta (\cos^2 \theta)^2 \cos \theta d\theta = \int \sin^3 \theta (1 - \sin^2 \theta)^2 \cos \theta d\theta$$

$$= \int u^3(1-u^2)^2 du \quad \left[ \begin{array}{l} u = \sin \theta, \\ du = \cos \theta d\theta \end{array} \right] = \int u^3(1-2u^2+u^4) du$$

$$= \int (u^3 - 2u^5 + u^7) du = \frac{1}{4}u^4 - \frac{2}{6}u^6 + \frac{1}{8}u^8 + C = \frac{1}{4}\sin^4 \theta - \frac{1}{3}\sin^6 \theta + \frac{1}{8}\sin^8 \theta + C$$

14. Let  $u = 1 + x^2$ , so that  $du = 2x dx$ . Then


$$\int \frac{x^3}{\sqrt{1+x^2}} dx = \int \frac{x^2}{\sqrt{1+x^2}} (x dx) = \int \frac{u-1}{u^{1/2}} \left( \frac{1}{2} du \right) = \frac{1}{2} \int (u^{1/2} - u^{-1/2}) du = \frac{1}{2} \left( \frac{2}{3} u^{3/2} - 2u^{1/2} \right) + C$$

$$= \frac{1}{3}(1+x^2)^{3/2} - (1+x^2)^{1/2} + C \quad [\text{or } \frac{1}{3}(x^2-2)\sqrt{1+x^2} + C]$$

15. Let  $x = \sin \theta$ , where  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ . Then  $dx = \cos \theta d\theta$  and  $(1-x^2)^{1/2} = \cos \theta$ ,

so

$$\int \frac{dx}{(1-x^2)^{3/2}} = \int \frac{\cos \theta d\theta}{(\cos \theta)^3} = \int \sec^2 \theta d\theta = \tan \theta + C = \frac{x}{\sqrt{1-x^2}} + C.$$



$$16. \int_0^{\sqrt{2}/2} \frac{x^2}{\sqrt{1-x^2}} dx = \int_0^{\pi/4} \frac{\sin^2 \theta}{\cos \theta} \cos \theta d\theta \quad \left[ \begin{array}{l} u = \sin \theta, \\ du = \cos \theta d\theta \end{array} \right]$$

$$= \int_0^{\pi/4} \frac{1}{2}(1 - \cos 2\theta) d\theta = \frac{1}{2} \left[ \theta - \frac{1}{2} \sin 2\theta \right]_0^{\pi/4} = \frac{1}{2} \left[ \left( \frac{\pi}{4} - \frac{1}{2} \right) - (0 - 0) \right] = \frac{\pi}{8} - \frac{1}{4}$$

$$\begin{aligned}
 17. \int x \sin^2 x \, dx & \left[ \begin{array}{l} u = x, \quad dv = \sin^2 x \, dx, \\ du = dx \quad v = \int \sin^2 x \, dx = \int \frac{1}{2}(1 - \cos 2x) \, dx = \frac{1}{2}x - \frac{1}{2} \sin x \cos x \end{array} \right] \\
 &= \frac{1}{2}x^2 - \frac{1}{2}x \sin x \cos x - \int \left( \frac{1}{2}x - \frac{1}{2} \sin x \cos x \right) dx \\
 &= \frac{1}{2}x^2 - \frac{1}{2}x \sin x \cos x - \frac{1}{4}x^2 + \frac{1}{4} \sin^2 x + C = \frac{1}{4}x^2 - \frac{1}{2}x \sin x \cos x + \frac{1}{4} \sin^2 x + C
 \end{aligned}$$

Note:  $\int \sin x \cos x \, dx = \int s \, ds = \frac{1}{2}s^2 + C$  [where  $s = \sin x$ ,  $ds = \cos x \, dx$ ].

A slightly different method is to write  $\int x \sin^2 x \, dx = \int x \cdot \frac{1}{2}(1 - \cos 2x) \, dx = \frac{1}{2} \int x \, dx - \frac{1}{2} \int x \cos 2x \, dx$ . If we evaluate the second integral by parts, we arrive at the equivalent answer  $\frac{1}{4}x^2 - \frac{1}{4}x \sin 2x - \frac{1}{8} \cos 2x + C$ .

$$18. \text{ Let } u = e^{2t}, du = 2e^{2t} dt. \text{ Then } \int \frac{e^{2t}}{1 + e^{4t}} dt = \int \frac{\frac{1}{2}(2e^{2t}) dt}{1 + (e^{2t})^2} = \int \frac{\frac{1}{2} du}{1 + u^2} = \frac{1}{2} \tan^{-1} u + C = \frac{1}{2} \tan^{-1}(e^{2t}) + C.$$

$$19. \text{ Let } u = e^x. \text{ Then } \int e^{x+e^x} dx = \int e^{e^x} e^x dx = \int e^u du = e^u + C = e^{e^x} + C.$$

$$20. \text{ Since } e^2 \text{ is a constant, } \int e^2 dx = e^2 x + C.$$

$$21. \text{ Let } t = \sqrt{x}, \text{ so that } t^2 = x \text{ and } 2t \, dt = dx. \text{ Then } \int \arctan \sqrt{x} \, dx = \int \arctan t (2t \, dt) = I. \text{ Now use parts with}$$

$$u = \arctan t, dv = 2t \, dt \Rightarrow du = \frac{1}{1+t^2} dt, v = t^2. \text{ Thus,}$$

$$\begin{aligned}
 I &= t^2 \arctan t - \int \frac{t^2}{1+t^2} dt = t^2 \arctan t - \int \left( 1 - \frac{1}{1+t^2} \right) dt = t^2 \arctan t - t + \arctan t + C \\
 &= x \arctan \sqrt{x} - \sqrt{x} + \arctan \sqrt{x} + C \quad \left[ \text{or } (x+1) \arctan \sqrt{x} - \sqrt{x} + C \right]
 \end{aligned}$$

$$22. \text{ Let } u = 1 + (\ln x)^2, \text{ so that } du = \frac{2 \ln x}{x} dx. \text{ Then}$$

$$\int \frac{\ln x}{x \sqrt{1 + (\ln x)^2}} dx = \frac{1}{2} \int \frac{1}{\sqrt{u}} du = \frac{1}{2} (2\sqrt{u}) + C = \sqrt{1 + (\ln x)^2} + C.$$

$$23. \text{ Let } u = 1 + \sqrt{x}. \text{ Then } x = (u-1)^2, dx = 2(u-1) du \Rightarrow$$

$$\int_0^1 (1 + \sqrt{x})^8 dx = \int_1^2 u^8 \cdot 2(u-1) du = 2 \int_1^2 (u^9 - u^8) du = \left[ \frac{1}{10} u^{10} - 2 \cdot \frac{1}{9} u^9 \right]_1^2 = \frac{1024}{5} - \frac{1024}{9} - \frac{1}{5} + \frac{2}{9} = \frac{4097}{45}.$$

$$24. \text{ Let } u = \ln(x^2 - 1), dv = dx \Leftrightarrow du = \frac{2x}{x^2 - 1}, v = x. \text{ Then}$$

$$\begin{aligned}
 \int \ln(x^2 - 1) dx &= x \ln(x^2 - 1) - \int \frac{2x^2}{x^2 - 1} dx = x \ln(x^2 - 1) - \int \left[ 2 + \frac{2}{(x-1)(x+1)} \right] dx \\
 &= x \ln(x^2 - 1) - \int \left[ 2 + \frac{1}{x-1} - \frac{1}{x+1} \right] dx = x \ln(x^2 - 1) - 2x - \ln|x-1| + \ln|x+1| + C
 \end{aligned}$$

$$25. \frac{3x^2 - 2}{x^2 - 2x - 8} = 3 + \frac{6x + 22}{(x-4)(x+2)} = 3 + \frac{A}{x-4} + \frac{B}{x+2} \Rightarrow 6x + 22 = A(x+2) + B(x-4). \text{ Setting}$$

$x = 4$  gives  $46 = 6A$ , so  $A = \frac{23}{3}$ . Setting  $x = -2$  gives  $10 = -6B$ , so  $B = -\frac{5}{3}$ . Now

$$\int \frac{3x^2 - 2}{x^2 - 2x - 8} dx = \int \left( 3 + \frac{23/3}{x-4} - \frac{5/3}{x+2} \right) dx = 3x + \frac{23}{3} \ln|x-4| - \frac{5}{3} \ln|x+2| + C.$$

$$26. \int \frac{3x^2 - 2}{x^3 - 2x - 8} dx = \int \frac{du}{u} \quad \left[ \begin{array}{l} u = x^3 - 2x - 8, \\ du = (3x^2 - 2) dx \end{array} \right] = \ln|u| + C = \ln|x^3 - 2x - 8| + C$$

$$27. \text{ Let } u = 1 + e^x, \text{ so that } du = e^x dx = (u-1) dx. \text{ Then } \int \frac{1}{1+e^x} dx = \int \frac{1}{u} \cdot \frac{du}{u-1} = \int \frac{1}{u(u-1)} du = I. \text{ Now}$$

$$\frac{1}{u(u-1)} = \frac{A}{u} + \frac{B}{u-1} \Rightarrow 1 = A(u-1) + Bu. \text{ Set } u = 1 \text{ to get } 1 = B. \text{ Set } u = 0 \text{ to get } 1 = -A, \text{ so } A = -1.$$

$$\text{Thus, } I = \int \left( \frac{-1}{u} + \frac{1}{u-1} \right) du = -\ln|u| + \ln|u-1| + C = -\ln(1+e^x) + \ln e^x + C = x - \ln(1+e^x) + C.$$

*Another method:* Multiply numerator and denominator by  $e^{-x}$  and let  $u = e^{-x} + 1$ . This gives the answer in the form  $-\ln(e^{-x} + 1) + C$ .

$$\begin{aligned} 28. \int \sin \sqrt{at} dt &= \int \sin u \cdot \frac{2}{a} u du \quad [u = \sqrt{at}, u^2 = at, 2u du = a dt] = \frac{2}{a} \int u \sin u du \\ &= \frac{2}{a} [-u \cos u + \sin u] + C \quad [\text{integration by parts}] = -\frac{2}{a} \sqrt{at} \cos \sqrt{at} + \frac{2}{a} \sin \sqrt{at} + C \\ &= -2 \sqrt{\frac{t}{a}} \cos \sqrt{at} + \frac{2}{a} \sin \sqrt{at} + C \end{aligned}$$

$$\begin{aligned} 29. \int_0^5 \frac{3w-1}{w+2} dw &= \int_0^5 \left( 3 - \frac{7}{w+2} \right) dw = \left[ 3w - 7 \ln|w+2| \right]_0^5 = 15 - 7 \ln 7 + 7 \ln 2 \\ &= 15 + 7(\ln 2 - \ln 7) = 15 + 7 \ln \frac{2}{7} \end{aligned}$$

$$30. x^2 - 4x < 0 \text{ on } [0, 4], \text{ so}$$

$$\begin{aligned} \int_{-2}^2 |x^2 - 4x| dx &= \int_{-2}^0 (x^2 - 4x) dx + \int_0^2 (4x - x^2) dx = \left[ \frac{1}{3}x^3 - 2x^2 \right]_{-2}^0 + \left[ 2x^2 - \frac{1}{3}x^3 \right]_0^2 \\ &= 0 - \left( -\frac{8}{3} - 8 \right) + \left( 8 - \frac{8}{3} \right) - 0 = 16 \end{aligned}$$

$$31. \text{ As in Example 5,}$$

$$\int \sqrt{\frac{1+x}{1-x}} dx = \int \frac{\sqrt{1+x}}{\sqrt{1-x}} \cdot \frac{\sqrt{1+x}}{\sqrt{1+x}} dx = \int \frac{1+x}{\sqrt{1-x^2}} dx = \int \frac{dx}{\sqrt{1-x^2}} + \int \frac{x dx}{\sqrt{1-x^2}} = \sin^{-1} x - \sqrt{1-x^2} + C.$$


*Another method:* Substitute  $u = \sqrt{(1+x)/(1-x)}$ .

$$\begin{aligned}
 32. \int \frac{\sqrt{2x-1}}{2x+3} dx &= \int \frac{u \cdot u du}{u^2+4} \quad \left[ \begin{array}{l} u = \sqrt{2x-1}, 2x+3 = u^2+4, \\ u^2 = 2x-1, u du = dx \end{array} \right] = \int \left( 1 - \frac{4}{u^2+4} \right) du \\
 &= u - 4 \cdot \frac{1}{2} \tan^{-1} \left( \frac{1}{2} u \right) + C = \sqrt{2x-1} - 2 \tan^{-1} \left( \frac{1}{2} \sqrt{2x-1} \right) + C
 \end{aligned}$$

$$33. 3 - 2x - x^2 = -(x^2 + 2x + 1) + 4 = 4 - (x+1)^2. \text{ Let } x+1 = 2 \sin \theta,$$

where  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ . Then  $dx = 2 \cos \theta d\theta$  and

$$\begin{aligned}
 \int \sqrt{3-2x-x^2} dx &= \int \sqrt{4-(x+1)^2} dx = \int \sqrt{4-4\sin^2 \theta} 2 \cos \theta d\theta \\
 &= 4 \int \cos^2 \theta d\theta = 2 \int (1 + \cos 2\theta) d\theta \\
 &= 2\theta + \sin 2\theta + C = 2\theta + 2 \sin \theta \cos \theta + C \\
 &= 2 \sin^{-1} \left( \frac{x+1}{2} \right) + 2 \cdot \frac{x+1}{2} \cdot \frac{\sqrt{3-2x-x^2}}{2} + C \\
 &= 2 \sin^{-1} \left( \frac{x+1}{2} \right) + \frac{x+1}{2} \sqrt{3-2x-x^2} + C
 \end{aligned}$$



$$\begin{aligned}
 34. \int_{\pi/4}^{\pi/2} \frac{1+4 \cot x}{4-\cot x} dx &= \int_{\pi/4}^{\pi/2} \left[ \frac{(1+4 \cos x / \sin x)}{(4-\cos x / \sin x)} \cdot \frac{\sin x}{\sin x} \right] dx = \int_{\pi/4}^{\pi/2} \frac{\sin x + 4 \cos x}{4 \sin x - \cos x} dx \\
 &= \int_{3/\sqrt{2}}^4 \frac{1}{u} du \quad \left[ \begin{array}{l} u = 4 \sin x - \cos x, \\ du = (4 \cos x + \sin x) dx \end{array} \right] \\
 &= \left[ \ln |u| \right]_{3/\sqrt{2}}^4 = \ln 4 - \ln \frac{3}{\sqrt{2}} = \ln \frac{4}{3/\sqrt{2}} = \ln \left( \frac{4}{3} \sqrt{2} \right)
 \end{aligned}$$

35. Because  $f(x) = x^8 \sin x$  is the product of an even function and an odd function, it is odd.

Therefore,  $\int_{-1}^1 x^8 \sin x dx = 0$  [by (5.5.7)(b)].

36.  $\sin 4x \cos 3x = \frac{1}{2}(\sin x + \sin 7x)$  by Formula 7.2.2(a), so

$$\int \sin 4x \cos 3x dx = \frac{1}{2} \int (\sin x + \sin 7x) dx = \frac{1}{2} \left[ -\cos x - \frac{1}{7} \cos 7x \right] + C = -\frac{1}{2} \cos x - \frac{1}{14} \cos 7x + C.$$

$$37. \int_0^{\pi/4} \cos^2 \theta \tan^2 \theta d\theta = \int_0^{\pi/4} \sin^2 \theta d\theta = \int_0^{\pi/4} \frac{1}{2}(1 - \cos 2\theta) d\theta = \left[ \frac{1}{2}\theta - \frac{1}{4} \sin 2\theta \right]_0^{\pi/4} = \left( \frac{\pi}{8} - \frac{1}{4} \right) - (0 - 0) = \frac{\pi}{8} - \frac{1}{4}$$

$$\begin{aligned}
 38. \int_0^{\pi/4} \tan^5 \theta \sec^3 \theta d\theta &= \int_0^{\pi/4} (\tan^2 \theta)^2 \sec^2 \theta \cdot \sec \theta \tan \theta d\theta = \int_1^{\sqrt{2}} (u^2 - 1)^2 u^2 du \quad \left[ \begin{array}{l} u = \sec \theta, \\ du = \sec \theta \tan \theta d\theta \end{array} \right] \\
 &= \int_1^{\sqrt{2}} (u^6 - 2u^4 + u^2) du = \left[ \frac{1}{7} u^7 - \frac{2}{5} u^5 + \frac{1}{3} u^3 \right]_1^{\sqrt{2}} \\
 &= \left( \frac{8}{7} \sqrt{2} - \frac{8}{5} \sqrt{2} + \frac{2}{3} \sqrt{2} \right) - \left( \frac{1}{7} - \frac{2}{5} + \frac{1}{3} \right) = \frac{22}{105} \sqrt{2} - \frac{8}{105} = \frac{2}{105} (11 \sqrt{2} - 4)
 \end{aligned}$$

39. Let  $u = \sec \theta$ , so that  $du = \sec \theta \tan \theta d\theta$ . Then  $\int \frac{\sec \theta \tan \theta}{\sec^2 \theta - \sec \theta} d\theta = \int \frac{1}{u^2 - u} du = \int \frac{1}{u(u-1)} du = I$ . Now

$$\frac{1}{u(u-1)} = \frac{A}{u} + \frac{B}{u-1} \Rightarrow 1 = A(u-1) + Bu. \text{ Set } u = 1 \text{ to get } 1 = B. \text{ Set } u = 0 \text{ to get } 1 = -A, \text{ so } A = -1.$$

$$\text{Thus, } I = \int \left( \frac{-1}{u} + \frac{1}{u-1} \right) du = -\ln|u| + \ln|u-1| + C = \ln|\sec \theta - 1| - \ln|\sec \theta| + C \text{ [or } \ln|1 - \cos \theta| + C].$$

40.  $4y^2 - 4y - 3 = (2y-1)^2 - 2^2$ , so let  $u = 2y-1 \Rightarrow du = 2 dy$ . Thus,

$$\begin{aligned} \int \frac{dy}{\sqrt{4y^2 - 4y - 3}} &= \int \frac{dy}{\sqrt{(2y-1)^2 - 2^2}} = \frac{1}{2} \int \frac{du}{\sqrt{u^2 - 2^2}} \\ &= \frac{1}{2} \ln|u + \sqrt{u^2 - 2^2}| \quad [\text{by Formula 20 in the table in this section}] \\ &= \frac{1}{2} \ln|2y-1 + \sqrt{4y^2 - 4y - 3}| + C \end{aligned}$$

41. Let  $u = \theta$ ,  $dv = \tan^2 \theta d\theta = (\sec^2 \theta - 1) d\theta \Rightarrow du = d\theta$  and  $v = \tan \theta - \theta$ . So

$$\begin{aligned} \int \theta \tan^2 \theta d\theta &= \theta(\tan \theta - \theta) - \int (\tan \theta - \theta) d\theta = \theta \tan \theta - \theta^2 - \ln|\sec \theta| + \frac{1}{2}\theta^2 + C \\ &= \theta \tan \theta - \frac{1}{2}\theta^2 - \ln|\sec \theta| + C \end{aligned}$$

42. Let  $u = \tan^{-1} x$ ,  $dv = \frac{1}{x^2} dx \Rightarrow du = \frac{1}{1+x^2} dx$ ,  $v = -\frac{1}{x}$ . Then

$$I = \int \frac{\tan^{-1} x}{x^2} dx = -\frac{1}{x} \tan^{-1} x - \int \left( -\frac{1}{x(1+x^2)} \right) dx = -\frac{1}{x} \tan^{-1} x + \int \left( \frac{A}{x} + \frac{Bx+C}{1+x^2} \right) dx$$

$$\frac{1}{x(1+x^2)} = \frac{A}{x} + \frac{Bx+C}{1+x^2} \Rightarrow 1 = A(1+x^2) + (Bx+C)x \Rightarrow 1 = (A+B)x^2 + Cx + A, \text{ so } C = 0, A = 1,$$


and  $A+B=0 \Rightarrow B=-1$ . Thus,

$$\begin{aligned} I &= -\frac{1}{x} \tan^{-1} x + \int \left( \frac{1}{x} - \frac{x}{1+x^2} \right) dx = -\frac{1}{x} \tan^{-1} x + \ln|x| - \frac{1}{2} \ln|1+x^2| + C \\ &= -\frac{\tan^{-1} x}{x} + \ln \left| \frac{x}{\sqrt{x^2+1}} \right| + C \end{aligned}$$

Or: Let  $x = \tan \theta$ , so that  $dx = \sec^2 \theta d\theta$ . Then  $\int \frac{\tan^{-1} x}{x^2} dx = \int \frac{\theta}{\tan^2 \theta} \sec^2 \theta d\theta = \int \theta \csc^2 \theta d\theta = I$ . Now use parts

with  $u = \theta$ ,  $dv = \csc^2 \theta d\theta \Rightarrow du = d\theta$ ,  $v = -\cot \theta$ . Thus,

$$\begin{aligned} I &= -\theta \cot \theta - \int (-\cot \theta) d\theta = -\theta \cot \theta + \ln|\sin \theta| + C \\ &= -\tan^{-1} x \cdot \frac{1}{x} + \ln \left| \frac{x}{\sqrt{x^2+1}} \right| + C = -\frac{\tan^{-1} x}{x} + \ln \left| \frac{x}{\sqrt{x^2+1}} \right| + C \end{aligned}$$



43. Let  $u = 1 + e^x$ , so that  $du = e^x dx$ . Then  $\int e^x \sqrt{1 + e^x} dx = \int u^{1/2} du = \frac{2}{3} u^{3/2} + C = \frac{2}{3} (1 + e^x)^{3/2} + C$ .

Or: Let  $u = \sqrt{1 + e^x}$ , so that  $u^2 = 1 + e^x$  and  $2u du = e^x dx$ . Then

$$\int e^x \sqrt{1 + e^x} dx = \int u \cdot 2u du = \int 2u^2 du = \frac{2}{3} u^3 + C = \frac{2}{3} (1 + e^x)^{3/2} + C.$$

44. Let  $u = \sqrt{1 + e^x}$ . Then  $u^2 = 1 + e^x$ ,  $2u du = e^x dx = (u^2 - 1) dx$ , and  $dx = \frac{2u}{u^2 - 1} du$ , so

$$\begin{aligned} \int \sqrt{1 + e^x} dx &= \int u \cdot \frac{2u}{u^2 - 1} du = \int \frac{2u^2}{u^2 - 1} du = \int \left( 2 + \frac{2}{u^2 - 1} \right) du = \int \left( 2 + \frac{1}{u - 1} - \frac{1}{u + 1} \right) du \\ &= 2u + \ln|u - 1| - \ln|u + 1| + C = 2\sqrt{1 + e^x} + \ln(\sqrt{1 + e^x} - 1) - \ln(\sqrt{1 + e^x} + 1) + C \end{aligned}$$

45. Let  $t = x^3$ . Then  $dt = 3x^2 dx \Rightarrow I = \int x^5 e^{-x^3} dx = \frac{1}{3} \int t e^{-t} dt$ . Now integrate by parts with  $u = t$ ,  $dv = e^{-t} dt$ :

$$I = -\frac{1}{3} t e^{-t} + \frac{1}{3} \int e^{-t} dt = -\frac{1}{3} t e^{-t} - \frac{1}{3} e^{-t} + C = -\frac{1}{3} e^{-x^3} (x^3 + 1) + C.$$

$$\begin{aligned} 46. \frac{1 + \sin x}{1 - \sin x} &= \frac{1 + \sin x}{1 - \sin x} \cdot \frac{1 + \sin x}{1 + \sin x} = \frac{1 + 2\sin x + \sin^2 x}{1 - \sin^2 x} = \frac{1 + 2\sin x + \sin^2 x}{\cos^2 x} = \frac{1}{\cos^2 x} + \frac{2\sin x}{\cos^2 x} + \frac{\sin^2 x}{\cos^2 x} \\ &= \sec^2 x + 2\sec x \tan x + \tan^2 x = \sec^2 x + 2\sec x \tan x + \sec^2 x - 1 = 2\sec^2 x + 2\sec x \tan x - 1 \end{aligned}$$

Thus, 
$$\int \frac{1 + \sin x}{1 - \sin x} dx = \int (2\sec^2 x + 2\sec x \tan x - 1) dx = 2\tan x + 2\sec x - x + C$$

47. Let  $u = x - 1$ , so that  $du = dx$ . Then

$$\begin{aligned} \int x^3 (x - 1)^{-4} dx &= \int (u + 1)^3 u^{-4} du = \int (u^3 + 3u^2 + 3u + 1) u^{-4} du = \int (u^{-1} + 3u^{-2} + 3u^{-3} + u^{-4}) du \\ &= \ln|u| - 3u^{-1} - \frac{3}{2} u^{-2} - \frac{1}{3} u^{-3} + C = \ln|x - 1| - 3(x - 1)^{-1} - \frac{3}{2} (x - 1)^{-2} - \frac{1}{3} (x - 1)^{-3} + C \end{aligned}$$

$$48. \text{ Let } u = x^2. \text{ Then } du = 2x dx \Rightarrow \int \frac{x dx}{x^4 - a^4} = \int \frac{\frac{1}{2} du}{u^2 - (a^2)^2} = \frac{1}{4a^2} \ln \left| \frac{u - a^2}{u + a^2} \right| + C = \frac{1}{4a^2} \ln \left| \frac{x^2 - a^2}{x^2 + a^2} \right| + C.$$

49. Let  $u = \sqrt{4x + 1} \Rightarrow u^2 = 4x + 1 \Rightarrow 2u du = 4 dx \Rightarrow dx = \frac{1}{2} u du$ . So

$$\begin{aligned} \int \frac{1}{x \sqrt{4x + 1}} dx &= \int \frac{\frac{1}{2} u du}{\frac{1}{4} (u^2 - 1) u} = 2 \int \frac{du}{u^2 - 1} = 2 \left( \frac{1}{2} \right) \ln \left| \frac{u - 1}{u + 1} \right| + C \quad [\text{by Formula 19}] \\ &= \ln \left| \frac{\sqrt{4x + 1} - 1}{\sqrt{4x + 1} + 1} \right| + C \end{aligned}$$

50. As in Exercise 49, let  $u = \sqrt{4x+1}$ . Then  $\int \frac{dx}{x^2 \sqrt{4x+1}} = \int \frac{\frac{1}{2}u du}{\left[\frac{1}{4}(u^2-1)\right]^2 u} = 8 \int \frac{du}{(u^2-1)^2}$ .

$$\text{Now } \frac{1}{(u^2-1)^2} = \frac{1}{(u+1)^2(u-1)^2} = \frac{A}{u+1} + \frac{B}{(u+1)^2} + \frac{C}{u-1} + \frac{D}{(u-1)^2} \Rightarrow$$

$$1 = A(u+1)(u-1)^2 + B(u-1)^2 + C(u-1)(u+1)^2 + D(u+1)^2. \quad u=1 \Rightarrow D = \frac{1}{4}, \quad u=-1 \Rightarrow B = \frac{1}{4}.$$

Equating coefficients of  $u^3$  gives  $A+C=0$ , and equating coefficients of  $1$  gives  $1 = A+B-C+D \Rightarrow$

$$1 = A + \frac{1}{4} - C + \frac{1}{4} \Rightarrow \frac{1}{2} = A - C. \text{ So } A = \frac{1}{4} \text{ and } C = -\frac{1}{4}. \text{ Therefore,}$$

$$\begin{aligned} \int \frac{dx}{x^2 \sqrt{4x+1}} &= 8 \int \left[ \frac{1/4}{u+1} + \frac{1/4}{(u+1)^2} + \frac{-1/4}{u-1} + \frac{1/4}{(u-1)^2} \right] du \\ &= \int \left[ \frac{2}{u+1} + 2(u+1)^{-2} - \frac{2}{u-1} + 2(u-1)^{-2} \right] du \\ &= 2 \ln|u+1| - \frac{2}{u+1} - 2 \ln|u-1| - \frac{2}{u-1} + C \\ &= 2 \ln(\sqrt{4x+1}+1) - \frac{2}{\sqrt{4x+1}+1} - 2 \ln|\sqrt{4x+1}-1| - \frac{2}{\sqrt{4x+1}-1} + C \end{aligned}$$