

$$1. r = \theta^2, 0 \leq \theta \leq \frac{\pi}{4}. \quad A = \int_0^{\pi/4} \frac{1}{2} r^2 d\theta = \int_0^{\pi/4} \frac{1}{2} (\theta^2)^2 d\theta = \int_0^{\pi/4} \frac{1}{2} \theta^4 d\theta = \left[\frac{1}{10} \theta^5 \right]_0^{\pi/4} = \frac{1}{10} \left(\frac{\pi}{4} \right)^5 = \frac{1}{10,240} \pi^5$$

$$2. r = e^{\theta/2}, \pi \leq \theta \leq 2\pi. \quad A = \int_{\pi}^{2\pi} \frac{1}{2} (e^{\theta/2})^2 d\theta = \int_{\pi}^{2\pi} \frac{1}{2} e^{\theta} d\theta = \frac{1}{2} \left[e^{\theta} \right]_{\pi}^{2\pi} = \frac{1}{2} (e^{2\pi} - e^{\pi})$$

$$3. r = \sin \theta, \frac{\pi}{3} \leq \theta \leq \frac{2\pi}{3}.$$

$$\begin{aligned} A &= \int_{\pi/3}^{2\pi/3} \frac{1}{2} \sin^2 \theta d\theta = \frac{1}{4} \int_{\pi/3}^{2\pi/3} (1 - \cos 2\theta) d\theta = \frac{1}{4} \left[\theta - \frac{1}{2} \sin 2\theta \right]_{\pi/3}^{2\pi/3} = \frac{1}{4} \left[\frac{2\pi}{3} - \frac{1}{2} \sin \frac{4\pi}{3} - \frac{\pi}{3} + \frac{1}{2} \sin \frac{2\pi}{3} \right] \\ &= \frac{1}{4} \left[\frac{2\pi}{3} - \frac{1}{2} \left(-\frac{\sqrt{3}}{2} \right) - \frac{\pi}{3} + \frac{1}{2} \left(\frac{\sqrt{3}}{2} \right) \right] = \frac{1}{4} \left(\frac{\pi}{3} + \frac{\sqrt{3}}{2} \right) = \frac{\pi}{12} + \frac{\sqrt{3}}{8} \end{aligned}$$

$$4. r = \sqrt{\sin \theta}, 0 \leq \theta \leq \pi. \quad A = \int_0^{\pi} \frac{1}{2} (\sqrt{\sin \theta})^2 d\theta = \int_0^{\pi} \frac{1}{2} \sin \theta d\theta = \left[-\frac{1}{2} \cos \theta \right]_0^{\pi} = \frac{1}{2} + \frac{1}{2} = 1$$

$$5. r = \sqrt{\theta}, 0 \leq \theta \leq 2\pi. \quad A = \int_0^{2\pi} \frac{1}{2} r^2 d\theta = \int_0^{2\pi} \frac{1}{2} (\sqrt{\theta})^2 d\theta = \int_0^{2\pi} \frac{1}{2} \theta d\theta = \left[\frac{1}{4} \theta^2 \right]_0^{2\pi} = \pi^2$$

$$6. r = 1 + \cos \theta, 0 \leq \theta \leq \pi.$$

$$\begin{aligned} A &= \int_0^{\pi} \frac{1}{2} (1 + \cos \theta)^2 d\theta = \frac{1}{2} \int_0^{\pi} (1 + 2 \cos \theta + \cos^2 \theta) d\theta = \frac{1}{2} \int_0^{\pi} \left[1 + 2 \cos \theta + \frac{1}{2} (1 + \cos 2\theta) \right] d\theta \\ &= \frac{1}{2} \int_0^{\pi} \left(\frac{3}{2} + 2 \cos \theta + \frac{1}{2} \cos 2\theta \right) d\theta = \frac{1}{2} \left[\frac{3}{2} \theta + 2 \sin \theta + \frac{1}{4} \sin 2\theta \right]_0^{\pi} = \frac{1}{2} \left(\frac{3}{2} \pi + 0 + 0 \right) - \frac{1}{2} (0) = \frac{3\pi}{4} \end{aligned}$$

$$7. r = 4 + 3 \sin \theta, -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}.$$

$$\begin{aligned} A &= \int_{-\pi/2}^{\pi/2} \frac{1}{2} ((4 + 3 \sin \theta)^2) d\theta = \frac{1}{2} \int_{-\pi/2}^{\pi/2} (16 + 24 \sin \theta + 9 \sin^2 \theta) d\theta = \frac{1}{2} \int_{-\pi/2}^{\pi/2} (16 + 9 \sin^2 \theta) d\theta \quad [\text{by Theorem 5.5.7(b)}] \\ &= \frac{1}{2} \cdot 2 \int_0^{\pi/2} [16 + 9 \cdot \frac{1}{2} (1 - \cos 2\theta)] d\theta \quad [\text{by Theorem 5.5.7(a)}] \\ &= \int_0^{\pi/2} \left(\frac{41}{2} - \frac{9}{2} \cos 2\theta \right) d\theta = \left[\frac{41}{2} \theta - \frac{9}{4} \sin 2\theta \right]_0^{\pi/2} = \left(\frac{41\pi}{4} - 0 \right) - (0 - 0) = \frac{41\pi}{4} \end{aligned}$$

$$8. r = \sin 2\theta, 0 \leq \theta \leq \frac{\pi}{2}.$$


$$A = \int_0^{\pi/2} \frac{1}{2} \sin^2 2\theta d\theta = \frac{1}{2} \int_0^{\pi/2} \frac{1}{2} (1 - \cos 4\theta) d\theta = \frac{1}{4} \left[\theta - \frac{1}{4} \sin 4\theta \right]_0^{\pi/2} = \frac{1}{4} \left(\frac{\pi}{2} \right) = \frac{\pi}{8}$$

$$9. \text{ The area above the polar axis is bounded by } r = 3 \cos \theta \text{ for } \theta = 0$$


to $\theta = \pi/2$ [not π]. By symmetry,

$$\begin{aligned} A &= 2 \int_0^{\pi/2} \frac{1}{2} r^2 d\theta = \int_0^{\pi/2} (3 \cos \theta)^2 d\theta = 3^2 \int_0^{\pi/2} \cos^2 \theta d\theta \\ &= 9 \int_0^{\pi/2} \frac{1}{2} (1 + \cos 2\theta) d\theta = \frac{9}{2} \left[\theta + \frac{1}{2} \sin 2\theta \right]_0^{\pi/2} = \frac{9}{2} \left[\left(\frac{\pi}{2} + 0 \right) - (0 + 0) \right] = \frac{9\pi}{4} \end{aligned}$$

Also, note that this is a circle with radius $\frac{3}{2}$, so its area is $\pi \left(\frac{3}{2} \right)^2 = \frac{9\pi}{4}$.




$$\begin{aligned}
 10. A &= \int_0^{2\pi} \frac{1}{2} r^2 d\theta = \int_0^{2\pi} \frac{1}{2} [3(1 + \cos \theta)]^2 d\theta \\
 &= \frac{9}{2} \int_0^{2\pi} (1 + 2 \cos \theta + \cos^2 \theta) d\theta \\
 &= \frac{9}{2} \int_0^{2\pi} \left[1 + 2 \cos \theta + \frac{1}{2}(1 + \cos 2\theta) \right] d\theta \\
 &= \frac{9}{2} \left[\frac{3}{2}\theta + 2 \sin \theta + \frac{1}{4} \sin 2\theta \right]_0^{2\pi} = \frac{27}{2} \pi
 \end{aligned}$$



11. The curve goes through the pole when $\theta = \pi/4$, so we'll find the area for


$0 \leq \theta \leq \pi/4$ and multiply it by 4.

$$\begin{aligned}
 A &= 4 \int_0^{\pi/4} \frac{1}{2} r^2 d\theta = 2 \int_0^{\pi/4} (4 \cos 2\theta) d\theta \\
 &= 8 \int_0^{\pi/4} \cos 2\theta d\theta = 4 [\sin 2\theta]_0^{\pi/4} = 4
 \end{aligned}$$



12. To find the area that the curve encloses, we'll double the area to the left of the vertical axis.

$$\begin{aligned}
 A &= 2 \int_{\pi/2}^{3\pi/2} \frac{1}{2} (2 - \sin \theta)^2 d\theta = \int_{\pi/2}^{3\pi/2} (4 - 4 \sin \theta + \sin^2 \theta) d\theta \\
 &= \int_{\pi/2}^{3\pi/2} \left[4 - 4 \sin \theta + \frac{1}{2}(1 - \cos 2\theta) \right] d\theta = \int_{\pi/2}^{3\pi/2} \left(\frac{9}{2} - 4 \sin \theta - \frac{1}{2} \cos 2\theta \right) d\theta \\
 &= \left[\frac{9}{2}\theta + 4 \cos \theta - \frac{1}{4} \sin 2\theta \right]_{\pi/2}^{3\pi/2} = \left(\frac{27\pi}{4} \right) - \left(\frac{9\pi}{4} \right) = \frac{9\pi}{2}
 \end{aligned}$$




Or: We could have doubled the area to the right of the vertical axis and integrated from $-\pi/2$ to $\pi/2$.

Or: We could have integrated from 0 to 2π [simpler arithmetic].


13. One-sixth of the area lies above the polar axis and is bounded by the curve

$r = 2 \cos 3\theta$ for $\theta = 0$ to $\theta = \pi/6$.


$$\begin{aligned}
 A &= 6 \int_0^{\pi/6} \frac{1}{2} (2 \cos 3\theta)^2 d\theta = 12 \int_0^{\pi/6} \cos^2 3\theta d\theta \\
 &= \frac{12}{2} \int_0^{\pi/6} (1 + \cos 6\theta) d\theta \\
 &= 6 \left[\theta + \frac{1}{6} \sin 6\theta \right]_0^{\pi/6} = 6 \left(\frac{\pi}{6} \right) = \pi
 \end{aligned}$$




$$\begin{aligned}
 14. A &= \int_0^{2\pi} \frac{1}{2} (2 + \cos 2\theta)^2 d\theta = \frac{1}{2} \int_0^{2\pi} (4 + 4 \cos 2\theta + \cos^2 2\theta) d\theta \\
 &= \frac{1}{2} \int_0^{2\pi} \left(4 + 4 \cos 2\theta + \frac{1}{2} + \frac{1}{2} \cos 4\theta \right) d\theta \\
 &= \frac{1}{2} \left[\frac{9}{2}\theta + 2 \sin 2\theta + \frac{1}{8} \sin 4\theta \right]_0^{2\pi} = \frac{1}{2} (9\pi) = \frac{9\pi}{2}
 \end{aligned}$$



$$\begin{aligned}
 15. A &= \int_0^{2\pi} \frac{1}{2} (1 + 2 \sin 6\theta)^2 d\theta = \frac{1}{2} \int_0^{2\pi} (1 + 4 \sin 6\theta + 4 \sin^2 6\theta) d\theta \\
 &= \frac{1}{2} \int_0^{2\pi} \left[1 + 4 \sin 6\theta + 4 \cdot \frac{1}{2} (1 - \cos 12\theta) \right] d\theta \\
 &= \frac{1}{2} \int_0^{2\pi} (3 + 4 \sin 6\theta - 2 \cos 12\theta) d\theta \\
 &= \frac{1}{2} \left[3\theta - \frac{2}{3} \cos 6\theta - \frac{1}{6} \sin 12\theta \right]_0^{2\pi} \\
 &= \frac{1}{2} \left[\left(6\pi - \frac{2}{3} - 0 \right) - \left(0 - \frac{2}{3} - 0 \right) \right] = 3\pi
 \end{aligned}$$




$$\begin{aligned}
 16. A &= \int_0^{\pi/2} \frac{1}{2} (2 \sin \theta + 3 \sin 9\theta)^2 d\theta = 2 \int_0^{\pi/2} \frac{1}{2} (2 \sin \theta + 3 \sin 9\theta)^2 d\theta \\
 &= \int_0^{\pi/2} (4 \sin^2 \theta + 12 \sin \theta \sin 9\theta + 9 \sin^2 9\theta) d\theta \\
 &= \int_0^{\pi/2} \left[2(1 - \cos 2\theta) + 12 \cdot \frac{1}{2} (\cos(\theta - 9\theta) - \cos(\theta + 9\theta)) + \frac{9}{2} (1 - \cos 18\theta) \right] d\theta \\
 &\quad \text{[integration by parts could be used for } \int \sin \theta \sin 9\theta d\theta \text{]} \\
 &= \int_0^{\pi/2} \left(2 - 2 \cos 2\theta + 6 \cos 8\theta - 6 \cos 10\theta + \frac{9}{2} - \frac{9}{2} \cos 18\theta \right) d\theta \\
 &= \left[\frac{13}{2} \theta - \sin 2\theta + \frac{3}{4} \sin 8\theta - \frac{3}{5} \sin 10\theta - \frac{1}{4} \sin 18\theta \right]_0^{\pi/2} = \frac{13\pi}{4}
 \end{aligned}$$




17. The shaded loop is traced out from $\theta = 0$ to $\theta = \pi/2$.

$$\begin{aligned}
 A &= \int_0^{\pi/2} \frac{1}{2} r^2 d\theta = \frac{1}{2} \int_0^{\pi/2} \sin^2 2\theta d\theta \\
 &= \frac{1}{2} \int_0^{\pi/2} \frac{1}{2} (1 - \cos 4\theta) d\theta = \frac{1}{4} \left[\theta - \frac{1}{4} \sin 4\theta \right]_0^{\pi/2} \\
 &= \frac{1}{4} \left(\frac{\pi}{2} \right) = \frac{\pi}{8}
 \end{aligned}$$



$$\begin{aligned}
 18. A &= \int_0^{\pi/3} \frac{1}{2} (4 \sin 3\theta)^2 d\theta = 8 \int_0^{\pi/3} \sin^2 3\theta d\theta \\
 &= 4 \int_0^{\pi/3} (1 - \cos 6\theta) d\theta \\
 &= 4 \left[\theta - \frac{1}{6} \sin 6\theta \right]_0^{\pi/3} = \frac{4\pi}{3}
 \end{aligned}$$





$$19. r = 0 \Rightarrow 3 \cos 5\theta = 0 \Rightarrow 5\theta = \frac{\pi}{2} \Rightarrow \theta = \frac{\pi}{10}.$$

$$A = \int_{-\pi/10}^{\pi/10} \frac{1}{2} (3 \cos 5\theta)^2 d\theta = \int_0^{\pi/10} 9 \cos^2 5\theta d\theta = \frac{9}{2} \int_0^{\pi/10} (1 + \cos 10\theta) d\theta = \frac{9}{2} \left[\theta + \frac{1}{10} \sin 10\theta \right]_0^{\pi/10} = \frac{9\pi}{20}$$

$$20. r = 0 \Rightarrow 2 \sin 6\theta = 0 \Rightarrow 6\theta = 0 \text{ or } \pi \Rightarrow \theta = 0 \text{ or } \frac{\pi}{6}.$$

$$A = \int_0^{\pi/6} \frac{1}{2} (2 \sin 6\theta)^2 d\theta = \int_0^{\pi/6} 2 \sin^2 6\theta d\theta = 2 \int_0^{\pi/6} \frac{1}{2} (1 - \cos 12\theta) d\theta = \left[\theta - \frac{1}{12} \sin 12\theta \right]_0^{\pi/6} = \frac{\pi}{6}$$

21.



This is a limaçon, with inner loop traced out between $\theta = \frac{7\pi}{6}$ and $\frac{11\pi}{6}$ [found by solving $r = 0$].


$$\begin{aligned} A &= 2 \int_{7\pi/6}^{3\pi/2} \frac{1}{2} (1 + 2 \sin \theta)^2 d\theta = \int_{7\pi/6}^{3\pi/2} (1 + 4 \sin \theta + 4 \sin^2 \theta) d\theta = \int_{7\pi/6}^{3\pi/2} \left[1 + 4 \sin \theta + 4 \cdot \frac{1}{2} (1 - \cos 2\theta) \right] d\theta \\ &= \left[\theta - 4 \cos \theta + 2\theta - \sin 2\theta \right]_{7\pi/6}^{3\pi/2} = \left(\frac{9\pi}{2} \right) - \left(\frac{7\pi}{2} + 2\sqrt{3} - \frac{\sqrt{3}}{2} \right) = \pi - \frac{3\sqrt{3}}{2} \end{aligned}$$

22. To determine when the strophoid $r = 2 \cos \theta - \sec \theta$ passes through the pole, we solve

$$r = 0 \Rightarrow 2 \cos \theta - \frac{1}{\cos \theta} = 0 \Rightarrow 2 \cos^2 \theta - 1 = 0 \Rightarrow \cos^2 \theta = \frac{1}{2} \Rightarrow$$


$$\cos \theta = \pm \frac{1}{\sqrt{2}} \Rightarrow \theta = \frac{\pi}{4} \text{ or } \theta = \frac{3\pi}{4} \text{ for } 0 \leq \theta \leq \pi \text{ with } \theta \neq \frac{\pi}{2}.$$

$$\begin{aligned} A &= 2 \int_0^{\pi/4} \frac{1}{2} (2 \cos \theta - \sec \theta)^2 d\theta = \int_0^{\pi/4} (4 \cos^2 \theta - 4 + \sec^2 \theta) d\theta \\ &= \int_0^{\pi/4} \left[4 \cdot \frac{1}{2} (1 + \cos 2\theta) - 4 + \sec^2 \theta \right] d\theta = \int_0^{\pi/4} (-2 + 2 \cos 2\theta + \sec^2 \theta) d\theta \\ &= \left[-2\theta + \sin 2\theta + \tan \theta \right]_0^{\pi/4} = \left(-\frac{\pi}{2} + 1 + 1 \right) - 0 = 2 - \frac{\pi}{2} \end{aligned}$$




$$23. 2 \cos \theta = 1 \Rightarrow \cos \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3} \text{ or } \frac{5\pi}{3}.$$

$$\begin{aligned} A &= 2 \int_0^{\pi/3} \frac{1}{2} [(2 \cos \theta)^2 - 1^2] d\theta = \int_0^{\pi/3} (4 \cos^2 \theta - 1) d\theta \\ &= \int_0^{\pi/3} \left\{ 4 \left[\frac{1}{2} (1 + \cos 2\theta) \right] - 1 \right\} d\theta = \int_0^{\pi/3} (1 + 2 \cos 2\theta) d\theta \\ &= \left[\theta + \sin 2\theta \right]_0^{\pi/3} = \frac{\pi}{3} + \frac{\sqrt{3}}{2} \end{aligned}$$




$$24. 1 - \sin \theta = 1 \Rightarrow \sin \theta = 0 \Rightarrow \theta = 0 \text{ or } \pi \Rightarrow$$

$$\begin{aligned} A &= \int_{\pi}^{2\pi} \frac{1}{2} [(1 - \sin \theta)^2 - 1] d\theta = \frac{1}{2} \int_{\pi}^{2\pi} (\sin^2 \theta - 2 \sin \theta) d\theta \\ &= \frac{1}{4} \int_{\pi}^{2\pi} (1 - \cos 2\theta - 4 \sin \theta) d\theta = \frac{1}{4} \left[\theta - \frac{1}{2} \sin 2\theta + 4 \cos \theta \right]_{\pi}^{2\pi} \\ &= \frac{1}{4} \pi + 2 \end{aligned}$$




25. To find the area inside the lemniscate $r^2 = 8 \cos 2\theta$ and outside the circle $r = 2$, we first note that the two curves intersect when $r^2 = 8 \cos 2\theta$ and $r = 2$, that is, when $\cos 2\theta = \frac{1}{2}$. For $-\pi < \theta \leq \pi$, $\cos 2\theta = \frac{1}{2} \Leftrightarrow 2\theta = \pm\pi/3$ or $\pm 5\pi/3 \Leftrightarrow \theta = \pm\pi/6$ or $\pm 5\pi/6$. The figure shows that the desired area is 4 times the area between the curves from 0 to $\pi/6$. Thus,



$$\begin{aligned} A &= 4 \int_0^{\pi/6} \left[\frac{1}{2}(8 \cos 2\theta) - \frac{1}{2}(2)^2 \right] d\theta = 8 \int_0^{\pi/6} (2 \cos 2\theta - 1) d\theta \\ &= 8 \left[\sin 2\theta - \theta \right]_0^{\pi/6} = 8(\sqrt{3}/2 - \pi/6) = 4\sqrt{3} - 4\pi/3 \end{aligned}$$

26. To find the shaded area A , we'll find the area A_1 inside the curve $r = 2 + \sin \theta$ and subtract $\pi(\frac{3}{2})^2$ since $r = 3 \sin \theta$ is a circle with radius $\frac{3}{2}$.




$$\begin{aligned} A_1 &= \int_0^{2\pi} \frac{1}{2}(2 + \sin \theta)^2 d\theta = \frac{1}{2} \int_0^{2\pi} (4 + 4 \sin \theta + \sin^2 \theta) d\theta \\ &= \frac{1}{2} \int_0^{2\pi} \left[4 + 4 \sin \theta + \frac{1}{2}(1 - \cos 2\theta) \right] d\theta \\ &= \frac{1}{2} \int_0^{2\pi} \left(\frac{9}{2} + 4 \sin \theta - \frac{1}{2} \cos 2\theta \right) d\theta \\ &= \frac{1}{2} \left[\frac{9}{2}\theta - 4 \cos \theta - \frac{1}{4} \sin 2\theta \right]_0^{2\pi} = \frac{1}{2}[(9\pi - 4) - (-4)] = \frac{9\pi}{2} \end{aligned}$$

So $A = A_1 - \frac{9\pi}{4} = \frac{9\pi}{2} - \frac{9\pi}{4} = \frac{9\pi}{4}$.


27. $3 \cos \theta = 1 + \cos \theta \Leftrightarrow \cos \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$ or $-\frac{\pi}{3}$.

$$\begin{aligned} A &= 2 \int_0^{\pi/3} \frac{1}{2}[(3 \cos \theta)^2 - (1 + \cos \theta)^2] d\theta \\ &= \int_0^{\pi/3} (8 \cos^2 \theta - 2 \cos \theta - 1) d\theta = \int_0^{\pi/3} [4(1 + \cos 2\theta) - 2 \cos \theta - 1] d\theta \\ &= \int_0^{\pi/3} (3 + 4 \cos 2\theta - 2 \cos \theta) d\theta = [3\theta + 2 \sin 2\theta - 2 \sin \theta]_0^{\pi/3} \\ &= \pi + \sqrt{3} - \sqrt{3} = \pi \end{aligned}$$




28. $3 \sin \theta = 2 - \sin \theta \Rightarrow 4 \sin \theta = 2 \Rightarrow \sin \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{6}$ or $\frac{5\pi}{6}$.

$$\begin{aligned} A &= 2 \int_{\pi/6}^{\pi/2} \frac{1}{2}[(3 \sin \theta)^2 - (2 - \sin \theta)^2] d\theta \\ &= \int_{\pi/6}^{\pi/2} (9 \sin^2 \theta - 4 + 4 \sin \theta - \sin^2 \theta) d\theta \\ &= \int_{\pi/6}^{\pi/2} (8 \sin^2 \theta + 4 \sin \theta - 4) d\theta \\ &= 4 \int_{\pi/6}^{\pi/2} \left[2 \cdot \frac{1}{2}(1 - \cos 2\theta) + \sin \theta - 1 \right] d\theta \\ &= 4 \int_{\pi/6}^{\pi/2} (\sin \theta - \cos 2\theta) d\theta = 4 \left[-\cos \theta - \frac{1}{2} \sin 2\theta \right]_{\pi/6}^{\pi/2} \\ &= 4 \left[(0 - 0) - \left(-\frac{\sqrt{3}}{2} - \frac{\sqrt{3}}{4} \right) \right] = 4 \left(\frac{3\sqrt{3}}{4} \right) = 3\sqrt{3} \end{aligned}$$




$$29. \sqrt{3} \cos \theta = \sin \theta \Rightarrow \sqrt{3} = \frac{\sin \theta}{\cos \theta} \Rightarrow \tan \theta = \sqrt{3} \Rightarrow \theta = \frac{\pi}{3}.$$

$$\begin{aligned} A &= \int_0^{\pi/3} \frac{1}{2} (\sin \theta)^2 d\theta + \int_{\pi/3}^{\pi/2} \frac{1}{2} (\sqrt{3} \cos \theta)^2 d\theta \\ &= \int_0^{\pi/3} \frac{1}{2} \cdot \frac{1}{2} (1 - \cos 2\theta) d\theta + \int_{\pi/3}^{\pi/2} \frac{1}{2} \cdot 3 \cdot \frac{1}{2} (1 + \cos 2\theta) d\theta \\ &= \frac{1}{4} \left[\theta - \frac{1}{2} \sin 2\theta \right]_0^{\pi/3} + \frac{3}{4} \left[\theta + \frac{1}{2} \sin 2\theta \right]_{\pi/3}^{\pi/2} \\ &= \frac{1}{4} \left[\left(\frac{\pi}{3} - \frac{\sqrt{3}}{4} \right) - 0 \right] + \frac{3}{4} \left[\left(\frac{\pi}{2} + 0 \right) - \left(\frac{\pi}{3} + \frac{\sqrt{3}}{4} \right) \right] \\ &= \frac{\pi}{12} - \frac{\sqrt{3}}{16} + \frac{\pi}{8} - \frac{3\sqrt{3}}{16} = \frac{5\pi}{24} - \frac{\sqrt{3}}{4} \end{aligned}$$




$$\begin{aligned} 30. A &= 4 \int_0^{\pi/2} \frac{1}{2} (1 - \cos \theta)^2 d\theta = 2 \int_0^{\pi/2} (1 - 2 \cos \theta + \cos^2 \theta) d\theta \\ &= 2 \int_0^{\pi/2} \left[1 - 2 \cos \theta + \frac{1}{2} (1 + \cos 2\theta) \right] d\theta \\ &= 2 \int_0^{\pi/2} \left(\frac{3}{2} - 2 \cos \theta + \frac{1}{2} \cos 2\theta \right) d\theta = \int_0^{\pi/2} (3 - 4 \cos \theta + \cos 2\theta) d\theta \\ &= \left[3\theta - 4 \sin \theta + \frac{1}{2} \sin 2\theta \right]_0^{\pi/2} = \frac{3\pi}{2} - 4 \end{aligned}$$




$$31. \sin 2\theta = \cos 2\theta \Rightarrow \frac{\sin 2\theta}{\cos 2\theta} = 1 \Rightarrow \tan 2\theta = 1 \Rightarrow 2\theta = \frac{\pi}{4} \Rightarrow \theta = \frac{\pi}{8}$$

$$\begin{aligned} A &= 8 \cdot 2 \int_0^{\pi/8} \frac{1}{2} \sin^2 2\theta d\theta = 8 \int_0^{\pi/8} \frac{1}{2} (1 - \cos 4\theta) d\theta \\ &= 4 \left[\theta - \frac{1}{4} \sin 4\theta \right]_0^{\pi/8} = 4 \left(\frac{\pi}{8} - \frac{1}{4} \cdot 1 \right) = \frac{\pi}{2} - 1 \end{aligned}$$




$$32. 3 + 2 \cos \theta = 3 + 2 \sin \theta \Rightarrow \cos \theta = \sin \theta \Rightarrow \theta = \frac{\pi}{4} \text{ or } \frac{5\pi}{4}.$$

$$\begin{aligned} A &= 2 \int_{\pi/4}^{5\pi/4} \frac{1}{2} (3 + 2 \cos \theta)^2 d\theta = \int_{\pi/4}^{5\pi/4} (9 + 12 \cos \theta + 4 \cos^2 \theta) d\theta \\ &= \int_{\pi/4}^{5\pi/4} \left[9 + 12 \cos \theta + 4 \cdot \frac{1}{2} (1 + \cos 2\theta) \right] d\theta \\ &= \int_{\pi/4}^{5\pi/4} (11 + 12 \cos \theta + 2 \cos 2\theta) d\theta = \left[11\theta + 12 \sin \theta + \sin 2\theta \right]_{\pi/4}^{5\pi/4} \\ &= \left(\frac{55\pi}{4} - 6\sqrt{2} + 1 \right) - \left(\frac{11\pi}{4} + 6\sqrt{2} + 1 \right) = 11\pi - 12\sqrt{2} \end{aligned}$$




$$33. \sin 2\theta = \cos 2\theta \Rightarrow \tan 2\theta = 1 \Rightarrow 2\theta = \frac{\pi}{4} \Rightarrow \theta = \frac{\pi}{8}$$

$$\begin{aligned} A &= 4 \int_0^{\pi/8} \frac{1}{2} \sin 2\theta d\theta \quad [\text{since } r^2 = \sin 2\theta] \\ &= \int_0^{\pi/8} 2 \sin 2\theta d\theta = [-\cos 2\theta]_0^{\pi/8} \\ &= -\frac{1}{2} \sqrt{2} - (-1) = 1 - \frac{1}{2} \sqrt{2} \end{aligned}$$



34. Let $\alpha = \tan^{-1}(b/a)$. Then


$$\begin{aligned} A &= \int_0^\alpha \frac{1}{2} (a \sin \theta)^2 d\theta + \int_\alpha^{\pi/2} \frac{1}{2} (b \cos \theta)^2 d\theta \\ &= \frac{1}{4} a^2 \left[\theta - \frac{1}{2} \sin 2\theta \right]_0^\alpha + \frac{1}{4} b^2 \left[\theta + \frac{1}{2} \sin 2\theta \right]_\alpha^{\pi/2} \\ &= \frac{1}{4} a^2 (\alpha^2 - b^2) + \frac{1}{8} \pi b^2 - \frac{1}{4} (a^2 + b^2) (\sin \alpha \cos \alpha) \\ &= \frac{1}{4} (a^2 - b^2) \tan^{-1}(b/a) + \frac{1}{8} \pi b^2 - \frac{1}{4} ab \end{aligned}$$




35. The darker shaded region (from $\theta = 0$ to $\theta = 2\pi/3$) represents $\frac{1}{2}$ of the desired area plus $\frac{1}{2}$ of the area of the inner loop.

From this area, we'll subtract $\frac{1}{2}$ of the area of the inner loop (the lighter shaded region from $\theta = 2\pi/3$ to $\theta = \pi$), and then double that difference to obtain the desired area.

$$\begin{aligned} A &= 2 \left[\int_0^{2\pi/3} \frac{1}{2} \left(\frac{1}{2} + \cos \theta \right)^2 d\theta - \int_{2\pi/3}^\pi \frac{1}{2} \left(\frac{1}{2} + \cos \theta \right)^2 d\theta \right] \\ &= \int_0^{2\pi/3} \left(\frac{1}{4} + \cos \theta + \cos^2 \theta \right) d\theta - \int_{2\pi/3}^\pi \left(\frac{1}{4} + \cos \theta + \cos^2 \theta \right) d\theta \\ &= \int_0^{2\pi/3} \left[\frac{1}{4} + \cos \theta + \frac{1}{2} (1 + \cos 2\theta) \right] d\theta \\ &\quad - \int_{2\pi/3}^\pi \left[\frac{1}{4} + \cos \theta + \frac{1}{2} (1 + \cos 2\theta) \right] d\theta \\ &= \left[\frac{\theta}{4} + \sin \theta + \frac{\theta}{2} + \frac{\sin 2\theta}{4} \right]_0^{2\pi/3} - \left[\frac{\theta}{4} + \sin \theta + \frac{\theta}{2} + \frac{\sin 2\theta}{4} \right]_{2\pi/3}^\pi \\ &= \left(\frac{\pi}{6} + \frac{\sqrt{3}}{2} + \frac{\pi}{3} - \frac{\sqrt{3}}{8} \right) - \left(\frac{\pi}{4} + \frac{\pi}{2} \right) + \left(\frac{\pi}{6} + \frac{\sqrt{3}}{2} + \frac{\pi}{3} - \frac{\sqrt{3}}{8} \right) \\ &= \frac{\pi}{4} + \frac{3}{4} \sqrt{3} = \frac{1}{4} (\pi + 3\sqrt{3}) \end{aligned}$$



36. $r = 0 \Rightarrow 1 + 2 \cos 3\theta = 0 \Rightarrow \cos 3\theta = -\frac{1}{2} \Rightarrow 3\theta = \frac{2\pi}{3}, \frac{4\pi}{3}$ [for $0 \leq 3\theta \leq 2\pi$] $\Rightarrow \theta = \frac{2\pi}{9}, \frac{4\pi}{9}$. The darker shaded region (from $\theta = 0$ to $\theta = 2\pi/9$) represents $\frac{1}{2}$ of the desired area plus $\frac{1}{2}$ of the area of the inner loop. From this area, we'll subtract $\frac{1}{2}$ of the area of the inner loop (the lighter shaded region from $\theta = 2\pi/9$ to $\theta = \pi/3$), and then double that difference to obtain the desired area.



$$A = 2 \left[\int_0^{2\pi/9} \frac{1}{2} (1 + 2 \cos 3\theta)^2 d\theta - \int_{2\pi/9}^{\pi/3} \frac{1}{2} (1 + 2 \cos 3\theta)^2 d\theta \right]$$

Now

$$\begin{aligned} r^2 &= (1 + 2 \cos 3\theta)^2 = 1 + 4 \cos 3\theta + 4 \cos^2 3\theta = 1 + 4 \cos 3\theta + 4 \cdot \frac{1}{2} (1 + \cos 6\theta) \\ &= 1 + 4 \cos 3\theta + 2 + 2 \cos 6\theta = 3 + 4 \cos 3\theta + 2 \cos 6\theta \end{aligned}$$

and $\int r^2 d\theta = 3\theta + \frac{4}{3} \sin 3\theta + \frac{1}{3} \sin 6\theta + C$, so


$$\begin{aligned} A &= \left[3\theta + \frac{4}{3} \sin 3\theta + \frac{1}{3} \sin 6\theta \right]_0^{2\pi/9} - \left[3\theta + \frac{4}{3} \sin 3\theta + \frac{1}{3} \sin 6\theta \right]_{2\pi/9}^{\pi/3} \\ &= \left[\left(\frac{2\pi}{3} + \frac{4}{3} \cdot \frac{\sqrt{3}}{2} + \frac{1}{3} \cdot \frac{-\sqrt{3}}{2} \right) - 0 \right] - \left[(\pi + 0 + 0) - \left(\frac{2\pi}{3} + \frac{4}{3} \cdot \frac{\sqrt{3}}{2} + \frac{1}{3} \cdot \frac{-\sqrt{3}}{2} \right) \right] \\ &= \frac{4\pi}{3} + \frac{4}{3}\sqrt{3} - \frac{1}{3}\sqrt{3} - \pi = \frac{\pi}{3} + \sqrt{3} \end{aligned}$$

37. The pole is a point of intersection.

$$1 + \sin \theta = 3 \sin \theta \Rightarrow 1 = 2 \sin \theta \Rightarrow \sin \theta = \frac{1}{2} \Rightarrow$$

$$\theta = \frac{\pi}{6} \text{ or } \frac{5\pi}{6}.$$

The other two points of intersection are $(\frac{3}{2}, \frac{\pi}{6})$ and $(\frac{3}{2}, \frac{5\pi}{6})$.




38. The pole is a point of intersection.

$$1 - \cos \theta = 1 + \sin \theta \Rightarrow -\cos \theta = \sin \theta \Rightarrow -1 = \tan \theta \Rightarrow$$

$$\theta = \frac{3\pi}{4} \text{ or } \frac{7\pi}{4}.$$

The other two points of intersection are $(1 + \frac{\sqrt{2}}{2}, \frac{3\pi}{4})$ and $(1 - \frac{\sqrt{2}}{2}, \frac{7\pi}{4})$.




39. $2 \sin 2\theta = 1 \Rightarrow \sin 2\theta = \frac{1}{2} \Rightarrow 2\theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{13\pi}{6}, \text{ or } \frac{17\pi}{6}.$

By symmetry, the eight points of intersection are given by

$(1, \theta)$, where $\theta = \frac{\pi}{12}, \frac{5\pi}{12}, \frac{13\pi}{12}, \text{ and } \frac{17\pi}{12}$, and

$(-1, \theta)$, where $\theta = \frac{7\pi}{12}, \frac{11\pi}{12}, \frac{19\pi}{12}, \text{ and } \frac{23\pi}{12}.$

[There are many ways to describe these points.]




40. Clearly the pole lies on both curves. $\sin 3\theta = \cos 3\theta \Rightarrow \tan 3\theta = 1 \Rightarrow$

$3\theta = \frac{\pi}{4} + n\pi$ [n any integer] $\Rightarrow \theta = \frac{\pi}{12} + \frac{\pi}{3}n \Rightarrow$

$\theta = \frac{\pi}{12}, \frac{5\pi}{12}, \text{ or } \frac{3\pi}{4}$, so the three remaining intersection points are $(\frac{1}{\sqrt{2}}, \frac{\pi}{12})$,

$(-\frac{1}{\sqrt{2}}, \frac{5\pi}{12})$, and $(\frac{1}{\sqrt{2}}, \frac{3\pi}{4})$.




41. The pole is a point of intersection. $\sin \theta = \sin 2\theta = 2 \sin \theta \cos \theta \Leftrightarrow$

$\sin \theta (1 - 2 \cos \theta) = 0 \Leftrightarrow \sin \theta = 0 \text{ or } \cos \theta = \frac{1}{2} \Rightarrow$

$\theta = 0, \pi, \frac{\pi}{3}, \text{ or } -\frac{\pi}{3} \Rightarrow$ the other intersection points are $(\frac{\sqrt{3}}{2}, \frac{\pi}{3})$

and $(\frac{\sqrt{3}}{2}, \frac{2\pi}{3})$ [by symmetry].




42. Clearly the pole is a point of intersection. $\sin 2\theta = \cos 2\theta \Rightarrow$



$\tan 2\theta = 1 \Rightarrow 2\theta = \frac{\pi}{4} + 2n\pi$ [since $\sin 2\theta$ and $\cos 2\theta$ must be

positive in the equations] $\Rightarrow \theta = \frac{\pi}{8} + n\pi \Rightarrow \theta = \frac{\pi}{8} \text{ or } \frac{9\pi}{8}.$

So the curves also intersect at $(\frac{1}{\sqrt{2}}, \frac{\pi}{8})$ and $(\frac{1}{\sqrt{2}}, \frac{9\pi}{8})$.




43.



From the first graph, we see that the pole is one point of intersection. By zooming in or using the cursor, we find the θ -values of the intersection points to be $\alpha \approx 0.88786 \approx 0.89$ and $\pi - \alpha \approx 2.25$. (The first of these values may be more easily estimated by plotting $y = 1 + \sin x$ and $y = 2x$ in rectangular coordinates; see the second graph.) By symmetry, the total area contained is twice the area contained in the first quadrant, that is,

$$\begin{aligned} A &= 2 \int_0^\alpha \frac{1}{2} (2\theta)^2 d\theta + 2 \int_\alpha^{\pi/2} \frac{1}{2} (1 + \sin \theta)^2 d\theta = \int_0^\alpha 4\theta^2 d\theta + \int_\alpha^{\pi/2} [1 + 2\sin \theta + \frac{1}{2}(1 - \cos 2\theta)] d\theta \\ &= \left[\frac{4}{3} \theta^3 \right]_0^\alpha + \left[\theta - 2\cos \theta + \left(\frac{1}{2} \theta - \frac{1}{4} \sin 2\theta \right) \right]_\alpha^{\pi/2} = \frac{4}{3} \alpha^3 + \left[\left(\frac{\pi}{2} + \frac{\pi}{4} \right) - \left(\alpha - 2\cos \alpha + \frac{1}{2} \alpha - \frac{1}{4} \sin 2\alpha \right) \right] \approx 3.4645 \end{aligned}$$

44.



We need to find the shaded area A in the figure. The horizontal line representing the front of the stage has equation $y = 4 \Leftrightarrow$

$r \sin \theta = 4 \Rightarrow r = 4 / \sin \theta$. This line intersects the curve

$$r = 8 + 8 \sin \theta \text{ when } 8 + 8 \sin \theta = \frac{4}{\sin \theta} \Rightarrow$$

$$8 \sin \theta + 8 \sin^2 \theta = 4 \Rightarrow 2 \sin^2 \theta + 2 \sin \theta - 1 = 0 \Rightarrow$$

$$\sin \theta = \frac{-2 \pm \sqrt{4+8}}{4} = \frac{-2 \pm 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2} \quad [\text{the other value is less than } -1] \Rightarrow \theta = \sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right).$$


This angle is about 21.5° and is denoted by α in the figure.

$$\begin{aligned} A &= 2 \int_{\alpha}^{\pi/2} \frac{1}{2} (8 + 8 \sin \theta)^2 d\theta - 2 \int_{\alpha}^{\pi/2} \frac{1}{2} (4 \csc \theta)^2 d\theta = 64 \int_{\alpha}^{\pi/2} (1 + 2 \sin \theta + \sin^2 \theta) d\theta - 16 \int_{\alpha}^{\pi/2} \csc^2 \theta d\theta \\ &= 64 \int_{\alpha}^{\pi/2} \left(1 + 2 \sin \theta + \frac{1}{2} - \frac{1}{2} \cos 2\theta\right) d\theta + 16 \int_{\alpha}^{\pi/2} (-\csc^2 \theta) d\theta = 64 \left[\frac{3}{2}\theta - 2 \cos \theta - \frac{1}{4} \sin 2\theta\right]_{\alpha}^{\pi/2} + 16 [\cot \theta]_{\alpha}^{\pi/2} \\ &= 16 [6\theta - 8 \cos \theta - \sin 2\theta + \cot \theta]_{\alpha}^{\pi/2} = 16 [(3\pi - 0 - 0 + 0) - (6\alpha - 8 \cos \alpha - \sin 2\alpha + \cot \alpha)] \\ &= 48\pi - 96\alpha + 128 \cos \alpha + 16 \sin 2\alpha - 16 \cot \alpha \end{aligned}$$

$$\text{From the figure, } x^2 + (\sqrt{3}-1)^2 = 2^2 \Rightarrow x^2 = 4 - (3 - 2\sqrt{3} + 1) \Rightarrow$$

$$x^2 = 2\sqrt{3} = \sqrt{12}, \text{ so } x = \sqrt{2\sqrt{3}} = \sqrt[4]{12}. \text{ Using the trigonometric relationships}$$

for a right triangle and the identity $\sin 2\alpha = 2 \sin \alpha \cos \alpha$, we continue:



$$\begin{aligned} A &= 48\pi - 96\alpha + 128 \cdot \frac{\sqrt[4]{12}}{2} + 16 \cdot 2 \cdot \frac{\sqrt{3}-1}{2} \cdot \frac{\sqrt[4]{12}}{2} - 16 \cdot \frac{\sqrt[4]{12}}{\sqrt{3}-1} \cdot \frac{\sqrt{3}+1}{\sqrt{3}+1} \\ &= 48\pi - 96\alpha + 64 \sqrt[4]{12} + 8 \sqrt[4]{12} (\sqrt{3}-1) - 8 \sqrt[4]{12} (\sqrt{3}+1) = 48\pi + 48 \sqrt[4]{12} - 96 \sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right) \\ &\approx 204.16 \text{ m}^2 \end{aligned}$$

$$\begin{aligned} 45. L &= \int_a^b \sqrt{r^2 + (dr/d\theta)^2} d\theta = \int_0^{\pi/3} \sqrt{(3 \sin \theta)^2 + (3 \cos \theta)^2} d\theta = \int_0^{\pi/3} \sqrt{9(\sin^2 \theta + \cos^2 \theta)} d\theta \\ &= 3 \int_0^{\pi/3} d\theta = 3[\theta]_0^{\pi/3} = 3\left(\frac{\pi}{3}\right) = \pi. \end{aligned}$$

As a check, note that the circumference of a circle with radius $\frac{3}{2}$ is $2\pi\left(\frac{3}{2}\right) = 3\pi$, and since $\theta = 0$ to $\pi = \frac{\pi}{3}$ traces out $\frac{1}{3}$ of the circle (from $\theta = 0$ to $\theta = \pi$), $\frac{1}{3}(3\pi) = \pi$.

$$\begin{aligned} 46. L &= \int_a^b \sqrt{r^2 + (dr/d\theta)^2} d\theta = \int_0^{2\pi} \sqrt{(e^{2\theta})^2 + (2e^{2\theta})^2} d\theta = \int_0^{2\pi} \sqrt{e^{4\theta} + 4e^{4\theta}} d\theta = \int_0^{2\pi} \sqrt{5e^{4\theta}} d\theta \\ &= \sqrt{5} \int_0^{2\pi} e^{2\theta} d\theta = \frac{\sqrt{5}}{2} [e^{2\theta}]_0^{2\pi} = \frac{\sqrt{5}}{2} (e^{4\pi} - 1) \end{aligned}$$

$$\begin{aligned}
 47. L &= \int_a^b \sqrt{r^2 + (dr/d\theta)^2} d\theta = \int_0^{2\pi} \sqrt{(\theta^2)^2 + (2\theta)^2} d\theta = \int_0^{2\pi} \sqrt{\theta^4 + 4\theta^2} d\theta \\
 &= \int_0^{2\pi} \theta \sqrt{\theta^2 + 4} d\theta = \int_0^{2\pi} \theta \sqrt{\theta^2 + 4} d\theta
 \end{aligned}$$

Now let $u = \theta^2 + 4$, so that $du = 2\theta d\theta$ [$\theta d\theta = \frac{1}{2} du$] and

$$\int_0^{2\pi} \theta \sqrt{\theta^2 + 4} d\theta = \int_4^{4\pi^2+4} \frac{1}{2} \sqrt{u} du = \frac{1}{2} \cdot \frac{2}{3} \left[u^{3/2} \right]_4^{4\pi^2+4} = \frac{1}{3} [4^{3/2}(\pi^2 + 1)^{3/2} - 4^{3/2}] = \frac{8}{3} [(\pi^2 + 1)^{3/2} - 1]$$


$$\begin{aligned}
 48. L &= \int_a^b \sqrt{r^2 + (dr/d\theta)^2} d\theta = \int_0^{2\pi} \sqrt{\theta^2 + 1} d\theta \stackrel{21}{=} \left[\frac{\theta}{2} \sqrt{\theta^2 + 1} + \frac{1}{2} \ln(\theta + \sqrt{\theta^2 + 1}) \right]_0^2 \\
 &= \pi \sqrt{4\pi^2 + 1} + \frac{1}{2} \ln(2\pi + \sqrt{4\pi^2 + 1})
 \end{aligned}$$

49. The curve $r = 3 \sin 2\theta$ is completely traced with

$$0 \leq \theta \leq 2\pi.$$

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = (3 \sin 2\theta)^2 + (6 \cos 2\theta)^2 \Rightarrow$$

$$L = \int_0^{2\pi} \sqrt{9 \sin^2 2\theta + 36 \cos^2 2\theta} d\theta \approx 29.0653$$



50. The curve $r = 4 \sin 3\theta$ is completely traced with

$$0 \leq \theta \leq \pi.$$

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = (4 \sin 3\theta)^2 + (12 \cos 3\theta)^2 \Rightarrow$$

$$L = \int_0^\pi \sqrt{16 \sin^2 3\theta + 144 \cos^2 3\theta} d\theta \approx 26.7298$$

