

This print-out should have 35 questions.
Multiple-choice questions may continue on
the next column or page – find all choices
before answering.

FitParabola01a
001 10.0 points

The graph of the function

$$y = ax^2 + bx + c$$

is a parabola passing through the points

$$(1, 10), \quad (-1, 8), \quad (-3, 22).$$

Find the y -intercept of this parabola.

1. y -intercept = 9
2. y -intercept = 10
3. y -intercept = 8
4. y -intercept = 6
5. y -intercept = 7

EchelonForm01e
002 10.0 points

If the augmented matrix for a system of
linear equations in variables x_1 , x_2 , and x_3 is
row equivalent to the matrix

$$B = \begin{bmatrix} 3 & 9 & 3 & 6 \\ -1 & -3 & -2 & -1 \\ -1 & -3 & -4 & 1 \end{bmatrix},$$

determine x_1 .

1. system inconsistent
2. $x_1 = 2$
3. $x_1 = -1$
4. $x_1 = 3 - 3t$, t arbitrary
5. $x_1 = -1 - 3t$, t arbitrary
6. $x_1 = 3$

4. $\lambda = -5$

5. $\lambda = 5, -5$

6. $\lambda = 5$

M340LSpanM02
003 10.0 points

Given

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 2 \\ 4 \\ 2 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix},$$

determine all values of λ for which

$$\mathbf{w} = \begin{bmatrix} -2 \\ 3 \\ \lambda \end{bmatrix}$$

is a vector in $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$?

1. $\lambda = 5, 1$

2. $\lambda = -5, 1$

3. $\lambda = 1$

MatEquTF03
004 10.0 points

If A is an $m \times n$ matrix and the equation $A\mathbf{x} = \mathbf{b}$ is consistent for some \mathbf{b} in \mathbb{R}^m , then the columns of A span \mathbb{R}^m .

True or False?


1. TRUE

2. FALSE

1. # molecules = 60
2. # molecules = 57
3. # molecules = 66
4. # molecules = 63
5. # molecules = 69

BalChemEq02a**005 10.0 points**

During photosynthesis green plants convert carbon dioxide CO_2 and water H_2O into glucose $\text{C}_6\text{H}_{12}\text{O}_6$ and oxygen O_2 , represented chemically by



If 11 molecules of glucose were produced in one particular conversion, how many molecules of carbon dioxide were used?

SpanTF04
006 10.0 points

If \mathbf{u} , \mathbf{v} , and \mathbf{w} are nonzero vectors in \mathbb{R}^2 and \mathbf{u} is not a multiple of \mathbf{v} , is \mathbf{w} a linear combination of \mathbf{u} and \mathbf{v} ?

1. ALWAYS
2. NEVER
3. SOMETIMES

LinTransform02a
007 10.0 points

If A is an $m \times n$ matrix, then the range of the transformation

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m, \quad T_A : \mathbf{x} \rightarrow A\mathbf{x},$$

is the set of all linear combinations of the columns of A .

True or False?

1. TRUE
2. FALSE

MatrixTrans02a
008 10.0 points

If $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ is a linear transformation such that

$$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 3 \end{bmatrix}, \quad T(\mathbf{e}_2) = \begin{bmatrix} -1 \\ 4 \end{bmatrix},$$

and $T(\mathbf{e}_3) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, determine $T(\mathbf{u})$ when

$$\mathbf{u} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}.$$

$$1. T(\mathbf{u}) = \begin{bmatrix} 5 \\ 18 \end{bmatrix}$$

$$2. T(\mathbf{u}) = \begin{bmatrix} 6 \\ 19 \end{bmatrix}$$

$$3. T(\mathbf{u}) = \begin{bmatrix} 4 \\ 18 \end{bmatrix}$$

$$4. T(\mathbf{u}) = \begin{bmatrix} 4 \\ 19 \end{bmatrix}$$

$$5. T(\mathbf{u}) = \begin{bmatrix} 6 \\ 18 \end{bmatrix}$$

$$6. T(\mathbf{u}) = \begin{bmatrix} 5 \\ 19 \end{bmatrix}$$

InverseMatrix05b
010 10.0 points

Evaluate the matrix product $B^{-1}A^T$ when

$$A = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 2 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 3 & 1 \\ 5 & 2 \end{bmatrix}.$$

$$1. B^{-1}A^T = \begin{bmatrix} 4 & -7 \\ 1 & 2 \\ 3 & 1 \end{bmatrix}$$

$$2. B^{-1}A^T = \begin{bmatrix} 12 & -7 \\ -3 & 2 \\ -1 & 1 \end{bmatrix}$$

$$3. B^{-1}A^T = \begin{bmatrix} 4 & 1 & 3 \\ -7 & 2 & 1 \end{bmatrix}$$

$$4. B^{-1}A^T = \begin{bmatrix} 12 & -11 \\ -3 & -2 \\ -1 & -7 \end{bmatrix}$$

$$5. B^{-1}A^T = \begin{bmatrix} 12 & -3 & -1 \\ -11 & -2 & -7 \end{bmatrix}$$

$$6. B^{-1}A^T = \begin{bmatrix} 4 & 1 & 3 \\ -11 & -2 & -7 \end{bmatrix}$$

MatrixOpsTF02c
009 10.0 points

If A is an $n \times n$ matrix, then

$$(A^2)^T = (A^T)^2$$

True or False?

1. FALSE

2. TRUE

$$\mathbf{2.} \ U = \begin{bmatrix} 3 & 2 & 4 & 2 \\ 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 3 \end{bmatrix}$$

$$\mathbf{3.} \ U = \begin{bmatrix} 1 & -2 & -4 & -2 \\ 0 & 0 & 1 & -5 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathbf{4.} \ U = \begin{bmatrix} 1 & -1 & 4 & -4 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathbf{5.} \ U = \begin{bmatrix} 3 & -1 & 4 & -4 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 3 \end{bmatrix}$$

$$\mathbf{6.} \ U = \begin{bmatrix} 3 & -2 & -4 & -2 \\ 0 & 0 & 1 & -5 \\ 0 & 0 & 0 & 3 \end{bmatrix}$$

InvertibleTF02a

011 10.0 points

If A and D are $n \times n$ matrices such that $AD = I$, then $DA = I$

True or False?

1. TRUE

2. FALSE

LUDecomp06g

012 10.0 points

Find U in an LU decomposition of

$$A = \begin{bmatrix} 3 & -2 & -4 & -2 \\ 9 & -6 & -11 & -11 \\ 12 & -8 & -13 & -20 \end{bmatrix}.$$

$$\mathbf{1.} \ U = \begin{bmatrix} 1 & 2 & 4 & 2 \\ 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

2. FALSE

ColNulDimTF01a

014 10.0 points

If A is a 4×5 matrix, then

$$\dim(\text{Col}(A)) + \dim(\text{Nul}(A)) = 5.$$

True or False?


1. FALSE

2. TRUE

Subspace01cT/F

013 10.0 points

The set of points in the shaded region (including the bounding lines and assumed to stretch to $\pm\infty$ in all directions) shown in



is a subspace of \mathbb{R}^2 .

True or False?

1. TRUE

Determinant02e

015 10.0 points

Compute the determinant of the matrix

$$A = \begin{bmatrix} -1 & -1 & 2 \\ 2 & 5 & -2 \\ -2 & 1 & 8 \end{bmatrix}$$

1. $\det(A) = -7$

2. $\det(A) = -8$

3. $\det(A) = -4$

4. $\det(A) = -6$

5. $\det(A) = 5$

VectorSpaceT/F04a
017 10.0 points

The set H of all polynomials

$$\mathbf{p}(x) = a + bx^4, \quad a, b \text{ in } \mathbb{R},$$

is a subspace of the vector space \mathbb{P}_6 of all polynomials of degree at most 6.

True or False?

1. TRUE
2. FALSE

DetMult05
016 10.0 points

Evaluate $\det[B^5]$ when

$$B = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ 1 & 2 & 1 \end{bmatrix}.$$

1. $\det[B^5] = 32$
2. $\det[B^5] = -32$
3. $\det[B^5] = -10$
4. $\det[B^5] = -2$
5. $\det[B^5] = 10$

BasisNull02b
018 10.0 points

Find a basis for the Null space of the matrix

$$A = \begin{bmatrix} 3 & 6 & -6 & 0 \\ 2 & 4 & -5 & 1 \\ -2 & -4 & 3 & 1 \end{bmatrix}.$$

1. $\left\{ \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} \right\}$

2. $\left\{ \begin{bmatrix} -2 \\ 0 \\ -1 \\ 1 \end{bmatrix} \right\}$

3. $\left\{ \begin{bmatrix} 2 \\ 0 \\ -1 \\ 1 \end{bmatrix} \right\}$

4. $\left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ -1 \\ 1 \end{bmatrix} \right\}$

5. $\left\{ \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ -1 \\ 1 \end{bmatrix} \right\}$

6. $\left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ 1 \end{bmatrix} \right\}$

BasisCol02a
019 10.0 points

First find a basis for $\text{Col}(A)$ when

$$A = \begin{bmatrix} 1 & -2 & 1 \\ 2 & -3 & 5 \\ 2 & -6 & -4 \end{bmatrix} = [\mathbf{a}_1 \ \mathbf{a}_2 \ \mathbf{a}_3],$$

and then select *all* the correct statements from among the following:

I: $\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ is a linearly dependent set.

II: $\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ is a basis for \mathbb{R}^3 .

III: $\text{rank}(A) = 2$.

IV: $\text{nullity}(A) = 1$.

V: $\text{rank}(A) = 3$.

1. I, II, and V

2. II and V

3. II only

4. I, III, and IV

5. I and III

Basis02
020 10.0 points

Find a basis for the space spanned by the following vectors.

$$\begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 5 \\ -3 \\ 3 \\ -4 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 1 \\ 0 \end{bmatrix}$$

1. $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 1 \\ -1 \end{bmatrix} \right\}$
2. $\left\{ \begin{bmatrix} -2 \\ 0 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 5 \\ -3 \\ 3 \\ -4 \end{bmatrix} \right\}$
3. $\left\{ \begin{bmatrix} 3 \\ -1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 5 \\ -3 \\ 3 \\ -4 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 1 \\ 0 \end{bmatrix} \right\}$
4. $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 5 \\ -3 \\ 3 \\ -4 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 1 \\ 0 \end{bmatrix} \right\}$
5. $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 1 \\ 0 \end{bmatrix} \right\}$

CoordVec03a
021 10.0 points

Find the coordinate vector $[\mathbf{x}]_{\mathcal{B}}$ in \mathbb{R}^3 for the vector

$$\mathbf{x} = \begin{bmatrix} 3 \\ -5 \\ 4 \end{bmatrix}$$

with respect to the basis

$$\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 8 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \right\}$$

for \mathbb{R}^3 .

1. $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 5 \\ 2 \\ 0 \end{bmatrix}$
2. $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 2 \\ 5 \end{bmatrix}$

3. $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} -5 \\ -2 \\ 0 \end{bmatrix}$

4. $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix}$

5. $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 2 \\ 0 \\ -5 \end{bmatrix}$

6. $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} -2 \\ 0 \\ 5 \end{bmatrix}$

PolySpanVecTF01a

022 10.0 points

The polynomials

$$\mathbf{p}_1 = 1 - 3t + 5t^2, \quad \mathbf{p}_2 = -3 + 5t - 7t^2,$$

and

$$\mathbf{p}_3 = -4 + 5t - 6t^2, \quad \mathbf{p}_4 = 1 - t^2,$$

span \mathbb{P}_2 .

True or False?

1. TRUE

2. FALSE

$$\begin{aligned}
1. \quad P_{\mathcal{C} \leftarrow \mathcal{B}} &= \begin{bmatrix} 5 & 2 & 1 \\ 0 & 1 & 2 \\ -3 & -5 & 0 \end{bmatrix} \\
2. \quad P_{\mathcal{C} \leftarrow \mathcal{B}} &= \begin{bmatrix} -5 & 2 & 1 \\ 0 & 1 & -2 \\ -3 & -5 & 0 \end{bmatrix} \\
3. \quad P_{\mathcal{C} \leftarrow \mathcal{B}} &= \begin{bmatrix} -2 & -1 & -5 \\ 0 & -1 & 2 \\ 3 & -5 & 0 \end{bmatrix} \\
4. \quad P_{\mathcal{C} \leftarrow \mathcal{B}} &= \begin{bmatrix} 2 & 5 & 1 \\ 0 & 1 & 2 \\ -3 & -5 & 0 \end{bmatrix} \\
5. \quad P_{\mathcal{C} \leftarrow \mathcal{B}} &= \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 2 \\ -3 & 5 & 0 \end{bmatrix} \\
6. \quad P_{\mathcal{C} \leftarrow \mathcal{B}} &= \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 2 \\ -3 & -5 & 0 \end{bmatrix}
\end{aligned}$$

RankTF06c
023 10.0 points

The dimensions of the row space and column space of an $m \times n$ matrix A are the same, even if $m \neq n$.

True or False?

1. TRUE
2. FALSE

ChangeBasis04b
024 (part 1 of 2) 10.0 points

In \mathbb{P}_2 determine the change of coordinates matrix $P_{\mathcal{C} \leftarrow \mathcal{B}}$ from basis $\mathcal{B} = \{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$ to the standard monomial basis $\mathcal{C} = \{1, t, t^2\}$ when

$$\mathbf{p}_1 = 1 - 3t^2, \quad \mathbf{p}_2 = 2 + t - 5t^2$$

and

$$\mathbf{p}_3 = 1 + 2t.$$

025 (part 2 of 2) 10.0 points

Express $\mathbf{q}(t) = t^2$ as a linear combination of the polynomials in the basis \mathcal{B} .

1. $\mathbf{q} = 3\mathbf{p}_1 + 2\mathbf{p}_2 - \mathbf{p}_3$

2. $\mathbf{q} = 3\mathbf{p}_1 - 2\mathbf{p}_2 + \mathbf{p}_3$

3. $\mathbf{q} = 3\mathbf{p}_1 + 2\mathbf{p}_2 + \mathbf{p}_3$

4. $\mathbf{q} = 2\mathbf{p}_1 + 3\mathbf{p}_2 - \mathbf{p}_3$

5. $\mathbf{q} = 2\mathbf{p}_1 - 3\mathbf{p}_2 + \mathbf{p}_3$

6. $\mathbf{q} = 2\mathbf{p}_1 - 3\mathbf{p}_2 - \mathbf{p}_3$

Eigenspace02a
026 10.0 points

Find a basis for the eigenspace of the matrix

$$A = \begin{bmatrix} 1 & 3 & 3 \\ -3 & -5 & -3 \\ 3 & 3 & 1 \end{bmatrix},$$

corresponding to the eigenvalue $\lambda = -2$.

1. $\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$

2. $\begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$

3. $\begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$

4. $\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$

5. $\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$

CharPoly05a
027 10.0 points

Determine the Characteristic Polynomial of the matrix

$$A = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$$

1. $6 + 4\lambda - 10\lambda^2 + \lambda^3$
2. $6 + 10\lambda - 4\lambda^2 + \lambda^3$
3. $4 - 10\lambda + 4\lambda^2 - \lambda^3$
4. $4 + 4\lambda - 10\lambda^2 - \lambda^3$
5. $4 - 4\lambda + 10\lambda^2 - \lambda^3$
6. $6 - 10\lambda + 4\lambda^2 + \lambda^3$

Diagonalize02a
028 10.0 points

Find a matrix P and d_2, d_3 so that

$$P \begin{bmatrix} 5 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{bmatrix} P^{-1}, \quad d_1 \geq d_2 \geq d_3,$$

is a diagonalization of the matrix

$$A = \begin{bmatrix} 5 & -9 & -9 \\ 0 & 3 & 0 \\ 0 & -1 & 2 \end{bmatrix}.$$

1. $d_2 = -2, d_3 = -3,$

$$P = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$$

2. $d_2 = 3, d_3 = 2,$

$$P = \begin{bmatrix} 3 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & -1 & 0 \end{bmatrix}$$

3. $d_2 = -2, d_3 = -3,$

$$P = \begin{bmatrix} 3 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & -1 & 0 \end{bmatrix}$$

4. $d_2 = 3, d_3 = 2,$

$$P = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$$

5. $d_2 = 3, d_3 = 2,$

$$P = \begin{bmatrix} 1 & 3 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

6. $d_2 = -2, d_3 = -3,$

$$P = \begin{bmatrix} 1 & 3 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

CalC13c03a
029 10.0 points

Which of the following statements are true for all vectors \mathbf{a} , \mathbf{b} ?

- A. $|\mathbf{a} \cdot \mathbf{b}| \leq \|\mathbf{a}\| \|\mathbf{b}\|$,
 - B. $|\mathbf{a} \cdot \mathbf{b}| = \|\mathbf{a}\| \|\mathbf{b}\|$, $\mathbf{a} \neq 0$, $\mathbf{b} \neq 0 \implies \mathbf{a} \perp \mathbf{b}$,
 - C. $\|\mathbf{a} - \mathbf{b}\|^2 = \|\mathbf{a}\|^2 + 2\mathbf{a} \cdot \mathbf{b} + \|\mathbf{b}\|^2$.
1. none of them
 2. B and C only
 3. A and B only
 4. A only
 5. C only
 6. all of them
 7. A and C only
 8. B only

OrthoBasis01b
030 10.0 points

Determine c_2 so that

$$\mathbf{y} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + c_3 \mathbf{u}_3$$

when

$$\mathbf{y} = \begin{bmatrix} -4 \\ -2 \\ 0 \end{bmatrix}$$

and

$$\mathbf{u}_1 = \begin{bmatrix} 0 \\ 4 \\ -4 \end{bmatrix}, \quad \mathbf{u}_2 = \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix}, \quad \mathbf{u}_3 = \begin{bmatrix} -4 \\ 3 \\ 3 \end{bmatrix}.$$

1. $c_2 = \frac{16}{17}$
2. No value of c_2 exists.
3. $c_2 = 0$
4. $c_2 = -\frac{16}{17}$
5. $c_2 = -\frac{8}{17}$
6. $c_2 = \frac{8}{17}$

6. $\text{dist} = 2\sqrt{5}$

DistanceMC01
031 10.0 points

Find the distance from \mathbf{y} to the plane in \mathbb{R}^3 spanned by \mathbf{u}_1 and \mathbf{u}_2 when

$$\mathbf{y} = \begin{bmatrix} 5 \\ -9 \\ 5 \end{bmatrix}, \quad \mathbf{u}_1 = \begin{bmatrix} -3 \\ -5 \\ 1 \end{bmatrix}, \quad \mathbf{u}_2 = \begin{bmatrix} -3 \\ 2 \\ 1 \end{bmatrix}.$$

1. $\text{dist} = 8$

2. $\text{dist} = 6$

3. $\text{dist} = 4$

4. $\text{dist} = \sqrt{6}$

5. $\text{dist} = 2\sqrt{10}$

GramSchmidt01a
032 10.0 points

Use the fact that

$$A = \begin{bmatrix} 1 & -4 & 9 & -7 \\ -1 & 2 & -4 & 1 \\ 5 & -6 & 10 & 7 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 0 & -1 & 5 \\ 0 & -2 & 5 & -6 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

to determine an orthogonal basis for $\text{Col}(A)$.

1. $\begin{bmatrix} -4 \\ 2 \\ -6 \end{bmatrix}, \quad \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix}$

2. $\begin{bmatrix} -4 \\ 2 \\ -6 \end{bmatrix}, \quad \begin{bmatrix} 1 \\ -1 \\ 5 \end{bmatrix}$

3. $\begin{bmatrix} 1 \\ -1 \\ 5 \end{bmatrix}, \quad \begin{bmatrix} -4 \\ 1 \\ 1 \end{bmatrix}$

4. $\begin{bmatrix} 1 \\ -1 \\ 5 \end{bmatrix}, \quad \begin{bmatrix} 1 \\ -4 \\ -1 \end{bmatrix}$

LeastSquares02c

033 10.0 points

Find the least-squares solution of $A\mathbf{x} = \mathbf{b}$ when

$$A = \begin{bmatrix} 1 & -2 \\ 0 & -1 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} -1 \\ 2 \\ 7 \end{bmatrix}.$$

1. $\begin{bmatrix} 6 \\ 1 \end{bmatrix}$

2. $\begin{bmatrix} -1 \\ 7 \end{bmatrix}$

3. $\frac{1}{2} \begin{bmatrix} 8 \\ 5 \end{bmatrix}$

4. $\begin{bmatrix} -1 \\ 2 \end{bmatrix}$

5. $\frac{1}{2} \begin{bmatrix} 22 \\ 19 \end{bmatrix}$

RegressionLine03c**034 10.0 points**

Find the Least Squares Regression line $y = mx + b$ that best fits the data points

$$(-1, 1), \quad (0, -2), \quad (1, 3), \quad (2, 2).$$

OrthogDiag02a**035 10.0 points**

When

$$A = \begin{bmatrix} 1 & 8 \\ 8 & -11 \end{bmatrix}$$

find matrices D and P in an orthogonal diagonalization of A given that $\lambda_1 > \lambda_2$.

