

Universal Calculus Derivations for the Millennium Prize Problems

Formal Review Manuscript

Nicholas Kleban

June 8, 2026

Abstract. This manuscript defines Universal Calculus as a formal branched equalizer calculus and applies it to the seven Millennium Prize Problems. The primitive operator is the Kleban Operator $K^\beta = \delta_A \circ R^\beta$; the exact sector is $M_0^\beta = \ker K^\beta = \text{Eq}(R_0^\beta, R_1^\beta, R_2^\beta)$. For each Clay problem P , the native formulation is represented by a problem frame $(\Lambda_P, Y_P, R_{P,0}, R_{P,1}, R_{P,2}, \text{Adm}_P, \pi_P, T_P, Q_P)$, with exact admissible sector $M_{P,0} = \text{Adm}_P \cap \ker K_P$. Each part gives the native-to-Universal Calculus correspondence, the inverse extraction map, and the resulting certificate condition. Poincare is treated as solved natively by the Ricci-flow/geometrization route. For the six problems listed by the Clay Mathematics Institute as unsolved, the derivations identify the certificate whose native construction is required for Clay-level validity. CMI states that before it considers a proposed solution, the solution must be published in a qualifying outlet, at least two years must have passed, and the solution must have received general acceptance in the global mathematics community [4].

Contents

I	Universal Calculus	3
1	Formal substrate	3
2	Problem frames	4
3	Native-to-UC bijection condition	4
4	Certificate types	5
II	Birch and Swinnerton-Dyer	5
5	Native target	5
6	UC frame	5
7	Bijection derivation	6
8	Unique UC solution form	6
III	Hodge Conjecture	6
9	Native target	6
10	UC frame	7
11	Bijection derivation	7
12	Unique UC solution form	7
IV	Navier–Stokes Existence and Smoothness	8
13	Native target	8
14	UC frame	8
15	Bijection derivation	9
16	Unique UC solution form	9
V	P versus NP	10
17	Native target	10

18 UC frame	10
19 Bijection derivation	10
20 Unique UC solution form	11
VI Poincare Conjecture	11
21 Native target	11
22 UC frame	11
23 Bijection derivation	12
24 Unique UC solution form	12
VII Riemann Hypothesis	12
25 Native target	12
26 UC frame	13
27 Bijection derivation	13
28 Unique UC solution form	14
VIII Yang–Mills Existence and Mass Gap	14
29 Native target	14
30 UC frame	14
31 Bijection derivation	15
32 Unique UC solution form	15
Criteria for Clay-Level Validity	15

Part I

Universal Calculus

1 Formal substrate

Let $A = \mathbb{Z}_3 = \{0, 1, 2\}$ with cyclic convention. Let $V = \mathbb{R}^3$ or a problem-specific additive equality carrier. For each chart index n , let S_n be a raw state set and let T_n be an admissible family of transformations. For tolerance $\varepsilon \geq 0$, the stable comparison selector is

$$\text{Stab}_\varepsilon(S_n, T_n) = \{(a, b, \delta, \chi) : |\delta(a, b; \chi) - \delta(Ta, Tb; T\chi)| \leq \varepsilon \ \forall T \in T_n\}.$$

The latent comparison space is the structured quotient

$$\Lambda_n = \langle \text{Stab}_\varepsilon(S_n, T_n) \rangle / \sim.$$

The global phase is a single carrier $\theta \in \mathbb{R} \rightarrow S^1$. The branch set is

$$B = \{E, P, H\},$$

with elliptic fixed-radius, polar parent, and hyperbolic outward branches as defined in Universal Calculus [1].

For each axis $i \in A$, let $a_i : \Lambda_n \rightarrow \mathbb{R}$, $r_i : \Lambda_n \rightarrow \mathbb{R}_{\geq 0}$, and $\alpha_i : \Lambda_n \rightarrow \mathbb{R}$ be coordinate fields, and let $\beta : \Lambda_n \rightarrow B^A$ be the branch field. With cyclic insertion $J_i : \mathbb{R}^2 \rightarrow V$ and branch trace Γ_b , the axis reconstruction map is

$$R_i^\beta(\lambda, \theta) = a_i(\lambda)e_i + J_i\Gamma_{\beta_i(\lambda)}(\rho_{i,\beta_i}(\lambda), \theta + \alpha_i(\lambda), \sigma_i(\lambda)).$$

The reconstruction triple is

$$R^\beta = (R_0^\beta, R_1^\beta, R_2^\beta) : \Lambda_n \times \Theta \rightarrow V^A.$$

The axis coboundary is

$$\delta_A(R_0, R_1, R_2) = (R_0 - R_1, R_1 - R_2, R_2 - R_0).$$

The Kleban Operator is

$$K^\beta = \delta_A \circ R^\beta, \quad K^\beta(\lambda, \theta) = (R_0^\beta - R_1^\beta, R_1^\beta - R_2^\beta, R_2^\beta - R_0^\beta).$$

The exact and finite-tolerance sectors are

$$M_0^\beta = \ker K^\beta = \text{Eq}(R_0^\beta, R_1^\beta, R_2^\beta), \quad M_\varepsilon^\beta = \{\lambda : \|K^\beta(\lambda, \theta)\| \leq \varepsilon\}.$$

The equalizer identity follows by direct expansion:

$$K^\beta(\lambda, \theta) = 0 \iff R_0^\beta(\lambda, \theta) = R_1^\beta(\lambda, \theta) = R_2^\beta(\lambda, \theta).$$

When a differentiable approximation exists, the Kleban differential is DK^β . The normal defect metric is

$$g_N^\beta = (DK^\beta)^*\Xi(DK^\beta),$$

where Ξ is a bilinear contrast on the defect target. When a target contrast h is supplied, the tangential realspace metric is

$$g_T^\beta = (D\pi^\beta)^* h (D\pi^\beta).$$

The relative curvature is the axis coboundary of the gluing curvature,

$$\Omega^\beta = \delta_A d(d\alpha + \omega),$$

and the stationary-history condition is written $\delta S = 0$, with action dominated by squared Kleban defect plus regularity, boundary, and branch-transition costs [1].

2 Problem frames

A Clay problem frame in Universal Calculus is a tuple

$$P = (\Lambda_P, Y_P, R_{P,0}, R_{P,1}, R_{P,2}, \text{Adm}_P, \pi_P, T_P, Q_P),$$

where Λ_P is the comparison domain, Y_P is the equality carrier, $R_{P,i} : \Lambda_P \rightarrow Y_P$ are three reconstruction sectors, $\text{Adm}_P \subseteq \Lambda_P$ is the admissible set, π_P is the terminal projection, T_P is the native target class, and Q_P is the terminal predicate. The problem operator and exact admissible sector are

$$K_P = \delta_A(R_{P,0}, R_{P,1}, R_{P,2}), \quad M_{P,0} = \text{Adm}_P \cap \ker K_P.$$

The Universal Calculus resolution predicate is

$$Q_P(M_{P,0}) = Q_P(\text{Adm}_P \cap \ker K_P).$$

This is a formal equalizer representation. It is Clay-level only when the map between native mathematical data and the Universal Calculus frame is a bijection on the stated equivalence classes and preserves the terminal predicate.

3 Native-to-UC bijection condition

For each problem P , define $\mathcal{N}_P$ as the set of native objects, hypotheses, proof data, and equivalence relations appearing in the official mathematical formulation of the problem. Define $\mathcal{U}_P$ as the set of Universal Calculus data satisfying the corresponding frame syntax. A bijection is a pair

$$\Phi_P : \mathcal{N}_P \rightarrow \mathcal{U}_P, \quad \Psi_P : \mathcal{U}_P \rightarrow \mathcal{N}_P$$

such that

$$\Psi_P \circ \Phi_P = \text{id}_{\mathcal{N}_P}, \quad \Phi_P \circ \Psi_P = \text{id}_{\mathcal{U}_P}$$

on admissible equivalence classes, and

$$K_P(\Phi_P(N)) = 0 \iff N \text{ satisfies the native equations and compatibility laws,}$$

$$Q_P(\Phi_P(N)) \iff N \text{ satisfies the Clay terminal statement.}$$

The phrase “unique solution form” means the certificate form forced by these maps after the native target and equivalence relation are fixed. It does not mean uniqueness of possible proof strategies.

4 Certificate types

The certificate types used in the seven parts are:

$$\begin{aligned} \text{section: } & \sigma : T_P \rightarrow \text{Adm}_P, & \pi_P \sigma &= \text{id}_{T_P}, & K_P \sigma &= 0, \\ \text{separator: } & I : \text{Adm}_P \rightarrow W, & \lambda \in M_{P,0} &\Rightarrow I(\lambda) \in C, & \lambda \in B &\Rightarrow I(\lambda) \notin C, \\ \text{witness: } & \lambda_* \in \text{Adm}_P, & K_P(\lambda_*) &= 0, & m(\lambda_*) &\geq \Delta > 0. \end{aligned}$$

The derivation is immediate:

$$\pi_P \sigma = \text{id}_{T_P}, \quad K_P \sigma = 0 \Rightarrow \pi_P(M_{P,0}) = T_P.$$

If $\lambda \in M_{P,0} \cap B$, a separator gives $I(\lambda) \in C$ and $I(\lambda) \notin C$, hence $M_{P,0} \cap B = \emptyset$. If a witness satisfies $K_P(\lambda_*) = 0$ and $m(\lambda_*) \geq \Delta > 0$, then the exact sector contains an admissible object with positive terminal margin.

Part II

Birch and Swinnerton-Dyer

5 Native target

The Clay problem concerns the relation between rational points on an elliptic curve or abelian variety and the behavior near $s = 1$ of the associated L -function. CMI describes the conjecture as relating the size of the group of rational points to the behavior of an associated zeta function near $s = 1$ [5]; Wiles' official problem description is the Clay reference for the precise formulation [6]. For an elliptic curve $E/\mathbb{Q}$, the rank statement is

$$\text{rank } E(\mathbb{Q}) = \text{ord}_{s=1} L(E, s).$$

The refined statement is the leading-term identity

$$L^*(E, 1) = \frac{\Omega_E \text{Reg}_E |\text{Sha}(E)| \prod_p c_p}{|E(\mathbb{Q})_{\text{tors}}|^2},$$

with period, regulator, Tate-Shafarevich group, Tamagawa factors, and torsion interpreted in the standard arithmetic-geometric sense.

6 UC frame

Let

$$\Lambda_{BSD} = \{(E, r_{\text{alg}}, r_{\text{an}}, L^*, A, D_{\text{loc}}, D_{\text{fun}}, D_{\text{Sha}}) : E/\mathbb{Q} \text{ elliptic curve with comparison data}\}.$$

Set

$$\begin{aligned} r_{\text{alg}}(E) &= \text{rank } E(\mathbb{Q}), & r_{\text{an}}(E) &= \text{ord}_{s=1} L(E, s), \\ A(E) &= \frac{\Omega_E \text{Reg}_E |\text{Sha}(E)| \prod_p c_p}{|E(\mathbb{Q})_{\text{tors}}|^2}. \end{aligned}$$

Use equality carrier

$$Y_{BSD} = \mathbb{Z} \oplus \mathbb{C} \oplus Y_{loc} \oplus Y_{fun} \oplus Y_{Sha}.$$

Define

$$R_0(\lambda) = (r_{alg}, L^*, D_{loc}, D_{fun}, D_{Sha}), \quad R_1(\lambda) = (r_{an}, A, 0, 0, 0), \quad R_2(\lambda) = R_1(\lambda).$$

Then

$$K_{BSD}(\lambda) = 0 \iff r_{alg} = r_{an}, \quad L^* = A, \quad D_{loc} = D_{fun} = D_{Sha} = 0.$$

Thus $\ker K_{BSD}$ is the Universal Calculus equalizer encoding of the BSD rank equality, refined leading coefficient formula, local compatibility, functorial compatibility, and Tate-Shafarevich admissibility.

7 Bijection derivation

Let $\mathcal{N}_{BSD}$ consist of native BSD proof data

$$(E, \mathcal{D}_E),$$

where $\mathcal{D}_E$ proves the rank equality, refined leading-term identity, local factor compatibility, and finiteness conditions required in the chosen BSD formulation. Let $\mathcal{U}_{BSD}$ consist of $\lambda \in \text{Adm}_{BSD}$ with $\pi(\lambda) = E$ and $K_{BSD}(\lambda) = 0$.

Define $\Phi_{BSD}(E, \mathcal{D}_E)$ by inserting the arithmetic and analytic quantities certified by $\mathcal{D}_E$ into λ . Define $\Psi_{BSD}(\lambda)$ by reading the equalities forced by $K_{BSD}(\lambda) = 0$. Then

$$\Psi_{BSD}\Phi_{BSD}(E, \mathcal{D}_E) = (E, \mathcal{D}_E)$$

provided proof annotations are identified up to the native equivalence relation, and

$$\Phi_{BSD}\Psi_{BSD}(\lambda) = \lambda$$

provided λ contains no data beyond the listed BSD equality carrier. The terminal equivalence is

$$\pi(\text{Adm}_{BSD} \cap \ker K_{BSD}) = \{E/\mathbb{Q} \text{ in the target class}\} \iff \text{BSD holds for all } E \text{ in that class.}$$

8 Unique UC solution form

The forced certificate is a section

$$\Sigma_{BSD} : T_{BSD} \rightarrow \text{Adm}_{BSD}, \quad E \mapsto \Sigma_{BSD}(E),$$

with

$$\pi\Sigma_{BSD}(E) = E, \quad K_{BSD}\Sigma_{BSD}(E) = 0 \quad \forall E \in T_{BSD}.$$

The Clay-level construction required by this certificate is a native proof of the rank equality, the refined leading-term identity, and the relevant finiteness and compatibility conditions for every admissible curve or abelian variety in the target formulation.

Part III

Hodge Conjecture

9 Native target

CMI states that the Hodge conjecture asserts that, for projective algebraic varieties, Hodge cycles are rational linear combinations of algebraic cycles [8]; Deligne's official problem description supplies

the Clay reference formulation [9]. For a smooth projective complex variety X and codimension p , the standard formulation is

$$\text{Hdg}^p(X) = H^{2p}(X, \mathbb{Q}) \cap H^{p,p}(X),$$

with cycle class map

$$\text{cl}_X : \text{CH}^p(X)_{\mathbb{Q}} \rightarrow H^{2p}(X, \mathbb{Q}).$$

The target statement is

$$\text{Hdg}^p(X) \subseteq \text{im}(\text{cl}_X) \quad \text{for all } X, p.$$

10 UC frame

Let

$$\Lambda_H = \{(X, p, \gamma, Z) : X \text{ smooth projective over } \mathbb{C}, \gamma \in H^{2p}(X, \mathbb{Q}), Z \in \text{CH}^p(X)_{\mathbb{Q}}\}.$$

Let $\Pi^{p,p}$ denote the Hodge-type projection. Define

$$R_0(X, p, \gamma, Z) = \gamma, \quad R_1(X, p, \gamma, Z) = \Pi^{p,p}\gamma, \quad R_2(X, p, \gamma, Z) = \text{cl}_X(Z).$$

The equality carrier is $H^{2p}(X, \mathbb{Q})$ after embedding the three terms in a common rational cohomology target. Then

$$K_H(\lambda) = 0 \iff \gamma = \Pi^{p,p}\gamma = \text{cl}_X(Z).$$

Therefore $K_H = 0$ is equivalent to: γ is a rational Hodge class and is represented by an algebraic cycle.

11 Bijection derivation

Let $\mathcal{N}_H$ consist of native data $(X, p, \gamma, Z, \mathcal{D})$ such that $\gamma \in \text{Hdg}^p(X)$ and $\mathcal{D}$ verifies $\text{cl}_X(Z) = \gamma$. Let $\mathcal{U}_H$ consist of $\lambda = (X, p, \gamma, Z) \in \text{Adm}_H$ satisfying $K_H(\lambda) = 0$.

The map Φ_H sends $(X, p, \gamma, Z, \mathcal{D})$ to (X, p, γ, Z) . The map Ψ_H sends $\lambda \in \mathcal{U}_H$ to the native assertions

$$\gamma \in H^{p,p}(X), \quad \gamma = \text{cl}_X(Z).$$

The inverse identities hold on equivalence classes of rational cycles modulo rational equivalence, because the only native object retained by the Universal Calculus frame is the cycle class $\text{cl}_X(Z)$ together with the chosen rational cycle representative. The terminal equivalence is

$$\forall \gamma \in \text{Hdg}^p(X) \exists Z \in \text{CH}^p(X)_{\mathbb{Q}} : K_H(X, p, \gamma, Z) = 0$$

if and only if the Hodge conjecture holds for (X, p) .

12 Unique UC solution form

The forced certificate is a cycle section

$$\Sigma_H(X, p) : \text{Hdg}^p(X) \rightarrow \text{CH}^p(X)_{\mathbb{Q}}$$

satisfying

$$\text{cl}_X(\Sigma_H(X, p)(\gamma)) = \gamma \quad \forall \gamma \in \text{Hdg}^p(X).$$

Equivalently, for

$$\lambda_\gamma = (X, p, \gamma, \Sigma_H(X, p)(\gamma)),$$

one has

$$K_H(\lambda_\gamma) = 0, \quad \pi_\gamma(\lambda_\gamma) = \gamma.$$

The Clay-level construction required by this certificate is a native algebraic-cycle construction for every rational Hodge class, with rational coefficient control and compatibility with the equivalence relation in $\text{CH}^p(X)_\mathbb{Q}$.

Part IV

Navier–Stokes Existence and Smoothness

13 Native target

CMI states that the Navier–Stokes problem asks for proof of existence and uniqueness for the equations governing fluids, or a qualifying counterexample in the official formulation [11]; Fefferman’s official problem description supplies the Clay mathematical statement [12]. In the standard incompressible setting on $\Omega = \mathbb{R}^3$ or $\mathbb{T}^3$,

$$\partial_t u + (u \cdot \nabla)u - \nu \Delta u + \nabla p = f, \quad \nabla \cdot u = 0, \quad \nu > 0,$$

with smooth divergence-free initial datum

$$u(0) = u_0.$$

Here ν denotes viscosity.

14 UC frame

Let

$$\Lambda_{NS} = \{(u_0, u, p, T) : u_0 \in C^\infty, \nabla \cdot u_0 = 0, (u, p) \text{ candidate history on } [0, T]\}.$$

Define the native defect vector

$$D_{NS}(u, p; u_0) = (\partial_t u + (u \cdot \nabla)u - \nu \Delta u + \nabla p - f, \nabla \cdot u, D_0(u), D_{bdry}),$$

where $D_0(u)$ denotes $u(0) - u_0$ and D_{bdry} denotes the required decay, periodic, or boundary condition. Set

$$R_0 = D_{NS}, \quad R_1 = 0, \quad R_2 = 0.$$

Then

$$K_{NS}(u, p; u_0) = 0 \iff D_{NS}(u, p; u_0) = 0.$$

For s above the classical smoothness threshold, define the regularity monitor

$$\mathcal{R}_s(u; T) = \sup_{0 \leq t \leq T} \|u(t)\|_{H^s}.$$

15 Bijection derivation

Let $\mathcal{N}_{NS}^+$ consist of native global smooth solution data $(u_0, u, p, \mathcal{D})$ with uniqueness proof and $\mathcal{R}_s(u; T) < \infty$ for every finite T . Let $\mathcal{U}_{NS}^+$ consist of $\lambda \in \text{Adm}_{NS}$ satisfying $K_{NS}(\lambda) = 0$, global H^s bounds, and uniqueness in the exact sector.

The map Φ_{NS}^+ sends a native solution to its Universal Calculus history. The map Ψ_{NS}^+ reads from $K_{NS} = 0$ the PDE, incompressibility, initial condition, boundary condition, and global regularity bound. Thus

$$\mathcal{N}_{NS}^+ \cong \mathcal{U}_{NS}^+.$$

Let $\mathcal{N}_{NS}^-$ consist of native counterexample data $(u_0, u, p, T_*, \mathcal{D})$ satisfying the exact equations up to $T_* < \infty$ and finite-time loss of smoothness. Let $\mathcal{U}_{NS}^-$ consist of exact witnesses with

$$K_{NS}(u, p; u_0) = 0, \quad \limsup_{t \rightarrow T_*^-} \|u(t)\|_{H^s} = \infty.$$

Then

$$\mathcal{N}_{NS}^- \cong \mathcal{U}_{NS}^-.$$

The Clay terminal predicate is the disjunction

$$Q_{NS} = Q_{NS}^+ \vee Q_{NS}^-.$$

The two branches correspond to the two kinds of acceptable Clay resolution in the official problem statement.

16 Unique UC solution form

The forced positive certificate is

$$C_{NS}^+ = (\Sigma_{NS}, U_{NS}, B_{NS}),$$

where

$$\Sigma_{NS} : u_0 \mapsto (u, p), \quad K_{NS}(\Sigma_{NS}(u_0)) = 0, \quad \mathcal{R}_s(u; T) < \infty \quad \forall T < \infty.$$

Here U_{NS} is a uniqueness separator and B_{NS} is a continuation barrier:

$$\mathcal{R}_s(u; T) < \infty \Rightarrow \text{the exact history extends past } T.$$

The forced negative certificate is a witness

$$C_{NS}^- = (u_0, u, p, T_*),$$

with

$$D_0(u) = 0, \quad K_{NS}(u, p; u_0) = 0, \quad T_* < \infty, \quad \limsup_{t \rightarrow T_*^-} \|u(t)\|_{H^s} = \infty.$$

A completed Clay resolution selects one of C_{NS}^+ or C_{NS}^- . The native construction required is either a global regularity and uniqueness proof or an exact finite-time singular witness satisfying the official

hypotheses.

Part V

P versus NP

17 Native target

CMI states the P versus NP question as whether every problem whose solution can be checked efficiently can also be solved efficiently [14]. Cook's official problem description supplies the Clay reference formulation [15]; Cook, Levin, and Karp supply the standard NP-completeness foundations [16, 17, 18]. Let $L \subseteq \Sigma^*$ be an NP language with polynomial verifier $V(x, w)$ and witness bound $p(|x|)$. The native condition is

$$x \in L \iff \exists w, \quad |w| \leq p(|x|), \quad V(x, w) = 1.$$

The question is whether there exists a deterministic polynomial-time decider for every such L .

18 UC frame

Let

$$\Lambda_{P/NP} = \{(L, V, p, A, x) : L \in NP, \text{ } V \text{ verifier, } A \text{ deterministic decider candidate, } x \in \Sigma^*\}.$$

Define Boolean-valued sectors

$$\begin{aligned} R_0(L, x) &= \chi_L(x), \\ R_1(V, p, x) &= \bigvee_{|w| \leq p(|x|)} V(x, w), \\ R_2(A, x) &= A(x). \end{aligned}$$

The equality carrier is $\{0, 1\}^{\Sigma^*}$, and

$$K_{P/NP}(L, V, p, A) = 0 \iff \chi_L(x) = \bigvee_{|w| \leq p(|x|)} V(x, w) = A(x) \quad \forall x.$$

The verifier equality $R_0 = R_1$ is the native NP definition. The equality $R_1 = R_2$ is exact polynomial-time decision when A is in the polynomial-time admissible class.

19 Bijection derivation

Let $\mathcal{N}_{P/NP}^=$ consist of native data assigning to every verifier pair (V, p) a deterministic polynomial-time decider $A_{V,p}$ deciding the same language. Let $\mathcal{U}_{P/NP}^=$ consist of sections

$$\Sigma_{P/NP} : (L, V, p) \mapsto A_{L,V,p}$$

such that $A_{L,V,p} \in P$ and $K_{P/NP}(L, V, p, A_{L,V,p}) = 0$ on all inputs. Direct expansion of $K_{P/NP} = 0$ gives

$$\mathcal{N}_{P/NP}^= \cong \mathcal{U}_{P/NP}^=.$$

Thus this branch is equivalent to $P = NP$.

Let $\mathcal{N}_{P/NP}^\#$ consist of a native lower-bound proof for a language $L_* \in NP$ proving that no deterministic polynomial-time decider decides L_* . Let $\mathcal{U}_{P/NP}^\#$ consist of L_* and a separator

$$I_{P/NP} : \text{Adm}_{P/NP} \rightarrow W, \quad C \subseteq W,$$

with

$$\begin{aligned} \lambda \in M_{P/NP,0}(L_*) &\Rightarrow I_{P/NP}(\lambda) \in C, \\ A \in P \text{ as a candidate for } L_* &\Rightarrow I_{P/NP}(A) \notin C. \end{aligned}$$

Then an exact polynomial-time decider would be both in and outside C , a contradiction. Hence

$$\mathcal{N}_{P/NP}^\# \cong \mathcal{U}_{P/NP}^\#$$

provided $I_{P/NP}$ is a native lower-bound invariant and is not invalidated by standard barrier phenomena such as relativization, natural proofs, or algebrization [19, 20, 21].

20 Unique UC solution form

The forced equality certificate is

$$\Sigma_{P/NP} : (L, V, p) \mapsto A_{L,V,p}, \quad A_{L,V,p} \in P, \quad K_{P/NP}(L, V, p, A_{L,V,p}) = 0 \quad \forall x.$$

The forced separation certificate is

$$(L_*, I_{P/NP}, C),$$

with the separator equations above. A completed Clay resolution constructs exactly one of these two mutually exclusive certificate branches in native complexity theory.

Part VI

Poincare Conjecture

21 Native target

CMI lists the Poincare conjecture as solved and describes the question as whether the three-dimensional sphere is characterized as the unique simply connected three-manifold; CMI also states that Perelman's proof gives geometrization of three-manifolds [22]. The native proof route is Hamilton's Ricci flow with Perelman's monotonicity, surgery, and geometrization analysis [24, 25, 26, 27, 28]. The terminal native statement is

$$M^3 \text{ closed and simply connected} \implies M^3 \cong S^3.$$

22 UC frame

Let

$$\begin{aligned} \Lambda_{Pc} = \{ & (M, g(t), S, \Pi) : M \text{ closed 3-manifold,} \\ & g(t) \text{ Ricci-flow candidate,} \\ & S \text{ surgery data,} \\ & \Pi \text{ terminal geometry classifier} \}. \end{aligned}$$

For the Poincare sector, impose $\pi_1(M) = 0$. Define

$$D_{Pc} = (D_{flow}, D_{surgery}, D_{terminal}), \quad D_{flow} = \partial_t g + 2 \text{Ric}(g).$$

Set

$$R_0 = D_{Pc}, \quad R_1 = 0, \quad R_2 = 0.$$

Then

$$K_{Pc} = 0 \iff \text{the Ricci-flow, surgery, and terminal-geometrization conditions are satisfied.}$$

23 Bijection derivation

Let $\mathcal{N}_{Pc}$ consist of native Hamilton–Perelman proof data: Ricci-flow history, monotonicity control, surgery admissibility, canonical neighborhood control, extinction or decomposition analysis, and terminal geometrization. Let $\mathcal{U}_{Pc}$ consist of exact Universal Calculus data $(M, g(t), S, \Pi)$ with $K_{Pc} = 0$ and $\Pi(M) = S^3$ in the simply connected closed sector.

The map Φ_{Pc} sends the native Ricci-flow/geometrization proof data to the corresponding Universal Calculus defect components. The map Ψ_{Pc} extracts from $K_{Pc} = 0$ the native flow equation, surgery admissibility, monotone control, and terminal classifier. The inverse identities hold after quotienting by the usual equivalences of Ricci-flow parametrization, surgery choices, and 3-manifold classification data. Therefore

$$\mathcal{N}_{Pc} \cong \mathcal{U}_{Pc}.$$

The terminal equivalence is

$$K_{Pc} = 0, \pi_1(M) = 0, M \text{ closed} \implies \Pi(M) = S^3 \iff M \cong S^3.$$

24 Unique UC solution form

The forced certificate is

$$C_{Pc} = (H_{RF}, I_{mono}, S_{surg}, \Pi_{geom}),$$

where H_{RF} is the Ricci-flow history, I_{mono} is monotone control excluding inadmissible singular behavior, S_{surg} is surgery data, and Π_{geom} is the terminal geometrization projection. The certificate equations are

$$\begin{aligned} K_{Pc}(H_{RF}, S_{surg}, \Pi_{geom}) &= 0, \\ M^3 \text{ closed, } \pi_1(M) = 0 &\Rightarrow \Pi_{geom}(M) = S^3. \end{aligned}$$

This is the one case in the manuscript where the native certificate is already supplied by accepted mathematics.

Part VII

Riemann Hypothesis

25 Native target

CMI states the Riemann Hypothesis as the assertion that all non-obvious zeros of the zeta function have real part $1/2$ [29]; Bombieri’s official problem description is the Clay reference formulation [30].

With

$$\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s), \quad \xi(s) = \xi(1-s),$$

the native target is

$$\zeta(s) = 0, \quad 0 < \operatorname{Re}(s) < 1 \implies \operatorname{Re}(s) = \frac{1}{2}.$$

26 UC frame

Let $s = 1/2 + \tau + it$. Let F_0, F_1, F_2 be three reconstructions of the same completed zeta object: additive analytic reconstruction, prime/Euler reconstruction after regularization, and functional-reflection reconstruction. Embed $\mathbb{C}$ in an additive equality carrier by

$$\Phi_\mu(w) = \left(\operatorname{Re} w, \Im w, \log \left(1 + \frac{|w|^2}{\mu^2} \right) \right).$$

Define

$$Z_i(s) = \Phi_\mu(F_i(s)), \quad R_i = Z_i.$$

Then

$$K_{RH} = 0 \iff Z_0(s) = Z_1(s) = Z_2(s).$$

The exact zero fiber is

$$\mathcal{Z}_\zeta = \{\lambda \in \operatorname{Adm}_{RH} \cap \ker K_{RH} : F_0(\lambda) = F_1(\lambda) = F_2(\lambda) = 0\}.$$

The bad sector is

$$B_{RH} = \{\lambda \in \mathcal{Z}_\zeta : \operatorname{Re} s(\lambda) \neq 1/2\}.$$

27 Bijection derivation

Let $\mathcal{N}_{RH}$ consist of native analytic zero data for $\xi(s)$, including analytic continuation, functional equation, Euler product where valid, and growth conditions. Let $\mathcal{U}_{RH}$ consist of admissible zeta reconstruction data satisfying $K_{RH} = 0$ and the same analytic constraints.

The map Φ_{RH} sends native zero data to the triple (F_0, F_1, F_2) and then to (Z_0, Z_1, Z_2) . The map Ψ_{RH} extracts from $K_{RH} = 0$ the equality of the three reconstructions and reads the common value as the completed zeta value. The bijection condition is

$$\mathcal{N}_{RH} \cong \mathcal{U}_{RH}$$

only if the reconstruction triple is complete: no zeros, poles, continuation data, or growth constraints are lost by the embedding Φ_μ .

Under this condition,

$$M_{RH,0} \cap B_{RH} = \emptyset \iff \text{there is no nontrivial zero off } \operatorname{Re}(s) = 1/2.$$

This is exactly the native Riemann Hypothesis.

28 Unique UC solution form

The forced certificate is a separator

$$I_{RH} : \text{Adm}_{RH} \rightarrow W_{RH}, \quad C_{line} \subseteq W_{RH},$$

with

$$\begin{aligned} \lambda \in \mathcal{Z}_\zeta \cap M_{RH,0} &\Rightarrow I_{RH}(\lambda) \in C_{line}, \\ \lambda \in B_{RH} &\Rightarrow I_{RH}(\lambda) \notin C_{line}. \end{aligned}$$

Then an off-line zero in the exact admissible sector would imply simultaneous membership and non-membership in C_{line} . The native construction required is a complete off-critical-line zero exclusion mechanism in analytic number theory.

Part VIII

Yang–Mills Existence and Mass Gap

29 Native target

CMI states that quantum Yang–Mills theory is mathematically foundationally unresolved and that the required mass gap has no proof [32]. Jaffe and Witten’s official problem description is the Clay reference formulation [33]. The problem asks, for each compact simple gauge group G , for a constructive quantum Yang–Mills theory on $\mathbb{R}^4$ satisfying the standard axioms of constructive quantum field theory and having positive mass gap

$$\Delta_G > 0.$$

Yang and Mills introduced the non-abelian gauge framework in 1954 [34]; the constructive formulation is conventionally expressed through Euclidean or Wightman/Osterwalder–Schrader type conditions [35, 36, 37].

30 UC frame

Let

$$\Lambda_{YM} = \{(G, \mathcal{A}, \mu, \mathcal{O}, H, \Delta) : \begin{aligned} &G \text{ compact simple,} \\ &\mathcal{A} \text{ gauge-field data,} \\ &\mu \text{ continuum measure,} \\ &\mathcal{O} \text{ observables,} \\ &H \text{ Hamiltonian, } \Delta > 0 \}. \end{aligned}$$

Define a defect vector

$$D_{YM} = (D_{gauge}, D_{OS}, D_{loc}, D_{ren}, D_{cont}, D_{nontriv}, D_{gap}),$$

where the entries encode gauge invariance, constructive quantum-field axioms, locality, renormalization consistency, continuum limit existence, nontriviality, and positive spectral gap. Let

$$D_{gap} = \inf(\text{Spec}(H) \setminus \{0\}) - \Delta.$$

Set

$$R_0 = D_{YM}, \quad R_1 = 0, \quad R_2 = 0.$$

Then

$$K_{YM}(\lambda) = 0 \iff D_{YM}(\lambda) = 0.$$

With $\Delta > 0$, this is the Universal Calculus encoding of constructive Yang–Mills existence and mass gap.

31 Bijection derivation

Let $\mathcal{N}_{YM}$ consist of native constructive Yang–Mills data for a compact simple group G : fields or Euclidean measures, gauge-invariant observables, reflection positivity or equivalent reconstruction data, locality, covariance, continuum limit, nontriviality, Hilbert-space reconstruction, Hamiltonian H , and spectral gap $\Delta_G > 0$. Let $\mathcal{U}_{YM}$ consist of $\lambda \in \text{Adm}_{YM}$ satisfying $K_{YM}(\lambda) = 0$ and $\Delta > 0$.

The map Φ_{YM} sends native constructive data to the defect vector D_{YM} and the Universal Calculus witness λ . The map Ψ_{YM} extracts from $K_{YM} = 0$ the vanishing of each constructive-QFT defect and reads $\inf(\text{Spec}(H) \setminus \{0\}) \geq \Delta > 0$. The bijection condition is

$$\mathcal{N}_{YM} \cong \mathcal{U}_{YM}$$

only if the regulator, continuum topology, gauge quotient, observable algebra, and Hilbert-space reconstruction are preserved without collapse to a trivial theory.

32 Unique UC solution form

The forced certificate is a witness family

$$\{(\lambda_G, \Delta_G) : G \text{ compact simple, } \lambda_G \in \text{Adm}_{YM}, \Delta_G > 0\}$$

with

$$K_{YM}(\lambda_G) = 0, \quad \inf(\text{Spec}(H_G) \setminus \{0\}) \geq \Delta_G > 0.$$

The native construction required is a rigorous four-dimensional continuum Yang–Mills theory satisfying constructive quantum-field criteria and a strictly positive mass gap for each compact simple gauge group in the Clay formulation.

Criteria for Clay-Level Validity

A Universal Calculus derivation is Clay-level only if the following conditions are verified in native mathematics.

1. **Faithful native domain.** The map Φ_P preserves all native hypotheses: quantifiers, regularity, boundary conditions, admissible equivalences, gauge quotients, coefficient fields, and limiting procedures.
2. **Bijection on equivalence classes.** The maps Φ_P and Ψ_P satisfy $\Psi_P \Phi_P = \text{id}$ and $\Phi_P \Psi_P = \text{id}$ after quotienting by the native equivalence relation. No native object may be lost, identified too coarsely, or duplicated by the Universal Calculus carrier.

3. **Kernel equivalence.** $K_P = 0$ must be proven equivalent to the native equations, compatibility laws, and admissibility constraints. It is insufficient for $K_P = 0$ to imply only a weaker formal identity.
4. **Terminal equivalence.** The terminal predicate Q_P must be exactly the Clay target predicate: rank/leading-term identity for BSD, algebraic-cycle generation for Hodge, global smoothness or exact blowup for Navier–Stokes, equality or separation for P versus NP, 3-sphere classification for Poincare, critical-line zero exclusion for RH, and constructive continuum theory with positive spectral gap for Yang–Mills.
5. **Constructed certificate.** The section, separator, or witness must be constructed by native mathematical proof. Naming Σ_P , I_P , or λ_* is not a construction.
6. **No circular admissibility.** The desired Clay conclusion must not be inserted into Adm_P , the equality carrier Y_P , the reconstruction maps $R_{P,i}$, or the terminal projection π_P .
7. **Limit stability.** If the derivation uses $M_{P,\varepsilon} \rightarrow M_{P,0}$, regulator removal, compactness, analytic continuation, weak-to-strong passage, or continuum reconstruction, the limiting operation must preserve the native terminal predicate.
8. **Problem-specific native obligations.**
 - a. BSD: global rank comparison, refined leading coefficient, local factors, and Tate-Shafarevich control.
 - b. Hodge: algebraic cycle construction for every rational Hodge class with rational coefficient control.
 - c. Navier–Stokes: critical regularity estimate and uniqueness, or an exact finite-time singular witness satisfying the official hypotheses.
 - d. P versus NP: a uniform polynomial-time solver for every NP verifier, or a lower-bound invariant proving separation and surviving known barriers.
 - e. Poincare: preservation of the accepted Hamilton–Perelman proof data under the Universal Calculus recoding.
 - f. RH: complete analytic zero exclusion off the critical line, preserving analytic continuation, functional equation, Euler data, and growth conditions.
 - g. Yang–Mills: constructive four-dimensional continuum existence, nontriviality, gauge-invariant observables, Hilbert-space reconstruction, and positive spectral gap.
9. **CMI procedural conditions.** CMI states that it does not accept direct submissions of proposed solutions and that, before CMI consideration, the solution must be published in a qualifying outlet, at least two years must have elapsed, and the solution must have received general acceptance in the global mathematics community [4].

References

- [1] Nicholas Kleban. *Universal Calculus: The Kleban Operator as Primitive Generator*. Uploaded source document, Kleban Operator Edition, June 8, 2026.
- [2] Nicholas Kleban. *Universal Calculus Certificates for the Millennium Prize Problems*. Uploaded source document, June 8, 2026.
- [3] Clay Mathematics Institute. The Millennium Prize Problems. Accessed June 8, 2026. <https://www.claymath.org/millennium-problems/>.
- [4] Clay Mathematics Institute. Rules for the Millennium Prize Problems. Revised rules adopted September 26, 2018; accessed June 8, 2026. <https://www.claymath.org/millennium-problems/rules/>.
- [5] Clay Mathematics Institute. Birch and Swinnerton-Dyer Conjecture. Accessed June 8, 2026. <https://www.claymath.org/millennium/birch-and-swinnerton-dyer-conjecture/>.
- [6] Andrew Wiles. The Birch and Swinnerton-Dyer Conjecture. Official Clay Mathematics Institute problem description, 2000; available through [5].
- [7] B. J. Birch and H. P. F. Swinnerton-Dyer. Notes on elliptic curves. II. *Journal für die reine und angewandte Mathematik*, 218:79–108, 1965.
- [8] Clay Mathematics Institute. Hodge Conjecture. Accessed June 8, 2026. <https://www.claymath.org/millennium/hodge-conjecture/>.
- [9] Pierre Deligne. The Hodge Conjecture. Official Clay Mathematics Institute problem description, 2000; available through [8].
- [10] W. V. D. Hodge. *The Topological Invariants of Algebraic Varieties*. Proceedings of the International Congress of Mathematicians, Cambridge, Massachusetts, 1950.
- [11] Clay Mathematics Institute. Navier–Stokes Equation. Accessed June 8, 2026. <https://www.claymath.org/millennium/navier-stokes-equation/>.
- [12] Charles L. Fefferman. Existence and Smoothness of the Navier–Stokes Equation. Official Clay Mathematics Institute problem description, 2000; available through [11].
- [13] Jean Leray. Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Mathematica*, 63:193–248, 1934.
- [14] Clay Mathematics Institute. P vs NP. Accessed June 8, 2026. <https://www.claymath.org/millennium/p-vs-np/>.
- [15] Stephen Cook. The P versus NP Problem. Official Clay Mathematics Institute problem description, 2000; available through [14].
- [16] Stephen A. Cook. The complexity of theorem-proving procedures. *Proceedings of the Third Annual ACM Symposium on Theory of Computing*, 151–158, 1971.
- [17] Leonid A. Levin. Universal sequential search problems. *Problems of Information Transmission*, 9(3):265–266, 1973.

- [18] Richard M. Karp. Reducibility among combinatorial problems. In R. E. Miller and J. W. Thatcher, editors, *Complexity of Computer Computations*, 85–103. Plenum, 1972.
- [19] Theodore Baker, John Gill, and Robert Solovay. Relativizations of the $P=?NP$ question. *SIAM Journal on Computing*, 4(4):431–442, 1975.
- [20] Alexander A. Razborov and Steven Rudich. Natural proofs. *Journal of Computer and System Sciences*, 55(1):24–35, 1997.
- [21] Scott Aaronson and Avi Wigderson. Algebrization: A new barrier in complexity theory. *ACM Transactions on Computation Theory*, 1(1):2, 2009.
- [22] Clay Mathematics Institute. Poincare Conjecture. Accessed June 8, 2026. <https://www.claymath.org/millennium/poincare-conjecture/>.
- [23] John Milnor. The Poincare Conjecture. Official Clay Mathematics Institute problem description, 2000; available through [22].
- [24] Richard S. Hamilton. Three-manifolds with positive Ricci curvature. *Journal of Differential Geometry*, 17(2):255–306, 1982.
- [25] Grigori Perelman. The entropy formula for the Ricci flow and its geometric applications. arXiv:math/0211159, 2002.
- [26] Grigori Perelman. Ricci flow with surgery on three-manifolds. arXiv:math/0303109, 2003.
- [27] Grigori Perelman. Finite extinction time for the solutions to the Ricci flow on certain three-manifolds. arXiv:math/0307245, 2003.
- [28] John Morgan and Gang Tian. *Ricci Flow and the Poincare Conjecture*. Clay Mathematics Monographs, vol. 3. American Mathematical Society and Clay Mathematics Institute, 2007.
- [29] Clay Mathematics Institute. Riemann Hypothesis. Accessed June 8, 2026. <https://www.claymath.org/millennium/riemann-hypothesis/>.
- [30] Enrico Bombieri. The Riemann Hypothesis. Official Clay Mathematics Institute problem description, 2000; available through [29].
- [31] Bernhard Riemann. Ueber die Anzahl der Primzahlen unter einer gegebenen Groesse. *Monatsberichte der Berliner Akademie*, 671–680, 1859.
- [32] Clay Mathematics Institute. Yang–Mills and the Mass Gap. Accessed June 8, 2026. <https://www.claymath.org/millennium/yang-mills-the-maths-gap/>.
- [33] Arthur Jaffe and Edward Witten. Quantum Yang–Mills Theory. Official Clay Mathematics Institute problem description, 2000; available through [32].
- [34] C. N. Yang and R. L. Mills. Conservation of isotopic spin and isotopic gauge invariance. *Physical Review*, 96(1):191–195, 1954.
- [35] Konrad Osterwalder and Robert Schrader. Axioms for Euclidean Green’s functions. *Communications in Mathematical Physics*, 31:83–112, 1973.
- [36] Konrad Osterwalder and Robert Schrader. Axioms for Euclidean Green’s functions II. *Communications in Mathematical Physics*, 42:281–305, 1975.

- [37] R. F. Streater and A. S. Wightman. *PCT, Spin and Statistics, and All That*. W. A. Benjamin, 1964.