

The Characteristic Equation

Geometric Algorithms

Lecture 19

Practice Problem

$$\begin{bmatrix} 5 & 2 & 3 & 0 \\ -1 & 2 & -3 & 1 \\ 2 & 4 & 10 & 0 \\ 1 & 2 & 3 & 5 \end{bmatrix}$$

Determine the dimension of the eigenspace of A for the eigenvalue 4.

(try not to do any row reductions)

Answer

$$\begin{bmatrix} 5 & 2 & 3 & 0 \\ -1 & 2 & -3 & 1 \\ 2 & 4 & 10 & 0 \\ 1 & 2 & 3 & 5 \end{bmatrix}$$

See A1 for answer

Objectives

1. Briefly recap eigenvalues and eigenvectors
2. Get a primer on determinants
3. Determine how to find eigenvalues (not just verify them)

Keyword

eigenvectors

eigenvalues

eigenspaces

eigenbases

determinant

characteristic equation


polynomial roots

triangular matrices

multiplicity

Recap


Recall: Eigenvalues/vectors



A *nonzero* vector \mathbf{v} in \mathbb{R}^n and real number λ are an **eigenvector** and **eigenvalue** for a $n \times n$ matrix A if

$$A\mathbf{v} = \lambda\mathbf{v}$$


Recall: Eigenvalues/vectors



A *nonzero* vector \mathbf{v} in \mathbb{R}^n and real number λ are an **eigenvector** and **eigenvalue** for a $n \times n$ matrix A if

$$A\mathbf{v} = \lambda\mathbf{v}$$

Recall: Eigenvalues/vectors



A *nonzero* vector \mathbf{v} in \mathbb{R}^n and real number λ are an **eigenvector** and **eigenvalue** for a $n \times n$ matrix A if

$$A\mathbf{v} = \lambda\mathbf{v}$$

\mathbf{v} is "just scaled" by A , not rotated

Recall: Verifying Eigenvectors

Recall: Verifying Eigenvectors

Question. Determine if \mathbf{v} is an eigenvector of A and determine the corresponding eigenvalues.

Recall: Verifying Eigenvectors

Question. Determine if \mathbf{v} is an eigenvector of A and determine the corresponding eigenvalues.

Solution. Easy. Work out the matrix–vector multiplication.

Recall: Verifying Eigenvectors

Question. Determine if \mathbf{v} is an eigenvector of A and determine the corresponding eigenvalues.

Solution. Easy. Work out the matrix–vector multiplication.

Example.

$$\begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} 6 \\ -5 \end{bmatrix} = \begin{bmatrix} -24 \\ 20 \end{bmatrix} = -4 \begin{bmatrix} 6 \\ -5 \end{bmatrix} \quad \checkmark$$

$$\begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 13 \\ 9 \end{bmatrix} \quad \times$$

Recall: Verifying Eigenvalues

Recall: Verifying Eigenvalues

Question. Find an eigenvector of A whose corresponding eigenvalue is λ .

Recall: Verifying Eigenvalues

Question. Find an eigenvector of A whose corresponding eigenvalue is λ .

Solution. Find a nontrivial solution to

$$(A - \lambda I)\mathbf{x} = \mathbf{0}$$

Recall: Verifying Eigenvalues

Question. Find an eigenvector of A whose corresponding eigenvalue is λ .

Solution. Find a nontrivial solution to

$$(A - \lambda I)\mathbf{x} = \mathbf{0}$$

*If we don't need the vector we can just show that $A - \lambda I$ is **not** invertible (by IMT).*

Recall: Finding Eigenspaces

Recall: Finding Eigenspaces

Question. Find a basis for the eigenspace of A corresponding to λ .

Recall: Finding Eigenspaces

Question. Find a basis for the eigenspace of A corresponding to λ .

Solution. Find a basis for $\text{Nul}(A - \lambda I)$.

Recall: Finding Eigenspaces

Question. Find a basis for the eigenspace of A corresponding to λ .

Solution. Find a basis for $\text{Nul}(A - \lambda I)$.

(we did this for our recap problem)

How do eigenvectors relate
to linear dynamical systems?

Recall: (Closed-Form) Solutions

Recall: (Closed-Form) Solutions

A (closed-form) solution of a linear dynamical system $\mathbf{v}_{i+1} = A\mathbf{v}_i$ is an expression for \mathbf{v}_k which is does **not** contain A^k or previously defined terms

Recall: (Closed-Form) Solutions

A **(closed-form) solution** of a linear dynamical system $\mathbf{v}_{i+1} = A\mathbf{v}_i$ is an expression for \mathbf{v}_k which is does **not** contain A^k or previously defined terms

In other word, it does not depend on A^k and is not **recursive**

Example

$$\mathbf{v}_k = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \mathbf{v}_{k-1} \quad \mathbf{v}_0 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Determine a closed form for the above linear dynamical system.

Solutions with Eigenvectors as Initial States

Solutions with Eigenvectors as Initial States

It's easy to give a closed-form solution if the initial state is an eigenvector:

$$\mathbf{v}_k = A^k \mathbf{v}_0 = \lambda^k \mathbf{v}_0$$

Solutions with Eigenvectors as Initial States

It's easy to give a closed-form solution if the initial state is an eigenvector:

$$\mathbf{v}_k = A^k \mathbf{v}_0 = \lambda^k \mathbf{v}_0$$

No dependence on A^k or \mathbf{v}_{k-1}

Solutions with Eigenvectors as Initial States

It's easy to give a closed-form solution if the initial state is an eigenvector:

$$\mathbf{v}_k = A^k \mathbf{v}_0 = \lambda^k \mathbf{v}_0$$

No dependence on A^k or \mathbf{v}_{k-1}

The Key Point. This is still true of sums of eigenvectors.

Solutions in terms of eigenvectors

Let's simplify $A^k \mathbf{v}$, given we have eigenvectors $\mathbf{b}_1, \mathbf{b}_2$ for A which span all of \mathbb{R}^2 :

Eigenvectors and Growth in the Limit

Theorem. For a linear dynamical system A with initial state \mathbf{v}_0 , if \mathbf{v}_0 can be written in terms of eigenvectors $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_k$ of A with eigenvalues

$$\lambda_1 > \lambda_2 \dots \geq \lambda_k$$

then $\mathbf{v}_k \sim \lambda_1^k c_1 \mathbf{b}_1$ for some constant c_1 (in other words, in the long term, the system grows exponentially in λ_1).

Verify:

Eigenbases

Eigenbases

Definition. An **eigenbasis** of \mathbb{R}^n for a $n \times n$ matrix A is a basis of \mathbb{R}^n made up entirely of eigenvectors of A .

Eigenbases

Definition. An **eigenbasis** of \mathbb{R}^n for a $n \times n$ matrix A is a basis of \mathbb{R}^n made up entirely of eigenvectors of A .

*We can represent vectors as **unique** linear combinations of eigenvectors.*

Eigenbases

Definition. An **eigenbasis** of \mathbb{R}^n for a $n \times n$ matrix A is a basis of \mathbb{R}^n made up entirely of eigenvectors of A .

*We can represent vectors as **unique** linear combinations of eigenvectors.*

Not all matrices have eigenbases.

Eigenbases and Growth in the Limit

Theorem. For a linear dynamical system A with initial state \mathbf{v}_0 , if A has an eigenbasis $\mathbf{b}_1, \dots, \mathbf{b}_k$, then

$$\mathbf{v}_k \sim \lambda_1^k c_1 \mathbf{b}_1$$

for some constant c_1 , where λ_1 is the **largest eigenvalue** of A and \mathbf{b}_1 is its **eigenvector**.

Eigenbases and Growth in the Limit

Theorem. For a linear dynamical system A with initial state \mathbf{v}_0 , if A has an eigenbasis $\mathbf{b}_1, \dots, \mathbf{b}_k$, then


$$\mathbf{v}_k \sim \lambda_1^k c_1 \mathbf{b}_1$$

for some constant c_1 , where λ_1 is the **largest eigenvalue of A** and \mathbf{b}_1 **is its eigenvector**.

The largest eigenvalue describes the long-term exponential behavior of the system.

Example: CS Major Growth

see the notes for more details



$$\mathbf{v}_0 = \begin{bmatrix} v_{0,1} \\ v_{0,2} \\ v_{0,3} \\ v_{0,4} \end{bmatrix} = \begin{bmatrix} \# \text{ of year 1 students enrolled in 2024} \\ \# \text{ of year 2 students enrolled in 2024} \\ \# \text{ of year 3 students enrolled in 2024} \\ \# \text{ of year 4 students enrolled in 2024} \end{bmatrix}$$

$$\mathbf{v}_k = A^k \mathbf{v}_0$$

(A is determined by least squares)

This is clearly exponential. If we want to "extract" the exponent, we need to look at the largest eigenvalue.

moving on...

Finding Eigenvalues

Finding Eigenvalues

Question. Determine the eigenvalues of A , along with their associated eigenspaces.

Finding Eigenvalues

Question. Determine the eigenvalues of A , along with their associated eigenspaces.

Solution (Idea). Can we somehow "solve for λ " in the equation

$$(A - \lambda I)\mathbf{x} = \mathbf{0}$$


Determinants

An Aside: Determinants are Mysterious

Determinants are
strangely polarizing

Some people love them,
some people hate them

We'll only scratch the
surface...



What kind of thing is the determinant?

What kind of thing is the determinant?

A determinant is a number associated with a matrix.

What kind of thing is the determinant?

A determinant is a number associated with a matrix.

Notation. We will write $\det(A)$ for the determinant of A .

What kind of thing is the determinant?

A determinant is a number associated with a matrix.

Notation. We will write $\det(A)$ for the determinant of A .

In broad strokes, it's a big sum of products of entries of A .

A Scary-Looking Definition (we won't use)

$$\det(A) = \sum_{\sigma \in S_n} (-1)^{\text{sgn}(\sigma)} A_{1\sigma(1)} A_{2\sigma(2)} \cdots A_{n\sigma(n)}$$

We can think of this function as a procedure:

```
1 FUNCTION det(A):  
2   total = 0  
3   FOR all matrix B we can get by swapping a bunch of rows of A:  
4     s = 1 IF (# of swaps necessary) is even ELSE -1  
5     total += s * (product of the diagonal entries of B)  
6   RETURN total
```

The Determinant of 2×2 Matrices

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$$

The Determinant of 2×2 Matrices

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \rightarrow^0 \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$(-1)^0 ad$$

The Determinant of 2×2 Matrices

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \rightarrow^1 \begin{bmatrix} c & d \\ a & b \end{bmatrix}$$

$$(-1)^1 cb$$

$$A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

The Determinant of 3×3 matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

The Determinant of 3×3 matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^0 \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$(-1)^0 aei$$

The Determinant of 3×3 matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^2 \begin{bmatrix} g & h & i \\ a & b & c \\ d & e & f \end{bmatrix}$$

$$(-1)^2 gbf$$

The Determinant of 3×3 matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^2 \begin{bmatrix} d & e & f \\ g & h & i \\ a & b & c \end{bmatrix}$$

$$(-1)^2 dhc$$

The Determinant of 3×3 matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^1 \begin{bmatrix} g & h & i \\ d & e & f \\ a & b & c \end{bmatrix}$$

$$(-1)^1 gec$$

The Determinant of 3×3 matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^1 \begin{bmatrix} d & e & f \\ a & b & c \\ g & h & i \end{bmatrix}$$

$$(-1)^1 dbi$$

The Determinant of 3×3 matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^1 \begin{bmatrix} a & b & c \\ g & h & i \\ d & e & f \end{bmatrix}$$

$$(-1)^1 ahf$$

Another Perspective

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Let's row reduce an arbitrary 2×2 matrix:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \sim \begin{bmatrix} a & b & | & 0 \\ 0 & ad-bc & | & 0 \end{bmatrix}$$

$$A\vec{x} = 0$$

sol'n set

\uparrow if $ad-bc = 0 \Rightarrow$ free variable

$\Rightarrow A\vec{x} = 0$ has nontrivial sol'n's

$$\det(A) = ad-bc = 0 \Leftrightarrow A \text{ not invertible}$$

$\Rightarrow A$ not invertible

Another Perspective

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

Let's row reduce an arbitrary 3×3 matrix:

$$\left[\begin{array}{ccc} \cancel{1} & \cancel{1} & \cancel{1} \\ 0 & \cancel{1} & * \\ 0 & 0 & a(\det A) \end{array} \right]$$

Determinants and Invertibility

Determinants and Invertibility

Theorem. A matrix is invertible if and only if $\det(A) \neq 0$.

Determinants and Invertibility

Theorem. A matrix is invertible if and only if $\det(A) \neq 0$.

So we can yet again extend the IMT:

Determinants and Invertibility

Theorem. A matrix is invertible if and only if $\det(A) \neq 0$.

So we can yet again extend the IMT:

- » A is invertible
- » $\det(A) \neq 0$
- » 0 is not an eigenvalue

$A\vec{v} = \vec{0}$ has some nontrivial sol'n

so $\text{Nul } A \neq \{\vec{0}\}$


These must be all true or all false.

so A not invertible

A Geometric Interpretation: Volume


$$|\det[\vec{v}_1, \vec{v}_2]|$$

$$\parallel \text{vol}(P)$$



$$|\det[\vec{v}_1, \vec{v}_2, \vec{v}_3]|$$

$$\parallel \text{vol}(P)$$



A not invertible $\Rightarrow \dim(\text{col } A) < n \Rightarrow P$ has zero volume
 $\Rightarrow \det(A) = 0$

(look up cofactor expansion also)

Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} U_{11} U_{22} \cdots U_{nn}$$

↙ echelon form

Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} U_{11} U_{22} \cdots U_{nn}$$

Defintion. The **determinant** of a matrix A is given by the above equation, where

Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} U_{11} U_{22} \cdots U_{nn}$$

Defintion. The **determinant** of a matrix A is given by the above equation, where

- U is an echelon form of A

Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} U_{11} U_{22} \cdots U_{nn}$$

Defintion. The **determinant** of a matrix A is given by the above equation, where

- U is an echelon form of A
- s is the number of row swaps used to get U

Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} U_{11} U_{22} \cdots U_{nn}$$

Defintion. The **determinant** of a matrix A is given by the above equation, where

- U is an echelon form of A
- s is the number of row swaps used to get U
- c is the product of all scalings used to get U

Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} \text{product of diagonal entries } U_{11}U_{22}\cdots U_{nn}$$

Defintion. The **determinant** of a matrix A is given by the above equation, where

- U is an echelon form of A
- s is the number of row swaps used to get U
- c is the product of all scalings used to get U

Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} \text{product of diagonal entries } U_{11}U_{22}\cdots U_{nn}$$

$c = 0$ if A is not invertible

Defintion. The **determinant** of a matrix A is given by the above equation, where

- U is an echelon form of A
- s is the number of row swaps used to get U
- c is the product of all scalings used to get U

Example

$$s = 0 + 1$$
$$c = 1 \cdot (-1/2)$$

$$R_2 \leftarrow R_2 - 2R_1$$

$$\begin{bmatrix} 1 & 5 & 0 \\ 2 & 4 & -1 \\ 0 & -2 & 0 \end{bmatrix}$$

Let's find the determinant of this matrix:

$$\begin{bmatrix} 1 & 5 & 0 \\ 0 & -6 & -1 \\ 0 & -2 & 0 \end{bmatrix} \xrightarrow[\text{② } R_2 \leftarrow -1/2 R_2]{\text{① swap } R_2, R_3} \begin{bmatrix} 1 & 5 & 0 \\ 0 & 1 & 0 \\ 0 & -6 & -1 \end{bmatrix} \xrightarrow{R_3 \leftarrow R_3 + 6R_2} \begin{bmatrix} 1 & 5 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

$$\det(A) = \frac{(-1)^1}{-1/2} (1)(1)(-1) = -2$$

Example (Again)

$$S = 0 + 1 + 1$$

$$C = 1$$

$$R_1 \leftrightarrow R_2$$

$$A = \begin{bmatrix} 1 & 5 & 0 \\ 2 & 4 & -1 \\ 0 & -2 & 0 \end{bmatrix}$$

Let's find the determinant of this matrix again but with a different sequence of row operations:

$$\begin{bmatrix} 2 & 4 & -1 \\ 1 & 5 & 0 \\ 0 & -2 & 0 \end{bmatrix}$$

$$\underline{R_2 \leftrightarrow R_2 - \frac{1}{2}R_1}$$

$$\begin{bmatrix} 2 & 4 & -1 \\ 0 & 3 & \frac{1}{2} \\ 0 & -2 & 0 \end{bmatrix}$$

$$\underline{R_2 \leftrightarrow R_3}$$

$$\begin{bmatrix} 2 & 4 & -1 \\ 0 & -2 & 0 \\ 0 & 3 & \frac{1}{2} \end{bmatrix}$$

$$\} R_3 + \frac{3}{2}R_2$$

$$\begin{bmatrix} 2 & 4 & -1 \\ 0 & -2 & 0 \\ 0 & 0 & \frac{1}{2} \end{bmatrix}$$

$$\det A = \frac{(-1)^2}{1} (2)(-2)\left(\frac{1}{2}\right) = -2$$

The definition holds no matter
which sequence of row
operations you use.

How To: Determinants

How To: Determinants

Question. Determine the determinant of a matrix A .

How To: Determinants

Question. Determine the determinant of a matrix A .

Solution.

How To: Determinants

Question. Determine the determinant of a matrix A .

Solution.

1. Convert A to an echelon form U .

How To: Determinants

Question. Determine the determinant of a matrix A .

Solution.

1. Convert A to an echelon form U .
2. Keep track of the number of row swaps you used, call this s , and the product of all scalings, call this c

How To: Determinants

Question. Determine the determinant of a matrix A .

Solution.

1. Convert A to an echelon form U .
2. Keep track of the number of row swaps you used, call this s , and the product of all scalings, call this c
3. Determine the product of entries along the diagonal of U , call this P .

How To: Determinants

Question. Determine the determinant of a matrix A .

Solution.

1. Convert A to an echelon form U .
2. Keep track of the number of row swaps you used, call this s , and the product of all scalings, call this c .
3. Determine the product of entries along the diagonal of U , call this P .
4. The determinant of A is $\frac{(-1)^s P}{c}$.

Question

$$S = 1$$
$$C = 1$$

$$A = \begin{bmatrix} 1 & 5 & -4 \\ -1 & -5 & 5 \\ -2 & -8 & 7 \end{bmatrix}$$

$$\det A = \frac{(-1)^1}{1} (1)(2)(1) = -2$$

Find the determinant of the above matrix.

$$\begin{bmatrix} 1 & 5 & -4 \\ 0 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \xrightarrow{R_2 \leftrightarrow R_3} \begin{bmatrix} 1 & 5 & -4 \\ 0 & 2 & -1 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer

The Shorter Version

Beyond small matrices, we'll just use a computer

With NumPy:

`numpy.linalg.det(A)`

Properties of Determinants

Properties of Determinants (1)

$$\det(AB) = \det(A) \det(B)$$

$$\det(E_k \cdots E_1 A) = \det(E_k) \cdots \det(E_1) \det(A)$$

It follows that AB is invertible if and only if A and B are invertible

(we won't verify this)

Example Question

$$\begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix}$$

Use the fact that $\det(AB) = \det(A)\det(B)$ to give an expression for $\det(A^{-1})$ in terms of $\det(A)$.

Hint. What is $\det(I)$? $\rightarrow 1$

$$\det(I) = \det(A \cdot A^{-1}) = \det(A)\det(A^{-1}) = 1$$
$$\det(A^{-1}) = \frac{1}{\det(A)}$$

Properties of Determinants (2)

$$\det(A^T) = \det(A)$$

It follows that A^T is invertible if and only if A is invertible.

(we also won't verify this)

Example Question

orthogonal matrices

If $A^{-1} = A^T$, then what are the possible values of $\det(A)$?

$$\det(A^T) = \det(A^{-1}) = \frac{1}{\det A}$$


$\det(A^T) \stackrel{||}{=} \det(A)$

$$\frac{1}{\det A} = \det A \Rightarrow (\det A)^2 = 1$$
$$\det A = \pm 1$$

Properties of Determinants (3)

Theorem. If A is triangular, then $\det(A)$ is the product of entries along the diagonal.

Verify:



$$A = \begin{pmatrix} a_{11} & & 0 \\ * & \ddots & \\ & & a_{nn} \end{pmatrix}$$

$$\det(A) = \det(A^T)$$

turns it into an upper triangular matrix

Answer

Characteristic Equation

What kind of thing is the determinant, really?

What kind of thing is the determinant, really?

The determinant of a matrix A is an arithmetic expression written in terms of the entries of A .

What kind of thing is the determinant, really?

The determinant of a matrix A is an arithmetic expression written in terms of the entries of A .

But a matrix may not have numbers as entries.

What kind of thing is the determinant, really?

The determinant of a matrix A is an arithmetic expression written in terms of the entries of A .

But a matrix may not have numbers as entries.

We might think of the matrix $A - \lambda I$ as having *polynomials* as entries.

What kind of thing is the determinant, really?


The determinant of a matrix A is an arithmetic expression written in terms of the entries of A .

But a matrix may not have numbers as entries.

We might think of the matrix $A - \lambda I$ as having *polynomials* as entries.


Then $\det(A - \lambda I)$ is a **polynomial**.

Reminder: Polynomial Roots



Reminder: Polynomial Roots


A **root** of a polynomial $p(x)$ is a value r such that $p(r) = 0$.



Reminder: Polynomial Roots

A **root** of a polynomial $p(x)$ is a value r such that $p(r) = 0$.

(A polynomial may have many roots)




Reminder: Polynomial Roots

A **root** of a polynomial $p(x)$ is a value r such that $p(r) = 0$.

(A polynomial may have many roots)

If r is a root of $p(x)$, then it is possible to find a polynomial $q(x)$ such that

$$p(x) = (x - r)q(x)$$



Characteristic Polynomial

Characteristic Polynomial

Definition. The characteristic polynomial of a matrix A is $\det(A - \lambda I)$ viewed as a polynomial in the variable λ .

Characteristic Polynomial

Definition. The characteristic polynomial of a matrix A is $\det(A - \lambda I)$ viewed as a polynomial in the variable λ .

This is a polynomial with the eigenvalues of A as roots.

Characteristic Polynomial

Definition. The characteristic polynomial of a matrix A is $\det(A - \lambda I)$ viewed as a polynomial in the variable λ .

This is a polynomial with the eigenvalues of A as roots.

So we can "solve" for the eigenvalues in the equation

$$\det(A - \lambda I) = 0$$

"Deriving" the characteristic polynomial

Q: When is λ an eigenvalue for A ?

A: When $(A - \lambda I)\vec{v} = 0$ has nontrivial solutions.

\Downarrow ($A - \lambda I$ not invertible)

$$\underline{\underline{\det(A - \lambda I) = 0}}$$

Hence, the characteristic polynomial

Example: 2×2 Matrix

$$rk \left(A - \frac{1+\sqrt{5}}{2} I \right) = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

Let's find the characteristic polynomial of this matrix:

$$A - \lambda I = \begin{bmatrix} 1-\lambda & 1 \\ 1 & -\lambda \end{bmatrix} \Rightarrow \det(A - \lambda I) = (1-\lambda)(-\lambda) - (1)(1) \\ = \boxed{\lambda^2 - \lambda - 1}$$

$$ax^2 + bx + c = 0$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\lambda = \frac{1 \pm \sqrt{1 - 4(1)(-1)}}{2} \\ = \frac{1 \pm \sqrt{5}}{2}$$

Example: Triangular matrix

$$\begin{bmatrix} 1 & -3 & 0 & 6 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 4 \end{bmatrix}$$

The characteristic polynomial of a triangular matrix comes pre-factored:

$$A - \lambda I = \begin{bmatrix} 1-\lambda & -3 & 0 & 6 \\ 0 & -\lambda & 1 & 1 \\ 0 & 0 & 1-\lambda & 2 \\ 0 & 0 & 0 & 4-\lambda \end{bmatrix} \Rightarrow \det(A - \lambda I) = (1-\lambda)^2 (-\lambda) (4-\lambda)$$

$\uparrow \quad \quad \uparrow \quad \quad \uparrow$

$$= (\lambda-1)^2 \lambda (\lambda-4)$$

How To: Finding Eigenvalues

How To: Finding Eigenvalues

Question. Find all eigenvalues of the matrix A .

How To: Finding Eigenvalues

Question. Find all eigenvalues of the matrix A .

Solution. Find the roots of the characteristic polynomial of A .

An Observation: Multiplicity

$$\lambda^1(\lambda - 1)^2(\lambda - 4)^1 \text{ multiplicities}$$

In the examples so far, we've seen a number appear as a root multiple times.

This is called the **multiplicity** of the root.

Is the multiplicity meaningful in this context?

Multiplicity and Dimension

Theorem. The dimension of the eigenspace of A for the eigenvalue λ is at most the multiplicity of λ in $\det(A - \lambda I)$.

The multiplicity is an upper bound on
"how large" the eigenspace is.

dimensionality

"Eigenspace $_{\lambda}$ " = $\text{Nul}(A - \lambda I)$
not notation

Example

$$A\vec{v} = 0\vec{v} = \vec{0}$$

Let A be a 5×5 matrix with characteristic polynomial $(x-1)^3(x-3)(x+5)$.

» What is $\text{rank}(A)$?

$$\text{rk } A \geq 3$$

← # distinct nonzero eigenvalues

» What is the minimum possible rank of $A - I$?

$$\text{rk } A - I + \text{nullity } A - I = 5$$

$$\boxed{\text{rk } A - I \geq 2}$$

$$\leq 3$$

$$1 \leq \dim(\text{Nu}(A - I)) \leq 3$$