

Matrix Inverse

Geometric Algorithms
Lecture 10

Practice Problem

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \mapsto \begin{bmatrix} 2x_1 - x_2 \\ 4x_2 \\ 3x_1 - 3x_2 \end{bmatrix}$$

Determine if the above transformation is onto, one-to-one, both, or neither

Answer

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 2 \\ 0 \\ 3 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} \mapsto \begin{bmatrix} -1 \\ 4 \\ -3 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 \\ 0 & 4 \\ 3 & -3 \end{bmatrix} \sim \begin{bmatrix} \boxed{1} & 0 \\ 0 & \boxed{1} \\ 0 & 0 \end{bmatrix} \left. \vphantom{\begin{bmatrix} 2 & -1 \\ 0 & 4 \\ 3 & -3 \end{bmatrix}} \right\} \begin{array}{l} \text{work} \\ \text{this} \\ \text{out} \end{array}$$

$$\vec{x} \mapsto \begin{bmatrix} 2 & -1 \\ 0 & 4 \\ 3 & -3 \end{bmatrix} \vec{x}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \mapsto \begin{bmatrix} 2x_1 - x_2 \\ 4x_2 \\ 3x_1 - 3x_2 \end{bmatrix}$$

onto $\Leftrightarrow$ 1 pivot / row

Not onto

1-1 $\Leftrightarrow$ 1 pivot / col $\Leftrightarrow$ cols L.I.

is 1-1

$\{\vec{v}_1, \vec{v}_2\}$ is L.D. $\Leftrightarrow \vec{v}_1 = c \vec{v}_2$

Objectives

» Motivate and define matrix inverses

Keywords

Matrix Transpose

Inner Product

Matrix Power

Square Matrix

Matrix Inverse

Invertible Transformation

1-1 Correspondence


`numpy.linalg.inv`

Determinant


Invertible Matrix Theorem

Recap


Recap: Shearing and Reflecting (Geometrically)



shear



reflect



Recap: The Key Fact

$$T(S(\vec{u} + \vec{v})) = T(S(\vec{u}) + S(\vec{v})) = T(S(\vec{u})) + T(S(\vec{v})) \quad (\text{additive})$$

Thm. The composition of two linear transformations is a linear transformation

This means the composition of two matrix transformations can be implemented by a *single* matrix

Matrix Multiplication

Df. For $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$ with columns $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_p$ the product $AB \in \mathbb{R}^{m \times p}$ is given by

$$AB = A \begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \dots & \mathbf{b}_p \end{bmatrix} = \begin{bmatrix} A\mathbf{b}_1 & A\mathbf{b}_2 & \dots & A\mathbf{b}_p \end{bmatrix}$$

Replace each column of B with A multiplied by that column

Recap: Matrix Multiplication and Composition

Thm. For any matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$ and any vector $\mathbf{v} \in \mathbb{R}^p$,

$$A(B\mathbf{v}) = (AB)\mathbf{v}$$

The matrix implementing the composition is the product of the two underlying matrices

Recap: Matrix "Interface"

multiplication just given

addition add entry-wise

scaling scale entry-wise


Recall: Transpose (Pictorially)

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 \end{bmatrix}$$



$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 2 & 2 \\ 3 & 3 & 3 & 3 \\ 4 & 4 & 4 & 4 \\ 5 & 5 & 5 & 5 \end{bmatrix}$$

Recall: Transpose (Pictorially)



Recall: Transpose

Df. For $A \in \mathbb{R}^{m \times n}$ the **transpose** of A , written A^T , is the $n \times m$ matrix such that

$$(A^T)_{ij} = A_{ji}$$

Ex.

$$\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}^T = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$$

Recall: Algebraic Properties

$$(A^T)^T = A$$

$$(A + B)^T = A^T + B^T$$

$$(cA)^T = cA^T \text{ (where } c \text{ is a scalar)}$$

$$(AB)^T = B^T A^T$$

Recall: Algebraic Properties

$$(A^T)^T = A$$

$$(A + B)^T = A^T + B^T$$

$$(cA)^T = cA^T \text{ (where } c \text{ is a scalar)}$$

$$(AB)^T = B^T A^T \text{ Important: the order reverses!}$$

Transposes and Inner Products

$$[u_1 \quad u_2 \quad u_3 \quad u_4] \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = u_1 v_1 + u_2 v_2 + u_3 v_3 + u_4 v_4$$

Transposes and Inner Products

$$[u_1 \quad u_2 \quad u_3 \quad u_4] \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = u_1 v_1 + u_2 v_2 + u_3 v_3 + u_4 v_4$$

Df. The **inner product** of two vectors **u** and **v** in $\mathbb{R}^n$ is $\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u} \cdot \mathbf{v} = \mathbf{u}^T \mathbf{v}$

Recall: Matrix Powers

Recall: Matrix Powers

Obs. If $A \in \mathbb{R}^{n \times n}$ then AA is well-defined

Recall: Matrix Powers

Obs. If $A \in \mathbb{R}^{n \times n}$ then AA is well-defined

Df. For $A \in \mathbb{R}^{n \times n}$ we write A^k for the k -fold product of A with itself

Recall: Matrix Powers

Obs. If $A \in \mathbb{R}^{n \times n}$ then AA is well-defined

Df. For $A \in \mathbb{R}^{n \times n}$ we write A^k for the k -fold product of A with itself

What should A^0 be? (we want $A^0 A^k = A^{0+k} = A^k$)

$$A^{-1} A^{-2} = 1, \quad A^{-1} A^{-1} = 1$$

$$A^0 = I$$

$$A^0 A^k = A^{0+k} = A^k$$

$$A^m A^n = A^{m+n}$$

Matrix Powers (Computationally)

Use `numpy.linalg.matrix_power`

This can be *much* faster than doing a sequence of matrix multiplications (*why?*)

$$A^8 = A A A A A A A A$$

$$A^8 = (A^4)^2 = ((A^2)^2)^2$$

Recall: Final Warnings

Recall: Final Warnings

1. AB isn't necessarily equal to BA , even if both are well-defined

Recall: Final Warnings

1. AB isn't necessarily equal to BA , even if both are well-defined
2. If $AB = AC$ then it isn't necessary that $B = C$ even if $A \neq 0$

Recall: Final Warnings

1. AB isn't necessarily equal to BA , even if both are well-defined
2. If $AB = AC$ then it isn't necessary that $B = C$ even if $A \neq 0$
3. If $AB = 0$ (the zero matrix) it isn't necessary that $A = 0$ or $B = 0$

Matrix Operations (So Far)

Matrix Operations (So Far)

transpose

$$A^T$$

Matrix Operations (So Far)

transpose

$$A^T$$

scaling

$$cA$$

Matrix Operations (So Far)

transpose

$$A^T$$

scaling

$$cA$$

addition (subtraction)

$$A + B$$

$$A + (-1)B = A - B$$

Matrix Operations (So Far)

transpose

$$A^T$$

scaling

$$cA$$

addition (subtraction)

$$A + B$$

$$A + (-1)B = A - B$$

multiplication (powers)

$$AB$$

$$A^k$$

Matrix Inverses

Recall: The Identity Matrix

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \boxed{1} & 0 \\ 0 & \boxed{0} & 1 \end{bmatrix} \quad (I_3)_{22} \quad (I_3)_{32}$$
$$(I_n)_{ij} = \begin{cases} 1 & i=j \\ 0 & i \neq j \end{cases}$$

$$A \in \mathbb{R}^{m \times n}$$

$$I_m A = A = A I_n$$

$$\vec{x} \mapsto I \vec{x} \quad \text{"do nothing" transformation}$$

Recall: The Identity Matrix

Recall: The Identity Matrix

Df. The $n \times n$ **identity matrix** is the matrix whose *diagonal* contains all 1s, and all other entries are 0s

$$I_{ij} = \begin{cases} 1 & i = j \\ 0 & \text{otherwise} \end{cases}$$

Recall: The Identity Matrix

Df. The $n \times n$ **identity matrix** is the matrix whose *diagonal* contains all 1s, and all other entries are 0s

$$I_{ij} = \begin{cases} 1 & i = j \\ 0 & \text{otherwise} \end{cases}$$

Ex.

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Recall: The Identity Matrix

Recall: The Identity Matrix

The identity matrix implements the "do nothing" transformation. For any $\mathbf{v}$,

$$I\mathbf{v} = \mathbf{v}$$

Recall: The Identity Matrix

The identity matrix implements the "do nothing" transformation. For any $\mathbf{v}$,

$$I\mathbf{v} = \mathbf{v}$$

It is the "1" of matrices. For any A

$$IA = AI = A$$

Recall: The Identity Matrix

The identity matrix implements the "do nothing" transformation. For any $\mathbf{v}$,

$$I\mathbf{v} = \mathbf{v}$$

It is the "1" of matrices. For any A

$$IA = AI = A$$

These may be different sizes

Recall: The Identity Matrix

$$\begin{array}{ccccc} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} & \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \end{bmatrix} & = & \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \end{bmatrix} & \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} & = & \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \end{bmatrix} \\ 2 \times 2 & 2 \times 4 & & 2 \times 4 & 4 \times 4 & & 2 \times 4 \end{array}$$

Basic Algebra

$$2x = 10$$

Basic Algebra

$$2x = 10$$

How do we solve this equation?

Basic Algebra

$$2x = 10$$

How do we solve this equation?

Divide on both sides by 2 to get $x = 5$

Basic Algebra

$$2x = 10$$

How do we solve this equation?

Divide on both sides by 2 to get $x = 5$

i.e., multiply each side by $\frac{1}{2}$ (a.k.a. 2^{-1})

Basic Algebra

$$\left(\frac{1}{2}\right)2x = 10 \quad \left(\frac{1}{2}\right)$$

$$\textcircled{x} = 1x = \textcircled{5}$$

How do we solve this equation?

Divide on both sides by 2 to get $x = 5$

i.e., multiply each side by $\frac{1}{2}$ (a.k.a. 2^{-1})

$\frac{1}{2}$ is the **reciprocal** or **multiplicative inverse** of 2

Wouldn't it be nice...

$$\mathbf{Ax} = \mathbf{b}$$

Wouldn't it be nice...

$$\mathbf{Ax} = \mathbf{b}$$

How do we solve this equation?

Wouldn't it be nice...

$$\mathbf{Ax} = \mathbf{b}$$

How do we solve this equation?

Multiply each side by A^{-1} to get $\mathbf{x} = A^{-1}\mathbf{b}$

Wouldn't it be nice...

$$\mathbf{Ax} = \mathbf{b}$$

How do we solve this equation?

Multiply each side by A^{-1} to get $\mathbf{x} = A^{-1}\mathbf{b}$

A^{-1} is the **multiplicative inverse** of A

Wouldn't it be nice...

$$A^{-1}A\mathbf{x} = A^{-1}\mathbf{b}$$

How do we solve this equation?

Multiply each side by A^{-1} to get $\mathbf{x} = A^{-1}\mathbf{b}$

A^{-1} is the **multiplicative inverse** of A

Wouldn't it be nice...

$$\mathbf{I}\mathbf{x} = A^{-1}\mathbf{b}$$

How do we solve this equation?

Multiply each side by A^{-1} to get $\mathbf{x} = A^{-1}\mathbf{b}$

A^{-1} is the **multiplicative inverse** of A

Wouldn't it be nice...

$$\mathbf{x} = A^{-1}\mathbf{b}$$

How do we solve this equation?

Multiply each side by A^{-1} to get $\mathbf{x} = A^{-1}\mathbf{b}$

A^{-1} is the **multiplicative inverse** of A

Do all matrices have
inverses?

Do all matrices have
inverses?

*No. If they did, then every linear
system would have a solution*

When does a matrix have
an inverse?

Square Matrices

Df. A matrix $A \in \mathbb{R}^{m \times n}$ is **square** if $m = n$

$$\begin{bmatrix} * & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{bmatrix}$$

i.e., it has same number of rows as columns

Square Matrices are Special

Square Matrices are Special

They are the only kind of matrices...

Square Matrices are Special

They are the only kind of matrices...

» that can have a pivot in every row and every column

Square Matrices are Special

They are the only kind of matrices...

» that can have a pivot in every row and every column

» whose transformations can be both 1-1 and onto

Square Matrices are Special

They are the only kind of matrices...

» that can have a pivot in every row and every column

» whose transformations can be both 1-1 and onto

» whose columns can have full span and be linearly independent

Square Matrices are Special

They are the only kind of matrices...

» that can have a pivot in every row and every column

» whose transformations can be both 1-1 and onto

» whose columns can have full span and be linearly independent

» *that can have inverses*

Matrix Inverses

Df. Given $A \in \mathbb{R}^{n \times n}$, a matrix $B \in \mathbb{R}^{n \times n}$ is
an **inverse** of A if:

$$AB = I_n$$

$$BA = I_n$$

Matrix Inverses

Matrix Inverses

Df. For a matrix $A \in \mathbb{R}^{n \times n}$, an **inverse** of A is a matrix $B \in \mathbb{R}^{n \times n}$ such that

$$AB = I_n \text{ and } BA = I_n$$

Matrix Inverses

Df. For a matrix $A \in \mathbb{R}^{n \times n}$, an **inverse** of A is a matrix $B \in \mathbb{R}^{n \times n}$ such that

$$AB = I_n \text{ and } BA = I_n$$

A is **invertible** if it has an inverse. Otherwise it is **singular**

Example


$$\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} = \begin{bmatrix} 1(1) + 0(-1) & 1(0) + 0(1) \\ 1(1) + 1(-1) & 1(0) + 1(1) \end{bmatrix}$$
$$= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \checkmark$$

$$\begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \checkmark$$

Example

$$\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$$



Example


Reflection across the x -axis is it's own inverse

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} \mapsto \begin{bmatrix} 0 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$



Non-Example

$$A \quad B = I$$
$$\begin{bmatrix} 1 & 2 & -1 \\ 0 & 3 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \vec{b}_1 & \vec{b}_2 & \vec{b}_3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$A \vec{b}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & -1 & 0 \\ 0 & 3 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

A has no inverse

Inverses are Unique

Thm. If B and C are inverses of A , then $B = C$

$$B = I B = (C A) B = C (A B) = C I = C$$

$CA = I$ since

C is an ~~inverse~~

$AB = I$ since

B is an ~~inverse~~

A^{-1} is the ~~inverse~~ of A (if it exists)

Solutions for Invertible Matrix Equations

Thm. For matrix $A \in \mathbb{R}^{n \times n}$, if A is invertible then the equation $A\mathbf{x} = \mathbf{b}$ has a unique solution for any choice of $\mathbf{b}$

$$A \vec{v}_1 = \vec{b}$$

$$A \vec{v}_2 = \vec{b}$$

$$A \vec{v}_1 = A \vec{v}_2 \quad \swarrow \text{uniqueness}$$

$$\vec{v}_1 = I \vec{v}_1 = A^{-1} A \vec{v}_1 = A^{-1} A \vec{v}_2 = I \vec{v}_2 = \vec{v}_2$$

$$A \vec{x} = \mathbf{b}$$

$$A \boxed{A^{-1} \vec{b}} = \mathbf{b} \quad \nwarrow \text{existence}$$

Unique Solutions

If $A\mathbf{x} = \mathbf{b}$ has a unique solution for any choice of $\mathbf{b}$, then it has

» exactly one solution for any choice of $\mathbf{b}$

Unique Solutions

If $A\mathbf{x} = \mathbf{b}$ has a unique solution for any choice of $\mathbf{b}$, then it has

» at least one solution for any choice of $\mathbf{b}$

» at most one solution for any choice of $\mathbf{b}$

Unique Solutions

If $A\mathbf{x} = \mathbf{b}$ has a unique solution for any choice of $\mathbf{b}$, then it has

» T is onto

» T is one-to-one


where T is implemented by A

Connection to Transformations

Df. A linear transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is **invertible** if there is a linear transformation S such that

$$S(T(\mathbf{v})) = \mathbf{v} \quad \text{and} \quad T(S(\mathbf{v})) = \mathbf{v}$$

for any $\mathbf{v}$ in $\mathbb{R}^n$



Connection to Transformations

Connection to Transformations

Thm. A matrix $A \in \mathbb{R}^{n \times n}$ is invertible if and only if the matrix transformation $\mathbf{x} \mapsto A\mathbf{x}$ is invertible

Connection to Transformations

Thm. A matrix $A \in \mathbb{R}^{n \times n}$ is invertible if and only if the matrix transformation $\mathbf{x} \mapsto A\mathbf{x}$ is invertible

A matrix is invertible if it's possible to "undo" its transformation without "losing information"


Non-Example

Projection onto the x -axis in $\mathbb{R}^2$

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} \mapsto \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ is
not invertible



Connection to Transformations

Connection to Transformations

Df. A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a **one-to-one correspondence** (bijection) if any vector $\mathbf{b}$ in $\mathbb{R}^n$ is the **image of exactly one vector** $\mathbf{v}$ in $\mathbb{R}^n$ (where $T(\mathbf{v}) = \mathbf{b}$)

Connection to Transformations

Df. A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a **one-to-one correspondence** (bijection) if any vector $\mathbf{b}$ in $\mathbb{R}^n$ is the **image of exactly one vector** $\mathbf{v}$ in $\mathbb{R}^n$ (where $T(\mathbf{v}) = \mathbf{b}$)

A transformation is a 1-1 correspondence if it is 1-1 and onto


Connection to Transformations

Df. A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a **one-to-one correspondence** (bijection) if any vector $\mathbf{b}$ in $\mathbb{R}^n$ is the **image of exactly one vector** $\mathbf{v}$ in $\mathbb{R}^n$ (where $T(\mathbf{v}) = \mathbf{b}$)


A transformation is a 1-1 correspondence if it is 1-1 and onto

Invertible transformations are 1-1 correspondences


Kinds of Transformations (Pictorially)




1-1 correspondence



onto, not 1-1



1-1 not onto



not 1-1, not onto

Computing Matrix Inverses

Fundamental Questions

How do we determine if A has an inverse?

How do we compute A^{-1} ?

In General

$$A \begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \mathbf{b}_3 \end{bmatrix} = I$$

Can we solve for each $\mathbf{b}_i$?

$$\left[\begin{array}{c} A \vec{b}_1 \\ A \vec{b}_2 \\ A \vec{b}_3 \end{array} \right]$$

In General

$$A\mathbf{b}_1 = \mathbf{e}_1$$

$$A\mathbf{b}_2 = \mathbf{e}_2$$

$$A\mathbf{b}_3 = \mathbf{e}_3$$

If we want a matrix B such that $AB = I$, then the above equation must hold (in the case B has 3 columns)

In General

$$A\mathbf{b}_1 = \mathbf{e}_1$$

$$A\mathbf{b}_2 = \mathbf{e}_2$$

$$A\mathbf{b}_3 = \mathbf{e}_3$$

If we want a matrix B such that $AB = I$, then the above equation must hold (in the case B has 3 columns)

We need to solve 3 matrix equations

How To: Matrix Inverses

Q. Find the inverse of invertible $A \in \mathbb{R}^{n \times n}$

A. Solve the equation $A\mathbf{x} = \mathbf{e}_i$ for every standard basis vector $\mathbf{e}_i$. Put those solutions $\mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_n$ into a single matrix

$$[\mathbf{s}_1 \quad \mathbf{s}_2 \quad \dots \quad \mathbf{s}_n]$$

How To: Matrix Inverses

Q. Find the inverse of invertible $A \in \mathbb{R}^{n \times n}$

A. Row reduce the matrix $[A \ I]$ to a matrix $[I \ B]$.
Then B is the inverse of A

This is really the same thing. It's a simultaneous reduction

demo

Special Case: 2×2 Matrice Inverses

Special Case: 2×2 Matrice Inverses

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Special Case: 2×2 Matrice Inverses

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

The **determinant** of a 2×2 matrix is the value $ad - bc$

Special Case: 2×2 Matrice Inverses

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

The **determinant** of a 2×2 matrix is the value $ad - bc$

The inverse is defined only if the determinant is nonzero

Special Case: 2×2 Matrice Inverses

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$


The **determinant** of a 2×2 matrix is the value $ad - bc$

The inverse is defined only if the determinant is nonzero

(see the notes on linear transformations for more information about determinants)

Example

$$\begin{bmatrix} -6 & 14 \\ 3 & -7 \end{bmatrix} \quad \infty$$


$$\frac{1}{42 - 42} \begin{bmatrix} -7 & -14 \\ -3 & -6 \end{bmatrix}$$

not invertible

Algebra of Matrix Inverses

Challenge Problem

Challenge Problem

Df. B is right inverse of A if $AB = I$

Challenge Problem

Df. B is right inverse of A if $AB = I$

Df. B is a left inverse of A if $BA = I$

Challenge Problem

Df. B is right inverse of A if $AB = I$

Df. B is a left inverse of A if $BA = I$

Thm. If A has a right inverse B , then B is also a left inverse A

Challenge Problem

Df. B is right inverse of A if $AB = I$

Df. B is a left inverse of A if $BA = I$

Thm. If A has a right inverse B , then B is also a left inverse A

We only need to show A has a right/left inverse to show it is invertible

Algebraic Properties (Matrix Inverses)

Thm. For an invertible matrix A , the matrix A^{-1} is invertible

Algebraic Properties (Matrix Inverses)

Thm. For an invertible matrix A , the matrix A^T is invertible

Algebraic Properties (Matrix Inverses)

Thm. For a invertible matrices $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times n}$, the matrix AB is invertible

Question

Suppose that A is a $n \times n$ invertible matrix such that $A = A^T$ and B is a $m \times n$ matrix

Simplify the expression $A(BA^{-1})^T$ using the algebraic properties we've seen

Answer: B^T

$$A(BA^{-1})^T$$

$$A = A^T$$

Invertible Matrix Theorem

The Point

Q. How do we know if a square matrix is invertible?

A. *Every* perspective we've taken so far can help us answer this question

Invertible Matrix Theorem

Thm. For any invertible matrix $A \in \mathbb{R}^{n \times n}$

1. A^T is invertible

Invertible Matrix Theorem

Thm. For any invertible matrix $A \in \mathbb{R}^{n \times n}$

- 2. $A\mathbf{x} = \mathbf{b}$ has at least one solution for every $\mathbf{b}$
- 3. $A\mathbf{x} = \mathbf{b}$ has at most one solution for every $\mathbf{b}$
- 4. $A\mathbf{x} = \mathbf{b}$ has at exactly one solution for every $\mathbf{b}$

Invertible Matrix Theorem

Thm. For any invertible matrix $A \in \mathbb{R}^{n \times n}$

5. A has a pivot in every column

6. A has a pivot in every row

7. A is row equivalent to I_n

Invertible Matrix Theorem

Thm. For any invertible matrix $A \in \mathbb{R}^{n \times n}$

- 8. The columns of A are linearly independent
- 9. The columns of A span $\mathbb{R}^n$
- 10. $A\mathbf{x} = \mathbf{0}$ has only the trivial solution

Invertible Matrix Theorem

Thm. For any invertible matrix $A \in \mathbb{R}^{n \times n}$

- 11. The linear transformation $\mathbf{x} \mapsto A\mathbf{x}$ is onto
- 12. $\mathbf{x} \mapsto A\mathbf{x}$ is one-to-one
- 13. $\mathbf{x} \mapsto A\mathbf{x}$ is a one-to-one correspondence
- 14. $\mathbf{x} \mapsto A\mathbf{x}$ is invertible

Taking Stock: IMT

The following are logically equivalent:

1. A is invertible
2. A^T is invertible
3. $A\mathbf{x} = \mathbf{b}$ has at least one solution for any $\mathbf{b}$
4. $A\mathbf{x} = \mathbf{b}$ has at most one solution for any $\mathbf{b}$
5. $A\mathbf{x} = \mathbf{b}$ has a unique solution for any $\mathbf{b}$
6. A has n pivots (per row and per column)
7. A is row equivalent to I
8. $A\mathbf{x} = \mathbf{0}$ has only the trivial solution
9. The columns of A are linearly independent
10. The columns of A span $\mathbb{R}^n$
11. The linear transformation $\mathbf{x} \mapsto A\mathbf{x}$ is onto
12. $\mathbf{x} \mapsto A\mathbf{x}$ is one-to-one
13. $\mathbf{x} \mapsto A\mathbf{x}$ is a one-to-one correspondence
14. $\mathbf{x} \mapsto A\mathbf{x}$ is invertible

These all express the
same thing

!! only for square matrices !!

We get a lot of information for free

We get a lot of information for free

Thm. If A is square, then

A **is 1-1** if and only if A **is onto**

We get a lot of information for free

Thm. If A is square, then

A **is 1-1** if and only if A **is onto**

We only need to check one of these

We get a lot of information for free

Thm. If A is square, then

A is 1-1 if and only if A is onto

We only need to check one of these

Warning. Remember this only applies square matrices

We get a lot of information for free

Theorem. If A is square, then

A is invertible $\equiv Ax = 0$ implies $x = 0$

Invertibility is completely determined by what A sends to 0

Question

If A is invertible, and B is row equivalent to A , is B also invertible?

Challenge Problem

If $[\mathbf{a}_1 \ \mathbf{a}_2 \ \mathbf{a}_3]$ is invertible, is $[(\mathbf{a}_1 + \mathbf{a}_2 - 2\mathbf{a}_3) \ (\mathbf{a}_2 + 5\mathbf{a}_3) \ \mathbf{a}_3]$ also invertible?

Summary

The algebra of matrices can help us simplify matrix expressions

The invertible matrix theorem connects all the perspectives we've taken so far