Subspaces

Geometric Algorithms

CAS CS 132

Practice Problem

$$A = \begin{bmatrix} 1 & 0 & -1 & 2 \\ 2 & 0 & 2 & 3 \\ 1 & 4 & 0 & 2 \end{bmatrix} \quad \begin{matrix} R_1 \leftarrow R_1 + R_2 \\ R_3 \leftarrow 2R_3 \\ R_2 \leftrightarrow R_3 \\ \longrightarrow \end{matrix} \quad \begin{bmatrix} 3 & 0 & 1 & 5 \\ 2 & 8 & 0 & 4 \\ 2 & 0 & 2 & 3 \end{bmatrix} = B$$

Consider the following pair of matrices A and B which are row equivalent. Find a matrix E such that $EA = B$

Answer

$$R_1 \leftarrow R_1 + R_2$$
$$R_3 \leftarrow 2R_3$$
$$R_2 \leftrightarrow R_3$$

$$A = \begin{bmatrix} 1 & 0 & -1 & 2 \\ 2 & 0 & 2 & 3 \\ 1 & 4 & 0 & 2 \end{bmatrix} \longrightarrow \begin{bmatrix} 3 & 0 & 1 & 5 \\ 2 & 8 & 0 & 4 \\ 2 & 0 & 2 & 3 \end{bmatrix} = B$$

Objectives

- Introduce **subspaces** and **bases**
- Extend our intuitions about planes in $\mathbb{R}^3$ to subspaces in $\mathbb{R}^n$
- Connected subspaces to matrices

Keywords

subspace

closed under addition

closed under scaling

column space


null space

basis

Subspaces


Subspaces

Subspaces



Subspaces


A plane in $\mathbb{R}^3$ looks like a tilted copy of $\mathbb{R}^2$



Subspaces

A plane in $\mathbb{R}^3$ looks like a tilted copy of $\mathbb{R}^2$

Subspaces *generalize* this




Subspaces

A plane in $\mathbb{R}^3$ looks like a tilted copy of $\mathbb{R}^2$

Subspaces *generalize* this

We can look at "tilted copies" of $\mathbb{R}^7$ sitting in $\mathbb{R}^{11}$



Subspaces (Algebraic Definition)

subspace has to "live" somewhere!

Df. Nonempty $H \subseteq \mathbb{R}^n$ is a subspace of $\mathbb{R}^n$

(1) if $\vec{u}, \vec{v} \in H$ then $\vec{u}+\vec{v} \in H$ (closure under addition)

(2) if $\vec{u} \in H$ then $c\vec{u} \in H$ (closure under scaling)

Note. $\vec{u}_1, \dots \vec{u}_k \in H \Rightarrow \sum_{i=1}^{k} c_i \vec{u}_k \in H$ (closure under lin comb.)

$\vec{0} \overset{?}{\in} H$ $\quad v \in H \Rightarrow 0\vec{v} \in H$

$0\vec{v} = \vec{0}$

Subspace (Algebraic Definition)

Df. A **subspace** of $\mathbb{R}^n$ is a set H of vectors in $\mathbb{R}^n$ such that

1. for every $\mathbf{u}$ and $\mathbf{v}$ in H, the vector $\mathbf{u}+\mathbf{v}$ is in H

2. for every $\mathbf{u}$ in H and scalar c, the vector $c\mathbf{u}$ is in H

Subspace (Algebraic Definition)

Df. A **subspace** of $\mathbb{R}^n$ is a set H of vectors in $\mathbb{R}^n$ such that

1. for every $\mathbf{u}$ and $\mathbf{v}$ in H, the vector $\mathbf{u}+\mathbf{v}$ is in H — H **is closed under addition**

2. for every $\mathbf{u}$ in H and scalar c, the vector $c\mathbf{u}$ is in H — H **is closed under scaling**

Subspace (Algebraic Definition)

Df. A **subspace** of $\mathbb{R}^n$ is a set H of vectors in $\mathbb{R}^n$ such that

1. for every $\mathbf{u}$ and $\mathbf{v}$ in H, the vector $\mathbf{u}+\mathbf{v}$ is in H — H **is closed under addition**


2. for every $\mathbf{u}$ in H and scalar c, the vector $c\mathbf{u}$ is in H — H **is closed under scaling**

!! Subspaces must "live" somewhere !!

How to Think About this

It's not possible to "leave" H by addition or scaling

(recall this is also how we discussed spans)



https://textbooks.math.gatech.edu/ila/spans.html

How To: Verifying Subspaces

How To: Verifying Subspaces

Q. Verify that H is a subspace of $\mathbb{R}^n$

How To: Verifying Subspaces

Q. Verify that H is a subspace of $\mathbb{R}^n$

A.

1. Show if $\mathbf{u}$ and $\mathbf{v}$ are in H then so is $\mathbf{u} + \mathbf{v}$

How To: Verifying Subspaces

Q. Verify that H is a subspace of $\mathbb{R}^n$

A.

1. Show if $\mathbf{u}$ and $\mathbf{v}$ are in H then so is $\mathbf{u} + \mathbf{v}$
2. Show if $\mathbf{u}$ is in H then so is $c\mathbf{u}$ for any scalar c

How To: Verifying Non-Subspaces

How To: Verifying Non-Subspaces

Q. Verify that H is *not* a subspace of $\mathbb{R}^n$

How To: Verifying Non-Subspaces

Q. Verify that H is *not* a subspace of $\mathbb{R}^n$

A.

Find $\mathbf{u}$ and $\mathbf{v}$ in H where $\mathbf{u} + \mathbf{v}$ is not in H

How To: Verifying Non-Subspaces

Q. Verify that H is *not* a subspace of $\mathbb{R}^n$


A.

Find $\mathbf{u}$ and $\mathbf{v}$ in H where $\mathbf{u}+\mathbf{v}$ is not in H

or

Find $\mathbf{u}$ in H where $c\mathbf{u}$ is not in H for *some* scalar c

Example: Lines in $\mathbb{R}^2$




Example: Lines in $\mathbb{R}^2$



Subspaces must include the origin

Fact. For any subspace H of $\mathbb{R}^n$, the zero vector is in H. In set notation: $\mathbf{0} \in H$



(this can also be used to verify non-subspace-ness)

Example: The Origin

H

Fact. The set $\{\mathbf{0}_n\}$ is a subspace of $\mathbb{R}^n$ ✓

$\vec{u}, \vec{v} \in H \Rightarrow \vec{u} = \vec{0}, \vec{v} = \vec{0} \Rightarrow \vec{u} + \vec{v} = \vec{0} + \vec{0} = \vec{0} \in H$

(closed under add.

ex. closure under scaling

Example: $\mathbb{R}^n$

Fact. The set $\mathbb{R}^n$ is a subspace of $\mathbb{R}^n$

$\vec{u}, \vec{v} \in \mathbb{R}^n \implies \vec{u} + \vec{v} \in \mathbb{R}^n$

$\vec{u} \in \mathbb{R}^n \implies c\vec{u} \in \mathbb{R}^n$

Example: Spans

Fact. For any set of vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$ in $\mathbb{R}^n$ the set $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n\}$ is a subspace of $\mathbb{R}^n$

H

$\vec{u}, \vec{w} \in H$

$$\vec{u} = \sum_{i=1}^{n} \alpha_i \vec{v}_i \qquad \vec{w} = \sum_{i=1}^{n} \beta_i \vec{v}_i$$

$$\vec{u} + \vec{w} = \sum_{i=1}^{n} \alpha_i \vec{v}_i + \sum_{i=1}^{n} \beta_i \vec{v}_i = \sum_{i=1}^{n} (\alpha_i \vec{v}_i + \beta_i \vec{v}_i)$$

Still lin. combo. (closure under add)

$$= \boxed{\sum_{i=1}^{n} (\alpha_i + \beta_i) \vec{v}_i} \in H$$

(ex. closure under scaling)

Subspace in $\mathbb{R}^3$ (Geometrically)


There are only 4 kinds of subspaces of $\mathbb{R}^3$:

1. $\{\mathbf{0}\}$ just the origin

2. lines (through the origin)

3. planes (through the origin)

4. All of $\mathbb{R}^3$




https://commons.wikimedia.org/wiki/File:Linear_subspaces_with_shading.svg

Subspace in $\mathbb{R}^3$ (Geometrically)

There are only 4 kinds of subspaces of $\mathbb{R}^3$:

1. $\{\mathbf{0}\}$ just the origin
2. lines (through the origin)
3. planes (through the origin)
4. All of $\mathbb{R}^3$




https://commons.wikimedia.org/wiki/File:Linear_subspaces_with_shading.svg

Subspace in $\mathbb{R}^3$ (Geometrically)

There are only 4 kinds of subspaces of $\mathbb{R}^3$:

1. $\{\mathbf{0}\}$ just the origin
2. lines (through the origin)
3. planes (through the origin)
4. All of $\mathbb{R}^3$




https://commons.wikimedia.org/wiki/File:Linear_subspaces_with_shading.svg

Subspace in $\mathbb{R}^3$ (Geometrically)

There are only 4 kinds of subspaces of $\mathbb{R}^3$:

1. $\{\mathbf{0}\}$ just the origin
2. lines (through the origin)
3. planes (through the origin)
4. All of $\mathbb{R}^3$




https://commons.wikimedia.org/wiki/File:Linear_subspaces_with_shading.svg

Subspace in $\mathbb{R}^3$ (Geometrically)

There are only 4 kinds of subspaces of $\mathbb{R}^3$:

1. $\{\mathbf{0}\}$ just the origin
2. lines (through the origin)
3. planes (through the origin)
4. All of $\mathbb{R}^3$



https://commons.wikimedia.org/wiki/File:Linear_subspaces_with_shading.svg

Non-Example: Bounded Sets

H


Fact. The set $\{(x, y) : x \geq 3\}$ is *not* a subspace of $\mathbb{R}^2$

$\begin{bmatrix} 0 \\ 0 \end{bmatrix} \notin H$

$H' = \{(x, y) : x \geq -4\}$

$\begin{bmatrix} -1 \\ 0 \end{bmatrix} \in H' \Rightarrow 5\begin{bmatrix} -1 \\ 0 \end{bmatrix} = \begin{bmatrix} -5 \\ 0 \end{bmatrix} \notin H'$

H' is <u>not</u> a subspace



https://brainly.com/question/14147114

Question

1. Show that the unit sphere $\{(x, y, z) : x^2 + y^2 + x^2 = 1\}$ is <u>not</u> a subspace of $\mathbb{R}^3$

also consider ≤ 1

2. Show that the range of a linear transformation $T : \mathbb{R}^m \to \mathbb{R}^n$ is a subspace of $\mathbb{R}^n$

Answer (1)

Answer (2)

$T(\vec{v}) = A\vec{v}$

$\text{ran}(T) = \text{span}\{\vec{a}_1, \dots, \vec{a}_n\} = [\vec{a}_1 \cdots \vec{a}_n]$ ✓

and spans are subspaces $\in \text{span}\{\vec{a}_1, \dots, \vec{a}_n\}$

closure under add: (ex. closure und scaling)

$\vec{u}, \vec{w} \in \text{ran}(T)$

$T(\vec{v}_1) = \vec{u}$


$T(\vec{v}_2) = \vec{w}$

$\vec{u} + \vec{w} =$

$T(\vec{v}_1) + T(\vec{v}_2) =$

$T(\vec{v}_1 + \vec{v}_2)$

$\vec{u} + \vec{w} \in \text{ran}(T)$



How To: Subspaces and Span

Q. Show that $\mathbf{v}$ lies in the subspace generated by $\mathbf{u}_1, \ldots, \mathbf{u}_k$

A. Show that $\mathbf{v}$ is in $\text{span}\{\mathbf{u}_1, \ldots, \mathbf{u}_k\}$

We will start using "subspace generated by" and "span of" interchangeably

Subspaces and Matrices

The Connection


Since matrices can be viewed as:

- collections of vectors
- implementing linear transformations

they have many associated subspaces

Today we'll look at:

- column space
- null space



https://mathworld.wolfram.com/FundamentalMatrixSubspaces.html

Column Space

Column Space

Df. The **column space** of a matrix A, written $\text{Col}\, A$, is the set of all linear combinations of the columns of A

Column Space

Df. The **column space** of a matrix A, written $\text{Col}\, A$, is the set of all linear combinations of the columns of A

Df. *The column space of a matrix is the span of its columns*

Column Space

Df. The **column space** of a matrix A, written $\text{Col } A$, is the set of all linear combinations of the columns of A

Df. *The column space of a matrix is the span of its columns*

Df. *The column space of a matrix is the range of the linear transformation it implements*

Subspace of What?

$A \in \mathbb{R}^{m \times n}$

$T \quad \vec{x} \mapsto A\vec{x} : \mathbb{R}^n \to \mathbb{R}^m$

$\mathbb{R}^m$ = codomain of T

$$m \left| \begin{bmatrix} | & | & \dots & | & | \\ \mathbf{a}_1 & \mathbf{a}_2 & \dots & \mathbf{a}_{n-1} & \mathbf{a}_n \\ | & | & \dots & | & | \end{bmatrix} \right. \quad n$$

$\mathrm{Col}(A)$

is a subspace of

$\mathbb{R}^m$

$c_1\mathbf{a}_1 + c_2\mathbf{a}_2 + \dots c_n\mathbf{a}_n$ **is a vector in** $\mathbb{R}^m$

Examples

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 0 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & -1 \end{bmatrix}$$

$a_1 \quad a_2 \quad a_3$

$$B = \begin{bmatrix} 1 & -1 & 4 \\ 2 & -2 & 5 \\ 3 & -3 & 6 \end{bmatrix}$$

$b_1 \quad b_2 \quad b_3$

$\text{Col } A = \text{span}\{a_1, a_2, a_3\} = \mathbb{R}^3$

$\Updownarrow$

$\vec{e}_1, \vec{e}_2, \vec{e}_3 \in \text{span}\{a_1, a_2, a_3\}$

$\text{Col } B = \text{span}\{b_1, b_2, b_3\} = \text{span}\{b_1, b_3\} \neq \mathbb{R}^3$

Null Space

$A \in \mathbb{R}^{m \times n}$ sol. set of $A\vec{x} = \vec{0}$

$N = \{\vec{v} : A\vec{v} = \vec{0}\}$

Claim. N is a subspace

$\vec{u}, \vec{v} \in N$, $A(\vec{u} + \vec{v}) = A\vec{u} + A\vec{v} = \vec{0} + \vec{0} = \vec{0}$

$\Rightarrow \vec{u} + \vec{v} \in N$ ✓ (closure under add.)

$\vec{u} \in N$, $A(c\vec{u}) = cA\vec{u} = c\vec{0} = \vec{0}$ ✓ (closure under scaling)

Null Space

Df. The **null space** of a matrix A, written $\text{Nul}\, A$, is the set of all solutions to the homogenous equation $A\mathbf{x} = \mathbf{0}$

Null Space

Df. The **null space** of a matrix A, written $\text{Nul}\, A$, is the set of all solutions to the homogenous equation $A\mathbf{x} = \mathbf{0}$

The null space of a matrix A is the set of all vectors that are mapped to the zero vector by A

Subspace of What?

$$\underset{m \text{ rows}}{\overset{n \text{ columns}}{A}}\mathbf{v} = \mathbf{0}$$

$m \times n \quad n \times 1 \qquad\qquad m \times 1$

$\mathbf{v}$ **is a vector in** $\mathbb{R}^n$

$$\text{Nul}(A)$$

is a subspace of

$$\mathbb{R}^n$$

The Null Space is a Subspace

Fact. For any $m \times n$ matrix A, the set $\text{Nul}(A)$ is a subspace of $\mathbb{R}^n$


Examples

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 0 & 0 \end{bmatrix} \qquad B = \begin{bmatrix} 1 & -1 & 4 \\ 2 & -2 & 5 \\ 3 & -3 & 6 \end{bmatrix}$$

$\mathrm{Nul}\,A = \{\vec{0}\}$ (by IMT)

$$\mathrm{Nul}(B) = \left\{ c\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} : c \in \mathbb{R} \right\} = \mathrm{span}\left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$$


Linear Transformations Perspective



Linear Algebra and its Applications (Lay, Lay, McDonald)

Linear Transformations Perspective

If A implements T then




Linear Algebra and its Applications (Lay, Lay, McDonald)

Linear Transformations Perspective

If A implements T then

» $\text{Col}(A)$ is the same as $\text{ran}(T)$, where vectors are "sent" by T




Linear Algebra and its Applications (Lay, Lay, McDonald)

Linear Transformations Perspective

If A implements T then

- $\text{Col}(A)$ is the same as $\text{ran}(T)$, where vectors are "sent" by T
- $\text{Nul}(A)$ is the set of vectors that are "zeroed out" by T, also called the **kernel** of T



Linear Algebra and its Applications (Lay, Lay, McDonald)

Bases

The idea behind a basis

The idea behind a basis

We've already said spans are subspaces, but the *converse* true too

The idea behind a basis

We've already said spans are subspaces, but the *converse* true too

Every subspace is the span of a set of vectors

The idea behind a basis

We've already said spans are subspaces, but the *converse* true too

Every subspace is the span of a set of vectors


A basis is a "minimal" choice of these vectors, a "concise representation" of a subspace

Recall: Standard Basis

Df. The *n-dimensional standard basis vectors* (or standard coordinate vectors) are the vectors $\mathbf{e}_1, \ldots, \mathbf{e}_n$ where

$$\mathbf{e}_i = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix} \begin{matrix} 1 \\ 2 \\ \vdots \\ i-1 \\ i \\ i+1 \\ \vdots \\ n-1 \\ n \end{matrix}$$


What's special about the standard basis?



What's special about the standard basis?

The n standard basis vectors in $\mathbb{R}^n$:

- are linearly independent
- span all of $\mathbb{R}^n$



What's special about the standard basis?


$$\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = a_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + a_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + a_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

The n standard basis vectors in $\mathbb{R}^n$:

- are linearly independent
- span all of $\mathbb{R}^n$

Their span is as "large" as possible while the set of vectors generating the span is as "small" as possible



Basis

Basis

Df. A **basis** for a subspace H of $\mathbb{R}^n$ is a linearly independent set $\{\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k\}$ of vectors that spans H

Basis

Df. A **basis** for a subspace H of $\mathbb{R}^n$ is a linearly independent set $\{\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k\}$ of vectors that spans H

A basis is a *minimal* set of vectors which spans all of H

Example: Standard basis

The standard basis is a basis of $\mathbb{R}^n$

$$\begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} = v_1\mathbf{e}_1 + v_2\mathbf{e}_2 + \ldots + v_n\mathbf{e}_n$$

Column vectors are just weights for a linear combination of the standard basis

Example: Column Space of Invertible Matrices

Fact. The columns of an invertible $n \times n$ matrix form a basis of $\mathbb{R}^n$

$A \sim I$

Subsets of Spanning Sets

Subsets of Spanning Sets

Thm. If the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ span a subspace H then a subset of them form a basis of H

Subsets of Spanning Sets

Thm. If the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ span a subspace H then a subset of them form a basis of H

We can *remove* vectors from a spanning set until we get a basis

Subsets of Spanning Sets

Thm. If the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ span a subspace H then a subset of them form a basis of H

We can *remove* vectors from a spanning set until we get a basis

How do we do this?

Subsets of Spanning Sets

Thm. If the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ span a subspace H then a subset of them form a basis of H

We can *remove* vectors from a spanning set until we get a basis

How do we do this?

As usual, by connecting back to matrices

Question

$$\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ -3 \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ 3 \end{bmatrix} \right\}$$

Is this set of vectors a basis for $\mathbb{R}^3$*?*

Bases of Column Space and Null Space

The Goal of this Last Section

Determine how to find bases for the **column space** and the **null space** of a given matrix

How to: Finding a basis for the null space

Q. Given a $m \times n$ matrix A find a basis for $\text{Nul}(A)$

How to: Finding a basis for the null space

Q. Given a $m \times n$ matrix A find a basis for $\mathrm{Nul}(A)$

The idea. Describe the solutions of $Ax = 0$ as linear combination of vectors

Example

$$A \sim \begin{bmatrix} 1 & -2 & 0 & -1 & 3 \\ 0 & 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Let's write down a general form solution for A:

$x_1 = 2x_2 + x_4 - 3x_5$

x_2 is free

$x_3 = -2x_4 + 2x_5$

x_4 is free

x_5 is free

Parametric Solutions

We can think of our general form solution as a *linear transformation*

$x_1 = 2x_2 + x_4 - 3x_5$

x_2 is free

$x_3 = (-2)x_4 + 2x_5$

x_4 is free

x_5 is free

$\equiv$

"given values for x_2, x_3, and x_4, I can give you a solution"

Parametric Solutions

We can think of our general form solution as a *linear transformation*

$x_1 = 2x_2 + x_4 - 3x_5$

x_2 is free

$x_3 = (-2)x_4 + 2x_5$

x_4 is free

x_5 is free

$\equiv$

$$\begin{bmatrix} s \\ t \\ u \end{bmatrix} \mapsto \begin{bmatrix} 2s + t - 3u \\ s \\ (-2)t + 2u \\ t \\ u \end{bmatrix}$$

Parametric Solutions

We can think of our general form solution as a *linear transformation*

!! this transformation is only linear !!
!! in the case of homogeneous equations !!

$x_1 = 2x_2 + x_4 - 3x_5$

x_2 is free

$x_3 = (-2)x_4 + 2x_5$

x_4 is free

x_5 is free

$$\equiv \quad \begin{bmatrix} s \\ t \\ u \end{bmatrix} \mapsto \begin{bmatrix} 2s + t - 3u \\ s \\ (-2)t + 2u \\ t \\ u \end{bmatrix}$$

Example

$$\begin{bmatrix} s \\ t \\ u \end{bmatrix} \mapsto \begin{bmatrix} 2s + t - 3u \\ s \\ (-2)t + 2u \\ t \\ u \end{bmatrix}$$

Let's find the matrix implementing this transformation:


$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \qquad \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 1 \\ 0 \\ -2 \\ 1 \\ 0 \end{bmatrix} \qquad \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \mapsto \begin{bmatrix} -3 \\ 0 \\ 2 \\ 0 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 & -3 \\ 1 & 0 & 0 \\ 0 & -2 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Example

$$\begin{bmatrix} 2 & 1 & -3 \\ 1 & 0 & 0 \\ 0 & -2 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Thm. The columns of this matrix are linearly independent



Example

$$\begin{bmatrix} 2 & 1 & -3 \\ 1 & 0 & 0 \\ 0 & -2 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Example

$$\begin{bmatrix} 2 & 1 & -3 \\ 1 & 0 & 0 \\ 0 & -2 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

The columns of this matrix *span* $\text{Nul}(A)$

Example

$$\begin{bmatrix} 2 & 1 & -3 \\ 1 & 0 & 0 \\ 0 & -2 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

The columns of this matrix *span* $\text{Nul}(A)$

The columns of this matrix are linearly independent

Example

$$\begin{bmatrix} 2 & 1 & -3 \\ 1 & 0 & 0 \\ 0 & -2 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

The columns of this matrix *span* $\text{Nul}(A)$

The columns of this matrix are linearly independent

The columns of this matrix form a basis for $\text{Nul}(A)$

Example

Alternatively, we can write a general form solution so that it is a linear combination of vectors with *free variables as weights*

$$x_1 = 2x_2 + x_4 - 3x_5$$
x_2 is free
$$x_3 = (-2)x_4 + 2x_5$$
x_4 is free
x_5 is free

$$x_2 \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} 1 \\ 0 \\ -2 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -3 \\ 0 \\ 2 \\ 0 \\ 1 \end{bmatrix}$$

How to: Finding a basis for the null space

Q. Given a $m \times n$ matrix A find a basis for $\text{Nul}(A)$

A.

1. Find a general form solution for $A\mathbf{x} = \mathbf{0}$
2. Write this solution as a linear combination of vectors *where the free variables are the weights*
3. The resulting vectors form a basis for $\text{Nul}(A)$

An Observation

The *number* of vectors in the basis we found is the same as the number of *free variables* in a general form solution

$x_1 = 2x_2 + x_4 - 3x_5$

x_2 is free

$x_3 = (-2)x_4 + 2x_5$

x_4 is free

x_5 is free

$\equiv$

$$\begin{bmatrix} s \\ t \\ u \end{bmatrix} \mapsto \begin{bmatrix} 2s + t - 3u \\ s \\ (-2)t + 2u \\ t \\ u \end{bmatrix}$$

Practice Problem

$$A \sim \begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{bmatrix}$$

Determine a basis for $\text{Nul}(A)$

Answer

$$\begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{bmatrix}$$

onto column space...

How To: Finding a basis for the column space

How To: Finding a basis for the column space

Question. Given a $m \times n$ matrix A, find a basis for $\text{Col}(A)$

How To: Finding a basis for the column space

Question. Given a $m \times n$ matrix A, find a basis for $\text{Col}(A)$

We already know the columns of A span $\text{Col}(A)$

How To: Finding a basis for the column space

Question. Given a $m \times n$ matrix A, find a basis for $\text{Col}(A)$

We already know the columns of A span $\text{Col}(A)$

So we also already know *some* subset of columns of A form a basis for $\text{Col}(A)$

How To: Finding a basis for the column space

Question. Given a $m \times n$ matrix A, find a basis for $\text{Col}(A)$

We already know the columns of A span $\text{Col}(A)$

So we also already know *some* subset of columns of A form a basis for $\text{Col}(A)$

Which vectors should we choose?

Column Space and Reduced Echelon form


$$[\mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_3 \quad \mathbf{a}_4 \quad \mathbf{a}_5] \sim \begin{bmatrix} 1 & -2 & 0 & -1 & 3 \\ 0 & 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Column Space and Reduced Echelon form

$$[\mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_3 \quad \mathbf{a}_4 \quad \mathbf{a}_5] \sim \begin{bmatrix} 1 & -2 & 0 & -1 & 3 \\ 0 & 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The idea. What if we cover up the non-pivot columns?


Column Space and Reduced Echelon form



The idea. What if we cover up the non-pivot columns?

Then we see $[\mathbf{a}_1 \ \ \mathbf{a}_3]$ has 2 pivots

Column Space and Reduced Echelon form



The idea. What if we cover up the non-pivot columns?


Then we see $[\mathbf{a}_1 \quad \mathbf{a}_3]$ has 2 pivots

So the pivot columns are *linearly independent*

Column Space and Reduced Echelon form


$$[\mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_3 \quad \mathbf{a}_4 \quad \mathbf{a}_5] \sim \begin{bmatrix} 1 & -2 & 0 & -1 & 3 \\ 0 & 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Column Space and Reduced Echelon form



Observation. $[2\ 1\ 0\ 0\ 0]^T$ is a solution to the system $A\mathbf{x} = \mathbf{0}$

Column Space and Reduced Echelon form



Observation. $[2\ 1\ 0\ 0\ 0]^T$ is a solution to the system $A\mathbf{x} = \mathbf{0}$

So $2\mathbf{a}_1 + \mathbf{a}_2 = \mathbf{0}$ and $\mathbf{a}_2 = (-2)\mathbf{a}_1$

Column Space and Reduced Echelon form

$$[\mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_3 \quad \mathbf{a}_4 \quad \mathbf{a}_5] \sim \begin{bmatrix} 1 & -2 & 0 & -1 & 3 \\ 0 & 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Observation. $[2\ 1\ 0\ 0\ 0]^T$ is a solution to the system $A\mathbf{x} = \mathbf{0}$

So $2\mathbf{a}_1 + \mathbf{a}_2 = \mathbf{0}$ and $\mathbf{a}_2 = (-2)\mathbf{a}_1$

In general, every non-pivot column of A can be written as a linear combination pivots in front of it

Column Space and Reduced Echelon form

$$[\mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_3 \quad \mathbf{a}_4 \quad \mathbf{a}_5] \sim \begin{bmatrix} 1 & -2 & 0 & -1 & 3 \\ 0 & 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Observation. $[2\ 1\ 0\ 0\ 0]^T$ is a solution to the system $A\mathbf{x} = \mathbf{0}$

So $2\mathbf{a}_1 + \mathbf{a}_2 = \mathbf{0}$ and $\mathbf{a}_2 = (-2)\mathbf{a}_1$

In general, every non-pivot column of A can be written as a linear combination pivots in front of it

This tells us that $\mathbf{a}_1$ and $\mathbf{a}_3$ *span* Col(A)

Column Space and Reduced Echelon form

$$[\mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_3 \quad \mathbf{a}_4 \quad \mathbf{a}_5] \sim \begin{bmatrix} 1 & -2 & 0 & -1 & 3 \\ 0 & 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Column Space and Reduced Echelon form

$$[\mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_3 \quad \mathbf{a}_4 \quad \mathbf{a}_5] \sim \begin{bmatrix} 1 & -2 & 0 & -1 & 3 \\ 0 & 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The takeaway. The pivot columns of A form a basis for $\mathrm{Col}(A)$

Column Space and Reduced Echelon form

$$[\mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_3 \quad \mathbf{a}_4 \quad \mathbf{a}_5] \sim \begin{bmatrix} 1 & -2 & 0 & -1 & 3 \\ 0 & 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The takeaway. The pivot columns of A form a basis for $\text{Col}(A)$

!! IMPORTANT !!

Choose the columns of A

($\mathbf{e}_1$ and $\mathbf{e}_2$ do not necessarily form a basis for $\text{Col}(A)$)

How To: Finding a basis for the column space

Q. Given a $m \times n$ matrix A, find a basis for $\text{Col}(A)$

A.

1. Find the pivot columns in an echelon form of A
2. The associated columns <u>in A</u> form a basis for $\text{Col}(A)$

Question

$$A = \begin{bmatrix} 1 & -2 & 19 & 0 & -4 \\ 1 & 0 & 9 & 1 & 1 \\ 1 & -1 & 14 & 1 & -1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 9 & 0 & 0 \\ 0 & 1 & -5 & 0 & 2 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}$$

Find a bases for the column space and null space of A

Answer

$$A = \begin{bmatrix} 1 & -2 & 19 & 0 & -4 \\ 1 & 0 & 9 & 1 & 1 \\ 1 & -1 & 14 & 1 & -1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 9 & 0 & 0 \\ 0 & 1 & -5 & 0 & 2 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}$$

Summary

Subspaces define "tilted versions" of $\mathbb{R}^k$ in $\mathbb{R}^n$ (where $k \leq n$)

Bases are compact representation of subspaces as minimal spanning sets

Matrices have useful associated subspaces like the column space and null space