

BOUNDARY RIGIDITY AND STABLE LEFT–RIGHT MODULI AT GENERIC DEGREE THREE

NATHANIEL MONSON

ABSTRACT. We exhibit a one-parameter family of polynomial Keller maps $\mathbb{A}_{\mathbb{C}}^3 \rightarrow \mathbb{A}_{\mathbb{C}}^3$, all of generic degree three, whose members remain pairwise inequivalent after adjoining arbitrarily many identity variables and applying arbitrary polynomial changes of source and target coordinates. The family is obtained from the quadratic cubic-frame data

$$A_{\alpha}(c) = c + \alpha c^2, \quad B_{\alpha,\beta}(c) = -2 - 4\alpha c + \beta c^2.$$

For $\alpha \neq 0$, the complete stable left–right invariant is

$$q = \beta/\alpha^2.$$

The line $\alpha = 0$ is a single ordinary left–right orbit, stably distinct from every $\alpha \neq 0$ orbit.

The invariant is recovered from the nonproperness divisor. After normalizing its singular discriminant component, one obtains an affine plane with two intrinsically marked curves. A unique-factorization argument shows that every automorphism of every affine cylinder preserving this marked pair fixes q . A separate singular-incidence argument treats the exceptional value $q = -2$. Exact symbolic calculations verify the discriminant, normalization, conductor, exceptional fiber, and the explicit gauge transformation on the line $\alpha = 0$.

More generally, on the squarefree coprime cubic-frame locus, the complete stable invariant is the deleted infinity scheme $V(A/c)$ decorated by the transverse derivative $B|_{V(A/c)}$. Stable and ordinary left–right equivalence coincide there. This yields stable-moduli spaces of arbitrarily large dimension, all at generic degree three and, for each family, fixed ordinary degree.

Working manuscript. The exact symbolic verifier has been rerun from the recovered source artifact. Independent implementation and specialist review remain submission requirements.

1. INTRODUCTION

The recent counterexample to the Jacobian conjecture gives a polynomial map $\mathbb{A}_{\mathbb{C}}^3 \rightarrow \mathbb{A}_{\mathbb{C}}^3$ with nonzero constant Jacobian determinant that is not an automorphism [1, 6]. We credit Akhil Mathew as the primary human source of the problem: Mathew prompted Levent Alpöge, Alpöge prompted Fable, Fable produced the example, and Alpöge announced it publicly [7]. The construction was quickly placed in larger families. In particular, public

Date: July 22, 2026.

accounts allow arbitrary one-variable deformation data and produce counterexamples of every generic degree [7, 3]. Varying generic degree already gives inequivalent maps.

This paper asks a narrower question:

Can a family of counterexamples of one fixed generic degree carry a modulus that survives arbitrary polynomial source and target changes, even after stabilization?

The answer is yes already at generic degree three.

Let

$$F, H: \mathbb{A}^n \longrightarrow \mathbb{A}^n$$

be polynomial maps. They are *polynomially left–right equivalent* if there are polynomial automorphisms Φ, Ψ of $\mathbb{A}^n$ such that

$$H \circ \Phi = \Psi \circ F.$$

They are *stably polynomially left–right equivalent* if, for some $m \geq 0$, the maps

$$F \times \text{id}_{\mathbb{A}^m}, \quad H \times \text{id}_{\mathbb{A}^m}$$

are polynomially left–right equivalent. We use this definition throughout; “stable equivalence” has other meanings in the Jacobian-conjecture literature.

For $\alpha, \beta \in \mathbb{C}$, consider

$$A_\alpha(C) = C + \alpha C^2, \quad B_{\alpha, \beta}(C) = -2 - 4\alpha C + \beta C^2. \quad (1.1)$$

These polynomials define a Keller map $G_{\alpha, \beta}$ by the cubic-frame construction in section 2. Every member has generic degree three and is nonproper. Our main theorem classifies the entire two-parameter family.

Theorem 1.1 (Complete stable classification). *The stable polynomial left–right orbit set of the family $\{G_{\alpha, \beta}\}$ is*

$$\{\mathcal{O}_0\} \sqcup \{\mathcal{O}_q : q \in \mathbb{C}\},$$

where

$$\mathcal{O}_0 = \{G_{0, \beta} : \beta \in \mathbb{C}\}$$

and

$$\mathcal{O}_q = \{G_{\alpha, \beta} : \alpha \neq 0, \beta/\alpha^2 = q\}.$$

More explicitly:

- (i) all maps $G_{0, \beta}$ are ordinarily left–right equivalent;
- (ii) no $G_{0, \beta}$ is stably equivalent to a member with $\alpha \neq 0$; and
- (iii) if $\alpha, \alpha' \neq 0$, then

$$G_{\alpha, \beta} \sim_{\text{stable}} G_{\alpha', \beta'} \iff \frac{\beta}{\alpha^2} = \frac{\beta'}{\alpha'^2}.$$

The weighted-scaling invariance of β/α^2 is immediate. The content of theorem 1.1 is completeness under *arbitrary* polynomial left–right equivalence and after arbitrary affine stabilization.

Our invariant comes from the set at which the map is not proper. For a dominant equidimensional polynomial map this set, if nonempty, is a hypersurface by Jelonek’s theorem [4]. In our normalized one-parameter slice it has a singular discriminant component D_q and a plane component P . The normalization of D_q is an affine plane containing two intrinsic marked curves:

$$L_q : 3c(c+1)t + qc^2 - 4c - 2 = 0, \quad M : c+1 = 0.$$

The pair (L_q, M) remembers q , even after taking a product with affine space. This gives the necessity in theorem 1.1.

The existence of left–right moduli for nonproper étale maps on other affine varieties has important precedent: Dubouloz and Palka constructed high-dimensional families on affine pseudo-planes [2]. Our theorem concerns polynomial self-maps of affine space arising directly from the new three-dimensional construction. The general marked-root framework and its unstabilized equivalence criterion are developed in the companion paper [5]. The proof below includes all of the special geometry needed for stable separation in the quadratic slice.

2. THE QUADRATIC CUBIC-FRAME FAMILY

Put

$$c = 2x - 3x^2y - x^3z, \quad t = y + \frac{1}{x}, \quad r = \frac{2}{x}. \quad (2.1)$$

For $A = A_\alpha$ and $B = B_{\alpha,\beta}$, let

$$h(T, C) = A(C)T^3 + B(C)T^2$$

and define

$$b = r - \frac{\partial h}{\partial T}(t, c), \quad 2a = h(t, c) + tb. \quad (2.2)$$

Although (2.1) uses $1/x$, the two expressions in (2.2) are polynomials. We write

$$G_{\alpha,\beta} = (a, b, c) : \mathbb{A}_{x,y,z}^3 \longrightarrow \mathbb{A}_{a,b,c}^3.$$

Proposition 2.1. *For every $\alpha, \beta \in \mathbb{C}$, the map $G_{\alpha,\beta}$ is polynomial,*

$$\det DG_{\alpha,\beta} = -2,$$

and its generic degree is three.

Proof. Set

$$\xi = 1 + xy, \quad d = 2 - 3xy - x^2z,$$

so that $t = \xi/x$ and $c = xd$. Clearing denominators in b gives

$$b = \frac{2 + 4\xi - 3d\xi^2}{x} - 3\alpha d^2\xi^2 + 8\alpha d\xi - 2\beta xd^2\xi.$$

The numerator of the displayed fraction vanishes at $x = 0$, hence is divisible by x . Similarly,

$$2a = \frac{2\xi - 2d\xi^3 + 2\xi^2}{x^2} + \frac{4\alpha d\xi^2 - 2\alpha d^2\xi^3}{x} - \beta d^2\xi^2.$$

The first numerator is divisible by x^2 , and the second by x ; this follows directly on substituting $\xi = 1 + xy$ and $d = 2 - 3xy - x^2z$. Thus a and b are polynomial.

On $x \neq 0$, the coordinate changes in (2.1) give

$$\det \frac{\partial(a, b, c)}{\partial(t, r, c)} = \frac{r}{2}, \quad \det \frac{\partial(t, r, c)}{\partial(x, y, z)} = -2x.$$

Since $r = 2/x$, their product is -2 . Polynomiality extends this identity across $x = 0$.

For a target point (a, b, c) , introduce

$$P_{a,b,c}(T) = A(c)T^3 + B(c)T^2 + bT - 2a. \quad (2.3)$$

The construction gives

$$P_{G_{\alpha,\beta}(x,y,z)}(t) = 0, \quad P'_{G_{\alpha,\beta}(x,y,z)}(t) = \frac{2}{x}.$$

Conversely, every finite simple root t reconstructs

$$x = \frac{2}{P'(t)}, \quad y = t - \frac{P'(t)}{2}, \quad z = \frac{2x - 3x^2y - c}{x^3}. \quad (2.4)$$

A general cubic (2.3) has three finite simple roots, giving three source points. $\square$

Remark 2.2. The determinant in proposition 2.1 is -2 , rather than 1, only because of our target normalization. A linear target rescaling gives the usual Keller normalization.

2.1. The ordinary scaling orbit. Suppose $\alpha, \alpha' \neq 0$, and put

$$u = \frac{\alpha}{\alpha'}.$$

If

$$\frac{\beta}{\alpha^2} = \frac{\beta'}{\alpha'^2},$$

then direct substitution gives

$$G_{\alpha',\beta'}\left(ux, \frac{y}{u}, \frac{z}{u^2}\right) = \left(\frac{a}{u^2}, \frac{b}{u}, uc\right) \quad \text{when} \quad (a, b, c) = G_{\alpha,\beta}(x, y, z). \quad (2.5)$$

Thus equality of $q = \beta/\alpha^2$ is sufficient for ordinary equivalence. Every nonzero- α orbit therefore has the normalized representative

$$G_q := G_{1,q},$$

with

$$A(c) = c(c+1), \quad B_q(c) = qc^2 - 4c - 2. \quad (2.6)$$

3. THE INTRINSIC NONPROPERNESS DIVISOR

For a dominant polynomial map F , write S_F for its reduced nonproperness set. Polynomial source automorphisms leave S_F unchanged, target automorphisms carry it isomorphically to the new nonproperness set, and

$$S_{F \times \text{id}_{\mathbb{A}^m}} = S_F \times \mathbb{A}^m. \quad (3.1)$$

The last identity follows either from the valuative criterion or directly from sequences escaping to infinity.

The discriminant of the normalized fiber cubic

$$P_{a,b,c}(T) = A(c)T^3 + B_q(c)T^2 + bT - 2a$$

is

$$\Delta_q = B_q^2 b^2 - 4Ab^3 + 8B_q^3 a - 108A^2 a^2 - 36AB_q ab. \quad (3.2)$$

For $q \neq -2$, let

$$D_q = V(\Delta_q).$$

For $q = -2$, the polynomial Δ_{-2} has content $c + 1$; in that case put

$$D_{-2} = V\left(\frac{\Delta_{-2}}{c+1}\right).$$

In every case let

$$P = V(c+1).$$

Proposition 3.1. *For every $q \in \mathbb{C}$,*

$$S_{G_q} = D_q \cup P.$$

These are the two irreducible components. For $q \neq -2$, the component D_q is singular and P is smooth.

Proof. For $q \neq -2$, the polynomials A and B_q are coprime. Regarding Δ_q as a quadratic in a , one computes

$$\text{Disc}_a(\Delta_q) = 64(B_q^2 - 3Ab)^3.$$

The right side is not a square in $\mathbb{C}(b, c)$, so Gauss's lemma gives the irreducibility of Δ_q .

Use instead the projective root incidence

$$\overline{X}_q = V(AU^3 + B_q U^2 V + bUV^2 - 2aV^3) \subset \mathbb{A}^3 \times \mathbb{P}^1.$$

The source is its simple-root locus with the deleted infinity section over $c = -1$ removed. The simple infinity root over $c = 0$ is retained: it is the image of $x = 0$, and the restriction there is a triangular isomorphism in (y, z) . The incidence is proper, so the image of its boundary is exactly the union of the projective discriminant and the deleted plane P . This proves the assertion for $q \neq -2$, including the retained infinity-root case. See appendix B for the general construction.

At $q = -2$, one has $B_{-2} = -2(c+1)^2$. Dividing the discriminant by its coefficient content $c+1$ gives the unique primitive nonplane component, while the same lost-root argument gives P . Irreducibility of the primitive

discriminant follows by applying the preceding quadratic argument over $\mathbb{C}(c)$ and then Gauss's lemma. This is also the special case of the all-multiplicity nonproperness theorem in [5].

Finally, for $q \neq -2$ the normalization calculation in proposition 4.1 shows that D_q is singular, while P is a plane. $\square$

4. NORMALIZATION AND THE MARKED PAIR

Assume throughout this section that $q \neq -2$. A repeated root t of the inverse cubic gives

$$\begin{aligned} b &= -3A(c)t^2 - 2B_q(c)t, \\ a &= -A(c)t^3 - \frac{1}{2}B_q(c)t^2. \end{aligned} \tag{4.1}$$

Let

$$\nu_q: \mathbb{A}_{c,t}^2 \longrightarrow D_q$$

be the map (4.1), and put

$$H_q(c, t) = 3c(c+1)t + qc^2 - 4c - 2. \tag{4.2}$$

Proposition 4.1. *The map ν_q is the normalization of D_q . Moreover,*

$$\nu_q^{-1}(\text{Sing } D_q) = L_q := V(H_q).$$

Proof. On the parametrization, t satisfies

$$3At^2 + 2B_q t + b = 0, \tag{4.3}$$

$$B_q t^2 + 2bt - 6a = 0. \tag{4.4}$$

Because $\gcd(A, B_q) = 1$, choose $\rho, \sigma \in \mathbb{C}[c]$ with $\rho A + \sigma B_q = 1$. The sum of $\rho/3$ times (4.3) and σ times (4.4) is a monic quadratic equation for t over $\mathbb{C}[D_q]$. Hence ν_q is finite.

Direct substitution gives

$$B_q^2 - 3Ab = H_q^2, \quad -18Aa - B_q b = 2tH_q^2. \tag{4.5}$$

Thus t is rationally recovered away from $H_q = 0$, so ν_q is birational. Since $\mathbb{A}^2$ is normal, it is the normalization.

The pullbacks of the three partial derivatives of Δ_q are

$$(\Delta_q)_a \circ \nu_q = 8H_q^3, \tag{4.6}$$

$$(\Delta_q)_b \circ \nu_q = -4tH_q^3, \tag{4.7}$$

$$(\Delta_q)_c \circ \nu_q = -4t^2(2cq + 2ct + t - 4)H_q^3. \tag{4.8}$$

The first equality shows that all three vanish simultaneously exactly on $H_q = 0$. $\square$

The plane component P marks a second curve upstairs. Write

$$d = q + 2.$$

On $c = -1$, formulas (4.1) become

$$a = -\frac{d}{2}t^2, \quad b = -2dt. \quad (4.9)$$

Since $d \neq 0$, the inverse image of $D_q \cap P$ is the line

$$M := V(c+1) \subset \mathbb{A}_{c,t}^2,$$

and $M \rightarrow D_q \cap P$ is an isomorphism. Notice also that

$$H_q|_M = q+2 \neq 0,$$

so $L_q \cap M = \emptyset$.

The intrinsic data recovered from S_{G_q} are therefore

$$(\mathbb{A}_{c,t}^2, L_q, M). \quad (4.10)$$

The next lemma is the heart of the stable argument.

Lemma 4.2 (Rigidity of the marked cylinder). *Let $q, q' \neq -2$ and $m \geq 0$. Suppose a polynomial automorphism of $\mathbb{A}^{2+m}$ carries*

$$L_q \times \mathbb{A}^m \text{ to } L_{q'} \times \mathbb{A}^m \quad \text{and} \quad M \times \mathbb{A}^m \text{ to } M \times \mathbb{A}^m.$$

Then $q = q'$.

Proof. Let C, T be the pullbacks of the first two coordinates on the target cylinder. Since the prime principal divisor $M \times \mathbb{A}^m$ is preserved, unique factorization gives

$$C+1 = \lambda(c+1) \quad (4.11)$$

for some $\lambda \in \mathbb{C}^*$. For $q \neq -2$, the polynomial H_q is primitive and linear in t , hence irreducible. Preservation of the other prime divisor similarly gives

$$H_{q'}(C, T) = \kappa H_q(c, t) \quad (4.12)$$

for some $\kappa \in \mathbb{C}^*$.

The right side of (4.12) is independent of the stabilization variables and linear in t . Since C is the nonconstant polynomial in (4.11), it follows that

$$T = \mu(c)t + h(c)$$

for polynomials $\mu, h \in \mathbb{C}[c]$. Comparing the coefficients of t in (4.12) and cancelling $c+1$ gives

$$\lambda(\lambda(c+1)-1)\mu(c) = \kappa c. \quad (4.13)$$

The linear polynomial $\lambda(c+1)-1$ must divide c . Therefore $\lambda = 1$, and (4.13) gives $\mu = \kappa$.

The constant term of (4.12) is now

$$3c(c+1)h(c) + B_{q'}(c) = \kappa B_q(c). \quad (4.14)$$

At $c = 0$, both B_q and $B_{q'}$ equal -2 , so $\kappa = 1$. At $c = -1$, equation (4.14) gives $q' + 2 = q + 2$. $\square$

Theorem 4.3. *If $q, q' \neq -2$, then G_q and $G_{q'}$ are stably polynomially left-right equivalent if and only if $q = q'$.*

Proof. Sufficiency is immediate. For necessity, suppose the stabilizations by $\mathbb{A}^m$ are left-right equivalent. By (3.1), the target automorphism identifies

$$(D_q \cup P) \times \mathbb{A}^m \quad \text{with} \quad (D_{q'} \cup P) \times \mathbb{A}^m.$$

The plane component is smooth and the discriminant component is singular, so the two components cannot be exchanged. We obtain an isomorphism

$$D_q \times \mathbb{A}^m \simeq D_{q'} \times \mathbb{A}^m$$

that preserves the intersection with the plane component.

Normalization commutes with adjoining polynomial variables. The isomorphism therefore lifts uniquely to an automorphism of $\mathbb{A}^2 \times \mathbb{A}^m$. The inverse image of the singular locus is $L_q \times \mathbb{A}^m$, and the inverse image of the plane intersection is $M \times \mathbb{A}^m$. The lifted automorphism preserves this marked pair, so lemma 4.2 gives $q = q'$. $\square$

5. THE EXCEPTIONAL VALUE

At $q = -2$,

$$B_{-2}(c) = -2(c+1)^2$$

and Δ_{-2} acquires the factor $c+1$. Let

$$\Delta_{-2}^{\text{res}} = \frac{\Delta_{-2}}{c+1}, \quad D_{-2} = V(\Delta_{-2}^{\text{res}}).$$

Proposition 5.1. *The map G_{-2} is not stably left-right equivalent to G_q for any $q \neq -2$.*

Proof. On the plane $c = -1$, exact substitution gives

$$\Delta_{-2}^{\text{res}}|_{c=-1} = 4b^3.$$

Thus the reduced intersection $D_{-2} \cap P$ is the line $b = 0$. At the point

$$p = (a, b, c) = (0, 0, -1),$$

all three first partial derivatives of Δ_{-2}^{res} vanish. Hence

$$\text{Sing}(D_{-2}) \cap P \neq \emptyset.$$

For $q \neq -2$, by contrast,

$$\left. \frac{\partial \Delta_q}{\partial a} \right|_{c=-1} = 8(q+2)^3 \neq 0.$$

Consequently D_q is smooth along its entire intersection with P , and

$$\text{Sing}(D_q) \cap P = \emptyset.$$

Whether the singular locus of the unique nonplane component meets the plane component is intrinsic to the reduced nonproperness divisor. The same is true after taking a product with affine space. Equation (3.1) therefore separates G_{-2} from every other G_q stably. $\square$

Corollary 5.2. *For all $q, q' \in \mathbb{C}$,*

$$G_q \sim_{\text{stable}} G_{q'} \iff q = q'.$$

Proof. Combine theorem 4.3 and proposition 5.1. $\square$

6. THE LINE $\alpha = 0$ IS GAUGE

It remains to understand the parameter line not covered by the normalization $\alpha = 1$. When $\alpha = 0$,

$$A(C) = C, \quad B(C) = -2 + \beta C^2.$$

Put

$$s = \frac{\beta}{3}, \quad c(x, y, z) = 2x - 3x^2y - x^3z,$$

and define

$$\Theta_s(x, y, z) = \left(x, y + sc, z - 3s\frac{c}{x} \right). \quad (6.1)$$

This is polynomial because

$$\frac{c}{x} = 2 - 3xy - x^2z.$$

It preserves c , has Jacobian determinant one, and has inverse Θ_{-s} .

Set

$$\ell_s(c) = 3s^2c^3 - 4sc, \quad \delta_s(c) = s^3c^4 - 2s^2c^2, \quad (6.2)$$

and define the triangular target automorphism

$$\Xi_s(a, b, c) = \left(a - \frac{1}{2}\delta_s(c) - \frac{sc}{2}b, b + \ell_s(c), c \right). \quad (6.3)$$

Proposition 6.1. *For every $\beta \in \mathbb{C}$,*

$$G_{0,\beta} = \Xi_{\beta/3} \circ G_{0,0} \circ \Theta_{\beta/3}.$$

In particular, the entire line $\alpha = 0$ is one ordinary left-right orbit.

Proof. Because Θ_s fixes c , it sends the framed root $t = y + 1/x$ to $t + sc$. Substitute this translation in

$$cT^3 - 2T^2$$

and compare the coefficients of T with

$$cT^3 + (-2 + 3sc^2)T^2.$$

The residual linear and constant terms are precisely the target shears ℓ_s and δ_s in (6.2). Substitution in (2.2) gives the displayed identity. All expressions are polynomial, and Θ_s, Ξ_s are triangularly invertible. $\square$

Proposition 6.2. *No member with $\alpha = 0$ is stably left-right equivalent to a member with $\alpha \neq 0$.*

Proof. For $\alpha = 0$, the polynomial $A(C) = C$ has only the marked root. The reduced nonproperness set therefore consists solely of its irreducible primitive discriminant component. For $\alpha \neq 0$, the second root $-1/\alpha$ contributes a plane component in addition to the primitive discriminant component, as in proposition 3.1. Thus the two nonproperness sets have respectively one and two irreducible components. Products with $\mathbb{A}^m$ preserve this component count, and (3.1) makes it a stable left-right invariant. $\square$

Proof of theorem 1.1. The first assertion is proposition 6.1; the second is proposition 6.2. For nonzero α, α' , equality of β/α^2 gives the ordinary equivalence (2.5). Conversely, normalize both maps to $\alpha = 1$ using the same formula and apply corollary 5.2. $\square$

7. EXACT VERIFICATION AND PROOF BOUNDARY

The accompanying verifier uses exact SymPy arithmetic. It checks:

- (1) the discriminant (3.2) and its discriminant as a quadratic in a ;
- (2) the normalization formulas (4.1);
- (3) both rational-recovery identities (4.5);
- (4) all three gradient factorizations (4.6)–(4.8);
- (5) the marked-line formulas (4.9);
- (6) the factorization and singular-incidence calculation at $q = -2$;
- (7) the coefficient identity used in lemma 4.2;
- (8) the identity that gauges away β when $\alpha = 0$; and
- (9) the Jacobian identity $\det D\Theta_s = 1$.

The unchanged recovered verifier was rerun in a fresh environment with SymPy declared explicitly. It printed

ALL q-ORBIT CHECKS PASSED,

and regenerated a JSON audit byte-for-byte identical to the recovered reference output.

The principal verifier is `code/main-modulus/verify_q_full_orbit.py`. The companion source also includes separate exact checks for the reciprocal family, the grading-preserving and based moduli problems, conductor arrangements, weighted-lift critical-root invariants, and boundary-residue constructions. They are indexed by mathematical role in `COMPUTATION.md`; their purpose is to make the displayed invariant formulas reproducible without conflating the narrower framed equivalence problems with the stable left–right theorem in the body.

These computations verify the displayed polynomial identities. They do not replace the geometric arguments that the nonproperness divisor is intrinsic, that normalization lifts an isomorphism, or that the two marked curves are recovered as asserted. Those are conventional proofs in sections 3 and 4. A genuinely independent CAS implementation and expert review of the stable-cylinder argument remain release gates.

8. CONSEQUENCES AND QUESTIONS

Theorem 1.1 gives an affine line of stable left–right orbits at one fixed generic degree, together with one additional degenerate orbit. The modulus is not imposed as part of a framing: it is reconstructed from the intrinsic boundary geometry of the map.

The proof deliberately leaves broader claims outside its scope. In particular, it does not classify all cubic-frame counterexamples, all generic-degree-three Keller maps, or the full stable moduli functor.

Question 8.1. *Which larger finite-dimensional cubic-frame families admit complete stable invariants recovered from their normalized nonproperness divisors?*

Question 8.2. *Can the marked-cylinder rigidity argument be formulated as a general cancellation theorem for pairs $(\mathbb{A}^2, L, M)$?*

Question 8.3. *What scheme or stack represents deformations of the maps in theorem 1.1 modulo stable polynomial left–right equivalence?*

Question 8.4 (Compactification across boundary collisions). *Can the all-multiplicity finite-root classification of theorem B.5 be compactified across strata where deleted roots escape to $c = \infty$? The elementary one-root transition on the resultant-open locus is the contracted gauge chart of proposition B.7. The desired object must also handle simultaneous escapes and the nonunit resultant boundary while retaining the weighted relative-Jacobian data as the number of finite marked planes changes.*

APPENDIX A. BOUNDARY RESIDUES FOR GENERAL SQUAREFREE CUBIC FRAMES

The one-parameter invariant is the first member of a general construction. We record the squarefree statement here; the full standalone sequel source, including expanded calculations and examples, is retained under [supplements/](#).

Definition A.1 (Admissible frame). A pair $A, B \in \mathbb{C}[c]$ is admissible if

$$A(0) = 0, \quad A'(0) = 1, \quad B(0) = -2, \quad B'(0) = -2A''(0).$$

Using

$$c = 2x - 3x^2y - x^3z, \quad t = y + \frac{1}{x}, \quad r = \frac{2}{x},$$

define $G_{A,B} = (a, b, c)$ by

$$b = r - 3A(c)t^2 - 2B(c)t, \quad 2a = A(c)t^3 + B(c)t^2 + tb.$$

Proposition A.2 (Universal cubic frame). *For every admissible pair, $G_{A,B}$ is polynomial and*

$$\det DG_{A,B} = -2.$$

If $A \neq 0$, its generic degree is three.

Proof. The admissibility jets cancel the possible poles at $x = 0$. On $x \neq 0$, the change $(x, y, z) \mapsto (t, r, c)$ has determinant $-2x$, while $(t, r, c) \mapsto (a, b, c)$ has determinant $r/2$. Since $r = 2/x$, the product is -2 , and polynomiality extends it across $x = 0$. Finite simple roots of

$$A(c)T^3 + B(c)T^2 + bT - 2a$$

recover the source, giving generic degree three. $\square$

Assume now that A is squarefree and $\gcd(A, B) = 1$. Put

$$Z_A^\circ = V(A/c) \subset \mathbb{G}_m.$$

Homogenizing the inverse cubic gives

$$A(c)U^3 + B(c)U^2V + bUV^2 - 2aV^3 = 0.$$

At a root s of A , in the local coordinate V/U , the transverse derivative of the infinity section is $B(s)$. The root over $c = 0$ is retained in the affine source; the roots in Z_A° are deleted. Thus the natural decorated infinity datum is

$$(Z_A^\circ, \sigma_B = B|_{Z_A^\circ}).$$

The discriminant component is normalized by

$$\nu_{A,B}: \mathbb{A}_{c,t}^2 \longrightarrow D_{A,B},$$

$$b = -3A(c)t^2 - 2B(c)t, \quad a = -A(c)t^3 - \frac{1}{2}B(c)t^2.$$

The preimage of its singular locus is

$$L_{A,B} = V(3A(c)t + B(c)),$$

and the deleted infinity components lift to the vertical lines

$$M_s = V(c - s), \quad s \in Z_A^\circ.$$

Theorem A.3 (Stable boundary Torelli on the squarefree locus). *Let (A, B) and $(\tilde{A}, \tilde{B})$ be admissible squarefree coprime pairs. The following are equivalent:*

- (i) $G_{A,B}$ and $G_{\tilde{A},\tilde{B}}$ are stably polynomially left-right equivalent;
- (ii) they are ordinarily polynomially left-right equivalent;
- (iii) there is $u \in \mathbb{C}^*$ such that

$$\tilde{A}(uc) = uA(c), \quad \tilde{B}(uc) - B(c) \in cA(c)\mathbb{C}[c];$$

- (iv) multiplication by u identifies the decorated finite schemes

$$(Z_A^\circ, \sigma_B) \quad \text{and} \quad (Z_{\tilde{A}}^\circ, \sigma_{\tilde{B}}).$$

Proof. A stable equivalence preserves the reduced nonproperness divisor. Its unique singular nonplane component is the discriminant component, while its other components are the planes $c = s$, $s \in Z_A^\circ$. Passing to normalizations gives an automorphism of a cylinder over $\mathbb{A}_{c,t}^2$ preserving $L_{A,B}$ and the union of the vertical lines.

Write $Q_A = A/c$, and let C, T be the pullbacks of the normalization coordinates. Unique factorization gives

$$Q_{\tilde{A}}(C) = \lambda Q_A(c).$$

Hence C is algebraic over $\mathbb{C}(c)$. Since $\mathbb{C}(c)$ is algebraically closed in the purely transcendental cylinder function field, $C \in \mathbb{C}(c)$. Polynomiality of the automorphism and its inverse then force

$$C = uc + v.$$

The omitted vertical roots account for every root except the distinguished retained root, so the latter is preserved and $v = 0$. The normalization $A'(0) = \tilde{A}'(0) = 1$ yields

$$\tilde{A}(uc) = uA(c).$$

Preservation of the irreducible singular-preimage curve gives

$$3\tilde{A}(C)T + \tilde{B}(C) = \kappa(3A(c)t + B(c)).$$

The value at $c = 0$ gives $\kappa = 1$, and the admissibility jets imply

$$\tilde{B}(uc) - B(c) \in cA(c)\mathbb{C}[c].$$

This proves (i) $\Rightarrow$ (iii); (iii) and (iv) are equivalent by squarefreeness.

Conversely, after the diagonal change

$$(x, y, z) \mapsto (ux, y/u, z/u^2), \quad (a, b, c) \mapsto (a/u^2, b/u, uc),$$

condition (iii) reduces to

$$\tilde{B} - B = 3A\phi, \quad \phi \in c\mathbb{C}[c].$$

The source root translation

$$(x, y, z) \mapsto \left(x, y + \phi(c), z - 3\frac{\phi(c)}{x} \right)$$

is polynomial, preserves c , and shifts t by $\phi(c)$. Expanding the cubic supplies a triangular target automorphism, proving ordinary left–right equivalence. $\square$

Corollary A.4 (Arbitrarily large fixed-degree stable moduli). *If $\deg A = N + 1$, the generic squarefree orbit space is*

$$\frac{\{(s_i, v_i)_{i=1}^N \in (\mathbb{G}_m \times \mathbb{G}_m)^N : s_i \neq s_j\}}{\mathbb{G}_m \times S_N},$$

where $\mathbb{G}_m$ scales the s_i and fixes the v_i . Its generic dimension is $2N - 1$. Fixing a root configuration with no scaling symmetry leaves an N -dimensional family of pairwise stably inequivalent maps. With the canonical interpolating representative for B , every map in such a family has

$$\begin{aligned} \det DG &= -2, & \mu(G) &= 3, \\ (\deg G_1, \deg G_2, \deg G_3) &= (4N + 7, 4N + 6, 4). \end{aligned}$$

Remark A.5 (The quadratic modulus). For

$$A = c + \alpha c^2, \quad B = -2 - 4\alpha c + \beta c^2,$$

the unique deleted infinity point is $-1/\alpha$, and

$$B(-1/\alpha) = 2 + \frac{\beta}{\alpha^2} = q + 2.$$

Thus q is the normalized transverse derivative at the deleted infinity section. Equivalently, it is the principal part of the rational Tschirnhaus translation needed to identify the cubic frames. At $q = -2$ that derivative vanishes and the deleted section becomes ramified.

Proof boundary. The exact verifier checks all displayed polynomial identities, the frame Jacobian, the marked infinity section, the root-translation gauge, and an explicit three-parameter interpolation family. The four load-bearing geometric steps—identification of the affine open in the finite completion, equality with the nonproperness locus, lifting stable equivalences through normalization, and rigidity of the marked vertical cylinder—still require independent specialist review.

APPENDIX B. ALL MULTIPLICITIES VIA THE RELATIVE JACOBIAN BLOWUP

The squarefree boundary-residue argument extends to arbitrary common roots once the absolute singular locus is replaced by the relative Jacobian over the intrinsic affine c -line. Let (A, B) be an admissible cubic frame, put $Q = A/c$, and write

$$P_{a,b,c}(T) = A(c)T^3 + B(c)T^2 + bT - 2a.$$

B.1. Projective incidence and the primitive discriminant. Consider

$$\overline{X}_{A,B} = V\left(AU^3 + BU^2V + bUV^2 - 2aV^3\right) \subset \mathbb{A}_{a,b,c}^3 \times \mathbb{P}_{[U:V]}^1.$$

The marked-root map extends regularly by

$$(x, y, z) \longmapsto (G_{A,B}(x, y, z), [1 + xy : x]).$$

If $\overline{X}^{\text{simp}}$ is the simple-projective-root locus, then

$$\mathbb{A}^3 \simeq \overline{X}^{\text{simp}} \setminus V(V, Q(c)). \quad (\text{B.1})$$

On the finite-root chart the inverse is the usual reconstruction formula. At $x = 0$, the retained infinity root lies over $c = 0$; writing

$$A = c + a_2c^2 + a_3c^3 + \cdots, \quad B = -2 - 4a_2c + b_2c^2 + \cdots,$$

its restriction is the triangular map

$$b = y + 4a_2, \quad a = z + 4y^2 + 2a_2y - 2b_2 - 8a_3.$$

Thus the retained infinity section belongs to the affine source.

The cubic discriminant is

$$\Delta = B^2b^2 - 4Ab^3 + 8B^3a - 108A^2a^2 - 36ABab.$$

Proposition B.1 (Primitive discriminant). *Its coefficient content in a, b is*

$$\chi(c) = \gcd(A(c), B(c)^2),$$

normalized by $\chi(0) = 1$. The polynomial $\delta = \Delta/\chi$ is irreducible, and

$$S_{G_{A,B}} = V(\delta) \cup \bigcup_{s \in V(Q)_{\text{red}}} V(c - s).$$

Proof. At s , the minimum coefficient valuation of Δ is

$$\min\{\text{ord}_s A, 2 \text{ord}_s B\},$$

which gives the content formula. Over $\mathbb{C}(c)$,

$$\text{Disc}_a(\Delta) = 64(B^2 - 3Ab)^3,$$

which is not a square in $\mathbb{C}(c, b)$; Gauss's lemma gives irreducibility after removing content. Finally, equation (B.1) identifies the boundary of a proper incidence morphism: its two kinds of images are nonsimple roots and the deleted simple infinity roots. $\square$

Proposition B.2 (Intrinsic common planes). *For a deleted root s of A ,*

$$B(s) = 0 \iff V(\delta) \cap V(c - s) \text{ is scheme-theoretically nonreduced.}$$

Proof. Put $u = c - s$, $A = u^r \alpha$, and $B = u^k \beta$, with the displayed units at $u = 0$. If $k = 0$, the restriction is the reduced smooth conic

$$B(s)^2(b^2 + 8B(s)a).$$

If $k > 0$, its primitive restriction is, according as $r < 2k$, $r = 2k$, or $r > 2k$,

$$-4\alpha(0)b^3, \quad b^2(\beta(0)^2 - 4\alpha(0)b), \quad \beta(0)^2b^2.$$

$\square$

B.2. The relative Gauss chart. The pencil of deleted planes recovers c up to affine change. On $D = V(\delta)$, define

$$\mathfrak{J}_{D/\mathbb{A}^1} = \text{Fitt}_1 \Omega_{D/\mathbb{A}^1} = (\delta_a, \delta_b).$$

Let $\mathfrak{B}_D = \text{Bl}_{\mathfrak{J}} D$. Write

$$g = \gcd(A, B), \quad A = gA_0, \quad B = gB_0,$$

and put

$$H_0 = 3A_0t + B_0, \quad \rho = g^3/\chi.$$

Proposition B.3 (Finite-root chart and weighted divisor). *The δ_a -chart of $\mathfrak{B}_D$ is canonically $\mathbb{A}_{c,t}^2$. On it,*

$$\mathfrak{J}\mathcal{O}_{\mathbb{A}^2} = (\rho(c)H_0(c, t)^3).$$

Proof. Under the repeated-root parametrization

$$b = -3At^2 - 2Bt, \quad a = -At^3 - \frac{1}{2}Bt^2,$$

one obtains

$$\delta_a = 8\rho H_0^3, \quad \delta_b = -4t\rho H_0^3.$$

Hence $t = -2\delta_b/\delta_a$, so adjoining the blowup ratio adjoins exactly t ; conversely a, b are polynomial in c, t . $\square$

This chart is intrinsic. Indeed, on the other chart put $q = \delta_a/\delta_b = -2/t$. Then

$$B(c) = qb + \frac{3}{2}q^2a, \quad A(c) = \frac{1}{4}q^2b + \frac{1}{2}q^3a.$$

The complement $q = 0$ therefore maps precisely into the curve intersections with the common deleted planes characterized by proposition B.2; removing the prime divisors above those curves leaves the chart of proposition B.3.

B.3. Multiplicity recovery and stable Torelli. At a deleted root s , let

$$r = \text{ord}_s A, \quad k = \text{ord}_s B, \quad m = \min(r, k).$$

The intrinsic chart records

$$p = \text{ord}_s A_0 = r - m, \quad d = \text{ord}_s \rho = 3m - \min(r, 2k).$$

Lemma B.4 (Multiplicity recovery). *The common-factor multiplicity is*

$$m = \begin{cases} d, & p \geq d, \\ (p + d)/2, & p < d. \end{cases}$$

Theorem B.5 (All-multiplicity fixed-frame Torelli). *Let (A, B) and $(\tilde{A}, \tilde{B})$ be admissible cubic frames, with A/c and $\tilde{A}/c$ nonconstant. The following are equivalent:*

- (i) $G_{A,B}$ and $G_{\tilde{A},\tilde{B}}$ are stably left-right equivalent;
- (ii) they are ordinarily polynomially left-right equivalent;
- (iii) there is $u \in \mathbb{C}^\times$ such that

$$\tilde{A}(uc) = uA(c), \quad \tilde{B}(uc) - B(c) \in cA(c)\mathbb{C}[c].$$

No squarefreeness or coprimality hypothesis is required.

Proof. The intrinsic finite-root chart, its marked vertical lines, and the divisor $\text{div}(\rho H_0^3)$ all survive a product with affine space. A stable equivalence preserves this data. A marked vertical line forces the pulled-back base coordinate to have the form $C = uc + v$. The unique nonvertical divisor forces

$$3\tilde{A}_0(C)T + \tilde{B}_0(C) = \kappa(3A_0(c)t + B_0(c)).$$

Thus $T = \mu(c)t + h(c)$. The cylinder Jacobian makes μ a nonzero constant and excludes dependence on stabilization variables.

The horizontal divisor gives every p_s , while the vertical multiplicity gives every d_s . By lemma B.4, all multiplicities of A are therefore preserved. The deleted planes account for every root except the retained root zero, so $v = 0$. Comparing the t -coefficient and using the admissible first jets gives

$$\tilde{A}(uc) = uA(c).$$

The constant coefficient then gives

$$\tilde{B}(uc) - B(c) = -3uA(c)h(c).$$

Differentiating at zero shows $h(0) = 0$, which is condition (iii). Conversely, the explicit root translation used in the ordinary classification constructs a polynomial left–right equivalence and does not use squarefreeness or coprimality. $\square$

Equivalently, scaling identifies the decorated Artin schemes

$$(\operatorname{Spec} \mathbb{C}[c]/(A/c), B \bmod (A/c)).$$

Proposition B.6 (The exceptional weighted stratum). *For the exceptional quadratic member $A = c(c+1)$, $B = -2(c+1)^2$, one has*

$$\rho = (c+1)^2, \quad H_0 = 3ct - 2(c+1).$$

Thus its special behavior is the intrinsic vertical multiplicity two, not an ad hoc extra orbit invariant.

Proof. Here $g = c+1$, $A_0 = c$, $B_0 = -2(c+1)$, and $\chi = c+1$. Substitution in proposition B.3 gives

$$\mathfrak{J}\mathcal{O}_{\mathbb{A}^2} = ((c+1)^2(3ct - 2(c+1))^3),$$

so the vertical multiplicity is exactly two. $\square$

B.4. The elementary cross-length boundary chart. The preceding theorem works at fixed degree. There is nevertheless a simple codimension-one transition between successive fixed-degree normal forms. Write the length- N coefficient chart as

$$Q = 1 + q_1c + \cdots + q_Nc^N, \quad R = r_0 + r_1c + \cdots + r_{N-1}c^{N-1},$$

with

$$A = cQ, \quad B = -2 - 4q_1c + c^2R.$$

The divisor $q_N = 0$ in its coefficient closure is the locus where one root of Q has escaped to infinity.

Proposition B.7 (One-root transition). *On the open part*

$$q_N = 0, \quad q_{N-1} \operatorname{Res}(Q, B) \neq 0,$$

put

$$\kappa = \frac{r_{N-1}}{q_{N-1}}, \quad r'_j = r_j - \kappa q_j \quad (0 \leq j \leq N-2), \quad q_0 = 1.$$

Then the coefficient change is invertible and equivariant, with κ of weight two. The boundary chart is the length- $(N-1)$ chart times $\mathbb{A}_\kappa^1$, and the map to polynomial left–right orbit data contracts the second factor.

Proof. Set $R' = R - \kappa Q$. Its top coefficient vanishes, so $\deg R' \leq N-2$, while

$$B - B' = \kappa c^2 Q = \kappa c A = 3A \left(\frac{\kappa c}{3} \right).$$

The last expression is exactly the allowed polynomial root-translation gauge. Conversely,

$$r_j = r'_j + \kappa q_j \quad (0 \leq j \leq N-2), \quad r_{N-1} = \kappa q_{N-1},$$

so this is an invertible coordinate change on $q_{N-1} \neq 0$. Under the weighted scaling, q_j has weight j , r_j has weight $j+2$, and hence κ has weight two and r'_j has weight $j+2$. Finally, with the displayed degrees in the Sylvester determinant,

$$\text{Res}(Q, B) = (-1)^{N-1} q_{N-1} \text{Res}(Q, B').$$

The multiplier is a unit on this chart, so the resultant-open condition is unchanged. $\square$

The first two instances are

transition	κ	R'
$2 \longrightarrow 1$	r_1/q_1	$r_0 - r_1/q_1$
$3 \longrightarrow 2$	r_2/q_2	$(r_0 - r_2/q_2) + (r_1 - q_1 r_2/q_2)c$

Thus the first boundary phenomenon is not an additional modulus: it is a gauge line over the lower-length chart.

Question B.8 (Gluing at infinity). *Can proposition B.7 be extended across simultaneous root escapes, the nonunit resultant boundary, and the weighted relative-Jacobian marking to give a full compactification? This also includes comparison with the $\alpha = 0$ stratum and remains separate from the finite-root theorem above.*

APPENDIX C. EXACT CONDUCTORS AND GROWING STABLE MODULI

We return to the cubic frame with

$$h(T, C) = A(C)T^3 + B(C)T^2.$$

The target fiber cubic is

$$P(T) = A(c)T^3 + B(c)T^2 + vT - 2u,$$

with discriminant

$$\Delta_{A,B} = B^2 v^2 - 4A v^3 + 8B^3 u - 108A^2 u^2 - 36ABuv.$$

Assume first that $\gcd(A, B) = 1$.

C.1. The conductor.

Theorem C.1 (Exact discriminant conductor). *The map*

$$\nu : \mathbb{A}_{c,t}^2 \longrightarrow V(\Delta_{A,B}),$$

$$v = -3A(c)t^2 - 2B(c)t, \quad u = -A(c)t^3 - \frac{1}{2}B(c)t^2$$

is the normalization. If

$$H(c, t) = 3A(c)t + B(c),$$

then the conductor in $\mathbb{C}[c, t]$ is exactly

$$H^2\mathbb{C}[c, t].$$

Its contraction to the discriminant coordinate ring is generated by

$$B^2 - 3Av, \quad -18Au - Bv.$$

Proof. Irreducibility follows from

$$\text{Disc}_u(\Delta_{A,B}) = 64(B^2 - 3Av)^3,$$

since $B^2 - 3Av$ is not a square in $\mathbb{C}(c, v)$. Bézout coefficients for A, B combine the equations

$$3At^2 + 2Bt + v = 0, \quad Bt^2 + 2vt - 6u = 0$$

into a monic quadratic equation for t , proving finiteness. The identities

$$B^2 - 3Av = H^2, \quad -18Au - Bv = 2tH^2$$

give birationality and show that $H^2\mathbb{C}[c, t]$ lies in the conductor.

At the generic point of $H = 0$, use H as transverse coordinate. Over $K = \mathbb{C}(c)$, the inclusion is

$$K[H^2, H^3] \subset K[H],$$

whose conductor is $H^2K[H]$. Every global conductor element therefore has H -valuation at least two. This proves equality and the stated contraction. $\square$

Thus the reduced conductor support

$$L_{A,B} = V(3A(c)t + B(c))$$

is intrinsic to the discriminant surface, not merely detected by a chosen gradient calculation.

C.2. Marked arrangements. Assume A is squarefree and a nonempty subset of its roots marks planes on which an infinity sheet is missing. In the normalization these planes pull back to vertical lines $M_\rho = (c = \rho)$.

Theorem C.2 (Stable conductor-arrangement classification). *Two marked arrangements*

$$\left(\mathbb{A}^2, L_{A,B}, \{M_\rho\}_{\rho \in S}\right) \quad \text{and} \quad \left(\mathbb{A}^2, L_{A',B'}, \{M_{\rho'}\}_{\rho' \in S'}\right)$$

become isomorphic after multiplication by an affine space if and only if there exist

$$\gamma(c) = \lambda c + \mu, \quad \lambda \neq 0, \quad \nu, \kappa \in \mathbb{C}^*, \quad h(c) \in \mathbb{C}[c],$$

such that γ carries the marked roots to the marked roots and

$$A'(\gamma(c))\nu = \kappa A(c),$$

$$B'(\gamma(c)) + 3A'(\gamma(c))h(c) = \kappa B(c).$$

Stabilization introduces no additional transformations.

Proof. If a cylinder automorphism has first coordinate C , mapping one marked line to another gives equality of principal prime ideals and hence

$$C - \rho' = \lambda(c - \rho).$$

Thus $C = \lambda c + \mu$, independent of t and of all stabilization variables. The conductor curves are irreducible principal divisors, so

$$3A'(C)T + B'(C) = \kappa(3A(c)t + B(c)).$$

It follows that $T = \nu(c)t + h(c)$. Factoring the Jacobian determinant of the cylinder automorphism shows that $\nu(c)$ is a unit, hence constant. Coefficient comparison gives the two equations. Conversely, they define the required triangular automorphism without using any stabilization variable. $\square$

For squarefree A , the complete data are therefore the affine configuration of its roots, with represented/missing markings, and the projective vector

$$[B(\rho)]_{\rho \in Z(A)}$$

modulo the finite permutations induced by affine symmetries. Equivalently, after compactifying the t -direction, the boundary tangent at a root ρ is

$$\tau_\rho = -\frac{3A'(\rho)}{B(\rho)},$$

and the projective vector of these slopes is intrinsic.

C.3. Consequences for the q -family. For

$$A(c) = c(c+1), \quad B_q(c) = qc^2 - 4c - 2,$$

the roots 0 and -1 are respectively represented and missing, and

$$B_q(0) = -2, \quad B_q(-1) = q + 2.$$

Thus, for $q \neq -2$,

$$q = 2 \frac{\tau_0}{\tau_{-1}} - 2.$$

At $q = -2$, the discriminant gains a plane factor and the residual component is singular at $(u, v, c) = (0, 0, -1)$; this separates the exceptional member intrinsically. These calculations give another proof of the complete stable q -classification in the body.

C.4. A three-dimensional degree-fifteen family. Let

$$s \in \mathbb{C} \setminus \{0, 1\}, \quad d_1, d_2 \in \mathbb{C}^*,$$

and set

$$A_s(c) = \frac{c(c+1)(c+s)}{s}.$$

There is a unique cubic B_{s,d_1,d_2} satisfying

$$B(0) = -2, \quad B'(0) = -2A_s''(0), \quad B(-1) = d_1, \quad B(-s) = d_2.$$

Explicitly,

$$B = -2 - \frac{4(s+1)}{s}c + b_2c^2 + b_3c^3,$$

where

$$b_2 = \frac{d_1s^3 - d_2 - 2s^3 - 4s^2 + 4s + 2}{s^2(s-1)}, \quad b_3 = \frac{d_1s^2 - d_2 - 2s^2 + 2}{s^2(s-1)}.$$

Theorem C.3 (Degree-fifteen stable moduli). *The associated maps are polynomial Keller maps of component degrees $(15, 14, 4)$, generic degree three, and generic monodromy S_3 . Their stable equivalence relation is exactly*

$$(s, d_1, d_2) \sim (1/s, d_2, d_1).$$

Consequently degree fifteen contains a three-dimensional stable-moduli family, modulo this finite involution.

Proof. The jet conditions at $c = 0$ are precisely the pole-cancellation conditions in the cubic frame. The represented root is 0, while the two missing roots are $-1, -s$. By theorem C.2, an equivalence must fix the represented root and either fix or interchange the missing roots. This gives the identity or the displayed involution. The involution is realized by

$$(x, y, z) \mapsto (x/s, sy, s^2z)$$

and

$$(a, b, c) \mapsto (s^2a, sb, c/s).$$

□

Theorem C.4 (Growing stable moduli). *For every $n \geq 2$, the cubic frame contains a $(2n-3)$ -dimensional family of pairwise stably inequivalent generic-degree-three Keller maps of ordinary degree $4n + 3$.*

Proof. Take squarefree A of degree n , with one represented root and $n - 1$ missing roots. Normalize the represented root to 0, impose

$$A'(0) = 1, \quad B(0) = -2, \quad B'(0) = -2A''(0),$$

and prescribe arbitrary nonzero values of B at the other roots. Hermite interpolation produces B of degree at most n . The affine root configuration contributes $n - 2$ parameters, and the projective conductor-value vector contributes $n - 1$. The classification theorem identifies only a finite permutation quotient, giving $(n - 2) + (n - 1) = 2n - 3$ moduli. The leading terms have ordinary degree $4n + 3$. □

Theorem C.5 (Degree threshold inside the cubic frame). *Every polynomial member of ordinary degree at most ten in the full cubic frame is polynomially source-target equivalent to the degree-seven base map. Degree eleven is the first degree at which genuine moduli occur in this frame.*

Proof. If $m = \deg A$ and $n = \deg B$, their unique leading contributions have degrees

$$(4m + 3, 4m + 2) \quad \text{and} \quad (4n + 2, 4n + 1)$$

in the first two coordinates. Degree at most ten forces

$$\deg A = 1, \quad \deg B \leq 2.$$

After centering and scaling,

$$A(C) = C, \quad B(C) = -2 + b_1 C + b_2 C^2.$$

The coefficient b_1 is removed by the source translation responsible for the w_{20} -gauge. The b_2 -family is removed by the explicit locally nilpotent flow already described in the classification of the $\alpha = 0$ line. The q -family supplies inequivalent degree-eleven members. $\square$

C.5. Formal arcs and global separation. For fixed q , take

$$A_s(c) = c + sc^2, \quad B_{s,q}(c) = -2 - 4sc + qs^2c^2.$$

The special fiber $s = 0$ is the base map, and every $s \neq 0$ fiber lies in the stable orbit indexed by q . The coefficientwise formal source trivialization has, through order three, initial degrees

$$(19, 17, 19), \quad (37, 35, 37), \quad (55, 53, 55).$$

The first coefficient is independent of q , and q first occurs in the second.

Proposition C.6 (Formal-local versus stable separation). *All q -arcs have the same first-order formal tangent and are source-trivial through the three computed formal orders, while their punctured fibers remain pairwise stably inequivalent. Thus the stable conductor modulus is invisible to this local formal data.*

Proof. The formal inverse recursion gives the displayed coefficients directly. Stable inequivalence of the punctured fibers follows from the intrinsic ratio of conductor values. No assertion is made here about a minimal degree formula for all higher formal orders. $\square$

APPENDIX D. THE BOUNDED COEFFICIENT GROUPOID AND ITS QUOTIENTS

This appendix separates three objects that coincide on the generic finite-root locus but behave differently on the boundary: the affine categorical quotient, the fppf orbit sheaf, and a separated space retaining degeneration directions.

D.1. A single bounded orbit map. Fix an ambient degree bound N , and write

$$\widehat{Q}(z) = z^N + q_1 z^{N-1} + \cdots + q_N, \quad P \in \mathbb{C}[z]/(\widehat{Q}).$$

Let E_N be the tautological rank- N bundle on $\text{Hilb}^N(\mathbb{A}^1)$, and let Z denote multiplication by z .

Proposition D.1 (Bounded orbit map). *The degree-preserving root translations form the kernel pair of the map*

$$\Theta_N: \text{Tot}(E_N) \longrightarrow \text{Tot}(E_N), \quad (\widehat{Q}, P) \longmapsto (\widehat{Q}, Z^N P).$$

If $\widehat{Q} = z^m Q_d$, with $Q_d(0) \neq 0$, then uniquely

$$P = z^m P_d + Q_d S, \quad \deg P_d < d, \quad \deg S < m.$$

The finite-root decoration is P_d ; root translation removes precisely the principal part S/z^m supported at infinity.

This formulation uses polynomial remainder by the monic polynomial $\widehat{Q}$, and therefore remains defined on the nonunit-resultant and nonreduced boundary.

D.2. Affine categorical quotient. Let R be a $\mathbb{Q}$ -algebra, E a finite projective R -module, $M: E \rightarrow E$, and

$$K = \ker(\mathbf{V}(M)) \curvearrowright X = \mathbf{V}(E)$$

the translation action. Put $A = \text{Sym}_R(E^\vee)$.

Theorem D.2 (Differential description). *The invariant algebra is*

$$A^K = \ker \left(A \xrightarrow{d_{X/R}} A \otimes_R E^\vee \longrightarrow A \otimes_R \text{coker}(M^\vee) \right).$$

Equivalently, in homogeneous degree r , it is the kernel of one finite syzygy map

$$\text{Sym}^r(E^\vee) \longrightarrow \text{Sym}^{r-1}(E^\vee) \otimes_R \text{coker}(M^\vee).$$

Proof. The linear term of $f(p+s) - f(p)$ gives necessity. Conversely, if $df = M^\vee h$, then for $F(t) = f(p+ts)$ one has

$$F'(t) = \langle h(p+ts), Ms \rangle.$$

Coefficientwise integration from zero to one, valid over a $\mathbb{Q}$ -algebra, shows $f(p+s) - f(p)$ lies in the ideal defining K . $\square$

For the cubic-frame problem $M = Z^N$, and the only height-one support of $\text{coker}(M^\vee)$ is the one-root wall $q_N = 0$.

Theorem D.3 (One-wall control). *Inside the generic polynomial algebra,*

$$A_N^{K_N} = A_N \cap (A_N^{K_N})_{(q_N)}.$$

Thus simultaneous escapes impose no additional divisorial conditions on global invariant functions.

Proof idea. The differential sequence has free middle terms, and its cokernel is supported on $q_N = 0$. The resulting homogeneous invariant module is the intersection of the free module with its localization at that one height-one prime. This Cohen–Macaulay support argument replaces an earlier claim that all homogeneous invariant modules were reflexive; that stronger claim was unnecessary and is not used here. $\square$

D.3. Why the fppf quotient is not algebraic.

Theorem D.4 (Rank-one wall obstruction). *Let R be a DVR with uniformizer π , and let*

$$K_e = \operatorname{Spec} R[s]/(\pi^e s)$$

act on $\mathbb{A}_R^1$ by translation. Its fppf orbit sheaf is not an algebraic space, and the quotient stack is not an algebraic stack. Consequently the bounded cubic-frame fppf quotient has the same failure along its generic one-root wall.

Proof idea. The orbit sheaf on an R -algebra B is

$$B/\operatorname{Ann}_B(\pi^e) \simeq \pi^e B.$$

For the standard pushout

$$B = R/(\pi^{e+1}) \times_{R/(\pi^e)} R/(\pi^{e+1}),$$

its value maps diagonally into $k \oplus k$, rather than surjectively. This violates the strong Rim–Schlessinger condition for algebraic stacks. The action is free, so an algebraic quotient stack would be an algebraic space. At the generic cubic-frame wall, Smith normal form gives this rank-one action times an unaffected affine factor. $\square$

D.4. Separated degeneration spaces. For a simple wall, the graph closure of the generic quotient coordinate is the blowup

$$\operatorname{Bl}_{(\epsilon^{N+2}, y)} \operatorname{Spec} B[\epsilon, y].$$

For a simultaneous escape of length m , the unique decomposition in proposition D.1 supplies weighted principal part coordinates. Their projectivization has weights

$$(1, 2, \dots, m, m+1, m, \dots, 2)$$

and gives a finite-type projective separated model of degeneration directions.

These graph closures do not represent the orbit sheaf and do not by themselves prove that stable left–right equivalence is intrinsically reconstructed at infinity. A global complete-weighted-collineation space is a concrete candidate for that last compactification, not a theorem of this appendix.

COMPANION RESULTS AND RESEARCH REGISTER

This reader edition is accompanied by a separate *Results and Research Register*. The register preserves secondary proved results, conditional developments, open questions, corrections, and computational leads without making them part of this paper’s theorem spine. Proof-bearing technical developments omitted from this reader PDF remain in the versioned source tree and in the complete version-8 archival edition. The v9 primary-home map distinguishes the paper responsible for a result from other papers that use or motivate it.

AI AND COMPUTATIONAL DISCLOSURE

AI systems were used extensively in exploration, proof drafting, symbolic program development, source organization, and manuscript editing. They are not authors. The mathematical evidence for the computer-checked identities is the exact verifier and reproducible output described in section 7, not the output of a language model. Human specialist review has not yet been completed.

REFERENCES

1. Levent Alpöge, *Post announcing an explicit counterexample to the jacobian conjecture*, X post.
2. Adrien Dubouloz and Karol Palka, *The jacobian conjecture fails for pseudo-planes*, *Advances in Mathematics* **339** (2018), 248–284.
3. Alexis Gallagher, *The jacobian counterexample, explained*, The page credits substantial mathematical, expository, and computational assistance to GPT-5.6-sol.
4. Zbigniew Jelonek, *The set of points at which a polynomial map is not proper*, *Annales Polonici Mathematici* **58** (1993), no. 3, 259–266.
5. Nathaniel Monson, *Cubic marked-root covers: Normal forms, equivalence, and stable uniqueness*, Companion manuscript.
6. Terence Tao, *A digestion of the jacobian conjecture counterexample*.
7. Ulam AI, *A counterexample to the jacobian conjecture*.