

Top level comment: Try to remove all references to things being either “interesting” or “surprising” unless it is in the discussion and you can point to a specific reference or paper that predicts something different!

SUBMITTED TO THE ASTROPHYSICAL JOURNAL

Preprint typeset using L<sup>A</sup>T<sub>E</sub>X style emulateapj v. 12/16/11

TOP LEVEL comment: I don’t think we’re ever talking about AGN, exactly. At some point we should clarify the difference between our black-hole feedback and AGN feedback and be super explicit and careful not to use the two terms interchangeably...

## Not So Heavy Metals: Black Hole Feedback Enriches The Circumgalactic Medium

### NOT SO HEAVY METALS: BLACK HOLE TRANSPORTED OXYGEN TO THE CIRCUMGALACTIC MEDIUM

N. NICOLE SANCHEZ<sup>1</sup>, JESS WERK<sup>1</sup>, MICHAEL TREMMEL<sup>2</sup>, ANDREW PONTZEN<sup>3</sup>, CHARLOTTE CHRISTENSEN<sup>4</sup>, TOM QUINN<sup>1</sup>, AKAXIA CRUZ<sup>1</sup>, AND [Order subject to change]

<sup>1</sup>Astronomy Department, University of Washington, Seattle, WA 98195, US, sanchenn@uw.edu

<sup>2</sup>Yale Center for Astronomy & Astrophysics, Physics Department, P.O. Box 208120, New Haven, CT 06520, USA

<sup>3</sup>Department of Physics & Astronomy, University College London, 132 Hampstead Road, London, NW1 2PS, United Kingdom and

<sup>4</sup>Physics Department, Grinnell College, 1116 Eighth Ave., Grinnell, IA 50112, United States

Submitted to *The Astrophysical Journal*

#### ABSTRACT

[UNDER CONSTRUCTION] We examine the effects of supermassive black hole (SMBH) feedback on the circumgalactic medium (CGM) using a cosmological hydrodynamic simulation (ROMULUS 25 Tremmel et al. 2017) and a set of zoom-in “genetically modified” Milky Way-mass galaxies sampling different evolutionary paths. We compare the column densities of OVI in Milky Way-mass galaxies and compare them with observations from the COS-Halos Survey; contrary to previous simulations which underpredicted the CGM column densities of OVI, these simulations are consistent with COS-Halos observations. We determine that a galaxy’s star formation history and accretion rate have little effect on the appearance of OVI in its CGM while column densities of OVI are more closely tied to galaxy halo mass. The set of zoom-in, genetically modified Milky Way-mass galaxies further confirm this result and demonstrate the effect of AGN feedback on the CGM’s OVI. We find that the SMBH acts as the physical mechanism for transporting metals out into its host halo thereby significantly impacting the appearance of OVI found in the CGM.

AGN?? SMBH.

*Subject headings:* Gas physics – Galaxies: circumgalactic medium – Galaxies: spiral – Galaxies: kinematics and dynamics – Methods: Numerical

#### 1. INTRODUCTION

maybe not  
“unexplored...” is  
richly structured  
with the by-  
products of galaxy  
evolution. The circumgalactic medium (CGM), the extended re-  
gion of gas surrounding galaxies out to their virial radii,  
is a rich and vast yet mostly unexplored area of astron-  
omy. Due to its diffuse nature, the CGM has remained  
one of the most difficult regions to observe. However, due  
to technological advances like the Cosmic Origins Spec-  
trograph (COS) on the HST, researchers have finally be-  
gun observing this mysterious component of all galaxies.  
Observers have found it to be a structurally complex,  
multiphase medium ((Tumlinson et al. 2011; Werk et al.  
2012, 2013, 2016; Tumlinson et al. 2017). Examinations  
like Werk et al. (2014) show that most of the “missing  
baryons” of galaxies likely reside in this diffuse region,  
implying that the CGM may play a key role in the growth  
of galaxies and the build up of their disks. Therefore, it  
is clear that understanding the CGM is crucial for un-  
derstanding the complex nature of galaxy evolution and  
growth.

In particular, COS-Halos finds a

Several studies have been done to examine the effect of  
galaxy evolution on the CGM. The COS-Halo observa-  
tions show a correlation between the column densities of  
OVI out to 150 kpc and the specific SFR of their observed  
galaxies. Higher abundances of OVI are found around SF  
galaxies compared to their passive counter parts. Oppenheimer 2016 argue that this bimodality arises due to  
the OVI acting as a proxy for the virial temperature of  
gas in these galaxy halos. Therefore, the more massive  
galaxies in the COS-Halo sample show less OVI in their  
CGM due to the intrinsically higher virial temperature  
of these massive red ellipticals. In contrast, Suresh et al.  
(2017) argue that the OVI is built up by AGN feedback,  
which can physically modify the CGM via outflows or  
heat it to the appropriate temperature for ionizing OVI.

In both cases, each argument implies an intrinsic link  
between the CGM and its host galaxy’s evolution.

Additionally, Galaxy evolution has been shown to be strongly tied  
to the evolution of its central supermassive black hole  
(SMBH). Through relations like the M- $\sigma$  and the bulge  
mass-BH mass correlation (Ferrarese & Merritt 2000;  
McConnell & Ma 2013), recent studies indicate that the  
SMBH and its host galaxy halo co-evolve (Gebhardt et  
al. 2000a; Ferrarese & Merritt 2000; McConnell & Ma  
2013; Kormendy & Ho 2013, Reines 2015 and references  
therein). It is therefore unsurprising that the AGN is  
thought to leave its marks on the CGM. However, the  
direct mechanisms by which the AGN readily impacts  
the CGM are still hotly debated.

AGN may effect the CGM in a variety of ways. First,  
feedback from the active SMBH may inject energy into  
the surrounding material, raising temperatures, and ionizing  
metals in the gas [CITE]. Additionally, massive  
outflows of gas from the AGN may physically push gas  
out of the galaxy (some of which may end up falling  
back into the galaxy as part of the “recycling” of the  
CGM (Tumlinson et al. 2017, MORE), enriching CGM  
gas with metals from the center of the galaxy, and fur-  
ther enriching the IGM as gas is expelled from the galaxy  
halo. Cosmological hydrodynamic simulations have be-  
come a powerful tool for examining the physics driving  
the multiphase nature of the CGM [CITE].

Simulators have long been examining the underlying  
physics of AGN activity in galaxies; however, examining  
these effects in the diffuse region of the CGM is still  
a fairly new field ripe for discovery. Toward this  
end, we utilize a cutting edge set of simulations: the  
cosmological volume, ROMULUS25 (Tremmel et al.  
2017) and three “genetically modified” variations of

In a paper best to keep from such judgments about your own simulation :

search and replace COS-Halo with COS-Halos! :)

If you’re going to introduce Oppenheimer here, you need to be explicit...i.e. more massive halos have  
more of their oxygen in a phase traced by OVII and OVIII...

an isolated, zoom-in Milky-Way (MW) mass galaxy (Roth et al. 2016; Pontzen et al. 2017) simulated with and without the implementation of advanced BH physics (Tremmel et al. 2015). With these simulations, we plan to examine two specific questions with our study.

- How do the star formation and accretion history of the galaxy co-evolve with the CGM?
- How does AGN activity imprint itself on the CGM?

**not really AGN?**

**Second sentence:** we want to add a sentence saying we examine several high-resolution cosmological zooms and choose a MW-analog based on its halo mass

Both the ROMULUS25 simulation and our set of high resolution, zoom in simulations with genetic modifications will allow us to investigate the first question in detail. First, we examine the CGM in a range of MW-mass galaxies with varied morphologies from ROMULUS25. Additionally, from our isolated MW-mass galaxy and its subsequent “genetic modifications”, we quantify the effect of AGN feedback on the CGM by examining a set of galaxies run both with and without BH physics. Using these isolated, zoom-in simulations in tandem with the Romulus25 simulation, we illuminate the roles that stellar evolution and AGN feedback play in setting the properties of the CGM of MW-mass galaxies.

## 2. SIMULATION PARAMETERS

### 2.1. ChaNGa Physics

Both ROMULUS25 (hereafter R25) and our set of zoom-in galaxies were run using the smoothed particle hydrodynamics (SPH) N-body tree code, Charm N-body GrAvity solver [ChaNGa] (Menon et al. 2015). ChaNGa includes the same models for a cosmic UV background, star formation, ‘blastwave’ SN feedback, and low temperature metal line cooling as previously used in GASOLINE (Wadsley et al. 2004, 2008; Stinson et al. 2006; Shen et al. 2010). ChaNGa includes an improved SPH formalism which includes a geometric density approach in the force expression:  $(P_i + P_j)/(\rho_i \rho_j)$  instead of  $P_i/\rho_i^2 + P_j/\rho_j^2$  where  $P_i$  and  $\rho_i$  are the particle’s pressure and density, respectively. This update to the hydrodynamic treatment includes thermal diffusion (Shen et al. 2010) and reduces artificial surface tension allowing for better resolution of fluid instabilities (Ritchie & Thomas 2001; Menon et al. 2015; Governato et al. 2015).

Additional improvements have been made to the BH formation, accretion, and feedback models as well an improved prescription for dynamical friction (Tremmel et al. 2015, 2017).

Our simulations were run with a  $\Lambda$ CDM cosmology from the most recent Planck collaboration utilizing  $\sigma_0 = 0.3086$ ,  $\alpha = 0.6914$ ,  $h = 0.67$ ,  $\sigma_8 = 0.77$  and have Plummer equivalent force softening lengths of 250 pc. For simulating the cosmic reionization energy, both simulations enact a UV background at  $z \sim 9$  (Haardt & Madau 2012).

### 2.2. Romulus 25 Cosmological Volume

The ROMULUS25 (Tremmel et al. 2017) simulation is a cosmological volume which includes galaxy halos within the mass range  $10^8 - 10^{13} M_\odot$ . R25 has a mass resolution of  $3.4 \times 10^5 M_\odot$  and  $2.1 \times 10^5 M_\odot$  for DM and gas particles, respectively. For our study, we focus on galaxies within the MW-mass halo regime,  $5 \times 10^{11} M_\odot - 2 \times$

we examine the effects of both environmental and internal galaxy processes on the physical state and content of the CGM. Specifically we address:




FIG. 1.— R25 is an SPH simulated cosmological volume with galaxy halos spanning the orders of  $10^8 - 10^{13} M_\odot$ . For clarity, we’ve plotted a histogram of only the galaxies in the volume with masses between  $2 \times 10^{11} M_\odot - 2 \times 10^{13} M_\odot$  (The latter is the upper limit of galaxy masses in the volume.) The shaded region indicates the “MW” mass range,  $5 \times 10^{11} M_\odot - 2 \times 10^{12} M_\odot$ , from which we select our 25 galaxies.

$10^{12} M_\odot$  (Figure 1). We examine all the galaxies within the specified mass range and exclude any galaxies within twice the virial radius of another galaxy, thereby removing galaxies that might be satellites of a larger galaxy. With these selection criteria in place, our sample includes 25 galaxies.

Perhaps say a bit more about Romulus? Highlight results? Why is it valid?

### 2.3. Zoom-In Galaxies: Patient 0 and its Genetic Modifications

more specific? MW analog?

To select our galaxy, we ran initial uniform-volume,  $50 h^{-1}$  Mpc on a side, dark matter-only cosmological volume. From this volume a MW-mass halo at  $z=0$  was selected as our “Patient 0” and then re-simulated at a higher resolution. For the subsequent, “genetically modified” (GM) zoom-in runs, we utilize the same method of modification as Pontzen et al. (2017). These GM simulations keep the large scale structure and cosmological conditions  $\Lambda$ CDM consistent (as in Patient 0), while allowing for modifications of their accretion histories (Roth et al., 2015). Patient 0 (and its 3 GM simulations) have a mass resolutions of  $1.4 \times 10^5 M_\odot$  and  $2.1 \times 10^5 M_\odot$  for DM and gas particles, respectively. The DM field in these galaxies is simulated at twice the gas linear resolution to reduce noise in the potential near the galactic center (Pontzen et al. 2017) and more accurately trace black hole dynamics (Tremmel et al. 2015).

Give MW halo mass. Give analog halo mass. Were there other candidates we ruled out? how/when are baryons added? More details need to be included summarizing what GM is.

3.1. Galaxies with BH Physics should we show an image of P0?

At  $z = 0$ , our “Patient 0” (hereafter P0) galaxy is a star forming galaxy with a disk (Figure 2a). P0 has an incoming satellite at  $z = 1$  with an original mass of  $7.34 \times 10^{10} M_\odot$  (mass ratio,  $q = 0.12$ ). For each GM galaxy simulation, we systematically shrink this satellite halo’s mass prior to its merger with the main halo. (Table 1) GM1 results in a similar disked, star forming galaxy as P0, while GM2 and GM3 become quenched at  $z \sim 1$  (Figure 2a).

are generated using the same method as While the GM galaxies utilize the same process as Pontzen et al. (2017), their study examines a different set

TABLE 1  
ZOOM-IN GALAXIES PROPERTIES *with BHs* AT  $z = 0.17$  *Values that have been updated are italicized*

Sim	Total Halo Mass ( $M_{\odot}$ )	Total Gas Mass ( $M_{\odot}$ )	Total Stellar Mass ( $M_{\odot}$ )	CGM Gas Mass ( $M_{\odot}$ )	$R_{vir}$ (kpc)	$T_{vir}$ (K)	Satellite Mass ( $M_{\odot}$ )
P0	$1.08 \times 10^{12}$	$1.09 \times 10^{11}$	$5.71 \times 10^{10}$	$9.32 \times 10^{10}$	268.9	$5.12 \times 10^5$	<b><math>7.34 \times 10^{10}</math></b>
GM1	$1.07 \times 10^{12}$	$1.01 \times 10^{11}$	$6.43 \times 10^{10}$	$8.39 \times 10^{10}$	269.2	$5.12 \times 10^5$	<b><math>5.86 \times 10^{10}</math></b>
GM2	$8.69 \times 10^{11}$	$7.41 \times 10^{10}$	$1.38 \times 10^{10}$	$6.89 \times 10^{10}$	254.1	$X \times 10^5$	<b><math>3.97 \times 10^{10}</math></b>
GM3	$7.76 \times 10^{11}$	$6.36 \times 10^{10}$	$1.04 \times 10^{10}$	$5.02 \times 10^{10}$	241.7	$4.14 \times 10^5$	<b><math>2.45 \times 10^{10}</math></b>

TABLE 2  
ZOOM-IN GALAXIES PROPERTIES *without BHs* AT  $z = 0.17$

Sim	Total Halo Mass ( $M_{\odot}$ )	Total Gas Mass ( $M_{\odot}$ )	Total Stellar Mass ( $M_{\odot}$ )	CGM Gas Mass ( $M_{\odot}$ )	$R_{vir}$ (kpc)	$T_{vir}$ (K)	Satellite Mass ( $M_{\odot}$ )
P0noBH	$8.36 \times 10^{11}$	$7.0 \times 10^{10}$	$7.37 \times 10^{10}$	$X \times 10^{10}$	262.8	$X \times 10^5$	<b><math>7.34 \times 10^{10}</math></b>
GM1noBH	$8.38 \times 10^{11}$	$7.07 \times 10^{10}$	$7.34 \times 10^{10}$	$X \times 10^{10}$	263.0	$X \times 10^5$	<b><math>5.86 \times 10^{10}</math></b>
GM2noBH	$8.42 \times 10^{11}$	$7.08 \times 10^{10}$	$7.33 \times 10^{10}$	$X \times 10^{10}$	263.4	$X \times 10^5$	<b><math>3.97 \times 10^{10}</math></b>
GM3noBH	$8.43 \times 10^{11}$	$7.19 \times 10^{10}$	$7.28 \times 10^{10}$	$X \times 10^{10}$	263.5	$X \times 10^5$	<b><math>2.45 \times 10^{10}</math></b>

of galaxies. The three galaxies in Pontzen et al. (2017) were run to  $z = 2$  and have  $M_{Halo} \sim 10^{12}$ . They each have incoming satellites whose masses are both increased and decreased prior to merging with the main galaxy, as in our galaxies; however, the resulting effect of their modifications were different from ours, as we explore in Section 4.

### 2.3.2. Galaxies without BH Physics

split into two sentences.

One key benefit of the individual zoom-in galaxies include the ability to remove or adjust the physical parameters affecting our galaxies to test different theoretical models which would be too computationally expensive to do with a large volume like R25. We utilize this benefit to further understand the effects of the AGN. To isolate the effect of the AGN on the CGM, all four of the zoom-in simulations (P0 and its 3 GMs) were re-simulated at the same resolution and with all the same physics *excluding* BH formation, feedback, and dynamical friction (Table 2). Black hole seed formation was disabled and the BH feedback and accretion efficiency parameters were set to 0. search and replace utilize with ANY OTHER WORD :)

In particular, we may exploit this utility to understand directly the effects of quenching in GM2 and GM3

break into two sentences.

Particularly of note, the top panel of Figure 2, which shows star formation histories of the four zoom-in galaxies with BH physics included, clearly shows that unlike P0 and GM1 which remain star forming throughout their history, GM2 and GM3 become quenched at  $z \sim 1$ . This immediate quenching just after the merger of the satellite with the main halo is particularly interesting because it does not take effect in the set of zoom-in galaxies without BH physics. Contrastingly, the lower panel of Figure 2 shows the star formation histories of the four zoom-in galaxies without BH physics and all four of their histories are nearly identical. The stark differences between the GM2 and GM3 galaxies with and without BHs imply that some interplay between the satellite's mass and the AGN feedback must play a pivotal role in quenching these galaxies so thoroughly.

We further examine the effects of the BH by looking to the mass buildup and accretion rate of the BHs. The accretion rates

Put redshift on top axes?





FIG. 2.— The star formation histories for the zoom-in galaxies: Patient 0 and its 3 GM galaxies with BH physics (*Upper*) and *Lower* without BH physics. In the galaxies including BH physics, P0 and GM1 remain star forming throughout their histories while GM2 and GM3 become quenched at  $z \sim 1$ . Without BH physics, all four galaxies remain star forming until  $z = 0$ . This is an effect we will explore in future work.

upper panel in Figure 3 shows the SMBH mass as a func-

AGN Feedback does NOT equal SMBH feedback...at least not as far as I'm aware. Let's make sure we're not using them interchangeably.

Overall, this new section 2.3.3 raises more questions than it answers, and we need to address \*some\* of them. a) we need to more clearly summarize results of Pontzen et al. (2017) earlier on, or first thing in this section. b) we really need to figure out why the black holes in GM2 and 3 grow so much more quickly than that in P0 and GM1 \*before\* the merger. c) we need to make sure satellite masses of all the GMs are really changed in exactly the same way as before when you turn off the black hole physics. and if so, we need to comment more on how disturbingly similar all of the SFHs are in GMs without black holes and what this means.





FIG. 3.— SMBH mass (*Upper*) and SMBH accretion rates (*Lower*) for our 4 zoom-in galaxies. Colors as in Figure 7. The SMBH growth of GM2 and GM3 occurs quicker than the growth of the SMBH in the two star forming zoom-in galaxies. In particular, GM3, which has the most significant modification to its satellite's mass, has a SMBH that grows quickest. Both quenched galaxies also have a sharp peak in accretion rate around the time of the merger with the satellite ( $z \sim 1$ ,  $t \sim 6$  Gyr), indicated by the dashed grey line.

tion of time. Here we see that the mass growth in the quenched galaxies, GM2 and GM3, occurs earlier in comparison to the star forming galaxies, especially in the case of GM3. A similar result can also be seen in lower panel of Figure 3 which depicts the SMBH accretion rate as a function of time. It is clear from this figure that an increase of accretion occurs near the time of the merger,  $z \sim 1$  or  $t \sim 6$  Gyr. It is clear from the BH's activity and growth that the effect of the mass of the incoming satellite has profound affect on the assembly history of this galaxy. Pontzen et al. (2017) previously explored the relationship between BH feedback and mergers and its effect on quenching, using the same genetic modification technique as we use for the GM galaxies in our study. They determine that AGN feedback is critical to quenching a galaxy, which is consistent with our finding that the only quenched galaxies are in simulations that include SMBHs. Pontzen et al. (2017) argues that the merger can disrupt the cold disk of the galaxy, which then allows the feedback of the AGN to have a farther reaching effect on the star forming gas of the disk thereby keeping

the galaxy in a state of quiescence. We note however, that the genetic modifications performed on the galaxies of (Pontzen et al. 2017) was different from the ones implemented here. In their case, it was an increase of the satellite's mass that resulted in a quenched galaxy, rather than a shrinking as we affect here. implement

This set of galaxies, which were produced from very similar initial conditions but which illustrate very different star formation and accretion histories, allows us to directly examine how assembly history may imprint itself on the CGM. However, the specific effects of the AGN's accretion on the assembly history of the galaxy is beyond the scope of this paper. We leave an examination of the quenching mechanisms of these galaxies to future work.

reword this...We're not really

### 3. RESULTS

#### 3.1. Qualitative CGM Properties

Individual halos in the ROMULUS25 cosmological volume and in the individual zoom-in galaxies are extracted using the Amiga Halo Finder (AHF) (Knollmann & Knebe 2009) and central SMBH positions and velocities are defined relative to the center position and inner 1 kpc center-of-mass velocity of their host halo, respectively. From R25, we specifically examine Milky Way-mass halos which are defined as halos between  $5 \times 10^{11}$  and  $2 \times 10^{12} M_{\odot}$  and are at least twice a virial radii from their nearest neighbor (to exclude any satellites). All zoom-in galaxies are isolated with a minor merger at  $z \sim 1$ .

The CGM of each individual galaxy halo (within the R25 galaxies and our zoom-ins) is defined as the mass enclosed in an annulus from 10 kpc from the center position out to a virial radius defined as the radius at which 200 times the critical density,  $\rho_c$ , where  $\rho/\rho_c = 200$ . Figure 4 shows the phase diagrams of the CGMs of the 4 zoom-in galaxies with and without BH physics. Examining the CGM phase diagrams for the GMs that include BH physics (*Left Column*), we note the following key differences: less overall mass from the uppermost (P0) to lowermost (GM3) figure; variations in the amount of cool, dense gas ( $T < 10^{4.5}$ ,  $n_H > 10^{-3}$ ); and the lack of hot, dense gas ( $T > 10^{5.5}$ ,  $n_H > 10^{-3}$ ) in the phase diagrams of GM2 and GM3, our quenched galaxies. Similarly, examining the CGM phase diagrams that exclude BH physics (*Right Column*), we don't see any significant differences. I am disturbed slightly. comment more on this.

The first of the differences between CGM phase diagrams with BH physics is unsurprising, as it correlates directly with the final halo mass of the galaxies. P0 results in the most massive halo at  $z \sim 0$ , while GM3 results in the least massive (Table 1). The second of these differences [FINISH ANALYSIS ON COOL DENSE TAIL].

We are most interested in the last difference: the lack of the hot, dense gas in the quenched galaxies that is present in the star forming galaxies, P0 and GM1. We note that this feature is also present in the CGM diagrams of the galaxies without BH physics, which all result in star forming, disked galaxies. Figure 5 further examines this difference with the same CGM phase diagrams of P0 and GM2 colored by mass, metallicity, and distance from the center of the galaxy, with (*Two Upper Rows*) and without (*Two Lower Rows*) BH physics. The hot, dense gas in P0 with BH physics (*Upper Row*) appears to

let's reference someone on halo mass of MW. Actually, maybe Gwen Eadie is a good person to ask!!

Do people think MW had a minor merger at  $z \sim 1$ ?

Let's say something of MW accretion / assembly history as well.

CGM gas mass correlates with galaxy total mass. not necessarily unsurprising!

Say something like: BH feedback is \*necessary\* but not sufficient on its own to cause a galaxy to quench.

not really  
unclear  
what  
you  
mean  
here.

italics




FIG. 4.— Phase diagrams of the temperature and density of the two star forming zoom-in galaxies, P0 (*Top row*) and GM1 (*Second row*), and the two quenched galaxies, GM2 (*Third row*) and GM3 (*Bottom row*). The phase diagrams of galaxies with BH hole physics vary quite widely between the star forming (P0 and GM1) and quenched cases (GM2 and GM3), particularly in the highest temperature and density gas. However, the phase diagrams of the galaxies without BH physics appear nearly identical, as do their star formation histories (Figure 2 lower panel).

Nice figure, nice caption. Don't need to refer to Fig 2 at end though.

be mostly comprised of high metallicity gas that is close to the disk ( $R < 50$  kpc). Further examining this gas, we find that 3 % of the CGM gas has metallicity  $Z > Z_{\odot}$  at  $z = 0.17$ . Furthermore, of this 3 %, 23 % is between 10 and 20 kpc from the center of the galaxies. For GM1, the CGM is comprised of 7.6 % gas with  $Z > Z_{\odot}$  with nearly 50 % of that gas between 10 and 20 kpc. Contrastingly, a negligible amount of the CGM of both GM2 and GM3 have  $Z > Z_{\odot}$  at  $z = 0.17$ . The CGMs of the four galaxies without BH physics also have small amounts of gas with  $Z > Z_{\odot}$ , from 0.1 % in P0noBH to 0.04 % in GM3noBH. These percentages of high metallicity gases in P0 and GM1 with BH physics point to metal exchange in the galaxy that is strongly dependent on the SMBH (which we discuss further in Section 3.3). It's interesting to note that the removal of the BH physics in all four zoom-ins makes little difference in the final gas phase properties of the main halo (Table 2). Nevertheless, the differences we see in the CGMs of these galaxies, both with and without BH physics, don't seem to have a significant effect on the OVI that we see in the galaxies, as we explore below.

nice.

this sentence  
is confusing.  
reward  
and/or  
break into  
two.

AND  
collisions  
I  
ionization  
equilibriu  
m!

Actually, it's  
TOO  
WEAK.  
start with  
Kollmeier  
et al. 2014  
of OVI as a function of radius (Need to change this  
then move to impact parameter; Current profiles are created  
to a Shull when galaxies are in a face-on orientation.) for our  
et al. 2015  
paper.  
Oh boy.  
This is a  
can of  
worms!  
Just say  
OVI is  
predomi  
nantly  
CI in  
our  
sims  
and  
choice  
of UVB  
does  
not  
impact  
our  
results  
directly.

**COS-Halos!**

The **COS-Halo Survey** dataset is plotted on top in black, with squares and circles distinguishing between elliptical and spiral galaxies. Upper and lower limits are designated with arrows and unfilled markers. The R25 galaxies well match the observations from the COS-Halo Survey; however, we note that they are systematically higher than the upper limits of the more massive ellipticals in the survey. (See 4) We further compare the column densities of OVI in the R25 galaxies to the 4 zoom-in galaxies with BH physics and find that these galaxies also well match the observed column densities of COS-Halo and fall within the range of the R25 galaxies (Figure 6b).

Figures 6 and 7 make it clear that our simulations reproduce the column densities of OVI in the CGM; however, this conclusion is not the only important feature of these plots. In addition, we note that the column densities of OVI in the CGMs of these galaxies does not seem to depend on the assembly history of the galaxy. We see this in both R25, which in addition to providing evidence

For the discussion — I think you will want to comment on metal budgets of galaxies (ala Peebles et al. 2014).  
Actually, it would be great to compare TOTAL MASS in OXYGEN in CGM AND in disk for BH cases vs no BH cases.

for this initial result also gives cosmological credence to our suite of GM galaxies, and our four GM galaxies that include BH physics. We confirm our result with Patient 0 and its GMs, which include two star forming galaxies and two quenched galaxies, all of which well match the observations of **COS-Halo** despite differing formation histories. (Figure 7) assembly stick with the same term for clarity.

Figure 8 shows the average ionization fractions for all the ions of oxygen within three mass ranges: low mass ( $5 \times 10^{10} - 5 \times 10^{11} M_{\odot}$ ), Milky Way-mass ( $5 \times 10^{11} - 2 \times 10^{12} M_{\odot}$ ), and high mass ( $2 \times 10^{12} - 2 \times 10^{13} M_{\odot}$ ). The OVI fractions decrease from the MW-mass range to the high mass regime due to the increase in virial temperature which moves from a value close to the ionization peak for OVI,  $\sim 10^{5.5} K$  to  $10^{6.3} K$ . From this study, we determine that morphological evolution of the galaxy doesn't correlate with the evolution of the CGM, as described by the column densities of OVI. Instead, it appears that the mass of the galaxy, and connectedly its virial temperature [Table 1], plays a more significant role in determining the amount of OVI seen in the CGM.

### 3.3. Metal Transport by the SMBH

We examine the column densities of OVI in the CGMs of our 4 zoom-in galaxies *without* BH physics and compare them to the cases where BH physics is included. Figure 7 shows the column densities of OVI in the CGM of all four of our zoom-in galaxies with BH physics (solid lines) and without (dashed lines). We can see that in the cases where BH physics is not included, the values of  $N_{OVI}$  are significantly lower implying that the presence of the AGN must play an important role in populating OVI in the CGM. We look to the temperature, mass, density, and metallicity of the CGM to investigate the cause of this decrease in OVI. (Figure 9)

$N_{OVI}$

The difference between the CGMs of these two cases appears to come directly from the change in metallicity due to the lack of black hole activity. (Figure 9) We examine the metallicity of the disk to look for further clues about how the lack of AGN activity is affecting the galaxy. Figure 10 shows that, in the galaxies without BH physics, the metals produced in the disk aren't driven into the CGM due to the lack of AGN feedback. This result is also consistent with what we see in Figure 5. The lack of high metallicity ( $Z > Z_{\odot}$ ) in the CGM phase diagrams of the galaxies with no BH physics (Right Column) confirms the fact that metals are not being driven out of the disk. Surprisingly, however, the feedback doesn't seem to play a role in significantly heating or excavating the CGM gas, but it appears the AGN's feedback is pivotal in transporting the metals from the center of the galaxy out into the CGM. The AGN plays a significant role in physically driving the metals out of the disk and into the outer regions of the CGM.

[Include discussion about metal flux into and out of galaxy (once complete); comparisons between observations and amount of total metals in disk (plus metal gradients)]

## 4. DISCUSSION

Questions to expound on: Limitations of the simulations Comparisons with other groups: Oppenheimer, Suresh, Hummels(Fire)? What questions remain/are raised by results

Maybe on this leftmost column, instead of M\_CGM, do M\_oxygen? See comment above on “metal budgets.”




FIG. 5.— Phase diagrams of the temperature and density of the star forming zoom-in galaxy, P0 (*Top row* with BH physics, *Middle row* without), and the quenched galaxy, GM2 (*Bottom row*). *Left:* As we noted, the phase diagrams of galaxies with BH hole physics show stark differences between the star forming (P0 and GM1) and quenched cases (GM2 and GM3), particularly in the highest temperature and density gas. *Middle:* The same phase diagram showing temperature and density, however, the colorbar is weighted by the average metallicity of the star in each bin. We note that the high density, high temperature gas we see in the star forming P0, is also the highest metallicity gas in the CGM. *Right:* Similarly, a phase diagram with the colorbar now weighted by the average distance from the center of the galaxy of the gas particles in each bin. The concentration of gas at high density and temperature *and* high metallicity also appears to be the gas closest to the center of the galaxy. However, this gas is also not present in the noBH physics case. We conclude that due to the active star formation in P0 in tandem with the AGN physics at play, this excess of gas comes from gas that has been ejected from the disk due to AGN feedback. Therefore explaining it’s significant absence from the phase diagrams of P0 without BH physics and GM2.

No conclusions and/or analysis in a figure caption!




FIG. 6.— *Left:* Mean column densities of OVI as a function of radius for all Milky Way mass halos in the R25 simulation. Blue and red lines distinguish between disk dominated spirals and quenched elliptical galaxies within the R25 simulation. *Right:* Column densities of OVI in our 4 zoom-in galaxies. Grey lines indicate R25 MW galaxy column densities from *Left*. Blue solid lines describe our two star forming galaxies, P0 and GM1. GM4-7, our passive galaxies, are in solid red. Filled circles and squares indicate spirals and ellipticals from the COS-Halo Survey dataset. Unfilled squares indicate upper limits.




FIG. 7.— Column density profiles of OVI in our 4 zoom-in galaxies with (solid lines) and without (dashed lines) BH physics. P0 and GM1, our two star forming galaxies are colored as light blue and dark blue, respectively. Our quenched galaxies, GM2 and GM3, are labeled in dark red and pink, respectively.

## 5. CONCLUSION

### Result 1: OVI as a Tracer for Virial Temperature of the Halo

The combined, consistent results of the cosmological R25 and our 4 zoom-in galaxies (which include BH physics) imply a mechanism by which column densities of OVI are set by the virial temperature of the CGMs host galaxy. They aren't affected by the evolution of a disk. Their phase diagrams also lack significant difference in their overall assembly history, except where more gas is clearly present in the higher mass galaxies. Therefore, we surmise that the differences in the CGM are not determined by whether or not a galaxy quenches but rather these conditions for OVI are primarily set by the virial temperature of the galaxy.

These results are consistent with those of Oppenheimer et. al. 2016 who used a suite of EAGLE simulated galaxies to examine the bimodality of OVI column densities (further discussed in [Tumlinson 2011]) in star forming




FIG. 8.— Oxygen ion fractions in the CGM of our 25 MW-mass galaxies and 11 high mass galaxies from R25. Log  $M_{\text{halo}}$  for each halo are labeled in white text. [Under construction]

and quenched galaxies. They argue that the star forming galaxies ( $10^{11} - 10^{12}$ ), which were found to have a higher fraction of OVI, were at the right virial temperature to maximize OVI production, while their quenched galaxies ( $10^{12} - 10^{13}$ ) had high enough virial temperatures such that the dominant ionization state was not OVI but rather OVII or above. Oppenheimer et. al. 2016 argues that the OVI content was not a tracer of star formation directly, but rather a more direct thermometer for the temperature of the halo.

We note that the quenched galaxies in our sample are smaller in mass than our star forming galaxies, unlike those in Oppenheimer, explaining the lack of bimodality that we observe. While all the GMs are in the mass range to have virial temperatures which optimize OVI, we further examine the R25 simulation's higher mass, passive galaxies in addition to the MW-mass galaxies (which have virial temperatures spanning  $5.8 \times 10^5$  K to  $1.1 \times 10^6$  K) to see if the bimodality appears. (Figure 8) We determine that the OVI still provides a direct thermometer for the temperature of the halo. [Under

JW stopped detailed comments here. You have enough to work on before revisiting these conclusions :)




FIG. 9.— Temperature, Total Mass, Total Density, and Metallicity profiles of the CGM of our 4 zoom-in galaxies with and without BH physics. Colors and line styles as in Figure 7.




FIG. 10.— Metallicity profile of the gas within the disk of our 4 zoom-in galaxies with and without BH physics. Colors and line styles as in Figure 9. Without the black hole physics, metals remain trapped near the center of the disk with no mechanism to propagate out into the CGM.

#### construction]

Furthermore, examining galaxies with masses larger than our MW-mass GMs ( $>2 \times 10^{12}$ ) from the R25 suite,

we see that the column densities of OVI decrease as the ionization peak of OVI is surpassed by these halos. Since the virial temperature is higher, the oxygen is likely to be ionized to a higher ionizations state (OVII or OVII), which we show is the case in Figure 8. [Under construction]

#### Result 2: AGN as driver for metals in the CGM

Our result that the AGN acts a physical driver for metals in the CGM has interesting consequences. Previous studies have examined the effect of heating on the CGM as the AGNs energy input may put the gas into phases which optimize the production of OVI. (Suresh et al. 2017) Others have proposed that the feedback from AGN may physically drive outflows of gas out of the galaxy, resulting in a lower density CGM and therefore lower densities of OVI. Neither of these cases are what we see. Instead, we see a suite of CGM which rely on the AGN for the propagation of metal mass (but not total gas mass) into the outer galaxy and OVI columns which depend on the virial temperature of the galaxy. (Figure 10)

#### Acknowledgements

#### REFERENCES

- Governato, F., Weisz, D., Pontzen, A., et al. 2015, Monthly Notices of the Royal Astronomical Society, 448, 792 [2.1](#)
- Haardt, F., & Madau, P. 2012, The Astrophysical Journal, 746, 125 [2.1](#), [3.2](#)
- Knollmann, S. R., & Knebe, A. 2009, The Astrophysical Journal Supplement Series, 182, 608 [3.1](#)
- Mcconnell, N. J., & Ma, C.-P. 2013, The Astrophysical Journal, 764, 184 [1](#)
- Menon, H., Wesolowski, L., Zheng, G., et al. 2015, Computational Astrophysics and Cosmology, 2, 1 [2.1](#)
- Pontzen, A., Tremmel, M., Roth, N., et al. 2017, Monthly Notices of the Royal Astronomical Society, 465, 547 [1](#), [2.3](#), [2.3.1](#), [2.3.3](#)
- Ritchie, B. W., & Thomas, P. A. 2001, Monthly Notices of the Royal Astronomical Society, 323, 743 [2.1](#)
- Roth, N., Pontzen, A., & Peiris, H. V. 2016, Monthly Notices of the Royal Astronomical Society, 455, 974 [1](#)
- Shen, S., Wadsley, J., & Stinson, G. 2010, Monthly Notices of the Royal Astronomical Society, 407, 1581 [2.1](#)
- Stinson, G., Seth, A., Katz, N., et al. 2006, Monthly Notices of the Royal Astronomical Society, 373, 1074 [2.1](#)
- Stinson, G. S., Brook, C., Prochaska, J. X., et al. 2012, Monthly Notices of the Royal Astronomical Society, 425, 1270 [3.2](#)
- Suresh, J., Rubin, K. H. R., Kannan, R., et al. 2017, Monthly Notices of the Royal Astronomical Society, 465, 2966 [1](#), [5](#)
- Tremmel, M., Governato, F., Volonteri, M., & Quinn, T. R. 2015, Monthly Notices of the Royal Astronomical Society, 451, 1868 [1](#), [2.1](#), [2.3](#)
- Tremmel, M., Karcher, M., Governato, F., et al. 2017, Monthly Notices of the Royal Astronomical Society, 470, 1121 [\(document\)](#), [1](#), [2.1](#), [2.2](#)
- Tumlinson, J., Peebles, M. S., & Werk, J. K. 2017, Annual Review of Astronomy and Astrophysics, 55, 389 [1](#)
- Tumlinson, J., Thom, C., Werk, J. K., et al. 2011, Science, 334, 948 [1](#)
- Wadsley, J., Stadel, J., & Quinn, T. 2004, New Astronomy, 9, 137 [2.1](#)
- Wadsley, J. W., Veeravalli, G., & Couchman, H. M. P. 2008, Monthly Notices of the Royal Astronomical Society, 387, 427 [2.1](#)
- Werk, J. K., Prochaska, J. X., Thom, C., et al. 2012, The Astrophysical Journal Supplement Series, 198, 3 [1](#)
- . 2013, The Astrophysical Journal Supplement Series, 204, 17 [1](#)
- Werk, J. K., Prochaska, J. X., Tumlinson, J., et al. 2014, The Astrophysical Journal, 792, 8 [1](#)
- Werk, J. K., Prochaska, J. X., Cantalupo, S., et al. 2016, 24 [1](#)