

Early Black-Hole Growth Constraints from JWST

Nicole Jiang

Draft: 11 September 2026

Abstract

Aims. Massive black holes in the early Universe must grow within limited cosmic time. We identify which JWST black-hole candidates most strongly constrain early growth models and which assumptions control these constraints. **Methods.** We compile 350 literature measurement records from 32 source families at $z \geq 4$. We analyse 224 objects with broad hydrogen-line mass estimates and include 10 additional objects in a broader comparison. We propagate reported mass uncertainties and test mass-scale shifts and alternative growth assumptions. **Results.** For $100 M_{\odot}$ seeds formed at $z = 30$, radiative efficiency $\epsilon = 0.1$, and no merger boost, 12 of the 224 objects require a lifetime-average Eddington ratio $\overline{f_{\text{Edd,req}}} > 1$. Of these, 6 exceed this threshold in at least 95% of trials sampling their quoted mass errors under fixed growth assumptions. For UNCOVER-20466, COSMOS3D-13852 and RUBIES-EGS-55604, at least 95% of trials still require an average Eddington ratio above one after reducing each mass by 0.5 dex (a factor of approximately 3.2). Uncertainties in the mass estimators also affect these constraints. Delaying seeding to $z = 20$ makes GN-z11 the most demanding object. For A2744-QSO1, an independent dynamical mass estimate raises the required average Eddington ratio to approximately one. **Conclusions.** At $\epsilon = 0.05719$, the ideal thin-disc efficiency for a non-spinning black hole, all 234 central mass estimates are reachable from $100 M_{\odot}$ seeds formed at $z = 30$ without lifetime-average accretion above the Eddington limit.

Key words. black hole physics – accretion, accretion disks – galaxies: active – galaxies: high-redshift – quasars: supermassive black holes

1 Introduction

Massive black holes observed in the early Universe must have assembled within the available cosmic time. The strength of this constraint depends on the observed black-hole mass and redshift and on assumptions about seed formation, radiative efficiency, and the duration of active accretion. Different seed and efficiency assumptions can therefore give different required accretion rates for the same object.

We use a literature-based catalogue of JWST-identified high-redshift accreting black holes and candidates to test which combinations of seed mass, formation time, radiative efficiency and accretion rate can reproduce the observed masses by the observed redshifts. The atlas covers $z \geq 4$, extending from the first billion years to a cosmic age of approximately 1.54 Gyr at its lower-redshift boundary. We calculate the average accretion rates required across a range of seed and growth assumptions, incorporating reported mass uncertainties and accounting for differences in detection evidence and mass-estimation methods. We retain published measurements, object identity links and mass-estimation sources so that measurements from different studies can be compared.

Analytic seed-growth comparisons already establish the trade-off between seed mass and sustained accretion. For example, [P. Dayal \(2024\)](#) considered light and heavy seeds at $z_{\text{seed}} = 25$ and explored a primordial-seeding interpretation. We apply the same growth equations to a larger catalogue and test how each object’s constraints depend on its published mass, measurement method and growth assumptions. We calculate whether each black hole could reach its observed mass while accreting, on average, at or

below the Eddington limit. We repeat this calculation using masses drawn from the published uncertainty intervals, uniform shifts in the mass estimates, and alternative published measurements. These tests show how the inferred accretion rate depends on mass uncertainty and the method used to estimate the mass. For $100 M_{\odot}$ seeds formed at $z = 30$, radiative efficiency $\epsilon = 0.1$ and no mass added by mergers, 12 objects in our Balmer-line sample require an average Eddington ratio above one. We list all 12, describe the limitations reported for their mass estimates, and propose observations to test those estimates. For four objects also studied by Dayal, we calculate separately how the required accretion rates change when we update the measured masses and redshifts and when we start growth earlier. For A2744-QSO1, we repeat the calculation with an independent dynamical mass estimate. We keep the original mass estimate in the catalogue and report the dynamical comparison separately. The JADES broad-line census (I. Juodžbalis et al. 2026a) measures current accretion and host scaling relations, and the narrow-line census (J. Scholtz et al. 2025) identifies candidates for obscured activity. We use these measurements to calculate the cumulative growth needed to reach the adopted masses. The contributing studies targeted different kinds of objects and reached different detection limits. The probability that a black hole would be selected and detected therefore varies between studies and is unknown for the combined sample. The source-family inventory is given in Appendix A.

1.1 Theoretical framework

We use the luminosity-based Eddington ratio $f_{\text{Edd}} = L_{\text{bol}}/L_{\text{Edd}}$ and a fixed radiative efficiency ϵ , the fraction of accreted rest-mass energy emitted as radiation. For the available growth interval,

$$\Delta t = t(z_{\text{obs}}) - t(z_{\text{seed}}) > 0, \quad \overline{f_{\text{Edd}}} = \frac{1}{\Delta t} \int_{t(z_{\text{seed}})}^{t(z_{\text{obs}})} f_{\text{Edd}}(t) dt. \quad (1)$$

The analytic growth relation adopted by P. Dayal (2024) gives

$$M_{\text{BH}}(t_{\text{obs}}) = B_{\text{merge}} M_{\text{seed}} \exp \left[\frac{1 - \epsilon}{\epsilon} \frac{\overline{f_{\text{Edd}}} \Delta t}{t_{\text{Edd}}} \right], \quad (2)$$

where $t_{\text{Edd}} \simeq 0.45$ Gyr. The merger multiplier is $B_{\text{merge}} = 1$ for the reference calculation and 2 for a factor-of-two mass boost. We represent mergers by multiplying the seed mass by a constant factor before calculating accretion. This approximation omits the timing and sequence of mergers, their mass ratios, and mass lost through gravitational radiation. The required lifetime-average ratio is

$$\overline{f_{\text{Edd}}}_{\text{req}} = \max \left[0, \frac{\epsilon}{1 - \epsilon} \frac{t_{\text{Edd}}}{\Delta t} \ln \left(\frac{M_{\text{BH}}}{B_{\text{merge}} M_{\text{seed}}} \right) \right]. \quad (3)$$

If $B_{\text{merge}} M_{\text{seed}}$ exceeds the observed black-hole mass, the logarithm is negative and we set the required average Eddington ratio to zero. Even with zero accretion, this seed and merger contribution predict a mass larger than observed. Reproducing the observed mass would require a smaller seed, a smaller merger contribution, or a model that includes mass loss. This requirement averages accretion over the full growth interval. Inferring a duty cycle also requires assumptions about the rates during active and quiet phases.

We calculate cosmic ages in a spatially flat Λ CDM model, using the Planck-based parameters quoted by P. Dayal (2024): $H_0 = 67.3 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.315$ and $\Omega_{\Lambda} = 1 - \Omega_m = 0.685$. The age calculation includes matter and a cosmological constant and neglects radiation. Spatial flatness fixes $\Omega_{\Lambda} = 1 - \Omega_m$ in this approximation. Appendix D gives the supporting relations, implementation and inverse seed-mass diagnostic. Equations 2–3 assume constant radiative efficiency. The compatibility calculations use fixed spin values and the efficiency prescription in Appendix C.

2 Catalogue construction

We compile 350 literature measurements from 32 source families at $z \geq 4$. Each adopted mass is linked to its source, method and reported uncertainty. We select one published measurement per object using the source assessments and retain the alternatives for sensitivity tests.

Growth analysis requires a numerical mass and redshift, an identified mass method and a resolved lensing treatment. Upper limits, indirect model ranges and masses inferred by assuming an Eddington ratio are excluded from numerical inference. We also exclude all six records associated with three unresolved identity groups, including three records with eligible masses. This leaves 224 primary objects and an inclusive exploratory sample of 234 objects.

For the main comparison, we use masses estimated from broad $H\alpha$ or $H\beta$ emission lines in objects with secure or probable evidence for an accreting black hole. We exclude objects whose mass estimate depends on an unresolved interpretation of the broad emission as a gravitationally bound broad-line region. We refer to the resulting 224 objects as the primary sample. These single-epoch virial estimators combine the line width, which estimates gas velocity, with a luminosity-based estimate of the broad-line region’s radius. Their calibration is based on reverberation mapping of nearby AGN (Y. Shen 2013). Using the same line family and mass-estimation framework reduces differences between methods in the comparison. Applying these calibrations at high redshift still requires assessing gas dynamics, geometry, obscuration and lensing for each object. We also analyse an expanded sample of 234 objects, labelled exploratory in the tables and figures. It includes the 224 primary objects, nine additional candidates, and GN-z11. GN-z11’s mass is inferred from rest-frame ultraviolet emission lines using local virial relations (R. Maiolino et al. 2024). We include it in the expanded comparison because its lines and calibration differ from those of the Balmer-line sample. The sample division reflects the evidence for accretion and the methods being compared; the reliability of each mass estimate requires an individual assessment. When a paper gives a mass without an uncertainty, we can calculate the average Eddington ratio needed to reach that mass. We cannot calculate the probability that this ratio exceeds one from the published information, because no mass uncertainty distribution is available. We therefore include these objects in calculations at their published masses and omit them from probability calculations. The mass estimates retained in each sample still use different calibrations, and the contributing studies have different target-selection criteria and detection limits. Appendix A gives the source inventory, identity decisions and sample accounting (Figure 6).

3 Growth-model comparison

3.1 Reference model

We calculate the required lifetime-average Eddington ratio using Eq. 3, with $M_{\text{seed}} = 10^2 M_{\odot}$, $z_{\text{seed}} = 30$, $\epsilon = 0.1$ and $B_{\text{merge}} = 1$ as the reference history. We rank objects by this required average Eddington ratio. Figure 1 illustrates tracks at four fixed efficiencies; tracks are selected for proximity to the observed masses. Appendix D.6 specifies the display selection.

3.2 Uncertainty and sensitivity

We use Monte Carlo sampling to propagate reported asymmetric log-mass errors, with 10,000 draws per object. Each draw has equal probability of falling above or below the central mass, with its distance drawn from a half-normal distribution scaled by the corresponding upper or lower error. Redshifts are held fixed. For each sampled mass, we calculate the average Eddington ratio needed to grow from the assumed seed to that mass. The fraction of the 10,000 Monte Carlo draws with a required ratio above one estimates the

probability that growth requires an average rate above the Eddington limit, given the adopted mass-error distribution and growth assumptions. The draws use the quoted errors only. Probability calculations include 212 primary and 222 expanded-sample objects; the 12 objects without reported errors enter only calculations at their published masses.

We test mass-scale sensitivity by shifting all adopted log masses and their error draws by offsets of -0.5 , -0.3 , 0 , $+0.3$ and $+0.5$ dex, keeping the sample and growth assumptions fixed. These fixed offsets are stress tests at the uncertainty scale of single-epoch mass estimates (Y. Shen 2013). For each offset, we report the fraction of shifted Monte Carlo draws that require an average Eddington ratio above one. Each probability assumes that particular mass shift and the stated seed mass, starting redshift, radiative efficiency and merger contribution. We assign no probability to the offsets and report their results separately. Sampling and conversion details are given in Appendix D.6.

3.3 Compatibility across growth assumptions

For each assumed accretion rate, efficiency, starting redshift and merger contribution, we solve Eq. 15 for the initial seed mass that would grow to the observed black-hole mass. We then check whether that seed falls in each of four mass ranges: $10\text{--}100 M_\odot$ (light), $10^3\text{--}10^4 M_\odot$ (intermediate), $10^4\text{--}10^6 M_\odot$ (heavy), and $10^2\text{--}10^6 M_\odot$ (broad). We count an object as compatible with a range when some seed within it can reproduce the observed mass under these assumptions. The grid uses $z_{\text{seed}} = 30$, three spin/efficiency cases, merger multipliers 1 and 2, and average rates 0.3, 1 and 2. These maps use the supercritical efficiency prescription in Appendix C.

The seed-mass ranges overlap and are tested separately, without probability weights. The broad range spans several possible seed-formation channels. If the seed mass needed to reproduce an object is smaller than the lower bound of a tested range, even the smallest seed in that range would grow larger than the observed black hole at the assumed accretion rate. That object is therefore excluded from the count for that range and set of growth assumptions. For each set of assumptions, we report the percentage of catalogue objects whose observed masses can be reproduced from a seed in the tested range. These percentages describe this literature sample. Estimating the corresponding percentages among all high-redshift black holes would require accounting for how each contributing study selected and detected its objects.

4 Results


4.1 Required accretion under the reference assumptions

For a $10^2 M_\odot$ seed under the reference history, the 224-object primary sample has median required $f_{\text{Edd}} = 0.578$ (interquartile range 0.487–0.682), with 12 point requirements above unity and none above two. The inclusive 234-object sample has median 0.574 (0.488–0.679) and 14 above unity. UNCOVER-20466 is most demanding: its point requirement is 1.452 and its reported-error median is 1.453 (16th–84th percentiles 1.354–1.549). Table 1 lists the 12 primary threshold objects; their ordering by point requirement differs from ordering by quoted-error probability.

The shorter growth time at $z = 10.603$ places GN-z11 second in the required-rate ranking despite its smaller mass. Its UV virial estimator places it in the exploratory sample.

Growth tracks across radiative-efficiency assumptions

$z_{\text{seed}} = 30$; bands span $B = 1$ (thick edge) to $B = 2$ (thin edge). Seed mass is encoded by colour; exact rates are listed above each panel.



Grey circles: primary (224); purple triangles: exploratory only (10). Outlined targets are named in panel (a).

Figure 1: Growth tracks with 224 primary and ten exploratory-only objects. Panels show four constant radiative efficiencies at $z_{\text{seed}} = 30$; colours distinguish seeds of 10^2 , 10^4 and $10^5 M_{\odot}$, and legends list the selected average Eddington ratios. Shaded bands span merger boosts $B = 1$ – 2 , with thick and thin edges respectively. The bands show the effect of the assumed merger boost. Six named objects are outlined in every panel. The reversed redshift axis spans $z = 11.5$ to $z = 3$. Excluded identities and objects without eligible masses are omitted.

4.2 Robustness to reported mass uncertainty

Of the 12 primary point requirements above unity, eight remain above it at the 16th percentile and six have conditional $P(f_{\text{Edd}} > 1) \geq 0.95$ (Fig. 2). The exploratory counts are 14 at the median, ten at the 16th percentile and eight at $P \geq 0.95$; 15 exceed unity by the 84th percentile. Probabilities follow Section 3.2 and use 212 primary and 222 exploratory objects with reported errors. The 12 primary objects without errors enter only point counts.

4.3 Sensitivity to the virial-mass scale

Downward mass shifts of 0.3 and 0.5 dex reduce the primary point/16th-percentile/ $P \geq 0.95$ counts from 12/8/6 to 9/6/4 and 6/4/3, respectively (Table 2). UNCOVER-20466, COSMOS3D-13852 and RUBIES-

Table 1: The 12 primary objects above the reference threshold, ordered by their point requirements. Masses and errors are in $\log_{10}(M/M_{\odot})$; all estimators are single-epoch virial. f_0 is the reference point requirement, f_{16} its reported-error 16th percentile, and $P_0 = P(\overline{f_{\text{Edd}_{\text{req}}}} > 1)$. f_- and P_- apply the fixed -0.5 dex offset. All probabilities are conditional on the mass-error model and growth assumptions. Source-specific caveats and proposed observations are in Table 6.

Object/source	Line	Log mass	f_0	f_{16}	P_0	f_-	P_-
UNCOVER-20466 J. E. Greene et al. (2024)	H β	$8.17^{+0.42}_{-0.42}$	1.452	1.354	> 0.999	1.335	> 0.999
GS-20057765 I. Juodžbalis et al. (2026a)	H β	$7.33^{+0.62}_{-0.70}$	1.355	1.177	0.980	1.228	0.901
COSMOS3D-13852 X. Lin et al. (2026)	H β	$8.60^{+0.35}_{-0.33}$	1.314	1.247	> 0.999	1.214	> 0.999
RUBIES-EGS-55604 D. D. Kocevski et al. (2025)	H β	$9.31^{+0.10}_{-0.10}$	1.267	1.250	> 0.999	1.180	> 0.999
CEERS-1019 R. L. Larson et al. (2023)	H β	$6.95^{+0.37}_{-0.37}$	1.205	1.115	0.988	1.083	0.822
GS-20030333 I. Juodžbalis et al. (2026a)	H β	$7.42^{+0.65}_{-0.48}$	1.133	1.033	0.907	1.029	0.613
GS-164055 I. Juodžbalis et al. (2026a)	H β	$7.63^{+0.59}_{-0.66}$	1.065	0.941	0.696	0.970	0.395
GN-38509 I. Juodžbalis et al. (2026a)	H α	$8.57^{+0.37}_{-0.38}$	1.065	1.005	0.860	0.984	0.392
J0910_2028_12910 C. J. Baccus & X. Xu (2026)	H α	$8.58^{+0.01}_{-0.01}$	1.053	1.051	> 0.999	0.973	< 0.001
CEERS-00717 Y. Harikane et al. (2023)	H α	$7.99^{+0.16}_{-0.17}$	1.027	0.998	0.821	0.942	0.018
ZS7 H. Übler et al. (2024)	H β	$7.70^{+0.40}_{-0.40}$	1.023	0.954	0.625	0.934	0.173
GN-4685 I. Juodžbalis et al. (2026a)	H β	$7.36^{+0.45}_{-0.42}$	1.017	0.938	0.585	0.922	0.181

EGS-55604 retain $P \geq 0.95$ at -0.5 dex. Independent mass estimates would test these results. Biases specific to each source may exceed this offset (Section 5.2), and the required rates still depend on seed mass, formation time and radiative efficiency.

Table 2: Fixed mass-scale stress tests under the reference growth history. Triplets give counts above unity at the point estimate, at the 16th percentile, and with conditional $P(\overline{f_{\text{Edd}_{\text{req}}}} > 1) \geq 0.95$. The probability-based counts use 212 primary and 222 exploratory objects with reported errors; point counts use all 224 and 234 objects.

Mass offset (dex)	Primary	Exploratory
-0.5	6 / 4 / 3	7 / 5 / 4
-0.3	9 / 6 / 4	10 / 7 / 5
0	12 / 8 / 6	14 / 10 / 8
$+0.3$	15 / 13 / 12	17 / 15 / 14
$+0.5$	19 / 16 / 13	21 / 18 / 15

Replacing the redshifts, masses and errors of 44 Baccus counterparts matched by identifier and confirmed by position with published Table 1 values (C. J. Baccus & X. Xu 2026) preserves the primary 12/8/6 and exploratory 14/10/8 counts and both top-five rankings (Table 3). The 32 changed central masses shift by -0.10 to $+0.04$ dex. Of five unmatched records, one is identity-excluded; retaining or omitting the other four leaves these results unchanged (220/230 objects when omitted). The test updates the matched

Objects above the reference growth threshold




Figure 2: The 12 primary objects whose reference point requirements exceed unity. Circles and bars show reported-error medians and 16th–84th percentile intervals; diamonds show point requirements after a -0.5 dex mass shift. The vertical line is the reference threshold. The selected objects all have reported errors; these intervals do not include additional estimator systematics.

measurements while keeping all six identity exclusions, method classes, growth parameters and sampling rules fixed.

Table 3: Effect of adopting the published Baccus measurements after excluding unresolved identities. Every row excludes all open identity groups. Counts denote point requirements above unity, 16th percentiles above unity, and conditional $P \geq 0.95$, respectively. The adopted catalogue values define the baseline.

Sample	Measurement treatment	Objects	Counts
Primary	Frozen values	224	12/8/6
Primary	Published; retain unmatched	224	12/8/6
Primary	Published; omit unmatched	220	12/8/6
Exploratory	Frozen values	234	14/10/8
Exploratory	Published; retain unmatched	234	14/10/8
Exploratory	Published; omit unmatched	230	14/10/8

4.4 Model compatibility and measurement choice

Increasing the seed mass to $10^3 M_\odot$ reduces the primary count above unity from 12 to four; $10^5 M_\odot$ seeds remove all point exceedances. Delaying seeding to $z = 20$ raises the primary count to 22, while $\epsilon = 0.05719$ removes all point exceedances in both samples (Table 4). Each test changes one assumption: seed mass (P. Dayal 2024), available growth time, or the efficiency of an ideal thin disc around a non-spinning black hole (Eq. 7). We assign no population weights to these scenarios.

Removing the six ambiguous records from the original 227/237-object sets preserves threshold counts and

Table 4: Point-estimate requirements above $\overline{f_{\text{Edd}}} = 1$ under fixed histories. Each row changes only the named parameter from the reference ($M_{\text{seed}} = 10^2 M_{\odot}$, $z_{\text{seed}} = 30$, $\epsilon = 0.1$, $B_{\text{merge}} = 1$). Counts use central mass estimates and provisional object identities.

Scenario	Primary (224)	Exploratory (234)
Reference	12	14
$M_{\text{seed}} = 10^3 M_{\odot}$	4	5
$M_{\text{seed}} = 10^5 M_{\odot}$	0	0
$z_{\text{seed}} = 20$	22	24
$\epsilon = 1 - \sqrt{8/9}$	0	0

top-five rankings in every listed scenario, including the reference probability counts. The primary median changes from 0.574 to 0.578; the objects above the reference threshold remain the same.

Figure 3 shows the maximum fixed efficiency allowing $\overline{f_{\text{Edd}}}_{\text{req}} \leq 1$ for five targets as a function of seed mass. The $z_{\text{seed}} = 20$ and 30 curves show GN-z11’s greater sensitivity to the starting time. These boundaries use central mass estimates and assume no merger boost. Sustaining the required rate also depends on the fuel supply.




Figure 3: Seed-mass–efficiency boundaries for Eddington-limited average growth at two starting redshifts. Each curve uses an adopted point mass and no merger boost. Below a curve the required average is at most unity. Horizontal guides mark $\epsilon = 0.1$ (solid grey) and $1 - \sqrt{8/9}$ (dotted grey). GN-z11 (dashed) is exploratory; the other objects are primary. Each curve holds efficiency constant and has no assigned probability weight.

The primary and exploratory samples show similar compatibility patterns (Fig. 4). For heavy seeds at $a = +1$ and $f = 1$, changing B_{merge} from one to two raises compatibility from 57 to 70 per cent in the primary sample (59 to 71 per cent inclusive). At higher accretion rates, even the smallest seed in a tested mass range can grow larger than the observed black hole. Such objects reduce the percentage reproduced from seeds in that range. These fractions use central mass estimates and the efficiency prescription in Appendix C. They describe compatibility within the selected sample; estimating the frequency of each formation channel would require uncertainty propagation and a population selection model.

Compatibility across growth assumptions




Figure 4: Constant-state compatibility fractions in the primary sample (left) and the inclusive exploratory sample (right). The exploratory sample includes all primary objects. Each column specifies merger multiplier B and fixed luminosity ratio f ; rows specify ideal spin-dependent efficiency, with the supercritical prescription in Appendix C. Seed intervals overlap and have no assigned probability weights. The broad mass interval spans several possible seed-formation channels.

Seven alternate-measurement comparisons across six objects (two for CEERS-2782) span mass shifts of -0.600 to $+0.949$ dex and required- f_{Edd} shifts of -0.083 to $+0.121$ (Fig. 5). Both estimates remain below unity in every pair. This comparison covers six objects below the threshold. Testing the objects above it requires additional independent mass estimates.

4.5 Separating measurement and starting-time changes

For four unambiguous matches to Table 1 of P. Dayal (2024), Table 5 first replaces the earlier masses and redshifts with our adopted values at $z_{\text{seed}} = 25$, then changes seeding to $z = 30$. All columns use our cosmology, $\epsilon = 0.1$, $10^2 M_\odot$ seeds and no merger boost. Earlier seeding lowers the requirements more than the measurement updates change them. The comparison isolates these two effects for the four matched objects using our growth calculation.

Growth requirements from alternate mass estimates




Figure 5: Preferred (circles) and alternate (diamonds) point requirements for seven measurement pairs across six primary objects. Pair numbers follow the released alternate-measurement table, which supplies stable measurement IDs; RUBIES-EGS-50052 (CEERS-2782) has two alternatives. All pairs remain below the reference threshold.

Table 5: Measurement and starting-time comparison for four objects shared with the Dayal study. Columns give required average $\bar{f}_{\text{Edd}}$; the first two share $z_{\text{seed}} = 25$. The analysis files provide the input measurements and their source locations.

Object	Earlier inputs, 25	Adopted inputs, 25	Adopted inputs, 30
GN-z11	1.577	1.578	1.437
A2744-QSO1	0.974	0.974	0.929
UNCOVER-20466	1.540	1.548	1.452
CEERS-00717	1.074	1.076	1.027

4.6 External dynamical-mass check: A2744-QSO1

For A2744-QSO1, we compare the virial mass adopted in the catalogue $\log_{10}(M/M_{\odot}) = 7.3 \pm 0.2$ (J. E. Greene et al. 2024) with the independent MOKA3D dynamical estimate 7.7 ± 0.3 (I. Juodžbalis et al. 2026b), using $z = 7.04$ and the reference history. We retain the values from [arXiv:2508.21748v2](#); the reference lists the subsequent journal publication. The dynamical estimate depends on lens reconstruction and inclination. We propagate the quoted two-sided mass error; the separate inclination-free lower limit is a one-sided constraint. Using the central mass estimates, the required average Eddington ratio rises from 0.929 to 1.000. With a symmetric normal approximation to the quoted log-mass errors, the exact 16th–84th percentile intervals are 0.895–0.964 and 0.947–1.052, respectively. For the dynamical mass, the unrounded required ratio is just below one. Its uncertainty interval includes average accretion rates both below and above the Eddington limit. With this mass-error distribution and the growth assumptions held fixed, the probability that the required average Eddington ratio exceeds one is 0.497. This comparison tests the mass of one catalogue object. Calibrating the virial sample would require independent measurements for more objects.

5 Discussion

5.1 Dependence on seed and growth assumptions

The objects above the reference threshold lie at $z = 6.6214\text{--}10.603$; most catalogue objects lie at lower redshift and serve as the comparison sample. Their growth requirements constrain combinations of seed mass, starting time, efficiency and accretion. Several formation histories satisfy these constraints.

At fixed seed, growth interval and merger factor, required f_{Edd} scales as $\epsilon/(1 - \epsilon)$. Lowering ϵ from 0.1 to 0.05719 multiplies all requirements by 0.54594: the largest becomes 0.793. The requirement for super-Eddington growth therefore depends on the assumed efficiency. Several spin and seed combinations can reproduce the same mass. Larger seed masses, earlier growth onset and merger contributions also reduce the required accretion rate.

Table 6 lists observations that could test the mass estimates of the objects above the reference threshold. Independent masses and constraints on scattering, outflows, obscuration and lensing can distinguish these interpretations. We propose these observations based on the known limitations of each mass estimate. The quoted-error probabilities omit any unreported estimator bias.

5.2 Estimator credibility and observational tests

Some mass-estimator biases may exceed the -0.5 dex shift tested above. UNCOVER-20466’s demagnified $H\beta$ mass depends on lensing and broad-line interpretation (J. E. Greene et al. 2024); RUBIES-EGS-55604’s dust-corrected $H\beta$ mass has a separately recorded 0.5 dex calibration scale (D. D. Kocevski et al. 2025). For COSMOS3D-13852, scattering could inflate the $H\beta$ mass by 1–2 dex (X. Lin et al. 2026). Applying downward offsets of 1 and 2 dex gives point requirements of 1.115 and 0.916, respectively. These stress tests use the bias range discussed by the source; the actual correction remains uncertain.

J0910_2028_12910 has a 0.01 dex quoted error and a requirement of 1.053, giving $P > 0.999$, yet its separately recorded 0.5 dex calibration scale (C. J. Baccus & X. Xu 2026) allows it below unity in the negative-offset test. ZS7’s quoted 0.4 dex already includes calibration scatter (H. Übler et al. 2024). The probabilities therefore reflect different combinations of measurement and calibration uncertainty across the sample.

5.3 Growth constraints for individual objects

UNCOVER-20466 and GS-20057765 have reference requirements of 1.452 and 1.355, with 16th percentiles above unity. Their point requirements remain above unity for $10^3 M_\odot$ seeds (1.217 and 1.101), but fall to 0.746 and 0.592 for $10^5 M_\odot$ seeds. At $\epsilon = 0.05719$, light seeds require 0.793 and 0.740. For Eddington-limited active phases separated by zero-accretion phases, these imply active fractions of about 79 and 74 per cent of the available time, requiring a sustained fuel supply.

GN-z11’s reference requirement is 1.437. Seeding at $z = 20$ raises it to 1.880, above UNCOVER-20466’s 1.733, reversing their order. A mass measured with a method comparable to the Balmer-line sample would test this timing constraint. GN-z11 meets the primary evidence criterion and is placed in the exploratory sample because it uses a UV mass estimator.

5.4 Limitations

The contributing studies use different selection criteria and detection limits. Inferring population demographics would require a combined completeness model. Virial calibration and geometry errors are

Table 6: Mass-estimator limitations and proposed observations for all 12 primary objects above the reference threshold. Caveats summarize the source records cited in Table 1; proposed observations target the stated limitations; their feasibility has not been assessed.

Object	Mass-estimator caveat	Proposed observation
UNCOVER-20466	Hbeta; mass depends on lens model and BLR interpretation.	Independent lens model and broad-line profile constraints.
GS-20057765	Hbeta; extinction and calibration uncertainties are incompletely quantified.	Deeper line profiles and an extinction constraint.
COSMOS3D-13852	Hbeta; source warns of possible 1–2 dex scattering bias.	Separate scattering wings from virial broadening.
RUBIES-EGS-55604	Hbeta; dust-corrected mass; separate 0.5 dex calibration scale.	Test dust correction and broad-line kinematics.
CEERS-1019	Hbeta; broad component has comparatively low significance.	Confirm the broad component with deeper spectroscopy.
GS-20030333	Hbeta; large quoted errors and uncertain extinction and calibration.	Improve line profile and extinction constraints.
GS-164055	Hbeta; reported interval straddles the reference threshold.	Reduce mass uncertainty with deeper spectroscopy.
GN-38509	Halpha; separate 0.3 dex calibration scale.	Independent estimator or calibration constraint.
J0910_2028_12910	Halpha; 0.01 dex formal error excludes 0.5 dex mass scale.	Measure the mass with an independent calibration.
CEERS-00717	Halpha; local calibration; no numeric systematic supplied.	Cross-check the virial estimator and line decomposition.
ZS7	Hbeta; quoted 0.4 dex already includes calibration scatter.	Spatially separate nuclear kinematics and interacting components.
GN-4685	Hbeta; reported interval straddles the reference threshold.	Improve line profile and extinction constraints.

incompletely reported. The common mass offsets test a uniform shift; biases specific to individual sources, correlated errors and selection-dependent scatter remain unmodelled.

Two duplicate pairs have been reconciled using the source papers; the identities of three groups remain unresolved (Appendix A.2). Excluding all six affected records preserves the threshold results. Reinstating these records requires checking their spectra, masses, images or apertures against the source papers. Counts of unique objects and hosts remain provisional.

Equation 2 describes growth through an average accretion rate, a fixed merger multiplier and constant radiative efficiency. Feedback and individual accretion histories remain unspecified. The probabilities hold these assumptions fixed. Averaging over them would require justified parameter distributions and correlations; identifying seed origins would also require formation models and population information. The age relation neglects radiation, so its extension to $z_{\text{seed}} = 3400$ is an extrapolation.

Independent source checks cover all admitted mass/error fields, 350 central redshifts and 323 available coordinate pairs; 27 rows lack coordinates. Validation of redshift errors, full posteriors and auxiliary observables is incomplete. The catalogue covers the declared sources through 3 September 2026; other JWST discoveries may fall outside this source list. The external A2744-QSO1 test (Section 4.6) leaves this frozen membership intact.

6 Conclusions

For the adopted masses and growth assumptions, we find:

- The conservative 224-object primary sample contains 12 reference point requirements above unity and six with conditional $P \geq 0.95$ under quoted errors.
- UNCOVER-20466, COSMOS3D-13852, and RUBIES-EGS-55604 retain the probability threshold after a -0.5 dex mass shift. Their mass estimates require tests of lensing, scattering and calibration, respectively.
- Seed formation time changes both threshold membership and the most demanding object; lowering the fixed efficiency to 0.05719 removes all point requirements above unity.
- The external A2744-QSO1 dynamical estimate increases its reference requirement from 0.929 to approximately unity, with an error interval spanning the threshold.

Excluding the six records with unresolved identities preserves the threshold counts. Counts of unique objects in the full catalogue remain provisional. The proposed observations test the mass estimates of the most demanding objects under the stated growth assumptions.

Appendix A Source-family inventory

Manuscript sample accounting

338 catalogue object records




Figure 6: Sample accounting for 338 catalogue object records. Mutually exclusive categories comprise 224 primary objects, ten additional exploratory objects, 98 retained without eligible masses and six identity exclusions. The inclusive exploratory numerical sample contains 234 objects.

Table 7: Catalogue size and analysis samples.

Product or subset	Rows or objects
Measurements / provisional objects / hosts	350 / 338 / 337
Growth-eligible measurements / objects	244 / 237
Method-eligible primary measurements / objects	234 / 227
Manuscript primary / exploratory objects	224 / 234
Identity-excluded object records	6
Catalogue-only object records	101

The full catalogue contains 247 broad-line AGN, 49 narrow-line candidates, 30 photometric candidates, ten high-ionization candidates, and two X-ray candidates. Preferred-row evidence labels comprise 214 secure, 41 probable, 82 candidate, and one disputed classification. These full-catalogue counts precede the manuscript exclusions.

Table 8 gives every admitted source family. We count measurements and preferred objects separately. Each preferred object is counted once under its adopted source. Auxiliary coordinates are attributed in the registry to THRILS (T. A. Hutchison et al. 2025), UHZ1 spectroscopy (A. D. Goulding et al. 2023), and the versioned JADES DR3 GOODS-S prism catalogue. The machine-readable companion retains full selection-channel and caveat tags alongside source URLs. “Context” and “Limit” denote values retained outside numerical growth inference.

Table 8: Source-family inventory. Counts are admitted measurements / eligible mass measurements / preferred primary objects after identity exclusion (M/G/P). The columns sum to 350/244/224; G includes alternatives, whereas P counts each selected object once. “Balmer” denotes single-epoch virial masses.

Source	M/G/P	Mass	Selection channel; principal caveat
C. J. Baccus & X. Xu (2026)	49/49/48	Balmer	NIRSpec broad lines; Frozen preprint masses; calibration
Ó. A. Chávez Ortiz et al. (2026)	1/0/0	Context	UV/optical diagnostics; Model-dependent mass
J. Chisholm et al. (2024)	1/0/0	None	High-ionization lines; No canonical mass
K. Davis et al. (2026) / THRILS	6/6/6	Balmer	EELG-selected spectroscopy; Virial calibration
Q. Fei et al. (2026) / GLIMPSE	10/10/8	Balmer	Lensed broad lines; Lensing corrections
J. E. Greene et al. (2024) / UNCOVER	9/9/9	Balmer	Lensed broad lines; Lens model; BLR interpretation
Y. Harikane et al. (2023)	10/10/5	Balmer	NIRSpec broad lines; Local virial calibration
I. Juodžbalis et al. (2026a) / JADES	23/23/21	Balmer	Broad Balmer lines; Extinction; calibration
M. Killi et al. (2024)	1/1/1	Balmer	Broad Halpha; Linear mass-error conversion
D. D. Kocevski et al. (2025) / LRDs	6/6/6	Balmer	Broad Balmer lines; Dust correction; calibration
R. L. Larson et al. (2023) / CEERS-1019	1/1/1	Balmer	Broad Hbeta; Broad-component significance
X. Lin et al. (2024) / ASPIRE	16/16/16	Balmer	Broad Halpha; Virial calibration
X. Lin et al. (2026) / COSMOS-3D	13/13/13	Balmer	NIRCam grism broad lines; Possible scattering bias
J. Lyu et al. (2024) / SMILES	19/0/0	None	MIRI SED selection; SED-dependent AGN identification
R. Maiolino et al. (2024) / GN-z11	1/1/0	UV	UV lines; UV estimator outside primary
S. Mascia et al. (2026)	8/0/0	None	Compact blue broad emitters; No canonical mass
J. Matthee et al. (2024) / EIGER-FRESCO	20/20/19	Balmer	Broad Halpha; Dust; virial calibration
G. Mazzolari et al. (2025)	25/0/0	None	Narrow-line diagnostics; Star-formation overlap
G. C. K. Leung et al. (2026) / MEOW	12/0/0	None	MIRI SED selection; SED-dependent AGN fraction
R. P. Naidu et al. (2026) / MoM-BH*-1	1/0/0	Context	Broad Hbeta / SED; Scattering; stellar alternative
L. Napolitano et al. (2025a) / GHZ9	1/0/0	Context	UV lines / X-ray evidence; Mass assumes Eddington rate
L. Napolitano et al. (2025b) / GHZ4,7	2/0/0	None	Narrow / high-ionization lines; Tentative diagnostics
W. Ren et al. (2025) / ALPINE-CRISTAL	7/7/1	Balmer	Host-selected broad Halpha; Conditional BLR interpretations
J. Scholtz et al. (2025) / JADES	21/0/0	None	Narrow / high-ionization lines; Tentative classifications
D. D. Kocevski et al. (2026) / Skyfire (CEERS)	22/22/22	Balmer	Broad Halpha; Virial calibration
M. Tang et al. (2025)	3/0/0	None	High-ionization lines; Ionizing-source ambiguity
A. J. Taylor et al. (2025) / CEERS-RUBIES	37/37/35	Balmer	Broad Halpha; Slit loss; dust; calibration
H. Treiber et al. (2025) / UNCOVER	2/0/0	None	UV-line diagnostics; Stellar alternatives

Table 8 continued

Source	M/G/P	Mass	Selection channel; principal caveat
H. Übler et al. (2024) / ZS7	1/1/1	Balmer	Offset broad Hbeta; Error includes calibration scatter
Á. Bogdán et al. (2024); F. Zou et al. (2026) / UHZ1	2/0/0	Context	X-rays / SED; Later detection disputed
Z. Zhang et al. (2026) / LRDs	5/0/0	Limit	Narrow lines / SED; Mass upper limits
M.-Y. Zhuang et al. (2026) / NEXUS	15/12/12	Balmer	NIRCam WFSS broad lines; Twelve masses lack errors

A.1 Source-admission checks

The v3 additions include 15 NEXUS measurements from NIRCam/WFSS (M.-Y. Zhuang et al. 2026), 13 COSMOS-3D NIRCam-grism measurements (X. Lin et al. 2026), and GHZ4/GHZ7 (L. Napolitano et al. 2025b). These contribute 30 measurements and 29 object records. NX10835 lies 0.066 arcsec from an existing Mascia identifier at consistent redshift, so the new mass-bearing measurement becomes preferred for the same object.

The Scholtz et al. paper reports 42 objects, but its published table contains 41 rows. Exactly 21 table rows satisfy $z \geq 4$; all are admitted. One is identity-linked to a broad-line object, yielding 20 distinct narrow-line candidates from that source. JADES-NS-GS00099671 is retained with its published host and bolometric observables but without a black-hole mass estimate.

A.2 Preferred measurements and identity exclusions

Measurement, object and host identifiers link each value to its location in the source, publication version and quoted uncertainty. We choose preferred measurements using the source assessments. The CEERS-2782 group uses RUBIES-EGS-50052 because the source prefers its higher-signal-to-noise spectrum over the slit-loss measurement. NX10835 uses the newly available NEXUS mass; UHZ1 uses the later full-dataset evidence assessment and remains outside numerical growth inference. The measurement–object link table records each choice and its reason.

Before the conservative identity exclusions, there are 244 eligible measurements and 237 preferred objects: 236 Balmer-line and one UV single-epoch virial masses. The method-defined primary subset has 234 measurements and 227 objects. The six excluded records link to seven admitted measurements; the three with eligible masses are GDS_1210_9515, GS-8083 and GS-10013704. All three otherwise pass the primary cuts. The other three excluded records lack eligible masses. These records remain excluded until their identities can be resolved. The open groups are Baccus GDS_1210_9515 versus JADES GS-8083, GS-10013704 versus Scholtz 99671, and Scholtz 16745 versus 208643. The first requires target/spectral and preferred-mass reconciliation; the latter two require imaging and aperture checks. Seven of the nine additional candidate objects have conditional broad-line interpretations. All original measurements and catalogue flags remain available.

Appendix B Catalogue navigation score

The point navigation score is

$$S = \min \left[100, 100 \max \left(C \left(\frac{f_2 - 0.3}{1.2} \right), C \left(\frac{s_{0.3} - 4}{2.8} \right) \right) + 8C \left(\frac{z - 6}{4} \right) \right], \quad (4)$$

where $C(x) = \min(1, \max(0, x))$, f_2 is the required f_{Edd} for a $10^2 M_\odot$ seed, and $s_{0.3}$ is the required $\log_{10}(M_{\text{seed}}/M_\odot)$ at $f_{\text{Edd}} = 0.3$. This score combines two growth diagnostics and a redshift term to help browse the catalogue. It has no probabilistic calibration. Ties use decreasing f_2 , decreasing redshift, then stable identifier. The manuscript's required- f_{Edd} table is ordered directly by f_2 .

Appendix C Efficiency prescription for compatibility scenarios

For the illustrative spin-dependent maps, the ideal Kerr thin-disk efficiency is the ISCO binding efficiency (J. M. Bardeen et al. 1972):

$$\epsilon_a = 1 - \sqrt{1 - \frac{2}{3r_{\text{ISCO}}}}, \quad (5)$$

$$r_{\text{ISCO}} = 3 + Z_2 - \text{sgn}(a)\sqrt{(3 - Z_1)(3 + Z_1 + 2Z_2)}, \quad (6)$$

$$Z_1 = 1 + (1 - a^2)^{1/3}[(1 + a)^{1/3} + (1 - a)^{1/3}], \quad Z_2 = \sqrt{3a^2 + Z_1^2}. \quad (7)$$

The dimensionless spin is $a = cJ/(GM_{\text{BH}}^2)$, where J is the signed angular momentum relative to the disk, and r_{ISCO} is in units of GM_{BH}/c^2 . The ideal efficiencies at $a = -1, 0, +1$ are approximately 0.038, 0.057, and 0.423, respectively, neglecting photon capture and stress at the ISCO. Logarithmic luminosity growth at supercritical inflow is motivated by advection and wind models (J. Poutanen et al. 2007). We use a simplified logarithmic relation with coefficient one and omit radial mass loss. For Eddington ratios above unity, the maps adopt

$$f_{\text{Edd}} = 1 + \ln \dot{m}, \quad \dot{m} = \frac{\dot{M}_{\text{in}}}{L_{\text{Edd}}/(\epsilon_a c^2)}, \quad \epsilon_{\text{eff}} = \begin{cases} \epsilon_a, & f_{\text{Edd}} \leq 1, \\ \epsilon_a f_{\text{Edd}} \exp(1 - f_{\text{Edd}}), & f_{\text{Edd}} > 1. \end{cases} \quad (8)$$

The logarithmic luminosity relation applies only for super-Eddington accretion. Each compatibility cell holds both the rate and its effective efficiency constant over the interval. The grid tests constant accretion states with fixed spin and omits relativistic slim-disc dynamics. For time-varying efficiency, growth depends on $\int [(1 - \epsilon(t))/\epsilon(t)] f_{\text{Edd}}(t) dt$, which requires the joint time dependence of efficiency and accretion rate.

Figure 4 gives the full scenario grid. The broad 10^2 – $10^6 M_\odot$ interval is stored under the original code key `pbh_origin_hypothesis`. This key labels the mass interval; a primordial interpretation requires additional assumptions about seed formation. The descriptive fractions use point masses and the inclusive interval criterion in Section 3.3, without error integration. Appendix D.4 explains the conditional interpretation of this interval for pre-existing primordial black holes.

Appendix D Growth-model derivations and implementation

D.1 Accretion and exponential growth

The growth constraints follow from the relation between accretion luminosity, retained mass, and available cosmic time. For electron-scattering opacity in ionized hydrogen, the Eddington luminosity and luminosity-based Eddington ratio are

$$L_{\text{Edd}} = \frac{4\pi GM_{\text{BH}} m_p c}{\sigma_T}, \quad f_{\text{Edd}} = \frac{L_{\text{bol}}}{L_{\text{Edd}}}, \quad t_{\text{Edd}} = \frac{M_{\text{BH}} c^2}{L_{\text{Edd}}} = \frac{c \sigma_T}{4\pi G m_p} \simeq 0.45 \text{ Gyr}. \quad (9)$$

Here G , c , m_p , and σ_T denote the gravitational constant, speed of light, proton mass, and Thomson cross section. With radiative efficiency $0 < \epsilon < 1$ and rest-mass inflow rate $\dot{M}_{\text{in}}$,

$$L_{\text{bol}} = \epsilon \dot{M}_{\text{in}} c^2, \quad \dot{M}_{\text{BH}} = (1 - \epsilon) \dot{M}_{\text{in}} = \frac{1 - \epsilon}{\epsilon} \frac{f_{\text{Edd}}}{t_{\text{Edd}}} M_{\text{BH}}. \quad (10)$$

The inflow rate is defined at the radiating flow; earlier mass loss is omitted. Dividing Eq. 10 by M_{BH} and integrating at fixed ϵ gives

$$\ln \left(\frac{M_{\text{BH}}}{B_{\text{merge}} M_{\text{seed}}} \right) = \frac{1 - \epsilon}{\epsilon} \frac{1}{t_{\text{Edd}}} \int_{t(z_{\text{seed}})}^{t(z_{\text{obs}})} f_{\text{Edd}}(t) dt = \frac{1 - \epsilon}{\epsilon} \frac{\overline{f_{\text{Edd}}} \Delta t}{t_{\text{Edd}}}. \quad (11)$$

Exponentiating yields Eq. 2; solving for the average rate and applying the non-negative floor yields Eq. 3. For a constant positive f_{Edd} , the e-folding time is

$$t_{\text{efold}} = \frac{\epsilon}{1 - \epsilon} \frac{t_{\text{Edd}}}{f_{\text{Edd}}}. \quad (12)$$

It is 0.05 Gyr at $\epsilon = 0.1$ and $f_{\text{Edd}} = 1$.

D.2 Merger multiplier and inverse seed mass

The implementation evaluates Eq. 2 in log mass:

$$\log_{10} \left(\frac{M_{\text{BH}}}{M_{\odot}} \right) = \log_{10} \left(\frac{M_{\text{seed}}}{M_{\odot}} \right) + \log_{10} B_{\text{merge}} + \frac{1 - \epsilon}{\epsilon} \frac{\overline{f_{\text{Edd}}} \Delta t}{t_{\text{Edd}} \ln 10}. \quad (13)$$

We evaluate $B_{\text{merge}} = 1$ (no boost) and 2 (a factor-of-two mass boost). At fixed accretion history,

$$\frac{M_{\text{BH}}(B_{\text{merge}} = 2)}{M_{\text{BH}}(B_{\text{merge}} = 1)} = 2, \quad \Delta \log_{10} M_{\text{BH}} = \log_{10} 2 \simeq 0.3010. \quad (14)$$

Solving the same relation for seed mass gives

$$\log_{10} \left(\frac{M_{\text{seed,req}}}{M_{\odot}} \right) = \log_{10} \left(\frac{M_{\text{BH}}}{M_{\odot}} \right) - \log_{10} B_{\text{merge}} - \frac{1 - \epsilon}{\epsilon} \frac{\overline{f_{\text{Edd}}} \Delta t}{t_{\text{Edd}} \ln 10}. \quad (15)$$

Here $M_{\text{seed,req}}$ is the seed mass required for a specified average rate, efficiency, growth interval and merger multiplier.

D.3 Cosmic time

Cosmic ages use the flat matter-plus-Lambda relation

$$t(z) = \frac{2}{3H_0\sqrt{\Omega_{\Lambda}}} \operatorname{asinh} \left[\sqrt{\frac{\Omega_{\Lambda}}{\Omega_m}} (1+z)^{-3/2} \right], \quad (16)$$

with $H_0 = 67.3 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.315$, and $\Omega_{\Lambda} = 0.685$. The Hubble constant is converted to inverse time; ages and growth intervals are in Gyr. Radiation is neglected, including in the supplementary extrapolation to $z_{\text{seed}} = 3400$.

D.4 Earlier growth onset and primordial seed interpretations

The growth equations take the initial mass and subsequent accretion history as inputs. The seed-formation mechanism is left unspecified. A primordial black hole (PBH) already present at $z = 30$ can therefore be represented by its effective mass at that epoch if it lies within the broad 10^2 – $10^6 M_\odot$ interval and follows the adopted subsequent growth prescription. This mass need not equal its primordial formation mass. This interval covers a specified range of effective masses; inferring a primordial origin requires a separate formation model. In P. Dayal (2024), $z \simeq 3400$ marks matter–radiation equality and the start of the modelled evolution of PBHs that formed earlier.

Moving the start of the same constant-rate, constant-efficiency history from $z = 30$ to 3400 adds $\delta t = t(30) - t(3400) = 0.09990$ Gyr in our matter-plus-Lambda extrapolation. At fixed seed mass and merger multiplier, the resulting shift is

$$\Delta \log_{10} M_{\text{BH}} = \frac{1 - \epsilon}{\epsilon} \frac{\overline{f_{\text{Edd}}} \delta t}{t_{\text{Edd}} \ln 10}. \quad (17)$$

For $\overline{f_{\text{Edd}}} = 1$, this is 0.868 dex (a factor of 7.37) at $\epsilon = 0.1$, or 1.589 dex (a factor of 38.85) at the ideal non-spinning thin-disc efficiency. The tracks shift upward by these amounts (Figure 7). at fixed observed mass and seed mass requiring positive accretion, the ratio of required average rates is $[t(z_{\text{obs}}) - t(30)]/[t(z_{\text{obs}}) - t(3400)]$; the earlier start lowers the requirement by 6.5, 13.1 and 22.9 per cent at $z_{\text{obs}} = 4, 7$ and 10.603, respectively.




Figure 7: Growth from $z_{\text{seed}} = 30$ (a) and 3400 (b), with matched seed masses and the fixed assumptions shown. Earlier onset adds 99.90 Myr and shifts each track upward by 0.868 dex. Here z_{seed} denotes the assumed start of accretion. The $z = 3400$ comparison extrapolates the matter-plus-Lambda cosmology and assumes uninterrupted accretion. Radiation, gas supply, halo assembly and feedback are omitted.

These comparisons quantify the additional growth time. Applying them to PBHs requires an effective mass at the chosen starting epoch and a model for the subsequent gas supply and accretion history.

D.5 Two-state duty cycle

For a two-state history with the same efficiency in both states,

$$\overline{f_{\text{Edd}}} = D f_{\text{burst}} + (1 - D) f_{\text{quiet}}, \quad D_{\text{req}} = \frac{\overline{f_{\text{Eddreq}}} - f_{\text{quiet}}}{f_{\text{burst}} - f_{\text{quiet}}}. \quad (18)$$

Here D is the active-state time fraction, $f_{\text{burst}} > f_{\text{quiet}} \geq 0$, and the diagnostic requires $\overline{f_{\text{Eddreq}}} \geq f_{\text{quiet}}$. We retain values $D_{\text{req}} > 1$ to flag histories that would require more active time than is available. The inferred duty cycle depends on the assumed burst and quiet-state rates.

D.6 Sampling, mass conversions and track selection

Draws use seed 20260808 and a stable identifier-based random stream. For 10,000 draws, the binomial Monte Carlo standard error of a probability is at most 0.005 (approximately 0.0022 at probability 0.95). This numerical error measures the precision from a finite number of draws. The twelve mass-bearing NEXUS objects have no published mass errors: repeated point values produce zero-width stored intervals, while probabilities and uncertainty ranks remain unavailable. Quoted source errors may include systematic contributions. For ZS7, the published 0.4-dex error already includes virial-calibration scatter. We record separately reported method systematics and use them in sensitivity tests where possible. Their distributions and correlations are generally unspecified.

The mass-scale tests add the same fixed offset to every original mass draw. We calculate the results separately for each offset, using the original scatter. The final published Killi mass $8_{-0.4}^{+0.5} \times 10^8 M_{\odot}$ is converted by transforming both linear bounds, giving $\log_{10}(M/M_{\odot}) = 8.90309_{-0.02228}^{+0.02633}$.

For Figure 1, candidate average rates span 0.1–2.0 in steps of 0.1. For each seed mass and fixed efficiency, we retain the two rates with the most primary objects within 0.5 dex of either the $B = 1$ or $B = 2$ prediction, requiring at least five such objects. Ties favour smaller median absolute residuals, then smaller rates. Most objects selected by this 0.5-dex plotting tolerance lie at $z \simeq 4$ –6. Baseline rankings, duty-cycle diagnostics and seed-redshift–mass maps retain $\epsilon = 0.1$. The score in Appendix B is used for catalogue browsing; scientific conclusions use the required-rate ranking.

Appendix E Data and code availability

The catalogue, source registries, code, and manuscript analysis are available at <https://github.com/nnickels/highz-accretion-atlas>. The frozen v3 catalogue baseline is identified by Git commit [a40a0d2](#); version manifests under `releases/` record the dataset artifact hashes. The comparison results and generated tables are in `paper/analysis/`, with explicit external inputs in `paper/review_inputs.json`. An archival DOI is pending.

Five ordered notebooks reproduce the catalogue, science products, figures, and atlas using pinned dependencies. Reproduction checks compare regenerated numerical products, figures, and manuscript table fragments with an independent stored baseline, then verify their internal consistency. Independent cosmic-age quadrature and growth integration supplement the regression checks. The source-family inventory in Appendix A links measurements, methods, selection channels, and major caveats. Full identifiers, source-native observables, alternative measurements, and per-object maps remain available.

These checks verify transcription and numerical reproduction. The validity of the mass estimators and unresolved object identities require observational assessment. All records in the three unresolved groups listed in Section 5.4 are excluded from manuscript inference.

References

- Baccus, C. J., & Xu, X. 2026, *ApJ*, 1006, 165, doi: [10.3847/1538-4357/ae7de7](https://doi.org/10.3847/1538-4357/ae7de7)
- Bardeen, J. M., Press, W. H., & Teukolsky, S. A. 1972, *ApJ*, 178, 347, doi: [10.1086/151796](https://doi.org/10.1086/151796)
- Bogdán, Á., Goulding, A. D., Natarajan, P., et al. 2024, *Nature Astronomy*, 8, 126, doi: [10.1038/s41550-023-02111-9](https://doi.org/10.1038/s41550-023-02111-9)
- Chávez Ortiz, Ó. A., Finkelstein, S. L., Plat, A., et al. 2026, *ApJ*, 1008, 227, doi: [10.3847/1538-4357/ae96a4](https://doi.org/10.3847/1538-4357/ae96a4)
- Chisholm, J., Berg, D. A., Endsley, R., et al. 2024, *MNRAS*, 534, 2633, doi: [10.1093/mnras/stae2199](https://doi.org/10.1093/mnras/stae2199)
- Davis, K., Brooks, M., Trump, J. R., et al. 2026, *ApJ*, submitted. <https://arxiv.org/abs/2602.23310v1>
- Dayal, P. 2024, *A&A*, 690, A182, doi: [10.1051/0004-6361/202451481](https://doi.org/10.1051/0004-6361/202451481)
- Fei, Q., Fujimoto, S., Naidu, R. P., et al. 2026, *ApJ*, 1003, 244, doi: [10.3847/1538-4357/ae6248](https://doi.org/10.3847/1538-4357/ae6248)
- Goulding, A. D., Greene, J. E., Setton, D. J., et al. 2023, *ApJL*, 955, L24, doi: [10.3847/2041-8213/acf7c5](https://doi.org/10.3847/2041-8213/acf7c5)
- Greene, J. E., Labbe, I., Goulding, A. D., et al. 2024, *ApJ*, 964, 39, doi: [10.3847/1538-4357/ad1e5f](https://doi.org/10.3847/1538-4357/ad1e5f)
- Harikane, Y., Zhang, Y., Nakajima, K., et al. 2023, *ApJ*, 959, 39, doi: [10.3847/1538-4357/ad029e](https://doi.org/10.3847/1538-4357/ad029e)
- Hutchison, T. A., Larson, R. L., Arrabal Haro, P., et al. 2025, *arXiv e-prints*. <https://arxiv.org/abs/2512.12509v1>
- Juodžbalis, I., Maiolino, R., Baker, W. M., et al. 2026a, *MNRAS*, 546, stag086, doi: [10.1093/mnras/stag086](https://doi.org/10.1093/mnras/stag086)
- Juodžbalis, I., Marconcini, C., D'Eugenio, F., et al. 2026b, *Nature*, 653, 1017, doi: [10.1038/s41586-026-10579-4](https://doi.org/10.1038/s41586-026-10579-4)
- Killi, M., Watson, D., Brammer, G., et al. 2024, *A&A*, 691, A52, doi: [10.1051/0004-6361/202348857](https://doi.org/10.1051/0004-6361/202348857)
- Kocevski, D. D., Finkelstein, S. L., Barro, G., et al. 2025, *ApJ*, 986, 126, doi: [10.3847/1538-4357/adbc7d](https://doi.org/10.3847/1538-4357/adbc7d)
- Kocevski, D. D., Taylor, A. J., Onoue, M., et al. 2026, *ApJ*, submitted. <https://arxiv.org/abs/2609.00112v1>
- Larson, R. L., Finkelstein, S. L., Kocevski, D. D., et al. 2023, *ApJL*, 953, L29, doi: [10.3847/2041-8213/ace619](https://doi.org/10.3847/2041-8213/ace619)
- Leung, G. C. K., Eilers, A.-C., Endsley, R., et al. 2026, *ApJ*, submitted. <https://arxiv.org/abs/2607.02666v1>
- Lin, X., Fan, X., Wang, F., et al. 2026, *ApJ*, 996, 93, doi: [10.3847/1538-4357/ae1b9b](https://doi.org/10.3847/1538-4357/ae1b9b)
- Lin, X., Wang, F., Fan, X., et al. 2024, *ApJ*, 974, 147, doi: [10.3847/1538-4357/ad6565](https://doi.org/10.3847/1538-4357/ad6565)
- Lyu, J., Alberts, S., Rieke, G. H., et al. 2024, *ApJ*, 966, 229, doi: [10.3847/1538-4357/ad3643](https://doi.org/10.3847/1538-4357/ad3643)
- Maiolino, R., Scholtz, J., Witstok, J., et al. 2024, *Nature*, 627, 59, doi: [10.1038/s41586-024-07052-5](https://doi.org/10.1038/s41586-024-07052-5)
- Mascia, S., Matthee, J., Torralba, A., et al. 2026, *A&A*, submitted. <https://arxiv.org/abs/2608.25021v1>
- Matthee, J., Naidu, R. P., Brammer, G., et al. 2024, *ApJ*, 963, 129, doi: [10.3847/1538-4357/ad2345](https://doi.org/10.3847/1538-4357/ad2345)
- Mazzolari, G., Scholtz, J., Maiolino, R., et al. 2025, *A&A*, 700, A12, doi: [10.1051/0004-6361/202451860](https://doi.org/10.1051/0004-6361/202451860)
- Naidu, R. P., Matthee, J., Katz, H., et al. 2026, *Nature*, 656, 329, doi: [10.1038/s41586-026-10846-4](https://doi.org/10.1038/s41586-026-10846-4)
- Napolitano, L., Castellano, M., Pentericci, L., et al. 2025a, *ApJ*, 989, 75, doi: [10.3847/1538-4357/ade706](https://doi.org/10.3847/1538-4357/ade706)
- Napolitano, L., Castellano, M., Pentericci, L., et al. 2025b, *A&A*, 693, A50, doi: [10.1051/0004-6361/202452090](https://doi.org/10.1051/0004-6361/202452090)
- Poutanen, J., Lipunova, G., Fabrika, S., Butkevich, A. G., & Abolmasov, P. 2007, *MNRAS*, 377, 1187, doi: [10.1111/j.1365-2966.2007.11668.x](https://doi.org/10.1111/j.1365-2966.2007.11668.x)

- Ren, W., Silverman, J. D., Faisst, A. L., et al. 2025, *MNRAS*, 544, 211, doi: [10.1093/mnras/staf1709](https://doi.org/10.1093/mnras/staf1709)
- Scholtz, J., Maiolino, R., D'Eugenio, F., et al. 2025, *A&A*, 697, A175, doi: [10.1051/0004-6361/202348804](https://doi.org/10.1051/0004-6361/202348804)
- Shen, Y. 2013, *Bulletin of the Astronomical Society of India*, 41, 61. <https://arxiv.org/abs/1302.2643v1>
- Tang, M., Stark, D. P., Plat, A., et al. 2025, *ApJ*, 991, 217, doi: [10.3847/1538-4357/adfd57](https://doi.org/10.3847/1538-4357/adfd57)
- Taylor, A. J., Finkelstein, S. L., Kocevski, D. D., et al. 2025, *ApJ*, 986, 165, doi: [10.3847/1538-4357/add15b](https://doi.org/10.3847/1538-4357/add15b)
- Treiber, H., Greene, J. E., Weaver, J. R., et al. 2025, *ApJ*, 984, 93, doi: [10.3847/1538-4357/adc38f](https://doi.org/10.3847/1538-4357/adc38f)
- Übler, H., Maiolino, R., Pérez-González, P. G., et al. 2024, *MNRAS*, 531, 355, doi: [10.1093/mnras/stae943](https://doi.org/10.1093/mnras/stae943)
- Zhang, Z., Jiang, L., Liu, W., Ho, L. C., & Inayoshi, K. 2026, *ApJ*, 998, 170, doi: [10.3847/1538-4357/ae33c4](https://doi.org/10.3847/1538-4357/ae33c4)
- Zhuang, M.-Y., Li, J., Shen, Y., et al. 2026, *ApJ*, 999, 31, doi: [10.3847/1538-4357/ae3612](https://doi.org/10.3847/1538-4357/ae3612)
- Zou, F., Gallo, E., Zuo, Z., et al. 2026, *ApJ*, in press. <https://arxiv.org/abs/2603.24893v2>