Fast Regression of the Tritium Breeding Ratio in Tokamak Fusion Reactors

G Van Goffrier¹ and P Mánek^{1,2}, V Gopakumar³, N Nikolau¹, J Shimwell³, I Waldmann¹

- Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK
- ² Institute of Experimental and Applied Physics, Czech Technical University, Husova 240/5, Prague 110 00, Czech Republic
- ³ UK Atomic Energy Authority, Culham Science Centre, OX14 3DB Abingdon, UK

E-mail: graham.vangoffrier.19@ucl.ac.uk, petr.manek.19@ucl.ac.uk

Abstract. The tritium breeding ratio (TBR) is an essential quantity for the design of modern and next-generation Tokamak nuclear fusion reactors. Representing the ratio between tritium fuel generated in breeding blankets and fuel consumed during reactor runtime, the TBR depends on reactor geometry and material properties in a complex manner. In this work, we explored the training of surrogate models to produce a cheap but high-quality approximation for a Monte Carlo TBR model in use at the UK Atomic Energy Authority. We investigated possibilities for dimensional reduction of its feature space, reviewed 9 families of surrogate models for potential applicability, and performed hyperparameter optimisation. Here we present the performance and scaling properties of these models, the fastest of which, an artificial neural network, demonstrated $R^2 = 0.985$ and a mean prediction time of 0.898 us, representing a relative speedup of $8 \cdot 10^6$ with respect to the expensive MC model. We further present a novel adaptive sampling algorithm, Quality-Adaptive Surrogate Sampling, capable of interfacing with any of the individually studied surrogates. Our preliminary testing on a toy TBR theory has demonstrated the efficacy of this algorithm for accelerating the surrogate modelling process.

Index terms— magnetic moment, solar neutrinos, astrophysics Submitted to: J.

- 1. Introduction
- 2. Data Exploration
- 3. Methodology
- 4. Results
- 5. Conclusion
- 6. Acknowledgements
- 7. References