

Fourier Series

Suppose we have an arbitrary periodic function $f(t)$ that repeats itself every Time period T with a frequency $1/T$. Under certain conditions the function can be expressed as the sum of an infinite number of sin & cosine functions known as a Fourier series. The Fourier equation is defined by:

$$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos\left(\frac{2n\pi t}{T}\right) + b_n \sin\left(\frac{2n\pi t}{T}\right) \right)$$

Where the amplitudes or coefficients of each term are defined as:

$$a_0 = \frac{2}{T} \int_0^T f(t) dt$$

$$a_n = \frac{2}{T} \int_0^T f(t) \cos\left(\frac{2n\pi t}{T}\right) dt$$


$$b_n = \frac{2}{T} \int_0^T f(t) \sin\left(\frac{2n\pi t}{T}\right) dt$$

The coefficients of the Fourier series integrate an area of the function and factorise each area by the Time period T . The first coefficient a_0 calculates the area of the cyclic function that is being converted to trigonometric functions, the coefficients a_n (for an even function of $f(t)$) & b_n (for an odd function of $f(t)$) calculates the amplitude & number of components of $f(t)$ and represents each component with an area of a sin or cos term, each term is linearised by the period to a component value for $f(t)$. The component values of $f(t)$ are represented in the Fourier series as the amplitude of sin & cos terms added together to give the value of $f(t)$ for any t from 0 to T . The sum of components is from $n=1$ to ∞ as each term is normalised by $1/n$ for each n th term.

In figure 1 is an example of a triangular wave function with a period of $T=5$:

$$f(t) = t + 2.5 \quad -2.5 < t < 0$$

$$f(t) = -t + 2.5 \quad 0 < t < 2.5$$



$$a_0 = \frac{2}{5} \left(\int_0^{2.5} (-t + 2.5) dt + \int_{-2.5}^0 (t + 2.5) dt \right)$$

$$a_0 = \frac{2}{5} \left(\left[-\frac{t^2}{2} + 2.5t \right]_0^{2.5} + \left[\frac{t^2}{2} + 2.5t \right]_{-2.5}^0 \right)$$

$$a_0 = \frac{5}{2}$$

$$a_n = \frac{2}{5} \left(\int_0^{2.5} (-t + 2.5) \cos\left(\frac{2\pi nt}{T}\right) dt + \int_{-2.5}^0 (t + 2.5) \cos\left(\frac{2\pi nt}{T}\right) dt \right)$$

$$a_n = \frac{2}{5} \left(\left[\frac{(-t + 2.5)T \sin\left(\frac{2\pi nt}{T}\right)}{2\pi n} - \left(\frac{T}{2\pi n}\right)^2 \cos\left(\frac{2\pi nt}{T}\right) \right]_0^{2.5} + \left[\frac{(t + 2.5)T \sin\left(\frac{2\pi nt}{T}\right)}{2\pi n} + \left(\frac{T}{2\pi n}\right)^2 \cos\left(\frac{2\pi nt}{T}\right) \right]_{-2.5}^0 \right)$$

$$a_n = \frac{8}{5} \left(\frac{T}{2\pi n} \right)^2$$

For the case of half range cosine expansion:

$$a_n = \frac{8}{5} \left(\frac{T}{(2n-1)\pi} \right)^2$$

$$f(t) = \frac{5}{4} + \sum_{n=1}^{\infty} \left(\frac{10}{\pi^2 (2n-1)^2} \cos\left(\frac{(2n-1)\pi t}{5}\right) \right)$$