

vv214: Orthogonality. Gram-Schmidt Orthogonalization. Least-squares solutions.

Dr.Olga Danilkina

UM-SJTU Joint Institute



June 19, 2020

1. Norms. Normed linear spaces. Convergence.
2. Inner product. Inner product spaces.
3. Natural norm.
4. Banach and Hilbert spaces.
5. The Cauchy-Schwarz inequality.
6. Orthogonal and orthonormal elements of inner product spaces.
7. Orthogonal complements and direct sums.
8. Formal definition of a Fourier series.
9. Correlation.
10. Construction of orthonormal bases. Gram-Schmidt process.
 QR - factorization.

Norms

Definition: Let X be a linear space over a scalar field \mathbb{K} . A real-valued function $\|\cdot\|: X \rightarrow \mathbb{R}$ defined on X is called a **norm** provided

Norms

Definition: Let X be a linear space over a scalar field \mathbb{K} . A real-valued function $\|\cdot\|: X \rightarrow \mathbb{R}$ defined on X is called a **norm** provided

1. $\|x\| \geq 0 \quad \forall x \in X$ and $\|x\| = 0$ iff $x = 0$

Norms

Definition: Let X be a linear space over a scalar field \mathbb{K} . A real-valued function $\|\cdot\|: X \rightarrow \mathbb{R}$ defined on X is called a **norm** provided

1. $\|x\| \geq 0 \quad \forall x \in X$ and $\|x\| = 0$ iff $x = 0$
2. $\|\alpha x\| = |\alpha| \|x\| \quad \forall x \in X \forall \alpha \in \mathbb{K}$

Norms

Definition: Let X be a linear space over a scalar field \mathbb{K} . A real-valued function $\|\cdot\|: X \rightarrow \mathbb{R}$ defined on X is called a **norm** provided

1. $\|x\| \geq 0 \quad \forall x \in X$ and $\|x\| = 0$ iff $x = 0$
2. $\|\alpha x\| = |\alpha| \|x\| \quad \forall x \in X \forall \alpha \in \mathbb{K}$
3. $\|x + y\| \leq \|x\| + \|y\| \quad \forall x, y \in X$

Norms

Definition: Let X be a linear space over a scalar field \mathbb{K} . A real-valued function $\|\cdot\|: X \rightarrow \mathbb{R}$ defined on X is called a **norm** provided

1. $\|x\| \geq 0 \quad \forall x \in X$ and $\|x\| = 0$ iff $x = 0$
2. $\|\alpha x\| = |\alpha| \|x\| \quad \forall x \in X \forall \alpha \in \mathbb{K}$
3. $\|x + y\| \leq \|x\| + \|y\| \quad \forall x, y \in X$

Definition: A linear space with a norm $(X, \|\cdot\|)$ is called a **normed linear space**.

Convergence in a normed linear space is defined as convergence w.r.t norm:

Norms

Definition: Let X be a linear space over a scalar field \mathbb{K} . A real-valued function $\|\cdot\|: X \rightarrow \mathbb{R}$ defined on X is called a **norm** provided

1. $\|x\| \geq 0 \quad \forall x \in X$ and $\|x\| = 0$ iff $x = 0$
2. $\|\alpha x\| = |\alpha| \|x\| \quad \forall x \in X \forall \alpha \in \mathbb{K}$
3. $\|x + y\| \leq \|x\| + \|y\| \quad \forall x, y \in X$

Definition: A linear space with a norm $(X, \|\cdot\|)$ is called a **normed linear space**.

Convergence in a normed linear space is defined as convergence w.r.t norm:

$$\{x_n\} \rightarrow x \text{ as } n \rightarrow \infty \Rightarrow \|x_n - x\| \rightarrow 0 \text{ as } n \rightarrow \infty$$

Norms: Examples

1. \mathbb{R} : $||x|| = |x|$

Norms: Examples

1. \mathbb{R} : $||x|| = |x|$

2. \mathbb{R}^n : $||\bar{x}||_{\infty} = \max_{i=1..n} |x_i|$, $||\bar{x}||_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}$, $p \geq 1$

Norms: Examples

1. \mathbb{R} : $\|x\| = |x|$

2. \mathbb{R}^n : $\|\bar{x}\|_\infty = \max_{i=1..n} |x_i|$, $\|\bar{x}\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}$, $p \geq 1$

3. $C[a, b]$: $\|f(t)\| = \max_{t \in [a, b]} |f(t)|$, $\|f(t)\| = \int_a^b |f(t)| dt$

Norms: Examples

1. \mathbb{R} : $\|x\| = |x|$

2. \mathbb{R}^n : $\|\bar{x}\|_\infty = \max_{i=1..n} |x_i|$, $\|\bar{x}\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}$, $p \geq 1$

3. $C[a, b]$: $\|f(t)\| = \max_{t \in [a, b]} |f(t)|$, $\|f(t)\| = \int_a^b |f(t)| dt$

4. l^∞ : $\|\bar{x}\| = \sup_{i=1..\infty} |x_i|$

Norms: Examples

1. \mathbb{R} : $\|x\| = |x|$

2. \mathbb{R}^n : $\|\bar{x}\|_\infty = \max_{i=1..n} |x_i|$, $\|\bar{x}\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}$, $p \geq 1$

3. $C[a, b]$: $\|f(t)\| = \max_{t \in [a, b]} |f(t)|$, $\|f(t)\| = \int_a^b |f(t)| dt$

4. l^∞ : $\|\bar{x}\| = \sup_{i=1..\infty} |x_i|$

5. l^p : $\|\bar{x}\|_p = \left(\sum_{i=1}^{\infty} |x_i|^p \right)^{1/p}$, $p \geq 1$

Inner Product

Definition: Let X be a linear space over a scalar field \mathbb{K} . A complex-valued function of two variables $(\cdot, \cdot): X \times X \rightarrow \mathbb{C}$ is called an **inner product** (dot product) provided

Inner Product

Definition: Let X be a linear space over a scalar field \mathbb{K} . A complex-valued function of two variables $(\cdot, \cdot): X \times X \rightarrow \mathbb{C}$ is called an **inner product** (dot product) provided

1. $(x, x) \geq 0 \quad \forall x \in X$ and $(x, x) = 0$ iff $x = 0$

Inner Product

Definition: Let X be a linear space over a scalar field \mathbb{K} . A complex-valued function of two variables $(\cdot, \cdot): X \times X \rightarrow \mathbb{C}$ is called an **inner product** (dot product) provided

1. $(x, x) \geq 0 \quad \forall x \in X$ and $(x, x) = 0$ iff $x = 0$
2. $(\alpha x + \beta y, z) = \alpha(x, z) + \beta(y, z) \quad \forall x, y, z \in X \forall \alpha, \beta \in \mathbb{K}$

Inner Product

Definition: Let X be a linear space over a scalar field \mathbb{K} . A complex-valued function of two variables $(\cdot, \cdot): X \times X \rightarrow \mathbb{C}$ is called an **inner product** (dot product) provided

1. $(x, x) \geq 0 \quad \forall x \in X$ and $(x, x) = 0$ iff $x = 0$
2. $(\alpha x + \beta y, z) = \alpha(x, z) + \beta(y, z) \quad \forall x, y, z \in X \forall \alpha, \beta \in \mathbb{K}$
3. $(x, y) = \overline{(y, x)} \quad \forall x, y \in X$

Inner Product

Definition: Let X be a linear space over a scalar field \mathbb{K} . A complex-valued function of two variables $(\cdot, \cdot): X \times X \rightarrow \mathbb{C}$ is called an **inner product** (dot product) provided

1. $(x, x) \geq 0 \quad \forall x \in X$ and $(x, x) = 0$ iff $x = 0$
2. $(\alpha x + \beta y, z) = \alpha(x, z) + \beta(y, z) \quad \forall x, y, z \in X \forall \alpha, \beta \in \mathbb{K}$
3. $(x, y) = \overline{(y, x)} \quad \forall x, y \in X$

Examples:

Inner Product

Definition: Let X be a linear space over a scalar field \mathbb{K} . A complex-valued function of two variables $(\cdot, \cdot): X \times X \rightarrow \mathbb{C}$ is called an **inner product** (dot product) provided

1. $(x, x) \geq 0 \quad \forall x \in X$ and $(x, x) = 0$ iff $x = 0$
2. $(\alpha x + \beta y, z) = \alpha(x, z) + \beta(y, z) \quad \forall x, y, z \in X \forall \alpha, \beta \in \mathbb{K}$
3. $(x, y) = \overline{(y, x)} \quad \forall x, y \in X$

Examples:

$$1. \mathbb{R}^n: (\bar{x}, \bar{y}) = \sum_{i=1}^n x_i y_i, \quad \mathbb{C}^n: (\bar{x}, \bar{y}) = \sum_{i=1}^n x_i \bar{y}_i$$

Inner Product

Definition: Let X be a linear space over a scalar field \mathbb{K} . A complex-valued function of two variables $(\cdot, \cdot): X \times X \rightarrow \mathbb{C}$ is called an **inner product** (dot product) provided

1. $(x, x) \geq 0 \quad \forall x \in X$ and $(x, x) = 0$ iff $x = 0$
2. $(\alpha x + \beta y, z) = \alpha(x, z) + \beta(y, z) \quad \forall x, y, z \in X \forall \alpha, \beta \in \mathbb{K}$
3. $(x, y) = \overline{(y, x)} \quad \forall x, y \in X$

Examples:

1. $\mathbb{R}^n: (\bar{x}, \bar{y}) = \sum_{i=1}^n x_i y_i, \quad \mathbb{C}^n: (\bar{x}, \bar{y}) = \sum_{i=1}^n x_i \bar{y}_i$
2. $C[a, b]: (f(t), g(t)) = \int_a^b f(t)g(t) dt$

Inner Product

Definition: Let X be a linear space over a scalar field \mathbb{K} . A complex-valued function of two variables $(\cdot, \cdot): X \times X \rightarrow \mathbb{C}$ is called an **inner product** (dot product) provided

1. $(x, x) \geq 0 \quad \forall x \in X$ and $(x, x) = 0$ iff $x = 0$
2. $(\alpha x + \beta y, z) = \alpha(x, z) + \beta(y, z) \quad \forall x, y, z \in X \forall \alpha, \beta \in \mathbb{K}$
3. $(x, y) = \overline{(y, x)} \quad \forall x, y \in X$

Examples:

1. $\mathbb{R}^n: (\bar{x}, \bar{y}) = \sum_{i=1}^n x_i y_i, \quad \mathbb{C}^n: (\bar{x}, \bar{y}) = \sum_{i=1}^n x_i \bar{y}_i$
2. $C[a, b]: (f(t), g(t)) = \int_a^b f(t)g(t) dt$
3. $l^2: (\bar{x}, \bar{y}) = \sum_{i=1}^{\infty} x_i \bar{y}_i$

Inner Product

Definition: Let X be a linear space over a scalar field \mathbb{K} . A complex-valued function of two variables $(\cdot, \cdot): X \times X \rightarrow \mathbb{C}$ is called an **inner product** (dot product) provided

1. $(x, x) \geq 0 \quad \forall x \in X$ and $(x, x) = 0$ iff $x = 0$
2. $(\alpha x + \beta y, z) = \alpha(x, z) + \beta(y, z) \quad \forall x, y, z \in X \forall \alpha, \beta \in \mathbb{K}$
3. $(x, y) = \overline{(y, x)} \quad \forall x, y \in X$

Examples:

1. $\mathbb{R}^n: (\bar{x}, \bar{y}) = \sum_{i=1}^n x_i y_i, \quad \mathbb{C}^n: (\bar{x}, \bar{y}) = \sum_{i=1}^n x_i \bar{y}_i$
2. $C[a, b]: (f(t), g(t)) = \int_a^b f(t)g(t) dt$
3. $l^2: (\bar{x}, \bar{y}) = \sum_{i=1}^{\infty} x_i \bar{y}_i$
4. $M_{n \times n}: (A, B) = \text{tr}(\bar{A}^T B)$ **Frobenius inner product**

Inner Product

Definition: Let X be a linear space over a scalar field \mathbb{K} . A complex-valued function of two variables $(\cdot, \cdot): X \times X \rightarrow \mathbb{C}$ is called an **inner product** (dot product) provided

1. $(x, x) \geq 0 \quad \forall x \in X$ and $(x, x) = 0$ iff $x = 0$
2. $(\alpha x + \beta y, z) = \alpha(x, z) + \beta(y, z) \quad \forall x, y, z \in X \quad \forall \alpha, \beta \in \mathbb{K}$
3. $(x, y) = \overline{(y, x)} \quad \forall x, y \in X$

Examples:

1. $\mathbb{R}^n: (\bar{x}, \bar{y}) = \sum_{i=1}^n x_i y_i, \quad \mathbb{C}^n: (\bar{x}, \bar{y}) = \sum_{i=1}^n x_i \bar{y}_i$
2. $C[a, b]: (f(t), g(t)) = \int_a^b f(t)g(t) dt$
3. $l^2: (\bar{x}, \bar{y}) = \sum_{i=1}^{\infty} x_i \bar{y}_i$
4. $M_{n \times n}: (A, B) = \text{tr}(\bar{A}^T B)$ **Frobenius inner product**
5. $\mathbb{R}^2: (\bar{x}, \bar{y}) = x_1 y_1 - x_2 y_1 - x_1 y_2 + 4x_2 y_2$

The Cauchy-Schwarz Inequality

Lemma (the Cauchy-Schwarz inequality): Let X be a linear space with an inner product.

$$|(x, y)| \leq \sqrt{(x, x)} \cdot \sqrt{(y, y)}$$

The Cauchy-Schwarz Inequality

Lemma (the Cauchy-Schwarz inequality): Let X be a linear space with an inner product.

$$|(x, y)| \leq \sqrt{(x, x)} \cdot \sqrt{(y, y)}$$

Lemma (triangle inequality):

$$\|x + y\| \leq \|x\| + \|y\|$$

Natural Norm

Definition: If X is a linear space with a norm $\|\cdot\|$, i.e. a normed linear space, then the norm is a **natural norm induced by the inner product** if

$$\|x\| = \sqrt{(x, x)} \quad \forall x \in X$$

Natural Norm

Definition: If X is a linear space with a norm $\|\cdot\|$, i.e. a normed linear space, then the norm is a **natural norm induced by the inner product** if

$$\|x\| = \sqrt{(x, x)} \quad \forall x \in X$$

Examples: In \mathbb{R}^n , $\|x\| = \left(\sum_{i=1}^n |x_i|^2 \right)^{1/2}$ is the natural norm.

In $C[a, b]$, $\|f\| = \left(\int_a^b f^2(t) dt \right)^{1/2}$ is the natural norm.

Natural Norm

Definition: If X is a linear space with a norm $\|\cdot\|$, i.e. a normed linear space, then the norm is a **natural norm induced by the inner product** if

$$\|x\| = \sqrt{(x, x)} \quad \forall x \in X$$

Examples: In \mathbb{R}^n , $\|x\| = \left(\sum_{i=1}^n |x_i|^2 \right)^{1/2}$ is the natural norm.

In $C[a, b]$, $\|f\| = \left(\int_a^b f^2(t) dt \right)^{1/2}$ is the natural norm.

Remark: To verify that a norm $\|\cdot\|$ is a natural norm, check whether the **Parallelogram Equality**

$$\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2)$$

holds.

In every parallelogram, the sum of the squares of the lengths of the diagonals equals the sum of the squares of the lengths of the four sides.

Natural Norm

Important Definitions

Definition: A sequence $\{x_n\} \subset X$ is said to be a **Cauchy sequence** if

$$\|x_n - x_m\| \rightarrow 0 \text{ as } n, m \rightarrow \infty$$

Important Definitions

Definition: A sequence $\{x_n\} \subset X$ is said to be a **Cauchy sequence** if

$$\|x_n - x_m\| \rightarrow 0 \text{ as } n, m \rightarrow \infty$$

Definition: A normed linear space X is said to be **complete** if ANY Cauchy sequence converges w.r.t norm in X .

Important Definitions

Definition: A sequence $\{x_n\} \subset X$ is said to be a **Cauchy sequence** if

$$\|x_n - x_m\| \rightarrow 0 \text{ as } n, m \rightarrow \infty$$

Definition: A normed linear space X is said to be **complete** if ANY Cauchy sequence converges w.r.t norm in X .

Definition: A complete normed linear space is called a **Banach space**.

A complete inner product space is called a **Hilbert space**.

Orthonormality

Definition: Let X be a linear space. Two elements $x, y \in X$ are said to be **orthogonal** (perpendicular) if $(x, y) = 0$.

Orthonormality

Definition: Let X be a linear space. Two elements $x, y \in X$ are said to be **orthogonal** (perpendicular) if $(x, y) = 0$.

Definition: If elements x, y are orthogonal and $\|x\| = \|y\| = 1$, then x and y are called **orthonormal**.

Orthonormality

Definition: Let X be a linear space. Two elements $x, y \in X$ are said to be **orthogonal** (perpendicular) if $(x, y) = 0$.

Definition: If elements x, y are orthogonal and $\|x\| = \|y\| = 1$, then x and y are called **orthonormal**.

Remark: For any $x \in X, x \neq 0$, the element $y = \frac{x}{\|x\|}$ has the norm $\|y\| = 1$. Any element $x \in X$ with $\|x\| = 1$ is a **unit** element.

Examples of orthonormal elements:

Orthonormality

Definition: Let X be a linear space. Two elements $x, y \in X$ are said to be **orthogonal** (perpendicular) if $(x, y) = 0$.

Definition: If elements x, y are orthogonal and $\|x\| = \|y\| = 1$, then x and y are called **orthonormal**.

Remark: For any $x \in X, x \neq 0$, the element $y = \frac{x}{\|x\|}$ has the norm $\|y\| = 1$. Any element $x \in X$ with $\|x\| = 1$ is a **unit** element.

Examples of orthonormal elements:

1. \mathbb{R}^n : $\bar{e}_1, \bar{e}_2, \dots, \bar{e}_n$

Orthonormality

Definition: Let X be a linear space. Two elements $x, y \in X$ are said to be **orthogonal** (perpendicular) if $(x, y) = 0$.

Definition: If elements x, y are orthogonal and $\|x\| = \|y\| = 1$, then x and y are called **orthonormal**.

Remark: For any $x \in X, x \neq 0$, the element $y = \frac{x}{\|x\|}$ has the norm $\|y\| = 1$. Any element $x \in X$ with $\|x\| = 1$ is a **unit** element.

Examples of orthonormal elements:

1. \mathbb{R}^n : $\bar{e}_1, \bar{e}_2, \dots, \bar{e}_n$
2. \mathbb{R}^2 : $\begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}, \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix}$

Orthonormality

Definition: Let X be a linear space. Two elements $x, y \in X$ are said to be **orthogonal** (perpendicular) if $(x, y) = 0$.

Definition: If elements x, y are orthogonal and $\|x\| = \|y\| = 1$, then x and y are called **orthonormal**.

Remark: For any $x \in X, x \neq 0$, the element $y = \frac{x}{\|x\|}$ has the norm $\|y\| = 1$. Any element $x \in X$ with $\|x\| = 1$ is a **unit** element.

Examples of orthonormal elements:

1. \mathbb{R}^n : $\bar{e}_1, \bar{e}_2, \dots, \bar{e}_n$
2. \mathbb{R}^2 : $\begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}, \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix}$
3. $C[-\pi, \pi]$: $\frac{1}{\sqrt{2\pi}}, \frac{1}{\sqrt{\pi}} \cos t, \frac{1}{\sqrt{\pi}} \sin t, \frac{1}{\sqrt{\pi}} \cos 2t, \frac{1}{\sqrt{\pi}} \sin 2t, \dots$

Orthonormality

Definition: Let X be a linear space. Two elements $x, y \in X$ are said to be **orthogonal** (perpendicular) if $(x, y) = 0$.

Definition: If elements x, y are orthogonal and $\|x\| = \|y\| = 1$, then x and y are called **orthonormal**.

Remark: For any $x \in X, x \neq 0$, the element $y = \frac{x}{\|x\|}$ has the norm $\|y\| = 1$. Any element $x \in X$ with $\|x\| = 1$ is a **unit** element.

Examples of orthonormal elements:

1. \mathbb{R}^n : $\bar{e}_1, \bar{e}_2, \dots, \bar{e}_n$
2. \mathbb{R}^2 : $\begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}, \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix}$
3. $C[-\pi, \pi]$: $\frac{1}{\sqrt{2\pi}}, \frac{1}{\sqrt{\pi}} \cos t, \frac{1}{\sqrt{\pi}} \sin t, \frac{1}{\sqrt{\pi}} \cos 2t, \frac{1}{\sqrt{\pi}} \sin 2t, \dots$
4. \mathbb{R}^4 : $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}), (\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}), (\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}),$
 $(-\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

Orthonormal Basis

Lemma (Pythagorean Theorem): If $x, y \in X$ are orthogonal then

$$\|x + y\|^2 = \|x\|^2 + \|y\|^2$$

Orthonormal Basis

Lemma (Pythagorean Theorem): If $x, y \in X$ are orthogonal then

$$\|x + y\|^2 = \|x\|^2 + \|y\|^2$$

Lemma (orthogonal decomposition): Let $x, y \in X$, $y \neq 0$.

$$x = cy + w,$$

$$\text{where } (w, y) = 0 \text{ and } c = \frac{(x, y)}{\|y\|^2} \Rightarrow w = x - \frac{(x, y)}{\|y\|^2} y$$

Orthonormal Basis

Lemma (Pythagorean Theorem): If $x, y \in X$ are orthogonal then

$$||x + y||^2 = ||x||^2 + ||y||^2$$

Lemma (orthogonal decomposition): Let $x, y \in X$, $y \neq 0$.

$$x = cy + w,$$

$$\text{where } (w, y) = 0 \text{ and } c = \frac{(x, y)}{||y||^2} \Rightarrow w = x - \frac{(x, y)}{||y||^2}y$$

Remark: Prove the Cauchy-Schwarz inequality using an orthogonal decomposition.

Theorem: Orthonormal elements are linearly independent.

Orthonormal Basis

Lemma (Pythagorean Theorem): If $x, y \in X$ are orthogonal then

$$||x + y||^2 = ||x||^2 + ||y||^2$$

Lemma (orthogonal decomposition): Let $x, y \in X, y \neq 0$.

$$x = cy + w,$$

$$\text{where } (w, y) = 0 \text{ and } c = \frac{(x, y)}{||y||^2} \Rightarrow w = x - \frac{(x, y)}{||y||^2} y$$

Remark: Prove the Cauchy-Schwarz inequality using an orthogonal decomposition.

Theorem: Orthonormal elements are linearly independent.

Remark: Any n orthonormal vectors $\bar{u}_1, \dots, \bar{u}_n \in \mathbb{R}^n$ form a basis of \mathbb{R}^n .

Orthonormal Basis

Definition: Let $V \subset X$ be a linear subspace. The set

$$V^\perp = \{x \in X : (x, v) = 0 \quad \forall v \in V\}$$

is called the **orthogonal complement** of V .

Is V^\perp a linear space as well?

Orthonormal Basis

Definition: Let $V \subset X$ be a linear subspace. The set

$$V^\perp = \{x \in X : (x, v) = 0 \quad \forall v \in V\}$$

is called the **orthogonal complement** of V .

Is V^\perp a linear space as well? **Yes.**

Orthonormal Basis

Definition: Let $V \subset X$ be a linear subspace. The set

$$V^\perp = \{x \in X : (x, v) = 0 \quad \forall v \in V\}$$

is called the **orthogonal complement** of V .

Is V^\perp a linear space as well? **Yes.**

Theorem: If V is a linear subspace of X , $\dim V = n$, then

$$X = V \oplus V^\perp$$

Orthonormal Basis

Definition: Let $V \subset X$ be a linear subspace. The set

$$V^\perp = \{x \in X : (x, v) = 0 \quad \forall v \in V\}$$

is called the **orthogonal complement** of V .

Is V^\perp a linear space as well? **Yes.**

Theorem: If V is a linear subspace of X , $\dim V = n$, then

$$X = V \oplus V^\perp$$

Remark: If V is a linear subspace of \mathbb{R}^n with an orthonormal basis $\bar{u}_1, \dots, \bar{u}_m$, then

$$\text{proj}_V \bar{x} = (\bar{x}, \bar{u}_1)\bar{u}_1 + \dots + (\bar{x}, \bar{u}_m)\bar{u}_m$$

Orthonormal Basis

Definition: Let $V \subset X$ be a linear subspace. The set

$$V^\perp = \{x \in X : (x, v) = 0 \quad \forall v \in V\}$$

is called the **orthogonal complement** of V .

Is V^\perp a linear space as well? **Yes.**

Theorem: If V is a linear subspace of X , $\dim V = n$, then

$$X = V \oplus V^\perp$$

Remark: If V is a linear subspace of \mathbb{R}^n with an orthonormal basis $\bar{u}_1, \dots, \bar{u}_m$, then

$$\text{proj}_V \bar{x} = (\bar{x}, \bar{u}_1)\bar{u}_1 + \dots + (\bar{x}, \bar{u}_m)\bar{u}_m$$

Remarks:

1. $V \cap V^\perp = \{0\}$
2. $V = (V^\perp)^\perp$
3. $n = \dim X = \dim V^\perp + \dim V$

Orthonormal Basis

How to convert an arbitrary basis $\{\bar{v}_1, \dots, \bar{v}_m\}$ of a linear subspace $V \subset \mathbb{R}^n$ into an orthonormal one, say $\{\bar{u}_1, \dots, \bar{u}_m\}$?

Case1: $\dim V = 1 \Rightarrow V$ is a line $\Rightarrow V = \text{span } \bar{v}_1 \Rightarrow \bar{u}_1 = \frac{\bar{v}_1}{\|\bar{v}_1\|}$

Orthonormal Basis

How to convert an arbitrary basis $\{\bar{v}_1, \dots, \bar{v}_m\}$ of a linear subspace $V \subset \mathbb{R}^n$ into an orthonormal one, say $\{\bar{u}_1, \dots, \bar{u}_m\}$?

Case 1: $\dim V = 1 \Rightarrow V$ is a line $\Rightarrow V = \text{span } \bar{v}_1 \Rightarrow \bar{u}_1 = \frac{\bar{v}_1}{\|\bar{v}_1\|}$

Case 2: $\dim V = 2 \Rightarrow V$ is a plane $\Rightarrow V = \text{span}(\bar{v}_1, \bar{v}_2)$

$$\bar{u}_1 = \frac{\bar{v}_1}{\|\bar{v}_1\|} \quad \text{AND} \quad \bar{u}_2 \perp \bar{u}_1$$

$$\text{Let } L = \text{span } \bar{u}_1 \Rightarrow \bar{u}_2 \perp \text{proj}_L \bar{v}_2$$

$$\bar{v}_2 = \underbrace{\text{proj}_L \bar{v}_2}_{\perp L} + \underbrace{\bar{v}}_{\perp L} \Rightarrow \bar{u}_2 \|\bar{v} = \bar{v}_2 - \underbrace{\text{proj}_L \bar{v}_2}_{=(\bar{v}_2, \bar{u}_1)\bar{u}_1}$$

$$\text{Denote } \bar{v} = \bar{u}'_2. \text{ Then } \bar{u}_2 = \frac{\bar{u}'_2}{\|\bar{u}'_2\|}$$

Orthonormal Basis

$$V = \text{span} \left(\underbrace{\begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}}_{\tilde{v}_1}, \underbrace{\begin{pmatrix} 1 \\ 9 \\ 9 \\ 1 \end{pmatrix}}_{\tilde{v}_2} \right)$$

Orthonormal Basis

$$V = \text{span} \left(\underbrace{\begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}}_{\tilde{v}_1}, \underbrace{\begin{pmatrix} 1 \\ 9 \\ 9 \\ 1 \end{pmatrix}}_{\tilde{v}_2} \right)$$

$$\bar{u}_1 = \begin{pmatrix} 1/2 \\ 1/2 \\ 1/2 \\ 1/2 \end{pmatrix},$$

Orthonormal Basis

$$V = \text{span} \left(\underbrace{\begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}}_{\tilde{v}_1}, \underbrace{\begin{pmatrix} 1 \\ 9 \\ 9 \\ 1 \end{pmatrix}}_{\tilde{v}_2} \right)$$

$$\bar{u}_1 = \begin{pmatrix} 1/2 \\ 1/2 \\ 1/2 \\ 1/2 \end{pmatrix}, \quad \bar{u}_2 = \begin{pmatrix} -1/2 \\ 1/2 \\ 1/2 \\ -1/2 \end{pmatrix}$$

$$\bar{u}'_2 = \begin{pmatrix} 1 \\ 9 \\ 9 \\ 1 \end{pmatrix} - (1/2 \cdot 1 + 1/2 \cdot 9 + 1/2 \cdot 9 + 1/2 \cdot 1) \begin{pmatrix} 1/2 \\ 1/2 \\ 1/2 \\ 1/2 \end{pmatrix}$$

Orthonormal Basis

Case 3: $\dim V = 3 \Rightarrow V = \text{span}(\bar{v}_1, \bar{v}_2, \bar{v}_3)$

$$\bar{u}_1 = \frac{\bar{v}_1}{\|\bar{v}_1\|}, \quad \bar{u}_2 = \frac{\bar{u}'_2}{\|\bar{u}'_2\|}, \quad \bar{u}'_2 = \bar{v}_2 - (\bar{v}_2, \bar{u}_1)\bar{u}_1$$

Denote $E = \text{span}(\bar{u}_1, \bar{u}_2)$ and find $\text{proj}_E \bar{v}_3$

Orthonormal Basis

Case 3: $\dim V = 3 \Rightarrow V = \text{span}(\bar{v}_1, \bar{v}_2, \bar{v}_3)$

$$\bar{u}_1 = \frac{\bar{v}_1}{\|\bar{v}_1\|}, \quad \bar{u}_2 = \frac{\bar{u}'_2}{\|\bar{u}'_2\|}, \quad \bar{u}'_2 = \bar{v}_2 - (\bar{v}_2, \bar{u}_1)\bar{u}_1$$

Denote $E = \text{span}(\bar{u}_1, \bar{u}_2)$ and find $\text{proj}_E \bar{v}_3 = \bar{v}_3 - \underbrace{\bar{v}}_{\perp E}$

$$\bar{u}'_3 = \bar{v} = \bar{v}_3 - \text{proj}_E \bar{v}_3 = \bar{v}_3 - (\bar{v}_3, \bar{u}_1)\bar{u}_1 - (\bar{v}_3, \bar{u}_2)\bar{u}_2$$

Gram-Schmidt Theorem

Theorem: Let $\{\bar{v}_1, \dots, \bar{v}_m\}$ be a basis of a linear subspace $V \subset \mathbb{R}^n$. Since

$$\bar{v}_j = \bar{v}_j^{\parallel} + \bar{v}_j^{\perp} \quad \forall j = 2, 3, \dots$$

$$\bar{v}_j^{\parallel} \in \text{span}(\bar{v}_1, \dots, \bar{v}_{j-1}) \text{ and } \bar{v}_j^{\perp} \perp \text{span}(\bar{v}_1, \dots, \bar{v}_{j-1}),$$

so

$$\bar{u}_1 = \frac{\bar{v}_1}{\|\bar{v}_1\|}, \bar{u}_2 = \frac{\bar{v}_2^{\perp}}{\|\bar{v}_2^{\perp}\|}, \dots, \bar{u}_m = \frac{\bar{v}_m^{\perp}}{\|\bar{v}_m^{\perp}\|}$$

is the orthonormal basis of V .

$$\forall j > 2 \quad \bar{v}_j^{\perp} = \bar{v}_j - (\bar{v}_j, \bar{u}_1)\bar{u}_1 - \dots - (\bar{v}_j, \bar{u}_{j-1})\bar{u}_{j-1}$$

Gram-Schmidt Theorem

Gram-Schmidt Theorem

Gram-Schmidt Theorem

Gram-Schmidt Theorem: Remarks

Remark 1:

Using the Gram-Schmidt process, we change the basis

$\mathfrak{B}_1 = \{\bar{v}_1, \dots, \bar{v}_m\}$ to the basis $\mathfrak{B}_2 = \{\bar{u}_1, \dots, \bar{u}_m\}$

$$\underbrace{(\bar{v}_1 \ \bar{v}_2 \ \dots \ \bar{v}_m)}_M = \underbrace{(\bar{u}_1 \ \bar{u}_2 \ \dots \ \bar{u}_m)}_Q \underbrace{\quad}_R$$

the change of basis matrix

QR-factorization of M : $M = QR$

Gram-Schmidt Theorem: Remarks

Remark 1:

Using the Gram-Schmidt process, we change the basis

$\mathfrak{B}_1 = \{\bar{v}_1, \dots, \bar{v}_m\}$ to the basis $\mathfrak{B}_2 = \{\bar{u}_1, \dots, \bar{u}_m\}$

$$\underbrace{(\bar{v}_1 \ \bar{v}_2 \ \dots \ \bar{v}_m)}_M = \underbrace{(\bar{u}_1 \ \bar{u}_2 \ \dots \ \bar{u}_m)}_Q \underbrace{R}_{\text{the change of basis matrix}}$$

QR-factorization of M : $M = QR$

Remark 2: How to find the change of basis matrix R .

$\forall j > 2 \quad \bar{v}_j^\perp = \bar{v}_j - (\bar{u}_1, \bar{v}_j)\bar{u}_1 - \dots - (\bar{u}_{j-1}, \bar{v}_j)\bar{u}_{j-1}$ and $\bar{v}_j^\perp = \|\bar{v}_j^\perp\| \bar{u}_j$

$$\bar{v}_j = \underbrace{(\bar{u}_1, \bar{v}_j)}_{r_{1j}} \bar{u}_1 + \underbrace{(\bar{u}_2, \bar{v}_j)}_{r_{2j}} \bar{u}_2 \dots + \underbrace{(\bar{u}_{j-1}, \bar{v}_j)}_{r_{j-1,j}} \bar{u}_{j-1} + \underbrace{\|\bar{v}_j^\perp\|}_{r_{j,j}} \bar{u}_j$$

$$R = (r_{ij}) = \begin{cases} (\bar{u}_i, \bar{v}_j) & i < j \\ 0 & i > j, \\ \|\bar{v}_j^\perp\| & i = j, j > 2 \\ \|\bar{v}_1\| & i = j = 1 \end{cases}$$

QR-factorization. Example

Theorem (QR-factorization):

If the columns $\bar{v}_1, \bar{v}_2, \dots, \bar{v}_m$ of the matrix $M_{n \times m}$ are linearly independent, then there exists a matrix $Q_{n \times m}$ with orthogonal columns $\bar{u}_1, \bar{u}_2, \dots, \bar{u}_m$ and an upper triangular matrix R with positive diagonal entries such that

$$M = QR$$

QR-factorization. Example

Theorem (QR-factorization):

If the columns $\bar{v}_1, \bar{v}_2, \dots, \bar{v}_m$ of the matrix $M_{n \times m}$ are linearly independent, then there exists a matrix $Q_{n \times m}$ with orthogonal columns $\bar{u}_1, \bar{u}_2, \dots, \bar{u}_m$ and an upper triangular matrix R with positive diagonal entries such that

$$M = QR$$

Example 1:

$$M = \begin{pmatrix} 0 & -3 & 0 \\ 0 & 0 & 0 \\ 2 & 0 & 0 \\ 0 & 0 & 4 \end{pmatrix}$$

QR-factorization. Example

Theorem (QR-factorization):

If the columns $\bar{v}_1, \bar{v}_2, \dots, \bar{v}_m$ of the matrix $M_{n \times m}$ are linearly independent, then there exists a matrix $Q_{n \times m}$ with orthogonal columns $\bar{u}_1, \bar{u}_2, \dots, \bar{u}_m$ and an upper triangular matrix R with positive diagonal entries such that

$$M = QR$$

Example 1:

$$M = \begin{pmatrix} 0 & -3 & 0 \\ 0 & 0 & 0 \\ 2 & 0 & 0 \\ 0 & 0 & 4 \end{pmatrix} = \underbrace{\begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_Q$$

QR-factorization. Example

Theorem (QR-factorization):

If the columns $\bar{v}_1, \bar{v}_2, \dots, \bar{v}_m$ of the matrix $M_{n \times m}$ are linearly independent, then there exists a matrix $Q_{n \times m}$ with orthogonal columns $\bar{u}_1, \bar{u}_2, \dots, \bar{u}_m$ and an upper triangular matrix R with positive diagonal entries such that

$$M = QR$$

Example 1:

$$M = \begin{pmatrix} 0 & -3 & 0 \\ 0 & 0 & 0 \\ 2 & 0 & 0 \\ 0 & 0 & 4 \end{pmatrix} = \underbrace{\begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_Q \underbrace{\begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{pmatrix}}_R$$

QR-factorization. Example

Example 2:

$$M = \begin{pmatrix} 2 & 2 \\ 1 & 7 \\ -2 & -8 \end{pmatrix}_{3 \times 2} \rightarrow QR$$

QR-factorization. Example

Example 2:

$$M = \begin{pmatrix} 2 & 2 \\ 1 & 7 \\ -2 & -8 \end{pmatrix}_{3 \times 2} \rightarrow QR$$

$$\bar{v}_1 = \begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix} \Rightarrow \bar{u}_1 = \begin{pmatrix} 2/3 \\ 1/3 \\ -2/3 \end{pmatrix},$$

QR-factorization. Example

Example 2:

$$M = \begin{pmatrix} 2 & 2 \\ 1 & 7 \\ -2 & -8 \end{pmatrix}_{3 \times 2} \rightarrow QR$$

$$\bar{v}_1 = \begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix} \Rightarrow \bar{u}_1 = \begin{pmatrix} 2/3 \\ 1/3 \\ -2/3 \end{pmatrix}, \bar{u}'_2 = \bar{v}_2 - (\bar{v}_2, \bar{u}_1)\bar{u}_1 =$$

$$= \begin{pmatrix} 2 \\ 7 \\ -8 \end{pmatrix} - \left(2 \cdot \frac{2}{3} + 7 \cdot \frac{1}{3} + 8 \cdot \frac{2}{3} \right) \begin{pmatrix} \frac{2}{3} \\ \frac{1}{3} \\ -\frac{2}{3} \end{pmatrix} =$$

$$= \begin{pmatrix} -4 \\ 4 \\ -2 \end{pmatrix} \Rightarrow$$

QR-factorization. Example

Example 2:

$$M = \begin{pmatrix} 2 & 2 \\ 1 & 7 \\ -2 & -8 \end{pmatrix}_{3 \times 2} \rightarrow QR$$

$$\bar{v}_1 = \begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix} \Rightarrow \bar{u}_1 = \begin{pmatrix} 2/3 \\ 1/3 \\ -2/3 \end{pmatrix}, \bar{u}'_2 = \bar{v}_2 - (\bar{v}_2, \bar{u}_1)\bar{u}_1 =$$

$$= \begin{pmatrix} 2 \\ 7 \\ -8 \end{pmatrix} - \left(2 \cdot \frac{2}{3} + 7 \cdot \frac{1}{3} + 8 \cdot \frac{2}{3} \right) \begin{pmatrix} \frac{2}{3} \\ \frac{1}{3} \\ -\frac{2}{3} \end{pmatrix} =$$

$$= \begin{pmatrix} -4 \\ 4 \\ -2 \end{pmatrix} \Rightarrow \bar{u}_2 = \begin{pmatrix} -2/3 \\ 2/3 \\ -1/3 \end{pmatrix} \Rightarrow$$

QR-factorization. Example

Example 2:

$$M = \begin{pmatrix} 2 & 2 \\ 1 & 7 \\ -2 & -8 \end{pmatrix}_{3 \times 2} \rightarrow QR$$

$$\bar{v}_1 = \begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix} \Rightarrow \bar{u}_1 = \begin{pmatrix} 2/3 \\ 1/3 \\ -2/3 \end{pmatrix}, \bar{u}'_2 = \bar{v}_2 - (\bar{v}_2, \bar{u}_1)\bar{u}_1 =$$

$$= \begin{pmatrix} 2 \\ 7 \\ -8 \end{pmatrix} - \left(2 \cdot \frac{2}{3} + 7 \cdot \frac{1}{3} + 8 \cdot \frac{2}{3} \right) \begin{pmatrix} \frac{2}{3} \\ \frac{1}{3} \\ -\frac{2}{3} \end{pmatrix} =$$

$$= \begin{pmatrix} -4 \\ 4 \\ -2 \end{pmatrix} \Rightarrow \bar{u}_2 = \begin{pmatrix} -2/3 \\ 2/3 \\ -1/3 \end{pmatrix} \Rightarrow Q = \frac{1}{3} \begin{pmatrix} 2 & -2 \\ 1 & 2 \\ -2 & -1 \end{pmatrix}$$

QR-factorization. Example

Example 2 (cont):

$$M = \begin{pmatrix} 2 & 2 \\ 1 & 7 \\ -2 & -8 \end{pmatrix}_{3 \times 2} \rightarrow QR, \quad Q = \frac{1}{3} \begin{pmatrix} 2 & -2 \\ 1 & 2 \\ -2 & -1 \end{pmatrix}$$

$$r_{11} = \|\bar{v}_1\| = 3, \quad r_{12} = (\bar{u}_1, \bar{v}_2) = 9, \quad r_{21} = 0, \quad r_{22} = \|\bar{u}'_2\| = 6$$

$$R = \begin{pmatrix} 3 & 9 \\ 0 & 6 \end{pmatrix}$$

$$M = \begin{pmatrix} 2 & 2 \\ 1 & 7 \\ -2 & -8 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 2 & -2 \\ 1 & 2 \\ -2 & -1 \end{pmatrix} \begin{pmatrix} 3 & 9 \\ 0 & 6 \end{pmatrix}$$


QR -factorization. Example

Useful Statements

Motivation: Let

$$V = \text{Im} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \Rightarrow V \text{ is a line spanned by } \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$$

$\text{Ker} \begin{pmatrix} 1 & 2 & 3 \end{pmatrix}$ is the plane $x_1 + 2x_2 + 3x_3 = 0$



Useful Statements

Theorem 1: $(\text{Im } A)^\perp = \text{Ker } (A^T)$

Useful Statements

Theorem 1: $(\text{Im } A)^\perp = \text{Ker } (A^T)$

Theorem 2: Let $A_{n \times m}: \mathbb{R}^m \rightarrow \mathbb{R}^n$. Then

$$1. \text{Ker } A = \text{Ker}(A^T A)$$

$$2. \text{Ker } A = \{\bar{0}\} \Rightarrow \exists (A^T A)^{-1}$$

Theorem

Theorem: Let $\bar{x} \in \mathbb{R}^n$ and $V \subset \mathbb{R}^n$ be a linear subspace. Then

$$\|\bar{x} - \text{proj}_V \bar{x}\| \leq \|\bar{x} - \bar{v}\|$$

Theorem

Theorem: Let $\bar{x} \in \mathbb{R}^n$ and $V \subset \mathbb{R}^n$ be a linear subspace. Then

$$\|\bar{x} - \text{proj}_V \bar{x}\| \leq \|\bar{x} - \bar{v}\|$$

Proof: Show that

$$\|\bar{x} - \bar{v}\|^2 = \|\bar{x} - \text{proj}_V \bar{x}\|^2 + \|\bar{v} - \text{proj}_V \bar{x}\|^2$$

Remark: The orthogonal projection of \bar{x} onto V is closest to \bar{x} .

Least-squares solution

Let a system $A\bar{x} = \bar{b}$ be *inconsistent*. Then \bar{b} is not in $Im(A)$.

We can try to find an **approximate solution**, i.e. a vector \bar{x}^* such that $A\bar{x}^*$ as close to \bar{b} as possible $\Rightarrow \min ||A\bar{x} - \bar{b}||$

Least-squares solution

Let a system $A\bar{x} = \bar{b}$ be *inconsistent*. Then \bar{b} is not in $Im(A)$.

We can try to find an **approximate solution**, i.e. a vector \bar{x}^* such that $A\bar{x}^*$ as close to \bar{b} as possible $\Rightarrow \min ||A\bar{x} - \bar{b}||$

Definition: A vector $\bar{x}^* \in \mathbb{R}^m$ is called a **least-squares solution** of the system $A_{n \times m} \bar{x} = \bar{b}$ if

$$||\bar{b} - A\bar{x}^*|| \leq ||\bar{b} - A\bar{x}|| \quad \forall \bar{x} \in \mathbb{R}^m$$

Remarks:

1. If the system is consistent then $||\bar{b} - A\bar{x}|| = 0$.
2. If $||\bar{b} - A\bar{x}^*|| \leq ||\bar{b} - A\bar{x}||$ then $A\bar{x}^* = proj_{Im A} \bar{b}$

Theorem

Theorem: The least-squares solutions of the system

$$A\bar{x} = \bar{b}$$

are the exact solutions of

$$A^T A\bar{x} = A^T \bar{b}$$

Theorem

Theorem: The least-squares solutions of the system

$$A\bar{x} = \bar{b}$$

are the exact solutions of

$$A^T A\bar{x} = A^T \bar{b}$$

Remarks:

1. If $\text{Ker } A = \{\bar{0}\}$, then there exists a unique solution

$$\bar{x} = \left(A^T A\right)^{-1} A^T \bar{b}$$

2. The system $A^T A\bar{x} = A^T \bar{b}$ is called the **normal equation** of $A\bar{x} = \bar{b}$
3. Let $V = \text{span}(\bar{v}_1, \dots, \bar{v}_m) \subset \mathbb{R}^n$ and $A = (\bar{v}_1 \ \bar{v}_2 \ \dots \ \bar{v}_m)$. Then the matrix of the orthonormal projection onto V is

$$A \left(A^T A\right)^{-1} A^T$$

$$\text{rank } A = \text{rank } A^T$$

$$1. \text{rank } A = \text{rank } A^T$$

$$\text{Let } A_{n \times m} \Rightarrow A: \mathbb{R}^m \rightarrow \mathbb{R}^n$$

$$a. \text{Im } A \subset \mathbb{R}^n \text{ is a linear subspace} \Rightarrow \dim \text{Im } A + \dim (\text{Im } A)^\perp = n$$

$$\dim (\text{Im } A)^\perp = n - \dim \text{Im } A = n - \text{rank } A$$

$$b. \text{ by the Rank-Nullity theorem,}$$

$$\dim \text{Ker } A + \dim \text{Im } A = m \Rightarrow \dim \underbrace{\text{Ker } A^T}_{(\text{Im } A)^\perp} + \dim \text{Im } A^T = n$$

$$\Rightarrow \dim (\text{Im } A)^\perp = n - \dim \text{Im } A^T$$

$$\Rightarrow \dim \text{Im } A^T = \text{rank } A$$

$$2. \text{rank } A = \text{rank } A^T A$$

$$\text{Ker } A = \text{Ker}(A^T A)$$

$$\dim \text{Ker } A = m - \dim \text{Im } A = m - \dim \text{Im}(A^T A) = \dim \text{Ker}(A^T A)$$

$$\dim \text{Im } A = \dim \text{Im}(A^T A) \Rightarrow \text{rank } A = \text{rank } A^T A$$

Correlation

Def: Let $\bar{x}, \bar{y} \in \mathbb{R}^n$. There is a **positive correlation** between \bar{x} and \bar{y} if and only if $(\bar{x}, \bar{y}) > 0$.

Correlation

Def: Let $\bar{x}, \bar{y} \in \mathbb{R}^n$. There is a **positive correlation** between \bar{x} and \bar{y} if and only if $(\bar{x}, \bar{y}) > 0$.

Def: The **correlation coefficient** r of two vectors \bar{x} and \bar{y} is

$$r = \cos(\bar{x}, \bar{y}) = \frac{(\bar{x}, \bar{y})}{|\bar{x}||\bar{y}|}$$

Remark: By the Cauchy-Schwarz inequality,

$$|(\bar{x}, \bar{y})| \leq |\bar{x}||\bar{y}| \Rightarrow -1 \leq r \leq 1$$

Correlation: Example

Consider meat consumption and incidence of cancer rate in the following countries:

Country	Consumption	Rate	Deviation: Cons	Deviation: Rate
Japan	26	7.5	-122	-10.7
Finland	101	9.8	-47	-8.4
Israel	124	16.4	-24	-1.8
GB	205	23.3	57	5.1
US	284	34	136	15.8
Mean	148	18.2		

The correlation coefficient is

$$r = \frac{122 \cdot 10.7 + 47 \cdot 8.4 + 24 \cdot 1.8 + 57 \cdot 5.1 + 136 \cdot 15.8}{198.53 \cdot 21.539} \approx 0.9782$$

Problems

1. Consider

$$\bar{u}_1 = (1/2, 1/2, 1/2, 1/2), \bar{u}_2 = (1/2, 1/2, -1/2, -1/2),$$

$$\bar{u}_3 = (1/2, -1/2, 1/2, -1/2)$$

in \mathbb{R}^4 . Can you find a vector \bar{u}_4 such that the vectors $\bar{u}_1, \bar{u}_2, \bar{u}_3, \bar{u}_4$ are orthonormal?

Problems

1. Consider

$$\bar{u}_1 = (1/2, 1/2, 1/2, 1/2), \bar{u}_2 = (1/2, 1/2, -1/2, -1/2),$$

$$\bar{u}_3 = (1/2, -1/2, 1/2, -1/2)$$

in \mathbb{R}^4 . Can you find a vector \bar{u}_4 such that the vectors $\bar{u}_1, \bar{u}_2, \bar{u}_3, \bar{u}_4$ are orthonormal?

2. Let

$$W = \text{span}((1, 2, 3, 4); (5, 6, 7, 8))$$

Find a basis for W^\perp .

Problems

1. Consider

$$\bar{u}_1 = (1/2, 1/2, 1/2, 1/2), \bar{u}_2 = (1/2, 1/2, -1/2, -1/2),$$

$$\bar{u}_3 = (1/2, -1/2, 1/2, -1/2)$$

in \mathbb{R}^4 . Can you find a vector \bar{u}_4 such that the vectors $\bar{u}_1, \bar{u}_2, \bar{u}_3, \bar{u}_4$ are orthonormal?

2. Let

$$W = \text{span}((1, 2, 3, 4); (5, 6, 7, 8))$$

Find a basis for W^\perp .

3. Let

$$V = \text{Im} \begin{pmatrix} 1 & 1 \\ 1 & -1 \\ 1 & -1 \\ 1 & 1 \end{pmatrix}$$

Find $\text{proj}_V \bar{x}$, $\bar{x} = (1, 3, 1, 7)$

Problems

4. Let $L = \text{span} \left(1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots \right) \subset l^2$. Find the orthogonal projection of $(1, 0, 0, \dots)$ onto L .

Problems

4. Let $L = \text{span} \left(1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots \right) \subset l^2$. Find the orthogonal projection of $(1, 0, 0, \dots)$ onto L .
5. Among all the vectors in \mathbb{R}^n whose components add up to 1, find the vector of minimal length.

Problems

4. Let $L = \text{span} \left(1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots \right) \subset l^2$. Find the orthogonal projection of $(1, 0, 0, \dots)$ onto L .
5. Among all the vectors in \mathbb{R}^n whose components add up to 1, find the vector of minimal length.
6. Among all the unit vectors in \mathbb{R}^n , find the one for which the sum of the components is maximal.

Problems

4. Let $L = \text{span} \left(1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots \right) \subset l^2$. Find the orthogonal projection of $(1, 0, 0, \dots)$ onto L .
5. Among all the vectors in \mathbb{R}^n whose components add up to 1, find the vector of minimal length.
6. Among all the unit vectors in \mathbb{R}^n , find the one for which the sum of the components is maximal.
7. There are three exams in your linear algebra class, and you theorize that your score in each exam (out of 100) will be numerically equal to the number of hours you study for that exam. The three exams count 20, 30, and 50, respectively, toward the final grade. If your (modest) goal is to score 76 in the course, how many hours a , b and c should you study for each of the three exams to minimize quantity $a^2 + b^2 + c^2$?