

Vv557 Methods of Applied Mathematics II

Green Functions and Boundary Value Problems

Assignment 5

Date Due: 12:55 PM, Wednesday, the 31st of March 2021

Discussion Class Preparation

Please (re-)view Video files 29-31 and/or finish reading the section “Adjoint BVPs and Higher-Order Equations” in the lecture slides. You should be able to answer the following questions:

- i) How are adjoint boundary conditions defined?
- ii) What is the adjoint boundary value problem to a given BVP?
- iii) What is the adjoint Green function?
- iv) Explain the role of the conjunct in constructing the adjoint BVP.
- v) Compare the solution formula using the conjunct with the solution formula obtained previously.

Exercises (15 Marks)

Exercise 5.1

We want to find a fundamental solution of the stationary equation for a simply supported beam, i.e., a function $g(x, \xi)$ satisfying

$$\frac{d^4 g}{dx^4} = \delta(x - \xi), \quad 0 < x, \xi < 1,$$

with boundary conditions

$$g(0, \xi) = g''(0, \xi) = g(1, \xi) = g''(1, \xi) = 0.$$

- i) Find a causal fundamental solution, i.e., a function E satisfying

$$\frac{d^4 E}{dx^4} = \delta(x - \xi), \quad 0 < x, \xi < 1,$$

and $E(x) = 0$ for $x < \xi$.

(3 Marks)

- ii) Add a solution of the homogeneous equation $\frac{d^4 u}{dx^4} = 0$ to E to obtain a function that satisfies the boundary conditions.

(2 Marks)

Exercise 5.2

Consider again the stationary equation for a traveling wave with wavenumber k ,

$$-\frac{d^2 g}{dx^2} - k^2 g = \delta(x - \xi), \quad 0 < x, \xi < 1,$$

this time with boundary conditions

$$g(0, \xi) = g(1, \xi) = 0.$$

Use the fundamental solution obtained in class via the Fourier transform to find a function that satisfies the boundary condition. Do this by adding a suitable solution of the homogeneous equation $-\frac{d^2 u}{dx^2} - k^2 u = 0$.

(2 Marks)



JOINT INSTITUTE

交大密西根学院

Exercise 5.3

Consider the boundary value problem operator given by

$$L = \frac{d^2}{dx^2}, \quad 0 < x < 1, \quad B_1 u = u(0).$$

Characterize M^* by three boundary functionals.

(2 Marks)

Exercise 5.4

Consider the boundary value operator given by

$$L = \frac{d^4}{dx^4}, \quad 0 < x < 1, \quad B_1 u = u(0), \quad B_2 u = u'''(0), \quad B_3 = u(1), \quad B_4 = u''(1)$$

- i) Find $g(x, \xi)$.
(2 Marks)
- ii) It is obvious that $L = L^*$. Find the adjoint boundary conditions and calculate $g^*(x, \xi)$.
(3 Marks)
- iii) Show that $g(x, \xi) \neq g(\xi, x)$.
(1 Mark)